A collaborative approach to improving representation in viral genomic surveillance

Paul Y. Kim[1]*, Audrey Y. Kim[1], Jamie J. Newman[2], Eleonora Cella[3], Thomas C. Bishop[4], Peter

J. Huwe[5], Olga N. Uchakina[5], Robert J. McKallip[5], Vance L. Mack[6], Marnie P. Hill[6], Ifedayo

Victor Ogungbe[7], Olawale Adeyinka[7], Samuel Jones[8], Gregory Ware[9], Jennifer Carroll[9], Jarrod

F. Sawyer[9], Kenneth H. Densmore[9], Michael Foster[2], Lescia Valmond[1], John Thomas[1], Taj

Azarian[3], Krista Queen[9], Jeremy P. Kamil[9,10]*

[1] Department of Biological Sciences, Grambling State University, Grambling, LA

[2] School of Biological Sciences, Louisiana Tech University, Ruston, LA

[3] Burnett School of Biomedical Sciences, University of Central Florida, Orlando, FL

[4] Physics and Chemistry Programs, Louisiana Tech University, Ruston, LA

[5] Mercer University School of Medicine, Macon, GA

[6] Mercer Medicine, Macon, GA

[7] Department of Chemistry, Jackson State University, Jackson, MS

[8] Health Services Center, Jackson State University, Jackson, MS

[9] Center of Excellence for Emerging Viral Threats, Louisiana State University Health Shreveport,

Shreveport, LA

[10] Department of Microbiology and Immunology, Louisiana State University Health Shreveport,

Shreveport, LA

* Corresponding authors

Email: kimp@gram.edu (PYK), jeremy.kamil@lsuhs.edu (JPK)

Abstract

The lack of routine viral genomic surveillance delayed the initial detection of SARS-CoV-2, allowing the virus to spread unfettered at the outset of the U.S. epidemic. Over subsequent months, poor surveillance enabled variants to emerge unnoticed. Against this backdrop, long-standing social and racial inequities have contributed to a greater burden of cases and deaths among minority groups. To begin to address these problems, we developed a new variant surveillance model geared toward building microbial genome sequencing capacity at universities in or near rural areas and engaging the participation of their local communities. The resulting genomic surveillance network has generated more than 1,000 SARS-CoV-2 genomes to date, including the first confirmed case in northeast Louisiana of Omicron, and the first and sixth confirmed cases in Georgia of the emergent BA.2.75 and BQ.1.1 variants, respectively. In agreement with other studies, significantly higher viral gene copy numbers were observed in Delta variant samples compared to those from Omicron BA.1 variant infections, and lower copy numbers were seen in asymptomatic infections relative to symptomatic ones. Collectively, the results and outcomes from our collaborative work demonstrate that establishing genomic surveillance capacity at smaller academic institutions in rural areas and fostering relationships between academic teams and local health clinics represent a robust pathway to improve pandemic readiness.

Author summary

Genomic surveillance involves decoding a pathogen's genetic code to track its spread and evolution. During the pandemic, genomic surveillance programs around the world provided valuable data to scientists, doctors, and public health officials. Knowing the complete SARS-CoV-2 genome has helped detect the emergence of new variants, including ones that are more

transmissible or cause more severe disease, and has supported the development of

diagnostics, vaccines, and therapeutics. The impact of genomic surveillance on public health

depends on representative sampling that accurately reflects the diversity and distribution of

populations, as well as rapid turnaround time from sampling to data sharing. After a slow start,

SARS-CoV-2 genomic surveillance in the United States grew exponentially. Despite this, many

rural regions and ethnic minorities remain poorly represented, leaving significant gaps in the

data that informs public health responses. To address this problem, we formed a network of

universities and clinics in Louisiana, Georgia, and Mississippi with the goal of increasing SARS-

CoV-2 sequencing volume, representation, and equity. Our results demonstrate the advantages

of rapidly sequencing pathogens in the same communities where the cases occur and present a

model that leverages existing academic and clinical infrastructure for a powerful decentralized

genomic surveillance system.

38 Introduction

In late 2019, a newly emergent betacoronavirus, severe acute respiratory syndrome

coronavirus 2 (SARS-CoV-2), was identified as the etiological agent associated with an

outbreak of viral pneumonia in Wuhan, China [1]. The outbreak quickly grew to become a global

public health emergency, now commonly referred to as the COVID-19 pandemic. Between

November 2020 and January 2021, the first variants of concern (VOC) Alpha, Beta, and

Gamma, were detected in England, South Africa, and Brazil and Japan, respectively. All three

carried convergent Spike (S) N501Y substitutions, while Beta and Gamma shared substitutions

S:K417N/T and S:E484K. It is now appreciated that SARS-CoV-2 variants can be more

transmissible [2], differ in pathogenicity [3,4], and/or escape immunity afforded by vaccination or

by infection with earlier variants [5,6]. Although the Omicron lineage has displaced Delta [7],

new Omicron sub-lineages continue to emerge [8], certain of which harbor concerning mutations

such as S:F486V that escape broadly neutralizing antibodies [9,10]. Overall, the pandemic

remains largely uncontrolled in virtually all geographies, highlighting the need for ongoing

surveillance to inform public health responses.

New SARS-CoV-2 variants are detected and tracked using viral genomic surveillance,

an approach wherein the genomes from circulating viruses are sequenced, usually from patient

nasal swabs, and data is rapidly shared using global platforms, such as GISAID

(www.gisaid.org) [11] or NCBI. These practices collectively enable scientists around the world to

track viral evolution across space and time while keeping tabs on mutations that can

compromise the utility of diagnostic assays or negatively impact the clinical efficacy of vaccines

or monoclonal antibody therapeutics. Effective surveillance relies on prompt viral whole genome

sequencing of recent samples across representative geographies and populations and low

latency sharing of resulting data. Accordingly, investments in viral whole genome sequencing

have increased greatly during the pandemic, with vastly more whole genome sequences of

SARS-CoV-2 being shared than any virus in history.

At the outset of the pandemic, SARS-CoV-2 genome sequencing efforts were patchy at

best and in many geographies non-existent. Although the U.K. and South Africa set up

exemplary viral genomic surveillance at the outset, much of the initial U.S. viral genomic

surveillance data came from individuals who generated it at their own initiative, typically relying

on discretionary funds, and in some cases, support from private foundations and philanthropies.

Once the first variants of concern emerged, the U.S. Centers for Disease Control and the

National Institutes of Health began to invest in the development of a SARS-CoV-2 genomic

surveillance system. Despite this, many states in the Midwest and Southern U.S. remain poorly

represented in the genomic surveillance data [12] even though their residents have been

disproportionately impacted by COVID-19 [13–15]. This rural-urban disparity was compounded

during the pandemic for racial and ethnic minorities [16,17] who already bore a greater burden

of disease [18]. The failure to adequately track emerging variants in a representative fashion

may have exacerbated existing disparities in allocation of public health resources, such as rapid

antigen tests and monoclonal antibody therapies.

To address this lack of representation, Grambling State University (GSU), Louisiana

Tech University (Tech), and Louisiana State University Health Shreveport (LSUHS) established

a viral genomic surveillance hub to reach several vulnerable and underserved populations in

north Louisiana [19]. Soon after, we partnered with Mercer University School of Medicine

(MUSM) in Georgia, Jackson State University (JSU) in Mississippi, as well as with community

health centers in each of our areas to increase SARS-CoV-2 genome sequencing volume and

representativeness. These efforts resulted in a new viral genomic surveillance network serving a

total of 15 counties or parishes across these 3 southern U.S. states (Figure 1).

Fig 1. Geographic coverage of SARS-CoV-2 genomic surveillance in Louisiana, Georgia,

and Mississippi. Map of the surveillance region with parishes (Louisiana) or counties (Georgia,

Mississippi) where at least one specimen was sequenced by the network indicated in red.

Our primary goals were to improve sampling from rural communities and minorities, to

empower underrepresented minorities conducting the scientific work of viral genomic

surveillance, and to provide individuals who donate samples for viral genome sequencing the

opportunity to do so with consent and an understanding of how their samples would be used to

track viral variants. The pandemic has underscored the importance of contextual metadata

accompanying genomic data. When metadata is incomplete, it hinders comparative analyses

and data-driven policymaking [20,21]. Thus, an additional objective was to collect more

complete metadata to maximize the usefulness of the shared sequences. Here we used patient

metadata to assess the relationships between viral gene copies (as an index of viral load) and

variant type, vaccination status, and symptomatology. We report on the results of our work in

Louisiana, Georgia, and Mississippi.

Results and discussion

Improving representativeness in genomic surveillance

In Louisiana, our six partner clinics collected 405 rapid antigen-positive clinical

specimens from the residents of nine parishes (Allen, Bienville, Franklin, Jackson, Lincoln,

Morehouse, Ouachita, Union, and Webster) between 22 July 2021 and 12 April 2022 (Fig 1).

We worked closely with clinicians at the collection sites to gain informed consent from patients

and collected metadata on patient demographics and vaccination status for more than 90% of

the specimens (Table 1).

Table 1. Characteristics of study donors in Louisiana (22 July 2021 – 12 April 2022).

CHARACTERISTICS	NUMBER OF DONORS (%)* n = 405
Median age (range)	22 (1 – 86) years
Sex	
Male	180 (44.4)
Female	215 (53.21)
Not disclosed	10 (2.5)
Race/ethnicity	
White	99 (24.4)
Black	205 (50.6)
Hispanic	34 (8.4)
Asian	4 (<1)
Pacific Islander	1 (<1)
Alaska Native	1 (<1)
Other	1(<1)
Not disclosed	66 (16.2)
PCR	
Negative	15 (3.7)
Positive	389 (96)
Indeterminate	1 (<1)
Vaccination	
No	209 (51.6)

Yes	170 (42)
Partial †	26
Full ‡	113
Booster §	18
Not specified ¶	13
Not disclosed	26 (6.4)
Previous infection	
No	321 (79.3)
Yes	41 (10.1)
Not disclosed	43 (10.6)
Symptomatic	
No	52 (12.8)
Yes	336 (83)
Not disclosed	17 (4.2)

*unless otherwise indicated
Vaccination:
† one dose mRNA
‡ two dose mRNA or one dose Janssen
§ three dose mRNA
¶ specific type and date(s) of vaccination unknown
119

In these nine parishes, between 24.2% (Ouachita) to 82.9% (Union) of the residents live

in rural areas as defined by the U.S. Census Bureau [22]. Rural areas may be especially

vulnerable to the pandemic because they tend to have less capacity for diagnostic testing and

healthcare services [23,24] while being home to a poorer, older, health-compromised population

[25]. Bienville and Franklin parishes in particular ranked in the top 20 of the 300 poorest

counties with the highest COVID-19 mortality rates in a recent analysis of 3,200 U.S. counties

and the majority of the other parishes in this region were likewise characterized as low income-

high mortality [26].

The disproportionate impact of COVID-19 on racial and ethnic minorities has also been

well documented [27–29]. More than 60% of our donors self-identified as non-white or as two or

more races compared to approximately 41% overall in the sampled parishes. In our view, this

oversampling supports equitable genomic surveillance by compensating for known biases in

sampling [30] from both rural and minority populations. Notably, 8.4% of our specimens were

from Hispanic donors who make up just 3.1% of the northeast Louisiana population but were far

more likely to contract COVID-19, accounting for 13.8% of overall cases in the region (up to

60% in some parishes) according to the Louisiana Department of Health. Our partnership with

The Health Hut mobile clinic staffed with Spanish-speaking translators improved

representativeness in this population. Of the 405 specimens that were collected in Louisiana,

389 had detectable SARS-CoV-2 RNA in the CDC real-time RT-PCR diagnostic assay and

underwent sequencing, with genomes ultimately recovered from 272 or 70% of PCR-positive

specimens. This represents 1.32% of the 20,617 genomes in GISAID from Louisiana during the

same period.

Accelerating regional genomic surveillance

After successfully establishing relationships with local clinics and developing a pipeline

for exchanging specimens and data across our network in Louisiana, we felt confident sharing

our model with other peer institutions across the south. Mercer University School of Medicine

(MUSM) in Georgia and Jackson State University (JSU) in Mississippi both joined our efforts,

building on relationships to engage their own communities in genomic surveillance.

In Georgia, MUSM specimens were collected from Mercer Medicine (MM) Student

Health services in Bibb, Chatham, DeKalb, and Fulton counties and an MM community clinic

located in Peach county (Fig 1). Roughly 81% of the Georgia specimens were obtained from

counties outside of the Atlanta metropolitan area. Approximately 48% of MM donors did not self-

identify as white. After testing positive via PCR analysis, specimens were sent to the Center of

Excellence for Emerging Viral Threats (CEVT) at LSUHS for Illumina sequencing.

Among other achievements, Mercer's work identified two emerging subvariants of BA.5

carrying immune escape mutations [31,32]: the first case of BA.2.75 in Georgia, which was only

the 40th documented instance of the variant in the U.S., and the sixth case of BQ.1.1 in

Georgia, one of the most antibody-evasive variants to date [31]. MM generated 703 SARS-CoV-

2 genomes representing 1.8% of the total data in GISAID from Georgia between 13 September

2021, when the first sample was collected by MM to 10 October 2022.

In Mississippi, the JSU Health Services Center served as the collection site (Fig 1). The

Center serves as the primary COVID-19 testing site for students, staff, and visitors of the

university and supports the surrounding community's testing and vaccination efforts. The

engagement with JSU extended the network's reach into a predominantly black community.

One hundred percent of the 33 samples submitted for sequencing at the CEVT were from non-

white donors.

168 Building NGS capacity for local genomic surveillance

The specimens collected in Louisiana were initially submitted to the CEVT for

sequencing until we built local sequencing capacity at GSU and Tech. By sequencing locally,

we achieved a median latency, or turnaround time from sample collection to release on GISAID,

of 5 days at GSU and 12 days at Tech when processing contemporary rather than previously

stored specimens (Fig 2B). Through this approach, we were able to report the first confirmed

case of Omicron in northeast Louisiana, a region comprising 12 parishes and 323,000 residents,

within 2 days of sample collection [33]. We then closed the surveillance loop by sharing the

variant data along with weekly wastewater surveillance results and other COVID-19-related

health information through a dashboard developed for the north Louisiana community (www.nla-

health.com/dashboard).

Fig 2. SARS-CoV-2 genomic surveillance in Louisiana. (A) Number of variants sequenced

over time in Louisiana overlaid with number of confirmed COVID-19 cases in the U.S. (B)

Number of sequences uploaded and latency for new sequencing entities established at

Grambling State University (GSU) and Louisiana Tech University (Tech).

The results of sequencing, including comparison of Nanopore to Illumina, investigation of

viral genome coverage and copy number, and phylogenetic analyses are described below.

Nanopore concordance with Illumina

Among the various approaches to whole genome sequencing of viruses [34], multiplex

PCR amplification of the viral genome followed by amplicon sequencing is the most widely

adopted for SARS-CoV-2 genome sequencing. The second-generation Illumina platform

dominates the next generation sequencing (NGS) field, but third-generation Oxford Nanopore

technology has attracted attention during the pandemic in large part for its low costs of entry,

real-time analysis, and amenability to low-throughput use [35].

We first validated the performance of Nanopore MinION sequencing at GSU and Tech

by sequencing a matched subset of 16 SARS-CoV-2 positive specimens using a well-

established Illumina workflow at the CEVT. The results confirmed the accuracy of the Nanopore

sequencing protocol and analysis pipeline in our setting with 100% consensus sequence identity

in 13/16 of the samples and 99.9% sequence identity in 3/16 samples. The one discordant

basecall was identical in the three samples: T by Illumina sequencing and G by Nanopore

sequencing at 17,259, within the gene encoding the replicase accessory protein NSP13. The

reference SARS-CoV-2 genome also contains a G at this position; the T mutation identified by

Illumina sequencing results in a glutamic acid rather than an aspartic acid. The E1264D

mutation identified in the samples sequenced with the Illumina protocol is well supported.

Nanopore basecalling accuracy decreases in homopolymeric tracts [36] but the reason for the

discordance observed in this region is unclear.

Importantly, all sequences were 100% concordant for the Pango lineages and GISAID

clade assignments. The 16 samples sequenced by both protocols spanned a wide range of C_q

values (15.6 – 32.4) but almost all samples, Nanopore or Illumina, generated > 98% genome

coverage. Illumina sequencing resulted in an average of 285 bases more sequence data than

Nanopore, but this was mainly confined to the extreme 5' and 3' ends of the genome.

Viral genome coverage

Plotting the breadth of coverage for Nanopore sequencing reveals a threshold of

approximately C_q 30 for reliably recovering a full genome in our setting and an upper limit of

approximately C_q 35 (Fig 3). These coverage thresholds are congruent with values from the

literature of different NGS methodologies for SARS-CoV-2 genome sequencing [37], although

some groups using the Nanopore platform have reported high coverage consensus genome

sequences from specimens with C_q values up to 39 [38]. Variability in primer design and

amplification parameters, library preparation method, and sequencing run time can influence

results. For example, longer amplicons reduce the likelihood of primer interactions [39] and

primer mismatches due to mutations [40], generally outperforming shorter amplicons [41,42].

Shorter amplicons, however, can recover more complete genomes from specimens where the

viral load is low or the RNA is degraded [43,44].

Fig 3. Decreasing genome breadth of coverage in samples with high C_q values. Higher C_q

values were observed in Omicron (n = 134) than in Delta (n = 27). All but one Delta specimen

was amplified for 32 cycles and all Omicron specimens were amplified for 35 – 40 cycles.

Specimens carrying low viral loads can have uneven genome coverage due to poor

amplification of the less efficient amplicons, which can be addressed to some extent by

optimizing the PCR and sequencing protocols. Lagerborg *et al.* found that increasing PCR

cycles from 35 to 40 resulted in more uniform coverage for Illumina-based ARTIC sequencing of

low viral load specimens while still avoiding PCR artifacts [44]. Compared to the Nanopore rapid

barcoding kit used for library preparation in our study, the more labor-intensive Nanopore

ligation sequencing kit can yield greater sequencing depth and better (*de novo*) assembly of the

SARS-CoV-2 genome [45]. Of course, longer run times can produce more mapped reads to

help recover complete genomes.

240   Viral gene copies

The fast-spreading Omicron variant displaced the Delta variant in our sampling with little

overlap in prevalence (Fig 2A). If we infer that low coverage, unclassified sequences are the

predominant variant at the time of sample collection, we find that the mean C_q value of Omicron

specimens (29.87, 95% CI 29.07 – 30.67) was significantly higher than that of Delta specimens

(25.55, 95% CI 23.38 – 27.72). This corroborates early reports of lower viral gene copy number

determined by RT-qPCR [46,47] or lower infectious viral load determined by focus forming

assay [48] in Omicron infections compared to Delta infections.

One possible explanation for the observed difference in gene copy number/imputed viral

load is that during the initial Omicron surge, a greater proportion of the population may have

acquired some degree of immunity through vaccination or prior infection. Some reports support

the idea of attenuated viral load with vaccination [49–51] while others suggest no significant

difference between unvaccinated and vaccinated individuals [52–54].

Having collected vaccination and prior infection metadata from 379 and 362 of the 405

donors respectively, we compared immune history with viral gene copy numbers. We found that

the mean C_q value of unvaccinated individuals with no previous infection (n = 82) was 1.06

cycles lower than that of fully vaccinated individuals with or without a previous infection (n = 81),

but this difference was not statistically significant (p = 0.218) (Fig 4A).

Fig 4. Viral gene copies by vaccination status and symptomatology. (A) No statistically

significant differences (p = 0.218) in viral gene copies were observed between unvaccinated

donors without a previous infection (n = 82) and vaccinated donors with or without a previous

infection (n = 81). (B) Viral gene copies were lower (p = 0.0039) in asymptomatic donors (n =

30) compared to symptomatic donors (n = 164).

Several groups have found no correlation between RT-qPCR quantification of viral

genes and symptomatology [55–58]. Two studies reported higher C_t values in asymptomatic

individuals (n = 5 or 3) compared to symptomatic individuals (n = 14 or 9) but the sample sizes

were limited [59,60]. In a larger study, asymptomatic individuals (n = 2179) exhibited higher C_t

values compared to symptomatic individuals (n = 739), but the authors concluded that the

statistically significant difference of 0.71 cycles was not clinically meaningful [61]. In our dataset,

the mean C_q value of asymptomatic donors (31.36, 95% CI 29.7 – 33.01; n = 30) was nearly 3

cycles higher than that of symptomatic donors (28.38, 95% CI 27.56 – 29.19; n = 164) (Fig 4B).

The asymptomatic donors in this study were mostly college students who tested positive

during baseline surveillance on the GSU campus, so the age distribution is highly skewed

compared to symptomatic donors. There are conflicting reports on the relationship between

patient age and viral load, but one of the largest cross-sectional studies found that viral load

tends to increase with age [62]. To account for a possible age effect, we compared

asymptomatic donors who were age 30 or younger (n = 28) with symptomatic donors in the

same age group (n = 93) and found that the asymptomatic group still had lower mean viral gene

copy number (31.78, 95% CI 30.17 – 33.39 vs 28.55, 95% CI 27.39 – 29.7).

Only C_q values derived at GSU were used in these analyses to avoid potential

confounding by interlaboratory variation in PCR assays [63]. These analyses should be

considered carefully within the limitations of our study which include a relatively small sample

size and lack of longitudinal information on peak copy number, and also within the overarching

context that copy number is an imprecise index of viral load [64].

Phylogenetic analysis of northeast Louisiana SARS-CoV-2

genomes

We performed independent phylogenetic analysis for Delta and BA.1.x SARS-CoV-2

viral populations from Louisiana and Georgia in the context of representative global samples.

Time-scaled maximum likelihood phylogenies revealed that Delta and BA.1.x isolates from

Louisiana and Georgia are interspersed throughout the global population suggesting multiple

independent introductions (Fig 5). Often, these introductions resolve into distinct subclades

indicative of ongoing transmission within a new location. Further, putative introductions were

spatially diverse and included Europe and South America as well as other states throughout the

US. These local epidemics were nested within the larger North American epidemic for both

Delta and BA.1.x waves.

Fig 5. Time-resolved Maximum Likelihood (ML) phylogenies of Delta and Omicron

(BA.1.x) sublineages. The genomes generated in this study are colored in red for Georgia and

blue for Louisiana. Tree tips are colored according to their location (continent) and the color

legend is on the top of the figure.

Conclusion

NGS at scale

Nanopore sequencing has been used for real-time genomic surveillance during outbreaks of Ebola in West Africa [65], Zika [66] and Yellow Fever [67] in Brazil, and Lassa fever in Nigeria [68]. During the COVID-19 pandemic, laboratories around the world applied Nanopore sequencing to rapidly share SARS-CoV-2 genomic data. The Nanopore platform is especially suitable for smaller institutions due to its low capital and operational cost, accessible infrastructure requirements, and high degree of scalability. A modestly equipped molecular biology laboratory with the appropriate biosafety containment can implement the Nanopore rapid barcoding workflow to go from swab to data for less than $3500 USD at the time of this writing. The operational cost (excluding RNA extraction) has been estimated at £18.91 per ligated barcode for 12 barcodes on a MinION flow cell when using a wash kit to remove the previous library [69]. This corresponds with our cost analysis (*including* RNA extraction) of $20 – $25 USD per tagmented barcode for 12 barcodes when the flow cell is washed and reused.

Working with NGS data can demand significant computational resources and bioinformatics expertise, but Nanopore-based SARS-CoV-2 sequencing can be performed on a consumer-class PC using prepared workflows including point-and-click pipelines in the cloud. Although higher throughput laboratories can multiplex 96 barcodes for less than £10 per barcode [69], the platform does allow small-volume laboratories to cost-effectively conduct genomic surveillance in areas of low sequencing capacity.

The democratization of genomics is nowhere better demonstrated than in Africa where technological advances in NGS coupled with major investments by Africa CDC and partners rapidly expanded genomic surveillance across the continent [70]. The efforts in Africa highlight some parallels from our experience in the rural Southern U.S., that local sequencing programs

328   can close surveillance blind spots while significantly decreasing turnaround time. This is

329   exemplified by our sequencing the first confirmed case of Omicron in northeast Louisiana and

330   detecting the emergence of BA.2.75 and BQ.1.1 in Georgia.

332   Collaborative community-centered genomic surveillance networks

As part of a broader initiative supported by the Rockefeller Foundation's Pandemic

Prevention Institute, we established a genomic surveillance network in our community focused

on equitable sequencing among a more vulnerable, marginalized population. We formed

collaborations between R1 institutions, primarily undergraduate universities, a graduate medical

school, Historically Black Colleges and Universities, and community health centers to collect

and sequence respiratory specimens from patients in rural Louisiana, Georgia, and Mississippi.

These three states rank 41st, 43rd, and 45th respectively among U.S. states in the percentage

of cumulative cases sequenced [12]. With a modest investment from the Rockefeller Foundation

and internal support from our institutions, we increased sequencing volume from these states so

that sequenced samples better represent the diversity of the U.S. population. We collected

enriched metadata including vaccination status and county/parish-level geographical location,

which is particularly important when emerging variants such as BA.2.75 and BQ.1.1 are

detected. In both the GSU and Tech initiatives, health care workers at partnering clinics briefly

educated patients about how viral genomic surveillance is used to track emerging variants and

provided patients with the opportunity to provide consent for their positive sample to be used for

whole viral genome sequencing.

Rural residents face significant barriers rooted in geographic isolation, scarcity of

services, lower socioeconomic status and other social determinants of health, and even cultural

constraints that impede them from getting the healthcare that they need [71]. During the

pandemic, this disparity manifested as disproportionately high mortality rates in rural America,

particularly among minorities [72]. In rural areas, so-called regional universities represent

essential infrastructure that can provide not only educational opportunities but also access to

testing and vaccination. These universities are uniquely positioned to gain trust and encourage

participation in genomic surveillance because they often function as community anchors, vital to

the local economy, civics, and culture [73]. A viral genomic surveillance strategy that leverages

these strong community ties also offers a much-needed on-ramp for students from diverse

groups to gain genomics research experience that will prepare them to combat misinformation

in their own communities and participate in the genomics workforce.

Our work demonstrates the impact that universities have within their local communities,

the power of the relationships that can be built between academia and clinics, and the positive

influence of these partnerships on public health. We propose that these types of nimble, local

networks should continue to play an important role in national efforts to modernize viral genomic

surveillance of common respiratory illnesses by filling in critical gaps in genomic surveillance

coverage, consent, *and* participation.

Methods

Specimen collection

Patients testing positive (n = 405) on the BinaxNOW rapid antigen test (Abbott

Laboratories; Abbott Park, IL) at the Foster Johnson Health Center in Grambling State

University (Grambling, LA), The Health Hut (Ruston, LA), Louisiana Tech University (Ruston,

LA), Michael Brooks Family Clinic (Ruston, LA), Minden Family Care Center (Minden, LA), and

Serenity Springs Specialty Hospital (Ruston, LA) between 23 August 2021 and 12 April 2022

were recruited through an informed consent process. Remnant anterior nasal swabs used in the

rapid test (The Health Hut, Michael Brooks Family Clinic, Louisiana Tech University, Minden

Family Care Center, Serenity Springs Specialty Hospital) or freshly collected mid-turbinate nasal

swabs (Foster Johnson Health Center) were preserved in 3 mL of viral transport media (VTM) at

4 °C and later in the study in 1 mL of DNA/RNA Shield at room temperature. The work at GSU

was conducted under 45 CFR 46.102(l)(2) as a public health surveillance activity and exempt

from IRB review. The work at Tech was conducted under IRB protocol HUC 21-106. LDT

COVID-19 IMD$_A$ specimen collection at Mercer University (Macon, GA) was conducted under

IRB protocol H2110206 using previously collected anterior nasopharyngeal swabs or mid-

turbinate nasal swabs, and the samples were sequenced at LSUHS CEVT. Specimen collection

at Jackson State University (Jackson, MS) was conducted under IRB protocol 0097-22 using

either anterior nasal swabs or mid-turbinate nasal swabs, and the samples were sequenced at

LSUHS CEVT. Sequencing at LSUHS CEVT was conducted under IRB protocol #

STUDY00001445.

Illumina sequencing (LSUHS CEVT)

Total RNA was isolated from VTM within 72 hours of specimen collection using the

Omega Bio-Tek Viral RNA Xpress kit (Norcross, GA) on an automated extraction platform.

Presence of SARS-CoV-2 RNA was determined by using CDC primers and probes with

LunaScript RT Supermix Kit (NEB) run on BioRad (Hercules, CA) C1000 real-time PCR

machines. cDNA, PCR amplification, and library preparation was performed using the NEBNext

ARTIC SARS-CoV-2 FS kit (NEB) according to the manufacturer's instructions. Libraries were

quantified using Invitrogen Qubit BR dsDNA kit and size determined on an Agilent Tapestation

2200 using high sensitivity d1000 tapes (Santa Clara, CA). Libraries were loaded at 10.5 pM on

an Illumina (San Diego, CA) MiSeq version 2, 300 cycle kit on an Illumina MiSeq instrument.

Consensus sequences were generated with an in-house data analysis pipeline using bwa,

samtools, and ivar.

Nanopore sequencing (GSU and Tech)

Total RNA was extracted from VTM or DNA/RNA Shield within 48 hours of specimen collection via the QIAmp Viral RNA Minikit (Qiagen; Germantown, MD) or Zymo Quick-DNA/RNA Viral MagBead Kit (Zymo Research; Irvine, CA) according to the manufacturers' protocols. Briefly, 500 uL of VTM was centrifuged at 5,000 g for 10 minutes to pellet cells and debris and either 140 uL of supernatant was extracted in 560 uL of Buffer AVL containing 560 ug of carrier RNA (QIAmp) or 200 uL of supernatant was extracted in 400 uL of Viral DNA/RNA Buffer followed by addition of 10 uL MagBinding Beads (Zymo). After washing the spin column or beads with the manufacturers' respective wash buffers, total RNA was eluted from the column in 60 uL of Buffer AVE (QIAmp) or from the beads in 30 uL of DNase/RNase-free water (Zymo) and immediately used for RT-qPCR and library preparation. The presence of SARS-CoV-2 was confirmed using the SARS-CoV-2 Research Use Only qPCR Primer and Probe Kit (Integrated DNA Technologies, IDT; Coralville, IA) and qScript XLT One-Step RT-qPCR ToughMix (Quantabio; Beverly, MA) according to the CDC 2019-nCoV Real-Time RT-PCR Diagnostic Panel procedure.

The sequencing library was prepared using the 1200 bp amplicon "midnight" primer set (versions 1 or 2) [72] following the protocol by Freed and Silander [38]. Briefly, the RNA was reverse transcribed using the LunaScript RT Supermix Kit (New England BioLabs, NEB; Ipswich, MA). Tiled 1200 bp amplicons were generated using midnight primers (IDT) with Q5 Hot Start High-Fidelity 2X Master Mix (NEB) for 32 cycles (LA-GSU1 to LA-GSU19) or up to 40 cycles (LA-GSU20 to LA-GSU148 and LA-TECH1 to LA-TECH68) of multiplex PCR amplification. The two overlapping amplicon pools for each specimen were combined and quantified using the Qubit dsDNA HS Assay Kit (Invitrogen; Carlsbad, CA). A total of 75 – 150 ng of PCR product per specimen was barcoded using the Nanopore Rapid Barcoding Kits SQK-

RBK004 or SQK-RBK110.96 (Oxford Nanopore Technologies, ONT; Oxford, UK). The barcoded

samples were pooled and purified using AMPure XP beads at a 1:1 ratio of sample:beads

(Beckman Coulter; Brea, CA). The final libraries (350 – 800 ng DNA) were sequenced on an

R9.4.1 flow cell using a MinION Mk1B or Mk1C device, basecalled in real-time via MinKNOW,

and analyzed using the wf-artic workflow (medaka, minimap2, bcftools, samtools, nextclade,

artic, pangolin) according to the ONT protocol.

434 Phylogenetic analysis

SARS-CoV-2 genomes were downloaded from GISAID (www.gisaid.org) excluding the

low coverage and incomplete records up to April 30th 2022, and subsampling carried out using

subsampler to obtain a representative dataset [73]. Subsampling 4,199,772 (Delta sublineages)

and 2,895,902 (BA.1.x sublineages) genomic sequences resulted in 5,882 (Delta sublineages)

and 6,843 (BA.1.x sublineages) total sequences. Viral sequences were aligned using ViralMSA

with default parameters [74] using Wuhan-1 (MN908947.3) reference and then manually

curated with Aliview to designate the start site and trim terminal regions [75]. A maximum

likelihood (ML) phylogeny was inferred using IQTREE v1.6.10 with the best-fit nucleotide

substitution model identified by the Model Finder function [76]. Statistical support for nodes was

determined using 1,000 bootstrap replicates. TreeTime [77] was used to transform this ML tree

topologies into a dated tree using a constant mean rate of 8.0×10^{-4} nucleotide substitutions per

site per year, after the exclusion of outlier sequences. The phylogeny was visualized using the

ggtree package in RStudio with tips colored by collection source (continent).

449 Statistical analysis

450   Statistical analysis was performed using Prism version 7.05 (GraphPad; San Diego, CA).

451   Continuous variables between two groups were tested for normal distribution (D'Agostino &

452 Pearson test) and homogeneity of variance (F test) then compared using an independent t-test

453 with two-tails and alpha set to 0.05.

Data availability

456   The 1,053 complete SARS-CoV-2 genomes that have been sequenced to date by this

457 initiative using Illumina or Nanopore sequencing are available on GISAID and accessible at

458 doi.org/10.55876/gis8.221011tg. The phylogenetic analyses are based on sequences available

459 on GISAID and accessible at doi.org/10.55876/gis8.220901cu.

Acknowledgements

The authors deeply thank the healthcare professionals at Foster Johnson Health Center

(Dawn Holmes), The Health Hut (Jackie White, Chelsea Streets), Michael Brooks Family Clinic

(Natalie Wise, Taylor Schutzman, Whitney Walker), Minden Family Care Center (Beau Burton),

Serenity Springs Specialty Hospital (Tonya Monk), Mercer Medicine (Certified Medical

Assistants), and Jackson State University's Health Services Center (Kimberly Bullock) for

recruiting patients, gaining informed consent, and/or collecting the clinical specimens. We thank

Maarten Van Diest for assistance with data curation. We are grateful to the Louisiana

Department of Health for their efforts to facilitate enhanced viral genomic surveillance as a

public health activity. We gratefully acknowledge all data contributors, i.e., the Authors and their

Originating laboratories responsible for obtaining the specimens, and their Submitting

laboratories for generating the genetic sequence and metadata and sharing via the GISAID

Initiative, on which this research is based. The full acknowledgement table can be found in

Supplemental Table 1 and 2. We gratefully acknowledge the contribution of the GISAID Data

Curation Team to enable rapid sharing of genomic data. We acknowledge the Louisiana

Biomedical Research Network and its Bioinformatics, Biostatistics, and Computational Biology

Core for providing computational resources.

References

1.   Zhu N, Zhang D, Wang W, Li X, Yang B, Song J, et al. A Novel Coronavirus from

Patients with Pneumonia in China, 2019. N Engl J Med. 2020;382: 727–733.

2.   Brown CM, Vostok J, Johnson H, Burns M, Gharpure R, Sami S, et al. Outbreak of

SARS-CoV-2 Infections, Including COVID-19 Vaccine Breakthrough Infections, Associated with

Large Public Gatherings - Barnstable County, Massachusetts, July 2021. MMWR Morb Mortal

Wkly Rep. 2021;70: 1059–1062.

3.   Sheikh A, McMenamin J, Taylor B, Robertson C, Public Health Scotland and the EAVE II

Collaborators. SARS-CoV-2 Delta VOC in Scotland: demographics, risk of hospital admission,

and vaccine effectiveness. Lancet. 2021;397: 2461–2462.

4.   Twohig KA, Nyberg T, Zaidi A, Thelwall S, Sinnathamby MA, Aliabadi S, et al. Hospital

admission and emergency care attendance risk for SARS-CoV-2 delta (B.1.617.2) compared

with alpha (B.1.1.7) variants of concern: a cohort study. Lancet Infect Dis. 2022;22: 35–42.

5.   Planas D, Veyer D, Baidaliuk A, Staropoli I, Guivel-Benhassine F, Rajah MM, et al.

Reduced sensitivity of SARS-CoV-2 variant Delta to antibody neutralization. Nature. 2021;596:

276–280.

6.   Planas D, Saunders N, Maes P, Guivel-Benhassine F, Planchais C, Buchrieser J, et al.

Considerable escape of SARS-CoV-2 Omicron to antibody neutralization. Nature. 2022;602:

671–675.

7.   Viana R, Moyo S, Amoako DG, Tegally H, Scheepers C, Althaus CL, et al. Rapid

epidemic expansion of the SARS-CoV-2 Omicron variant in southern Africa. Nature. 2022;603:

679–686.

8.   Tegally H, Moir M, Everatt J, Giovanetti M, Scheepers C, Wilkinson E, et al. Emergence

of SARS-CoV-2 Omicron lineages BA.4 and BA.5 in South Africa. Nat Med. 2022;28: 1785–

1790.

9.   Cao Y, Yisimayi A, Jian F, Song W, Xiao T, Wang L, et al. BA.2.12.1, BA.4 and BA.5

escape antibodies elicited by Omicron infection. Nature. 2022;608: 593–602.

10.   Wang Q, Guo Y, Iketani S, Nair MS, Li Z, Mohri H, et al. Antibody evasion by SARS-

CoV-2 Omicron subvariants BA.2.12.1, BA.4 and BA.5. Nature. 2022;608: 603–608.

11.   Elbe S, Buckland-Merrett G. Data, disease and diplomacy: GISAID's innovative

contribution to global health. Glob Chall. 2017;1: 33–46.

12.   Centers for Disease Control and Prevention. COVID Data Tracker. In: COVID Data

Tracker [Internet]. Atlanta, GA; [cited 1 Oct 2022]. Available: https://covid.cdc.gov/covid-data-

tracker

13.   Cuadros DF, Branscum AJ, Mukandavire Z, Miller FD, MacKinnon N. Dynamics of the

COVID-19 epidemic in urban and rural areas in the United States. Ann Epidemiol. 2021;59: 16–

20.

14.   Matthews KA, Ullrich F, Gaglioti AH, Dugan S, Chen MS, Hall DM. Nonmetropolitan

COVID-19 Incidence and Mortality Rates Surpassed Metropolitan Rates Within the First 24

Weeks of the Pandemic Declaration: United States, March 1-October 18, 2020. J Rural Health.

2021;37: 272–277.

15.   Mueller JT, McConnell K, Burow PB, Pofahl K, Merdjanoff AA, Farrell J. Impacts of the

COVID-19 pandemic on rural America. Proc Natl Acad Sci U S A. 2021;118: 2019378118.

16.   Cheng KJG, Sun Y, Monnat SM. COVID-19 Death Rates Are Higher in Rural Counties

With Larger Shares of Blacks and Hispanics. J Rural Health. 2020;36: 602–608.

17.   Henning-Smith C, Tuttle M, Kozhimannil KB. Unequal Distribution of COVID-19 Risk

Among Rural Residents by Race and Ethnicity. J Rural Health. 2021;37: 224–226.

18. National Academies of Sciences, Engineering, and Medicine, Health and Medicine

Division, Board on Population Health and Public Health Practice, Committee on Community-

Based Solutions to Promote Health Equity, Baciu A, Negussie Y, et al. The State of Health

Disparities in the United States. National Academies Press (US); 2017.

19. Houtman J, Glassman R, Shultz L, Rivera JM, Bass E, Bright R. Genomic Surveillance

Is Essential To Track Covid-19 Variants in Both Unvaccinated and Vaccinated Populations. In:

Rockefeller Foundation [Internet]. 1 Dec 2021 [cited 8 Jun 2022]. Available:

https://www.rockefellerfoundation.org/blog/genomic-surveillance-is-essential-to-track-covid-19-

variants-in-both-unvaccinated-and-vaccinated-populations/

20. Gozashti L, Corbett-Detig R. Shortcomings of SARS-CoV-2 genomic metadata. BMC

Res Notes. 2021;14: 189.

21. Schriml LM, Chuvochina M, Davies N, Eloe-Fadrosh EA, Finn RD, Hugenholtz P, et al.

COVID-19 pandemic reveals the peril of ignoring metadata standards. Scientific Data. 2020;7:

1–4.

22. Ratcliffe M, Burd C, Holder K, Fields A. Defining Rural at the U.S. Census Bureau. U.S.

Census Bureau; 2016. Report No.: ACSGEO-1.

23. Iglehart JK. The Challenging Quest to Improve Rural Health Care. N Engl J Med.

2018;378: 473–479.

24. Johnston KJ, Wen H, Joynt Maddox KE. Lack Of Access To Specialists Associated With

Mortality And Preventable Hospitalizations Of Rural Medicare Beneficiaries. Health Aff .

2019;38: 1993–2002.

25. Peters DJ. Community Susceptibility and Resiliency to COVID-19 Across the Rural-

Urban Continuum in the United States. J Rural Health. 2020;36: 446–456.

26. A Poor People's Pandemic Report: Mapping the Intersections of Poverty, Race, and

COVID-19. In: Poor Peoples Campaign [Internet]. Apr 2022 [cited 8 Jun 2022]. Available:

https://www.poorpeoplescampaign.org/pandemic-report/

27. Webb Hooper M, Nápoles AM, Pérez-Stable EJ. COVID-19 and Racial/Ethnic

Disparities. JAMA. 2020;323: 2466–2467.

28. Gross CP, Essien UR, Pasha S, Gross JR, Wang S-Y, Nunez-Smith M. Racial and

Ethnic Disparities in Population-Level Covid-19 Mortality. Journal of General Internal Medicine.

2020. pp. 3097–3099. doi:10.1007/s11606-020-06081-w

29. Mackey K, Ayers CK, Kondo KK, Saha S, Advani SM, Young S, et al. Racial and Ethnic

Disparities in COVID-19-Related Infections, Hospitalizations, and Deaths : A Systematic

Review. Ann Intern Med. 2021;174: 362–373.

30. World Health Organization. Genomic sequencing of SARS-CoV-2: a guide to

implementation for maximum impact on public health. World Health Organization; 2021 Jan.

31. Cao Y, Jian F, Wang J, Yu Y, Song W, Yisimayi A, et al. Imprinted SARS-CoV-2

humoral immunity induces convergent Omicron RBD evolution. bioRxiv. 2022. p.

2022.09.15.507787. doi:10.1101/2022.09.15.507787

32. Sheward DJ, Kim C, Fischbach J, Muschiol S, Ehling RA, Björkström NK, et al. Omicron

sublineage BA.2.75.2 exhibits extensive escape from neutralising antibodies. bioRxiv. 2022. p.

2022.09.16.508299. doi:10.1101/2022.09.16.508299

33. Bergeron N. Omicron Variant Identified in Lincoln Parish. Ruston Daily Leader. 23 Dec

2021: 1.

34. Houldcroft CJ, Beale MA, Breuer J. Clinical and biological insights from viral genome

sequencing. Nat Rev Microbiol. 2017;15: 183–192.

35. Bull RA, Adikari TN, Ferguson JM, Hammond JM, Stevanovski I, Beukers AG, et al.

Analytical validity of nanopore sequencing for rapid SARS-CoV-2 genome analysis. Nat

Commun. 2020;11: 6272.

36. Delahaye C, Nicolas J. Sequencing DNA with nanopores: Troubles and biases. PLoS

One. 2021;16: e0257521.

37.   Baker DJ, Aydin A, Le-Viet T, Kay GL, Rudder S, de Oliveira Martins L, et al. CoronaHiT:

high-throughput sequencing of SARS-CoV-2 genomes. Genome Med. 2021;13: 21.

38.   Yan Y, Wu K, Chen J, Liu H, Huang Y, Zhang Y, et al. Rapid Acquisition of High-Quality

SARS-CoV-2 Genome via Amplicon-Oxford Nanopore Sequencing. Virol Sin. 2021;36: 901–

912.

39.   Itokawa K, Sekizuka T, Hashino M, Tanaka R, Kuroda M. Disentangling primer

interactions improves SARS-CoV-2 genome sequencing by multiplex tiling PCR. PLoS One.

2020;15: e0239403.

40.   Freed N, Silander O. SARS-CoV2 genome sequencing protocol (1200bp amplicon

"midnight" primer set, using Nanopore Rapid kit) v2. protocols.io. ZappyLab, Inc.; 2021.

doi:10.17504/protocols.io.bwyppfvn

41.   Brejová B, Boršová K, Hodorová V, Čabanová V, Gafurov A, Fričová D, et al. Nanopore

sequencing of SARS-CoV-2: Comparison of short and long PCR-tiling amplicon protocols. PLoS

One. 2021;16: e0259277.

42.   Liu H, Li J, Lin Y, Bo X, Song H, Li K, et al. Assessment of two-pool multiplex long-

amplicon nanopore sequencing of SARS-CoV-2. J Med Virol. 2022;94: 327–334.

43.   Charre C, Ginevra C, Sabatier M, Regue H, Destras G, Brun S, et al. Evaluation of NGS-

based approaches for SARS-CoV-2 whole genome characterisation. Virus Evol. 2020;6:

veaa075.

44.   Lagerborg KA, Normandin E, Bauer MR, Adams G, Figueroa K, Loreth C, et al. DNA

spike-ins enable confident interpretation of SARS-CoV-2 genomic data from amplicon-based

sequencing. bioRxiv. 2021. doi:10.1101/2021.03.16.435654

45.   González-Recio O, Gutiérrez-Rivas M, Peiró-Pastor R, Aguilera-Sepúlveda P, Cano-

Gómez C, Jiménez-Clavero MÁ, et al. Sequencing of SARS-CoV-2 genome using different

nanopore chemistries. Appl Microbiol Biotechnol. 2021;105: 3225–3234.

46.   Hay JA, Kissler SM, Fauver JR, Mack C, Tai CG, Samant RM, et al. Viral dynamics and

duration of PCR positivity of the SARS-CoV-2 Omicron variant. medRxiv. 2022.

doi:10.1101/2022.01.13.22269257

47.   Woodbridge Y, Amit S, Huppert A, Kopelman NM. Viral load dynamics of SARS-CoV-2

Delta and Omicron variants following multiple vaccine doses and previous infection. medRxiv.

2022. doi:10.1101/2022.03.20.22272549

48.   Puhach O, Adea K, Hulo N, Sattonnet P, Genecand C, Iten A, et al. Infectious viral load

in unvaccinated and vaccinated patients infected with SARS-CoV-2 WT, Delta and Omicron.

bioRxiv. 2022. doi:10.1101/2022.01.10.22269010

49.   Bramante CT, Proper JL, Boulware DR, Karger AB, Murray T, Rao V, et al. Vaccination

Against SARS-CoV-2 Is Associated With a Lower Viral Load and Likelihood of Systemic

Symptoms. Open Forum Infect Dis. 2022;9: ofac066.

50.   Elliott P, Haw D, Wang H, Eales O, Walters CE, Ainslie KEC, et al. Exponential growth,

high prevalence of SARS-CoV-2, and vaccine effectiveness associated with the Delta variant.

Science. 2021;374: eabl9551.

51.   Pritchard E, Matthews PC, Stoesser N, Eyre DW, Gethings O, Vihta K-D, et al. Impact of

vaccination on new SARS-CoV-2 infections in the United Kingdom. Nat Med. 2021;27: 1370–

1378.

52.   Acharya CB, Schrom J, Mitchell AM, Coil DA, Marquez C, Rojas S, et al. Viral Load

Among Vaccinated and Unvaccinated, Asymptomatic and Symptomatic Persons Infected With

the SARS-CoV-2 Delta Variant. Open Forum Infect Dis. 2022;9: ofac135.

53.   Chia PY, Ong SWX, Chiew CJ, Ang LW, Chavatte J-M, Mak T-M, et al. Virological and

serological kinetics of SARS-CoV-2 Delta variant vaccine breakthrough infections: a multicentre

cohort study. Clin Microbiol Infect. 2022;28: 612.e1–612.e7.

54.   Singanayagam A, Hakki S, Dunning J, Madon KJ, Crone MA, Koycheva A, et al.

Community transmission and viral load kinetics of the SARS-CoV-2 delta (B.1.617.2) variant in

vaccinated and unvaccinated individuals in the UK: a prospective, longitudinal, cohort study.

The Lancet Infectious Diseases. 2022. pp. 183–195. doi:10.1016/s1473-3099(21)00648-4

55. Lee S, Kim T, Lee E, Lee C, Kim H, Rhee H, et al. Clinical Course and Molecular Viral

Shedding Among Asymptomatic and Symptomatic Patients With SARS-CoV-2 Infection in a

Community Treatment Center in the Republic of Korea. JAMA Intern Med. 2020;180: 1447–

1452.

56. Ra SH, Lim JS, Kim G-U, Kim MJ, Jung J, Kim S-H. Upper respiratory viral load in

asymptomatic individuals and mildly symptomatic patients with SARS-CoV-2 infection. Thorax.

2021;76: 61–63.

57. Yang Q, Saldi TK, Gonzales PK, Lasda E, Decker CJ, Tat KL, et al. Just 2% of SARS-

CoV-2−positive individuals carry 90% of the virus circulating in communities. Proceedings of the

National Academy of Sciences. 2021. doi:10.1073/pnas.2104547118

58. Zuin M, Gentili V, Cervellati C, Rizzo R, Zuliani G. Viral Load Difference between

Symptomatic and Asymptomatic COVID-19 Patients: Systematic Review and Meta-Analysis.

Infect Dis Rep. 2021;13: 645–653.

59. Han MS, Seong M-W, Kim N, Shin S, Cho SI, Park H, et al. Viral RNA Load in Mildly

Symptomatic and Asymptomatic Children with COVID-19, Seoul, South Korea. Emerg Infect

Dis. 2020;26: 2497–2499.

60. Zhou R, Li F, Chen F, Liu H, Zheng J, Lei C, et al. Viral dynamics in asymptomatic

patients with COVID-19. Int J Infect Dis. 2020;96: 288–290.

61. Lennon NJ, Bhattacharyya RP, Mina MJ, Rehm HL, Hung DT, Smole S, et al. Cross-

sectional assessment of SARS-CoV-2 viral load by symptom status in Massachusetts

congregate living facilities. J Infect Dis. 2021. doi:10.1093/infdis/jiab367

62. Euser S, Aronson S, Manders I, van Lelyveld S, Herpers B, Sinnige J, et al. SARS-CoV-

2 viral-load distribution reveals that viral loads increase with age: a retrospective cross-sectional

cohort study. Int J Epidemiol. 2022;50: 1795–1803.

63.   Evans D, Cowen S, Kammel M, O'Sullivan DM, Stewart G, Grunert H-P, et al. The

Dangers of Using Cq to Quantify Nucleic Acid in Biological Samples: A Lesson From COVID-19.

Clin Chem. 2021;68: 153–162.

64.   Michalakis Y, Sofonea MT, Alizon S, Bravo IG. SARS-CoV-2 viral RNA levels are not

"viral load." Trends Microbiol. 2021;29: 970–972.

65.   Quick J, Loman NJ, Duraffour S, Simpson JT, Severi E, Cowley L, et al. Real-time,

portable genome sequencing for Ebola surveillance. Nature. 2016;530: 228–232.

66.   Quick J, Grubaugh ND, Pullan ST, Claro IM, Smith AD, Gangavarapu K, et al. Multiplex

PCR method for MinION and Illumina sequencing of Zika and other virus genomes directly from

clinical samples. Nat Protoc. 2017;12: 1261–1276.

67.   Faria NR, Kraemer MUG, Hill SC, Goes de Jesus J, Aguiar RS, Iani FCM, et al.

Genomic and epidemiological monitoring of yellow fever virus transmission potential. Science.

2018;361: 894–899.

68.   Kafetzopoulou LE, Pullan ST, Lemey P, Suchard MA, Ehichioya DU, Pahlmann M, et al.

Metagenomic sequencing at the epicenter of the Nigeria 2018 Lassa fever outbreak. Science.

2019;363: 74–77.

69.   Tyson JR, James P, Stoddart D, Sparks N, Wickenhagen A, Hall G, et al. Improvements

to the ARTIC multiplex PCR method for SARS-CoV-2 genome sequencing using nanopore.

bioRxiv. 2020. doi:10.1101/2020.09.04.283077

70.   Tegally H, San JE, Cotten M, Moir M, Tegomoh B, Mboowa G, et al. The evolving

SARS-CoV-2 epidemic in Africa: Insights from rapidly expanding genomic surveillance. Science.

2022; eabq5358.

71.   Douthit N, Kiv S, Dwolatzky T, Biswas S. Exposing some important barriers to health

care access in the rural USA. Public Health. 2015;129: 611–620.

72.   Lundberg DJ, Cho A, Raquib R, Nsoesie EO, Wrigley-Field E, Stokes AC. Geographic

and Temporal Patterns in Covid-19 Mortality by Race and Ethnicity in the United States from

March 2020 to February 2022. medRxiv. 2022. doi:10.1101/2022.07.20.22277872

73.   Orphan CM, McClure KR. An Anchor for the Region: Examining a Regional

Comprehensive University's Eff orts to Serve its Rural, Appalachian Community. J Res Rural

Educ. 2019;35: 1–19.

74.   Freed NE, Vlková M, Faisal MB, Silander OK. Rapid and inexpensive whole-genome

sequencing of SARS-CoV-2 using 1200 bp tiled amplicons and Oxford Nanopore Rapid

Barcoding. Biology Methods & Protocols. 2020;5. Available:

https://academic.oup.com/biomethods/article/5/1/bpaa014/5873518

75.   Alpert T, Brito AF, Lasek-Nesselquist E, Rothman J, Valesano AL, MacKay MJ, et al.

Early introductions and transmission of SARS-CoV-2 variant B.1.1.7 in the United States. Cell.

2021. pp. 2595–2604.e13. doi:10.1016/j.cell.2021.03.061

76.   Moshiri N. ViralMSA: massively scalable reference-guided multiple sequence alignment

of viral genomes. Bioinformatics. 2021;37: 714–716.

77.   Larsson A. AliView: a fast and lightweight alignment viewer and editor for large datasets.

Bioinformatics. 2014;30: 3276–3278.

78.   Nguyen L-T, Schmidt HA, von Haeseler A, Minh BQ. IQ-TREE: a fast and effective

stochastic algorithm for estimating maximum-likelihood phylogenies. Mol Biol Evol. 2015;32:

268–274.

79.   Sagulenko P, Puller V, Neher RA. TreeTime: Maximum-likelihood phylodynamic

analysis. Virus Evol. 2018;4: vex042.

Fig1

Fig2

Fig3

Fig4

Delta sublineages

0.1

BA.1.x sublineages

0.06

Fig5