1   **Metabolic evolution in response to interspecific competition in a eukaryote**

2   **Authors:** Giulia Ghedini[1,2]*, Dustin J. Marshall[1]

3   **Affiliations:**

4
5   [1] Centre for Geometric Biology, School of Biological Sciences, Monash University, 3800 Clayton,
6   VIC, Australia.

7   [2] Present address: Instituto Gulbenkian de Ciência, Rua Quinta Grande 6, 2780-156 Oeiras,
8   Portugal.

9   * Corresponding author. Email: gghedini@igc.gulbenkian.pt
10

Abstract

Competition can drive rapid evolution which, in turn, alters the trajectory of ecological communities.

The role of eco-evolutionary dynamics in ecological communities is increasingly well-appreciated, but

a mechanistic framework for identifying the types of traits that will evolve, and their trajectories, is

required. Metabolic theory makes explicit predictions about how competition should shape the

evolution of metabolism and size but these predictions have gone largely untested, particularly in

eukaryotes. We use experimental evolution of a eukaryotic phototroph to examine how metabolism,

size, and demography coevolve under both inter- and intra-specific competition. We find that the focal

species evolves a smaller body size in response to competition, reducing density-dependence and

maximizing carrying capacity. Metabolic theory successfully predicted most of these adaptations, but

we also find important departures from theory. Longer-term evolution (70 generations) led to Pareto

improvements in both population growth rate and carrying capacity, suggesting that classic r-K trade-

offs observed among species can be evaded within species. The evasion of this trade-off appeared to

arise due to the rapid evolution of enhanced metabolic plasticity: lineages exposed to competition

evolved more labile metabolisms that tracked resource availability more effectively than lineages that

were competition-free. We predict that rapid evolution in both size and metabolism may be a

ubiquitous feature of adaptation to changing resource regimes that occur *via* species invasions and

environmental change.

Keywords: energy flux, community, species interactions, fitness, homeostasis, resources

Introduction

When ecological and evolutionary processes occur on similar timescales, eco-evolutionary dynamics can shape the diversity and functioning of communities [1–3]. Studies in bacteria imply that competition from whole communities alters evolutionary trajectories in ways that are fundamentally different from simpler pairwise interactions [4–7]. Whether these results extend to eukaryotic communities, where species tend to have lower population densities and longer life cycles, and thus slower rates of evolution, remains unclear [8]. Macroevolutionary patterns in eukaryotes suggest that competition for resources can drive species to evolve differential resource use (e.g. Darwin's finches) [9–11] but such escapes from competition may not always be accessible [12,13]. In most cases, predicting trait evolution in species that share similar niches remains a formidable challenge [8,14,15].

Metabolic theory should help with identifying the eco-evolutionary trajectories of key phenotypic traits. The theory proposes that metabolism should drive competition by setting *per capita* resource demands [16,17]. For example, because smaller organisms have lower absolute metabolic rates, their populations should support higher maximum population densities than those of larger organisms [18]. Similarly, because small organisms usually have higher metabolism per unit mass (assuming that metabolism scales hypo-allometrically with body size), their populations should grow faster than those of larger organisms – producing classic r-K trade-offs often observed among species [17]. Therefore, evolved changes in size and metabolism should both affect demography and capacity to compete with other species [19,20] but whether size and metabolism actually coevolve in response to interspecific competition has not been explored.

We combined experimental evolution and metabolic theory to determine how size, metabolic fluxes, and demography coevolve in response to both intra- and inter-specific competition in phytoplankton. We base our assessment on a focal species, the marine eukaryotic microalga *Dunaliella tertiolecta*, which we evolved for ~70 generations in each of three environments: i) competition-free; ii) intraspecific competition; and iii) interspecific competition (three other phytoplankton species). To manipulate the environment, we used dialysis bags that physically isolated the evolving population from its competitors while exposing it to competition for light and nutrients [6,21–23] (Fig. 1). We treated the competition-free lineages as a reference control. We batch-transferred both the focal species and the competitors weekly standardizing biovolumes between treatments (~7 generations). We assessed how the focal populations had evolved at two points: at ~35 generations and ~70 generations. At each

of these time points, we placed subsamples of all the evolved lineages of the focal species into a

common environment, and left them for several generations such that any persistent phenotypic

differences that remained among treatments represented evolved responses to the environments [24].

We assessed metabolism (photosynthesis, respiration, and net daily energy production), morphology

(cell size and shape) and demography (growth rate, max. population density, and max. total biomass).

Figure 1. To test how metabolism, size and demography coevolve in response to competition, we

evolved the eukaryotic microalga *Dunaliella tertiolecta* in three environments: competition-free

("none"), intraspecific ("intra") and interspecific competition ("inter"). We enclosed the focal species in

dialysis bags that physically isolated the evolving population from competitors while exposing it to

competition for light and nutrients for 10 weeks corresponding to ~70 generations. We propagated the

focal lineages and the competitors performing weekly batch transfers. After ~35 and ~70 generations

we used common garden experiments to compare changes in metabolic fluxes, morphology, and

demography among the evolved focal populations.

Results

Metabolism evolves under competition

After ~35 generations of experiencing interspecific competition, algae evolved lower metabolic fluxes than algae that experienced no interspecific competition. Specifically, rates of both photosynthesis and dark respiration were lower in lineages exposed to interspecific competition (photosynthesis: competition × biovolume interaction, $F_{2,270} = 3.85$, p = 0.02; respiration: main competition effect, $F_{2,272} = 6.56$, p = 0.002; see Supplementary Information*:* Fig. S1, Table S1). These differences in metabolism were only apparent when we measured energy fluxes under resource limitation; the lineages were the same when resources were abundant (Fig. S1, Table S1). In other words, algae that experienced interspecific competition evolved plasticity to have 'thriftier' metabolisms when resources were scarce, reducing their rates of energy use, and presumably nutrient demands, while photosynthesising relative to other lineages.

After a further 35 generations of experimental evolution, interspecific competition drove more pronounced evolutionary changes in metabolism and metabolic plasticity. When we assessed metabolic fluxes under resource-limiting conditions, metabolic rates (both photosynthesis and respiration) were lowest in lineages that had experienced interspecific competition and highest in those that experienced no competition (competition effect on photosynthesis: $F_{2,47} = 6.75$, P = 0.003; Fig. 2b; respiration: $F_{2,47} = 15.83$, p < 0.0001, Fig. 2d; Table S1). Again, lineages that evolved in the presence of competition were thriftier when resources were scarce, particularly those that evolved under interspecific competition. But when we assessed metabolic fluxes when resources were abundant, the patterns were reversed: lineages exposed to competition had the highest rates of photosynthesis and respiration relative to competition-free lineages. In other words, lineages that experienced competition had evolved greater capacity to both fuel (via photosynthesis) and do (via respiration) biological work when resources were abundant (competition × biovolume effect on photosynthesis: $F_{2,210} = 3.59$, p = 0.03; Fig. 2a; competition effect on respiration: $F_{2,46} = 10.86$, p = 0.0001, Fig. 2c; Table S1). Overall then, competition-exposed lineages evolved higher levels of metabolic plasticity than competition-free lineages: competition-exposed lineages downregulated their resource demands when resources were scarce and strongly upregulated their demands during periods of resource abundance (Fig. 3). In contrast, competition-free lineages exhibited more modest metabolic plasticity in response to changes in resource availability. Together, these evolved metabolic

changes meant that competition-exposed lineages produced more energy on balance (photosynthesis

– respiration), thereby fuelling more biological work when resources were abundant (Fig. 2e), but

required fewer resources and produced less net energy when resource were scarce, relative to

competition-free lineages (Fig. 2f).

Figure 2. Populations exposed to interspecific competition for ~70 generations evolved greater

metabolic plasticity (as revealed by the relationship between population biovolume and metabolic

rate) than competition-free populations. Competition exposed populations had greater (a)

photosynthesis and (c) respiration rates when resources were abundant during the exponential

growth phase (left panel). But upon entering stationary phase and approaching carrying capacity

(right panel), populations that experienced competition from interspecifics had much lower metabolic

rates (b for photosynthesis, d for respiration). Integrating photosynthesis and respiration over 24

hours, we found that populations that had evolved in the presence of competitors produced more net

energy than competition-free populations when resources were abundant (e) but they also produced

less net energy when resources were scarce because they downregulated their metabolism relatively
more (f).

Figure 3. Populations exposed to interspecific competition for ~70 generations ("inter") evolved
greater metabolic plasticity relative to lineages that experienced no competition ("none") allowing
them to better track resource abundance. Cells with a history of competition increase their metabolic
rates (both photosynthesis and respiration) much more than control lineages when resources are
abundant (day 3, exponential growth phase) but downregulate them when resources are scarce (day
11, stationary phase).

Size evolution, metabolic scaling and predicting demography

Lineages exposed to competition evolved smaller cell sizes than lineages that were free from
competition (Fig. 4a for day 3 after 10 weeks; estimated marginal means for day 3: competition-free =
202; intra = 185; inter = 162 µm^3). After 70 generations of experimental evolution, lineages
experiencing interspecific competition were on average 13.4% smaller in cell volume than
competition-free lineages (ranging from 3.6% to 19.6% smaller during common garden experiments),
while those experiencing only intraspecific competition were intermediate in size (5.5% smaller on
average; ranging from 0.6% larger to 8.5% smaller in size than competition-free lineages)
(competition × time: $F_{24,564}$ = 3.65, p < 0.0001; Fig. S2 for all days, Table S2). Cell shape coevolved
with cell size: lineages exposed to competitors evolved rounder cells than competition-free lineages
(Fig. 4b for day 3; Fig. S2; competition × time: $F_{24,564}$ = 8.50, p < 0.0001, Table S3 for post-hoc
comparisons).

While metabolic rates often scale hypo-allometrically for eukaryotes, we found that both
photosynthesis and respiration rates scaled hyper-allometrically with cell size in the focal species

(photosynthesis: 1.31 [CI: 1.05; 1.58]; respiration: 1.28 [1.02; 1.55]; Table S4), similarly to what

previously found for this species [20]. But the scaling became progressively shallower as populations

grew denser over time (Table S4). Thus, under hyper-allometry, the smaller cells evolved under

interspecific competition should have lower resource demands *per capita,* in an absolute sense but

also relative to cell biovolume. Given the observed metabolic scalings during the exponential growth

phase ($B_{exponential}$ = 1.13) and stationary phase ($B_{stationary}$ = 0.60), metabolic theory would therefore

make the following predictions regarding the scaling of demographic parameters with cell size [19,20,25]:

(1) $r_{max} = M^{B_{exponential}}/M^1 = M^{0.13}$

(2) $K_{cells} = M^0/M^{B_{stationary}} = M^{-0.60}$

(3) $K_{bio} = M \times K_{cells} = M \times M^{-B} = M^{0.40}$

From these equations, we qualitatively predict that competition exposed lineages (which evolved

smaller sizes) should have lower intrinsic rates of increase (r_{max}) but higher carrying capacities (K_{cells})

than competition-free lineages. In a quantitative sense, we also predict that growth rate and biomass

carrying capacity (K_{bio}) should positive covary with cell size, while population carrying capacity should

negatively covary with cell size (K_{cells}). We tested both sets of predictions experimentally.

162
Figure 4. Both the size and shape of the microalga *Dunaliella tertiolecta* evolved in response to

competition. After ~70 generations, populations exposed to competitors evolved smaller cell sizes,

with a stronger decline in response to interspecific rather than intraspecific competitors (a, here

shown for day 3). Changes in cell size were accompanied by changes in shape: the smaller cells

exposed to intra- and inter-specifics competitors were also rounder than competition-free cells. See

Fig. S2 for the complete temporal series.

The evolution of greater carrying capacity

We found strong qualitative and quantitative support for our demographic predictions based on

metabolic theory. After ~35 generations, lineages that experienced competition had lower maximum

growth rates but greater carrying capacity (K_{cells}) than competition free lineages, displaying a classic r-

K trade-off (effect of competition on r_{max}: $F_{2,42} = 10.46$, $p < 0.0005$, Fig. 5a; effect on K_{cells}: $F_{2,42} =$

19.18, $p < 0.001$, Fig. 5b). After 70 generations of experimental evolution, we found the same evolved

difference in carrying capacity (K_{cells}; $F_{2,46} = 13.91$, $p < 0.001$, Fig. 5d) but the competition-exposed

lineages showed maximum population growth rates that were similar to the competition-free lineages

($F_{2,46} = 1.97$, $p = 0.15$, Fig. 5c; Table S5). The evolved demographic responses were similar

regardless of whether competition was intra- or inter-specific (Fig. S3, Table S5).

Figure 5. Populations that experienced competition had lower maximum population growth rates but

greater carrying capacity in terms of cell numbers after 5 weeks of experimental evolution (~35

generations, top row). Differences in carrying capacity persisted after a further 35 generations but with

no differences in growth rates between competition environments (bottom row). At either time, the

type of competition experienced (intra- or inter-specific) did not affect demographic parameters.

We compared our quantitative predictions derived from metabolic theory with the observed scaling of

each demographic parameter with cell size for the focal populations, and found remarkably strong

congruence between the two for most parameters. We find that K_{cells} declines (c.f.: observed = –0.56;

predicted = –0.60; $F_{1,48}$ = 22.4, p < 0.001; CI: –0.79; –0.32; Fig. 6b) and K_{bio} increases with size as

predicted by theory (c.f.: observed = 0.44; predicted = 0.40; $F_{1,48}$ = 14.1, p < 0.001; Fig. 6c). But we

find a slightly weaker than predicted relationship between r_{max} and size (c.f.: observed= 0.06;

predicted = 0.13); this relationship is non-significant as the confidence intervals overlap zero ($F_{1,48}$ =

0.02, p = 0.88; CI: –0.71; 0.83; Fig. 6a).

Figure 6. We find strong congruence between predictions from metabolic theory (broken lines) and

demographic observations (solid lines). The scalings of max. growth rate (r_{max}), max. population

density (K_{cells}) and max. biovolume (K_{bio}) with cell size observed from our data after 70 generations

are: r_{max} = observed 0.06 (CI: –0.71; 0.83); predicted 0.13; K_{cells} = observed –0.56 (CI: –0.79; –0.32);

predicted –0.60; K_{bio} = observed 0.44 (CI: 0.21; 0.68); predicted 0.40. Scaling exponents are

calculated for the 3 competition treatments together (blue = inter; green = intra; red = competition-

free). We calculated the expected scaling of r_{max} as the average scaling observed over the first 5 days

because population growth is determined early on (exponential growth phase); we calculated the

expected scaling of K_{cells} and K_{bio} as the average scaling over the last 3 sampling days (12 to 16)

when cultures were in stationary phase and because cell size stabilised from day 12 onwards (Fig.

S2). We used the same range of days to calculate the average cell size for each replicate shown

above (hence differences in size on x-axis between panel a and panels b-c).

Discussion

We found that intra- and inter-specific competition drove the evolution of cell size and metabolism in the eukaryote *Dunaliella tertiolecta* in ways that metabolic theory anticipates. Because metabolism scaled hyper-allometrically with cell size in the focal lineages, the evolution of smaller sizes in response to competition reduced *per capita* resource demands (both in absolute and relative terms), such that cells achieved higher population densities but lower total biomass. These increases in population carrying capacity initially came at the expense of population growth rates – after only 35 generations of experimental evolution, the focal lineages experienced the classic trade-off between population growth rate and carrying capacity. However, through further evolution (~70 generations), competition-exposed lineages achieved Pareto improvements in demography: their max. population growth rates were comparable to those of competition-free lineages while also maintaining higher carrying capacities (Fig. 5). The evolution of greater metabolic flexibility appeared to enable this change; competition-exposed lineages were able to better downregulate their energy fluxes when resources were scare but strongly upregulated their metabolism when resources were abundant, thereby increasing both population growth rates and carrying capacity. This finding represents a novel departure from theory as the evolution of enhanced metabolic plasticity was unanticipated by any theory base but, in hindsight, seems almost inevitable. We eagerly await further studies that test the role of competition in driving metabolic plasticity in other systems. Enhancements in plasticity might explain why within-species patterns may not always reflect the same trade-offs observed among species [14,26].

Metabolism sets resource use and fuels growth, so metabolic rate should be under selection when competition is intense [27]. Classic macroevolutionary patterns suggest that competition selects for efficiency which is often associated with larger body sizes and lower metabolic rates because metabolism scales hypo-allometrically in most metazoans [28–30]. In contrast, our species exhibits hyper-allometric scaling of both photosynthesis (1.31) and respiration (1.28) with cell size so, while larger cells produce relatively more energy *per capita*, they are also both absolutely and relatively more expensive to maintain. Therefore, lineages exposed to competition evolved smaller, more resource-efficient cells. Interestingly, Long Term Experimental Evolution (LTEE) in *E. coli* yielded the opposite evolutionary trajectory of cell size to produce the same outcome – cells became larger over

time to enhance resource-use efficiency [25]. The trajectories of cell size in both studies actually accord

with metabolic theory. In both, cells change size such that size-specific metabolic rates decrease

(metabolism scales hypo-allometrically with size in the LTEE). Metabolic theory also predicts the

relationships between size, density, and biomass in each study (based on species-specific metabolic

scaling relationships). Based on our results and those of Marshall et al. [25], we would therefore predict

that competition should select for smaller body size in species which show hyper-allometric metabolic

scaling, but larger body size in species that show hypo-allometric scaling. For metazoans, most

species show hypo-allometric metabolic scaling within species (the appropriate biological scale in this

instance), whereas intraspecific metabolic scaling in unicellular organisms remains largely unexplored

except for a few cases. We look forward to future studies that explore how metabolic scaling and

competition interact in other species, particularly other microbial species given their pivotal role in

driving biogeochemical cycles [31,32].

Our findings have some concerning implications regarding global change. Phytoplankton play a key

role in the global carbon budget, accounting for ~50% of all carbon fixed annually; thus, any change in

phytoplankton productivity will strongly influence global carbon fluxes [33]. Increased water stratification

due to ocean warming will intensify resource competition and decrease cell size within species [34,35].

Our competition-exposed lineages evolved smaller cells in response to stronger resource competition

– in that sense, our experimental evolution mimics the impacts of global change. We found that the

net energy fluxes of the competition-exposed lineages were ~15% lower in the stationary phase than

those of competition-free lineages (Fig. 2f). Hence the net carbon assimilation rate of our evolved

lineages was systematically lower. If natural phytoplankton populations also evolve smaller cell sizes

in response to warming and/or nutrient limitation (and initial studies suggest that they do) [34], then our

results would imply that rates of carbon assimilation and phytoplankton biomass will decline with

further global change, creating a positive feedback loop. While thermal adaptation might dampen the

negative effects of warming on phytoplankton productivity [36,37], species interactions can increase the

sensitivity of species to global change through resource competition [7,38] and alter their evolutionary

rates in ways that are poorly resolved [4,6]. Our results suggest that competition for resources might

compound the impacts of global change by further selecting for smaller cell sizes and thriftier

metabolisms that better tolerate competition but that may have undesirable impacts on carbon

fixation.

Evolution happened rapidly in our system (within 70 generations). Such rapid evolution was likely

fuelled by the very large population size of the founding populations (~5 million cells). These rates of

evolution are comparable to those observed in other experimental evolution studies of phytoplankton

where size and metabolic traits evolved within 100 generations [37,39,40], whereas it takes longer for

evolution to manifest when populations are derived from a single clone [41]. Overall, competition from a

community exerted stronger effects on the evolution of these traits than intraspecific competition,

possibly because diverse competitors consume available resources more rapidly and effectively [42,43].

Thus, by posing a stronger selective pressure, competition from a community speeded up adaptation

countering the negative effects of reduced population sizes due to stronger resource competition [44],

similarly to what observed in bacteria [4,5]. Still, we find remarkable consistency between the

evolutionary responses to intra- and inter-specific competition, with both selecting for smaller, more

efficient cells able to sustain greater maximum population densities. Carrying capacity (or equilibrium

population size, K), rather than growth rate, is thus the quantity maximised by competition as

predicted by life history theory for stable or low stochastic environments [45–47]. Selection for r-K traits

has been demonstrated in numerous systems within species [27,46,48] but few studies have

demonstrated the same principle in response to interspecific competition. MacArthur & Wilson first

hypothesized in 1967 that selection for K should also apply in a community when species compete for

similar resources [45] but, to our best knowledge, these results are a first experimental demonstration

that this demographic prediction of life history theory also holds under interspecific competition.

Since size and metabolism together determine resource access and use [34,49,50], changes in both traits

might be a common feature of how species cope with shifts in competition and resource regimes.

Both size and metabolic evolution might become more common, or rapid, as climate change, species

introductions and biodiversity loss alter competition for resources [51,52]. It seems likely that metabolic

evolution will underpin many human-induced changes in resource regimes as well as mortality risks

[53], and we look forward to future studies that search for signatures of shifts in metabolism associated

with changes in competition regimes.

Materials and Methods

Overview of experimental set up

We tested the effects of competition on the metabolic fluxes, morphology, and demography of the eukaryotic unicellular alga *Dunaliella tertiolecta*, which we obtained from the CSIRO Australian National Algae Culture Collection (ANACC, strain CS-14). To manipulate the competitive environment, we enclosed the focal species in dialysis bags which were then placed in one of three environments for 10 weeks (1st June to 10th August 2021, ~70 generations): 1) surrounded by media with no competitors (competition-free), 2) by a population of conspecifics from the same strain (intraspecific competition) or 3) by a community of three other phytoplankton species (interspecific competition). For this latter treatment, the species were chosen to represent different sizes and phytoplankton groups: *Amphidinium carterae* (CS-740), *Phaeodactylum tricornutum* (CS-29) and *Nannochloropsis oculata* (CS-179) (Fig. 1, phase 1). Every week we batch-transferred both the focal species and the competitors (~7 generations). After 5 and 10 weeks (~35 and ~70 generations, respectively), we quantified changes in the traits of the focal species in common garden experiments (phase 2).

We sourced all phytoplankton species as 20 ml populations from CSIRO algal collection, Australia. We grew each population individually in 2 L glass bottles with standard f/2 enriched seawater medium enriched with silica for a month prior to the experiment, transferring each population into fresh media every week. The same medium was used throughout the experiment and common gardens. The medium was prepared with 0.45 μm filtered seawater and autoclaved following the recipe of Guillard & Ryther, 1962 [54]. All experiments were performed in a temperature controlled room (20 ± 1°C) on a 14:10 hours light:dark cycle and the cultures were grown under a light intensity of 115 ± 5 μmol photons m^{-2} s^{-1}.

Phase 1: evolution with competitors

The dialysis bags in which we enclosed the focal species enabled competition for light and nutrients but prevented cell mixing among phytoplankton species and exchange of bacteria (MWCO 12-14000 Da, pore size 24 Angstrom). Each bag contained a volume of 35 ml and was placed at the centre of a 500 ml glass jar assigned to one of the three competition treatments above (n = 20 for each competition treatment).

We started the experiment with an initial biovolume of 9.6×10^8 µm³ of the focal species which

corresponded 5×10^6 cells. This large inoculum size meant that the founding populations likely

contained a large amount of standing genetic variation. The glass jars containing the competitors

were filled to 350 ml to completely submerge the dialysis bag. To maintain the same biovolume to

media ratio, the initial biovolume of the competitors was 10 times that of the focal species (i.e. $9.6 \times$

10^9 µm³). We added the three competitor species of the interspecific treatment in equal biovolumes.

The jars of the control treatment (competition-free) were filled with media only.

We quantified biovolume as the product of cell density and cell size obtained for each species from

two 10 µl samples, fixed with 1% Lugol's solution. For each sample, we took 20 photos equally

spaced around the cell-counting grid of a Neubauer counting chamber (ProSciTech, Australia) under

an Olympus light microscope at 400× magnification. We analysed the images with ImageJ and Fiji

software (version 2.0) [55] to quantify the number of cells (cells µl^{-1}) and their size (µm, length and

width; diameter for *Nannochloropsis*). We calculated cell volume by assigning to each species an

approximate geometric shape [56] (prolate spheroid *V = (pi/6) × width² × length* for all species except

Nannochloropsis which we treated as a sphere *V = (pi/6) × diameter³*). Cell circularity is calculated by

the Fiji software as *4 × pi × (area/perimeter²)* and ranges from 0 to 1 (perfect circle).

Each week we transferred the same initial biovolume of the focal species and the same initial

biovolume of the competitors to a new, sterilized set of dialysis bag and jar. Cell densities, size and

biovolume were determined as described above. We did not manipulate the relative abundance of

species in the interspecific treatment. In weeks 3, 5 and 6 the cell densities of the focal species were

very low in the interspecific treatment – had we reinoculated the same initial biovolume we would not

have been able to add fresh medium. To avoid this nutrient limitation, we reinoculated only half of the

initial biovolumes across all treatments. By week 6, we discarded 13 lineages of the focal species in

the interspecific treatment because they were contaminated by one of the interspecific competitors

(the diatom *Phaeodactylum tricornutum*). To continue the experiment, we established an additional

lineage from each of the remaining seven replicates. By the end of the experiment, we lost 4 other

replicates in the interspecific treatment because of *Phaeodactylum*'s contamination.

The batch-transfer approach meant that all populations in our experiment experienced fluctuating

densities, but those surrounded by competitors always faced greater densities and thus competition.

Shifts in densities are common in nature and mediate the balance between density-independent and

density-dependent selection (r- and K-selection) [45]. If stochasticity is low, life theory predicts that

selection for K should prevail even under fluctuating conditions [28,47,57].

Phase 2: common garden experiments

We quantified evolutionary changes in the metabolism, morphology and demography of the focal

species in two common garden experiments; the first after five weeks of evolution with competitors

(~35 generations) and the second after ten weeks (~70 generations). Before each common garden,

we grew the focal species in a neutral environment for two generations to reduce any environmental

conditioning (neutral selection). Both for the neutral selection and common garden, we inoculated an

equal biovolume of each lineage in cell culture flasks (n = 20 for the no competition and intraspecific

treatments, n = 6 and n = 10 for the interspecific treatment in the first and second common garden,

respectively). Biovolume was determined as before from the average cell size and cell density of each

lineage; we decided a priori to add 15 ml of the lineage with the limiting biovolume (~14 × 10^8 μm^3),

back calculated the volumes of the other lineages and added 100 ml of media (f/2 + silica, as above)

to all flasks. The two common gardens started on 8th July and 12th August 2021, respectively, and

lasted 16 days, which was the approximate time needed for the cultures to reach carrying capacity –

one of the key traits we wanted to phenotype. We discontinued the common garden after 16 days to

avoid further evolution or adaptation by the focal lineages to the common garden conditions. Each

day, we removed 10 ml from each culture flask for sampling and replaced it with fresh media. In the

first common garden we sampled every day except on days 6, 10, 13 and 15; in the second we

sampled every day except on days 7, 13 and 15.

Trait measurements

Each sampling day, we fixed 1 ml sample with 1% Lugol's solution to quantify the cell size (volume),

shape (circularity) and abundance of each lineage as described above. We assessed changes in cell

shape in addition to size because the shape of a cell can mediate access to resources and is thus an

important component of fitness in unicellular organisms [34,58–61].

We measured oxygen evolution rates in 5 ml samples using 24-channel optical fluorescence oxygen

readers (PreSens Sensor Dish Reader, SDR; AS-1 Scientific Wellington, New Zealand) following

established protocols [62,63]. Sensors were calibrated with 0% and 100% air saturation before the

experiment. Net photosynthesis (oxygen production) was measured under the light intensity at which

the cultures were grown for 20 minutes, followed by 1 hour in the dark to measure respiration rates.

Thirteen blanks were filled with the media obtained from centrifuged samples (spun at 2,500 rpm for

10 min to separate the algae from the supernatant) to correct for background microbial activity since

cultures were not axenic. Prior to measurements, samples were spiked with 50 µl of sodium

bicarbonate stock for a final concentration of 2 mM sodium bicarbonate to avoid carbon limitation.

The change in percentage oxygen saturation was calculated with linear regressions using the LoLinR

package [64]. The rate of photosynthesis or respiration of the whole sample (VO_2; units µmol O_2/min)

was then measured as $VO_2 = 1 \times ((m_a - m_b)/100 \times V\beta O_2)$ following [65], where m_a is the rate of change

of O_2 saturation in each sample (min^{-1}), m_b is the mean O_2 saturation across all blanks (min^{-1}), V is

the sample volume (0.005 L) and $V\beta O_2$ is the oxygen capacity of air-saturated seawater at 20°C and

35 ppt salinity (225 µmol O_2/L). The first three minutes of measurements in the light were discarded

for all samples. Respiration rates were calculated after 15 minutes of dark when oxygen levels started

to show a linear decline. Photosynthesis and respiration (µmol O_2/min) were converted to calorific

energy (J/min) using the conversion factor of 0.512 J/µmol O_2 to estimate energy production and

energy consumption, respectively [66].

Statistical analyses

All analyses and plots were done in RStudio (version 4.1.3), separately for the two common gardens,

using the main packages nlme [67], lme4 [68], car [69], and ggplot2 [70].

Morphology. We assessed differences in cell morphology (size and shape) with linear mixed models

including competition treatment (3 levels) and time (day) as categorical predictors, and lineage

identity as random intercept to account for repeated measures; time was considered categorical

because the relationship with cell morphology was non-linear. We take day 3, during the exponential

growth phase, as a reference to report post hoc results on differences in cell size and shape in the

main text, and we report post hoc results for each day in supplements.

Metabolism. While oxygen rates increased linearly with biovolume over the first few days, this

relationship broke down as biovolume increased (over ~3 $µm^3$/µl) (Fig. S4). Log-transformation did not

result in linearity across the entire range of biovolume. Therefore, we analysed oxygen rates

separately for the first part of common gardens during the exponential growth phase (from day 1 to

day 6 included), and for the second part during the stationary phase (day 7 to 16). We used linear

mixed models with biovolume (covariate) and competition treatment as predictors, and lineage identity

as random intercept for the data of the second common garden to account for repeated measures.

Because variances were heterogenous for the photosynthesis data during the growth phase we used

a generalised linear mixed model including competition-specific variances. Finally, we integrated

photosynthesis and respiration rates over a 24-hour period to estimate net energy production of the

whole sample (J/day), as 14 hr of energy produced through net photosynthesis minus 10 hr of

respiration using rates predicted from the mixed models above.

We could not use mixed models on the oxygen evolution data from the first common garden because

of a singular fit error likely due to the lower replication of the interspecific competition treatment ($n =$

6). Therefore, we used linear models with biovolume (covariate) and competition treatment as

predictors (i.e. without including lineage identity). While this is not ideal, we repeated the analyses for

the second common garden with the same, simple linear models and obtained the same results of

mixed models (results not reported here), suggesting that oxygen rates are not strongly affected by

lineage identity. Nonetheless, we report the results on oxygen rates for the first common garden as a

supplementary figure. For all analyses of oxygen rates, data were not transformed but biovolume was

rescaled to 10^{-5} μm^3 μl^{-1}. Interactions between biovolume and competition were removed when p >

0.25 or if the interaction was not significant and the model with interaction did not perform better than

the simpler model compared by AIC and anova.

<u>Demography</u>. To test differences in the maximum rates of increase (r_{max}) and maximum values (K_{cells})

of cell density we followed a three-step approach (described in detail in Ghedini et al. [63], adapted from

Malerba et al. [71]). First, we fitted four growth models to each individual replicate lineage and chose the

best-fitting model among the four candidates to describe changes in the cell density (cells μl^{-1}) of

each culture over time. We used AIC to determine which growth model best described the dynamics

of a culture and successful convergence was ensured for all best-fitting models. The four models

were: a logistic-type sinusoidal growth model with lower asymptote forced to 0 (i.e. three-parameter

logistic curve), a logistic-type sinusoidal growth model with non-zero lower asymptote (i.e. four-

parameter logistic curve), a Gompertz-type sinusoidal growth model (i.e. three-parameter Gompertz

curve) and a modified Gompertz-type sinusoidal growth model including population decline after

reaching a maximum (i.e. four-parameter Gompertz-like curve including mortality). Second, we used

the best-fitting model to estimate growth parameters (i.e. r_{max} and K_{cells}) for each culture. From each

nonlinear curve, we extracted the maximum predicted value of population density (K_{cells}, cells μl^{-1}).

From the first derivative of the curve, we extracted the maximum rate of population increase (r_{max},

unit: day^{-1}). Third, we used an analysis of covariance to evaluate the influence of competition on each

parameter, using a linear model including the initial cell density estimated from the previous step as a

covariate and competition environment as a factor (three levels). The estimates of K_{cells} of the first

common garden had heterogeneous variances among treatments, so we used generalized least

squares models (instead of linear models) including treatment-specific variance for each level of

competition treatment (varIdent function in R). We then estimated and plotted least square means and

95% confidence intervals using Tukey p-value adjustment for comparing three estimates.

<u>Scaling of demographic parameters with cell size</u>. We determined the scaling of metabolism with cell

size for our evolved populations after ten weeks of evolution (~70 generation). For this assessment

we combined the data across the three competition treatments to assess a wider range of size. We

calculated the scaling of respiration and photosynthesis, separately, from linear models including cell

size and experiment day as numerical predictors. Oxygen rates and cell size were log_{10}-transformed.

Results did not qualitatively change when we analysed each competition treatment separately, as

each of them scaled > 1 and declined over time (Table S4).

Based on these results (effects of size and experiment day), we then calculated the expected scaling

of respiration with size for any given day. According to metabolic theory, the cost of production is

directly proportional to cell size (scales with size at 1) [17,25], so demographic parameters should scale

with size at:

(1) $r_{max} = M^B_{exponential}/M^1 = M^{B-1}$

(2) $K_{cells} = M^0/M^B_{stationary} = M^{-B}$

(3) $K_{bio} = M \times K_{cells} = M \times M^{-B} = M^{1-B}$

We tested these predictions by comparing the predicted metabolic scaling values for the growth

phase and stationary phase to the observed metabolic scalings from our data. We used early (days 1-

5) respiratory scaling values for expectations about r_{max} because it is mostly determined by growth

rates early on ($B_{exponential} = 1.13$). We used late (days 12-16) respiratory scaling values for

expectations about K_{cells} and K_{bio} ($B_{stationary} = 0.60$) because cultures reached carrying capacity in the

latter part of the common garden and because cell size (which affects estimates of K_{bio}) stabilised

around day 12 (Fig. S2). We estimated maximum total biomass (K_{bio}) as the product of K_{cells} obtained from the population growth models and the average cell size for each replicate over the last three sampling days (day 12 to 16), when cell size stabilised. Finally, we determined the empirical scaling of each demographic parameter with cell size across the focal populations, using the average cell size calculated over the same intervals above (i.e., over the first 5 days for r_{max}, between day 12 and 16 for K_{cells} and K_{bio}).

Acknowledgements: We thank Mike McDonald and John DeLong for valuable comments on earlier versions of the manuscript. We also thank Jiaye Qin for assistance in data collection and the Quantitative Biology Unit at the Instituto Gulbenkian de Ciência for statistical advice. This work was supported by the Australian Research Council through a DECRA fellowship (DE190100660) and by a fellowship (LCF/BQ/PI21/11830001) from "la Caixa" Foundation (ID 100010434) and the European Union's Horizon 2020 research and innovation programme under the Marie Skłodowska Curie grant agreement No 847648 to GG.

References

1. Aubree, F. *et al.* How community adaptation affects biodiversity–ecosystem functioning relationships. *Ecol. Lett.* **23**, 1263–1275 (2020).

2. Post, D. M. & Palkovacs, E. P. Eco-evolutionary feedbacks in community and ecosystem ecology: interactions between the ecological theatre and the evolutionary play. *Philos. Trans. R. Soc. B Biol. Sci.* (2009) doi:10.1098/rstb.2009.0012.

3. Hart, S. P., Turcotte, M. M. & Levine, J. M. Effects of rapid evolution on species coexistence. *Proc. Natl. Acad. Sci.* **116**, 2112 (2019).

4. Jousset, A., Eisenhauer, N., Merker, M., Mouquet, N. & Scheu, S. High functional diversity stimulates diversification in experimental microbial communities. *Sci. Adv.* **2**, e1600124 (2016).

5. Lawrence, D. *et al.* Species Interactions Alter Evolutionary Responses to a Novel Environment. *PLOS Biol.* **10**, e1001330 (2012).

6. Scheuerl, T. *et al.* Bacterial adaptation is constrained in complex communities. *Nat. Commun.* **11**,
754 (2020).

7. García, F. C. *et al.* The temperature dependence of microbial community respiration is amplified
by changes in species interactions. *Nat. Microbiol.* **8**, 272–283 (2023).

8. Barraclough, T. G. How Do Species Interactions Affect Evolutionary Dynamics Across Whole
Communities? *Annu. Rev. Ecol. Evol. Syst.* **46**, 25–48 (2015).

9. Grant, P. R. & Grant, B. R. Evolution of Character Displacement in Darwin's Finches. *Science*
313, 224–226 (2006).

10. Rainey, P. B. & Travisano, M. Adaptive radiation in a heterogeneous environment. *Nature* **394**,
69–72 (1998).

11. Schluter, D. Experimental Evidence That Competition Promotes Divergence in Adaptive
Radiation. *Science* **266**, 798–801 (1994).

12. Abrams, P. A. Alternative Models of Character Displacement and Niche Shift. I. Adaptive Shifts in
Resource Use When There is Competition for Nutritionally Nonsubstitutable Resources. *Evolution*
41, 651–661 (1987).

13. Fox, J. W. & Vasseur, D. A. Character Convergence under Competition for Nutritionally Essential
Resources. *Am. Nat.* **172**, 667–680 (2008).

14. Gallego, I. & Narwani, A. Ecology and evolution of competitive trait variation in natural
phytoplankton communities under selection. *Ecol. Lett.* **25**, 2397–2409 (2022).

15. TerHorst, C. P. *et al.* Evolution in a Community Context: Trait Responses to Multiple Species
Interactions. *Am. Nat.* **191**, 368–380 (2018).

16. Brown, J. H., Gillooly, J. F., Allen, A. P., Savage, V. M. & West, G. B. Toward a metabolic theory
of ecology. *Ecology* **85**, 1771–1789 (2004).

17. Savage, V. M., Gillooly, J. F., Brown, J. H., West, G. B. & Charnov, E. L. Effects of body size and
temperature on population growth. *Am. Nat.* **163**, 429–441 (2004).

18. Damuth, J. Population-density and body size in mammals. *Nature* **290**, 699–700 (1981).

19. Isaac, N. J. B., Carbone, C. & McGill, B. Population and Community Ecology. in *Metabolic*
Ecology (eds. Sibly, R. M., Brown, J. H. & Kodric-Brown, A.) (2012).

20. Malerba, M. E. & Marshall, D. J. Size-abundance rules? Evolution changes scaling relationships
between size, metabolism and demography. *Ecol. Lett.* **22**, 1407–1416 (2019).

21. Berga, M., Székely, A. J. & Langenheder, S. Effects of disturbance intensity and frequency on

bacterial community composition and function. *PLoS ONE* **7**, e36959–e36959 (2012).

22. Pomati, F., Jokela, J., Castiglioni, S., Thomas, M. K. & Nizzetto, L. Water-borne pharmaceuticals

reduce phenotypic diversity and response capacity of natural phytoplankton communities. *PLoS*

ONE **12**, e0174207 (2017).

23. Poulson-Ellestad, K. L. *et al.* Metabolomics and proteomics reveal impacts of chemically

mediated competition on marine plankton. *Proc. Natl. Acad. Sci.* **111**, 9009 (2014).

24. de Villemereuil, P., Gaggiotti, O. E., Mouterde, M. & Till-Bottraud, I. Common garden experiments

in the genomic era: new perspectives and opportunities. *Heredity* **116**, 249–254 (2016).

25. Marshall, D. J. *et al.* Long-term experimental evolution decouples size and production costs in

Escherichia coli. *Proc. Natl. Acad. Sci.* **119**, e2200713119 (2022).

26. Bernhardt, J. R. *et al.* The evolution of competitive ability for essential resources. *Philos. Trans.*

R. Soc. B Biol. Sci. **375**, 20190247 (2020).

27. Bassar, R. D. *et al.* Experimental Evidence for Density-Dependent Regulation and Selection on

Trinidadian Guppy Life Histories. *Am. Nat.* **181**, 25–38 (2013).

28. Pianka, E. R. On r- and K-Selection. *Am. Nat.* **104**, 592–597 (1970).

29. Bierbaum, T. J., Mueller, L. D. & Ayala, F. J. Density-Dependent Evolution of Life-History Traits in

Drosophila Melanogaster. *Evolution* **43**, 382–392 (1989).

30. Mueller, L. D. & Ayala, F. J. Trade-off between r-selection and K-selection in Drosophila

populations. *Proc. Natl. Acad. Sci.* **78**, 1303–1305 (1981).

31. Gorter, F. A., Manhart, M. & Ackermann, M. Understanding the evolution of interspecies

interactions in microbial communities. *Philos. Trans. R. Soc. B Biol. Sci.* **375**, 20190256 (2020).

32. Falkowski, P. G., Fenchel, T. & Delong, E. F. The Microbial Engines That Drive Earth's

Biogeochemical Cycles. *Science* **320**, 1034–1039 (2008).

33. Field, C. B., Behrenfeld, M. J., Randerson, J. T. & Falkowski, P. Primary Production of the

Biosphere: Integrating Terrestrial and Oceanic Components. *Science* **281**, 237 (1998).

34. Hillebrand, H. *et al.* Cell size as driver and sentinel of phytoplankton community structure and

functioning. *Funct. Ecol.* **36**, 276–293 (2022).

35. Yvon-Durocher, G., Montoya, J. M., Trimmer, M. & Woodward, G. U. Y. Warming alters the size
spectrum and shifts the distribution of biomass in freshwater ecosystems. *Glob. Change Biol.* **17**,
1681–1694 (2011).

36. Barton, S. *et al.* Evolutionary temperature compensation of carbon fixation in marine
phytoplankton. *Ecol. Lett.* **23**, 722–733 (2020).

37. Padfield, D., Yvon-Durocher, G., Buckling, A., Jennings, S. & Yvon-Durocher, G. Rapid evolution
of metabolic traits explains thermal adaptation in phytoplankton. *Ecol. Lett.* **19**, 133–142 (2016).

38. Govaert, L., Gilarranz, L. J. & Altermatt, F. Competition alters species' plastic and genetic
response to environmental change. *Sci. Rep.* **11**, 23518 (2021).

39. Malerba, M. E., White, C. R. & Marshall, D. J. Eco-energetic consequences of evolutionary shifts
in body size. *Ecol. Lett.* **21**, 54–62 (2017).

40. Bell, G. Experimental Evolution of Heterotrophy in a Green Alga. *Evolution* **67**, 468–476 (2013).

41. Collins, S. Competition limits adaptation and productivity in a photosynthetic alga at elevated
CO2. *Proc Biol Sci* **278**, 247–55 (2011).

42. Maynard, D. S., Crowther, T. W. & Bradford, M. A. Competitive network determines the direction
of the diversity–function relationship. *Proc. Natl. Acad. Sci.* **114**, 11464 (2017).

43. Vallina, S. M., Cermeno, P., Dutkiewicz, S., Loreau, M. & Montoya, J. M. Phytoplankton functional
diversity increases ecosystem productivity and stability. *Ecol. Model.* **361**, 184–196 (2017).

44. Osmond, M. M. & de Mazancourt, C. How competition affects evolutionary rescue. *Philos Trans R
Soc Lond B Biol Sci* **368**, 20120085–20120085 (2013).

45. MacArthur, R. H. & Wilson, E. O. *The Theory of Island Biogeography*. (Princeton University
Press, 1967).

46. Sæther, B.-E., Visser, M. E., Grøtan, V. & Engen, S. Evidence for r- and K-selection in a wild bird
population: a reciprocal link between ecology and evolution. *Proc. R. Soc. B Biol. Sci.* **283**,
20152411 (2016).

47. Lande, R., Engen, S. & Sæther, B.-E. An evolutionary maximum principle for density-dependent
population dynamics in a fluctuating environment. *Philos. Trans. R. Soc. B Biol. Sci.* **364**, 1511–
1518 (2009).

48. Kentie, R., Clegg, S. M., Tuljapurkar, S., Gaillard, J.-M. & Coulson, T. Life-history strategy varies

with the strength of competition in a food-limited ungulate population. *Ecol. Lett.* **23**, 811–820

(2020).

49. Gallego, I., Venail, P. & Ibelings, B. W. Size differences predict niche and relative fitness

differences between phytoplankton species but not their coexistence. *ISME J.* **13**, 1133–1143

(2019).

50. Brandl, S. J. *et al.* Metabolic rates mirror morphological and behavioral differences in two sand-

dwelling coral reef gobies. *Mar. Ecol. Prog. Ser.* **684**, 79–90 (2022).

51. Alexander, J. M., Diez, J. M. & Levine, J. M. Novel competitors shape species' responses to

climate change. *Nature* **525**, 515–518 (2015).

52. Lancaster, L. T., Morrison, G. & Fitt, R. N. Life history trade-offs, the intensity of competition, and

coexistence in novel and evolving communities under climate change. *Philos. Trans. R. Soc. B*

Biol. Sci. **372**, 20160046 (2017).

53. White, C. R., Alton, L. A., Bywater, C. L., Lombardi, E. J. & Marshall, D. J. Metabolic scaling is the

product of life-history optimization. *Science* **377**, 834–839 (2022).

54. Guillard, R. R. L. & Ryther, J. H. Studies of marine planktonic diatoms. I. *Cyclotella nana* Hustedt

and *Detonula confervacea* Cleve. *Can. J. Microbiol.* **8**, 229–239 (1962).

55. Schindelin, J. *et al.* Fiji: an open-source platform for biological-image analysis. *Nat. Methods* **9**,

676–682 (2012).

56. Hillebrand, H., Dürselen, C.-D., Kirschtel, D., Pollingher, U. & Zohary, T. Biovolume calculation for

pelagic and benthic microalgae. *J. Phycol.* **35**, 403–424 (1999).

57. Lenski, R. E. Phenotypic and Genomic Evolution during a 20,000-Generation Experiment with the

Bacterium Escherichia coli. in *Plant Breeding Reviews* 225–265 (John Wiley & Sons, Ltd, 2003).

doi:10.1002/9780470650288.ch8.

58. Grant, N. A., Abdel Magid, A., Franklin, J., Dufour, Y. & Lenski, R. E. Changes in Cell Size and

Shape during 50,000 Generations of Experimental Evolution with Escherichia coli. *J. Bacteriol.*

203, e00469-20 (2021).

59. Ryabov, A. *et al.* Shape matters: the relationship between cell geometry and diversity in

phytoplankton. *Ecol. Lett.* **24**, 847–861 (2021).

60. Litchman, E. & Klausmeier, C. A. Trait-Based Community Ecology of Phytoplankton. *Annu. Rev.*

Ecol. Evol. Syst. **39**, 615–639 (2008).

61. Tan, H., Hirst, A. G., Atkinson, D. & Kratina, P. Body size and shape responses to warming and

resource competition. *Funct. Ecol.* **35**, 1460–1469 (2021).

62. Malerba, M. E., White, C. R. & Marshall, D. J. Phytoplankton size-scaling of net-energy flux

across light and biomass gradients. *Ecology* **98**, 3106–3115 (2017).

63. Ghedini, G., Marshall, D. J. & Loreau, M. Phytoplankton diversity affects biomass and energy

production differently during community development. *Funct. Ecol.* **36**, 446–457 (2022).

64. Olito, C., White, C. R., Marshall, D. J. & Barneche, D. R. Estimating monotonic rates from

biological data using local linear regression. *J. Exp. Biol.* **220**, 759 (2017).

65. White, C. R., Kearney, M. R., Matthews, P. G. D., Kooijman, S. A. L. M. & Marshall, D. J. A

Manipulative Test of Competing Theories for Metabolic Scaling. *Am. Nat.* **178**, 746–754 (2011).

66. Williams, P. J. le B. & Laurens, L. M. L. Microalgae as biodiesel & biomass feedstocks: Review &

analysis of the biochemistry, energetics & economics. *Energy Environ. Sci.* **3**, 554–590 (2010).

67. Pinheiro, J. & Bates, D. nlme: Linear and Nonlinear Mixed Effects Models. R package version

3.1-161. *R Core Team* (2022) doi:https://CRAN.R-project.org/package=nlme.

68. Bates, D., Mächler, M., Bolker, B. & Walker, S. Fitting Linear Mixed-Effects Models Using lme4.

2015 **67**, 48 (2015).

69. Fox, J. & Weisberg, S. An {R} Companion to Applied Regression, Third Edition. Thousand Oaks

CA: Sage. URL: https://socialsciences.mcmaster.ca/jfox/Books/Companion/. (2019).

70. Wickham, H. ggplot2: Elegant Graphics for Data Analysis. *Springer-Verl. N. Y.* **ISBN 978-3-319-**

24277-4, (2016).

71. Malerba, M. E., Palacios, M. M. & Marshall, D. J. Do larger individuals cope with resource

fluctuations better? An artificial selection approach. *Proc. R. Soc. B Biol. Sci.* **285**, 20181347

(2018).