Dual receptive fields underlying target and motion sensitivity in looming sensitive descending neurons

Sarah Nicholas[1], Yuri Ogawa[1] and Karin Nordström[1,2]*

[1]Flinders Health and Medical Research Institute, Flinders University, GPO Box 2100, Adelaide, SA, 5001, Australia; [2]Department of Medical Cell Biology, Uppsala University, 75123 Uppsala, Sweden

*email: karin.nordstrom@flinders.edu.au

ORCIDs

Sarah Nicholas: 0000-0002-5555-9421

Yuri Ogawa: 0000-0002-9708-7063

Karin Nordström: 0000-0002-6020-6348

Keywords

Target motion, insect vision, motion vision, hoverfly, descending neuron, wide-field motion

Abstract

When an animal is pursuing a target, such as a bird pursuing prey, the retinal projection of the target is initially very small. However, prior to successful capture, as the pursuer nears the target, the image becomes a rapidly expanding object. This suggests that different neural pathways underlie these two behavioural stages: one tuned to small retinal objects, and one to larger, looming stimuli. Indeed, in zebrafish, two different optic brain regions support the initial visualization of prey and the final stages of prey capture. The pursuit of small targets is not limited to vertebrates, with many invertebrates also utilizing these capabilities for prey capture, territorial defences, or courtship. Male *Eristalis tenax* hoverflies, which are highly territorial, have target tuned neurons that likely underlie the initial stages of target detection and pursuit. However, the underlying neuronal control of the final stages of prey capture is currently unknown. We here show a class of descending neurons with dual sensitivity, separated into two distinct receptive fields. The dorsal receptive field shows sensitivity to small targets whereas the ventral receptive field responds to larger objects and looming stimuli. We show that the two receptive fields have different pre-synaptic input, where the inputs are not linearly summed. This novel and unique arrangement could thus support the different stages of target pursuit behavior, where targets are initially detected in the dorsal receptive field, whilst the ventral receptive field is used during the final stages of pursuit and eventual capture.

Introduction

The ability to respond quickly to visual stimuli is vital for the survival of many animals. Indeed, visual input may be used for a variety of tasks, from navigating and avoiding obstacles, choosing a suitable surface to rest upon, and also for detecting predators or prey. For example, many predatory insects rely on vision to identify suitable prey, which they then pursue with astonishing precision and accuracy (e.g. 1, 2, 3). Some non-predatory insects also have superb target detecting capabilities, which they may use to defend their territory against intruders or to pursue potential mates for courtship (4, 5).

Blowflies, houseflies and hoverflies pursue not only conspecifics in the field, but also artificial targets in man-made arenas (6-8), thus allowing more precise quantification of the target image during pursuit. Such experiments have shown that male *Eristalis tenax* hoverflies initiate pursuit of targets that subtend about $0.9 – 1.7°$ moving at $20 – 170°/s$ ($25^{th} – 75^{th}$ percentiles) (7). Small target motion detector (STMD) neurons in the hoverfly optic lobe, and their presumed post-synaptic targets, the target selective descending neurons (TSDNs), have size and velocity tuning properties that match the target image at the start of the pursuit, suggesting that they could support these behaviours (9, 10). However, later during pursuit, as the hoverfly closes in on the target, its projected retinal image becomes larger, and eventually becomes a looming stimulus (7). This suggests that another subset of neurons, tuned to larger and looming objects, may be recruited in the later stages of target pursuit (see e.g. Discussion in Ref. 11). Indeed, in zebrafish larvae separate neuronal pathways control the initial detection of prey and initiation of pursuit, versus subsequent prey tracking and capture manoeuvres (12).

Neurons that respond well to rapidly looming stimuli exist in a range of species and visual structures, including the cat superior colliculus (13), the bullfrog optic tectum (14), and in

zebrafish retinal ganglion cells (15). In insects, there are looming neurons in the optic lobes

as well as in the descending pathways. For example, the locust LGMD/DCMD system (16),

the *Drosophila* Foma-1 neurons (17) and descending giant fiber (18), are likely used for rapid

avoidance of bird predators. Interestingly, many of these looming neurons also respond

strongly to small moving targets. For example, the locust LGMD/DCMD pathway was

originally thought to play a role in object tracking (19), and several looming sensitive

neurons in the locust central complex also respond to small moving targets (20). Conversely,

some dragonfly target selective descending neurons (TSDNs) respond to both targets and to

looming stimuli (21)(22).

Historically, looming neurons have been studied in the context of predator avoidance (e.g. 16,

18, 23). However, there is emerging evidence that such neurons also play a key role in pursuit

behaviours. For example, zebrafish larvae usually detect prey in the lateral visual field, but

then centre the prey image (24, 25), where it becomes a rapidly expanding stimulus. This last

part of the pursuit is likely controlled by loom sensitive, non-directional, inter-ocular neurons

in the anterior tectum (12, 26). Similarly, silencing *Drosophila* Foma-1 neurons does not only

affect the escape response (17), but also the ability of male flies to follow females during

courtship (27).

Taken together, this suggests that some neurons classically defined as either target or

looming selective (21, 28) respond to both, and this dual sensitivity could serve a useful role

in pursuit behaviours. Like locusts (28) and dragonflies (21), hoverflies have descending

neurons that respond to both looming stimuli and to small moving targets (9). We here

investigate this dual sensitivity in detail and find these neurons have two distinct receptive

fields, one dorsal that responds selectively to the motion of small targets, and one ventral that

responds to larger objects, including sinusoidal gratings and high-contrast edges. We show

that when the centre of the ventral grating receptive field is to the right of the visual midline,

it prefers motion to the right, and vice versa. However, there is no correlation between the

preferred direction of the dorsal target receptive field and the ventral grating receptive field.

We also show that the two receptive fields receive separate input, from the pre-synaptic target

pathway and the pre-synaptic widefield motion pathway, respectively. However, when

receiving dual input, these are not linearly summed. We hypothesize that the unique response

characteristics of these neurons make them highly suitable for the control of different stages

of target pursuit, with distinct receptive fields sensitive to both small targets and larger

rapidly looming objects.

Materials and Methods

Electrophysiology

We recorded from 102 looming sensitive descending neurons (9) in 99 male *Eristalis tenax*

hoverflies, reared and maintained in-house as described previously (29). At experimental

time the hoverfly was immobilized ventral side up, using a beeswax and resin mixture, before

an opening was made in the thoracic cavity. A small silver hook was used to elevate and

support the cervical connective and a silver wire inside the opening served as a reference

electrode.

Recordings were made from the cervical connective using a sharp polyimide-insulated

tungsten microelectrode (2 MOhm, Microprobes, Gaithersburg, USA). Signals were

amplified at 100x gain and filtered through a 10 – 3000 Hz bandwidth filter using a DAM50

differential amplifier (World Precision Instruments, Sarasota, USA), with 50 Hz noise

removed with a HumBug (Quest Scientific, North Vancouver, Canada). The data were

digitized via a Powerlab 4/30 and recorded at 40 kHz with LabChart 7 Pro software

(ADInstruments, Sydney, Australia). Single units were discriminated by amplitude and half-

width using Spike Histogram software (ADInstruments).

Visual stimuli

Hoverflies were positioned perpendicular to and 6.5 cm from the middle of a linearized Asus

LCD screen (Asus, Taipei, Taiwan) with a mean illuminance of 200 Lux, a refresh rate of

165 Hz and a spatial resolution of 2,560 × 1,440 pixels (59.5 x 33.5 cm), giving a projected

screen size of 155 × 138°. Visual stimuli were displayed using custom software written in

Matlab (R2019b, Mathworks) using the Psychophysics toolbox (30, 31). The stimuli were not

perspective corrected. When values are given in degrees, this corresponds to the retinal size

in the centre of the visual field. Velocities are given in pixels/s.

Potential looming sensitive descending neurons were initially identified based on their

response to a small, black, moving target (left, Fig. 1, and see Ref. 9). Those neurons that

also responded strongly to a looming stimulus compared to an appearance control (9) were

kept for further analysis. The looming stimulus was a black circle on a white background,

expanding over 1 s from 1° diameter to 117° (right, Fig. 1), with a 10 ms rate of expansion

(32), also referred to as l/|v|. The appearance control was a black circle with 117° diameter

that appeared and remained on the screen for 1 s.

We mapped the target receptive field (9) of each neuron by scanning a target horizontally and

vertically at 20 evenly spaced elevations and azimuths (9) to create a 20 x 20 grid (Fig. S1A).

The 15 x 15 pixel (3 x 3°) black, square target moved at a velocity of 900 pixels/s. There was

a minimum 0.5 s interval between each stimulation.

We mapped the grating receptive field (9) using local sinusoidal gratings (400 x 400 pixels,

71 x 71° in the visual field centre, Fig. S1B) where the internal pattern moved in a series of 8

different directions presented in a pseudorandom order for 0.36 s each. The gratings had a

wavelength of 75 pixels (13° for the central patch, 0.08 cpd) and drifted at 5 Hz. The local

gratings were placed in an overlapping tiling fashion so that 8 x 6 (width x height) squares

covered the majority of the screen (Fig. S1B). There was a minimum 1 s interval between

each stimulation.

To map the leading-edge receptive field, we scanned the entire height or width of the screen

with an OFF-contrast edge moving left, right, down or up, at 900 pixels/s (Fig. S3A-D).

For size tuning experiments a bar drifted at 900 pixels/s, in the preferred direction of each

neuron's target receptive field. The bar drifted either horizontally or vertically through the

centre of the target or grating receptive field, as specified. The bar side parallel to the

direction of travel was maintained at a fixed size of 15 pixels, i.e. 3°, whilst the perpendicular

side varied from 0.2° to 155° (1 pixel to the full height of the screen). When presented

simultaneously with a small target, the bar moved in the preferred horizontal direction

through the grating receptive field only. In this case the bar height was varied between 5.7°

and 106° (28 to 749 pixels), whilst a fixed size 3 x 3° target moved through the target

receptive field.

To determine the input mechanism of each receptive field, an OFF edge, an ON edge and a

discrete bar, with a width of 15 pixels (3°) drifted horizontally at 900 pixels/s in each

neuron's preferred direction, across the entire width of the screen. The height of these objects

was 3° (15 pixels) when drifted through the target receptive field, 84° (500 pixels) when

drifted through the grating receptive field, or the height of the screen (138°) to cover both

receptive fields.

All stimuli were presented in a random order, with the exception of the receptive field

stimuli, which were presented in a pseudo-random order.

Data analysis and statistics

All data analysis was performed in Matlab and GraphPad Prism 9.3.1 (GraphPad Software,

USA). The sample size, statistical test and P-values are indicated in each figure legend, where

n refers to the number of repetitions within one neuron and *N* to the number of neurons.

Neurons were initially identified based on the response to a small target, with data from all

neurons that passed our definition of a looming neuron (Ref. 9, and see above) included in

the analysis.

For target receptive field mapping we used the resulting 20 x 20 grid (Fig. S1A) to calculate

the local preferred direction and local motion sensitivity (Fig. S1C), assuming a neural delay

of 20 ms. We calculated the local average response to the four directions of motion (dotted

line, Fig. S1C) after subtracting the spontaneous rate, calculated in the 485 ms prior to

stimulus presentation. We interpolated this to 100 x 100 grid to generate receptive field maps

(Fig. S1E) using Matlab's *contour* function. We defined the centre of the receptive field (Fig.

S1E) as the centre of the 50% contour line using Matlab's *centroid* function. We fitted a

cosine function (9) to the response to the four directions of motion (10) and extracted its

preferred direction and amplitude (Fig. S1C). We calculated the preferred direction for each

neuron by averaging the local preferred directions from the locations where the local motion

sensitivity was above 50% of the maximum (blue, Fig. S1G).

For the grating receptive fields we used the resulting 8 x 6 grid (Fig. S1B) to quantify the

local mean response for each direction of motion, after removing the first 100 ms of the

response, to avoid any initial onset transients (33, 34). We calculated the local mean response

(dotted line, Fig. S1D) after subtracting the spontaneous rate, calculated for 800 ms preceding

stimulus presentation. We spatially interpolated this 10 times and calculated the centre from

the 50% contour line (Fig. S1F). For each spatial location we fitted a cosine function (Fig.

S1D) to the response to get the local preferred direction and local motion sensitivity (9). We

calculated the overall direction selectivity using the top 50% of the local preferred directions

(red, Fig. S1H).

We calculated the horizontal and vertical distance between the receptive field centres and the

midline and equator. The distance between the two receptive field centres was calculated

using the Euclidean distance.

For leading-edge receptive field mapping we first quantified the spike histogram for each

neuron, after smoothing the response with a 100 ms square-wave filter with 40 kHz

resolution (Fig. S3A-D). We identified the maximum response to any direction of motion

(purple, Fig. S3D). 50% maximum response was used as a threshold to determine the limits

of the leading-edge receptive field (cyan, Fig. S3A, C, D). If a neuron's response did not

reach threshold to one direction of motion (Fig. S3B), the opposite direction of motion

determined the receptive field outlines. If a neuron responded to both directions of motion

(e.g. up and down) the outer thresholds were used to delineate the receptive field (Fig. S3E).

From the resulting rectangular receptive field, we determined the centre, and the proximity to

the target and the grating receptive fields respectively (Fig. S3F), using the following

proximity index:

$$(d1 - d2) / (d1 + d2)$$

where $d1$ is the Euclidean distance between the leading edge and the target receptive field

centres, and $d2$ the Euclidean distance between the leading edge and the grating receptive

field centres (Fig. 4B). Thus, if the leading-edge receptive field centre was closer to the

grating receptive field centre, the proximity index was positive, but if it was closer to the

target receptive field centre the proximity index was negative.

For all stimuli other than receptive field mapping, quantification of responses was done by

averaging the spike rate within a 0.56 s analysis window centred on each neuron's target or

grating receptive field centre, as specified.

Results

Looming neurons have dorsal target receptive fields and ventral grating receptive fields

We recorded from 102 looming sensitive descending neurons in male *Eristalis tenax*

hoverflies. The neurons described here responded both to small target motion (left, Fig. 1)

and to looming stimuli (right, Fig. 1). The response to a looming stimulus (right, Fig. 1B, D)

started well before the stimulus reached its full size (right, Fig. 1C), and was much stronger

than the response to an appearance control (not shown, but see Ref. 9).

To investigate this dual sensitivity (Fig. 1) in more detail, we mapped the receptive fields

using two different methods. For this purpose we either scanned the visual monitor with a

small, black target (10) moving in four different directions (Fig. S1A, C), or we used a local

sinusoidal grating (Fig. S1B, D) where the internal pattern drifted in eight different directions

(9). The data from an example neuron shows two distinct receptive fields, with a dorsal target

receptive field (blue, Fig. 2A, S1E), and a ventral grating receptive field (red, Fig. 2B, S1F).

We used the 50% response contours to define the two receptive field centres (blue and red

circle, Fig. 2A, S1E, F). Across the 102 neurons, the target receptive fields cluster above the

visual equator (blue, Fig. 2B), whereas the grating receptive field centres cluster below the

equator (red, Fig. 2B), even if there are some exceptions (see also Fig. S2). We quantified the

vertical distance between each receptive field centre and the visual equator (41° and -34° in

the example neuron, Fig. 2A), and found a bimodal distribution with target receptive field

centres peaking 30-40° dorsal and grating receptive field centres peaking 30-40° ventral (Fig.

2B, C).

We next quantified the horizontal distance between each receptive field centre and the visual

midline (14° and 18° in the example neuron, Fig. 2A). Across neurons we found a bimodal

distribution with a gap along the visual midline and a peak distribution about 20° either side

of the visual midline (Fig. 2B, D). There was no clear lateral distribution difference between

the two receptive field types.

We noted that the target receptive field was often in the dorsal visual field, but that there

were some exceptions (Fig. 2A-C, S2). We next investigated if the target receptive field was

always dorsal to the grating receptive field and determined the Euclidean distance between

the two receptive field centres (black line, 69°, Fig. 2A). Across neurons we found that the

target receptive field was indeed most often dorsal to the grating receptive field (grey, Fig.

2E), and that the distance could be quite large. When the grating receptive field was more

dorsal, the two receptive field centres tended to be closer to each other (black, Fig. 2E).

The grating receptive field is sensitive to motion away from the midline

We next determined the local motion sensitivity and average preferred direction of each

neuron's target and grating receptive field (coloured arrows, Fig. S1G, H, S2). In the example

neuron, the preferred direction of the target receptive field is up and toward the visual

midline (blue arrows, Fig. 2A, S1G), whereas the preferred direction of the grating receptive

field is horizontally to the right (red arrows, Fig. 2A, S1H). For comparison across neurons

we colour coded the preferred direction into four cardinal directions, and plotted them as a

function of receptive field centre location. This analysis shows that the preferred direction of

the target receptive fields was weakly correlated with location (Fig. 3A), with a slight

preference for motion away from the visual midline (Fig. 3C). In contrast, the directionality

of the grating receptive fields was strongly correlated with centre location (Fig. 3B), with a

strong dominance for motion away from the visual midline (Fig. 3D).

We next quantified the difference between the preferred directions of the target and the

grating receptive fields. In the example neuron, this is 141° (bottom right pictogram, Fig. 2A,

see also Fig. S1G, H). Across neurons there was a slight preference for similar directionality

in the two receptive fields, albeit with neurons encompassing the entire span of possible

directionality differences (Fig. 3E, see also Fig. S2). There was no distribution difference

based on whether the two receptive field centres were on the same side (grey, Fig. 3E) or

opposite sides (black, Fig. 3E) of the visual midline.

Leading-edge sensitivity matches grating receptive field

These descending neurons thus have two receptive fields, one that responds to small target

motion (blue, Fig. 2, S1, S2) and one that responds to local sinusoidal gratings (red, Fig. 2,

S1, S2). Which one of these receptive fields is most likely to contribute to their looming

sensitivity (right, Fig. 1)? Looming sensitive neurons in *Drosophila,* including the giant fiber,

also respond strongly to high-contrast bars and sweeping high-contrast edges (35). We thus

used full screen OFF edges to map the looming receptive field (Fig. S3). An example neuron

shows strong responses to an OFF edge sweeping either left (Fig. S3A), down (Fig. S3C), or

up (Fig. S3D) across the visual field, but not right (Fig. S3B). The resulting leading-edge

receptive field (cyan, Fig. S3E, F) overlaps substantially with the grating receptive field (red,

Fig. S3F), but not the target receptive field (blue, Fig. S3F).

Across neurons we compared the location of the leading-edge, the target, and the grating

receptive field centres (circles, Fig. 4A, S3F). A qualitative analysis shows that the leading-

edge receptive field centres tend to cluster below the visual equator, just like the grating

receptive field centres (cyan and red, Fig. 4A). For quantification, we calculated a proximity

index (Fig. 4B). When the leading-edge receptive field centre is closer to the grating

receptive field centre the proximity index is positive, with values close to 100% indicating

that they are much closer than to the target receptive field centre. In the example neuron, the

proximity index is 83% (Fig. 4B, S3). Across neurons, we found that the leading-edge

receptive field centre was frequently closer to the grating receptive field centre (red, Fig. 4C).

In those neurons where it was closer to the target receptive field centre, the proximity index

was relatively low (blue, Fig. 4C). In summary, it is likely that these neurons get their

looming sensitivity predominantly within the ventral visual field, overlapping with the

location of the grating receptive field.

Different size response function in the two receptive fields

Our previous work showed that looming sensitive neurons have a peculiar size response

function, with one peak to bars of a few degrees height, similar to the size tuning of target

selective neurons, and a second peak to full-screen bars (9). To investigate if this size

sensitivity differs between the two receptive fields we scanned bars of fixed width across the

visual monitor, while varying the height. We used two different trajectories, one centred on

the target receptive field (blue, Fig. 5A, S4), and one on the grating receptive field (red, Fig.

5A, S4). We found that the neurons responded strongly to small bars moving through the

target receptive field (blue, Fig. 5A, Fig. S4A-E), but not through the grating receptive field

(red, Fig. 5A, Fig. S4A-E). For the middle sized bars, the neurons responded weakly whether

they traversed the target or the grating receptive field (Fig. 5A, S4F-I).

When the bars were extended to cover a large part of the screen, they traversed both receptive

fields (Fig. S4J, K), making it hard to determine which receptive field the strong response

came from (Fig. 5A). To bypass this, we scanned the bars vertically instead of horizontally,

so that they traversed the grating receptive field and the target receptive field at different

points in time (pictogram, Fig. 5B, S5). We found that the neurons responded strongly when

small bars moved through the target analysis window (blue, Fig. 5B, S5A-E), and strongly to

large bars when they moved through the grating analysis window (red, Fig. 5B, S5G-L). This

shows that the target receptive field is tuned to small targets, whereas the grating receptive

field responds better to full-screen bars.

Separate inputs to the two receptive fields

The data above show that looming neurons have two receptive fields (Fig. 2-4), with size

tuning, suggesting that they receive separate input (Fig. 5). What happens when they are

stimulated simultaneously? To investigate this, we first determined the response when the

two receptive fields were stimulated separately, by scanning a small target through the target

receptive field (blue, Fig. 6A), and a series of bars through the grating receptive field (red,

Fig. 6A, consistent with the data in Fig. 5). We compared this to the response to simultaneous

stimulation (black, Fig. 6B). Interestingly, the response to simultaneous presentation (black,

Fig. 6B) was smaller than the linear sum of the two independent presentations (purple, Fig.

6B). Importantly, this cannot be due to response saturation, as the linear sum to the smallest

bars (purple, Fig. 6B) is on par with the measured response to the largest bars (black, Fig.

6B).

An alternative hypothesis is that the neuron selectively attends to one of the objects (36),

either the small target in the target receptive field, or the bar in the grating receptive field. We

investigated this by comparing the biggest response for each neuron (green, Fig. 6C), whether

that would be to the target or the bar, with the response to the simultaneous presentation

(black, Fig. 6C). We found that while there was a significant dependence on bar size, there

was no significant difference between the two conditions (compare green and black, Fig. 6C,

2-way ANOVA), suggesting that the neuron could be selectively attending to one of the

objects and ignoring the other.

Target receptive field is based on 1-point correlator input, whereas the grating receptive

field uses 2-point correlator input

The data above show two independent receptive fields. What is the likely pre-synaptic input

to each? Both optic lobe and descending target tuned neurons (37, 38) generate their target

selectivity using 1-point correlators, which are based on the comparison of an OFF contrast

change immediately followed by an ON contrast change at a single point in space (39). These

correlators are fundamentally different from 2-point correlators, such as Hassenstein-

Reichard elementary motion detectors (40), in their response to high-contrast edges. For

example, a 1-point correlator will respond only weakly to either OFF or ON contrast edges,

compared with complete objects, whereas 2-point correlators respond equally well to single

edges and complete objects (inset, Fig. 7, data replotted from Ref. 37).

To investigate the potential input to the two receptive fields we first scanned an OFF edge

through the grating receptive field, then an ON edge, followed by a complete bar (red, Fig.

7). We found that the responses to single edges were similar to the response to a complete bar

(ns, red, Fig. 7). The response was consistent with an underlying 2-point correlator input

(compare red data with inset, Fig. 7). In contrast, when we scanned edges or targets through

the target receptive field, the response to a complete target was much stronger than to either

OFF or ON edges on their own ($P < 0.0001$, blue, Fig. 7). Indeed, the response was consistent

with an underlying 1-point correlator input (compare blue data with inset, Fig. 7). In response

to full-screen edges and bars, which cover both receptive fields, we found the strongest

response to the OFF edge (P = 0.0049 for OFF vs bar, ns for ON vs bar, black, Fig. 7).

Discussion

We have shown here a group of descending neurons in the hoverfly *Eristalis tenax* that are

sensitive to both targets and to looming stimuli (Fig. 1). We show that this dual sensitivity

originates from two discrete receptive fields, with different locations (Fig. 2, S1E, F, S2) and

preferred directions (Fig. 3, S1G, H, S2). We show that the looming sensitivity is likely

associated with the ventral receptive field (Fig. 4, S3). The size tuning (Fig. 5, S4, S5) and

sensitivity to OFF and ON contrast edges (Fig. 7) supports independent input to the two

receptive fields, using two fundamentally different pre-synaptic pathways. Interestingly, the

input from the two pathways is not linearly summed (Fig. 6B) but could be subject to

selective attention (Fig. 6C), as previously shown in dragonfly target neurons (36).

Dual receptive fields

The neurons that we describe here were classified as looming sensitive based on a strong

response to a looming stimulus (right, Fig. 1) compared with an appearance control (9).

However, as they also respond strongly to small moving targets (left, Fig. 1, Fig. 5), they

could have been classified as target selective descending neurons (TSDN). Indeed, the

dragonfly TSDN DIT3 responds strongly to both small targets and to looming stimuli (21). In

the locust, LGMD/DCMD neurons respond to both targets and to looming stimuli (16), and

some central complex looming sensitive neurons also respond to small moving targets (20).

Similarly, in *Drosophila,* some optic lobe and descending looming sensitive neurons also

respond to smaller objects (e.g. 17, 35, 41, 42).

However, as opposed to these examples (e.g. 16, 20, 21, 35), we show that the dual

sensitivity to small targets and to larger objects comes from two discrete receptive fields (Fig.

2 – 4, S1 – S3). It is currently unknown if the dual sensitivity described in other insects (e.g.

16, 20, 21, 35) also comes from different receptive fields. Interestingly, *Drosophila* Foma-1

responds to small flickering targets (10° diameter) in addition to looming stimuli (17). While

the visual field location of the looming sensitivity was not specified, target sensitivity was

specific to the dorsal visual field (17), similar to our data (blue, Fig. 2B, C).

Our recordings were done extracellularly (Fig. 1), meaning that neurons with no spontaneous

activity are difficult to discover without presenting a suitable stimulus. We used a small

moving target to initially identify visual neurons (left, Fig. 1), thus biasing our results

towards those looming sensitive descending neurons that also responded to small targets.

However, it is likely that there are looming neurons that do not respond to small objects, such

as found in e.g. *Drosophila* (e.g. 42, 43) and crabs (see e.g. 44). Additionally, our visual

monitor was placed in front of the animal, thus biasing our results to neurons with frontal

sensitivity. It is likely that there are additional looming sensitive descending neurons with

dorsal receptive fields, which could be useful for e.g. detecting predators approaching from

above, or lateral receptive fields, which could be useful for avoiding imminent collision. For

example, in the crab there are 16 retinotopically arranged looming sensitive MLGs that

underlie directional escape behaviours (45). While each receptive field is small, together the

16 neurons cover 360° of the visual field (45), and are thus able to encode directional escape

responses.

Neuronal mechanisms

We showed that the looming sensitive descending neurons likely receive distinct input to the

two receptive fields (Fig. 5-7). Indeed, the dorsal target receptive field is likely to use pre-

synaptic 1-point correlators (blue, Fig. 7), just like the TSDNs do (38). Furthermore, the size

tuning of the dorsal target receptive field (blue, Fig. 5, S4-5) is similar to the size tuning of

TSDNs (38, 46), and of the presumably pre-synaptic STMDs (47). This suggests that the

dorsal target receptive field could share input with the TSDNs.

In contrast, the ventral grating receptive field responded better to larger bars than to small

targets (red, Fig. 5, S4-5), similar to optic flow sensitive descending neurons (38). In

addition, the ventral grating receptive field is likely to use pre-synaptic 2-point correlators of

the EMD-type (red, Fig. 7), similar to optic flow sensitive neurons (48). Interestingly, the

looming sensitive LPLC2 neurons, which are pre-synaptic to the *Drosophila* giant fiber (43),

get their input from T4/T5 (42), which is consistent with a 2-point, EMD-type, correlator

input (see e.g. 49). As our leading-edge data suggests that looming sensitivity is associated

with the grating receptive field (Fig. 4, S3), this indicates that looming sensitivity could be

generated by 2-point correlation. Indeed, in the housefly, escapes can be triggered by

widefield gratings, even if not as efficiently as by looming stimuli (50).

We found that the directionality of the grating receptive field was strongly correlated with the

azimuthal location of the receptive field centre (Fig. 3B, D). However, the directionality of

the target receptive field was much less dependent on its visual field location (Fig. 3A, C). In

addition, we found no correlation between the direction tuning of the two receptive fields

(Fig. 3E). We also found that when both receptive fields were stimulated, the response could

not be explained by a linear summation of the responses to the two stimuli on their own (Fig.

6B), again highlighting that the two receptive fields likely receive independent input.

Similarly, the locust LGMD (51) displays non-linear interactions when stimuli are placed in

different parts of the visual field, suggesting that the details of its receptive fields may be

worth investigating in the future.

Behavioural role

Previous work has shown that the same stimulus displayed in different parts of the visual field can elicit different behavioural output. For example, when crabs living in mudflats see a small dummy moved at ground level they initiate prey pursuit behaviour, but when the same dummy is moved above the crab, they try to escape it (52). In flying *Drosophila*, a looming stimulus in the lateral visual field leads to an escape response, whereas a looming stimulus in the frontal visual field leads to landing behaviour (53). While we did not stimulate the lateral visual field in our set-ups, the strong responses to frontal looming stimuli (right, Fig. 1), likely associated with the ventral receptive field (Fig. 4), suggests that this could be used during landing behaviours on e.g. flowers. Indeed, bees adjust their body angle when landing so the landing surface ends up in the ventral visual field (54).

However, the neurons that we described here could be even more useful in pursuit behaviours. Indeed, when a hoverfly is pursuing a target, during most of the pursuit it will be projected as a small object on the pursuer's eye (55). If the hoverfly were below the target during this part of the pursuit, having a dorsal target receptive field would be appropriate (blue, Fig. 2A-C). This part of the pursuit could thus be supported by either the neurons described here, or by TSDNs without looming sensitivity (9, 38, 46).

During later stages of the pursuit, when the hoverfly gets closer to the target, this will be seen as a looming object (7). Indeed, it has been suggested that this part of the pursuit cannot be subserved by classic target tuned neurons, but instead requires neurons that respond to larger objects and looming stimuli (see e.g. Discussion in Ref. 11), like in zebrafish larvae (12). Furthermore, during the final stages before capture, the pursuer would need to orient itself to grab the target with its legs. During this stage the target would be seen as a large object in the ventral visual field, which would make the more ventral receptive field useful (red, Fig. 2A-

C). In *Drosophila*, looming sensitive Foma-1 is not only involved in escape responses if the fly is stationary (17), but also in courtship, where its silencing results in male flies not being able to follow females (27). Future work investigating where the neurons described here project to, and which behaviours they are thus likely to control, will help elucidate this.

Acknowledgements

We thank past and current lab members for constructive feedback, and the Botanic Gardens of Adelaide for their ongoing support. This research was funded by the US Air Force Office of Scientific Research (AFOSR, FA9550-19-1-0294), the Australian Research Council (ARC, DP180100144, DP210100740 and FT180100289).

References

1. R. M. Olberg, R. C. Seaman, M. I. Coats, A. F. Henry, Eye movements and target fixation during dragonfly prey-interception flights. *J Comp Physiol A* **193**, 685-693 (2007).

2. S. T. Fabian, M. E. Sumner, T. J. Wardill, S. Rossoni, P. T. Gonzalez-Bellido, Interception by two predatory fly species is explained by a proportional navigation feedback controller. *Journal of The Royal Society Interface* **15** (2018).

3. V. Nityananda *et al.*, Insect stereopsis demonstrated using a 3D insect cinema. *Sci Rep* **6**, 18718 (2016).

4. S. M. Fitzpatrick, W. G. Wellington, Contrasts in the territorial behavior of three species of hover flies (Diptera: Syrphidae). *Can Entomol* **115**, 559-566 (1983).

5. J. Zeil, The territorial flight of male houseflies (*Fannia canicularis* L.). *Behav Ecol Sociobiol* **19**, 312-219 (1986).

6. N. Boeddeker, R. Kern, M. Egelhaaf, Chasing a dummy target: smooth pursuit and velocity control in male blowflies. *Proc Biol Sci* **270**, 393-399 (2003).

7. M. Thyselius *et al.*, Male hoverflies (*Eristalis tenax*) do not use heuristic rules based on retinal size and speed during indoor pursuits of artificial targets. *bioRxiv* 10.1101/2022.07.27.501787, 2022.2007.2027.501787 (2022).

8. L. Varennes, H. G. Krapp, S. Viollet, Two pursuit strategies for a single sensorimotor control task in blowfly. *Sci Rep* **10**, 20762 (2020).

9. S. Nicholas, R. Leibbrandt, K. Nordström, Visual motion sensitivity in descending neurons in the hoverfly. *J Comp Physiol A* **206**, 149-163 (2020).

10. K. Nordström, P. D. Barnett, D. C. O'Carroll, Insect detection of small targets moving in visual clutter. *PLoS Biol* **4**, 378-386 (2006).

468 11. Z. M. Bagheri, S. D. Wiederman, B. S. Cazzolato, S. Grainger, D. C. O'Carroll,

469   Properties of neuronal facilitation that improve target tracking in natural pursuit

470   simulations. *J Roy Soc Interface* **12**, 20150083 (2015).

471 12. P. M. Henriques, N. Rahman, S. E. Jackson, I. H. Bianco, Nucleus Isthmi Is Required

472   to Sustain Target Pursuit during Visually Guided Prey-Catching. *Current Biology* **29**,

473   1771-1786.e1775 (2019).

474 13. Y. J. Liu, Q. Wang, B. Li, Neuronal responses to looming objects in the superior

475   colliculus of the cat. *Brain Behav Evol* **77**, 193-205 (2011).

476 14. H. Nakagawa, K. Hongjian, Collision-sensitive neurons in the optic tectum of the

477   bullfrog, Rana catesbeiana. *J Neurophysiol* **104**, 2487-2499 (2010).

478 15. I. Temizer, J. C. Donovan, H. Baier, J. L. Semmelhack, A visual pathway for

479   looming-evoked escape in larval zebrafish. *Curr Biol* **25**, 1823-1834 (2015).

480 16. R. D. Santer, F. C. Rind, P. J. Simmons, Predator versus prey: locust looming-

481   detector neuron and behavioural responses to stimuli representing attacking bird

482   predators. *PLoS One* **7**, e50146 (2012).

483 17. S. E. de Vries, T. R. Clandinin, Loom-sensitive neurons link computation to action in

484   the *Drosophila* visual system. *Curr Biol* **22**, 353-362 (2012).

485 18. H. Fotowat, A. Fayyazuddin, H. J. Bellen, F. Gabbiani, A novel neuronal pathway for

486   visually guided escape in *Drosophila melanogaster*. *J Neurophysiol* **102**, 875-885

487   (2009).

488 19. F. C. Rind, P. J. Simmons, Orthopteran DCMD neuron: A re-evaluation of responses

489   to moving objects. I. Selective responses to approaching objects. *J Neurophysiol* **68**,

490   1654-1666 (1992).

491 20. R. Rosner, U. Homberg, Widespread sensitivity to looming stimuli and small moving

492   objects in the central complex of an insect brain. *J Neurosci* **33**, 8122-8133 (2013).

21.   P. T. Gonzalez-Bellido, H. Peng, J. Yang, A. P. Georgopoulos, R. M. Olberg, Eight

pairs of descending neurons in the dragonfly give wing motor centres accurate

population vector of prey direction. *Proc Natl Acad Sci USA* **110**, 676-701 (2013).

22.   M. A. Frye, R. M. Olberg, Visual receptive field properties of feature detecting

neurons in the dragonfly. *J Comp Physiol A* **177**, 569-576 (1995).

23.   C. R. von Reyn *et al.*, Feature Integration Drives Probabilistic Behavior in the

Drosophila Escape Response. *Neuron* **94**, 1190-1204.e1196 (2017).

24.   D. S. Mearns, J. C. Donovan, A. M. Fernandes, J. L. Semmelhack, H. Baier,

Deconstructing hunting behavior reveals a tightly coupled stimulus-response loop.

Curr Biol **30**, 54-69.e59 (2020).

25.   B. W. Patterson, A. O. Abraham, M. A. MacIver, D. L. McLean, Visually guided

gradation of prey capture movements in larval zebrafish. *J Exp Biol* **216**, 3071-3083

(2013).

26.   D. Förster *et al.*, Retinotectal circuitry of larval zebrafish is adapted to detection and

pursuit of prey. *eLife* **9**, e58596 (2020).

27.   P. Coen, M. Xie, J. Clemens, M. Murthy, Sensorimotor transformations underlying

variability in song intensity during *Drosophila* courtship. *Neuron* **89**, 629-644 (2016).

28.   R. D. Santer, Y. Yamawaki, F. C. Rind, P. J. Simmons, Preparing for escape: an

examination of the role of the DCMD neuron in locust escape jumps. *J Comp Physiol*

A Neuroethol Sens Neural Behav Physiol **194**, 69-77 (2008).

29.   S. Nicholas, M. Thyselius, M. Holden, K. Nordström, Rearing and long-term

maintenance of *Eristalis tenax* hoverflies for research studies. *JoVE*

doi:10.3791/57711, e57711 (2018).

30.   D. H. Brainard, The Psychophysics toolbox. *Spatial Vision* **10**, 433-436 (1997).

31.   D. G. Pelli, The VideoToolbox software for visual psychophysics: Transforming

518   numbers into movies. *Spatial Vision* **10**, 437-442 (1997).

519   32.   H. Fotowat, F. Gabbiani, Relationship between the phases of sensory and motor

520   activity during a looming-evoked multistage escape behavior. *J Neurosci* **27**, 10047-

521   10059 (2007).

522   33.   K. Nordström, I. M. Moyer de Miguel, D. C. O'Carroll, Rapid contrast gain reduction

523   following motion adaptation. *J Exp Biol* **214**, 4000-4009 (2011).

524   34.   K. Nordström, D. C. O'Carroll, The motion after-effect: Local and global

525   contributions to contrast sensitivity. *Proc R Soc Lond B* **276**, 1545-1554 (2009).

526   35.   J. M. Ache, S. Namiki, A. Lee, K. Branson, G. M. Card, State-dependent decoupling

527   of sensory and motor circuits underlies behavioral flexibility in *Drosophila*. *Nat*

528   *Neurosci* **22**, 1132-1139 (2019).

529   36.   S. D. Wiederman, D. C. O'Carroll, Selective attention in an insect visual neuron. *Curr*

530   *Biol* **23**, 156-161 (2013).

531   37.   S. D. Wiederman, P. A. Shoemaker, D. C. O'Carroll, Correlation between OFF and

532   ON Channels Underlies Dark Target Selectivity in an Insect Visual System. *J*

533   *Neurosci* **33**, 13225-13232 (2013).

534   38.   S. Nicholas, K. Nordström, Facilitation of neural responses to targets moving against

535   optic flow. *Proc Natl Acad Sci U S A* **118** (2021).

536   39.   S. D. Wiederman, P. A. Shoemaker, D. C. O'Carroll, A model for the detection of

537   moving targets in visual clutter inspired by insect physiology. *PLoS ONE* **3**, e2784

538   (2008).

539   40.   B. Hassenstein, W. Reichardt, Systemtheoretische Analyse der Zeit, Reihenfolgen

540   und Vorzeichenauswertung Bei der Bewegungsperzeption des Rüsselkafers

541   *Chlorophanus*. *Z Naturforsch* **11**, 513-524 (1956).

41.   S. Namiki, M. H. Dickinson, A. M. Wong, W. Korff, G. M. Card, The functional

organization of descending sensory-motor pathways in *Drosophila*. *eLife* **7**, e34272

(2018).

42.   N. C. Klapoetke *et al.*, Ultra-selective looming detection from radial motion

opponency. *Nature* **551**, 237-241 (2017).

43.   J. M. Ache *et al.*, Neural basis for looming size and velocity encoding in the

Drosophila Giant Fiber escape pathway. *Curr Biol* **29**, 1073-1081.e1074 (2019).

44.   A. Cámera, M. A. Belluscio, D. Tomsic, Multielectrode Recordings From Identified

Neurons Involved in Visually Elicited Escape Behavior. *Front Behav Neurosci* **14**,

592309 (2020).

45.   V. Medan, M. Beron De Astrada, F. Scarano, D. Tomsic, A network of visual motion-

sensitive neurons for computing object position in an arthropod. *J Neurosci* **35**, 6654-

6666 (2015).

46.   S. Nicholas, J. Supple, R. Leibbrandt, P. T. Gonzalez-Bellido, K. Nordström,

Integration of small- and wide-field visual features in Target-Selective Descending

Neurons of both predatory and non-predatory dipterans. *J Neurosci* **38**, 10725-10733

(2018).

47.   K. Nordström, Neural specializations for small target detection in insects. *Curr Opin*

Neurobiol **22**, 272-278 (2012).

48.   R. A. Harris, D. C. O'Carroll, S. B. Laughlin, Adaptation and the temporal delay filter

of fly motion detectors. *Vision Res* **39**, 2603-2613. (1999).

49.   E. Salazar-Gatzimas *et al.*, Direct Measurement of Correlation Responses in

Drosophila Elementary Motion Detectors Reveals Fast Timescale Tuning. *Neuron* **92**,

227-239 (2016).

50.   M. H. Holmqvist, M. V. Srinivasan, A visually evoked escape response of the

567   housefly. *J Comp Physiol A* **169**, 451-459 (1991).

568   51.   H. G. Krapp, F. Gabbiani, Spatial Distribution of Inputs and Local Receptive Field

569   Properties of a Wide-Field, Looming Sensitive Neuron. *Journal of Neurophysiology*

570   **93**, 2240-2253 (2005).

571   52.   D. Tomsic, J. Sztarker, M. Berón de Astrada, D. Oliva, E. Lanza, The predator and

572   prey behaviors of crabs: from ecology to neural adaptations. *Journal of Experimental*

573   *Biology* **220**, 2318-2327 (2017).

574   53.   L. F. Tammero, M. H. Dickinson, Collision-avoidance and landing responses are

575   mediated by separate pathways in the fruit fly, *Drosophila melanogaster*. *J Exp Biol*

576   **205**, 2785-2798 (2002).

577   54.   C. Evangelista, P. Kraft, M. Dacke, J. Reinhard, M. V. Srinivasan, The moment

578   before touchdown: landing manoeuvres of the honeybee Apis mellifera. *Journal of*

579   *Experimental Biology* **213**, 262-270 (2010).

580   55.   P. T. Gonzalez-Bellido, S. T. Fabian, K. Nordström, Target detection in insects:

581   optical, neural and behavioral optimizations. *Curr Opin Neurobiol* **41**, 122-128

582   (2016).

Figure Legends

Figure 1. Looming sensitive descending neurons respond robustly to looming stimuli and to small targets. **A)** Pictograms of the 3° × 3° square target moving horizontally (*Left*) and the looming stimulus with an l/|v| of 10 ms and a final size of 117° (*Right*), as projected on the frontal visual monitor. **B)** Raw data trace from an extracellular recording of a looming sensitive descending neuron in response to a small target (*Left*) or a looming stimulus (*Right*). **C)** The position of the target on the visual monitor (*Left*) and the diameter of the looming stimulus (*Right*), time aligned with the data in panel B. **D)** Spike histograms of the responses in panel B using 20 ms bins.

Figure 2. Two different receptive fields. **A)** The location of the two receptive fields of an example neuron as projected onto the visual monitor. The outlines show the 50% response and the small circle the centre of the target receptive field (blue) and the grating receptive field (red). Euclidean distance between the receptive field centres (black line and value) and the distance of each receptive field centre to the equator and the visual midline (coloured lines and values) are indicated. Bottom right pictogram indicates the preferred direction of the target (blue arrow) and the grating receptive field (red arrow), and the difference between the two (black arc and value). **B)** Location of target (blue) and grating receptive field centres (red) across 102 neurons. **C)** Vertical distance between target (blue) and grating receptive field centres (red) and the visual equator (10° bins). **D)** Horizontal distance from the visual midline (10° bins). **E)** Euclidean distance between each neuron's two receptive field centres (10° bins). Grey data come from neurons where the target receptive field was dorsal to the grating receptive field centre (N = 90) and black data from neurons where the grating receptive field was dorsal to the target receptive field centre (N = 12).

Figure 3. The preferred direction depends on the receptive field centre location. **A)** The

target receptive field centres, colour coded according to their preferred direction (pictogram

bottom right). **B)** The grating receptive field centres, colour coded according to their

preferred direction. **C)** Preferred direction of target receptive fields with centres in either the

left or the right visual field (30° bins). **D)** Preferred direction of grating receptive fields

centres in either the left or the right visual field (30° bins). **E)** Preferred direction difference

between the target and grating receptive field of each neuron (10° bins). Grey data show

neurons where the two receptive field centres were on the same side of the visual midline

(N = 93), whereas black data show neurons with receptive fields on opposite sides of the

visual midline (N = 9).

Figure 4. The leading-edge receptive field is closer to the grating receptive field. **A)** The

location of the target (blue), grating (red) and leading-edge (cyan) receptive field centres in

15 neurons. **B)** The receptive field centres for one example neuron (*Left*), with distances

(black lines) between the leading-edge and the target receptive field centre (*d1*), or the

grating receptive field centre (*d2*), used to calculate the proximity index (*Right*). **C)** Leading

edge proximity index across neurons (N = 15). The leading-edge receptive field was closer to

the grating receptive field centre (red) in more neurons (N = 12) than to the target receptive

field (blue, N = 3). The distribution was significantly different from 0 (P < 0.01, one sample t

and Wilcoxon signed rank test).

Figure 5. The two receptive fields have different size response functions. **A)** The pictograms

indicate the bar trajectory as it moved horizontally across the screen, subtending either the

target receptive field (*Left*, blue dashed line and arrow) or the grating receptive field (*Right*,

red dashed line and arrow, example bar height is 84°). Typical target and grating receptive

fields for an example neuron are shown. The grey shading shows the analysis window used to

calculate the mean response rate, which is the same for both trajectories. The graph shows

that responses to small bars are significantly stronger when passing through the target

receptive field (blue) compared to the grating receptive field (red, mean ± SEM, N = 8). **B)**

The pictogram indicates the analysis windows used to calculate the response to a bar of

varying width as it moved vertically along the screen (trajectory in black) subtending the

target receptive field (blue) or the grating receptive field (red). The graph shows that

responses to narrow bars are significantly stronger within the target analysis window (AW,

blue), while responses to wider bars are significantly stronger within the grating analysis

window (red, mean ± SEM, N = 10). Statistical test was a two-way ANOVA followed by

Sidak's multiple comparisons, with ****$P < 0.0001$, ***$P < 0.001$, **$P < 0.01$ and *$P <$

0.05.

Figure 6. Response to simultaneous stimulation of the two receptive fields. **A)** Responses to

a small target traversing the target receptive field (blue), or bars of varying heights traversing

the grating receptive field (red). The pictograms at the top show the screen position of each

trajectory in relation to the receptive fields for an example neuron. Grey shading indicated the

analysis window used to calculate the mean response rate. **B)** Pictogram showing the screen

position of simultaneously presented target and bar traversing the target and grating receptive

fields. The graph shows that the responses to simultaneous presentation (black) are

significantly lower than the sum of the responses to the same stimuli presented on their own

(purple). **C)** The responses to simultaneous stimuli (black) are not significantly different from

the strongest response evoked by either the target or the bar on its own (green). For all panels

the data show mean ± SEM, for the same N = 5. Statistical analysis was done using two-way

ANOVA, with ****$P < 0.0001$, **$P < 0.01$ and ns indicating $P > 0.05$.

Figure 7. These looming neurons get input from both 1- and 2-point correlators. The

response to a leading OFF edge, a trailing ON edge, or a complete bar, all with a height of

84°, traversing the grating receptive field (red, N = 9), a height of 3° traversing the target

receptive field (blue, N = 10), or the full height of the screen (black, N = 7). The stimuli

moved horizontally at a velocity of 900 pixels/s. In all cases the response from each neuron

was normalized to the sum of the response to all three stimuli from the same trajectory (i.e.

OFF edge only, ON edge only, or complete bar). The inset shows the predicted response of a

motion detector that compares luminance changes over one point (also referred to as an

elementary STMD) or two points in space (often referred to as an EMD). The inset

pictograms are replotted from (37).

A

N = 15

80 degrees

Equator

Midline

- ○ Leading edge centre
- ○ Target RF centre
- ○ Grating RF centre

B

Midline

N = 1

Equator

80 degrees

d1

d2

Proximity index
leading edge

$$= \frac{d1 - d2}{d1 + d2} \times 100\%$$

C

Target RF closer   Grating RF closer

Number of neurons

Proximity index leading edge (%)

A

Target RF trajectory   Grating RF trajectory

B

Target AW

Grating AW

A

Target RF trajectory   Grating RF trajectory

Target only   Bar height (degrees)

Target RF trajectory
Grating RF trajectory

B

Simultaneous

****

**

Bar height (degrees)

Sum of responses
Simultaneous

C

**

ns

Bar height (degrees)

Biggest response
Simultaneous

Supplementary Information

Figure S1. Receptive field mapping method. **A)** The resulting 20 x 20 analysis windows after

scanning the entire monitor along 20 evenly spaced horizontal and vertical target trajectories.

The blue boxes highlight the positions of the analysis windows (P1 – 4) shown in panel C. **B)**

The resulting 8 x 6 grid of overlapping patches of sinusoidal gratings. The red boxes

highlight the grid locations (P1 – 4) shown in panel D. **C)** The responses (open symbols) to 4

directions of target motion within the analysis windows highlighted in panel A. We fit a

cosine function (black) to the responses and used this to extract the local preferred direction

(LPD, filled triangle), and local motion sensitivity (LMS, vertical bar). The dotted line shows

the mean response. **D)** The responses (open symbols) to 8 different directions within the

positions highlighted in panel B, the resulting cosine function (black), the local preferred

direction (LPD, filled triangle), local motion sensitivity (LMS, vertical bar), and mean

response (dotted line). **E)** The target receptive field's position, shown as contour lines (blue

and grey). The centre position (blue circle) was defined as the centroid of the 50% contour

line. **F)** The grating receptive field's position, shown as contour lines (red and grey), and its

centre (red circle). **G)** The arrows show the local motion sensitivity (arrow length) and local

preferred direction (arrow angle) of the target receptive field. Blue arrows indicate the

preferred direction in positions where the local motion sensitivity was greater than 50% of the

maximum for this neuron. Bottom right pictogram indicates the resulting average preferred

direction. **H)** Local motion sensitivity and preferred direction of the grating receptive field.

The bottom right pictogram indicates the average preferred direction (red). The data in this

figure comes from the example neuron shown in Figure *2A*.

Figure S2. Looming sensitive descending neurons are a heterogeneous group of neurons.

A) Target (blue) and grating (red) receptive field of an example neuron. The contour lines

show the 50% response, and the circles the centres of each receptive field. The arrows show the local preferred direction for positions where the response was >50% of the maximum local motion sensitivity. The bottom right pictogram indicates the average preferred direction for the target receptive field (blue) and grating receptive field (red), as well as the difference between the two (black arc and value). The Euclidean distance between the two receptive field centres (black line and value) is indicated on the left. **B)** Example of a looming sensitive neuron with small target and grating receptive fields, and a large distance between the centres. **C)** Example neuron with a large difference between their preferred directions. **D)** Example neuron where the grating receptive field is dorsal to the target receptive field. **E)** Example neuron where the grating receptive field is dorsal to the equator and overlapping with the target receptive field. **F)** Example neuron where the grating and target receptive fields are large and partially overlapping.

Figure S3. Leading edge receptive field method. **A)** Example response from a single neuron (mean +/- SEM) to a full screen OFF edge moving left across the entire width of the screen. Dashed purple line indicates 50% of the maximum response achieved to any direction of leading-edge motion in this neuron. Thresholds (cyan lines) are defined as the time when the neuronal response passes 50% of the maximal response. **B)** Neuronal response to an OFF edge moving right. As the response did not reach 50% maximum, thresholds were not achieved in this direction. **C)** Neuronal response to an OFF edge moving down. **D)** The example neuron showed maximal response (solid purple line) to an OFF edge moving up. **E)** Leading edge receptive field (shaded cyan region), based on screen position of stimulus when thresholds were reached (dashed cyan lines). **F)** Screen position for grating (red), target (blue) and leading edge (cyan) receptive fields, for this example neuron. The data in panels A-D have been smoothed with a 100 ms square-wave filter and is shown at 40 kHz resolution.

Figure S4. Response to small bars is stronger in the dorsal, target receptive field. Response

from a single example neuron to a bar of varying height traversing the entire width of the

screen (mean ± SEM, n = 4) across either the target receptive field (blue) or the grating

receptive field (red). The 50% contour lines and centres for the target receptive field (blue)

and grating receptive field (red) for this neuron are shown. Grey shading indicates the

analysis window used to calculate the mean response for this neuron in Figure 5A. Bar

heights for each panel (A) 0.2°, (B) 0.8°, (C) 1.6°, (D) 3.1°, (E) 5.7°, (F) 11.2°, (G) 21.2°, (H)

39.7°, (I) 69.7°, (J) 106.5°, (K) 137.6°. The data have been smoothed with a 100 ms square-

wave filter, and is shown at 40 kHz resolution.

Figure S5. Sensitivity to bar width depends on which receptive field it is in. Response from

an example neuron to a bar of varying width traversing the entire height of the screen

(mean ± SEM, n = 4). Pictograms indicate the position of the bar trajectory as it moved

vertically up the screen (black dashed line and arrow), traversing first the grating receptive

field (red) and then the target receptive field (blue). Shading indicates the two analysis

windows used to calculate the mean neuronal response for this neuron in Figure 5B. Bar

width for each panel (A) 0.2°, (B) 0.8°, (C) 1.6°, (D) 3.1°, (E) 5.7°, (F) 11.2°, (G) 21.2°, (H)

39.7°, (I) 69.7°, (J) 106.5°, (K) 137.6°, (L) 155.4°. The data have been smoothed with a 100

ms square-wave filter, and is shown at 40 kHz resolution.