Mating activates neuroendocrine pathways signaling hunger in *Drosophila* females

Meghan Laturney^, Gabriella R. Sterne, and Kristin Scott^,#

University of California, Berkeley, United States;

^Corresponding authors: meghan.laturney@berkeley.edu, kscott@berkeley.edu

Lead contact

Summary

Mated females reallocate resources to offspring production, causing changes in nutritional requirements and challenges to energy homeostasis. Although observed in most species, the neural and endocrine mechanisms that regulate the nutritional needs of mated females are not well understood. Here, we investigate the neural circuitry that regulates sugar appetite in mated *Drosophila melanogaster* females. During copulation, a male-derived sex peptide is transferred to females, silencing the mating status circuit to elicit many postmating behavioral changes[1-3]. We find that increased sucrose consumption is a postmated female behavior and show that it is mediated by the mating status circuit. We discovered that sexually dimorphic insulin receptor (Lgr3) neurons integrate mating status and nutritional state signals to adjust sucrose consumption. Lgr3+ cells receive inhibitory input from the mating status circuit via female specific pCd-2 neurons. In mated females, the inhibition of Lgr3 cells from pCd-2 is attenuated, transforming the mated signal into a long-term hunger signal that promotes sugar intake. Our results thus demonstrate that the mating circuit alters nutrient sensing centers in

females to promote sugar consumption, providing a mechanism to increase intake in

anticipation of the energetic costs associated with reproduction.

Main

Female mating status impacts feeding decisions across species. In *Drosophila*, mated

females increase total food intake[4] and develop specific appetites for protein and salt[5-7]. Sugar,

another macronutrient, is vitally important for flies and is their main energy source[8]. In mated

females, increased activity[9] results in elevated metabolic rates[10] driving the need for additional

calories. Moreover, females pack eggs with lipids[11] synthesized from dietary carbohydrates[10].

Together, this predicts that mated females require significantly more sugar than virgins.

However, as excessive sugar negatively impacts fly health[12], sugar intake must be tightly

regulated. Although the sugar to protein ratio influences reproductive output and longevity[8], the

absolute levels of sugar intake of mated females remain untested.

In *Drosophila*, mated related changes in females are orchestrated by sex peptide (SP), a

36-amino acid peptide that is produced in the male seminal fluid and transferred to females

during copulation[1,2]. SP binds to its receptor (SPR)[13] which is expressed in sensory neurons

(SPSNs) in the uterus[14,15]. SPSNs convey mating status to sex peptide abdominal ganglia (SAG)

neurons that ascend to the central brain[3] and synapse onto pC1 neurons[16]. This female-specific

SPSN-SAG-pC1 circuit is active in virgin females and silenced after mating[3,16]. Reception of SP

and silencing of this circuit initiates the long-term postmating response in females, which

includes changes in egg laying and sexual receptivity[3,16,17]. The second-order SAG neurons also

drive changes in salt and protein appetites[7], demonstrating that the mating status circuit alters

food consumption in mated females. However, how the mating circuit influences nutritional state

circuits or feeding circuits remains a central question. Here, we use automated behavioral assays

for measuring consumption, powerful genetic tools, connectomics, and functional imaging

approaches to examine neural mechanisms for appetite changes in mated females.

To investigate the impact of female mating status on sugar intake, we monitored

individual feeding bouts over time using a high-throughput, automated feeding platform (FLIC[18],

Fig. 1a). We compared consumption of virgins and two types of mated females (1-hr or 72-hr

postmated). We found that 72-hr postmated females consumed for a significantly longer duration

on sucrose solution than virgins or 1-hr postmated females (Fig. 1b). To further evaluate the

onset of increased sucrose consumption, we monitored consumption of virgin, 24-hr postmated

females, and 72-hr psotmated females and established a postmated phenotype in both mated

groups (Fig. 1c). These studies demonstrate that mated females consume more sugar than virgins

and that changes in sucrose consumption manifest 6-24hr after copulation.

To explore differences in feeding dynamics, virgin and mated females were allowed to

feed on different sucrose concentrations and the number and length of feeding bouts were

examined. Mated females consumed for longer durations than virgins for all sucrose

concentrations below 500mM (Fig. 1d). Investigation into the microstructure of feeding revealed

that the postmated increase in sucrose consumption is due to an elongation in feeding bout

duration rather than an increased number of bouts (Fig. 1e,f). Thus, like hungry flies[19], mated

females consume more by engaging in longer feeding times rather than by initiating more

feeding events.

We tested two predictions suggested by the mating-induced changes in sugar feeding

dynamics. First, longer overall feeding durations suggest that a greater amount of sugar is

consumed. We directly tested this using a capillary feeder (CAFE[20], Fig. 1g) and found that

mated females consume significantly more sucrose than virgins (Fig. 1h). Second, virgin and

mated females execute a similar number of feeding bouts, suggesting that females do not differ

in the probability of initiating a feeding event. To directly test this, we monitored proboscis

extension upon sugar taste detection (Fig. 1i) and found that virgin and mated females have

similar feeding initiation propensities (Fig. 1j,k). These results further corroborate that increased

sugar consumption in mated females is due to longer meal duration rather than more frequent

feeding bouts.

Hunger increases the sensitivity of gustatory neurons to sugar[21]. To test whether changes

in sugar sensory sensitivity underlie increased sucrose consumption in mated females, we

monitored taste-induced neural activity in sugar gustatory neurons in mated and virgin females

with the GCaMP6s calcium indicator (Fig. 1l). Stimulation of the fly proboscis with sugar

elicited similar responses in gustatory axons regardless of mating status (Fig. 1m-o),

demonstrating that gustatory sensitivity is not altered by mating and not likely to contribute to

changes in feeding.

As mating increases egg production, we reasoned that the increase in sucrose

consumption could be driven by a need-dependent mechanism. In this scenario, mating induces

egg production, depleting energy stores and driving a need for sucrose consumption to restore

homeostasis. To test this, we quantified sucrose consumption of virgin and mated eggless

females with both FLIC and CAFE (Fig. 2a and Extended Data Fig.1a, respectively). We found

that despite a lack of egg production[7,22,23], mated females consumed more sucrose than virgins,

demonstrating that the change in postmated sucrose feeding is not driven by egg production

itself. Instead, this result suggests that sucrose feeding changes are anticipatory in nature and a

consequence of a need-independent mechanism.

Postmated changes in female behavior are elicited by sex peptide (SP)[1,2,24], a male-

derived seminal fluid peptide and require female uterine primary sensory neurons (SPSNs)[13-15]

(Fig. 2b). To test the involvement of the SPSNs in postmated sucrose feeding changes, we

genetically accessed SPSNs using three different genetic driver combinations (see methods). All

genetic intersections consistently labeled at least 4 SPSNs (Fig. 2c, Extended Data Fig.1b,c) with

no overlap of off-target neurons (Extended Data 1d), making them useful tools to investigate the

role of the SPSNs in postmating feeding changes. We silenced the SPSNs in virgin females using

constitutive silencers (either the potassium inward rectifier, KIR2.1, or the tetanus toxin light

chain, TNT) to mimic the mated state and measured sucrose consumption. Virgin females with

SPSNs silenced consumed significantly more sucrose than virgin controls (Fig. 2d, Extended

data 1e,f), recapitulating the mated phenotype. This data argues that mated females increase

sucrose consumption after mating via the SPSNs.

The canonical sex peptide pathway for postmated changes in female behavior is a three-

layer circuit from SPSNs to SAGs to pC1[3,16,17]. As with the SPSNs, SAG and pC1 neurons are

active in a virgin female and silenced in a mated female[3,16]. To precisely manipulate SAGs, we

generated a new split-GAL4 line (SAG-SS3, Fig. 2m). Next, we acutely silenced SAGs by

driving the expression of the green-light gated anion channelrhodopsin (GtACR1) with SAG-

SS3. Monitoring consumption over time, we found that acutely silencing SAGs significantly

increased sucrose consumption (Fig. 2n, Extended Data Fig. 2a). Although an off-target

ascending neuron is also labeled in SAG-SS3, it is not responsible for changes in sucrose

consumption (Extended Data Fig. 3), arguing that the increased sucrose consumption observed

with SAG-SS3 is due to the activity of SAG neuron itself. Downstream of SAG, we found that

acutely silencing pC1 also significantly increased sucrose consumption (Fig. 2o,p, Extended

Data Fig. 2b-d), demonstrating that pC1 neurons regulate postmating sucrose feeding. Taken

together, we conclude that the increased sugar consumption after mating is part of the repertoire

of behavioral changes induced after copulation by sex peptide acting on the SPSN-SAG-pC1

mating status circuit.

Postsynaptic to pC1, the mating status circuit splits into separate neural pathways that

mediate different behavioral subprograms, including circuits that elicit changes in sexual

receptivity[17] or egg-laying[16]. We hypothesized that the neural circuit supporting the postmated

increase in sucrose consumption could either follow one of the established circuits or diverge

after pC1. To test if sucrose feeding is executed by previously identified postmating subcircuits,

we optogenetically manipulated the activity of the descending neurons (DNs) for sexual

receptivity (vpoDNs) or egg laying (oviDNs) to mimic the mated state (Fig. 3a) and measured

sucrose consumption. We found no difference in sucrose consumption upon manipulation of

vpoDNs or oviDNs (Fig. 3b,c,f,g, Extended Data Fig. 2e-h). We note that we measured sucrose

intake in immobilized flies for oviDN manipulations, as activating oviDNs caused females to

stop walking (Fig. 3d,e), likely a natural component of egg laying. These studies argue that

neither of the previously identified circuits responsible for a single postmating response, egg

laying and sexual receptivity, mediate postmating sucrose feeding.

To identify novel pathways that mediate changes in mated feeding behavior, we

examined other major outputs of the mating status circuit. We identified postsynaptic partners of

SAG and pC1 using automated analysis[25,26] of a fly brain electron microscopy (EM)

connectome[27,28]. This approach identified three pCd-2 neurons[29,30], pCd-2a, pCd-2b, and pCd-2c

(Fig. 4a, Extended Data Fig. 4a), that are strongly connected to SAG (39 synapses) and pC1 (158

synapses). These neurons are promising candidates to mediate postmating feeding changes

because their processes descend into the prow region of the Subesophageal Zone (SEZ, Fig.

4b,c), a brain region implicated in energy homeostasis and feeding[31]. As pC1 and SAG are

cholinergic[3,16], pCd-2 cells likely receive excitatory signals from these two upstream synaptic

partners and therefore would be predicted to be less active in mated females (Fig. 4d). We

generated split-GAL4 lines to genetically access pCd-2a and pCd-2b (Fig. 4e,f) and found that

both neurons are female-specific (Extended Data Fig. 4b), consistent with a role in female

postmating behavior. Optogenetically silencing either pCd-2a or pCd-2b significantly increased

sucrose consumption, recapitulating the mated state (Fig. 4h,I, Extended Data Fig. 5c-j).

Importantly, silencing pCd-2a or pCd-2b did not influence egg laying (Fig. 4g, Extended Data

Fig. 5a,b), demonstrating separation of circuits for increased sucrose consumption from other

behaviors downstream of pC1.

To examine how pC1 influences sucrose consumption, we used EM analyses and

identified the major downstream targets of pCd-2 as t-shaped cells (711 synapses) having

processes in the SEZ prow region and a neuroendocrine center, the pars intercerebralis, with

terminal arbors similar to insulin producing cells and other median secretory cells[32] (Fig. 5a,

Extended Data Fig. 6). Based on anatomical similarity, these cells are Lgr3+ (leucine-rich repeat

G-protein-coupled receptor[33]) medial bundle neurons (mBNs) (Fig. 5b,c). Lgr3 expression is

sexually dimorphic, found specifically in the central brain and abdominal ganglia of females[34],

consistent with a role in female behavior. Moreover, Lgr3 is a receptor for the *Drosophila*

insulin-like peptide 8 (Dilp8[35-37]), and both Lgr3 and Dilp8 are known to regulate feeding[38].

These findings suggest that the Lgr3+ mBNs may represent the site where mating status circuits

interact with hunger/satiety circuits to regulate feeding.

To test this hypothesis, we first examined whether Lgr3+ mBNs are functionally

connected to pC1, pCd-2a, or pCd-2b by optogenetically activating these populations while

simultaneously measuring calcium responses in Lgr3+ neurons. In all cases, neural activation

caused significant calcium decreases in Lgr3+ neurons (Fig. 5d-f), validating Lgr3+ mBNs as

targets of the mating status pathway. As pC1 is inhibited in the mated state, our studies argue

that Lgr3+ mBNs are more active in mated females (Fig. 5g). To examine whether activity in

Lgr3+ mBNs influences consumption, we targeted this population with a genetic intersectional

approach (R19B09∩fru[34], Fig. 5c), optogenetically activated Lgr3+ mBNs, and monitored

sucrose consumption. We found that increased activity in Lgr3+ mBNs promotes sucrose

consumption, mimicking the mated state (Fig. 5h). Together, these studies reveal that Lgr3+

mBNs receive inputs from the canonical postmating pathway and are part of a neuroendocrine

pathway that regulates sucrose consumption in postmated females.

In conclusion, we report a new postmated response of increased sucrose consumption,

which independent of caloric deficiencies caused by egg production and anticipatory in

nature. Our findings provide a neural mechanism for the coordination of mating state and

sucrose consumption (Fig. 5g). In virgins, increased activity of SAG and pC1 potentiates pCd-2-

mediated inhibition of Lgr3+ cells within the endocrine center of the brain, resulting in low sugar

intake. After mating, the activity of SAG and pC1 is reduced[3,16], decreasing the activity of pCd-2

and alleviating the inhibition from *Lgr3+* cells, increasing sucrose intake.

The sexually dimorphic Lgr3+ neurons are the nexus integrating mating status and

neuroendocrine hunger signaling. Lgr3 is a leucine-rich repeat containing G-protein coupled

receptor, which binds Dilp8 to regulate feeding in flies. Expression levels of Lgr3 and Dilp8 rise

in fed flies and mutations in either gene increase feeding, arguing that Dilp8 and Lgr3 are satiety

factors[38]. Our calcium imaging and behavioral studies of Lgr3+ mBNs indicate that they are

more active in a mated state and promote sucrose consumption. Together, these data suggest a

model for how Lgr3+ mBN neurons integrate signals from the mating status and hunger/satiety

systems. In virgin or satiated female flies, circulating DILP8 and pCd-2 activity, respectively,

inhibit Lgr3+ mBNs (Fig. 5g), reducing sugar feeding. Conversely, in mated or hungry flies,

Lgr3+ mBNs lack the inhibitory signal from pCd-2 neurons or circulating Dilp8, respectively,

resulting in increased sugar consumption. Hence, the mated state may be viewed mechanistically

as an additional "hunger" signal within this system.

Thus, we report that an elevated appetite for sucrose is an important behavioral

subprogram elicited by mating and executed by female-specific circuitry, shifting the physiology

of a mated female. The activation of hunger centers by the mating status circuit provides a neural

mechanism that anticipates the large energetic demands associated with offspring production and

increases caloric intake, promoting reproductive success.

References

1. Chen, P. S., Stumm-Zollinger, E., Aigaki, T., Balmer, J., Bienz, M. & Böhlen, P. A male accessory gland peptide that regulates reproductive behavior of female D. melanogaster. *Cell* **54**, 291-298 (1988).
2. Liu, H. & Kubli, E. Sex-peptide is the molecular basis of the sperm effect in Drosophila melanogaster. *Proc. Natl. Acad. Sci. USA* **100**, 9929-9933 (2003).
3. Feng, K., Palfreyman, M. T., Häsemeyer, M., Talsma, A. & Dickson, B. J. Ascending SAG neurons control sexual receptivity of Drosophila females. *Neuron* **83**, 135-148 (2014).
4. Carvalho, G. B., Kapahi, P., Anderson, D. J. & Benzer, S. Allocrine modulation of feeding behavior by the sex peptide of Drosophila. *Curr. Biol.* **16**, 692-696 (2006).
5. Vargas, M. A., Luo, N., Yamaguchi, A. & Kapahi, P. A role for S6 kinase and serotonin in postmating dietary switch and balance of nutrients in D. melanogaster. *Curr. Biol.* **20**, 1006-1011 (2010).
6. Ribeiro, C. & Dickson, B. J. Sex peptide receptor and neuronal TOR/S6K signaling modulate nutrient balancing in Drosophila. *Curr. Biol.* **20**, 1000-1005 (2010).
7. Walker, S. J., Corrales-Carvajal, V. M. & Ribeiro, C. Postmating circuitry modulates salt taste processing to increase reproductive output in Drosophila. *Curr. Biol.* **25**, 2621-2630 (2015).

8. Simpson, S. J., Le Couteur, D. G. & Raubenheimer, D. Putting the balance back in diet. *Cell* **161**, 18-23 (2015).

9. Isaac, R. E., Li, C., Leedale, A. E. & Shirras, A. D. Drosophila male sex peptide inhibits siesta sleep and promotes locomotor activity in the post-mated female. *Proc. Biol. Sci.* **277**, 65-70 (2010).

10. Brown, E. B., Klok, J. & Keene, A. C. Measuring metabolic rate in single flies during sleep and waking states via indirect calorimetry. *J. Neurosci. Methods* **376**, doi: 10.1016/j.jneumeth.2022.109606 (2022).

11. Sieber, M. H & Spradling, A. C. Steroid signaling establishes a female metabolic state and regulates SREBP to control ooctyte lipid accumulation. *Curr. Biol.* **25**, 993-1004 (2015).

12. Baenas, N. & Wagner, A. E. Drosophila melanogaster as a model organism for obesity and type-2 diabetes mellitus by applying high-sugar and high-fat diets. *Biomolecules* **12**, doi: 10.3390/biom12020307 (2022).

13. Yapici, N., Kim, Y. J., Ribeiro, C. & Dickson, B. J. A receptor that mediates the post-mating switch in Drosophila melanogaster reproductive behavior. *Nature* **451**, 33-37 (2008).

14. Häsemeyer, M., Yapici, N., Heberlein, U. & Dickson, B. J. Sensory neurons in the Drosophila genital tract regulate female reproductive behavior. *Neuron* **61**,511-518 (2009).

15. Yang, C. H., Rumpf, S., Xiang, Y., Gordon, M. D., Song, W., Jan, L. Y. & Jan, Y. N. Control of the postmating behavioral switch in Drosophila females be internal sensory neurons. *Neuron* **61**, 519-526 (2009).

16. Wang, F., Wang, K., Forknall, N., Patrick, C., Yang, T., Parekh, R., Bock, D. & Dickson, B. J. Neural circuitry linking mating and egg laying in Drosophila females. *Nature* **579**, 101-105 (2020).

17. Wang, K., Wang, F., Forknall, N., Yang, T., Patrick, C., Parekh, R. & Dickson, B. J. Neural circuit mechanisms of sexual receptivity in Drosophila females. *Nature* **589**, 577-581 (2021).

18. Ro, J., Harvanek, Z. M. & Pletcher, S. D. FLIC: high-throughput, continuous analysis of feeding behaviors in Drosophila. *PLoS One* **9**, doi: 10.1371/journal.pone.0101107 (2014).

19. Itskov, P., Moreira, J. M., Vinnik, E., Lopes, G., Safarik, S, Dickinson, M. H. & Ribeiro, C. Automated monitoring and quantitative analysis of feeding behaviour in Drosophila. *Nat. Commun.* **5**, doi: 10.1038/ncomms5560 (2014).

20. Ja, W. W., Carvalho, G. B., Mak, E. M., de la Rosa, N. N., Fang, A. Y., Liong, J. C., Brummel, T. & Benzer, S. Prandiology of Drosophila and the CAFE assay. *Proc. Natl. Acad. Sci. USA* **104**, 8253-8256 (2007).

21. Inagaki, H. K., Panse, K. M. & Anderson, D. J. Independent, reciprocal neuromodulatory control of sweet and bitter taste sensitivity during starvation in Drosophila. *Neuron* **84**, 806-820 (2014).

22. Ohlstein, B. & McKearin, D. Ectopic expression of the Drosophila Bam protein eliminates oogenic germline stem cells. *Development* **124**, 3651-3662 (1997).

23. Mével-Ninio, M., Fouilloux, E., Guénal, I & Vincent, A. The three dominant female-sterile mutations of the Drosophila ovo gene are point mutations that create new translation-initiator AUG codons. *Development* **122**, 4131-4138 (1996).

24. Laturney, M. & Billeter, J. C. Neurogenetics of female reproductive behaviors in
Drosophila melanogaster. *Adv. Genet.* **85**, 1-108 (2015).

25. Buhmann, J. et al. Automatic detection of synaptic partners in a whole-brain Drosophila
electron microscopy data set. *Nat. Methods***7**, 771-774 (2021).

26. Heinrich, L., Funke, J., Pape, C., Nunez-Iglesias, J. & Saalfeld, S. Synaptic cleft
segmentation in non-isotropic volume electron microscopy of the complete Drosophila
brain. arXiv:180502718 [csCV]. (2018).

27. Zheng, Z. et al. A complete electron microscopy volume of the brain of adult Drosophila
melanogaster. *Cell***174**, 730-743 (2018).

28. Dorkenwald, S. et al. FlyWire: online community for whole-brain connectomics. *Nat.*
*Methods***1**, 119-128 (2022).

29. Kimura, K., Sato, C., Koganezawa, M. & Yamamoto, D. Drosophila ovipositor extension
in mating behavior and egg deposition involves distinct sets of brain interneurons. *PLoS*
*ONE***10**, e0126445 (2015).

30. Nojima, T., Rings, A., Allen, A. M., Otto, N., Verschut, T. A., Billeter, J. C., Neville, M.
C. & Goodwin, S. F. A sex-specific switch between visual and olfactory inputs underlies
adaptive sex differences in behavior. *Curr. Biol.* **31**, 1175-1192 (2021).

31. Shiu, P. K., Sterne, G. R., Engert, S., Dickson, B. J. & Scott, K. Taste quality and hunger
interactions in a feeding sensorimotor circuit. *Elife***11**, doi: 10.7554/eLife.79887 (2022).

32. Liao, S. & Nässel, D. R. Drosophila insulin-like peptide 8 (DILP8)in ovarian follicle cells
regulates ovulation and metabolism. *Front. Endocrinol.* **11**, doi:
10.3389/fendo.2020.00461 (2020).

33. Dolan. J. et al. The extracellular leucine-rich repeat superfamily; a comparative survey
and analysis of evolutionary relationships and expression patterns. *BMC Genomic***8**, doi:
10.1186/1471-2164-8-320 (2007).

34. Meissner, G. W., Luo, S. D., Dias, B. G., Texada, M. J. & Baker, B. S. Sex-specific
regulation of Lgr3 in Drosophila neurons. *Proc. Natl. Acad. Sci. USA***113**, 1256-1265
(2016).

35. Colombani, J., Andersen, D. S., Boulan, L., Boone, E., Romero, N., Virolle, V., Texada,
M. & Léopold, P. Drosophila Lgr3 couples organ growth with maturation and ensures
developmental stability. *Curr. Biol.* **25**, 2723-2729 (2015).

36. Garelli, A. et al. Dilp8 requires the neuronal relaxin receptor Lgr3 to couple growth to
developmental timing. *Nat. Commun.* **6**, doi: 10.1038/ncomms9732 (2015).

37. Vallejo, D. M., Juarez-Carreño, S., Bolivar, J., Morante, J. & Dominguez, M. A brain
circuit that synchronizes growth and maturation revealed through Dilp8 binding to Lgr3.
*Science***350**, doi: 10.1126/science.aac6767 (2015).

38. Yeom, E. et al. Tumour-derived Dlip8/INSL3 induces cancer anorexia by regulating
feeding neuropeptides via Lgr3/8 in the brain. *Nat. Cell Biol.* **23**, 172-183 (2021).

Fig. 1. Mated females consume more sucrose than virgins by elongating feeding events.
a, Schematic of fly feeding behavior (left) and the FLIC signal that monitors food contact over time (right). **b**, Cumulative drinking time in FLIC of virgin (white circles, n=34), 1-hr postmated (pm) (light pink triangles, n=30), and 72-hr pm (dark pink squares, n=28) females. Line graph shows mean and s.e.m. over time. Kruskal-Wallis was used to compare groups at hour intervals, Dunn's posthoc. **c**, Total drinking time in FLIC in 20-min assay of virgin (v; n=47), 24-hr pm (n=47), and 72-hr pm (n=22) females, Kruskal-Wallis, Dunn's posthoc. **d-f**, Schematic and plots of feeding behavior of virgin (v) and mated (m) females presented with sucrose of varying concentrations in FLIC 20-min assay (n=33-36) examining total drinking time (**d**), number of drinking bouts (**e**), and average bout duration (**f**). Box plot show first to third quartile, whiskers span 10-90 percentile, Mann Whitney. **g**, Schematic of female drinking from capillary containing sucrose solution in CAFE assay. **h**, Total volume consumed in CAFE assay per fly in 24 hours by virgin (v, n=21) and mated females (m, n=21), unpaired t-test. **i**, Schematic of proboscis extension response (PER) assay. **j-k**, Percentage of proboscis extensions observed per fly upon three presentations of sucrose of indicated concentration in virgin (v) and mated (m) females in starved (n=17-18, **j**) and fed state (n=16, **k**), Mann Whitney. **l**, Schematic of in vivo calcium imaging experiment monitoring taste-induced activity in gustatory axons. **m-o**, Changes in GCaMP6s signal in virgin (grey) and mated (dark pink) females, shown as representative fluorescent changes in single brains (**m**), changes over time (**n**), and maximum changes (**o**), n=10. Blue bar represents proboscis sucrose stimulation, Wilcoxen Rank test. *p<0.05, **p<0.01, ***p<0.001, ns = not significant. Scatter plot show mean +/- s.e.m. (**c,h,o**). Column graph show mean +/- s.e.m. (**j,k**). For full genotypes, see Supplementary Data Table 1.

Fig. 2. Postmated increases in sucrose consumption are induced by the canonical mating status circuit and are independent of egg-production.
a, Total drinking time in FLIC in 20-min assay of eggless virgin (v, n=16) and mated (m, n=15) females, Mann Whitney. **b**, Schematic of female with the mating circuit, showing first order (SPSN, pink-orange), second order (SAG, dark yellow) and third order (pC1, green) neurons. **c**, Confocal image of uterus stained to reveal SPSNs (UAS-CD8:GFP, green, arrowhead indicating cell bodies) and muscle tissue (phalloidin, blue). **d**-**e**, Total drinking time in FLIC in 20-min assay of virgin females of indicated genotype (n=17-28), Kruskal-Wallis, Dunn's posthoc. **f,h** Confocal images of brain (top) and ventral nerve cord (bottom) of SAG-SS3 (**f**) and pC1-SS1 (**h**) females, stained to reveal neural projection pattern (UAS-mVenus, green) and all synapses (nc82, blue). **g-i**, Drinking time of females exposed to light normalized to dark controls of indicated genotype in FLIC over 4-hrs (each data point is the drinking time of an individual in the light condition minus the daily average drinking time of all females in dark condition within genotype, n=17-39), Kruskal-Wallis, Dunn's posthoc at hour 4, #p<0.05, ###p<0.001. Line graphs show mean and s.e.m. over time. *p<0.05, **p<0.01, ***p<0.001 (**a**,**d**,**e**,**g**,**i**). Scatter plot show mean +/- s.e.m. (**a**,**d**,**e**). Scale bar 50 μm (**c**,**f**,**h**). For full genotypes see Supplementary Table 1.

Fig. 3. Descending neurons that regulate egg laying and sexual receptivity do not influence sucrose consumption.

a, Schematic of mating circuit with neural outputs of pC1 and resulting postmating response. Green circles indicate active neurons, salmon circles indicate silenced neurons. **b**-**c**, Difference in drinking time of indicated genotype in FLIC over 4 hrs. Each data point represents normalized drinking time of a single female in light condition (drinking time minus the daily average drinking time of all females in dark condition within genotype, n=20-24), Kruskal-Wallis. Line graph shows mean and s.e.m. over time. **d**, Velocity heatmap upon transient activation in individual walking flies. Red bars indicate light ON (n=8-16). **e**, Total distance traveled in locomotor assay (**d**). ON condition (red) includes 3 30-sec light ON conditions. OFF condition (white) includes 3-30sec light OFF conditions immediately preceding each ON condition, Mann Whitney, ***p<0.001. Aligned dot plot show mean +/- s.e.m. **f**-**g**, Drinking time of females exposed to light normalized to dark controls of indicated genotype in FLIC over 4-hrs (n=19-26), 1way ANOVA, Bonferroni's posthoc. ns = not significant. Scatter plot show mean +/- s.e.m. (**e**-**g**), For full genotypes see Supplementary Data Table 1.

Fig. 4. pCd-2 neural outputs of SAG and pC1 mediate postmating sucrose consumption.
a, Schematic of neural connectivity of the mating status circuit and pCd-2 cells. Arrows represent direct connection and number indicates number of synapses. **b**-**c**, Electron microscopy reconstruction of pCd-2a,b neurons. **d**, Model of circuit activity in virgin and mated state. Green circles indicate active neurons, salmon circles indicate silenced neurons. **e**-**f**, Confocal images of pCd-2a-SS1 and pCd-2b-SS1 brains, stained to reveal pCd-2 projection pattern (UAS-GFP, green) and all synapses (nc82, blue). Scale bar 50 μm. **g**, Difference in eggs laid in 24-hr of females exposed to light normalized to dark controls (n=10-20). For "UAS-GtACR1 mated" group, data set represents eggs laid by mated females normalized to virgin females of the same genotype. Scatter plot show mean +/- s.e.m., one sample t-test. **h**-**i**, Drinking time of females exposed to light normalized to dark controls of indicated genotype in FLIC over 4-hrs (n=22-26). Line graph show mean and s.e.m. over time, Kruskal-Wallis, Dunn's posthoc at hour 4, ##$p < 0.01$, ###$p < 0.001$. *$p < 0.05$, **$p < 0.01$, ***$p < 0.001$. For full genotypes see Supplementary Table 1.

Fig. 5. Sexually dimorphic neuroendocrine Lgr3+ cells receive mating status signals via pCd-2 neurons and regulate sucrose consumption in mated females.

a, Downstream neuronal cell types of pCd-2 cells identified by electron microscopy reconstruction, showing the number of pCd-2 output synapses. T-shape cell type represented with turquoise circles, other cell types represented with white circles (a-j). **b**, Electron microscopy reconstruction of t-shape neurons. **c**, Confocal image of fru+ Lgr3-expressing cells in medial brain (UAS-CD8:GFP, green) and all synapses (nc82, blue). Scale bar 50 μm. **d-f**, Calcium responses of Lgr3-expressing cell bodies in the prow region of the brain to activation of upstream pC1 (**d**), pCd-2a (**e**), or pCd-2b (**f**) neurons (n=7-8) and the analysis of area under the curve, Wilcoxon Rank Test, scatter plot show mean +/- s.e.m. (grey bar). **g**, Neural model for the coordination of mating status and sucrose intake. Green circles indicate active neurons, salmon circles indicate silenced neurons. **h**, Drinking time of females exposed to light normalized to dark controls of indicated genotype in FLIC over 4-hrs (n=10-15). Line graph shows mean and s.e.m. over time, Kruskal-Wallis, Dunn's posthoc at hour 4, #p<0.05, ##p<0.01. *p<0.05, **p<0.01, ***p<0.001. For full genotypes see Supplementary Table 1.

a

Volume consumed / fly (μl)

virgin   mated

*

b VT3280 ∩ ppk

SPSNs
Phalloidin

c fru ∩ VT3280

SPSNs
Phalloidin

d

| fru ∩ ppk | VT3280 ∩ ppk | fru ∩ VT3280 |

fru ∩ ppk
nc82

VT3280 ∩ ppk
nc82

fru ∩ VT3280
nc82

fru ∩ ppk
nc82

VT3280 ∩ ppk
nc82

fru ∩ VT3280
nc82

e

Total contact duration (s)

**   *

ppk-gal4   +   +   +
VT3280-LexA   +   +
UAS-TNT   +   +
UAS-KIR

Total contact duration (s)

*   *

+   +   +
     +   +

UAS-TNT   +
UAS-KIR   +   +

f

Total contact duration (s)

***
    **

VT3280-gal4   +   +   +
fru-LexA   +   +
UAS-TNT   +   +
UAS-KIR

Total contact duration (s)

***
     ns

+   +   +
     +   +

             +   +

Extended Data Fig. 1. Three independent SPSN lines regulate postmated sucrose consumption.
a, Total volume of 50mM sucrose consumed per fly in CAFE assay by eggless virgin (n=21) and eggless mated females (n=20). *p<0.05, unpaired t-test. **b-c**, Confocal image of uterus stained to reveal SPSNs (UAS-CD8:GFP, green, arrowhead indicating cell bodies) and muscle tissue (phalloidin, blue). **d**, Confocal images of brain (top) and ventral nerve cord (bottom) of genetic intersections accessing SPSNs stained to reveal off-target neurons in the central nervous system (UAS-GFP; green) and all synapses (nc82; blue). **e-f**, Total drinking time in FLIC in 20-min assay of virgin females of indicated genotype (n=19-41). Kruskal-Wallis with Dunn's Multiple Comparison Test revealed differences between silenced females and controls. Scatter plot show mean +/- s.e.m. (**a**,**e-f**). Scale bar 50 μm (**b-d**).

Extended Data Fig. 2.Contact duration times of virgin and mated females upon optogenetic manipulation of neurons in the mating status circuit.

a-b, Total drinking time in FLIC in 4-hr assay of females of indicated genotype and light condition. White circles are dark condition, green circles are green light condition, n=20-32. Data was analyzed with 2way ANOVA and Bonferroni posttests, ***p<0.001. **c**, Drinking time of females exposed to light normalized to dark controls of indicated genotype in FLIC over 4-hrs, n=20-24. Line graphs show mean and s.e.m. over time, Kruskal-Wallis, Dunn's posthoc at hour 4, ##p<0.01, ns = not significant. **d-f**, Total drinking time in FLIC in 4-hr assay of females of indicated genotype and light condition. White circles are dark condition, green circles are green light condition, n=20-32, 2way ANOVA and Bonferroni posttests. **g-h**, Total drinking time in TCA of females of indicated genotype and light condition. White circles are dark condition, red circles are red light condition, n=20-32, 2way ANOVA and Bonferroni posttests, *p<0.05, **p<0.01. Scatter plot show mean +/- s.e.m. (**a**-**b**,**d**-**h**).

a

EM reconstruction

pCd-2c

b

	female	male		female	male
pCd-2a-SS1	nc82 mVenus	nc82 mVenus	pCd-2b-SS1	nc82 mVenus	nc82 mVenus
pCd-2a-SS2	nc82 mVenus	nc82 mVenus	pCd-2b-SS2	nc82 mVenus	nc82 mVenus
			pCd-2b-SS3	nc82 mVenus	nc82 mVenus

Extended Data Fig. 5. Independent pCd-2a and pCd-2b lines regulate postmated sucrose consumption.

a-b, Number of eggs laid in 24 hours by virgin females of indicated genotype while exposed to green light (green) or no light (white). 2way ANOVA analysis revealed no significant differences between groups (n=9-20). **c-e**, Drinking time of females exposed to light normalized to dark controls of indicated genotype in FLIC over 4-hrs (n=19-36), Kruskal-Wallis, Dunn's posthoc at hour 4, ###$p<0.001$. Line graph shows mean and s.e.m. over time. **f-j** Total drinking time in FLIC in 4-hr assay (n=20-24). Analyzed with a 2way ANOVA and Bonferroni posttests (n=19-36). *$p<0.05$, **$p<0.01$, ***$p<0.001$, ns = not significant. Scatter plot show mean +/- s.e.m. (**a-b,f-j**).

Fly Wire IDs from left to right
720575940613895934
720575940628750718
720575940628422555
720575940616987426
720575940610815510
720575940606550706
720575940620156513
720575940615716753
720575940632607127
720575940623115499
720575940628121143
720575940625126192
720575940632733792
720575940630247784
720575940618115669
720575940625960360
720575940632879842
720575940641886605
720575940634391950
720575940620201084
720575940616044834
720575940613988242
720575940627466746
720575940629491275
720575940620064321
720575940629944407
720575940615979067
720575940613234591
720575940622386056
720575940625850212
720575940613943714

Extended Data Fig. 6. Post-synapatic targets of pCd-2.
Schematic of neural connectivity of pCd-2 cells and downstream t-shape neurons identified by electron microscopy reconstruction. Number indicates number of synapses.

Methods

Rearing conditions and strains: Flies were reared on standard cornmeal-agar-molasses medium, at 25°C, 65% humidity on a 12-hr-light-dark cycle. Flies used in optogenetic assays were reared on food containing 0.25mM all-trans-retinal (Sigma-Aldrich) in darkness, before and after eclosion.

Flies were collected under CO_2 1-8h after eclosion and housed in same-sex groups. To generate mated females, 15 virgin females and 5 males were paired 3-5 days post-eclosion and group housed for 72 hours. To generate recently mated females, a single virgin female was paired with a single male for 2 hours. If mating was observed, the recently mated female was then immediately placed into indicated assay. To generate 24-hr mated females, a single virgin female was paired with a single male for 2 hours. If mating was observed, the recently mated female was then group housed with other recently mated females until testing. To generate egg-less females we used either ovoD mutant females[23] or hsp70-bam females[22] following the heat-shock protocol[7].

Capillary feeding (CAFE) assay: Capillary feeding arenas were modified from the original[20]. Briefly, we equipped a plastic vial, containing a wet kimwipe, with an altered rubber stopper lid housing two truncated 200-ml pipette tips whereby the capillaries can be inserted through. The capillaries (calibrated glass micropipettes, 5 ml) were loaded with 50mM sucrose solution containing 0.25 mg/ml blue dye. To measure sucrose consumption, 5 virgin or mated females were gently aspirated into an arena for 24 hours, the displacement of the meniscus was measured, and the volume per fly was calculated (1mm = 0.038ml).

Fly liquid interaction counter (FLIC): Females of indicated genotype or mating status were first wet-starved for 12-18 hours. Following this, females were then gently aspirated into the fly liquid food interaction counter (FLIC[18]), which was pre-filled with sucrose solution of indicated concentration. The voltage from each well was continuously recorded and a .csv file was produced. Significant changes in the voltage indicated that the fly was contacting the liquid. Contact duration acted as a proxy for consumption time. Based on a video analysis of flies drinking in FLIC in our lab, we modified the parameters in R to better model consumption. The new parameters are as follows: Feeding.Interval.Minimum = 10, Feeding.Threshold.Value = 10, Tasting.Threshold.Interval = c(01,02), Feeding.Minevents = 6, Signal.Threshold = 10.
For optogenetic control of neural activity prior to FLIC, we placed vials containing flies into a custom made light box (20cm x 10cm x 30cm) equipped with a string of 150 LED lights programmed with an Arduino to have green light constantly on.
For optogenetic control of neural activity while flies were in the FLIC, we designed and constructed a 12-chamber lid equipped with either 3 red LEDs (617nm, Luxeon Star, catalog number SP-01-E6) or 3 green LEDs (530nm, Luxeon Star, catalog number SP-01-G4). The 12-chamber cover was made from half of a 24-well plate, which was trimmed to fit. Lights were attached onto half of the 24-well plate lid and fitted atop the chamber cover.

Proboscis extension response (PER) assay: Females of indicated mating status were either wet-starved for 12-18 hours (starved) or kept on standard food (fed). Flies were then anesthetized using CO_2 and fixed to a glass slide with No More Nails polish and then allowed to recover for 2 hours in a humidified box. Immediately before testing, flies were given water until they no

longer responded to 5 consecutive presentations. Flies were then presented 3 times with sucrose of an indicated concentration and proboscis extension was recorded.

Temporal consumption assay (TCA): Flies were starved for 18 hours in darkness, anesthetized using CO_2 and then fixed to a glass slide with No More Nails polish with limited light and allowed to recover for 2 hours in a humidified red light LED box or a humidified dark box. Immediately before testing, flies were given water until they no longer responded to 5 consecutive presentations. In testing, flies were presented with 250 mM sucrose and consumption time was recorded. Each trial ended when the fly was presented with the tastant and no longer responded to 5 consecutive presentations. Flies in the dark condition were given water and tested in low light conditions. Flies in red light condition were given water and tested in the presence of red light.

Egg laying assay: 7-day old females were placed in a vial with standard food and placed into a custom-made light box (20cm x 10cm x 30cm) that was equipped with a string of 150 LED lights programmed with an Arduino. In neural activation tests, red light was transiently on (1 second on/1 second off). In neural silencing tests, green light was constantly on. Half of the females of each genotype were placed in vials wrapped in foil to function as no-light condition controls. After 24 hours, females were removed and eggs were immediately counted.

Locomotor assay: Locomotor assays were conducted as described[39].

In vivo calcium imaging with taste stimulation: Protocol was adapted[31]. Virgin and mated females aged 14-20 days and mated females were co-housed with Canton-S males. Females were food-deprived for 18-24 hours prior to imaging. A window was made in the head of the fly to allow for visualization of the SEZ and a drop of ~250 mOsmo AHL solution (https://www.sciencedirect.com/science/article/pii/S0092867403000047) was added to the head and imaging was immediately performed on a fixed-stage 3i spinning disk confocal microscope with a piezo drive and a 20x water objective (1.6x optical zoom). A 250 mM sucrose solution was delivered to the proboscis using a glass capillary. To analyze these images, a maximum intensity projection encompassing the arbors of Gr64f neurons was made using Fiji. Large ROIs were drawn manually corresponding to the GCaMP expression and a large ROI was drawn in an adjacent region to measure background autofluorescence. Mean fluorescence levels from the background ROI were subtracted from the Gr64f ROIs at each time point, resulting in the fluorescence trace over time: $F(t)$. delta F/F was measured as follows: $(F(t) – F(0)) / F(0)$, where $F(0)$ was the average of 10 time points before stimulation with sucrose and $F(t)$ was the average background subtracted increases in fluorescence during sucrose stimulation. Area under the curve for 10 frames before stimulation and during stimulation was approximated with the trapezoidal rule in Python using the NumPy.trapz function. Statistical analysis was carried out in Python using the SciPy package, version 1.7.3 (https://www.nature.com/articles/s41592-019-0686-2). deltaF/F0 images were created as described[40].

In vivo functional connectivity experiments: In vivo sample preparation for calcium imaging was performed as described[31]. Prior to imaging, female flies were aged for 2 weeks and then food-deprived in a vial containing a distilled water-soaked kimwipe for 18–24 hours prior to imaging. Female flies were housed with Canton-S males throughout aging and food-deprivation.

Following dissection, samples were bathed in ~250 mOsmo AHL solution (https://www.sciencedirect.com/science/article/pii/S0092867403000047) and imaged immediately using a 2-photon microscope. Volumetric images of the prow region were acquired with 920 nm light at 2.9 Hz with resonant scanning and a piezo-driven objective. During imaging, a custom ScanImage plugin was used to deliver two 10-second pulses of 660 nm light through the objective at a 10-second interval to excite CsChrimson.

Image and statistical analysis of functional connectivity experiments was performed using Fiji (https://www.nature.com/articles/nmeth.2019), CircuitCatcher (a customized python program by Daniel Bushey), and Python (Python Software Foundation, https://www.python.org/). Using Fiji, image stacks for each time point were first maximum intensity projected. Then, using CircuitCatcher, an ROI containing Lrg3+ mBN cell bodies and a background ROI of brain tissue were selected and the average fluorescence intensity for each ROI at each timepoint was retrieved. Then, in Python, background subtraction was carried out for each timepoint (F_t) and initial fluorescence intensity ($F_{initial}$) was calculated as the mean corrected average fluorescence intensity for frames 1-15. Then, ΔF/F was calculated using the following formula: F_t-$F_{initial}$/$F_{initial}$. Area under the curve for before (off; frames 1-29) and during (on; frames 30-58) light stimulation was approximated with the trapezoidal rule in Python using the NumPy.trapz function. Statistical analysis of functional connectivity experiments was carried out in Python using the SciPy package, version 1.7.3 (https://www.nature.com/articles/s41592-019-0686-2).

Dissections and Immunohistochemistry: All CNS dissections and immunostaining (unless directly addressed) were performed following the detailed instructions found at https://www.janelia.org/project-team/flylight/protocols. To image split-GAL4 lines and intersections, we used the following primary antibodies: mouse α-Brp (nc82, DSHB, University of Iowa, USA) at 1:40, chicken α-GFP (Invitrogen, A10262) at 1:1000; and the following secondary antibodies: goat α-mouse AF647 (Invitrogen, A21236) at 1:500, goat α-chicken AF488 (Life Technologies, A11039) at 1:500. Multi-Color Flip-Out fly generation was performed following the protocol[41]. For imaging, we used the following primary antibodies: mouse α-Brp (nc82, DSHB, Univerisity of Iowa, USA) at 1:40, rabbit α-HA (rabbit α-HA (Cell Signaling Technology, C29F4) at 1:1000, rat α-Flag (Novus Biologicals, NBP1-06712) at 1:1000; and the following secondary antibodies: goat α-mouse AF647 (Invitrogen, A21236) at 1:500, goat α-rabbit AF488 (Invitrogen, A11034) at 1:500, and goat α-rat AF568 (Invitrogen, A11077) at 1:500.

Reproductive tract dissections and staining was performed as described[42]. As the primary antibody, we used chicken α-GFP (Invitrogen, A10262) at 1:1000 followed by goat α-chicken AF488 (Life Technologies, A11039) at 1:500 with the addition of Alexa Fluor 633 phalloidin (ThermoFisher Scientific) to visualize the muscle-lined reproductive tract (1:500).

Confocal imaging: Samples were imaged on a LSM710 confocal microscope (Zeiss) with a Plan-Apochromate 20x0.8 M27 objective and images were prepared in Fiji.

Electron Microscopy neural reconstructions and connectivity: SAG was previously partially reconstructed[16] in the Full Adult Female Brain (FAFB[27]) electron microscopy dataset using the CATMAID software[43]. We completed additional tracing of this neuron by continuing branches with a combination of both manual and assisted tracing. Manual tracing consisted of generating a skeleton of the neuron by following the neuron and marking the center of each branch. Assisted

tracing consisted of joining and proofreading pre-assembled skeleton fragments with automated
segmentation[44]. In addition to the skeleton tracing, new chemical synapses were also annotated
as previously described[27]. Finally, downstream synaptic targets of SAG were then traced out
from these additional locations using both manual and assisted tracing techniques as described
above.
Neurons traced in CATMAID, including SAG, pC1, egg-laying circuit neurons[16] and sexual
receptivity circuit neurons[17], were all located in Flywire (flywire.ai), which uses the same EM
electron microscopy dataset[27]. To identify synaptic partners, we used connectome annotation
versioning engine (CAVE[25,26]) using a cleft score cutoff of 100 to generate synapses of relatively
high confidence[26].
447
Intersectional method to access SPSNs:
Based on previous reports, three genetic drivers label the SPSN (VT003280[3], fru[14,15] and
ppk[14,15]) with driver lines available in two separate binary expression systems (UAS-GAL4
and LexA-LexAop). With the use of a conditional reporter line (UAS < stop < GFP) and in
combination with inducible FLP line (LexAopFLP) three combinations were produced: fru-
LexA∩ppk-GAL4, fru-LexA∩VT3280-GAL4, and VT3280-LexA∩ppk-GAL4. Images of the
female reproductive tract were used to confirm labeling of the SPSNs and images of the brain to
determine off-target neural expression. With the use of a conditional effector line (UAS < stop <
KIR or UAS < stop < TNT), SPSNs were silenced and females (along with genetic controls)
were tested for changes in sucrose consumption.
458
Split-GAL4 screening and stabilization
pCd-2 split generation:
Using the FlyEM Hemibrain V1.2.1 dataset via NeuPRINT, we queried "SAG" and explored the
output neuron list, identifying three pCd-2 neurons per hemisphere (SMP286, SMP287,
SMP288). From here, we used Neuron Bridge to manually explore the light microscopy matches
and generated a list of possible hemi-driver that labeled this cell type. The expression pattern of
the p65ADZp and ZpGAL4DBD combinations were examined by driving the expression of UAS
reporter (20xUAS-CsChrimson-mVenus in attP18). Combinations with specific expression in
pCd-2 neurons were stabilized.
468
SAGb split generation:
Using Neuron Bridge, we queried "VT050405" (the genetic construct used as the p65ADZp
hemidriver in SAG-SS3) and identified multiple "Body Ids" that morphologically matched the
non-SAG ascending cell type. From here we generated a "Color Depth Search" to create a list of
possible hemi-driver that labeled this cell type. The expression pattern of the p65ADZp and
ZpGAL4DBD combinations were examined by driving the expression of UAS reporter
(20xUAS-CsChrimson-mVenus in attP18) and combinations with specific expression in SAGb
neurons were stabilized.
477
478   **Statistical analysis:** Statistical tests for all experiments, with the exception of in vivo calcium
479   imaging and in vivo functional connectivity experiments, were performed in GraphPad Prism.
480   For two- and three-group comparisons, data was first tested for normality with the KS normality
481   test (alpha = 0.05). If all groups passed then groups would be compared with a parametric test (t-
482   test or One-Way ANOVA, respectively). If at least one group did not pass, groups were

compared with a non-parametric version (Mann-Whitney test or Kruskal-Wallis test, respectively). For all multi-factorial designs, a Two-Way ANOVA was performed with a Bonferroni post-hoc test. All statistical tests, significance levels, and number of data points (N) are specified in the figure legend.

Normalized data sets: Most datasets from optogenetic assays were normalized within each genotype. To generate this normalized dataset, data from females within the no light condition was averaged, creating a "no-light mean" for each genotype. This value was subtracted from each individual female within the light condition of the corresponding genotype. This dataset was then graphed and statistical analysis was performed as outlined above.

39. Bidaye, S. S., Laturney, M., Chang, A. K., Liu, Y., Bockemühl, T., Büschges, A. & Scott, K. Two brain pathways initiate distinct forward walking programs in Drosophila. *Neuron***108**, doi: 10.1016/j.neuron.2020.07.032 (2020).

40. Yao, Z. & Scott, K. Serotonergic neurons translate taste detection into internal nutrient regulation. *Neuron***110**, 1036-1050 (2022).

41. Nern, A., Pfeiffer, B. D. & Rubin, G. M. Optimized tools for multicolor stochastic labeling reveal diverse stereotyped cell arrangement in the fly visual system. *Proc. Natl. Acad. Sci. USA***112**, E2967-2976 (2015)

42. Billeter, J. C. & Goodwin, S. F. Characterization of Drosophila fruitless-gal4 transgenes reveals expression in male-specific fruitless neurons and innervation of male reproductive structures. *J. Comp. Neurol.* **475**, 270-287 (2004).

43. Saalfeld, S., Cardona, A., Hartenstein, V. & Tomancak, P. CATMAID: collaborative annotation toolkit for massive amounts of image data. *Bioinformatics***25**, 1984-1986 (2009).

44. Li, F., Lindsey, L. F., Januszewski, M., Tyka, M., Maitin-Shepard, J., Blakely, T. & Jain, V. Automated reconstruction of a serial-section EM Drosophila brain with flood-filling networks and local realignment. *Micro. Microanal.* **25**, 1364-1365 (2019).

Figure	short genotype	full genotype
1n-o	Gr65f-GAL4 > GCaMP	*UAS-CD8::tdTomato/+;20xUAS-IVS-GCaMP6s(attP40)/Gr64f-GAL4; 20xUAS-IVS-GCaMP6s(VK00005)/Gr64f-GAL4*
2a	hs-bam	*Hsp70(generic)-bam.O;+;+*
2c	fru-LexA ∩ ppk-GAL4 > GFP	*w*; 8xLexAop2-FLPL(attP40)/UAS(FRT.stop)Hsap\mcd8::GFP(VIE-19a); Fru-PI-LexA/ppk-GAL4*
2d	fru-LexA ∩ ppk-GAL4 > TNT	*w+; 8xLexAop2-FLPL(attP40)/w[+m*]=UAS(FRT.stop)Ctet\tetX(VIE-19A); Fru-PI-LexA/ppk-GAL4*
2d	fru-LexA, UAS-TNT (control)	*w+; w[+m*]=UAS(FRT.stop)Ctet\tetX(VIE-19A)/+; Fru-PI-LexA/+*
2e	fru-LexA ∩ ppk-GAL4 > KIR2.1	*w+;8xLexAop2-FLPL(attP40)/UAS(FRT.stop)Hsap\KCNJ2 (VIE-19A); Fru-PI-LexA/ppk-GAL4*
2e	fru-LexA, UAS-KIR (control)	*w+; UAS(FRT.stop)Hsap\KCNJ2(VIE-19A)/+; Fru-PI-LexA/+*
2d,e	ppk-GAL4 (control)	*w+;8xLexAop2-FLPL(attP40)/+;ppk-GAL4/+*
2f	SAG-SS3 > GFP	*w*;20xUAS-csChrimson::mVenus(attP18)/+; VT050405-p65.AD(attP40)/+;dsx^DBD/+*
2g	SAG-SS3 > GtACR1	*w+; VT050405-p65.AD(attP40)/+;20xUAS-IVS-GtACR1 EYFP(attP2)/dsx^DBD*
2g	SAG-SS3 (control)	*w+; VT050405-p65.AD(attP40)/+; dsx^DBD/+*
2h	pC1-SS1 > GFP	*w*;20xUAS-csChrimson::mVenus(attP18)/+; VT002064-p65AD (attP40)/+; VT008469-GAL4DBD(attP2)/+*
2i	pC1-SS1 > GtACR1	*w+; VT002064-p65AD(attP40)/+;20xUAS-IVS-GtACR1-EYFP (attP2)/VT008469-GAL4DBD(attP2)*
2i	pC1-SS1 (control)	*w+; VT002064-p65AD(attP40)/+; VT008469-GAL4DBD(attP2)/+*
2g,i	GtACR1 (control)	*w+; +; 20xUAS-IVS-GtACR1-EYFP(attP2)/+*
3b	vpoDN-SS1 > GtACR1	*w+; 31D07-p65AD(attP40)/+;20xUAS-IVS-GtACR1 EYFP(attP252F12-GAL4DBD(attP2)*
3b	vpoDN-SS1 (control)	*w+; 31D07-p65AD(attP40)/+;52F12-GAL4DBD(attP2)/+*
3c	vpoDN-SS2 > GtACR1	*w+; VT045670-p65AD(attP40)/+; 20xUAS-IVS-GtACR1-EYFP (attP2)/52F12-GAL4DBD(attP2)*
3c	vpoDN-SS2 (control)	*w+; VT045670-p65AD(attP40)/+;52F12-GAL4DBD(attP2)/+*
3b,c	GtACR1 (control)	*w+; +; 20xUAS-IVS-GtACR1-EYFP(attP2)/+*
3d-f	oviDN-SS1 > csChrimson	*w*,20xUAS-csChrimson::mVenus(attP18)/+; VT050660-p65.AD (attP40)/+; VT028160-GAL4DBD(attP2)/+*
3d-f	oviDN-SS1 (control)	*w*; VT050660-p65.AD(attP40)/+; VT028160-GAL4DBD(attP2)/+*
3d-g	csChrimson (control)	*w*,20xUAS-csChrimson::mVenus(attP18)/+;+*
3d,e,g	oviDN-SS2 > csChrimson	*w*,20xUAS-csChrimson::mVenus(attP18)/+; VT026873-p65.AD (attP40)/+; VT040574-GAL4DBD(attP2)/+*
3d,e,g	oviDN-SS2 (control)	*w*; VT026873-p65.AD(attP40)/+; VT040574-GAL4DBD(attP2)/+*
4e	pCd-2a-SS1 > GFP	*w*,20xUAS-csChrimson::mVenus(attP18)/+; VT027804-p65.AD (attP40)/+; dsx^DBD/+*
4f	pCd-2b-SS1 > GFP	*w*,20xUAS-csChrimson::mVenus(attP18)/+; R9B05-p65.AD (attP40)/+; R22D06-GAL4DBD(attP2)/+*
4g,h	pCd-2a-SS1 > GtACR1	*w+; VT027804-p65AD(attP40)/20xUAS-IVS-GtACR1-EYFP (attP2)/dsx^DBD*
4g,h	pCd-2a-SS1 (control)	*w+; VT027804-p65AD(attP40)/+; dsx^DBD/+*
4g,i	pCd-2b-SS1 > GtACR1	*w+; R9B05-p65AD(attP40)/20xUAS-IVS-GtACR1-EYFP(attP2)/ R22D06-GAL4DBD(attP2)/+*
4g,I	pCd-2b-SS1 (control)	*w+; R9B05-p65AD(attP40)/+; R22D06-GAL4DBD(attP2)/+*
4g-I	GtACR1 (control)	*w+; +; 20xUAS-IVS-GtACR1-EYFP(attP2)/+*
5d	pC1-SS1 > Chrimson, Lgr3 > GCaMP6s	*w*,LexAop2-syn21-opGCaMP6s,10x-UAS-syn21-Chrimson88-tdTomato; VT002064-p65AD(attP40)/R19B09-LexA(attp40); VT008469-GAL4DBD(attP2)/+*
5e	pCd-2a-SS1 > Chrimson, Lgr3 > GCaMP6s	*w*,LexAop2-syn21-opGCaMP6s,10x-UAS-syn21-Chrimson88-tdTomato; VT027804-p65AD(attP40)/R19B09-LexA(attp40);*

567			*dsx^{DBD}/+*
568	5f	pCd-2b-SS1 > Chrimson,	*w*,LexAop2-syn21-opGCaMP6s,10x-UAS-syn21-Chrimson88-*
569		Lgr3 > GCaMP6s	*tdTomato; R9B05-p65AD(attP40)/R19B09-LexA(attp40);*
570			*R22D06-GAL4DBD(attP2)/+*
571	5d-f	control	*w*,LexAop2-syn21-opGCaMP6s,10x-UAS-syn21-Chrimson88-*
572			*tdTomato; R19B09-LexA(attp40)/+;+*
573	5h	Lgr3 ∩ Fru > csChrimson	*w-,20xUAS(FRT.stop)csChrimson:mVenus(attP18)/+;*
574			*8xLexAop2-FLPL(attP40)/+; Fru-PI-LexA/R19B09-GAL4*

Supplementary Table 1 | Fly genotypes in figures

	Type	Name	Genotype	Reference/Source
579				
580	Wild-type	Canton-S	w+;+;+	N/A
581	Driver	Gr64f-Gal4	w-;Gr64-gal4;+	RRID: BDSC_57669
582	Driver	Gr64f-Gal4	w-;+;Gr64f-gal4	RRID: BDSC_57668
583	Reporter	UAS-GCaMP	w-;UAS-GCaMP6s(attP40);+	RRID: BDSC_42746
584	Reporter	UAS-GCaMP	w-;+;UAS-GCaMP6s(VK00005)	RRID: BDSC_42749
585	Reporter	UAS-tdTomato	UAS-CD8-tdTomato;+;+	[45]
586	Effector	ovoD	w+;+;ovoD1	[23]
587	Effector	hs-BAM	hs-bam,w1118;+;+	RRID: BDSC_24636
588	Driver	nsyb-Gal4	w-;+;nsyb-gal4	RRID: BDSC_51941
589	Effector	UAS-dcr2	w-;UAS-dcr-2;+	RRID: BDSC_24650
590	Effector	UAS-SPR-RNAi	w-;+;UAS-SPR-IR1	[13]
591	Driver	Ppk-gal4	w-;+;ppk-gal4	RRID: BDSC_32079
592	Driver	VT3280-Gal4	w-;+;VT003280-gal4 attP2	[3]
593	Driver	VT3280-LexA	w-;+;VT003280-LexA attP2	[3]
594	Driver	Fru-LexA	w-;+;Fru$^{P1-LexA}$	[46]
595	Effector	LexAop2-FLPL	w-;8xLexAop2-FLPL(attP40);+	RRID: BDSC_55820
596	Effector	UAS>stop>KIR	w+;UAS > stop > KIR	RRID: BDSC_67686
597	Effector	UAS>stop>TNT	w+;UAS > stop > TNT	RRID: BDSC_67690
598	Reporter	UAS>stop>GFP	w-;UAS > stop > CD8::GFP;+	RRID: BDSC_30125
599	Driver	SAG-SS3	VT050405-p65ADZp(attP40);	[47]
600			dsx-ZpGal4DBD	[48]
601	Driver	pC1-SS1	VT002064-p65ADZp(attP40);	[16]
602			VT008469-ZpGal4DBD(attP2)	
603	Driver	pC1-SS2	VT002064-p65ADZp(attP40);	[16] (modified)
604			dsx-ZpGal4DBD	
605	Driver	vpoDN-SS1	31D07-p65ADZp(attP40);	[17]
606			52F12-ZpGal4DBD(attP2)	
607	Driver	vpoDN-SS2	VT045670-p65ADZp(attP40);	[17]
608			52F12-ZpGal4DBD(attP2)	
609	Driver	oviIN-SS1	68A10-p65ADZp(attP40);	[16]
610			VT010054-ZpGal4DBD(attP2)	
611	Driver	oviIN-SS2	VT026347-p65ADZp(attP40);	[16]
612			VT026035-ZpGal4DBD(attP2)	
613	Driver	oviDN-SS1	VT050660-p65ADZp(attP40);	[16]
614			VT028160-ZpGal4DBD(attP2)	
615	Driver	oviDN-SS2	VT026873-p65ADZp(attP40);	[16]
616			VT040574-ZpGal4DBD(attP2)	
617	Effector	UAS-GtACR	w+;;20xUAS-IVS-GtACR1-EYFP9attP2)	[49]
618	Effector	UAS-csChrimson,	w-,20xUAS-ssChrimson::mVenus(attP18);+;+	RRID: BDSC_55134
619	Effector	MCFO		RRID: BDSC_
620	Driver	SAGb-SS1	VT004974-p65ADZp(attP40);	RRID: BDSC_71475
621			R69H11-ZpGal4DBD(attP2)	RRID: BDSC_69819
622	Driver	SAGb-SS2	VT023823-p65ADZp(attP40);	[47]
623			R69H11-ZpGal4DBD(attP2)	RRID: BDSC_69819
624	Driver	pCd-2a-SS1	VT027804-p65ADZp(attP40);	RRID: BDSC_72041
625			dsx-ZpGal4DBD	[48]
626	Driver	pCd-2a-SS2	VT057275-p65ADZp(attP40);	RRID: BDSC_71664
627			VT027804-ZpGal4DBD(attP2)	RRID: BDSC_71758
628	Driver	pCd-2b-SS1	R9B05-p65ADZp(attP40);	RRID: BDSC_70519
629			R22D06-ZpGal4DBD(attP2)	[50]
630	Driver	pCd-2b-SS2	R29G03-p65ADZp(attP40);	RRID: BDSC_75746
631			R22D06-ZpGal4DBD(attP2)	[50]
632	Driver	pCd-2b-SS3	R60C09-p65ADZp(attP40);	RRID: BDSC_69750
633			R22D06-ZpGal4DBD(attP2)	[50]

Driver	R19B09-gal4	R19B09-Gal4(attP2)	RRID: BDSC_48840
Driver	R19B09-LexA	R19B09-LexA(attP40)	RRID: BDSC_52539
Effector	UAS>stop>csChrimson w-,UAS > stop > csChrimsner(attP18)		[51]
Effector	UAS-Chrimson88 UAS-Chrimson88;+;+		[52]
Reporter	LexAop-GCaMP6s	LexAop-GCaMP6s;+;+	[53]

Supplementary Table 2 | Fly Stocks

Supplementary References

45. Thistle, R., Cameron, P., Ghorayshi, A., Dennison, L. & Scott, K. Contact chemoreceptors mediate male-male repulsion and male-female attraction during Drosophila courtship. *Cell* **149**, 1140-1151 (2012).

46. Mellert, D. J., Knapp, J. M., Manoli, D. S., Meissner, G. W. & Baker, B. S. Midline crossing by gustatory receptor neuron axons is regulated by fruitless, doublesex and theh Roundabout receptors. *Development* **137**, 323-332 (2010).

47. Tirian, L. & Dickson, B. J. The VT GAL4, LexA, and split-GAL4 driver line collections for targeted expression in the Drosophila nervous system. bioRxiv doi: https://doi.org/10.1101/198648.

48. Shirangi, T. R., Wong, A. M., Truman, J. W. & Stern, D. L. Doublesex regulates the connectivity of a neural circuit controlling Drosophila male courtship song. *Dev. Cell* **37**, 533-544 (2016).

49. Mohammad, F., Stewart, J. C., Ott, S., Chlebikova, K., Chua, J. Y., Koh, T. W., Ho, J & Claridge-Chang, A. Optogenetic inhibition of behavior with anion channelrhodopsins. Nat. *Methods* **14**, 271-274 (2017).

50. Dionne, H., Hibbard, K. L., Carvallaro, A., Kao, J. C. & Rubin, G. M. Genetic reagents for making split-GAL4 lines in Drosophila. *Genetics* **209**, 31-35 (2018).

51. Wu, M., Nern, A., Williamson, W. R., Morimoto, M. M., Reiser, M. B., Card, G. M. & Rubin, G. M. Visual projection neurons in the Drosophila lobula link feature detection to distinct behavioral programs. *Elife*, doi: 10.7554/eLife.21022.

52. Klapoetke, N. C. et al. Independent optical excitation of distinct neural population. *Nature Methods* **11**, 338-346 (2014).

53. Broussard, G. J. et al. In vivo measurement of afferent activity with axon-specific calcium imaging. *Nat. Neurosci.* **21**, 1272-1280 (2018).

Acknowledgements
Members of the Scott lab provided contributions to experimental design, data analysis, and manuscript preparation. Salil Bidaye assisted with the analyses of locomotor data. This work was supported by NIH R01DC013280 (K.S.). Neuronal reconstruction for this project took place in a collaborative CATMAID environment in which 27 labs are participating to build connectomes for specific circuits. Therefore, work done for published (Wang et al. 2020) and ongoing projects has proved useful to us. Development and administration of the FAFB tracing environment and analysis tools were funded in part by National Institutes of Health BRAIN Initiative grant 1RF1MH120679-01 to Davi Bock and Greg Jefferis, with software development effort and administrative support provided by Tom Kazimiers (Kazmos GmbH) and Eric Perlman (Yikes LLC). We thank Peter Li for sharing his automatic segmentation (citation is https://doi.org/10.1101/605634). Tracing in Cambridge was supported by Wellcome Trust (203261/Z/16/Z) and ERC (649111) awards to G. Jefferis. We thank the Princeton FlyWire team and members of the Murthy and Seung labs for development and maintenance of FlyWire (supported by BRAIN initiative grant MH117815 to Murthy and Seung). We found the partial tracing of SAG and pC1 in CATMAID useful (Wang et al. 2020). In addition to our tracing efforts, we are appreciative of the tracing accomplished by other labs. For help with tracing of pC1 in FlyWire, we thank the Murthy and Seung labs (77% of pC1a, 98% of pC1b, 92% of pC1c) and the Jefferis lab (7% of pC1a, 8% of pC1c). For help with tracing pCd-2 cells we would like to thank the Dickson lab (CATMAID: 76% of pCd-2a, 44% of pCd-2b, 28% of pCd-2c), the Jefferis lab (CATMAID: 20% of pCd-2a, 6% of pCd-2b, 3% of pCd-2c; FlyWire: 32% of pCd-2a, 8% of pCd-2b, 13% of pCd-2c), the Murthy and Seung lab (FlyWire: 15% of pCd-2a, 11% of pCd-2b, 6% of pCd-2c), and Janelia tracers (FlyWire: 52% of pCd-2a, 14% of pCd-2b). We would like to thank the Jefferis labs (46%) and the Murthy and Seung labs (50%) for their contribution to tracing the t-shaped Lgr3 cells. The tracers who contributed include Nora Forknall, Lily Talley, Katie Stevens, Shanice Bailey, Tansy Yang, Istvan Taisz, Greg Jefferis, Lucas Encarnacion-Rivera, Austin T. Burke, Yijie Yin, James Hebditch, Kyle Patrick Willie, Joshua Bañez, Philipp Schlegel, Dharini Sapkal, Irene Salgarella, Dhwani Patel, Ben Silverman, Bhargavi Parmar, Chitra Nair, Doug Bland, Laia Serratosa, Marta Costa, Nash Hadjerol, Regine Salem, Rey Adrian Candilada, Shirleyjoy Serona, Zairene Lenizo, Zeba Vohra, and Zepeng Yao.

Author contributions
M.L. and K.S. conceived and designed the study and wrote the manuscript. M.L. performed the experiments and analyzed the data with the exception of the functional imaging experiment, performed and analyzed by G.S.

Competing interests
The authors declare no competing interests.

Corresponence and requests for materials should be addressed to K.S. or M.L.