Genome size variation between pelagic and benthic communities across

prokaryotic taxonomy and environmental gradients in the Baltic Sea

<u>RUNNING TITLE</u>: Genome size in benthic and pelagic Baltic Sea

<u>AUTHORS</u>

Alejandro Rodríguez-Gijón[$1,2], Moritz Buck[3], Anders F. Andersson[2,4], Dandan Izabel-Shen[1],

Francisco J. A. Nascimento[1,5], Sarahi L. Garcia[$*1,2]

$ Correspondent authors

alejandro.rgijon@gmail.com

sarahi.garcia@su.se

1 Department of Ecology, Environment and Plant Sciences, Stockholm University,

Stockholm 106 91, Sweden

2 Science for Life Laboratory, Stockholm, Sweden

3 Department of Aquatic Sciences and Assessment, Swedish University of Agricultural

Sciences, Uppsala, Sweden

4 Department of Gene Technology, School of Engineering Sciences in Chemistry,

Biotechnology and Health, KTH Royal Institute of Technology, Stockholm, Sweden

5 Baltic Sea Centre, Stockholm University, Stockholm, Sweden

<u>Keywords</u>: microbial ecology, genome size, bacteria, Baltic Sea, streamlining theory

ABSTRACT (250 words)

Microbial genome size can be used as a predictor to explain the ecology and metabolism

of Bacteria and Archaea across major biomes. Despite their ecological significance, the

contribution of microbial genome size to differences in metabolic potential of benthic and

pelagic prokaryotes are poorly studied. Here, we investigated how taxonomy and microbial

genome size varies between benthic and pelagic habitats across environmental gradients of

the brackish Baltic Sea. We also explored the relationships between that variation, the

environmental heterogeneity, and microbial functions in these habitats. By analyzing Baltic

metagenomes and MAGs, we observe that pelagic brackish Bacteria and Archaea present

smaller genome sizes on average than pelagic marine and freshwater prokaryotes. Moreover,

we found that prokaryotic genome sizes in Baltic sediments (3.47 Mbp) are significantly

bigger than in the water column (2.96 Mbp). These differences in genome size persisted in

Bacteria from the phyla level to the order level. For pelagic prokaryotes, the smallest

genomes coded for a higher number of module steps per Mbp than bigger genomes for most

of the functions, such as amino acid metabolism and central carbohydrate metabolism.

However, we observed that nitrogen metabolism was almost absent in pelagic genomes and

was mostly present in benthic genomes. Finally, we also show that Bacteria inhabiting Baltic

sediments and water column not only differ in taxonomy, but also in their metabolic

potential, such as the Wood-Ljungdahl pathway or presence of different hydrogenases.

IMPORTANCE (150 words)

In Bacteria and Archaea, genome size is the result of strong selection pressures

shaping the ecology and metabolism of microbial lineages. Thanks to the development of

metagenomic analysis of environmental samples in the last three decades, we can now survey

the size of their genomes, quantify their functional capabilities, and infer the role they play in

the environment. Here, by using four publicly available datasets together with one new

unpublished dataset, we provide new insights on how genome size, metabolism and

taxonomy are linked and how they differ between sediments and water column in the Baltic

Sea. Moreover, we investigate differences in genome size of pelagic microorganisms across

ecosystems, from the brackish Baltic Sea to marine and freshwater Bacteria and Archaea. Our

results provide a comprehensive and contrasting view between the taxonomic and genetic

diversity of sediments and water column and its implications on nutrient cycling in the Baltic

Sea.

<u>INTRODUCTION</u>

Genome size in Bacteria and Archaea can be used as a predictor of microbial

metabolic complexity and niche occupation across Earth's biomes (Lynch, 2006; Maistrenko

et al., 2020; Rodríguez-Gijón et al., 2022; Swan et al., 2013). Over the last two decades,

different explanations have been postulated to find the mechanisms responsible for the

variations in prokaryotic genome length. Reported drivers of genome size are phylogenetic

history (Biller et al., 2015; Martinez-Gutierrez and Aylward, 2022), lifestyle such as free-

living, particle attached or host-associated (Giovannoni et al., 2014; Moran and Mira, 2001;

van Ham et al., 2003), or environment such as marine, freshwater, different types of

sediments or different hosts in host-associated microorganisms (Rodríguez-Gijón et al., 2022;

Simonsen, 2022). While many factors have been found to partially explain genome size,

ultimately, this implies that multiple evolutionary forces and their interactions shape genome

size in prokaryotes. Untangling how and why different lineages have dissimilar genome sizes

across different ecosystems can be key to understand the ecology and evolutionary history of

Bacteria and Archaea.

Aquatic microorganisms have been extensively sampled and have a large

representation in genomic and metagenomic datasets (Nayfach et al., 2021). Their genome

size spans from ~0.5 to ~15, averaging 3.1 million base pairs (Mbp) (Rodríguez-Gijón et al.,

2022). Aquatic environments are heterogeneous and many different abiotic factors, such as

salinity and depth, could be responsible for the microbial genome size variation. Within

marine environments, it has been postulated that pelagic microbes inhabiting deep marine

environments present bigger genome sizes than those in shallow marine waters

(Konstantinidis et al., 2009; Mende et al., 2017). Within freshwater ecosystems, we can

observe that isolates from the family Methylophilaceae (class Gammaproteobacteria) show a

decrease in genome size in the transition from sediments to pelagic forms (Salcher et al.,

2019). Additionally, several studies have shown that marine microbes have smaller genome

sizes than freshwater microorganisms (Cabello-Yeves et al., 2022; Chen et al., 2021;

Rodríguez-Gijón et al., 2022). Yet, genome size variation in the brackish realm remains

debated: Actinobacteria in the brackish Baltic Sea show bigger genome sizes than in

freshwaters (Hugerth et al., 2015), while picocyanobacteria show the opposite trend (Cabello-
Yeves et al., 2022). Further research must be done to understand how genome sizes vary
between and within different aquatic environments, particularly brackish water bodies
including pelagic and benthic microorganisms.

In pelagic Bacteria, many abundant and ubiquitous lineages present reduced genomes
(Hugerth et al., 2015). The Streamlining Theory has been proposed to explain this tendency
towards simplification in aquatic environments (Giovannoni et al., 2014). This evolutionary
theory indicates that abundant and ubiquitous aquatic microorganisms exhibit small cells and
reduced genomes, as observed for *Pelagibacter ubique* (class Alphaproteobacteria)
(Giovannoni, 2005; Grote et al., 2012) or *Prochlorococcus* spp. (phylum Cyanobacteria)
(Biller et al., 2015). In many cases, reduced genomes imply the absence of key metabolic
capabilities, such as the electron transport chains in Patescibacteria (Beam et al., 2020) or the
catalase-peroxidase in *Prochlorococcus* (Scanlan et al., 2009). The absence of essential
metabolic capabilities is explained by the Black Queen Hypothesis (Morris et al., 2012):
when the fitness cost of a function is higher than its benefit, microbes might stop performing
it and, instead, obtain benefit from leaky metabolites from neighboring cells, establishing
strong interdependencies. As a whole, genetic streamlining has significant consequences on
genome size, metabolism, adaptability, and lifestyle in microorganisms.

Environmental conditions strongly select which metabolic functions are present or
absent on microbial communities and, consequently, the length of the genome. Metagenomic
studies show that a vast majority of the genes in Bacteria and Archaea are specific to
particular environments, whereas very few genes are being shared between environments
(Coelho et al., 2022). This remarks how relevant is the relationship between niche specificity
and lineage specific functional traits to understand microbial ecology and evolution (Zhou et
al., 2020). These functional capabilities also differ between water column and sediments in
Bacteria and Archaea in both marine (Acinas et al., 2021; Louca et al., 2016) and freshwater
environments (Li et al., 2018). As a whole, gene repertoires and genome size are interrelated,
and researchers must consider them together with environmental gradients to better
understand niche-specificity and the ecology of different prokaryotic lineages.

In this research article, our main aim is to link the genome size of Bacteria and
Archaea with their ecology and functional potential across the environmental gradients in the

Baltic Sea in both pelagic and benthic habitats. Specifically, we investigate; i) how genome

size varies among brackish communities and taxonomic lineages of Bacteria and Archaea

from sediments and water column, ii) what relationship can be found between genome size

and the number of metabolic capabilities in Bacteria in the Baltic Sea, and iii) which taxa and

metabolic pathways contrast between pelagic and benthic Bacteria. To achieve this, we

selected three pelagic and one benthic publicly available metagenomic datasets and we

provide one new benthic dataset to study genome size distribution across sediments and water

column in the Baltic Sea (**Figure 1**). We use two different approaches to study genome size:

the estimated average genome size (AGS) per metagenomic sample, and the estimated

genome size of bacterial and archaeal metagenome-assembled genomes (MAGs). Our results

show that Bacteria and Archaea inhabiting brackish sediments present bigger genome sizes

than those in the water column. We also find that microorganisms inhabiting the Baltic

benthos have more metabolically-versatile genomes than pelagic prokaryotes, which mean

they code for a wider range of metabolic capabilities. Finally, we also find that some

functions, such as nitrogen metabolism, are mainly found in benthic Bacteria but are almost

absent in pelagic bacteria the Baltic Sea.

<u>RESULTS AND DISCUSSION</u>

Benthic metagenomes and MAGs show bigger estimated average genome sizes

than pelagic in the Baltic Sea. First, we calculated the average genome size (AGS) of

metagenomes across the latitudinal gradient of the Baltic Sea using two sediment and two

pelagic datasets. We observed that sediment-dwelling microbial communities present

significantly larger average genome sizes (mean AGS = 6.01 Mbp, n = 108) than pelagic

communities (mean AGS = 5.40 Mbp, n = 69) (t-test, $p < 0.01$) (**Figure 2A**). We then

evaluated the relationships between AGS of metagenomes and each of the environmental

variables (depth, salinity, temperature, and oxygen concentration) independently and their

interactions (ANOVA type II). The AGS in the pelagic metagenomes is significantly

associated only with depth and shows a negative correlation (**Supplementary Material 1** and

Figure S1). Previous marine analysis have found the opposite effect, bigger genome sizes in

deeper areas (Mende et al., 2017). However, our pelagic analysis only covers 5 meters of

depth and explains 19% of the genome size variation in pelagic metagenomes. AGS in

sediment metagenomes is significantly associated with both salinity and depth and most of

the interactions of these variables (**Supplementary Material 1**). Both depth and salinity have

a weak positive correlation with average genome size in the sediment samples (**Figure S1**).

To our knowledge, this is the first time where a positive correlation between genome size and

salinity is reported for brackish sediment microorganisms.

Although the ANOVA did not show a significant effect of water oxygen

concentration on genome size, we further investigate the effect of bottom water O_2

concentration in the AGS of metagenomes from sediments. We separated these metagenomes

into those from oxygen concentration 0 to 2 mg/L (mostly metagenomes from the Death

Zone) and those metagenomes with oxygen concentration 2 to 12.45 mg/L. We observe that

benthic metagenomes from lower oxygen concentration (mean AGS = 7.08 Mbp, n = 14)

present significantly higher genome sizes than those in sediments with oxygen concentration

higher than 2 mg/L (mean AGS = 5.85 Mbp, n = 94) (t-test, $p < 0.01$) (**Figure 2B**).

Complementarily, previous results show that the Dead Zone bacteria tend to be more

metabolically similar to each other when compared to bacteria from oxic sediments (Broman

et al., 2022). Our observation confirms a recent study that observed microaerophilic

microorganisms from diverse environments present smaller genome sizes than obligate

anaerobic (Nielsen et al., 2021). Moreover, previous studies show a positive correlation

between genome size and nutrient concentration (Allen et al., 2012; Aylward and Santoro,

2020; Mende et al., 2017). Dead Zones in the Baltic Sea are characterized by anthropogenic

eutrophication (Conley et al., 2009), which would also promote bigger genome sizes.

Altogether, these results indicate that high nutrient concentration and low oxygen

concentrations in Baltic Sea Dead Zones promote bigger genome sizes with similar metabolic

capabilities.

To compare genome size between pelagic and benthic Bacteria and Archaea, we

obtained metagenome-assembled genomes (MAGs; >75% completeness and <5%

contamination) from our metagenome datasets. Our dataset compiles 216 MAGs from the

sediments that dereplicated into 56 representative MAGs (95% average nucleotide identity).

Additionally, 1920 pelagic MAGs were dereplicated into 340 representative MAGs. We

observe that 12 phyla were detected in both sediments and water column, while 16 phyla

were specific to either habitat. Seven phyla were found specific to the sediment (14

representative MAGs) and nine phyla were specific to the water column (29 representative

MAGs) (**Table 1**). Interestingly, only one species-level representative MAG was binned from

both sediments and water column, belonging to genus *Mycobacterium* (phylum

Actinobacteriota). These differences on taxonomical composition in microbial communities
between water column and sediments altogether with latitudinal changes in microbial
biodiversity in the Baltic Sea (Herlemann et al., 2011) show how heterogeneous brackish
environments are.

Additionally, the eight largest representative genomes of the dataset belonged to
phylum Planctomycetota (family Planctomycetaceae) (7.95 – 9.69 Mbp) belonging to the
water column and not to sediments. It has been previously observed that Planctomycetota is
the phylum containing the aquatic MAG with the biggest known estimated genome size
(14.93 Mbp) (Rodríguez-Gijón et al., 2022). Further research to understand if extant genome
size in phylum Plactomycetota is the result of taxonomical divergence or ecological
adaptation to abiotic gradients would enlighten the evolution in this group. On the other side
of the genome size spectrum, the representative MAG with the smallest estimated genome
size belongs to class Alphaproteobacteria (family Pelagibacteraceae, 1.08 Mbp; **Table 1**).
Bacteria from this family has been widely reported to be streamlined, abundant and
ubiquitous (Giovannoni, 2017).

Altogether, representative MAGs from sediments presented an average estimated
genome size of 3.47 Mbp, which was significantly higher than for water column MAGs (2.96
Mbp) (t-test, $p < 0.01$) (**Figure 3A**). Bacteria from sediments presented the biggest estimated
genome size on average (3.67 Mbp), followed by pelagic bacteria (2.98 Mbp), pelagic
Archaea (1.97 Mbp) and sediment Archaea (1.43 Mbp) (**Figure 3B**). This is supported by
previous results, as Bacteria show bigger genome sizes than Archaea regardless of the
environment (Rodríguez-Gijón et al., 2022). Moreover, the average estimated genome size of
Baltic sediments (3.47 Mbp) is more similar to that of terrestrial microbial genomes (3.7
Mbp) (Rodríguez-Gijón et al., 2022). We also observe that pelagic microorganisms tend to be
more streamlined than those present in sediments, at the phyla (**Figure 3C**), class (data not
shown) and order level (**Figure 3D**). Our results corroborate previous findings that
streamlining is common in pelagic marine environments (Batut et al., 2014; Giovannoni,
2017, 2005) and pelagic brackish environments (Hugerth et al., 2015).

In our study, the average estimated AGS for the sediments (6.01 Mbp) and water
column (4.44 Mbp) is larger than the average estimated genome size of the MAGs assembled
and binned from the sediments (3.47 Mbp) and water column (2.96 Mbp), respectively

(**Figure 2A** and **3A**). We calculated the AGS for the metagenomic samples to estimate the
average genome size of the whole microbial community. We used MicrobeCensus as a robust
and accurate tool to calculate AGS (Pereira-Flores et al., 2019). However, this AGS of
metagenomes is biased by the viral and eukaryotic reads that might be present in the sample
(Kristensen et al., 2010; Lind and Pollard, 2021). On the other hand, there is a bias with
looking at the average estimated genome size of MAGs. This bias stems from all those
bacteria and archaea that are hard to assemble and bin due to high genomic intrapopulation
diversity (Ghurye, 2016). In our study, we use two methods that have different biases to
answer the same question; how is the genome size of microorganisms distributed in the Baltic
Sea. Irrespective of the method used, the average estimated genome size in sediments is
larger than in the water column.

In our available homogeneous metadata for all datasets, we could not find a clear
answer of why benthic genome size is bigger than pelagic genome size. While there is
accompanying information on DOC for pelagic metagenomes and organic carbon for benthic
metagenomes, they use methods and metrics that are non-comparable. Luckily a previous
study has compiled information on organic carbon stocks in the Baltic Sea (Scheffold and
Hense, 2020): by collecting information from many different studies and years, they show
that in average the top 10 cm of sediments contain between 2 and 4 times more organic
carbon per area than the water column in the Baltic Sea. More organic carbon available for
Bacteria and Archaea would also pose a lower pressure on the genome to streamline. From
the results of our study, we hypothesize that one of the reasons pelagic microbial genomes are
smaller than benthic microbial genomes is the difference in organic carbon available. After
all, a positive correlation between genome size and nutrient concentration has been shown
before (Allen et al., 2012; Aylward and Santoro, 2020; Mende et al., 2017).

Brackish pelagic microorganisms tend to show smaller genome sizes than marine
and freshwater. We compare the average estimated genome size of pelagic Baltic Sea
microorganisms to previously published genome size estimations (Rodríguez-Gijón et al.,
2022). This comparison includes all taxonomic groups found in three large datasets that
includes 4051 freshwater representative MAGs, 2118 marine representative MAGs and 341
pelagic representative MAGs from the Baltic Sea (completeness >75%) (Alneberg et al.,
2020; Buck et al., 2021; Nayfach et al., 2021).We found that the average estimated genome

size of the brackish pelagic MAGs (2.96 Mbp) is lower than marine MAGs (3.10 Mbp), and

these two smaller than freshwater MAGs (3.48 Mbp) (t-test, $p < 0.01$) (**Figure 4A**).

We then subset all MAGs into phyla and focus on the most common from aquatic

environments: phyla Actinobacteriota, Bacteroidota, Cyanobacteria and Proteobacteria

(classes Alpha and Gammaproteobacteria) (**Figure 4B-F**). We test if the genome size

differences are consistent across phyla. We observe that only phylum Bacteroidota follows

the same trend as the full dataset in the average estimated genome size (**Figure 4C**). On the

other hand, MAGs from phylum Actinobacteriota present the biggest genome sizes for

marine environments, while no difference on average genome size is observed between

freshwater and brackish (**Figure 4B**). This result updates previous observations on aquatic

Actinobacteriota genome size variation (Hugerth et al., 2015). Opposite to Actinobacteriota,

MAGs from phylum Cyanobacteria show the smallest average genome size for marine

environments, while we do not observe statistical differences between brackish and

freshwater MAGs (**Figure 4D**). Similar trends were observed for isolates and MAGs of

picocyanobacteria (Cabello-Yeves et al., 2022; Sánchez-Baracaldo et al., 2019). However, it

is important to remark that DNA extraction, assembly, binning and/or quality check of

aquatic cyanobacterial MAGs is still a big challenge that needs to be addressed

(**Supplementary Material 3**) (Alvarenga et al., 2017; Kim Tiam et al., 2019). All in all,

these results support that both environment (Rodríguez-Gijón et al., 2022) and taxonomy

(Martinez-Gutierrez and Aylward, 2022) impact on existent pelagic genome size

distributions.

Streamlining may affect certain functional categories in Bacteria. We selected

the bacterial MAGs with >90% completeness for metabolic annotation and analyze which

functional categories correlate with genome size in brackish sediments and water column

(**Figure 5 A-R** and **Supplementary Material 2**). Functional categories included different but

related metabolic pathways (KEGG modules), and each module comprises multiple specific

metabolic reactions (module steps) (Zhou et al., 2022). In our results, we observe two

different patterns: negative correlation between estimated genome size and number of module

steps per Mbp, or the almost absence of the module steps (**Figure 5**). For the core functions

amino acid metabolism, aminoacyl tRNAs and central carbohydrate metabolism, we observe

a higher number of module steps per Mbp at lower genome sizes in both environments both

in pelagic and benthic MAGs (**Figures 5A, D** & **J**). Genome size does not explain the

number of module steps per Mbp in all functional categories and metabolisms, such as drug
resistance and transport systems (**Supplementary Material 2F** & **N**). These results suggest
that streamlining of genomes select for specific functions and not the whole genome. This
can be explained by the Black Queen Hypothesis (Morris et al., 2012), where only functions
that are more costly than beneficial are lost. The loss of those specific functions has
consequential long-lasting metabolic partnerships within the community (Mas et al., 2016).

When considering the total number of module steps, we observe that sediments code
for a higher number of module steps than pelagic Bacteria in amino acid metabolism,
aminoacyl tRNAs, carbon fixation, nitrogen and sulfur metabolism (**Figures 5B, E, H, N &**
Q) (t-test, $p < 0.01$). This could be the result of a cofounding effect of genome size:
microorganisms from sediments have bigger genome sizes than those in the water column
(**Figure 2**) and therefore, code for a higher number of functions. Hence, the total number of
module steps per Mbp in all six categories was analyzed. If streamlining affects all functional
categories similarly, coding density trends would be similar for all functional categories.
Indeed, we can observe that pelagic Bacteria present a greater number of module steps per
Mbp than sediment Bacteria in amino acid metabolism (**Figure 5C**), aminoacyl tRNAs
(**Figure 5F**), carbon fixation (**Figure 5I**) and central carbohydrate metabolism (**Figure 5L**).
However, nitrogen metabolism shows the opposite trend: sediment Bacteria have a greater
number of module steps per Mbp than water column Bacteria (**Figure 5O**). Literature report
that Baltic sediments contain about 95% of the total pool of nitrogen while the water column
only 5%, hence the water column only carries a small part of the overall nitrogen cycle
(Lønborg and Markager, 2021). This indicates that benthic Bacteria potentially play a bigger
role than pelagic bacteria in nitrogen cycling of autotropic systems like the Baltic Sea (Albert
et al., 2021; Griffiths et al., 2017).

Moreover, although the majority of the MAGs presented at least 1 module step related
to drug resistance (96.97% sediment MAGs and 98.81% pelagic MAGs), bacteria from
sediments presented a higher number of antibiotic resistance module steps than pelagic
Bacteria. This applies both to total number of module steps and the number of module steps
per Mbp (**Supplementary Material 2**). Our results confirm that aquatic sediments are
reservoirs of antibiotic resistance genes (Guo et al., 2020; Marti et al., 2014). Just as
sediments harbor more than double the organic carbon than the water column, this allows

microbial genomes to have a bigger size and code for a higher number of genes. This allows
microbes to upkeep non-essential functions and allow metabolic reservoir in the sediments.

Baltic sediments and water column harbor various Bacteria with different
metabolic capabilities. From all >90% completeness MAGs, we selected only bacterial
phyla with five or more high-quality MAGs to observe how metabolism differs between
different taxa in Baltic sediments and water column (**Figure 6**).

We observed the presence of the genes *cdhH | cdhE | cooS* from the Wood-Ljungdhal
pathway exclusively in marine sediments (**Figure 6**), particularly in phyla Desulfobacterota,
Desulfobacterota E and Verrucomicrobiota (class Kiritimatiellae). This is a carbon fixation
pathway predominant in acetogenic bacteria found in anoxic conditions (Esposito et al.,
2019). Complementarily, we also find putative fermentation genes for acetogenesis (*acdA |*
ack | pta) to be widespread across taxa in Baltic sediments. This would explain the potential
success of acetogenic metabolism in brackish sediments (Lever, 2012). We also find that the
acs gene for acetate fermentation into acetyl-CoA is widely distributed in both sediments and
water column. These results support the common distribution of acetogens and the Wood-
Ljungdhal pathway in Baltic glacial sediments (Marshall et al., 2018).

No FeFe hydrogenases were detected, but different NiFe hydrogenases were spotted
to differ between environments: NiFe groups 3abd were detected mainly in pelagic bacteria,
while NiFe group 1 in sediments (**Figure 6**). NiFe group 1 hydrogenases could be playing a
vital role in nitrate (NO_3^-), sulfate (SO_4^{2-}) and iron (Fe^{3+}) reduction: these molecules can act
as acceptors of electrons coupled to H_2 oxidation in anoxic conditions (Peters et al., 2015).
Moreover, putative genes coding for the reduction of the above-mentioned molecules were
also detected on our sediment dataset (*napAB | narGH* for nitrate reduction, *aprA | sat* for
sulfate reduction, and iron reduction series genes). These results suggest a key role of
sediment Bacteria in sulfur, nitrogen, and iron cycling in the Baltic Sea. For example,
sulphate reducers such as Desulfobacterota found in our benthic MAGs collection, most
likely contribute to the release of Fe-bound phosphorus from sediments to the water column
(Sinkko, 2013).

Conclusions. Our analyses delve into genome size distributions across the sediments and
water column in the Baltic Sea, and its relationships with metabolic potential, taxonomic

identity and abiotic variables. Our key findings show that these three ecological and
evolutionary variables are linked to genome size and, while each of them may partially
explain the genome size patterns, it is the interplay between the three of them that drives
genome size variation in Bacteria in the Baltic Sea. We also provide the first comprehensive
overview of microbial MAGs comparing genome size variation across the three major pelagic
ecosystems, where brackish Bacteria have smaller estimated genome sizes than marine and
freshwater. Within the Baltic Sea, benthic Bacteria have the biggest estimated genome size.
As a future perspective, with the continuous progression of aquatic microbial ecology and the
development of new isolation, omics and bioinformatic techniques, we will be able to grasp a
more complete and unbiased view of genome sizes distribution in nature and, therefore,
microbial evolution and ecology.
382
383 <u>MATERIAL AND METHODS</u>
384
Baltic Sea metagenomes collection. For this study we compiled new and public
Baltic Sea metagenomes from the water column and the sediments. The final dataset
consisted of 219 metagenome samples from a wide range of locations in the Baltic Sea that
include 5 independent datasets (**Supplementary Table 1** and **Figure 1**). We compiled 108
sediment metagenomes, of which 59 were collected in 2019 and recently published (Broman
et al., 2022), and we collected 49 metagenomes from 2016 to 2018. The pelagic dataset
consists of 118 pelagic samples collected from 2011 to 2015 published in three different
studies (Alneberg et al., 2020, 2018; Larsson et al., 2014). All five datasets have abiotic
metadata of depth (m), salinity (PSU), temperature (C) and oxygen concentration (mg/L)
(**Supplementary Table 1**).
395
Environmental sampling. The top 2 cm of sediment was collected at soft bottom
clay-muddy habitats from 59 stations from north to south in the Baltic Sea in 2019, following
the sampling described in Broman et al., (2022) (**Supplementary Table 1** for coordinates).
Briefly, one sediment core was collected per station using a Kajak gravity corer (surface area:
50 cm^2, one core per station) and the top 0–2 cm layer was sliced into a 215 ml
polypropylene container (207.0215PP, Noax Lab, Sweden). The sediment was homogenized
and stored at -20°C on the boat, kept on an iced cooler without thawing for ~2 h during
transportation to the university, and finally stored again at -20 °C until DNA extraction.
Bottom water (~20 cm above the sediment surface) was collected at each station with a

Niskin bottle. This was followed by on deck measurements of bottom water temperature,

salinity, and dissolved O2 using a portable multimeter (HQ40D, Hach).

DNA extraction and sequencing. The sediment samples were thawed, homogenized,

and a subsample of 0.25 g was used for DNA extraction using the DNeasy PowerSoil kit

(Qiagen) according to the manufacturer's protocol. The quantity and quality of eluted DNA

were measured using NanoDrop One spectrophotometer and Qubit 2 (both by ThermoFisher

Scientific) to ensure that samples meet the minimum requirements for sequencing. The

samples were then sequenced at the Science for Life Laboratories facility on one NovaSeq

6000 S4 lanes using a 2×150 bp setup. Sequencing yielded on average 53.0 million reads

per sample.

MAGs collection. Assembling and binning of the 108 sediment metagenomes

resulted in 2248 bins. To obtain bins from metagenomes, we followed the

0053_metasssnake2 pipeline (https://github.com/moritzbuck/0053_metasssnake2) (v0.0.2). In

this pipeline we used Sourmash (Titus Brown and Irber, 2016) to create signatures, Megahit

(Li et al., 2016) to obtain single-sample assemblies and Metabat2 (Kang et al., 2019) for the

binning of the assemblies. We used default parameters throughout the whole pipeline. Quality

of the bins was assessed using CheckM (v1.1.3) (Parks et al., 2015): we used the typical

workflow (*lineage_wf*) with default parameters, and selected only bins with quality of

completeness >75% and contamination <5%. From all the bins, only 216 passed our quality

threshold and we named those MAGs (metagenome assembled genomes). All MAGs were

taxonomically classified using GTDB-tk (v1.5.0) (Chaumeil et al., 2019) according to the

GTDB classification (Parks et al., 2020). The quality of the MAGs belonging to phyla

Actinobacteriota and Patescibacteria were assessed separately using a custom set of marker

genes. Preliminary quality check of Actinobacteriota genomes in a publicly available

freshwater dataset (Buck et al., 2021) show that default parameters underestimate the quality

of the MAGs that are classified as Actinobacteriota compared to using a custom marker gene

set (**Supplementary Material 3**). We obtained the custom marker gene set using Ac1

circular genomes (Kang et al., 2017; Neuenschwander et al., 2018). In the case of phylum

Patescibacteria, we used a custom set of maker genes provided by CheckM (Brown et al.,

2015; Parks et al., 2015).

Complementarily, we used 1920 pelagic MAGs that were published (Alneberg et al.,

2020) and passed the >75% completeness and <5% contamination threshold. All high-quality

MAGs from the sediments and water column were dereplicated using fastANI (95% ANI

threshold as estimator of genetic unit) and mOTUlizer (v0.3.2) (Buck et al., 2022; Jain et al.,

2018). From the 216 >75% completeness MAGs, 56 were chosen as representatives. From

the 1920 pelagic MAGs, 340 were chosen as representatives. All genomic information for

pelagic and benthic MAGs is included in **Supplementary Table 2**.

Genome size analysis. We studied genome size in two different levels: entire

microbial community and bacterial/archaeal MAGs. To study differences in genome size

between Baltic sediments and water column at the community level, we first calculated the

average genome size (AGS) of the metagenomes using MicrobeCensus (v1.1.0) (Nayfach and

Pollard, 2015). MicrobeCensus estimates the AGS of a microbial community from

metagenomes by aligning reads to a set of single-copy genes that are widely distributed

across taxa to calculate their abundance, with highly accurate estimations (Pereira-Flores et

al., 2019). We excluded one of the pelagic datasets (Alneberg et al., 2018) due to the

presence of spike-in DNA. We used default parameters, but we set the number of reads

sampled to 10 million (*-n 10 000 000*). To estimate the genome size of the microbial MAGs,

we divided the MAGs assembly size by the completeness (provided by CheckM, ranging

from 0 to 1). To study genome size variation between our pelagic brackish dataset and other

major aquatic environments (freshwater and marine), we compiled the metagenomic

information from all pelagic MAGs from marine and freshwater environments (>75%

completeness and <5% contamination) from (Rodríguez-Gijón et al., 2022) (**Supplementary**

Table 3).

Metabolic annotation. To analyze the metabolic potential of sedimentary and pelagic

bacteria, we selected all MAGs with completeness >90% and contamination <5%. In total,

we obtained 99 MAGs from sediments and 1 241 MAGs from the water column. The

metabolic potential of sediment and pelagic MAGs was reconstructed using Prodigal

annotations (v2.6.3) (Hyatt et al., 2010). We used the resulting protein translation files to

predict biogeochemical and metabolic functional traits using METABOLIC (v4.0) (Zhou et

al., 2022). We used the METABOLIC-G script, using default settings.

Statistical analysis. We performed Wilcoxon non-parametric test to analyze if there were significant differences between pairs of boxplots (**Figures 2, 3** and **5**). Stars in boxplots indicate significant differences $p < 0.05$. In **Figures 3B** and **4** we performed Kruskal-Wallis test corrected with Benjamini-Hochberg, to test statistical differences between groups (different letters are the result of this non-parametric test; $p < 0.05$). We performed a ANOVA type II analysis to test the effect of abiotic factors and their interactions on the AGS, using the function *aov* from the R package *stats* v3.6.3 (R Core Team, 2020) (**Figure S1**). We obtained the correlation coefficients on scatterplots (**Figure S1** and **Figure 5**) to test the fit of our data to linear regressions using the function *stat_cor* from the R package *ggpubr* v0.4.0 (Kassambara, A., 2020).

Data availability. All metagenome datasets are available in public repositories under NCBI project accession number SRP077551 and ENA accession numbers PRJEB34883, PRJEB22997 and PRJEB41834. Specific accession numbers of all metagenomes are available in **Supplementary Table 1**. Pelagic MAGs can be found on project PRJEB34883 and benthic MAGs on project PRJNA891615. Assembly and binning of the dataset provided in this paper used scripts available at https://github.com/moritzbuck/0053_metasssnake2.

ACKNOWLEDGEMENTS

We thank Elias Broman for his help on the acquisition of the benthic dataset of metagenomes and metadata and Ola Svensson, Caroline Raymond and Jonas Gunnarsson for assistance during sampling. We also thank Juanita Gutiérrez-Valencia for her suggestions in statistical analysis.

This work was supported by SciLifeLab. The authors acknowledge support from the National Genomics Infrastructure in Stockholm funded by Science for Life Laboratory, the Knut and Alice Wallenberg Foundation and the Swedish Research Council, and SNIC/Uppsala Multidisciplinary Center for Advanced Computational Science for assistance with massively parallel sequencing and access to the UPPMAX computational infrastructure. Computational work and data handling were enabled by resources in the projects SNIC 2020/5-159, 2021/5-133, 2022/5-137, 2020-6-60 and 2022/6-77 provided by the Swedish National Infrastructure for Computing (SNIC) at UPPMAX, partially funded by the Swedish Research Council through grant agreement no. 2018-05973.

      AR-G and SLG conceptualized and designed research project. AR-G, MB and SLG

refined the project idea. AR-G, AFA and FJAN participated in data collection. AR-G and

DIS did the molecular work. AR-G and MB performed bioinformatic analysis to obtain

MAGs from sediment raw sequences. AR-G and SLG performed data analysis. AR-G and

SLG drafted the first manuscript. AR-G, DIS and SLG did literature searches. All authors

contributed to the writing and editing of the manuscript.

<u>CONFLICTS OF INTEREST</u>

The authors declare that there are no competing interests.

<u>REFERENCES</u>

Acinas, S.G., Sánchez, P., Salazar, G., Cornejo-Castillo, F.M., Sebastián, M., Logares, R.,
Royo-Llonch, M., Paoli, L., Sunagawa, S., Hingamp, P., Ogata, H., Lima-Mendez, G.,
Roux, S., González, J.M., Arrieta, J.M., Alam, I.S., Kamau, A., Bowler, C., Raes, J.,
Pesant, S., Bork, P., Agustí, S., Gojobori, T., Vaqué, D., Sullivan, M.B., Pedrós-Alió,
C., Massana, R., Duarte, C.M., Gasol, J.M., 2021. Deep ocean metagenomes
provide insight into the metabolic architecture of bathypelagic microbial communities.
Commun Biol 4, 604. https://doi.org/10.1038/s42003-021-02112-2
Albert, S., Bonaglia, S., Stjärnkvist, N., Winder, M., Thamdrup, B., Nascimento, F.J.A., 2021.
Influence of settling organic matter quantity and quality on benthic nitrogen cycling.
Limnol Oceanogr 66, 1882–1895. https://doi.org/10.1002/lno.11730
Allen, L.Z., Allen, E.E., Badger, J.H., McCrow, J.P., Paulsen, I.T., Elbourne, L.D.,
Thiagarajan, M., Rusch, D.B., Nealson, K.H., Williamson, S.J., Venter, J.C., Allen,
A.E., 2012. Influence of nutrients and currents on the genomic composition of
microbes across an upwelling mosaic. ISME J 6, 1403–1414.
https://doi.org/10.1038/ismej.2011.201
Alneberg, J., Bennke, C., Beier, S., Bunse, C., Quince, C., Ininbergs, K., Riemann, L.,
Ekman, M., Jürgens, K., Labrenz, M., Pinhassi, J., Andersson, A.F., 2020.
Ecosystem-wide metagenomic binning enables prediction of ecological niches from
genomes. Commun Biol 3, 119. https://doi.org/10.1038/s42003-020-0856-x
Alneberg, J., Sundh, J., Bennke, C., Beier, S., Lundin, D., Hugerth, L.W., Pinhassi, J.,
Kisand, V., Riemann, L., Jürgens, K., Labrenz, M., Andersson, A.F., 2018. BARM
and BalticMicrobeDB, a reference metagenome and interface to meta-omic data for
the Baltic Sea. Sci Data 5, 180146. https://doi.org/10.1038/sdata.2018.146
Alvarenga, D.O., Fiore, M.F., Varani, A.M., 2017. A Metagenomic Approach to
Cyanobacterial Genomics. Front. Microbiol. 8, 809.
https://doi.org/10.3389/fmicb.2017.00809
Aylward, F.O., Santoro, A.E., 2020. Heterotrophic Thaumarchaea with Small Genomes Are
Widespread in the Dark Ocean. mSystems 5, e00415-20,
/msystems/5/3/msys.00415-20.atom. https://doi.org/10.1128/mSystems.00415-20
Batut, B., Knibbe, C., Marais, G., Daubin, V., 2014. Reductive genome evolution at both
ends of the bacterial population size spectrum. Nat Rev Microbiol 12, 841–850.
https://doi.org/10.1038/nrmicro3331

Beam, J.P., Becraft, E.D., Brown, J.M., Schulz, F., Jarett, J.K., Bezuidt, O., Poulton, N.J.,
Clark, K., Dunfield, P.F., Ravin, N.V., Spear, J.R., Hedlund, B.P., Kormas, K.A.,
Sievert, S.M., Elshahed, M.S., Barton, H.A., Stott, M.B., Eisen, J.A., Moser, D.P.,
Onstott, T.C., Woyke, T., Stepanauskas, R., 2020. Ancestral Absence of Electron
Transport Chains in Patescibacteria and DPANN. Front. Microbiol. 11, 1848.
https://doi.org/10.3389/fmicb.2020.01848
Biller, S.J., Berube, P.M., Lindell, D., Chisholm, S.W., 2015. Prochlorococcus: the structure
and function of collective diversity. Nat Rev Microbiol 13, 13–27.
https://doi.org/10.1038/nrmicro3378
Broman, E., Izabel-Shen, D., Rodríguez-Gijón, A., Bonaglia, S., Garcia, S.L., Nascimento,
F.J.A., 2022. Microbial functional genes are driven by gradients in sediment
stoichiometry, oxygen, and salinity across the Baltic benthic ecosystem. Microbiome
10, 126. https://doi.org/10.1186/s40168-022-01321-z
Brown, C.T., Hug, L.A., Thomas, B.C., Sharon, I., Castelle, C.J., Singh, A., Wilkins, M.J.,
Wrighton, K.C., Williams, K.H., Banfield, J.F., 2015. Unusual biology across a group
comprising more than 15% of domain Bacteria. Nature 523, 208–211.
https://doi.org/10.1038/nature14486
Buck, M., Garcia, S.L., Fernandez, L., Martin, G., Martinez-Rodriguez, G.A., Saarenheimo,
J., Zopfi, J., Bertilsson, S., Peura, S., 2021. Comprehensive dataset of shotgun
metagenomes from oxygen stratified freshwater lakes and ponds. Sci Data 8, 131.
https://doi.org/10.1038/s41597-021-00910-1
Buck, M., Mehrshad, M., Bertilsson, S., 2022. mOTUpan: a robust Bayesian approach to
leverage metagenome-assembled genomes for core-genome estimation. NAR
Genomics and Bioinformatics 4, lqac060. https://doi.org/10.1093/nargab/lqac060
Cabello-Yeves, P.J., Callieri, C., Picazo, A., Schallenberg, L., Huber, P., Roda-Garcia, J.J.,
Bartosiewicz, M., Belykh, O.I., Tikhonova, I.V., Torcello-Requena, A., De Prado,
P.M., Puxty, R.J., Millard, A.D., Camacho, A., Rodriguez-Valera, F., Scanlan, D.J.,
2022. Elucidating the picocyanobacteria salinity divide through ecogenomics of new
freshwater isolates. BMC Biol 20, 175. https://doi.org/10.1186/s12915-022-01379-z
Chaumeil, P.-A., Mussig, A.J., Hugenholtz, P., Parks, D.H., 2019. GTDB-Tk: a toolkit to
classify genomes with the Genome Taxonomy Database. Bioinformatics btz848.
https://doi.org/10.1093/bioinformatics/btz848
Chen, M.-Y., Teng, W.-K., Zhao, L., Hu, C.-X., Zhou, Y.-K., Han, B.-P., Song, L.-R., Shu, W.-
S., 2021. Comparative genomics reveals insights into cyanobacterial evolution and
habitat adaptation. ISME J 15, 211–227. https://doi.org/10.1038/s41396-020-00775-z
Coelho, L.P., Alves, R., del Río, Á.R., Myers, P.N., Cantalapiedra, C.P., Giner-Lamia, J.,
Schmidt, T.S., Mende, D.R., Orakov, A., Letunic, I., Hildebrand, F., Van Rossum, T.,
Forslund, S.K., Khedkar, S., Maistrenko, O.M., Pan, S., Jia, L., Ferretti, P.,
Sunagawa, S., Zhao, X.-M., Nielsen, H.B., Huerta-Cepas, J., Bork, P., 2022.
Towards the biogeography of prokaryotic genes. Nature 601, 252–256.
https://doi.org/10.1038/s41586-021-04233-4
Conley, D.J., Björck, S., Bonsdorff, E., Carstensen, J., Destouni, G., Gustafsson, B.G.,
Hietanen, S., Kortekaas, M., Kuosa, H., Markus Meier, H.E., Müller-Karulis, B.,
Nordberg, K., Norkko, A., Nürnberg, G., Pitkänen, H., Rabalais, N.N., Rosenberg, R.,
Savchuk, O.P., Slomp, C.P., Voss, M., Wulff, F., Zillén, L., 2009. Hypoxia-Related
Processes in the Baltic Sea. Environ. Sci. Technol. 43, 3412–3420.
https://doi.org/10.1021/es802762a
Esposito, A., Tamburini, S., Triboli, L., Ambrosino, L., Chiusano, M.L., Jousson, O., 2019.
Insights into the genome structure of four acetogenic bacteria with specific reference
to the Wood–Ljungdahl pathway. MicrobiologyOpen 8.
https://doi.org/10.1002/mbo3.938
Ghurye, J.S., 2016. Metagenomic Assembly: Overview, Challenges and Applications 10.
Giovannoni, S.J., 2017. SAR11 Bacteria: The Most Abundant Plankton in the Oceans. Annu.
Rev. Mar. Sci. 9, 231–255. https://doi.org/10.1146/annurev-marine-010814-015934

Giovannoni, S.J., 2005. Genome Streamlining in a Cosmopolitan Oceanic Bacterium.
Science 309, 1242–1245. https://doi.org/10.1126/science.1114057
Giovannoni, S.J., Cameron Thrash, J., Temperton, B., 2014. Implications of streamlining
theory for microbial ecology. ISME J 8, 1553–1565.
https://doi.org/10.1038/ismej.2014.60
Griffiths, J.R., Kadin, M., Nascimento, F.J.A., Tamelander, T., Törnroos, A., Bonaglia, S.,
Bonsdorff, E., Brüchert, V., Gårdmark, A., Järnström, M., Kotta, J., Lindegren, M.,
Nordström, M.C., Norkko, A., Olsson, J., Weigel, B., Žydelis, R., Blenckner, T.,
Niiranen, S., Winder, M., 2017. The importance of benthic-pelagic coupling for
marine ecosystem functioning in a changing world. Glob Change Biol 23, 2179–2196.
https://doi.org/10.1111/gcb.13642
Grote, J., Thrash, J.C., Huggett, M.J., Landry, Z.C., Carini, P., Giovannoni, S.J., Rappé,
M.S., 2012. Streamlining and Core Genome Conservation among Highly Divergent
Members of the SAR11 Clade. mBio 3, e00252-12.
https://doi.org/10.1128/mBio.00252-12
Guo, X., Zhao, S., Chen, Y., Yang, J., Hou, L., Liu, M., Yang, Y., 2020. Antibiotic resistance
genes in sediments of the Yangtze Estuary: From 2007 to 2019. Science of The Total
Environment 744, 140713. https://doi.org/10.1016/j.scitotenv.2020.140713
Herlemann, D.P., Labrenz, M., Jürgens, K., Bertilsson, S., Waniek, J.J., Andersson, A.F.,
2011. Transitions in bacterial communities along the 2000 km salinity gradient of the
Baltic Sea. ISME J 5, 1571–1579. https://doi.org/10.1038/ismej.2011.41
Hugerth, L.W., Larsson, J., Alneberg, J., Lindh, M.V., Legrand, C., Pinhassi, J., Andersson,
A.F., 2015. Metagenome-assembled genomes uncover a global brackish
microbiome. Genome Biol 16, 279. https://doi.org/10.1186/s13059-015-0834-7
Hyatt, D., Chen, G.-L., LoCascio, P.F., Land, M.L., Larimer, F.W., Hauser, L.J., 2010.
Prodigal: prokaryotic gene recognition and translation initiation site identification.
BMC Bioinformatics 11, 119. https://doi.org/10.1186/1471-2105-11-119
Jain, C., Rodriguez-R, L.M., Phillippy, A.M., Konstantinidis, K.T., Aluru, S., 2018. High
throughput ANI analysis of 90K prokaryotic genomes reveals clear species
boundaries. Nat Commun 9, 5114. https://doi.org/10.1038/s41467-018-07641-9
Kang, D.D., Li, F., Kirton, E., Thomas, A., Egan, R., An, H., Wang, Z., 2019. MetaBAT 2: an
adaptive binning algorithm for robust and efficient genome reconstruction from
metagenome assemblies. PeerJ 7, e7359. https://doi.org/10.7717/peerj.7359
Kang, I., Kim, S., Islam, Md.R., Cho, J.-C., 2017. The first complete genome sequences of
the acI lineage, the most abundant freshwater Actinobacteria, obtained by whole-
genome-amplification of dilution-to-extinction cultures. Sci Rep 7, 42252.
https://doi.org/10.1038/srep42252
Kassambara, A., 2020. ggpubr: "ggplot2" Based Publication Ready Plots.
Kim Tiam, S., Comte, K., Dalle, C., Duval, C., Pancrace, C., Gugger, M., Marie, B.,
Yéprémian, C., Bernard, C., 2019. Development of a new extraction method based
on high-intensity ultra-sonication to study RNA regulation of the filamentous
cyanobacteria Planktothrix. PLoS ONE 14, e0222029.
https://doi.org/10.1371/journal.pone.0222029
Konstantinidis, K.T., Braff, J., Karl, D.M., DeLong, E.F., 2009. Comparative Metagenomic
Analysis of a Microbial Community Residing at a Depth of 4,000 Meters at Station
ALOHA in the North Pacific Subtropical Gyre. Appl Environ Microbiol 75, 5345–5355.
https://doi.org/10.1128/AEM.00473-09
Kristensen, D.M., Mushegian, A.R., Dolja, V.V., Koonin, E.V., 2010. New dimensions of the
virus world discovered through metagenomics. Trends in Microbiology 18, 11–19.
https://doi.org/10.1016/j.tim.2009.11.003
Larsson, J., Celepli, N., Ininbergs, K., Dupont, C.L., Yooseph, S., Bergman, B., Ekman, M.,
2014. Picocyanobacteria containing a novel pigment gene cluster dominate the
brackish water Baltic Sea. ISME J 8, 1892–1903.
https://doi.org/10.1038/ismej.2014.35

Lever, M.A., 2012. Acetogenesis in the Energy-Starved Deep Biosphere – A Paradox?
660   Front. Microbio. 2. https://doi.org/10.3389/fmicb.2011.00284
661   Li, D., Luo, R., Liu, C.-M., Leung, C.-M., Ting, H.-F., Sadakane, K., Yamashita, H., Lam, T.-
662   W., 2016. MEGAHIT v1.0: A fast and scalable metagenome assembler driven by
663   advanced methodologies and community practices. Methods 102, 3–11.
664   https://doi.org/10.1016/j.ymeth.2016.02.020
665   Li, M., Wei, G., Shi, W., Sun, Z., Li, H., Wang, X., Gao, Z., 2018. Distinct distribution patterns
666   of ammonia-oxidizing archaea and bacteria in sediment and water column of the
667   Yellow River estuary. Sci Rep 8, 1584. https://doi.org/10.1038/s41598-018-20044-6
668   Lind, A.L., Pollard, K.S., 2021. Accurate and sensitive detection of microbial eukaryotes from
669   whole metagenome shotgun sequencing. Microbiome 9, 58.
670   https://doi.org/10.1186/s40168-021-01015-y
671   Lønborg, C., Markager, S., 2021. Nitrogen in the Baltic Sea: Long-term trends, a budget and
672   decadal time lags in responses to declining inputs. Estuarine, Coastal and Shelf
673   Science 261, 107529. https://doi.org/10.1016/j.ecss.2021.107529
674   Louca, S., Parfrey, L.W., Doebeli, M., 2016. Decoupling function and taxonomy in the global
675   ocean microbiome. Science 353, 1272–1277.
676   https://doi.org/10.1126/science.aaf4507
677   Lynch, M., 2006. Streamlining and Simplification of Microbial Genome Architecture 25.
678   Maistrenko, O.M., Mende, D.R., Luetge, M., Hildebrand, F., Schmidt, T.S.B., Li, S.S.,
679   Rodrigues, J.F.M., von Mering, C., Pedro Coelho, L., Huerta-Cepas, J., Sunagawa,
680   S., Bork, P., 2020. Disentangling the impact of environmental and phylogenetic
681   constraints on prokaryotic within-species diversity. ISME J.
682   https://doi.org/10.1038/s41396-020-0600-z
683   Marshall, I.P.G., Karst, S.M., Nielsen, P.H., Jørgensen, B.B., 2018. Metagenomes from deep
684   Baltic Sea sediments reveal how past and present environmental conditions
685   determine microbial community composition. Marine Genomics 37, 58–68.
686   https://doi.org/10.1016/j.margen.2017.08.004
687   Marti, E., Variatza, E., Balcazar, J.L., 2014. The role of aquatic ecosystems as reservoirs of
688   antibiotic resistance. Trends in Microbiology 22, 36–41.
689   https://doi.org/10.1016/j.tim.2013.11.001
690   Martinez-Gutierrez, C.A., Aylward, F.O., 2022. Genome size distributions in bacteria and
691   archaea are strongly linked to evolutionary history at broad phylogenetic scales.
692   PLoS Genet 18, e1010220. https://doi.org/10.1371/journal.pgen.1010220
693   Mas, A., Jamshidi, S., Lagadeuc, Y., Eveillard, D., Vandenkoornhuyse, P., 2016. Beyond the
694   Black Queen Hypothesis. ISME J 10, 2085–2091.
695   https://doi.org/10.1038/ismej.2016.22
696   Mende, D.R., Bryant, J.A., Aylward, F.O., Eppley, J.M., Nielsen, T., Karl, D.M., DeLong,
697   E.F., 2017. Environmental drivers of a microbial genomic transition zone in the
698   ocean's interior. Nat Microbiol 2, 1367–1373. https://doi.org/10.1038/s41564-017-
699   0008-3
700   Moran, N.A., Mira, A., 2001. The process of genome shrinkage in the obligate symbiont
701   Buchnera aphidicola. Genome Biol 2, research0054.1. https://doi.org/10.1186/gb-
702   2001-2-12-research0054
703   Morris, J.J., Lenski, R.E., Zinser, E.R., 2012. The Black Queen Hypothesis: Evolution of
704   Dependencies through Adaptive Gene Loss. mBio 3, e00036-12.
705   https://doi.org/10.1128/mBio.00036-12
706   Nayfach, S., Pollard, K.S., 2015. Average genome size estimation improves comparative
707   metagenomics and sheds light on the functional ecology of the human microbiome.
708   Genome Biol 16, 51. https://doi.org/10.1186/s13059-015-0611-7
709   Nayfach, S., Roux, S., Seshadri, R., Udwary, D., Varghese, N., Schulz, F., Wu, D., Paez-
710   Espino, D., Chen, I.-M., Huntemann, M., Palaniappan, K., Ladau, J., Mukherjee, S.,
711   Reddy, T.B.K., Nielsen, T., Kirton, E., Faria, J.P., Edirisinghe, J.N., Henry, C.S.,
712   Jungbluth, S.P., Chivian, D., Dehal, P., Wood-Charlson, E.M., Arkin, A.P., Tringe,
713   S.G., Visel, A., IMG/M Data Consortium, Abreu, H., Acinas, S.G., Allen, E., Allen,

M.A., Alteio, L.V., Andersen, G., Anesio, A.M., Attwood, G., Avila-Magaña, V., Badis,
Y., Bailey, J., Baker, B., Baldrian, P., Barton, H.A., Beck, D.A.C., Becraft, E.D.,
Beller, H.R., Beman, J.M., Bernier-Latmani, R., Berry, T.D., Bertagnolli, A.,
Bertilsson, S., Bhatnagar, J.M., Bird, J.T., Blanchard, J.L., Blumer-Schuette, S.E.,
Bohannan, B., Borton, M.A., Brady, A., Brawley, S.H., Brodie, J., Brown, S., Brum,
J.R., Brune, A., Bryant, D.A., Buchan, A., Buckley, D.H., Buongiorno, J., Cadillo-
Quiroz, H., Caffrey, S.M., Campbell, A.N., Campbell, B., Carr, S., Carroll, J., Cary,
S.C., Cates, A.M., Cattolico, R.A., Cavicchioli, R., Chistoserdova, L., Coleman, M.L.,
Constant, P., Conway, J.M., Mac Cormack, W.P., Crowe, S., Crump, B., Currie, C.,
Daly, R., DeAngelis, K.M., Denef, V., Denman, S.E., Desta, A., Dionisi, H.,
Dodsworth, J., Dombrowski, N., Donohue, T., Dopson, M., Driscoll, T., Dunfield, P.,
Dupont, C.L., Dynarski, K.A., Edgcomb, V., Edwards, E.A., Elshahed, M.S.,
Figueroa, I., Flood, B., Fortney, N., Fortunato, C.S., Francis, C., Gachon, C.M.M.,
Garcia, S.L., Gazitua, M.C., Gentry, T., Gerwick, L., Gharechahi, J., Girguis, P.,
Gladden, J., Gradoville, M., Grasby, S.E., Gravuer, K., Grettenberger, C.L.,
Gruninger, R.J., Guo, J., Habteselassie, M.Y., Hallam, S.J., Hatzenpichler, R.,
Hausmann, B., Hazen, T.C., Hedlund, B., Henny, C., Herfort, L., Hernandez, M.,
Hershey, O.S., Hess, M., Hollister, E.B., Hug, L.A., Hunt, D., Jansson, J., Jarett, J.,
Kadnikov, V.V., Kelly, C., Kelly, R., Kelly, W., Kerfeld, C.A., Kimbrel, J., Klassen, J.L.,
Konstantinidis, K.T., Lee, L.L., Li, W.-J., Loder, A.J., Loy, A., Lozada, M., MacGregor,
B., Magnabosco, C., Maria da Silva, A., McKay, R.M., McMahon, K., McSweeney,
C.S., Medina, M., Meredith, L., Mizzi, J., Mock, T., Momper, L., Moran, M.A., Morgan-
Lang, C., Moser, D., Muyzer, G., Myrold, D., Nash, M., Nesbø, C.L., Neumann, A.P.,
Neumann, R.B., Noguera, D., Northen, T., Norton, J., Nowinski, B., Nüsslein, K.,
O'Malley, M.A., Oliveira, R.S., Maia de Oliveira, V., Onstott, T., Osvatic, J., Ouyang,
Y., Pachiadaki, M., Parnell, J., Partida-Martinez, L.P., Peay, K.G., Pelletier, D., Peng,
X., Pester, M., Pett-Ridge, J., Peura, S., Pjevac, P., Plominsky, A.M., Poehlein, A.,
Pope, P.B., Ravin, N., Redmond, M.C., Reiss, R., Rich, V., Rinke, C., Rodrigues,
J.L.M., Rodriguez-Reillo, W., Rossmassler, K., Sackett, J., Salekdeh, G.H., Saleska,
S., Scarborough, M., Schachtman, D., Schadt, C.W., Schrenk, M., Sczyrba, A.,
Sengupta, A., Setubal, J.C., Shade, A., Sharp, C., Sherman, D.H., Shubenkova,
O.V., Sierra-Garcia, I.N., Simister, R., Simon, H., Sjöling, S., Slonczewski, J., Correa
de Souza, R.S., Spear, J.R., Stegen, J.C., Stepanauskas, R., Stewart, F., Suen, G.,
Sullivan, M., Sumner, D., Swan, B.K., Swingley, W., Tarn, J., Taylor, G.T., Teeling,
H., Tekere, M., Teske, A., Thomas, T., Thrash, C., Tiedje, J., Ting, C.S., Tully, B.,
Tyson, G., Ulloa, O., Valentine, D.L., Van Goethem, M.W., VanderGheynst, J.,
Verbeke, T.J., Vollmers, J., Vuillemin, A., Waldo, N.B., Walsh, D.A., Weimer, B.C.,
Whitman, T., van der Wielen, P., Wilkins, M., Williams, T.J., Woodcroft, B., Woolet,
J., Wrighton, K., Ye, J., Young, E.B., Youssef, N.H., Yu, F.B., Zemskaya, T.I., Ziels,
R., Woyke, T., Mouncey, N.J., Ivanova, N.N., Kyrpides, N.C., Eloe-Fadrosh, E.A.,
2021. A genomic catalog of Earth's microbiomes. Nat Biotechnol 39, 499–509.
https://doi.org/10.1038/s41587-020-0718-6
Neuenschwander, S.M., Ghai, R., Pernthaler, J., Salcher, M.M., 2018. Microdiversification in
genome-streamlined ubiquitous freshwater Actinobacteria. ISME J 12, 185–198.
https://doi.org/10.1038/ismej.2017.156
Nielsen, D.A., Fierer, N., Geoghegan, J.L., Gillings, M.R., Gumerov, V., Madin, J.S., Moore,
L., Paulsen, I.T., Reddy, T.B.K., Tetu, S.G., Westoby, M., 2021. Aerobic bacteria and
archaea tend to have larger and more versatile genomes. Oikos 130, 501–511.
https://doi.org/10.1111/oik.07912
Parks, D.H., Chuvochina, M., Chaumeil, P.-A., Rinke, C., Mussig, A.J., Hugenholtz, P., 2020.
A complete domain-to-species taxonomy for Bacteria and Archaea. Nat Biotechnol
38, 1079–1086. https://doi.org/10.1038/s41587-020-0501-8
Parks, D.H., Imelfort, M., Skennerton, C.T., Hugenholtz, P., Tyson, G.W., 2015. CheckM:
assessing the quality of microbial genomes recovered from isolates, single cells, and
metagenomes. Genome Res. 25, 1043–1055. https://doi.org/10.1101/gr.186072.114

Pereira-Flores, E., Glöckner, F.O., Fernandez-Guerra, A., 2019. Fast and accurate average
genome size and 16S rRNA gene average copy number computation in
metagenomic data. BMC Bioinformatics 20, 453. https://doi.org/10.1186/s12859-019-
3031-y
Peters, J.W., Schut, G.J., Boyd, E.S., Mulder, D.W., Shepard, E.M., Broderick, J.B., King,
P.W., Adams, M.W.W., 2015. [FeFe]- and [NiFe]-hydrogenase diversity, mechanism,
and maturation. Biochimica et Biophysica Acta (BBA) - Molecular Cell Research
1853, 1350–1369. https://doi.org/10.1016/j.bbamcr.2014.11.021
R Core Team, 2020. R: A language and environment for statistical computing.
Rodríguez-Gijón, A., Nuy, J.K., Mehrshad, M., Buck, M., Schulz, F., Woyke, T., Garcia, S.L.,
2022. A genomic perspective across Earth's microbiomes reveals that genome size
in Archaea and Bacteria is linked to ecosystem type and trophic strategy. Frontiers in
Microbiology 12. https://doi.org/10.3389/fmicb.2021.761869
Salcher, M.M., Schaefle, D., Kaspar, M., Neuenschwander, S.M., Ghai, R., 2019. Evolution
in action: habitat transition from sediment to the pelagial leads to genome
streamlining in Methylophilaceae. ISME J 13, 2764–2777.
https://doi.org/10.1038/s41396-019-0471-3
Sánchez-Baracaldo, P., Bianchini, G., Di Cesare, A., Callieri, C., Chrismas, N.A.M., 2019.
Insights Into the Evolution of Picocyanobacteria and Phycoerythrin Genes (mpeBA
and cpeBA). Front. Microbiol. 10, 45. https://doi.org/10.3389/fmicb.2019.00045
Scanlan, D.J., Ostrowski, M., Mazard, S., Dufresne, A., Garczarek, L., Hess, W.R., Post,
A.F., Hagemann, M., Paulsen, I., Partensky, F., 2009. Ecological Genomics of
Marine Picocyanobacteria. MMBR 73, 249–299.
https://doi.org/10.1128/MMBR.00035-08
Scheffold, M.I.E., Hense, I., 2020. Quantifying Contemporary Organic Carbon Stocks of the
Baltic Sea Ecosystem. Front. Mar. Sci. 7, 571956.
https://doi.org/10.3389/fmars.2020.571956
Simonsen, A.K., 2022. Environmental stress leads to genome streamlining in a widely
distributed species of soil bacteria. ISME J 16, 423–434.
https://doi.org/10.1038/s41396-021-01082-x
Sinkko, H., 2013. Sediment bacterial communities in nutrient cycling and in the history of the
Baltic Sea (Doctoral dissertation). University of Helsinki, Helsinki, Finland.
Swan, B.K., Tupper, B., Sczyrba, A., Lauro, F.M., Martinez-Garcia, M., Gonzalez, J.M., Luo,
H., Wright, J.J., Landry, Z.C., Hanson, N.W., Thompson, B.P., Poulton, N.J.,
Schwientek, P., Acinas, S.G., Giovannoni, S.J., Moran, M.A., Hallam, S.J.,
Cavicchioli, R., Woyke, T., Stepanauskas, R., 2013. Prevalent genome streamlining
and latitudinal divergence of planktonic bacteria in the surface ocean. Proceedings of
the National Academy of Sciences 110, 11463–11468.
https://doi.org/10.1073/pnas.1304246110
Titus Brown, C., Irber, L., 2016. sourmash: a library for MinHash sketching of DNA. JOSS 1,
27. https://doi.org/10.21105/joss.00027
van Ham, R.C.H.J., Kamerbeek, J., Palacios, C., Rausell, C., Abascal, F., Bastolla, U.,
Fernández, J.M., Jiménez, L., Postigo, M., Silva, F.J., Tamames, J., Viguera, E.,
Latorre, A., Valencia, A., Morán, F., Moya, A., 2003. Reductive genome evolution in
Buchnera aphidicola. Proc. Natl. Acad. Sci. U.S.A. 100, 581–586.
https://doi.org/10.1073/pnas.0235981100
Zhou, Z., Tran, P.Q., Breister, A.M., Liu, Y., Kieft, K., Cowley, E.S., Karaoz, U.,
Anantharaman, K., 2022. METABOLIC: high-throughput profiling of microbial
genomes for functional traits, metabolism, biogeochemistry, and community-scale
functional networks. Microbiome 10, 33. https://doi.org/10.1186/s40168-021-01213-8
Zhou, Z., Tran, P.Q., Kieft, K., Anantharaman, K., 2020. Genome diversification in globally
distributed novel marine Proteobacteria is linked to environmental adaptation. ISME J
14, 2060–2077. https://doi.org/10.1038/s41396-020-0669-4
822

Figure 1. Overview of the sampling locations and average genome size (AGS) of

metagenomes (108 from sediments, in dark blue, and 111 from the water column, in light

blue). This figure shows the geographic location of all metagenomes used in this study. For

exact coordinates see Supplementary Table 1. Shape type indicates the reference and shape

size indicates the AGS of the given metagenome.

Figure 2. Boxplots showing the AGS distribution of Baltic metagenomes. **Panel A** indicates

the AGS distribution in both water column and sediments. **Panel B** indicates the AGS

distribution in metagenomes from sediments across the oxygen gradient (two groups, from 0

to 2 and from 2 to 12.45 mg/L). Stars in both panels indicate significant differences $p < 0.05$

(Wilcoxon non-parametric test).

Table 1. Summary of all 56 sediment and 340 water column representative MAGs (95%

average nucleotide identity) with >75% completeness. Table includes phyla, environment

(either water column or sediments), number of representative genomes (*n*), smallest and

largest estimated genome sizes (Mbp) observed for each phylum, average GC content (%)

and average coding density (%). When only one MAG is indicated, estimated genome size of

that MAG is expressed in the sixth column.

Figure 3. Overview of the estimated genome size in Bacteria and Archaea obtained from

Baltic Sea sediments (dark blue) and water column (light blue) using only the 397

representative MAGs (95% average nucleotide identity) with >75% completeness. **Panel A**

shows the genome size distribution of Archaea and Bacteria obtained from Baltic water

column and sediments for a total of 397 representative genomes. **Panel B** shows the

estimated genome size per domain and environment. Different letters indicate significant

differences $p < 0.05$ (Kruskal-Wallis non-parametric test; multiple testing corrected with

Benjamini-Hochberg). **Panel C** shows the estimated genome size per phylum. We selected

only phyla with at least 2 MAGs in each environment. **Panel D** shows the estimated genome

size per order. We selected only orders with at least 2 MAGs in each environment. Numbers

below the boxes indicate the number of MAGs per environment. Stars in Panel C and D

indicate significant differences $p < 0.05$ (Wilcoxon non-parametric test).

Figure 4. Overview of the estimated genome size of pelagic Bacteria and Archaea obtained
from Baltic Sea (blue), freshwater (green) and marine (yellow) using only representative
MAGs (calculates using 95% average nucleotide identity) with >75% completeness. We
compare all representative MAGs (**Panel A**), only phylum Actinobacteriota (**Panel B**),
phylum Bacteroidota (**Panel C**), phylum Cyanobacteria (**Panel D**), class Alphaproteobacteria
(**Panel E**) and class Gammaproteobacteria (**Panel F**). Different letters indicate significant
differences $p < 0.05$ (Kruskal-Wallis non-parametric test; multiple testing corrected with
Benjamini-Hochberg). Numbers below the boxes indicate the number of MAGs per
environment.

Figure 5. Overview of the presence of module steps for six metabolic categories: amino acid
metabolism (**Panels A-C**), aminoacyl tRNAs (**Panels D-F**), carbon fixation (**Panels G-I**),
central carbohydrate metabolism (**Panels J-L**), nitrogen metabolism (**Panels M-O**) and sulfur
metabolism (**Panels P-R**). We used all bacterial MAGs with very-high quality (>90%
completeness and <5% contamination), from both environments (99 MAGs from sediments
and 1215 MAGs from water column). Stars in boxplots indicate significant differences $p <$
0.05 (Wilcoxon non-parametric test). Panels B, E, H, K, N and Q indicate total number of
module steps. Panels C, F, I, L, O and R indicate the number of module steps per Mbp.

Figure 6. Metabolic potential of all high-quality MAGs (>90% completeness and <5%
contamination). We selected all phyla with at least 5 MAGs, and then divided by class.
Boxes on the top of the figure indicate environment (pelagic in light blue and sediments in
dark blue), and inner numbers indicate the number of MAGs per category. In the heatmap,
white squares indicate absence of a given gene in all MAGs, and the darkest purple indicates
presence in all of them (gradient scale on the bottom-left part of the figure for reference).

SUPPLEMENTARY MATERIAL

Figure S1. Overview of the average genome size (AGS) of metagenomes across gradients of
salinity and depth.

Supplementary Material 1. Statistical analysis of the effect of abiotic variables and their
interactions on the average genome size (AGS) of full Baltic microbial communities (**Figure**
1). Tables show the results of ANOVA type II analysis for both sediments and water column.

Supplementary Material 2. Overview of the presence of module steps for several metabolic

categories, complementing **Figure 3**. We also provide a list of all functional categories used

with the different module categories included, and the number of module steps used in the

analysis.

Supplementary Material 3. Table with all Actinobacteriota genomes used to calculate

marker genes to use in CheckM quality assessment. We include information on GenBank

Accession, lake, tribe, assembly size (Mbp), GC content (%) and reference. Two figures are

included. First a boxplot showing the differences on completeness of the of the 8 phyla with

most bins in the StratfreshDB database. We have included the completeness estimations

using both default CheckM parameters and specific set of marker genes in both phyla

Patescibacteria and Actinobacteriota. Stars indicate significant differences p < 0.05

(Wilcoxon non-parametric test). We also provide a figure showing the markers used by

CheckM with default parameters for the 8 analyzed phyla.

Supplementary Table 1. Table with information of all metagenomes used in this research

project, including: sample run in NCBI/ENA, estimated average genome size (Mbp), depth

(m), salinity (PSU), temperature (C) and oxygen concentration (mg/L), sample material

processing (μm), environment, latitude, longitude, study accession, online database, sample

accession and reference.

Supplementary Table 2. Table with information of all metagenome assembled genomes

(MAGs) used in this research project, including: bin id, mOTU, completeness (%),

contamination (%), GC (%), coding density (%), bin length, estimated genome size (Mbp),

environment, domain, phylum, class, order, family, genus and species.

Supplementary Table 3. Table with information of all pelagic metagenome assembled

genomes (MAGs) used to compare genome size across major pelagic environments (marine,

freshwater and brackish). All brackish MAGs belong (Alneberg et al., 2020), while the

genomic information for marine and freshwater MAGs was retrieved from a previous study

(Rodríguez-Gijón et al., 2022) that compiled them from (Buck et al., 2021; Nayfach et al.,

2021). We include bin id, completeness (%), contamination (%), bin length, estimated

genome size (Mbp), ecosystem type, domain, phylum, class, order, family, genus and species.

Aminoacid metabolism

Aminoacyl tRNAs

Carbon fixation

Central carbohydrate metabolism

Nitrogen metabolism

Sulfur metabolism

Phyla	Environment	n	Smallest estimated genome size (Mbp)	Largest estimated genome size (Mbp)	Average GC (%)	Average coding density (%)
Actinobacteriota	Sediments	6	2.84	3.7	67.55	91.63
	Water column	59	1.25	4.53	54.23	93.74
Bacteroidota	Sediments	2	3	4.61	41.08	89.69
	Water column	86	1.38	4.71	40.7	92.97
Chloroflexota	Sediments	3	2.38	5.19	63.57	91.14
	Water column	5	1.25	5.91	57.42	90.26
Desulfobacterota	Sediments	8	2.71	4.66	49.32	84.55
	Water column	1	-	2.73	54.4	87.06
Gemmatimonadota	Sediments	1	-	2.85	66.66	93.99
	Water column	1	-	4.32	62.47	91.21
Myxococcota	Sediments	3	4.95	6.74	63.54	90.84
	Water column	1	-	7.66	66.33	90.86
Planctomycetota	Sediments	1	-	5.09	66.06	90.33
	Water column	29	3.18	**9.69**	59.01	88.23
Proteobacteria (Alfa)	Sediments	1	-	3.43	59.14	91.26
	Water column	46	**1.08**	5.92	51.88	91.98
Proteobacteria (Gamma)	Sediments	10	1.93	4.8	59.56	90.96
	Water column	54	1.18	4.19	50.48	92.09
Verrucomicrobiota	Sediments	3	2.97	3.69	56.16	89.4
	Water column	25	1.98	6.82	57.36	90.41
Thermoplasmatota	Sediments	1	-	1.51	54.16	91.35
	Water column	2	2.35	2.68	41.63	93.32
Thermoproteota	Sediments	3	1.42	1.58	35.32	89.34
	Water column	2	1.42	1.48	31.86	89.96
Acidobacteriota	Sediments	5	4.13	5.64	65.98	91.6
Desulfobacterota D	Sediments	1	-	2.33	45.15	88.3
Desulfobacterota E	Sediments	2	2.35	2.74	64.01	91.94
Nitrospirota	Sediments	3	2.86	4	54.01	86.72
Omnitrophota	Sediments	1	-	1.47	41.17	91.99
Zixibacteria	Sediments	1	-	3.68	41.12	90.73
Micrarchaeota	Sediments	1	-	1.14	48.22	93.23
Bdellovibrionota	Water column	1	-	3.73	49.16	93.45
Bdellovibrionota C	Water column	1	-	3.33	47.41	85.46
Campylobacterota	Water column	2	2.6	3.3	37.02	93.25
Cyanobacteria	Water column	16	1.98	6.03	51.8	88.45

Firmicutes	Water column	3	1.09	1.16	38.25	93.69
Krumholzibacteriota	Water column	1	-	3.34	63.78	92.99
Marinisomatota	Water column	2	2.62	3.11	39.47	91.47
Nitrospinota	Water column	2	3.18	3.35	47.22	87.87
Nanoarchaeota	Water column	1	-	1.93	29.77	92.86