Linking prokaryotic genome size variation to metabolic potential and

environment

<u>RUNNING TITLE</u>: Genome size links to metabolic potential

<u>AUTHORS</u>

Alejandro Rodríguez-Gijón[$1,2], Moritz Buck[3], Anders F. Andersson[2,4], Dandan Izabel-Shen[1],

Francisco J. A. Nascimento[1,5], Sarahi L. Garcia[$1,2]

$ Correspondent authors

alejandro.rgijon@gmail.com

sarahi.garcia@su.se

1 Department of Ecology, Environment and Plant Sciences, Stockholm University,

Stockholm 106 91, Sweden

2 Science for Life Laboratory, Stockholm, Sweden

3 Department of Aquatic Sciences and Assessment, Swedish University of Agricultural

Sciences, Uppsala, Sweden

4 Department of Gene Technology, School of Engineering Sciences in Chemistry,

Biotechnology and Health, KTH Royal Institute of Technology, Stockholm, Sweden

5 Baltic Sea Centre, Stockholm University, Stockholm, Sweden

<u>Keywords</u>: microbial ecology, genome size, Bacteria, Baltic Sea

ABSTRACT (250 words)

While theories and models have appeared to explain genome size as a result of
evolutionary processes, little work has shown that genome sizes carry ecological signatures.
Our work delves into the ecological implications of microbial genome size variation in benthic
and pelagic habitats across environmental gradients of the brackish Baltic Sea. While depth is
significantly associated with genome size in benthic and pelagic brackish metagenomes,
salinity is only correlated to genome size in benthic metagenomes. Overall, we confirm that
prokaryotic genome sizes in Baltic sediments (3.47 Mbp) are significantly bigger than in the
water column (2.96 Mbp). While benthic genomes have a higher number of functions than
pelagic genomes, the smallest genomes coded for a higher number of module steps per Mbp
for most of the functions irrespective of their environment. Some examples of this functions
are amino acid metabolism and central carbohydrate metabolism. However, we observed that
nitrogen metabolism was almost absent in pelagic genomes and was mostly present in benthic
genomes. Finally, we also show that Bacteria inhabiting Baltic sediments and water column
not only differ in taxonomy, but also in their metabolic potential, such as the Wood-Ljungdahl
pathway or the presence of different hydrogenases. Our work shows how microbial genome
size is linked to abiotic factors in the environment, metabolic potential and taxonomic identity
of Bacteria and Archaea within aquatic ecosystems.

INTRODUCTION

Genomes in Bacteria and Archaea are information-rich (1), and known to range in size
from 0.1 to 16 million base pairs (Mbp) (2). They can vary over evolutionary time through
genomic expansions and contractions via genetic drift, selection, homologous recombination,
deletions and insertions (3–9). Moreover, evolutionary studies have revealed extremely rapid
and highly variable flux of genes (10) with evolutionary forces acting on individual genes (5).
With all these evolutionary forces acting on the genes, we can presume that gene content and,
by consequence, genome size has an ecological meaning. Indeed, genome size has been linked
to phylogenetic history (11,12), lifestyle such as free-living, particle attached or host-associated
(4,13,14), or environment such as marine, freshwater, different types of sediments or different
hosts in host-associated microorganisms (2,15,16). We aim to delve into the ecological
implications of genome size in aquatic microorganisms, with emphasis on metabolic potential
using a brackish environment as a model.

In the last decade, aquatic microorganisms have been extensively sampled and now have a large representation in genomic and metagenomic datasets (17). Their genome size spans from 0.5 to 15 Mbp with an average of 3.1 Mbp (2). Aquatic environments are heterogeneous and many different abiotic factors, such as salinity and depth, could be linked to microbial genome size variation. For example, pelagic microbes inhabiting deep marine environments are estimated to present bigger genome sizes than those in shallow marine waters (18,19). Within freshwater ecosystems, isolates from the family Methylophilaceae (class Gammaproteobacteria) show a smaller genome size for pelagic than for sediment dwellers (20). Additionally, two studies have shown that marine Cyanobacteria have smaller genome sizes than freshwater (21,22). This literature already provides some insights on how genome size is linked to the environmental heterogeneity of freshwater and marine ecosystems. Yet, the studies are limited either to a specific marine station, or specific microbial lineages and it remains a question if these findings are applicable more widely. Moreover, genome size variation in the brackish realm remains debated: Actinobacteria in the brackish Baltic Sea show bigger genome sizes than in freshwaters (23), while picocyanobacteria show the opposite trend (22). Additionally, further research must be done to elucidate the link between genome size and abiotic factors within aquatic environments, particularly brackish water bodies.

In the link between abiotic factors in the environment and genome size, gene content is selected accordingly. Metagenomic studies show that a vast majority of the genes in Bacteria and Archaea are specific to particular environments, whereas very few genes are being shared between environments (24). This remarks how relevant is the relationship between niche specificity and lineage specific functional traits (25). These functional capabilities also differ between water column and sediments in Bacteria and Archaea in both marine (26,27) and freshwater environments (28). Since gene repertoires and genome size are related, they must be considered together with environmental gradients to better understand niche-specificity and the ecology of different prokaryotic lineages.

In this research article, we provide a comprehensive analysis to show the ecological implications of genome size variation of Bacteria and Archaea in pelagic and benthic communities in the Baltic Sea. Specifically, we investigate; i) how genome size varies across abiotic factors and taxonomic lineages of Bacteria and Archaea from sediments and water column, ii) what relationship can be found between genome size and the number of metabolic

capabilities in Bacteria in the Baltic Sea, and iii) which taxa and metabolic pathways contrast
between pelagic and benthic Bacteria. To achieve this, we selected 111 pelagic and 59 benthic
metagenomic samples that were previously published, and we provide one new and
unpublished benthic dataset with 49 metagenomic samples. We use these 219 metagenomes to
study genome size distribution across sediments and water column in the Baltic Sea (**Figure 1**
and **Table S1**). For this, we use two different approaches to study genome size: the estimated
average genome size (AGS) per metagenomic sample, and the estimated genome size of
bacterial and archaeal metagenome-assembled genomes (MAGs). Our results show that
Bacteria and Archaea with larger genomes in the sediments present lower coding density than
the smaller genomes in the water column. We also find that microorganisms inhabiting the
Baltic benthos have more metabolically-versatile genomes than pelagic prokaryotes, which
mean they code for a wider range of metabolic capabilities. Finally, we also find that functions
involved in the nitrogen metabolism are disproportionately more detected in benthic bacteria
in the Baltic Sea.

RESULTS AND DISCUSSION

While depth is significantly associated with genome size in benthic and pelagic
metagenomes, salinity is only correlated to genome size in benthic metagenomes. First, we
calculated the average genome size (AGS) of metagenomes across the latitudinal gradient of
the Baltic Sea comparing benthic and pelagic metagenomes. We observed that sediment-
dwelling microbial communities present significantly larger AGS (mean = 6.01 Mbp, n = 108)
than pelagic communities (mean AGS = 5.40 Mbp, n = 69) (Wilcoxon test, $p < 0.01$) (**Figure**
2A). We then evaluated the relationships between AGS of metagenomes and each of the
environmental variables (depth, salinity, temperature, and oxygen concentration)
independently and their interactions (ANOVA type II). The AGS in the pelagic metagenomes
is significantly associated only with depth and shows a negative correlation (**Supplementary**
Material 1 and **Figure S1**). Previous marine analysis have found the opposite effect, bigger
genome sizes in deeper areas (19). However, our pelagic analysis only covers 5 meters of depth
and explains 19% of the genome size variation in pelagic metagenomes. AGS in sediment
metagenomes is significantly associated with both salinity and depth and most of the
interactions of these variables (**Supplementary Material 1**). Both depth and salinity have a
weak positive correlation with AGS in the sediment samples (**Figure S1**). To our knowledge,

this is the first time where a positive correlation between genome size and salinity is reported
for brackish sediment microorganisms.

Although the ANOVA did not show a significant effect of water oxygen concentration
on genome size, we further investigate the effect of bottom water O_2 concentration in the AGS
of metagenomes from sediments. We separated these metagenomes into those from oxygen
concentration 0 to 2 mg/L (mostly metagenomes from the dead zone) and those metagenomes
with oxygen concentration 2 to 12.45 mg/L. We observe that benthic metagenomes from lower
oxygen concentration (mean AGS = 7.08 Mbp, n = 14) present significantly bigger genome
sizes than those in sediments with higher oxygen concentration (mean AGS = 5.85 Mbp, n =
94) (Wilcoxon test, $p < 0.01$) (**Figure 2B**). Complementarily, previous results show that the
dead zone bacteria tend to be more metabolically similar to each other when compared to
bacteria from oxic sediments (29). Our observation confirms a recent study that observed
obligate anaerobes from diverse environments present bigger genome sizes than
microaerophilic microorganisms (30). Moreover, previous studies show a positive correlation
between genome size and nutrient concentration (19,31,32). Dead zones in the Baltic Sea are
characterized by anthropogenic eutrophication (33), which would also promote bigger genome
sizes. Altogether, these results indicate that high nutrient concentration and low oxygen
concentrations in Baltic Sea dead zones may select for prokaryotes with bigger genome sizes.

While average benthic estimated genome size is bigger than pelagic, the biggest
genomes in the Baltic Sea were found in the water column. To compare genome size
between pelagic and benthic bacteria and archaea, we used metagenome-assembled genomes
(MAGs; >75% completeness and <5% contamination) from our metagenome datasets. Our
dataset compiles 216 MAGs from the sediments that dereplicated into 56 representative
genomospecies (95% average nucleotide identity). Additionally, 1920 pelagic MAGs were
dereplicated into 340 representative genomospecies. We observe that 12 phyla were detected
in both sediments and water column, while 16 phyla were specific to either habitat. Seven phyla
were found specific to the sediment (14 representative MAGs) and nine phyla were specific to
the water column (29 representative MAGs) (**Table 1**). Interestingly, only one genomospecies
representative was binned from both sediments and water column. This genome representative
belongs to genus *Mycobacterium* (phylum Actinobacteriota, mOTU_124/pelagic and
mOTU_027/sediments in **Table S2**), a genus that is not commonly found on brackish surface
waters (34). However, this genus was found to be abundant in sediments in some regions of

the Baltic Sea, especially in anoxic areas close to Landsort (35). These differences on
taxonomical composition in microbial communities between water column and sediments
altogether with latitudinal changes in microbial biodiversity in the Baltic Sea (34) show how
heterogeneous is the microbial composition of brackish environments.

Additionally, the eight largest representative genomes of the dataset were observed in
the water column and belonged to phylum Planctomycetota (family Planctomycetaceae) (7.95
– 9.69 Mbp). It has been previously observed that Planctomycetota is the phylum containing
the aquatic MAG with the biggest known estimated genome size (14.93 Mbp) (2). These large
genomes contain large collections of genes that could be linked to the very complex cell
structures and chromosomes observed in this phylum (36). Still, further research is necessary
to understand if extant genome size in phylum Plactomycetota is the result of ecological
adaptation to abiotic gradients. On the other side of the genome size spectrum, the
representative MAG with the smallest estimated genome size belongs to class
Alphaproteobacteria (family Pelagibacteraceae, 1.08 Mbp; **Table 1**). Bacteria from this family
have been widely reported to be streamlined, abundant and ubiquitous across all salinity
gradients (37,38).

Altogether, representative MAGs from sediments presented an average estimated
genome size of 3.47 Mbp, which was significantly higher than for water column MAGs (2.96
Mbp) ($p < 0.01$) (**Figure 3A**). This was true also at the phyla, class, and order level (**Figure
3C** and **D**). Bacteria from sediments presented bigger estimated genome size on average (3.67
Mbp), followed by pelagic Bacteria (2.98 Mbp), pelagic Archaea (1.97 Mbp) and sediment
Archaea (1.43 Mbp) (**Figure 3B**). This is supported by previous results, as Bacteria show
bigger genome sizes than Archaea regardless of the environment (2). Moreover, the average
estimated genome size of Baltic sediments (3.47 Mbp) is more similar to that of terrestrial
microbial genomes (3.7 Mbp) (2). Additionally, genomes in the Baltic sediments have also
lower coding density than pelagic (**Figure S2**). Our results corroborate previous findings that
streamlining is common in pelagic marine environments (7,37,39,40) and pelagic brackish
environments (23).

In our study, the average estimated AGS for the sediments (6.01 Mbp) and water
column (4.44 Mbp) is larger than the average estimated genome size of the MAGs assembled
and binned from the sediments (3.47 Mbp) and water column (2.96 Mbp), respectively (**Figure

2A and **3A**). We calculated the AGS to estimate the average genome size of the whole
microbial community. We used MicrobeCensus as a robust and accurate tool to calculate AGS
(41). However, this AGS of metagenomes overestimates the genome size because of the viral
and eukaryotic reads that might be present in the sample (42,43). On the other hand, there are
two biases with looking at the average estimated genome size of MAGs. One, assembly and
binning biases make a MAG an average 3.7% smaller than genomes from isolates in the same
genomospecies (2). Second, a bias of overlooking all those bacteria and archaea that are hard
to assemble and bin due to high genomic intrapopulation diversity (44). For these reasons, in
our study, we use two methods that have different biases to answer the same question; how is
the genome size of microorganisms distributed in the Baltic Sea. Irrespective of the method
used, the average estimated genome size in sediments is larger than in the water column.

The available metadata for benthic and pelagic carbon concentration was not
comparable due to methods and metrics used for analysis. This made it not possible to look for
a clear ecological link between bigger genome size in benthic zones and nutrient concentration
(**Table S1**). Luckily a previous study has compiled information on organic carbon stocks in the
Baltic Sea (45): by collecting information from many different studies and years, they show
that in average the top 10 cm of sediments contain between 2 and 4 times more organic carbon
per area than the water column in the Baltic Sea. More organic carbon available for bacteria
and archaea would also pose a lower pressure on the genome to streamline. From the results of
our study, we hypothesize that one of the reasons pelagic microbial genomes are smaller than
benthic microbial genomes is the difference in organic carbon availability. Altogether, a
positive correlation between genome size and nutrient concentration has been shown before
(19,31,32).

Brackish pelagic microorganisms tend to show smaller genome sizes than marine
and freshwater. In further analysis of genome size variation across salinity gradients, we
compared the average estimated genome size of pelagic Baltic Sea MAGs to previously
published genome size estimations (2). This comparison includes all taxonomic groups found
in three large MAG-datasets (17,46,47) (completeness >75%) that includes 4051 freshwater
representative MAGs, 2118 marine representative MAGs and 340 pelagic representative
MAGs from the Baltic Sea. We found that the average estimated genome size of the brackish
pelagic MAGs (2.96 Mbp) is lower than in marine MAGs (3.10 Mbp) (Kruskal-Wallis test, p
< 0.01). Furthermore, we observed the largest average estimated genome size in freshwater

MAGs (3.48 Mbp) (Kruskal-Wallis test, $p < 0.01$) (**Figure 4A**). These observed differences in

genome sizes across different pelagic environments together with the previously observed

differential functions (48) suggests that genome size has a potential signature across aquatic

environments.

We then divided all MAGs into phyla and focus on the most common from aquatic

environments: phyla Actinobacteriota, Bacteroidota, Cyanobacteria and Proteobacteria

(classes Alpha and Gammaproteobacteria) (**Figure 4B-F**). We test if the genome size

differences are consistent across phyla. We observe that only phylum Bacteroidota follows the

same trend as the full dataset in the average estimated genome size (**Figure 4C**). On the other

hand, MAGs from phylum Actinobacteriota present the biggest genome sizes for marine

environments, while no difference on average genome size is observed between freshwater and

brackish (**Figure 4B**). This result updates previous observations on aquatic Actinobacteriota

genome size variation (23). Opposite to Actinobacteriota, MAGs from phylum Cyanobacteria

show the smallest average genome size for marine environments, while we do not observe

statistical differences between brackish and freshwater MAGs (**Figure 4D**). Similar trends

were observed for isolates and MAGs of picocyanobacteria (22,49). However, it is important

to remark that DNA extraction, assembly, binning and/or quality check of aquatic

cyanobacterial MAGs is still a big challenge that needs to be addressed (**Supplementary**

Material 2) (50,51). All in all, these results hint at the complicated ecological role of genome

size in pelagic bacterial groups, where environment (2,15,16), lifestyle (4,13,14) and taxonomy

(11,12) are intertwined.

Smaller bacterial genomes in the Baltic Sea tend to lack certain functional

categories. We selected the bacterial MAGs with >90% completeness for metabolic

annotation and analyze which functional categories correlate with genome size in brackish

sediments and water column (**Figure 5 A-R** and **Supplementary Material 3**). Functional

categories include different but related metabolic pathways (KEGG modules), and each module

comprises multiple specific metabolic reactions (module steps) (52). In our results, we observe

patterns of negative correlation between estimated genome size and number of module steps

per Mbp in most of the functional categories analyzed (**Figure 5**). For example, given the core

functions amino acid metabolism, aminoacyl tRNAs and central carbohydrate metabolism, we

observe a higher number of module steps per Mbp at lower genome sizes in both environments

both in pelagic and benthic MAGs (**Figures 5A, D** and **J**). However, genome size does not

explain the number of module steps per Mbp in all functional categories and metabolisms,
especially in the case of non-essential functions such as drug resistance and transport systems
(**Supplementary Material 3F** and **N**). These results suggest that streamlining of genomes
select for specific functions and not the whole genome. This could be explained by the Black
Queen Hypothesis (53); when the fitness cost of a function is higher than its benefit, microbes
might lose it and, instead, obtain benefit from leaky metabolites from neighboring cells,
establishing interdependencies. The loss of those specific functions in the Baltic Sea might
have consequential long-lasting metabolic partnerships within the community.

When considering the total number of module steps, we observe that genomes of
benthic bacteria code for a higher number of module steps than pelagic bacteria in amino acid
metabolism, aminoacyl tRNAs, carbon fixation, nitrogen and sulfur metabolism (**Figures 5B,**
E, H, N and **Q**) (Wilcoxon test, $p < 0.01$). This could be the result of a confounding effect of
genome size: microorganisms from sediments have bigger genome sizes than those in the water
column (**Figure 2**) and therefore, code for a higher number of functions. Hence, the total
number of module steps per Mbp in all six categories was analyzed. If streamlining affects all
functional categories similarly, coding density trends would be similar for all functional
categories. Indeed, we observe that pelagic bacteria present a greater number of module steps
per Mbp than sediment bacteria in amino acid metabolism (**Figure 5C**), aminoacyl tRNAs
(**Figure 5F**), carbon fixation (**Figure 5I**) and central carbohydrate metabolism (**Figure 5L**).
However, nitrogen metabolism shows the opposite trend: sediment bacteria have a greater
number of module steps per Mbp than water column bacteria (**Figure 5O**). Literature report
that Baltic sediments contain about 95% of the total pool of nitrogen while the water column
only 5%, hence the water column only carries a small part of the overall nitrogen cycle (54).
As mentioned above for carbon, a higher availability of resources, including nitrogen-derived
compounds, would also imply a lower evolutionary pressure for streamlined genome sizes in
Baltic sediments. These results indicate that benthic bacteria potentially play a bigger role than
pelagic bacteria in nitrogen cycling of autotropic systems like the Baltic Sea (55,56).

Although most of the MAGs presented at least 1 module step related to drug resistance
(96.97% sediment MAGs and 98.81% pelagic MAGs), bacteria from sediments presented a
higher number of antibiotic resistance module steps than pelagic bacteria. This applies both to
total number of module steps and the number of module steps per Mbp (**Supplementary**
Material 3). Our results confirm that aquatic sediments are reservoirs of antibiotic resistance

genes (57,58). Just as sediments harbor more than double the organic carbon than the water

column, this allows microbial genomes to have a bigger size and code for a higher number of

genes. This allows microbes to upkeep non-essential functions and allow metabolic reservoir

in the sediments.

Baltic sediments and water column harbor bacteria with different metabolic

capabilities. From all >90% completeness MAGs, we selected only bacterial phyla with five

or more high-quality MAGs to observe how metabolism differs between different taxa in Baltic

sediments and water column (**Figure 6**).

We observed the presence of the genes *cdhH | cdhE | cooS* from the Wood-Ljungdhal

pathway exclusively in marine sediments (**Figure 6**), particularly in phyla Desulfobacterota,

Desulfobacterota E and Verrucomicrobiota (class Kiritimatiellae). This is a carbon fixation

pathway predominant in acetogenic bacteria found in anoxic conditions (59).

Complementarily, we also find putative fermentation genes for acetogenesis (*acdA | ack | pta*)

to be widespread across taxa in Baltic sediments. This would explain the potential success of

acetogenic metabolism in brackish sediments (60). We also find that the *acs* gene for acetate

fermentation into acetyl-CoA is widely distributed in both sediments and water column. These

results support the common distribution of acetogens and the Wood-Ljungdhal pathway in

Baltic glacial sediments (61).

No FeFe hydrogenases were detected, but different NiFe hydrogenases were spotted to

differ between environments: NiFe groups 3abd were detected mainly in pelagic bacteria, while

NiFe group 1 in sediments (**Figure 6**). NiFe group 1 hydrogenases could be playing a vital role

in nitrate (NO_3^-), sulfate (SO_4^{2-}) and iron (Fe^{3+}) reduction: these molecules can act as acceptors

of electrons coupled to H_2 oxidation in anoxic conditions (62). Moreover, putative genes coding

for the reduction of the above-mentioned molecules were also detected on our sediment dataset

(*napAB | narGH* for nitrate reduction, *aprA | sat* for sulfate reduction, and iron reduction series

genes). These results suggest a key role of sediment bacteria in sulfur, nitrogen, and iron

cycling in the Baltic Sea. For example, sulphate reducers such as Desulfobacterota found in

our benthic MAGs collection, most likely contribute to the release of Fe-bound phosphorus

from sediments to the water column (63).

Conclusions. In this research article, we provide a comprehensive analysis to investigate

the ecological implications of microbial genome in the Baltic Sea. We show that genome size

in Bacteria and Archaea is linked to the environment (**Figures 2**, **3, 4** and **S1** and

Supplementary Material 1), taxonomic identity (**Table 1** and **Figure 3**) and metabolic

potential (**Figure 5** and **Supplementary Material 3**). We also provide some insights on how

distinct pelagic and benthic microbial communities in the Baltic Sea are: not only microbial

MAGs retrieved from these two environments are different in taxonomy (**Table 1**), but also in

genome size (**Figure 3**) and metabolism (**Figures 5** and **6**). This highlights water bodies are

highly heterogeneous biomes, with highly distinct microbial communities between micro-

niches. With the continuous progression of aquatic microbial ecology and the development of

new isolation, omics and bioinformatic techniques, future research should provide a more

complete and unbiased view of genome sizes distribution in nature and its ecological

implication in microbial populations.

<u>MATERIAL AND METHODS</u>

Baltic Sea metagenomes collection. For this study we compiled new and public Baltic

Sea metagenomes from the water column and the sediments. The final dataset consisted of 219

metagenome samples from a wide range of locations in the Baltic Sea that include 5

independent datasets (**Table S1** and **Figure 1**). We compiled 108 sediment metagenomes, of

which 59 were collected in 2019 and recently published (29), and we collected 49 metagenomes

from 2016 to 2018. The pelagic dataset consists of 118 pelagic samples collected from 2011 to

2015 published in three different studies (46,64,65). All five datasets have abiotic metadata of

depth (m), salinity (PSU), temperature (C) and oxygen concentration (mg/L) (**Table S1**).

Environmental sampling. The top 2 cm of sediment was collected at soft bottom clay-

muddy habitats from 59 stations from north to south in the Baltic Sea in 2019, following the

sampling described in Broman et al., (2022) (**Table S1** for coordinates). Briefly, one sediment

core was collected per station using a Kajak gravity corer (surface area: 50 cm^2, one core per

station) and the top 0–2 cm layer was sliced into a 215 ml polypropylene container

(207.0215PP, Noax Lab, Sweden). The sediment was homogenized and stored at -20°C on the

boat, kept on an iced cooler without thawing for ~2 h during transportation to the university,

and finally stored again at -20 °C until DNA extraction. Bottom water (~20 cm above the

sediment surface) was collected at each station with a Niskin bottle. This was followed by on

deck measurements of bottom water temperature, salinity, and dissolved O2 using a portable
multimeter (HQ40D, Hach).

DNA extraction and sequencing. The sediment samples were thawed, homogenized,
and a subsample of 0.25 g was used for DNA extraction using the DNeasy PowerSoil kit
(Qiagen) according to the manufacturer's protocol. The quantity and quality of eluted DNA
were measured using NanoDrop One spectrophotometer and Qubit 2 (both by ThermoFisher
Scientific) to ensure that samples meet the minimum requirements for sequencing. The samples
were then sequenced at the Science for Life Laboratories facility on one NovaSeq 6000 S4
lanes using a 2 × 150 bp setup. Sequencing yielded on average 53.0 million reads per sample.

MAGs collection. Assembling and binning of the 108 sediment metagenomes resulted
in 2248 bins. To obtain bins from metagenomes, we followed the 0053_metasssnake2 pipeline
(https://github.com/moritzbuck/0053_metasssnake2) (v0.0.2). In this pipeline we used
Sourmash (66) to create signatures, Megahit (67) to obtain single-sample assemblies and
Metabat2 (68) for the binning of the assemblies. We used default parameters throughout the
whole pipeline. Quality of the bins was assessed using CheckM (v1.1.3) (69): we used the
typical workflow (*lineage_wf*) with default parameters, and selected only bins with quality of
completeness >75% and contamination <5%. From all the bins, only 216 passed our quality
threshold and we named those MAGs (metagenome-assembled genomes). All MAGs were
taxonomically classified using GTDB-tk (v1.5.0) (70) according to the GTDB classification
(data version r202) (71). The quality of the MAGs belonging to phyla Actinobacteriota and
Patescibacteria were assessed separately using a custom set of marker genes. Preliminary
quality check of Actinobacteriota genomes in a publicly available freshwater dataset (47) show
that default parameters underestimate the quality of the MAGs that are classified as
Actinobacteriota compared to using a custom marker gene set (**Supplementary Material 2**).
We obtained the custom marker gene set using Ac1 circular genomes (72,73). In the case of
phylum Patescibacteria, we used a custom set of maker genes provided by CheckM (69,74).

Complementarily, we used 1920 pelagic MAGs that were published (46) and passed
the >75% completeness and <5% contamination threshold. All high-quality MAGs from the
sediments and water column were dereplicated using fastANI (95% ANI threshold as estimator
of genetic unit) and mOTUlizer (v0.3.2) (75,76). From the 216 with >75% completeness
MAGs, 56 were chosen as representatives. From the 1920 pelagic MAGs, 340 were chosen as

representatives. All genomic information for pelagic and benthic MAGs is included in **Table**
S2.

Genome size analysis. We studied genome size in two different levels: entire microbial
community and bacterial/archaeal MAGs. To study differences in genome size between Baltic
sediments and water column at the community level, we first calculated the average genome
size (AGS) of the metagenomes using MicrobeCensus (v1.1.0) (77). MicrobeCensus estimates
the AGS of a microbial community from metagenomes by aligning reads to a set of single-
copy genes that are widely distributed across taxa to calculate their abundance, with highly
accurate estimations (41). We excluded one of the pelagic datasets (65) due to the presence of
spike-in DNA. We used default parameters, but we set the number of reads sampled to 10
million (*-n 10 000 000*). To estimate the genome size of the microbial MAGs, we divided the
MAGs assembly size by the completeness (provided by CheckM, ranging from 0 to 1). To
study genome size variation between our pelagic brackish dataset and other major aquatic
environments (freshwater and marine), we compiled the metagenomic information from all
pelagic MAGs from marine and freshwater environments (>75% completeness and <5%
contamination) from (2) (**Table S3**).

Metabolic annotation. To analyze the metabolic potential of sedimentary and pelagic
bacteria, we selected all MAGs with completeness >90% and contamination <5%. In total, we
obtained 99 MAGs from sediments and 1241 MAGs from the water column. The metabolic
potential of sediment and pelagic MAGs was reconstructed using Prodigal annotations (v2.6.3)
(78). We used the resulting protein translation files to predict biogeochemical and metabolic
functional traits using METABOLIC (v4.0) (52). We used the METABOLIC-G script, using
default settings.

Statistical analysis. We performed Wilcoxon non-parametric test to analyze if there
were significant differences between pairs of boxplots (**Figures 2, 3** and **5** and **Supplementary**
Material 3). Asterisks in boxplots indicate significant differences $p < 0.05$. In **Figures 3B** and
4 we performed Kruskal-Wallis test corrected with Benjamini-Hochberg, to test statistical
differences between groups. Different letters are the result of this non-parametric test; $p < 0.05$.
We performed a ANOVA type II analysis to test the effect of abiotic factors and their
interactions on the AGS, using the function *aov* from the R package *stats* v3.6.3 (79)
(**Supplementary Material 1**). We obtained the correlation coefficients on scatterplots

(**Figures 5** and **S1**) to test the fit of our data to linear regressions using the function *stat_cor*
from the R package *ggpubr* v0.4.0 (80).

Data availability. All metagenome datasets are available in public repositories under
NCBI project accession number SRP077551 and ENA accession numbers PRJEB34883,
PRJEB22997 and PRJEB41834. Specific accession numbers of all metagenomes are available
in **Table S1**. Pelagic MAGs can be found on project PRJEB34883 and benthic MAGs on
project PRJNA891615. Assembly and binning of the dataset provided in this paper used scripts
available at https://github.com/moritzbuck/0053_metasssnake2. Supplemental material can
also be accessed 10.17044/scilifelab.21378294.

ACKNOWLEDGEMENTS

We thank Elias Broman for his help on the acquisition of the benthic dataset of
metagenomes and metadata and Ola Svensson, Caroline Raymond and Jonas Gunnarsson for
assistance during sampling. We also thank Juanita Gutiérrez-Valencia for her suggestions in
statistical analysis.

This work was supported by SciLifeLab. The authors acknowledge support from the
National Genomics Infrastructure in Stockholm funded by Science for Life Laboratory, the
Knut and Alice Wallenberg Foundation and the Swedish Research Council, and SNIC/Uppsala
Multidisciplinary Center for Advanced Computational Science for assistance with massively
parallel sequencing and access to the UPPMAX computational infrastructure. Computational
work and data handling were enabled by resources in the projects SNIC 2020/5-159, 2021/5-
133, 2022/5-137, 2020-6-60 and 2022/6-77 provided by the Swedish National Infrastructure
for Computing (SNIC) at UPPMAX, partially funded by the Swedish Research Council
through grant agreement no. 2018-05973.

AR-G and SLG conceptualized and designed research project. AR-G, MB and SLG
refined the project idea. AR-G, AFA and FJAN participated in data collection. AR-G and DIS
did the molecular work. AR-G and MB performed bioinformatic analysis to obtain MAGs from
sediment raw sequences. AR-G and SLG performed data analysis. AR-G and SLG drafted the
first manuscript. AR-G, DIS and SLG did literature searches. All authors contributed to the
writing and editing of the manuscript.

CONFLICTS OF INTEREST

477   The authors declare that there are no competing interests.

REFERENCES

1.   Kirchberger PC, Schmidt ML, Ochman H. The Ingenuity of Bacterial Genomes. Annu
Rev Microbiol. 2020 Sep 8;74(1):815–34.

2.   Rodríguez-Gijón A, Nuy JK, Mehrshad M, Buck M, Schulz F, Woyke T, et al. A genomic
perspective across Earth's microbiomes reveals that genome size in Archaea and
Bacteria is linked to ecosystem type and trophic strategy. Frontiers in Microbiology. 2022
Jan 5;12(761869).

3.   Lynch M. Streamlining and Simplification of Microbial Genome Architecture. 2006;25.

4.   Giovannoni SJ, Cameron Thrash J, Temperton B. Implications of streamlining theory for
microbial ecology. ISME J. 2014 Aug;8(8):1553–65.

5.   Kuo CH, Moran NA, Ochman H. The consequences of genetic drift for bacterial genome
complexity. Genome Research. 2009 Aug 1;19(8):1450–4.

6.   Wolf YI, Koonin EV. Genome reduction as the dominant mode of evolution. BioEssays.
2013 Sep;35(9):829–37.

7.   Batut B, Knibbe C, Marais G, Daubin V. Reductive genome evolution at both ends of the
bacterial population size spectrum. Nat Rev Microbiol. 2014 Dec;12(12):841–50.

8.   Bobay LM, Ochman H. Factors driving effective population size and pan-genome
evolution in bacteria. BMC Evol Biol. 2018 Dec;18(1):153.

9.   Konstantinidis KT, Tiedje JM. Trends between gene content and genome size in
prokaryotic species with larger genomes. Proceedings of the National Academy of
Sciences. 2004 Mar 2;101(9):3160–5.

10. Puigbò P, Lobkovsky AE, Kristensen DM, Wolf YI, Koonin EV. Genomes in turmoil:
quantification of genome dynamics in prokaryote supergenomes. 2014;19.

11. Biller SJ, Berube PM, Lindell D, Chisholm SW. Prochlorococcus: the structure and
function of collective diversity. Nat Rev Microbiol. 2015 Jan;13(1):13–27.

12. Martinez-Gutierrez CA, Aylward FO. Genome size distributions in bacteria and archaea
are strongly linked to evolutionary history at broad phylogenetic scales. Casadesús J,
editor. PLoS Genet. 2022 May 23;18(5):e1010220.

13. Moran NA, Mira A. The process of genome shrinkage in the obligate symbiont Buchnera
aphidicola. Genome Biol. 2001;2(12):research0054.1.

14. van Ham RCHJ, Kamerbeek J, Palacios C, Rausell C, Abascal F, Bastolla U, et al.
Reductive genome evolution in *Buchnera aphidicola*. Proc Natl Acad Sci USA. 2003 Jan
21;100(2):581–6.

15. Maistrenko OM, Mende DR, Luetge M, Hildebrand F, Schmidt TSB, Li SS, et al.
Disentangling the impact of environmental and phylogenetic constraints on prokaryotic
within-species diversity. ISME J [Internet]. 2020 Feb 11 [cited 2020 Apr 1]; Available
from: http://www.nature.com/articles/s41396-020-0600-z

16. Simonsen AK. Environmental stress leads to genome streamlining in a widely distributed
species of soil bacteria. ISME J. 2022 Feb;16(2):423–34.

17. Nayfach S, Roux S, Seshadri R, Udwary D, Varghese N, Schulz F, et al. A genomic
catalog of Earth's microbiomes. Nat Biotechnol. 2021 Apr;39(4):499–509.

18. Konstantinidis KT, Braff J, Karl DM, DeLong EF. Comparative Metagenomic Analysis of
a Microbial Community Residing at a Depth of 4,000 Meters at Station ALOHA in the
North Pacific Subtropical Gyre. Appl Environ Microbiol. 2009 Aug 15;75(16):5345–55.

19. Mende DR, Bryant JA, Aylward FO, Eppley JM, Nielsen T, Karl DM, et al. Environmental
drivers of a microbial genomic transition zone in the ocean's interior. Nat Microbiol. 2017
Oct;2(10):1367–73.

20. Salcher MM, Schaefle D, Kaspar M, Neuenschwander SM, Ghai R. Evolution in action:
habitat transition from sediment to the pelagial leads to genome streamlining in
Methylophilaceae. ISME J. 2019 Nov;13(11):2764–77.

21. Chen MY, Teng WK, Zhao L, Hu CX, Zhou YK, Han BP, et al. Comparative genomics
reveals insights into cyanobacterial evolution and habitat adaptation. ISME J. 2021
Jan;15(1):211–27.

22. Cabello-Yeves PJ, Callieri C, Picazo A, Schallenberg L, Huber P, Roda-Garcia JJ, et al.
Elucidating the picocyanobacteria salinity divide through ecogenomics of new freshwater
isolates. BMC Biol. 2022 Aug 8;20(1):175.

23. Hugerth LW, Larsson J, Alneberg J, Lindh MV, Legrand C, Pinhassi J, et al.
Metagenome-assembled genomes uncover a global brackish microbiome. Genome Biol.
2015 Dec;16(1):279.

24. Coelho LP, Alves R, del Río ÁR, Myers PN, Cantalapiedra CP, Giner-Lamia J, et al.
Towards the biogeography of prokaryotic genes. Nature. 2022 Jan 13;601(7892):252–6.

25. Zhou Z, Tran PQ, Kieft K, Anantharaman K. Genome diversification in globally
distributed novel marine Proteobacteria is linked to environmental adaptation. ISME J.
2020 Aug;14(8):2060–77.

26. Acinas SG, Sánchez P, Salazar G, Cornejo-Castillo FM, Sebastián M, Logares R, et al.
Deep ocean metagenomes provide insight into the metabolic architecture of
bathypelagic microbial communities. Commun Biol. 2021 Dec;4(1):604.

27. Louca S, Parfrey LW, Doebeli M. Decoupling function and taxonomy in the global ocean
microbiome. Science. 2016 Sep 16;353(6305):1272–7.

28. Li M, Wei G, Shi W, Sun Z, Li H, Wang X, et al. Distinct distribution patterns of ammonia-
oxidizing archaea and bacteria in sediment and water column of the Yellow River
estuary. Sci Rep. 2018 Dec;8(1):1584.

29. Broman E, Izabel-Shen D, Rodríguez-Gijón A, Bonaglia S, Garcia SL, Nascimento FJA.
Microbial functional genes are driven by gradients in sediment stoichiometry, oxygen,
and salinity across the Baltic benthic ecosystem. Microbiome. 2022 Aug 15;10(1):126.

30. Nielsen DA, Fierer N, Geoghegan JL, Gillings MR, Gumerov V, Madin JS, et al. Aerobic
bacteria and archaea tend to have larger and more versatile genomes. Oikos. 2021
Apr;130(4):501–11.

31. Allen LZ, Allen EE, Badger JH, McCrow JP, Paulsen IT, Elbourne LD, et al. Influence of
nutrients and currents on the genomic composition of microbes across an upwelling
mosaic. ISME J. 2012 Jul;6(7):1403–14.

32. Aylward FO, Santoro AE. Heterotrophic Thaumarchaea with Small Genomes Are
Widespread in the Dark Ocean. Bouskill N, editor. mSystems. 2020 Jun 16;5(3):e00415-
20, /msystems/5/3/msys.00415-20.atom.

33. Conley DJ, Björck S, Bonsdorff E, Carstensen J, Destouni G, Gustafsson BG, et al.
Hypoxia-Related Processes in the Baltic Sea. Environ Sci Technol. 2009 May
15;43(10):3412–20.

34. Herlemann DP, Labrenz M, Jürgens K, Bertilsson S, Waniek JJ, Andersson AF.
Transitions in bacterial communities along the 2000 km salinity gradient of the Baltic
Sea. ISME J. 2011 Oct;5(10):1571–9.

35. Thureborn P, Lundin D, Plathan J, Poole AM, Sjöberg BM, Sjöling S. A Metagenomics
Transect into the Deepest Point of the Baltic Sea Reveals Clear Stratification of
Microbial Functional Capacities. Gilbert JA, editor. PLoS ONE. 2013 Sep
23;8(9):e74983.

36. Seeger C, Dyrhage K, Mahajan M, Odelgard A, Lind SB, Andersson SGE. The
Subcellular Proteome of a Planctomycetes Bacterium Shows That Newly Evolved
Proteins Have Distinct Fractionation Patterns. Front Microbiol. 2021 May 4;12:643045.

37. Giovannoni SJ. SAR11 Bacteria: The Most Abundant Plankton in the Oceans. Annu Rev
Mar Sci. 2017 Jan 3;9(1):231–55.

38. Lanclos VC, Rasmussen AN, Kojima CY, Cheng C, Henson MW, Faircloth BC, et al.
Ecophysiology and genomics of the brackish water adapted SAR11 subclade IIIa. ISME
J [Internet]. 2023 Feb 4 [cited 2023 Feb 6]; Available from:
https://www.nature.com/articles/s41396-023-01376-2

39. Swan BK, Tupper B, Sczyrba A, Lauro FM, Martinez-Garcia M, Gonzalez JM, et al.
Prevalent genome streamlining and latitudinal divergence of planktonic bacteria in the
surface ocean. Proceedings of the National Academy of Sciences. 2013 Jul
9;110(28):11463–8.

40. Giovannoni SJ. Genome Streamlining in a Cosmopolitan Oceanic Bacterium. Science.
2005 Aug 19;309(5738):1242–5.

41. Pereira-Flores E, Glöckner FO, Fernandez-Guerra A. Fast and accurate average
genome size and 16S rRNA gene average copy number computation in metagenomic
data. BMC Bioinformatics. 2019 Dec;20(1):453.

42. Kristensen DM, Mushegian AR, Dolja VV, Koonin EV. New dimensions of the virus world
discovered through metagenomics. Trends in Microbiology. 2010 Jan;18(1):11–9.

43. Lind AL, Pollard KS. Accurate and sensitive detection of microbial eukaryotes from
whole metagenome shotgun sequencing. Microbiome. 2021 Dec;9(1):58.

44. Ghurye JS. Metagenomic Assembly: Overview, Challenges and Applications. 2016;10.

45. Scheffold MIE, Hense I. Quantifying Contemporary Organic Carbon Stocks of the Baltic
Sea Ecosystem. Front Mar Sci. 2020 Dec 17;7:571956.

46. Alneberg J, Bennke C, Beier S, Bunse C, Quince C, Ininbergs K, et al. Ecosystem-wide
metagenomic binning enables prediction of ecological niches from genomes. Commun
Biol. 2020 Dec;3(1):119.

47. Buck M, Garcia SL, Fernandez L, Martin G, Martinez-Rodriguez GA, Saarenheimo J, et
al. Comprehensive dataset of shotgun metagenomes from oxygen stratified freshwater
lakes and ponds. Sci Data. 2021 Dec;8(1):131.

48. Jurdzinski KT, Mehrshad M, Delgado LF, Deng Z, Bertilsson S, Andersson AF. Large-
scale phylogenomics of aquatic bacteria reveal molecular mechanisms for adaptation to
salinity [Internet]. Microbiology; 2022 Oct [cited 2023 Jan 17]. Available from:
http://biorxiv.org/lookup/doi/10.1101/2022.10.03.510577

49. Sánchez-Baracaldo P, Bianchini G, Di Cesare A, Callieri C, Chrismas NAM. Insights Into
the Evolution of Picocyanobacteria and Phycoerythrin Genes (mpeBA and cpeBA). Front
Microbiol. 2019 Jan 30;10:45.

50. Alvarenga DO, Fiore MF, Varani AM. A Metagenomic Approach to Cyanobacterial
Genomics. Front Microbiol. 2017 May 9;8:809.

51. Kim Tiam S, Comte K, Dalle C, Duval C, Pancrace C, Gugger M, et al. Development of a
new extraction method based on high-intensity ultra-sonication to study RNA regulation
of the filamentous cyanobacteria Planktothrix. Kalendar R, editor. PLoS ONE. 2019 Sep
6;14(9):e0222029.

52. Zhou Z, Tran PQ, Breister AM, Liu Y, Kieft K, Cowley ES, et al. METABOLIC: high-
throughput profiling of microbial genomes for functional traits, metabolism,
biogeochemistry, and community-scale functional networks. Microbiome. 2022
Dec;10(1):33.

53. Morris JJ, Lenski RE, Zinser ER. The Black Queen Hypothesis: Evolution of
Dependencies through Adaptive Gene Loss. mBio. 2012 Mar 23;3(2):e00036-12.

54. Lønborg C, Markager S. Nitrogen in the Baltic Sea: Long-term trends, a budget and
decadal time lags in responses to declining inputs. Estuarine, Coastal and Shelf
Science. 2021 Oct;261:107529.

55. Albert S, Bonaglia S, Stjärnkvist N, Winder M, Thamdrup B, Nascimento FJA. Influence
of settling organic matter quantity and quality on benthic nitrogen cycling. Limnol
Oceanogr. 2021 May;66(5):1882–95.

56. Griffiths JR, Kadin M, Nascimento FJA, Tamelander T, Törnroos A, Bonaglia S, et al.
The importance of benthic-pelagic coupling for marine ecosystem functioning in a
changing world. Glob Change Biol. 2017 Jun;23(6):2179–96.

57. Guo X pan, Zhao S, Chen Y ru, Yang J, Hou L jun, Liu M, et al. Antibiotic resistance
genes in sediments of the Yangtze Estuary: From 2007 to 2019. Science of The Total
Environment. 2020 Nov;744:140713.

58. Marti E, Variatza E, Balcazar JL. The role of aquatic ecosystems as reservoirs of
antibiotic resistance. Trends in Microbiology. 2014 Jan;22(1):36–41.

59. Esposito A, Tamburini S, Triboli L, Ambrosino L, Chiusano ML, Jousson O. Insights into
the genome structure of four acetogenic bacteria with specific reference to the Wood–
Ljungdahl pathway. MicrobiologyOpen [Internet]. 2019 Dec [cited 2022 Aug 26];8(12).
Available from: https://onlinelibrary.wiley.com/doi/10.1002/mbo3.938

60. Lever MA. Acetogenesis in the Energy-Starved Deep Biosphere – A Paradox? Front
Microbio [Internet]. 2012 Jan [cited 2022 Aug 26];2(284). Available from:
http://journal.frontiersin.org/article/10.3389/fmicb.2011.00284/abstract

61. Marshall IPG, Karst SM, Nielsen PH, Jørgensen BB. Metagenomes from deep Baltic
Sea sediments reveal how past and present environmental conditions determine
microbial community composition. Marine Genomics. 2018 Feb;37:58–68.

62. Peters JW, Schut GJ, Boyd ES, Mulder DW, Shepard EM, Broderick JB, et al. [FeFe]-
and [NiFe]-hydrogenase diversity, mechanism, and maturation. Biochimica et Biophysica
Acta (BBA) - Molecular Cell Research. 2015 Jun;1853(6):1350–69.

63. Sinkko H. Sediment bacterial communities in nutrient cycling and in the history of the
Baltic Sea [Doctoral dissertation]. [Helsinki, Finland]: University of Helsinki; 2013.

64. Larsson J, Celepli N, Ininbergs K, Dupont CL, Yooseph S, Bergman B, et al.
Picocyanobacteria containing a novel pigment gene cluster dominate the brackish water
Baltic Sea. ISME J. 2014 Sep;8(9):1892–903.

65. Alneberg J, Sundh J, Bennke C, Beier S, Lundin D, Hugerth LW, et al. BARM and
BalticMicrobeDB, a reference metagenome and interface to meta-omic data for the
Baltic Sea. Sci Data. 2018 Dec;5(1):180146.

66. Titus Brown C, Irber L. sourmash: a library for MinHash sketching of DNA. JOSS. 2016
Sep 14;1(5):27.

67. Li D, Luo R, Liu CM, Leung CM, Ting HF, Sadakane K, et al. MEGAHIT v1.0: A fast and
scalable metagenome assembler driven by advanced methodologies and community
practices. Methods. 2016 Jun;102:3–11.

68. Kang DD, Li F, Kirton E, Thomas A, Egan R, An H, et al. MetaBAT 2: an adaptive
binning algorithm for robust and efficient genome reconstruction from metagenome
assemblies. PeerJ. 2019 Jul 26;7:e7359.

69. Parks DH, Imelfort M, Skennerton CT, Hugenholtz P, Tyson GW. CheckM: assessing the
quality of microbial genomes recovered from isolates, single cells, and metagenomes.
Genome Res. 2015 Jul;25(7):1043–55.

70. Chaumeil PA, Mussig AJ, Hugenholtz P, Parks DH. GTDB-Tk: a toolkit to classify
genomes with the Genome Taxonomy Database. Hancock J, editor. Bioinformatics.
2019 Nov 15;btz848.

71. Parks DH, Chuvochina M, Chaumeil PA, Rinke C, Mussig AJ, Hugenholtz P. A complete
domain-to-species taxonomy for Bacteria and Archaea. Nat Biotechnol. 2020 Sep
1;38(9):1079–86.

72. Neuenschwander SM, Ghai R, Pernthaler J, Salcher MM. Microdiversification in
genome-streamlined ubiquitous freshwater Actinobacteria. ISME J. 2018 Jan;12(1):185–
98.

73. Kang I, Kim S, Islam MdR, Cho JC. The first complete genome sequences of the acI
lineage, the most abundant freshwater Actinobacteria, obtained by whole-genome-
amplification of dilution-to-extinction cultures. Sci Rep. 2017 Sep 27;7(1):42252.

74. Brown CT, Hug LA, Thomas BC, Sharon I, Castelle CJ, Singh A, et al. Unusual biology
across a group comprising more than 15% of domain Bacteria. Nature. 2015
Jul;523(7559):208–11.

75. Buck M, Mehrshad M, Bertilsson S. mOTUpan: a robust Bayesian approach to leverage
metagenome-assembled genomes for core-genome estimation. NAR Genomics and
Bioinformatics. 2022 Jul 9;4(3):lqac060.

76. Jain C, Rodriguez-R LM, Phillippy AM, Konstantinidis KT, Aluru S. High throughput ANI
analysis of 90K prokaryotic genomes reveals clear species boundaries. Nat Commun.
2018 Dec;9(1):5114.

77. Nayfach S, Pollard KS. Average genome size estimation improves comparative
metagenomics and sheds light on the functional ecology of the human microbiome.
Genome Biol. 2015 Dec;16(1):51.

78. Hyatt D, Chen GL, LoCascio PF, Land ML, Larimer FW, Hauser LJ. Prodigal: prokaryotic
gene recognition and translation initiation site identification. BMC Bioinformatics. 2010
Dec;11(1):119.

79. R Core Team. R: A language and environment for statistical computing. [Internet]. R
Foundation for Statistical Computing, Vienna, Austria; 2020. Available from:
https://www.R-project.org/.

80. Kassambara, A. ggpubr: 'ggplot2' Based Publication Ready Plots. [Internet]. 2020.
Available from: https://CRAN.R-project.org/package=ggpubr

<u>FIGURES LEGENDS</u>

Figure 1. Overview of the sampling locations and average genome size (AGS) of metagenomes

(108 from sediments in dark blue, and 111 from the water column in light blue). This figure

shows the geographic location of all metagenomes used in this study. For exact coordinates see

Table S1. Shape type indicates the reference and shape size indicates the AGS of the given

metagenome.

Figure 2. Boxplots showing the AGS distribution of Baltic metagenomes. **Panel A** indicates the AGS distribution in both water column and sediments. **Panel B** indicates the AGS distribution in metagenomes from sediments across the oxygen gradient (two groups, from 0 to 2 and from 2 to 12.45 mg/L). Stars in both panels indicate significant differences $p < 0.05$ (Wilcoxon non-parametric test).

Figure 3. Overview of the estimated genome size in bacteria and archaea obtained from Baltic Sea sediments (dark blue) and water column (light blue) using only the 397 representative MAGs (95% average nucleotide identity) with >75% completeness. **Panel A** shows the genome size distribution of archaea and bacteria obtained from Baltic water column and sediments for a total of 397 representative genomes. **Panel B** shows the estimated genome size per domain and environment. Different letters indicate significant differences $p < 0.05$ (Kruskal-Wallis non-parametric test; multiple testing corrected with Benjamini-Hochberg). **Panel C** shows the estimated genome size per phylum. We selected only phyla with at least 2 MAGs in each environment. **Panel D** shows the estimated genome size per order. We selected only orders with at least 2 MAGs in each environment. Numbers below the boxes indicate the number of MAGs per environment. Stars in Panel C and D indicate significant differences $p < 0.05$ (Wilcoxon non-parametric test).

Figure 4. Overview of the estimated genome size of pelagic Bacteria and Archaea obtained from Baltic Sea (blue), freshwater (green) and marine (yellow) using only representative MAGs (calculates using 95% average nucleotide identity) with >75% completeness. We compare all representative MAGs (**Panel A**), only phylum Actinobacteriota (**Panel B**), phylum Bacteroidota (**Panel C**), phylum Cyanobacteria (**Panel D**), class Alphaproteobacteria (**Panel E**) and class Gammaproteobacteria (**Panel F**). Different letters indicate significant differences $p < 0.05$ (Kruskal-Wallis non-parametric test; multiple testing corrected with Benjamini-Hochberg). Numbers below the boxes indicate the number of MAGs per environment.

Figure 5. Overview of the presence of module steps for six metabolic categories: amino acid metabolism (**Panels A-C**), aminoacyl tRNAs (**Panels D-F**), carbon fixation (**Panels G-I**), central carbohydrate metabolism (**Panels J-L**), nitrogen metabolism (**Panels M-O**) and sulfur metabolism (**Panels P-R**). We used all bacterial MAGs with very-high quality (>90% completeness and <5% contamination), from both environments (99 MAGs from sediments

and 1215 MAGs from water column). Stars in boxplots indicate significant differences $p < 0.05$

(Wilcoxon non-parametric test). **Panels B, E, H, K, N** and **Q** indicate total number of module

steps. **Panels C, F, I, L, O** and **R** indicate the number of module steps per Mbp.

Figure 6. Metabolic potential of all high-quality MAGs (>90% completeness and <5%

contamination). We selected all phyla with at least 5 MAGs, and then divided by class. Boxes

on the top of the figure indicate environment (pelagic in light blue and sediments in dark blue),

and inner numbers indicate the number of MAGs per category. In the heatmap, white squares

indicate absence of a given gene in all MAGs, and the darkest purple indicates presence in all

of them (gradient scale on the bottom-left part of the figure for reference).

Table 1. Summary of all 56 sediment and 340 water column representative MAGs (95%

average nucleotide identity) with >75% completeness. Table includes phyla, environment

(either water column or sediments), number of representative genomes (*n*), smallest and largest

estimated genome sizes (Mbp) observed for each phylum, average GC content (%) and average

coding density (%). When only one MAG is indicated, estimated genome size of that MAG is

expressed in the sixth column.

Aminoacid metabolism

Aminoacyl tRNAs

Carbon fixation

Central carbohydrate metabolism

Nitrogen metabolism

Sulfur metabolism

Phyla	Environment	n	Smallest estimated genome size (Mbp)	Largest estimated genome size (Mbp)	Average GC (%)	Average coding density (%)
Actinobacteriota	Sediments	6	2.84	3.7	67.55	91.63
	Water column	59	1.25	4.53	54.23	93.74
Bacteroidota	Sediments	2	3	4.61	41.08	89.69
	Water column	86	1.38	4.71	40.7	92.97
Chloroflexota	Sediments	3	2.38	5.19	63.57	91.14
	Water column	5	1.25	5.91	57.42	90.26
Desulfobacterota	Sediments	8	2.71	4.66	49.32	84.55
	Water column	1	-	2.73	54.4	87.06
Gemmatimonadota	Sediments	1	-	2.85	66.66	93.99
	Water column	1	-	4.32	62.47	91.21
Myxococcota	Sediments	3	4.95	6.74	63.54	90.84
	Water column	1	-	7.66	66.33	90.86
Planctomycetota	Sediments	1	-	5.09	66.06	90.33
	Water column	29	3.18	**9.69**	59.01	88.23
Proteobacteria (Alfa)	Sediments	1	-	3.43	59.14	91.26
	Water column	46	**1.08**	5.92	51.88	91.98
Proteobacteria (Gamma)	Sediments	10	1.93	4.8	59.56	90.96
	Water column	54	1.18	4.19	50.48	92.09
Verrucomicrobiota	Sediments	3	2.97	3.69	56.16	89.4
	Water column	25	1.98	6.82	57.36	90.41
Thermoplasmatota	Sediments	1	-	1.51	54.16	91.35
	Water column	2	2.35	2.68	41.63	93.32
Thermoproteota	Sediments	3	1.42	1.58	35.32	89.34
	Water column	2	1.42	1.48	31.86	89.96
Acidobacteriota	Sediments	5	4.13	5.64	65.98	91.6
Desulfobacterota D	Sediments	1	-	2.33	45.15	88.3
Desulfobacterota E	Sediments	2	2.35	2.74	64.01	91.94
Nitrospirota	Sediments	3	2.86	4	54.01	86.72
Omnitrophota	Sediments	1	-	1.47	41.17	91.99
Zixibacteria	Sediments	1	-	3.68	41.12	90.73
Micrarchaeota	Sediments	1	-	1.14	48.22	93.23
Bdellovibrionota	Water column	1	-	3.73	49.16	93.45
Bdellovibrionota C	Water column	1	-	3.33	47.41	85.46
Campylobacterota	Water column	2	2.6	3.3	37.02	93.25
Cyanobacteria	Water column	16	1.98	6.03	51.8	88.45

Firmicutes	Water column	3	1.09	1.16	38.25	93.69
Krumholzibacteriota	Water column	1	-	3.34	63.78	92.99
Marinisomatota	Water column	2	2.62	3.11	39.47	91.47
Nitrospinota	Water column	2	3.18	3.35	47.22	87.87
Nanoarchaeota	Water column	1	-	1.93	29.77	92.86