SARS-CoV2 E and 3a proteins are inducers of pannexin currents

Barbara Oliveira Mendes[1,#], Malak Alameh[1,2,#], Béatrice Ollivier[1], Jérôme Montnach[1], Nicolas Bidère[3,4], Frédérique Souazé[5], Nicolas Escriou[6], Flavien Charpentier[1], Isabelle Baró[1], Michel De Waard[1,2,§,*], Gildas Loussouarn[1,§,*]

[1]Nantes Université, CNRS, INSERM, l'institut du thorax, Nantes, France

[2]Labex Ion Channels, Science and Therapeutics, Valbonne, France

[3]Team SOAP, CRCI2NA, Nantes Université, Inserm, CNRS, Université d'Angers, Nantes, France

[4]Equipe Labellisée Ligue Contre le Cancer, Paris, France

[5]Nantes Université, INSERM, CNRS, CRCI2NA, Nantes, France

[6]Institut Pasteur, Université Paris Cité, Département de Santé Globale, Paris, France

# these authors contributed equally to this work.

§ Both senior authors codirected this work.

* Corresponding authors*: *michel.dewaard@univ-nantes.fr* *or* *gildas.loussouarn@univ-nantes.fr*; *Tel.: +33-228-080-076*

Short title: Ionic conductances triggered by SARS-CoV2 E and 3a proteins

Abstract

Controversial reports have suggested that SARS-CoV E and 3a proteins may be viroporins that conduct currents through the plasma membrane of the infected cells. If true, these proteins would represent accessible targets for the development of new antiviral drugs by using high-throughput patch-clamp techniques. Here we aimed at better characterizing the cell responses induced by E or 3a protein with a particular focus on the ion conductances measured at the cell surface. First, we show that expression of SARS-CoV-2 E or 3a protein in CHO cells gives rise to cells with newly-acquired round shape, tending to detach from the Petri dish. This suggests that cell death is induced upon expression of E or 3a protein. We confirmed this hypothesis by using flow cytometry, in agreement with earlier reports on other cell types. In adhering cells expressing E or 3a protein, whole-cell currents were in fact not different from the control condition indicating that E and 3a proteins are not plasma membrane viroporins. In contrast, recording currents on detached cells uncovered outwardly-rectifying currents, much larger than those observed in control. The current characteristics are reminiscent of what was previously observed in cells expressing SARS-CoV-1 E or 3a proteins. Herein, we illustrate for the first time that carbenoxolone blocks these outward currents suggesting that they are conducted by pannexin channels, mostly likely activated by cell morphology change and/or cell death. Alongside we also demonstrate that truncation of the C-terminal PDZ binding motifs reduces the proportion of dying cells but does not prevent pannexin currents suggesting distinct pathways for cell death and pannexin currents induced by E and 3a proteins. We conclude that SARS-CoV-2 E and 3a proteins are not acting as viroporins expressed at the plasma membrane.

Author Summary

A viroporin (or viral porin) is a class of proteins that is encoded by a virus genome. It is named porin because its biological role is to conduct ions through a pore that it created in a lipid membrane such as the one surrounding a human cell. if such viroporin is present at the external membrane of a human cell infected by a virus, it can be an easy target of an antiviral agent which thus does not have to enter the cell to be active. One example of viroporin is the flu M2 protein that is the target of amantadine, an antiviral agent used against flu. In previous studies, two proteins of SARS-CoV viruses, named E protein and 3a protein, have been suggested to be viroporins at the surface of infected human cells, potentially opening a new research avenue against SARS. Here we demonstrate that both proteins are not viroporins at the external membrane but they rather trigger changes in the cell shape and promote cell death. They only indirectly induce the activity of a porin that is encoded by the cell genome, named pannexin.

Introduction

SARS-CoV-2 is the third virus of the genus Beta-coronavirus of the Coronaviridae family to be responsible for a Severe Acute Respiratory Syndrome in this century, after SARS-CoV-1 in 2002-2003 (1) and MERS-CoV in 2012 (2). As a result, it is of great importance to best characterize coronaviruses and the associated pathophysiology, with the hope that new treatments will emerge to complement vaccine approaches for people who cannot access the vaccines or are not responsive to them. In addition to paxlovid which is already available but associated with bothersome side-effects (3), many potential anti-COVID-19 treatments are in development but it is too soon to tell how efficient they will be, namely with regard to the continuous emergence of new variants, and if the cost will be reasonable (4).

Viroporins, *i.e.* ion channels encoded by a virus genome, are potential targets by antiviral agents, as demonstrated by the case of amantadine which inhibits the acid-activated M2 channel of Influenza A virus (5). Several studies led to the suggestion that two proteins of SARS-CoV are viroporins. SARS-CoV-2 Envelop (E) protein, is a one-transmembrane-domain membrane protein (75 amino-acids) almost identical to SARS-CoV-1 Envelop protein (95% identity). The SARS-CoV-2 ORF3a (3a) protein is a larger three-transmembrane-domain membrane protein (275 amino-acids) relatively similar to the SARS-CoV-1 3a protein (73% identity).

Regarding the ion channel function of these proteins, there are clearly several contradicting studies: some of them raising intriguing issues and others not confirming these reports. Concerning *in-vitro* membrane incorporation of purified E or 3a protein in lipid bilayers, the presence of ion channel activity was reportedly associated with these viral proteins (6–11). However, a review article soundly outlined the lack of robust data and raised ethical concerns casting doubts on the validity of the scientific messages (12). Concerning viral protein expression in cells, the expression of SARS-CoV-1 E protein also led to conflicting results (13,14). Pervushin et al managed to identify plasma membrane currents generated by heterologous expression of SARS-CoV-1 E protein in HEK-293 cells (13) but not Nieto-Torres et al (14). More recently, expression of wild-type (WT) SARS-CoV-2 E protein did not lead to interpretable ionic currents in HEK-293S or *Xenopus laevis* oocytes (15) despite their high homology among SARS-CoV viruses. In an attempt to favor plasma membrane targeting and reveal a putative current, a C-terminal predicted ER retention signal of SARS-CoV-1 E protein was replaced by a Golgi export signal from Kir2.1 channel. Expression of this chimera could then be associated with the generation of a non-rectifying and cation-selective current. This current was thus quite different from the outwardly rectifying current observed by Pervushin and collaborators (15). Furthermore, another study using a membrane targeting sequence, fused to the N-terminus of the SARS-CoV-1 E protein, provided a non-rectifying current that was 100-fold larger than the one observed in the two previous studies (16). This suggests that such modification of either N- or C-terminus are too drastic to faithfully report the actual activity of the native proteins.

SARS-CoV-1 3a protein was also investigated and expression of the WT protein in HEK-293 cells (17) or *Xenopus laevis* oocytes (18–20) was associated with a poorly selective outwardly rectifying current in both models, resembling to the one observed upon expression of the E protein.

To summarize, there is no unequivocal evidence that SARS-CoV E and 3a proteins are viroporins active at the plasma membrane of the host cell. However, it was recently reported that SARS-

CoV-2 E and 3a proteins can promote cell death (21,22), on one hand. On the other hand,
apoptosis is associated with the increase of an outwardly-rectifying current conducted by
pannexins (23–25). This led us to reinvestigate the actual function(s) of SARS-CoV-2 E and 3a
proteins in mammalian cells in the frame of the cell toxicity of these proteins.

In this study, CHO cells expressing either CoV-2 E or 3a proteins tended to develop into a round-
shaped form with a tendency to detach from the Petri dish, a process exacerbated compared to
control conditions. This cell phenotype is consistent with cell death (26,27) and we confirmed
by flow cytometry experiments that expression of E or 3a proteins does indeed promote cell
death. Transfected cells, still attached to the Petri dish (adhering cells), had unchanged basal
currents indicating that E and 3a proteins are unlikely to act as plasma membrane channels. In
contrast, recording whole-cell currents on round-shaped and detached cells, we observed large
outwardly-rectifying currents only in E or 3a protein-expressing cells but not in control dying
cells. This current is reminiscent of those observed in previous publications using HEK-293 cells
and oocytes expressing the SARS-CoV-1 proteins (13,18–20). Application of carbenoxolone, a
pannexin channel inhibitor, suggests for the first time that these currents are pannexin-
mediated conductances, potentially activated in apoptotic cells. In conclusion, both SARS-CoV-2
E and 3a proteins are most likely triggers of endogenous conductance.

Results

We first focused on native E and 3a proteins. To maximize the chance of observing E and 3a
protein-induced ionic currents, we chose to use pIRES plasmids, in which the protein of interest
situated in the first cassette is more expressed than the eGFP reporter in the second cassette,
thereby guaranteeing expression of a high level of the protein of interest in fluorescent cells
(28,29). For the purpose of this study, we also selected CHO rather than HEK-293 cells because
they express minimal endogenous currents (30). We compared whole-cell currents recorded
during a ramp protocol, in cells transfected either with a control pIRES2-eGFP plasmid (pIRES),
or the same plasmid containing the cDNA of the SARS-CoV-2 E protein (pIRES - E) or 3a protein
(pIRES - 3a). Unexpectedly, we did not observe any difference in the currents recorded for the
SARS-CoV-2 proteins expressing cells compared to control pIRES condition (Figure 1A). However,
many cells transfected with either E- or 3a-encoding plasmids were developing altered
morphology, shifting from spindle-like cells to more round cells (Figure 1B), similar to what was
previously observed in MDCK cells heterologously expressing SARS-CoV-1 E protein (31).
Analysis with Fiji tool confirmed an increase in cell roundness (Figure 1C and suppl. Figure 1). In
particular, in patch-clamp experiments, some cells were coming off from the dish bottom by
losing adhesion. Cell counting indicated that slightly more cells were losing adhesion when E or
3a protein were expressed (3.4 ± 0.6 % in non-transfected cells, 5.2 ± 1.0% in pIRES condition,
6.6 ± 0.7% in pIRES - E, 6.0 ± 1.2% in pIRES - 3a, n=3-5). As classically performed, currents shown
in Figure 1A were recorded from adhering cells while non-adhering cells were disregarded in this
initial investigation. Noteworthy, in each condition, both spindle-like and round adhering cells
were studied (pIRES: 21 spindle-like and 17 round cells; pIRES E: 9 and 13, pIRES 3a: 18 and 11).

Figure 1: Expression of E and 3a protein is accompanied by altered cellular morphology but no
modification in whole-cell currents in adhering cells. **(A)** Average current densities (± sem)
recorded during the ramp protocol (inset) in adhering CHO cells expressing either eGFP (pIRES),
SARS-CoV-2 E and eGFP proteins (pIRES - E) or SARS-CoV-2 3a and eGFP proteins (pIRES - 3a). **(B)**
Superimposed brightfield and eGFP fluorescence images of CHO cells in the same 3 conditions
as shown in A. **(C)** Analysis of cell roundness in cell automatically detected by Fiji, cf. suppl. Figure
1. Plot of individual cells (pIRES, n = 1922; pIRES - E, n = 1198; and pIRES - 3a, n = 1283), median
± interquartile range. **** p<0.0001 as compared to pIRES control, Kruskal-Wallis test.

Suppl. Figure 1: Morphology analysis of cells transfected with control pIRES or pIRES - E
plasmids. **(A)** Left: example of superimposed brightfield and eGFP fluorescence images. Right:
Automatic particle analysis showing the roundness index **(B)** Distribution of cells roudnesses
from the analysis of 5 areas of 1.7 mm x 1.7 mm for cells transfected with control pIRES, pIRES -
E or pIRES- 3a plasmids.

Since both E and 3a proteins are promoting cell death (21,22), we hypothesized that the various
cell morphological patterns (spindle-shaped, round-adhering, and round non-adhering) may
correspond to the development of cell death, as described earlier in CHO and other cells (26,27).
Flow cytometry analysis performed on the eGFP-positive CHO cells (Figure 2) showed that
expression of E and 3a proteins increases the percentage of dying cells, more significantly late
cell death, revealed by propidium iodide permeability (suppl. Figure 2). The effect of 3a protein
was greater than the effect of the E protein. E protein-induced cell death could be reduced by
the pan-caspase inhibitor QVD-Oph, while 3a protein-induced cell death was not (suppl. Figures
3 & 4), suggesting that E protein induces apoptosis, while 3a protein activates non-conventional
caspase-independent cell death.

Both E and 3a proteins possess a C-terminal PDZ binding motif (PBM). E protein PBM has been
suggested to be a virulence factor (11) and binds to host cells PDZ domains, leading to abnormal
cellular distribution of the targeted proteins (31). 3a PBM interacts with at least five human PDZ-
containing proteins (TJP1, NHERF3 & 4, RGS3, PARD3B), suggesting that it also alters cellular
organization (32). We thus evaluated whether deletion of these domains impacts E and 3a
proteins propensity to trigger cell death. Two C-terminal deletions used in previous studies to
remove E protein PBM, Δ4 for the last 4 amino-acids (31) and Δ8 for the last 8 residues (11)
abolished the pro-apoptotic effect of E protein (Figure 2). When looking individually at early and
late cell death, we observed that both truncation of E and 3a protein decreased late cell death
(suppl. Figure 2).

Figure 2: Expression of E and 3a protein induces cell death. **(A)** Flow cytometric analysis of eGFP-positive CHO cells expressing only eGFP (pIRES), or co-expressing eGFP and one of the following SARS-CoV-2 cDNA: the full-length E protein (pIRES - E), C-terminally deleted E protein (pIRES - E Δ 4 or pIRES - E Δ 8), the full length 3a protein (pIRES - 3a) or the C-terminally deleted 3a protein (pIRES - 3a Δ 10). After 48 h of expression, cells were stained with annexin V AlexaFluor 647 (APC)/propidium iodide (PI, Perc-P). **(B)** Mean ± sem of the percentage of Annexin V+ cells among eGFP-positive CHO cells expressing only eGFP (pIRES), or eGFP and full-length or truncated E protein. *** p<0.001 as compared to pIRES control, one-way ANOVA, ## p<0.01 as compared to E protein, t-test. **(C)** Mean ± sem of the percentage of Annexin V+ cells among eGFP-positive CHO cells expressing only eGFP (pIRES), or eGFP and full-length or truncated 3a protein. ** p<0.01 as compared to pIRES control, one-way ANOVA.

Suppl. Figure 2: Effects of E and 3a protein expression on early (Annexin V+, PI- in A&B) and
late (Annexin V+, PI+ in C&D) cell death. Mean ± sem of the percentage of stained cells among
eGFP-positive CHO cells expressing only eGFP (pIRES), or eGFP and full-length or truncated E or
3a protein. * p<0.05, *** p<0.001 as compared to pIRES control, one-way ANOVA, ## p<0.01, #
p<0.05 as compared to full length protein, t-test.

Suppl. Figure 3: Caspase dependence of E and 3a protein-induced cell death. (**A**) Cell death
induced by E protein expression is reduced by the pan-caspase inhibitor QVD-Oph. ** p<0.01,
as compared to pIRES control, one-way ANOVA. # p<0.05 as compared to E protein, t-test. (**B**)
Cell death induced by 3a protein expression is not reduced by the pan-caspase inhibitor QVD-
Oph. * p<0.05, as compared to pIRES control, one-way ANOVA. Concentrations are indicated in
the method section.

Suppl. Figure 4: Test of the apoptosis inducers and inhibitor in CHO cells. **** p<0.0001, ***
p<0.001, as compared to pIRES control, one-way ANOVA, # p<0.05 as compared to pIRES+S63-
845+ABT737, t-test.

Since both E and 3a proteins promote cell death, we hypothesized that the cells starting to come
off the surface may express currents induced by cell death such as pannexin currents (24,25,33).
We thus compared patch-clamp recordings of adhering cells *vs*. non-adhering cells in the 3
conditions: control pIRES, pIRES - E and pIRES - 3a plasmids (Figure 3). In the control pIRES
condition, focusing on non-adhering cells in the 35-mm dish and using the ramp protocol, we
observed an outwardly rectifying current with a mean current density of 8.5 ± 3.1 pA/pF at +70
mV, slightly higher than in spindle- or round-shaped adhering cells (3.4 ± 0.8 pA/pF). On the
other hand, in non-adhering cells expressing either the E or 3a protein, currents were much
larger in the E protein condition (I_{+70mV} = 31 ± 9 pA/pF, two-way ANOVA test on the ramp-evoked
currents: p<0.0001), but also in the 3a protein condition (I_{+70mV} = 44 ± 13 pA/pF, two-way ANOVA
test on the ramp-evoked currents: p<0.0001), as compared to non-adhering cells in the control
pIRES condition. Noteworthy, only a fraction of the cells was exhibiting large rectifying currents,
as shown in Figure 3: 4 out of 43 in the control pIRES condition, 14 out of 46 in the E protein
condition and 16 out of 41 in the 3a protein condition. These experiments suggest that cell death
and/or change in morphology induced by expression of E and 3a proteins may lead to an
increased membrane permeability by enhancing the expression or activity of an endogenous ion
channel.

The outwardly rectifying currents that we observed resemble apoptosis-induced and stretch-
induced pannexin currents (23–25). Thus, we applied the pannexin inhibitor carbenoxolone
(CBX) on non-adhering cells that display large outwardly rectifying currents (Figure 4). We
observed that CBX, applied at 50 µmol/L, inhibits the observed current, restoring current
amplitudes similar to the ones observed in the control cells. Altogether, these observations
suggest that the current triggered by the expression of E and 3a protein is a pannexin current.

Figure 3. Expression of E and 3a protein is accompanied by outwardly rectifying currents in non-adhering CHO cells only. Left, average current densities (± sem) recorded during the ramp protocol in adhering (empty circles) or non-adhering (filled circles) CHO cells expressing either eGFP (**A**, pIRES), SARS-CoV-2 E and eGFP proteins (**B**, pIRES - E) or SARS-CoV-2 3a and eGFP proteins (**C**, pIRES - 3a). Right, plot of the individual adhering cells (darker color) or non-adhering cells (lighter color). **** p<0.0001, as compared to adhering cells, two-way ANOVA.

Figure 4. Currents due to E and 3a protein expression in non-adhering CHO cells are suppressed by the pannexin inhibitor carbenoxolone (CBX). **(A)** Recording of the current amplitude in a CHO cell expressing SARS-CoV-2 E and eGFP proteins that displays large outwardly rectifying currents, in absence and presence of CBX. **(B)** Left, average current densities recorded during the ramp protocol in non-adhering CHO cells, expressing SARS-CoV-2 E and eGFP proteins (pIRES - E), in absence (Ctrl, lighter color) and presence of CBX (darker color). Right, plot of the individual non-adhering cells in absence (lighter color) and presence of CBX (darker color). **(C)** Recording of the current amplitude in a CHO cell expressing SARS-CoV-2 3a and eGFP proteins that displays the outwardly rectifying currents, in absence and presence of CBX. **(D)** Left, average current densities recorded during the ramp protocol in non-adhering CHO cells, expressing SARS-CoV-2 3a and eGFP proteins (pIRES - E), in absence (Ctrl, lighter color) and presence of CBX (darker color). Right, plot of the individual non-adhering cells in absence (lighter color) and presence of CBX (darker color). *** $p<0.001$ or **** $p<0.0001$, as compared to Ctrl, two-way ANOVA with repeated measures.

We reported above (Figure 2) that deleting the last 4 amino-acids of E protein (Δ4) drastically
reduced its pro-apoptotic effect. Cells expressing the Δ4 E protein showed an average roundness
similar to cells expressing the WT protein, suggesting that deletion did not prevent its effect on
cell morphology (Figure 5A and suppl. Figure 5). Also, when focusing on round and non-adhering
Δ4 E protein expressing cells, we could still record large outwardly rectifying currents (5 out of
20 cells), suggesting that C-terminal deletion of E protein does not abolish the induction of
pannexin currents, despite prevention of apoptosis probed by flow cytometry experiments
(Figure 5B).

We also reported in Figure 2 that the deletion of the last 10 amino-acids of 3a protein (Δ10) also
decreased cell death albeit to a lesser extent. As for E protein deletion, cells expressing the
truncated 3a protein showed an average roundness similar to cells expressing the WT protein
(Figure 5A and suppl. Figure 5). Focusing on the non-adhering cells, we could still record large
outwardly rectifying currents (9 out of 14 cells), suggesting that deletion of the last 10 amino-
acids is not sufficient to completely abolish the induction of pannexin-like currents (Figure 5C).

Figure 5. C-terminal deletion of E and 3a protein does not prevent cell characteristic changes.
(A) Cell roundness measured in CHO cells expressing eGFP (pIRES) alone, or in combination with
WT (pIRES - E) or the C-terminally deleted E protein (pIRES - E Δ4) or in combination with WT
(pIRES - 3a) or the C-terminally deleted 3a protein (pIRES - 3a Δ10). Plot of individual cells,
median ± interquartile range. **** p<0.0001 as compared to pIRES control, Kruskal-Wallis test.
(B) Left, average current densities (± sem) recorded during the ramp protocol in adhering (empty
circles) or non-adhering (filed circles) CHO cells expressing the C-terminally deleted E protein
(pIRES - E Δ4). Right, plot of the individual adhering (darker color) or non-adhering cells (lighter
color). *** p<0.001, as compared to adhering cells, two-way ANOVA. **(C)** Left, average current
densities (± sem) recorded during the ramp protocol in adhering (empty circles) or non-adhering
(filled circles) CHO cells expressing the C-terminally deleted 3a protein (pIRES - 3a Δ10). Right,
plot of the individual adhering (darker color) or non-adhering cells (lighter color). **** p<0.0001,
as compared to adhering cells, two-way ANOVA.

281
282 **Suppl. Figure 5.** Distribution of cells roundness from the analysis of 5 areas of 1.7 mm x 1.7 mm
283 of transfected CHO cells (Cf. Figure 5A).

Discussion

The concept that SARS-CoV E or 3a proteins could be viroporins expressed at the plasma
membrane is a seducing one as it could help the identification of new therapeutic drugs against
COVID-19 by setting up a screening program on the channel activity. However, this concept was
controversial and part of the reasons that explain the controversy about the function of E and
3a proteins is likely linked to the fact that these proteins also trigger morphological alterations
and cell death. Cell death is associated to important changes in lipid composition and membrane
permeability which suffices to fuel the controversy. One could imagine for instance that cell
death would be a way to activate the function of E and 3a proteins at the plasma membrane,
but an alternative hypothesis could be simply that cell death triggers an endogenous cell
conductance unrelated to the cell function of E and 3a viral proteins. The only way to address
these issues was to confirm the cell toxicity, to characterize the impact of the expression of these
proteins on cell shape and behavior, to measure plasma membrane conductance, and to
characterize them to get an idea of their nature and probe a pharmacological agent that would
match the conductance identity. We manage to solve these issues by carefully characterizing
the membrane conductances triggered by both E and 3a proteins. The fact that both proteins
trigger the same conductance independently of each other was a first indication that they could
not be viroporins at the plasma membrane. The second hint was that adhering cells, whether
they had a round-shape or not, did not exhibit any outward conductance in spite of E or 3a
protein expression. Finally, the sensitivity to carbenoxolone of the outward currents triggered
by E or 3a proteins in non-adhering cells was an indication that these viral proteins trigger
cellular alterations, such as morphological changes and cell death that are inducers of pannexin-

like current. Globally, these observations remain consistent with previous observations that
both E and 3a proteins are mainly localized in intracellular compartments in various cell types
(14,34–38). Therefore, it is fair to mention that we cannot fully conclude on the viroporin nature
of these viral proteins as their localization in subcellular organelles prevents us for clearly testing
their intrinsic potential for channel activity.

We showed that expression of either of these two proteins in CHO cells induces an increase in
cell death, as quantified by flow cytometry experiments. It is likely, although we did not
investigate this point in details, that this cell death accompanies the change in cell morphology
and Petri dish detachment. As such, our observation that pannexin-like currents are mainly
observed in detached round-shaped cells indicates that major cell morphology changes, up to
level of surface detachment, is required for the induction of pannexin-like currents. Whatever
the exact mechanism, the upregulation of pannexin channels upon cell death was already
observed earlier (39). It is thus not so surprising in fact that other reports faced problems
reporting and identifying the conductances triggered by E and 3a viral proteins. The conditions
for observing them are indeed quite drastic and require examining cells that are in the combined
dying and detachment process, something that is not naturally pursued by researchers,
especially if one hopes to detect a viroporin activity. To reconcile our data with earlier
publications, we noticed that whole-cell currents observed by others after SARS-CoV-1 E or 3a
protein expression in HEK 293 cells (13,17) were also very similar to pannexin currents:
outwardly rectifying current, with a reversal potential close to 0 mV at physiological ion
concentrations indicating a poor ion selectivity, and amplitude of a few 100 pA.

One possibility is that the pannexin-like currents that we observed are due to the classical
caspase-induced cleavage of pannexin (40). Intriguingly, deletion of the C-terminal PBM of E
protein abolished its pro-apoptotic effect but cell morphology alteration and pannexin-like
currents were still present. Regarding the 3a protein, deletion of its PBM domain only decreased
and did not completely abolish the promoted cell death, but again cell morphology alteration
and pannexin-like currents were preserved. Altogether, these results suggest that cell
morphology modification and pannexin induction may be linked and these processes are not
necessarily accompanied by cell death. One has to keep in mind that pannexin currents are
activated by many stimuli in addition to cell death (40). In particular, pannexin currents are also
stretch-activated and may be enhanced in the detached cells which are undergoing major
morphological alterations (23). If pannexins are already activated by stretch, they would not be
overactivated by their cleavage by caspase, which would explain the fact that E and 3a proteins
truncation do no prevent pannexin current, but only cell death.

In conclusion, SARS-CoV-2 native E and 3a proteins, and probably SARS-CoV-1 ones, do not act
as plasma membrane ion channels, but trigger the activity of plasma membrane pannexin
channels, most likely through morphological alteration of the cells. However, our study does not
rule out potential channel activity in intracellular membranes leading to cell death. Future
studies will give more insights on the role of pannexin channels in COVID-19 physiopathology
and treatment (41–43).

Materials and Methods

Cell culture

The Chinese Hamster Ovary cell line, CHO, was obtained from the American Type Culture
Collection (CRL-11965) and cultured in Dulbecco's modified Eagle's medium (Gibco 41966-029,

USA) supplemented with 10% fetal calf serum (Eurobio, EU), 2 mM L-Glutamine and antibiotics
(100 U/mL penicillin and 100 µg/mL streptomycin, Corning, EU) at 5% CO_2, maintained at 37°C
in a humidified incubator. This cell line was confirmed to be mycoplasma-free (MycoAlert,
Lonza).

Drugs

Carbenoxolone disodium salt was purchased from Sigma and 100 mmol/L stock solution was
prepared in H_2O. Drugs used for flow cytometry experiments were QVD-Oph (#OPH001, R&D
Systems, 10 mmol/L stock solution in DMSO), S63-845 (Chemieteck, 10 mmol/L stock solution
in DMSO), ABT737 (Selleckchem, 10 mmol/L stock solution in DMSO).

Construction of E and 3a proteins encoding plasmids

SARS-CoV-2 E and 3a nucleotide sequences, containing a Kozak sequence added right before the
ATG (RefSeq NC_045512.2) were synthesized by Eurofins (Ebersberg, EU) and subcloned into
the pIRES2 vector with eGFP in the second cassette (Takara Bio Europe, EU). Truncated Δ4 and
Δ8 E proteins, as well as Δ10 3a protein constructs, lacking the last 12, 24 and 30 last nucleotides,
respectively, were also synthesized by Eurofins. Plasmid cDNAs were systematically re-
sequenced by Eurofins after each plasmid in-house midiprep (Qiagen, EU).

E and 3a cDNA transfection

The Fugene 6 transfection reagent (Promega, WI, USA) was used to transfect WT and mutant E
and 3a plasmids for patch-clamp, morphology analysis and flow cytometry experiments
according to the manufacturer's protocol. Cells were cultured in 35-mm dishes and transfected,
at a 20% confluence for patch clamp experiments and 50% confluence for flow cytometry assay,
with a pIRES plasmid (2 µg DNA) with the first cassette empty or containing wild-type or
truncated SARS-CoV-2 E or 3a protein sequence. For morphology analysis, cells were cultured in
ibidi µ-Slide 8 well dishes and transfected at a 20% confluence with the same plasmids. In
pIRES2-eGFP plasmids, the second cassette (eGFP) is less expressed than the first cassette,
guaranteeing expression of high level of the protein of interest in fluorescent cells (28,29).

Electrophysiology

Two days after transfection, CHO cells were mounted on the stage of an inverted microscope
and bathed with a Tyrode solution (in mmol/L: NaCl 145, KCl 4, $MgCl_2$ 1, $CaCl_2$ 1, HEPES 5, glucose
5, pH adjusted to 7.4 with NaOH) maintained at 22.0 ± 2.0°C. Patch pipettes (tip resistance: 2.0
to 2.5 MΩ) were pulled from soda-lime glass capillaries (Kimble-Chase, USA) with a Sutter P-30
puller (USA). A fluorescent cell was selected by epifluorecence. The pipette was filled with
intracellular medium containing (in mmol/L): KCl, 100; Kgluconate, 45; $MgCl_2$, 1; EGTA, 5; HEPES,
10; pH adjusted to 7.2 with KOH. Stimulation and data recording were performed with pClamp
10, an A/D converter (Digidata 1440A) and a Multiclamp 700B (all Molecular Devices) or a VE-2
patch-clamp amplifier (Alembic Instruments). Currents were acquired in the whole-cell
configuration, low-pass filtered at 10 kHz and recorded at a sampling rate of 50 kHz. First, series
of twenty 30-ms steps to -80 mV were applied from holding potential (HP) of alternatively -70
mV and -90 mV to subsequently off-line calculate Cm and Rs values from the recorded currents.
Currents were then recorded using a 1-s ramp protocol from -80 mV to +70 mV, every 4 s (cf.
Figure 1). Regarding non-adhering cells, we considered them as with large current density when
the current density measured at +70 mV was superior to mean + 2 x standard deviation of the
current density in adhering cells, in the same condition.

Cell morphology assay

Cell roundness was estimated using the *Analyze Particle* function of the Fiji software, as described in suppl. Figure 1.

Flow cytometry assay

Two days after transfection, CHO cells were prepared to cell death detection following the user guide (https://assets.thermofisher.com/TFS-Assets/LSG/manuals/mp13199.pdf), to measure annexin V binding and propidium iodide (PI) intake. The cells were washed with cold PBS, trypsinized, collected by centrifugation and gently resuspended in annexin-binding buffer (V13246, Invitrogen, USA) at 1×10^6 cells/mL. To each 300-µL cell suspension were added 0.5 µL of annexin V AlexaFluor 647 (A23204, Invitrogen) and 1 µL of propidium iodide (PI) at 100 µg/mL (P3566, Invitrogen). CHO cells were incubated at room temperature for 15 minutes in the dark, and then maintained on ice until flow cytometry analysis within one hour.

To study the cell death pathways induced by E and 3a protein expression, non-transfected or transfected CHO cells were treated with inhibitors or inducers of apoptosis (inhibitor: 5 µmol/L QVD-OPh incubated for 48 h; activators: 3 µmol/L S63-845 + 8 µmol/L ABT737 incubated for 3 h).

The cytometer BD FACSCanto (BD Biosciences, USA) was used to sample acquisition. CHO cells transfected with an empty plasmid were used to determine the population to be analyzed. Monolabeled cells were used to establish the photomultiplier voltage of each channel (PMT) and to proceed with fluorescence compensation after the acquisitions. In order to detect cell death, only eGFP-positive CHO cells (FITC) were analyzed to Annexin V AlexaFluor 647 (APC) and PI (Perc-P) labeling. Analyses were performed using FlowJo software.

Acknowledgements

We thank the Agence Nationale de la Recherche for its financial support to the Région Pays de la Loire (ANR FLASH Covid-19 - CoV2-E-TARGET) and the laboratory of excellence 'Ion Channels, Science and Therapeutics' (grant No. ANR-11-LABX-0015). We acknowledge the IBISA MicroPICell facility (Biogenouest), member of the national infrastructure France-Bioimaging supported by the French national research agency (ANR-10-INBS-04). We thank the Cytometry Facility Cytocell from Nantes for expert technical assistance. We thank Hugues Abriel and Jean-Sebastien Rougier (Institute of Biochemistry and Molecular Medicine, university of Bern) for fruitful discussions and support and critical reading of the manuscript.

References

1. Drosten C, Günther S, Preiser W, van der Werf S, Brodt HR, Becker S, et al. Identification of a Novel Coronavirus in Patients with Severe Acute Respiratory Syndrome. N Engl J Med. 15 mai 2003;348(20):1967-76.

2. Zaki AM, van Boheemen S, Bestebroer TM, Osterhaus ADME, Fouchier RAM. Isolation of a Novel Coronavirus from a Man with Pneumonia in Saudi Arabia. N Engl J Med. 8 nov 2012;367(19):1814-20.

3. Burki T. The future of Paxlovid for COVID-19. The Lancet Respiratory Medicine. juill 2022;10(7):e68.

4. Ledford H. Hundreds of COVID trials could provide a deluge of new drugs. Nature. 3 mars 2022;603(7899):25-7.

5. Ison MG. Antivirals and resistance: influenza virus. Current Opinion in Virology. déc 2011;1(6):563-73.

6. Wilson L, Mckinlay C, Gage P, Ewart G. SARS coronavirus E protein forms cation-selective ion channels. Virology. déc 2004;330(1):322-31.

7. Wilson L, Gage P, Ewart G. Hexamethylene amiloride blocks E protein ion channels and inhibits coronavirus replication. Virology. sept 2006;353(2):294-306.

8. Torres J, Maheswari U, Parthasarathy K, Ng L, Liu DX, Gong X. Conductance and amantadine binding of a pore formed by a lysine-flanked transmembrane domain of SARS coronavirus envelope protein. Protein Sci. sept 2007;16(9):2065-71.

9. Verdiá-Báguena C, Nieto-Torres JL, Alcaraz A, DeDiego ML, Torres J, Aguilella VM, et al. Coronavirus E protein forms ion channels with functionally and structurally-involved membrane lipids. Virology. oct 2012;432(2):485-94.

10. Nieto-Torres JL, Verdiá-Báguena C, Jimenez-Guardeño JM, Regla-Nava JA, Castaño-Rodriguez C, Fernandez-Delgado R, et al. Severe acute respiratory syndrome coronavirus E protein transports calcium ions and activates the NLRP3 inflammasome. Virology. nov 2015;485:330-9.

11. Castaño-Rodriguez C, Honrubia JM, Gutiérrez-Álvarez J, DeDiego ML, Nieto-Torres JL, Jimenez-Guardeño JM, et al. Role of Severe Acute Respiratory Syndrome Coronavirus Viroporins E, 3a, and 8a in Replication and Pathogenesis. Denison MR, éditeur. mBio. 5 juill 2018;9(3):e02325-17.

12. McClenaghan C, Hanson A, Lee SJ, Nichols CG. Coronavirus Proteins as Ion Channels: Current and Potential Research. Front Immunol. 9 oct 2020;11:573339.

13. Pervushin K, Tan E, Parthasarathy K, Lin X, Jiang FL, Yu D, et al. Structure and Inhibition of the SARS Coronavirus Envelope Protein Ion Channel. Baric RS, éditeur. PLoS Pathog. 10 juill 2009;5(7):e1000511.

14. Nieto-Torres JL, DeDiego ML, Álvarez E, Jiménez-Guardeño JM, Regla-Nava JA, Llorente M, et al. Subcellular location and topology of severe acute respiratory syndrome coronavirus envelope protein. Virology. juill 2011;415(2):69-82.

15. Cabrera-Garcia D, Bekdash R, Abbott GW, Yazawa M, Harrison NL. The envelope protein of
SARS-CoV-2 increases intra-Golgi pH and forms a cation channel that is regulated by pH. J
Physiol. juin 2021;599(11):2851-68.

16. Breitinger U, Farag NS, Sticht H, Breitinger HG. Viroporins: Structure, function, and their
role in the life cycle of SARS-CoV-2. The International Journal of Biochemistry & Cell
Biology. avr 2022;145:106185.

17. Chan CM, Tsoi H, Chan WM, Zhai S, Wong CO, Yao X, et al. The ion channel activity of the
SARS-coronavirus 3a protein is linked to its pro-apoptotic function. The International
Journal of Biochemistry & Cell Biology. nov 2009;41(11):2232-9.

18. Lu W, Zheng BJ, Xu K, Schwarz W, Du L, Wong CKL, et al. Severe acute respiratory
syndrome-associated coronavirus 3a protein forms an ion channel and modulates virus
release. Proc Natl Acad Sci USA. 15 août 2006;103(33):12540-5.

19. Schwarz S, Wang K, Yu W, Sun B, Schwarz W. Emodin inhibits current through SARS-
associated coronavirus 3a protein. Antiviral Research. avr 2011;90(1):64-9.

20. Schwarz S, Sauter D, Wang K, Zhang R, Sun B, Karioti A, et al. Kaempferol Derivatives as
Antiviral Drugs against the 3a Channel Protein of Coronavirus. Planta Med. 23 janv
2014;80(02/03):177-82.

21. Ren Y, Shu T, Wu D, Mu J, Wang C, Huang M, et al. The ORF3a protein of SARS-CoV-2
induces apoptosis in cells. Cell Mol Immunol. août 2020;17(8):881-3.

22. Xia B, Shen X, He Y, Pan X, Liu FL, Wang Y, et al. SARS-CoV-2 envelope protein causes acute
respiratory distress syndrome (ARDS)-like pathological damages and constitutes an
antiviral target. Cell Res. août 2021;31(8):847-60.

23. Bao L, Locovei S, Dahl G. Pannexin membrane channels are mechanosensitive conduits for
ATP. FEBS Letters. 13 août 2004;572(1-3):65-8.

24. Senthivinayagam S, Serbulea V, Upchurch CM, Polanowska-Grabowska R, Mendu SK, Sahu
S, et al. Adaptive thermogenesis in brown adipose tissue involves activation of pannexin-1
channels. Molecular Metabolism. févr 2021;44:101130.

25. Bhat EA, Sajjad N. Human Pannexin 1 channel: Insight in structure–function mechanism
and its potential physiological roles. Mol Cell Biochem. mars 2021;476(3):1529-40.

26. Uchida K, Emoto K, Daleke DL, Inoue K, Umeda M. Induction of Apoptosis by
Phosphatidylserine. Journal of Biochemistry. 1 juin 1998;123(6):1073-8.

27. Crea F, Sarti D, Falciani F, Al-Rubeai M. Over-expression of hTERT in CHO K1 results in
decreased apoptosis and reduced serum dependency. Journal of Biotechnology. janv
2006;121(2):109-23.

28. Mizuguchi H, Xu Z, Ishii-Watabe A, Uchida E, Hayakawa T. IRES-Dependent Second Gene
Expression Is Significantly Lower Than Cap-Dependent First Gene Expression in a
Bicistronic Vector. Molecular Therapy. avr 2000;1(4):376-82.

29. Choveau FS, Rodriguez N, Ali FA, Labro AJ, Rose T, Dahimène S, et al. KCNQ1 Channels
Voltage Dependence through a Voltage-dependent Binding of the S4-S5 Linker to the Pore
Domain. Journal of Biological Chemistry. janv 2011;286(1):707-16.

30. Yu SP, Kerchner GA. Endogenous voltage-gated potassium channels in human embryonic
kidney (HEK293) cells. J Neurosci Res. 1 juin 1998;52(5):612-7.

31. Teoh KT, Siu YL, Chan WL, Schlüter MA, Liu CJ, Peiris JSM, et al. The SARS Coronavirus E
Protein Interacts with PALS1 and Alters Tight Junction Formation and Epithelial
Morphogenesis. Nusrat A, éditeur. MBoC. 15 nov 2010;21(22):3838-52.

32. Caillet-Saguy C, Durbesson F, Rezelj VV, Gogl G, Tran QD, Twizere J, et al. Host PDZ-
containing proteins targeted by SARS-CoV-2. FEBS J. sept 2021;288(17):5148-62.

33. Poon IKH, Chiu YH, Armstrong AJ, Kinchen JM, Juncadella IJ, Bayliss DA, et al. Unexpected
link between an antibiotic, pannexin channels and apoptosis. Nature. 20 mars
2014;507(7492):329-34.

34. Yuan X, Li J, Shan Y, Yang Z, Zhao Z, Chen B, et al. Subcellular localization and membrane
association of SARS-CoV 3a protein. Virus Research. mai 2005;109(2):191-202.

35. Nal B, Chan C, Kien F, Siu L, Tse J, Chu K, et al. Differential maturation and subcellular
localization of severe acute respiratory syndrome coronavirus surface proteins S, M and E.
Journal of General Virology. 1 mai 2005;86(5):1423-34.

36. Liao Y, Yuan Q, Torres J, Tam JP, Liu DX. Biochemical and functional characterization of the
membrane association and membrane permeabilizing activity of the severe acute
respiratory syndrome coronavirus envelope protein. Virology. juin 2006;349(2):264-75.

37. Cohen JR, Lin LD, Machamer CE. Identification of a Golgi Complex-Targeting Signal in the
Cytoplasmic Tail of the Severe Acute Respiratory Syndrome Coronavirus Envelope Protein.
J Virol. 15 juin 2011;85(12):5794-803.

38. Kern DM, Sorum B, Mali SS, Hoel CM, Sridharan S, Remis JP, et al. Cryo-EM structure of
SARS-CoV-2 ORF3a in lipid nanodiscs. Nat Struct Mol Biol. juill 2021;28(7):573-82.

39. Chekeni FB, Elliott MR, Sandilos JK, Walk SF, Kinchen JM, Lazarowski ER, et al. Pannexin 1
channels mediate 'find-me' signal release and membrane permeability during apoptosis.
Nature. oct 2010;467(7317):863-7.

40. Navis KE, Fan CY, Trang T, Thompson RJ, Derksen DJ. Pannexin 1 Channels as a Therapeutic
Target: Structure, Inhibition, and Outlook. ACS Chem Neurosci. 5 août
2020;11(15):2163-72.

41. Swayne LA, Johnstone SR, Ng CS, Sanchez-Arias JC, Good ME, Penuela S, et al.
Consideration of Pannexin 1 channels in COVID-19 pathology and treatment. American
Journal of Physiology-Lung Cellular and Molecular Physiology. 1 juill 2020;319(1):L121-5.

42. Luu R, Valdebenito S, Scemes E, Cibelli A, Spray DC, Rovegno M, et al. Pannexin-1 channel
opening is critical for COVID-19 pathogenesis. iScience. déc 2021;24(12):103478.

43. Nadeali Z, Mohammad-Rezaei F, Aria H, Nikpour P. Possible role of pannexin 1 channels
and purinergic receptors in the pathogenesis and mechanism of action of SARS-CoV-2 and
therapeutic potential of targeting them in COVID-19. Life Sciences. mai 2022;297:120482.