**Gene Gain and Loss from the Asian Corn Borer W Chromosome**

Wenting Dai[1], Judith E. Mank[2], Liping Ban[1, *]

[1] Department of Grassland Resources and Ecology, College of Grassland Science and

Technology, China Agricultural University, Beijing, China

[2] Department of Zoology and Biodiversity Research Centre, University of British

Columbia, Vancouver, Canada

[*] To whom correspondence should be addressed: liping_ban@163.com (LB).

**Abstract**

We built a chromosome-level genome assembly of the Asian corn borer, *Ostrinia*

*furnacalis* Guenée (Lepidoptera: Pyralidae, Pyraloidea), an economically important

pest in corn, from a female, including both the Z and W chromosome. Despite deep

conservation of the Z chromosome across Lepidoptera, our chromosome-level W

assembly reveals little conservation with available W chromosome sequence in

related species or with the Z chromosome, consistent with a non-canonical origin of

the W chromosome. The W chromosome has accumulated significant repetitive

elements and experienced rapid gene gain from the remainder of the genome, with

most genes exhibiting pseudogenization after duplication to the W. The genes that

retain significant expression are largely enriched for functions in DNA recombination,

the nucleosome, chromatin and DNA binding, likely related to meiotic and mitotic

processes within the female gonad.

**Introduction**

The non-recombining, sex-specific portions of the genome, namely Y and W chromosomes, exhibit very different properties from the remainder of the genome[1]. Recent advances in sequencing have made it possible to assemble full, or nearly full, sequences of numerous Y chromosomes[2], and these efforts have revealed general patterns such as the retention of genes related to dosage-sensitivity[3,4] and male function[5-8] as one might expect from a chromosome limited in its inheritance to males.

Studying Y chromosomes alone makes it difficult to differentiate the effects of sex-limitation from the effects of limitation to males, and so W chromosomes can be a powerful contrast to reveal the overall convergence of genomic patterns of these unusual regions of the genome. Despite providing an important comparison, sequencing of W chromosomes has lagged somewhat, with the most complete assemblies in birds[3,9] suggesting that although W chromosomes share some similarities with Y chromosomes, namely an abundance of repetitive elements[10], they may differ from Y chromosomes in that they lack genes with female-specific functions[3,9]. Whether this is unique to birds, or is a broader pattern of W chromosomes, remains unclear.

Lepidoptera, butterflies and moths, provide a key additional female heterogametic system[11,12]. The conservation of the Z chromosome has been well established in Lepidoptera[13], however, the W chromosome in Lepidoptera is unusual in that it was recruited into the genome well after the origin of the Z chromosome[14], as the basal lineages in the clade are Z0/ZZ. Available evidence suggests that, at least in some lineages, the W chromosome bears no homology to the Z[13,15,16], and may actually have originated from a B chromosome[13]. Within the Lepidoptera, at the base of the Ditrysia group, complex sex chromosomes, including neo-W chromosomes are observed in many clades based on cytogenetic analysis[14].

Recently, third-generation sequencing advances have permitted partial or

chromosome-level assemblies of the W chromosome in a limited number of

Lepidoptera, including *Cydia pomonella* (Torticidae, Tortricoideae)[16], *Cnaphalocrocis*

*medinalis* (Crambidae, Pyraloidea)[17], *Spodoptera exigua* (Noctuidea, Noctuoidea)[18],

*Spodoptera frugiperda* (Noctuidea, Noctuoidea)[19], *Trichoplusia ni* (Noctuidea,

Noctuoidea)[20] and *Dryas iulia* (Nymphalidae, Papilionoidea)[15]. This work has revealed

no detectable homology between the Z and W in *C. pomonella*[16], *T. ni*[21] and *D. iulia*[15].

In contrast, substantial synteny is observed between the W chromosomes of *S.*

*exigua* and *C. medinalis*[17,22]. All this is consistent with a non-canonical origin of the W

chromosome in Lepidoptera, where the W has been recruited from a B chromosome

and is therefore not homologous to the deeply conserved corresponding Z

chromosome.

To answer questions about the conservation and gene content of the lepidopteran

W, we built a chromosome-level genome of the Asian corn borer *Ostrinia furnacalis*

Guenée (Lepidoptera: Pyralidae, Pyraloidea), is a major insect pest of corn,

widespread in the Asian-Western Pacific region, using a combination of PacBio HiFi

(CCS) sequencing and Hi-C. We couple this with extensive RNA-Seq analysis of

multiple developmental stages and tissues in both sexes. Our catalog of W gene

content reveals extensive duplications from all other chromosomes in the genome,

with a relatively low percent of genes with persistent gene activity, which are

enriched for functions in DNA recombination, the nucleosome, chromatin and DNA

binding. Our results suggest that the W chromosome retains gene related to meiotic

and mitotic functions within the female gonad, in contrast to the avian W

chromosome[3,9] and more consistent with findings from Y chromosomes[5-8].

**Methods**

***Samples***

*O. furnacalis* larvae were collected from a corn field at Beijing, China, in July 2020,

and fed with an artificial diet in the laboratory of China Agricultural University. The

incubator environment was set at 26 ± 1℃ and 50 ± 5% relative humidity on a

photoperiod (Light: Dark = 16:8).

### Genome sequencing and assembly

Genomic DNA was extracted from a female laboratory-fed pupa. From this single

individual, we obtained 32.96 Gb circular consensus sequencing (CCS) data from one

cell of the PacBio Sequel II platform at Biomarker Co., Ltd., Qingdao, China (Table S1).

High accuracy PacBio Hifi data were assembled using hifiasm (v0.12) software to

obtain a draft genome[23]. We used purge_haplotigs[24] to remove redundant contigs

and generate a non-redundant assembled genome.

### Hi-C scaffolding

We constructed Hi-C libraries (300-700 bp insert size), following Rao et al.[25],

and sequenced them with pair-end 150 Illumina Hiseq at Biomarker Co., Ltd.,

Qingdao, China. 53.05 Gb of clean data were produced after filtering adapter

sequences, primer sequences and low-quality data (Table S2). The resulting trimmed

reads were aligned to the draft assembly with BWA (v0.7.10-r789), retaining only

uniquely aligned read pairs with mapping quality >20 for further analysis. Invalid read

pairs, including dangling-end, self-cycle, re-ligation and dumped products, were

removed by HiC-Pro (v2.8.1)[26].

The 56.25% of unique mapped read pairs represent valid interaction pairs and were

used for correction, clustering, ordering and orientation of scaffolds into

chromosomes with LACHESIS[27]. Before chromosome assembly, we performed a

preassembly for error correction of contigs which required the splitting of contigs

into segments of 50 kb on average. Then Hi-C data were mapped to these segments

using BWA (v0.7.10-r789). The uniquely mapped data were retained for the

assembly, and any two segments which showed inconsistent connection with

information from the raw scaffolds were checked manually. Parameters for running

LACHESIS included: CLUSTER_MIN_RE_SITES = 27; CLUSTER_MAX_LINK_DENSITY = 2;

ORDER_MIN_N_RES_IN_TRUNK = 15; ORDER_MIN_N_RES_IN_SHREDS = 15. After

this step, placement and orientation errors exhibiting obvious discrete chromatin

interaction patterns were manually adjusted. Finally, we constructed a heatmap

based on the interaction signals of valid mapped read pairs between chromosomes.

***Genome annotation***

Transposable elements (TEs) were identified by a combination of homology-based

and de novo approaches. We first carried out a de novo repeat prediction using

RepeatModeler2 (v2.0.1)[28], which implements RECON (v1.0.8)[29] and RepeatScout

(v1.0.6)[30]. Then the predicted results were classified using RepeatClassifier[28] by

means of repbase (v19.06)[31], REXdb (v3.0)[32] and Dfam (v3.2)[33]. Secondly, we

performed a de novo prediction for long terminal repeats (LTRs) using LTR_retriever

(v2.8)[34] via LTRharvest (v1.5.9)[35] and LTR_finder (v2.8)[36]. A non-redundant species-

specific TE library was constructed by combining the de novo TE library above with

the known databases. Final TE sequences in the *O. furnacalis* genome were identified

and classified by homology search against the library using RepeatMasker (v4.10)[37].

Tandem repeats were annotated by Tandem Repeats Finder (TRF v409)[38] and

MIcroSAtellite identification tool (MISA v2.1)[39].

We integrated de novo prediction, homology search, and transcript-based assembly

to annotate protein-coding genes. The de novo gene models were predicted using

Augustus (v2.4)[40] and SNAP (2006-07-28)[41], both ab initio gene-prediction

approaches. For the homolog-based approach, we used GeMoMa (v1.7)[42] with

reference gene models from *Drosophila melanogaster*, *Bombyx mori*, *Chilo*

*suppressalis* and *Galleria mellonella*. For the transcript-based prediction, RNA from a

mixture sample containing eggs, larvae, pupae and adults whole body of females and

males was used for generating sequencing libraries and was sequenced on an

Illumina NovaSeq 6000 platform (25.82-fold coverage of the genome, Table S3).

RNA-sequencing data were mapped to the reference genome using Hisat (v2.0.4)[43]

and assembled with Stringtie (v1.2.3)[44]. GeneMarkS-T (v5.1)[45] was used to predict

genes based on the assembled transcripts. PASA (v2.0.2)[46] was used to predict genes

based on the unigenes assembled by Trinity (v2.11)[47]. Gene models from these

different approaches were combined with EVM (v1.1.1)[48] and updated by PASA

(v2.0.2)[46]. The final gene models were annotated by searching the GenBank Non-

Redundant (NR, 20200921), TrEMBL (202005), Pfam (v33.1)[49], SwissProt (202005)[50],

eukaryotic orthologous groups (KOG, 20110125), gene ontology (GO, 20200615) and

Kyoto Encyclopedia of Genes and Genomes (KEGG, 20191220) databases.

We used tRNAscan-SE (v1.3.1)[51] with default parameters to identify transfer RNAs

(tRNAs), and barrnap (v0.9) with default parameters to identify the ribosomal RNAs

(rRNAs) based on Rfam (v12.0)[52]. miRNAs were identified with miRbase[53]. Small

nucleolar RNA (snoRNAs) and small nuclear RNA (snRNAs) were identified with

Infernal (v1.1.1)[54], using the Rfam (v12.0) database[52].

Pseudogenes have similar sequences to functional genes, but may have lost their

biological function because of mutations. GenBlastA (v1.0.4)[55] was used to scan the

whole genome after masking predicted functional genes. Putative candidates were

then analyzed by searching for premature stop codon and frameshift mutations using

GeneWise (v2.4.1)[56].

***Comparative genomics and phylogenetic reconstruction***

Protein sequence alignments between *O. furnacalis* and seven other lepidopteran

species (*B. mori*, *C. suppressalis*, *C. pomonella*, *C. medinalis*, *S. exigua*, *Spodoptera*

*fugiperda,* and *T. ni*) were performed with diamond (-e < 1e-5), then alignment

results were analyzed and the homologous chromosomal regions were identified

with MCScanX (MATCH_SCORE: 50, MATCH_SIZE: 5, GAP_PENALTY: -1,

OVERLAP_WINDOW: 5, E_VALUE: 1e-05, MAX GAPS: 15). The synteny relationships

among chromosomes were displayed using circos (v0.69–9).

We used the protein sequences of *O. furnacalis and* ten other insect species (*S.*

*frugiperda, Spodoptera litura, B. mori, C. suppressalis, Danaus plexippus, Melitaea*

*cinxia, Papilio Xuthus, C. pomonella*, *Plutella xylostella* and *D. melanogaster*) for

phylogenetic analysis, keeping only the longest transcript of each gene for analysis,

and using OrthoFinder (v2.5.4)[57] with default settings to identify orthologues and

homologues. To infer the phylogeny of these insects, multiple sequence alignments

of single-copy orthologs were performed using MAFFT (v7.471)[58]. Then we extracted

conserved sequences with gblocks (v0.91b)[59] and concatenated them to a single

sequence alignment. The resulting sequence alignment was used to construct a

maximum likelihood phylogenetic tree using IQ-TREE (v1.6.12)[60] (outgroup: *D.*

*melanogaster*). Divergence times between various species were estimated by

MCMCtree in PAML (v4.9j)[61]. Three standard divergence time points from the

TimeTree database (http://timetree.org/) were used for calibration: *S. frugiperda* – S.

*litura*, 100-122 million years ago (Mya), *D. plexippus* – *M. cinxia*, 76-102 Mya, (*D.*

*plexippus*, *M. cinxia*) – *P. xuthus*, 76-146 Mya. The tree was visualized using FigTree

(v1.4) (http://tree.bio.ed.ac.uk/software/figtree/). The gene count table from

OrthoFinder (v2.5.4) was used as inputs to examine the expansion and contraction of

each gene family in cafe (v5.1.0)[62].

***Detection of sex chromosomes***

In order to identify the sex chromosomes, we performed genome resequencing with

five female and male pupae. High-quality clean data (24.95-28.80-fold coverage of

the genome) were obtained through the pair-end 150 Illumina Hiseq platform at

Biomarker Co., Ltd., Qingdao, China (Table S3). The clean data were aligned to

reference genome with Bwa-men (v0.7.17). We compared the coverage differences

between male and female samples[63] to distinguish the Z, W and autosomes.

Specifically, we used the genomecov and groupby in Bedtools (v2.30.0) to obtain a

per-base median coverage depth for each chromosome and normalized them by the

mean of all chromosome median coverages for each sample. Normalized coverage

depth was averaged by sex to produce a coverage depth per chromosome for each

sex. Then we compared coverage depth between sexes for each chromosome and

calculated $\log_2$ male:female (M:F) coverage ratio [$\log_2$(M:F coverage)]. Autosomes

have an equal coverage between sexes [$\log_2$ (M:F coverage) = 0], while the Z

chromosome should show approximately two-fold greater coverage in the males

[$\log_2$(M:F coverage) = 1]. The W chromosome should show a strong female-biased

coverage pattern. We also calculated the M:F median coverage ratio along each

chromosome with nonoverlapping 1000 bp windows.

***W-gene homologs search and calculation of synonymous divergence ($d_S$)***

Predicted protein sequences were used to detect reciprocal best hits between the W

chromosome and the remainder of the genome using getRBH.pl

(https://github.com/Computational-conSequences/SequenceTools) for *O. furnacalis*.

For each pair of orthologous genes, we deleted stop codes and aligned the coding

sequences using macse (v2.05)[64], then extracted conserved sequences with gblocks

(v0.91b)[59]. The resulting alignments were used as inputs of CODEML in PAML (v4.9j)[61]

to estimate pairwise synonymous divergence with settings runmode = -2, seqtype = 1

and CodonFreq = 2. Since divergence estimates are not reliable for saturated sites,

we excluded orthologs with $d_S > 3$[65].

The homolog search of *O. furnacalis* W in *C. pomonella*, *C. medinalis*, *S. exigua*, *S.*

*fugiperda* and *T.ni* genome also used getRBH.pl. We assessed enrichment of Gene

Ontology[66,67] terms for W gene content.

***W-gene expression level***

We used two different RNA-Seq datasets to determine W expression, the mixture

sample of RNA-Seq data from eggs, larvae, pupae and adults used for genome

annotation (described above), and data from female gonad, as sex-limited

chromosomes are often largely expressed in the gonad. For female gonad, RNA from

five gonad samples of adult females was sequenced on an Illumina NovaSeq 6000

platform (10.14-fold coverage of the genome, Table S3). For each dataset separately,

we used fastp (v0.20.0)[68] to filter out low quality reads and to remove adapters with

the default parameters. Then we mapped the clean reads to the *O. furnacalis*

reference genome using HISAT2 (v2.1.0)[43]. The FPKM (fragments per kilobase of exon

per million fragments mapped) of each gene was determined using Stringtie

(v2.1.4)[44] based on the annotated GFF file. For female gonad, we used averaged

FPKM of five samples.

***Copy number variation of W and Z/autosome genes***

The amino acid sequences of protein-coding genes from whole genome were used as

the input of the blastp mapping against the Swiss-Prot database to assign gene

symbols (abbreviations for the gene names). We BLASTed each W gene to the

remainder of the genome to identify the best hit and only retained orthologous

genes on the W chromosome and Z/autosomes that consistently mapped to the

same protein (reciprocal best BLAST hit) and calculated their copy number following

the protocol in Mueller et al.[69].

**Results**

***Chromosome-level genome assembly of the Asian Corn Borer***

Our chromosome assembly strategy employed PacBio HiFi (CCS) sequencing data to

assemble the draft genome and Hi–C data to detect chromosomal contact

information. These PacBio long-reads were self-corrected using Quiver and

assembled into a draft genome assembly with a total length of 493.10Mb, consisting

of 57 contigs with an N50 length of 15.69Mb (Table S4). The assembled genome size

is similar to that obtained from genome surveys (Supplementary Figure 1). Next, Hi–C

linking information further anchored, ordered, and oriented 46 contigs to 32

chromosomes (30 autosomes, with Z and W sex chromosomes), which contained

86.79% of the whole genome assembly (Table S4, Supplementary Figure 2). The

chromosome-level genome assembly was 492.57Mb with a scaffold N50 of 16.47Mb

(Table S4).

We evaluated the completeness of *O. furnacalis* genome assembly with the

Metazoan Benchmark of Universal Single-Copy Orthologs (BUSCO v4) and CEGMA

core genes (CEGMA v2.5) datasets. Our assembly contained 97.69% of BUSCO genes,

of which 97.06% were single copy and 0.63% were duplicates, and 96.07% of CEGMA

genes, of which 79.03% were highly conserved (Table S5). To further evaluate the

genome assembly quality, genomic DNA sequencing data from Illumina HiSeq were

mapped to the assembly scaffolds, with a 99.07% coverage rate (Table S5). Finally,

lepidopteran genomes typically exhibit very high levels of synteny. Whole-genome

alignment of the *O. furnacalis* assembly to the chromosomes of the *S. litura* revealed

that chromosomal linkage and ordering of genes are highly conserved

(Supplementary Figure 3). All these analyses indicated that the genome assembly is

both highly reliable and complete for subsequent analyses.

***Genome annotation***

In total, 860,391 repeat sequences spanning 204.4Mb were identified, constituting

41.45% of the *O. furnacalis* genome (Table S4, Figure 1a). We integrated the result of

*ab initio*, homology-based and RNA-seq methods to annotate protein-coding genes,

most of the which came from homology-based and RNA-seq methods, indicating the

high-quality of annotation (Supplementary Figure 4). Finally, we obtained 16,509

protein coding genes in the *O. furnacalis* genome, which has similar gene features to

other lepidopteran genomes (gene length, gene number, coding length, intron length

of each gene, exon length) (Supplementary Figure 4). Next, we identified different

types of noncoding RNAs (ncRNAs), including 7,710 transfer RNAs (tRNA), 73

ribosomal RNAs (rRNA), and 39 microRNAs (miRNA) (Table S6). In addition, we

annotated 167 pseudogenes in the *O. furnacalis* genome, defined as any genomic

sequence that is similar to another gene but is defective, such as containing a

premature stop codon or a frameshifts mutation[70].  Most pseudogenes were located

in the chromosome LG3 (Figure 1c, Table S6), the W chromosome (see below). We

performed a functional annotation for all predicted protein coding genes using NCBI

nonredundant, EggNOG, GO, KEGG, SWISS-PROT and Pfam databases, with 98.21% of

our predicted genes assigned to at least one of these databases (Table S6).

**Fig 1. Genomic characterization of *Ostrinia furnacalis*.** a. Circos plot depicting the genomic landscape of the 32 chromosomes (Chr1–Chr32 on a Mb scale). The denotation of each track is listed in the center of the circle.  Blue lines in LG3 (the W chromosome) represent collinearity within the W, due to repeat elements. b. Number and distribution of protein-coding genes. c. Number and distribution of pseudogenes. d. Number of expressed genes (FPKM > 0.5 in mixed sample). e. Average chromosomal expression level (excluding genes with FPKM ≤ 0.5 in mixed sample).

We used OrthoFinder to find orthologous genes among *O. furnacalis* and ten other insect species (see Materials and Methods). A total of 16231 orthologous groups with 1163 single-copy orthologous genes were identified. We inferred a phylogenetic tree and divergence time estimate using the end-to-end concatenated amino acid sequences of 1163 single-copy orthologous genes. The Crambidae lineage to which *O. furnacalis* belongs was estimated to diverge from Obtectomeran Lepidoptera approximately 77.3 Mya ago (Figure 2a).

**Fig 2. a.** Phylogenetic tree and gene orthology of *O. furnacalis* with nine lepidopteran genomes and *D. melanogaster*. Comparative analysis of synteny between *O. furnacalis* and **b.** *C. suppressalis*, **c.** *B. mori*, **d.** *C. medinalis*, **e.** *C. pomonella,* **f.** *S.*

*exigua,* **g.** *S. fugiperda*, and **h.** *T. ni.* The chromosomes of *O. furnacalis* are shown in
the left, and the other insects' chromosomes are shown in the right. The *O.*
*furnacalis* W chromosome is LG3, and the Z is LG1 (b-h).

**Synteny, karyotype evolution, and sex chromosomes**

We compared the syntenic relationships between *O. furnacalis* and other

lepidopterans*,* including *B. mori, C. suppressalis, C. medinalis*, *S. litura, S. exigua, S.*

*fugiperda, and T. ni* (Figure 2b-h)*.* In general*,* the *O. furnacalis* genome shows a high

level of synteny with other lepidopteran genomes, though with some fusion and

fission events. *O. furnacalis* LG1 is syntenic with the Z chromosome in all other

species, suggesting it is the Z chromosome in *O. furnicalis* as well, and this is also

evident from M:F coverage analysis (Figure 3a), which reveals two-fold greater

coverage in males (ZZ) compared to females (ZW).

*O. furnacalis* LG3, the W chromosome, exhibits female-biased coverage consistent

with a female-limited chromosome (Figure 3a). The M:F coverage is highly variable

across chromosome LG3 compared to all other chromosomes (Figure 3b), due in

large part to the abundance of TEs on this chromosome (Supplementary Figure 5,

Supplementary Figure 6). The W chromosome (LG3) is the second largest

chromosome (22.23 Mb) in our assembly (Figure 1a), with the largest number of

predicted protein coding genes (excluding pseudogenes) compared with other

chromosomes (Figure 1b). The W also has the largest number of pseudogenes (Figure

1c), and contains 43.1% of all pseudogenes annotated in the genome. Many genes

that are not technically pseudogenized were expressed below our 0.5 FPKM

threshold. Only 66 protein coding genes showed significant expression above this

threshold in our mixed sample (comprised of eggs, larvae, pupae and adults, see

Materials and Methods, Figure 1d, 1e). 48 W genes showed expression >0.5 FPKM in

adult female gonads.

Furthermore, LG3 has a notably greater repeat density and significantly different

repeat composition compared to other chromosomes (Supplementary Figure 5, Supplementary Figure 6) with a larger proportion of satellites, DIRS, LINE, Copia and Gypsy, and an enrichment of *Maverick* elements.

**Fig 3. Detection of sex chromosomes. a.** Male:female coverage ratios for each chromosome, plotted by chromosome length. Each point represents a single chromosome. The dashed gray line is the theoretical expectation for autosomes and the dashed red line shows the expectation for the Z chromosome. **b.** Male:female coverage ratios plotted in 500bp windows across scaffolds for Z, W, and a representative autosome (LG32).

***W homology***

Given that previous reports have found no evidence of homology between the Z and

W in some basal lepidopteran species[15,16,21], we next investigated the evolutionary

history of the *O. furnacalis* W chromosome. Using our W gene set, we first examined

homology between the *O. furnacalis* W chromosome and the W chromosome from

other lepidopterans. The *O. furnacalis* W shows substantial homology with the W in

*C. medinalis*, consistent with some form of shared ancestry within the family

Crambidae. However, we observed no discernable homology between the *O.*

*furnacalis* W and the W chromosome of any of the more distantly related

lepidopterans that we queried (Figure 2b-h).

Reciprocal best BLAST hits between the W coding catalog and the remainder of the

*O. furnicalis* genome reveals similarity throughout the genome (Figure 4), consistent

with a B chromosome origin of the W chromosome followed by gene duplication

from locations throughout the genome. Of the 482 W coding genes having

Z/autosome paralogs (Figure 4), 202 exhibited fewer or no introns in the W paralogs,

consistent with retrotransposition.

**Fig. 4 The reciprocal best hits search for W (LG3) genes.** W chromosome (LG3) is

marked in red and Z chromosome (LG1) in blue, mt represent the mitochondrial

genome of *O. furnacalis.* Light blue line: identity ≥ 90%; purple line: 80% ≤ identity <

90%; light grey line: identity < 80%.

We next examined synonymous divergence ($d_S$) for these paralogs in order to

determine the timing of gene duplication to the W chromosome. The density of $d_S$

(Figure 5a) shows that most paralogs with W genes have relatively low divergence,

and the distribution of paralog $d_S$ across the W chromosome suggests that

duplications occur randomly throughout the chromosome. Given the low percent of

genes on the W chromosome that exhibit expression >0.5 FPKM and the high

number of pseudogenes on the W (Figure 1), it is likely that most duplicates to the W

chromosome fail to maintain significant expression and are silenced relatively

quickly.

**Fig 5. Synonymous divergence between W gene and best BLAST hit gene from the**

**remainder of the genome**. a. Density plot, b. Distribution across the W chromosome.

Sex-limited chromosomes often contain many copies of some genes[5,71], and so we

examined copy number for genes on the W chromosome and their most similar

paralog (Figure 6, Table S7). Overall, we observed lower number of gene copies on

the W than for the autosomal/Z chromosome paralog.

**Fig. 6. Copy number for W and autosomal/Z chromosome paralogs.**

Given the rapid apparent decay of W chromosome paralogs (Figure 5), we examined

the Gene Ontology enrichment for W expressed genes (>0.5 FPKM) (Tables 1 and 2),

and observed statistical enrichment of terms related to functions in DNA

recombination, the nucleosome, chromatin and DNA binding. Together, these results

suggest that many genes retained with functional expression on the W chromosome

are related to mitotic and meiotic processes, much of it within the female gonad.

**Table 1. Functional Enrichment of Significantly Expressed Genes (FPKM > 0.5 from**
**mixed sample) on *O. furnicalus* W chromosome**

| PANTHER GO | FDR | Expressed W genes |
|---|---|---|
| **Biological** | | |
| Negative regulation of DNA recombination | 7.64 e-11 | 5 |

| | 8.35 e -10 | 5 |
|---|---|---|
| Nucleosome assembly | 8.35 e -10 | 5 |
| Chromosome condensation | 2.48 e-8 | 5 |
| Telomere maintenance | 2.67 e-02 | 2 |
| **Molecular** | | |
| Nucleosomal DNA binding | 1.16 e-11 | 5 |
| Structural constituent of chromatin | 1.13 e-10 | 5 |
| Double-stranded DNA binding | 1.66 e-7 | 7 |
| **Cellular** | | |
| Nucleosome | 1.68 e-10 | 5 |
| Chromatin | 6.12 e-7 | 6 |

**Table 2. Functional Enrichment of Significantly Expressed Genes (FPKM < 0.5 from female gonad) on *O. furnicalus* W chromosome**

| PANTHER GO Term | FDR | Expressed W genes |
|---|---|---|
| **Biological** | | |
| Negative regulation of DNA recombination | 2.04e-10 | 5 |
| Nucleosome assembly | 2.22e-9 | 2 |
| Chromosome condensation | 7.70e-08 | 5 |
| **Molecular** | | |
| Nucleosomal DNA binding | 3.09e-11 | 5 |
| Structural constituent of chromatin | 3.02e-10 | 5 |
| Double-stranded DNA binding | 8.28e-05 | 6 |
| **Cellular** | | |
| Nucleosome | 4.46e-10 | 5 |
| Chromatin | 3.19e-6 | 6 |

**Discussion**

Here we report a high quality, chromosome-level genome for *O. furnicalis*. Our

reference genome includes a single, contiguous sequence for the female-specific W

chromosome, allowing us to query the content of this unique region of the genome.

W and Y chromosomes are often enriched for pseudogenes, as the lack of

recombination in these regions leads to high rates of gene silencing[1]. At the same

time, these regions often accumulate repeat sequences[9,72], resulting in significantly

higher repeat content compared to other chromosomes[9,16,17]. Consistent with this,

we although we found that the number of annotated protein-coding genes of W

chromosome is indeed the largest of all the chromosomes in the genome (Figure 1),

most of these are pseudogenes (Figure 1c), and those genes that retain functional

coding sequence have low overall transcriptional activity (Figure 1d)[12].

We also found the W is enriched for repetitive elements in *O. furnicalis*

(Supplementary Figure 5 and Supplementary Figure 6), with the number of *Maverick*

elements particularly higher on the W chromosome compared to the rest of the

genome (Supplementary Figure 5). *Maverick*, also known as *Polinton*, is a DNA

transposon widespread in eukaryotes[73], and one of the most striking characteristics

of *Maverick* is the large size, typically 15–20 kb[73].

**Conservation of sex chromosomes across Lepidoptera species**

420    We searched orthologs to Z-linked genes of *O. furnacalis* in other lepidopterans, and

421    found that although the Z chromosome shows clear strong conservation[14] (Figure 2)

422    the *O. furnacalis* W chromosome is only conserved with *C. medinalis*, also a member

423    of Crambidae (Figure 2b-h). This is consistent with previous work suggesting that the

424    composition of the W varies even among species with the same family[74].

**Evolutionary history of the sex chromosomes in *O. furnacalis***

Karyotype work has revealed a complex history of the W chromosome, including the

repeated origin of neo-W chromosomes in many lepidopteran lineages. The

evolutionary history of the W chromosome in Lepidoptera differs from the canonical

model of sex chromosome formation in that it was recruited, possibly from a B

element, after the formation of the Z chromosome[14], at the common root of Ditrysia

and Tischerioidea[12,75]. Lineages ancestral to this recruitment exhibit Z0:ZZ

karyotypes[14]. The B-chromosome origin of the W is supported by the fact that in

some basal lineages, W chromosome bears no homology to the Z[13,15,16].

We observe little similarity in gene content between the W and Z chromosome

(Figure 4), a steady rate of gene duplication to the W chromosome from throughout

the genome (Figure 5), and reduced intron number for many W paralogs compared

to their most similar autosomal/Z-linked copy suggests that this likely often occurs through retrotransposition across the W chromosome. Genes retrotransposed or otherwise duplicated to the W chromosome will immediately experience the full degenerative effects of a non-recombining region[1,76], and consistent with this, we observe high levels of pseudogenization (Figure 1) on the W chromosome. However, those genes on the W that retain functional coding sequence and expression (e.g. non-pseudogenized) are enriched for mitotic and meiotic functions, and display Gene Ontology term enrichments related to recombination, chromosome packaging and replication (Tables 1 and 2). Although it has been suggested that W chromosomes may differ from Y chromosomes in that they do not acquire genes related to female gonadal function[3,9] in the same way that Y chromosomes retain genes related to male function[5-8], our results suggest that this may not be a generalized pattern of W chromosomes. Indeed, the W chromosome in *O. furnicalis* are enriched for functions related to female meiosis and mitosis.

**Authors' contributions**

LB conceived and designed the research. WD performed the experiments, analyzed and interpreted the data, and wrote the manuscript. JEM helped analyze the data and write the manuscript. All authors read and approved the final manuscript.

**Acknowledgments**

This work was supported by the Beijing Agriculture Innovation Consortium (BAIC02-2023) and National Key R&D Program of China (2022YFD1400500). JEM gratefully acknowledges support from NSERC and a Canada 150 Research Chair. This work was supported by the high-performance computing platform of China Agricultural University.

**Competing interests**

The authors declare that they have no competing interests.

**References**

1. Bachtrog, D. Y-chromosome evolution: emerging insights into processes of Y-chromosome degeneration. *Nature Reviews Genetics* **14**, 113-124 (2013).

2. Tomaszkiewicz, M., Medvedev, P. & Makova, K.D. Y and W chromosome assemblies: approaches and discoveries. *Trends in Genetics* **33**, 266-282 (2017).

3. Bellott, D.W. et al. Avian W and mammalian Y chromosomes convergently retained dosage-sensitive regulators. *Nature Genetics* **49**, 387-394 (2017).

4. Bellott, D.W. & Page, D.C. Dosage-sensitive functions in embryonic development drove the survival of genes on sex-specific chromosomes in snakes, birds, and mammals. *Genome Research* **31**, 198-210 (2021).

5. Rhie, A. et al. The complete sequence of a human Y chromosome. *bioRxiv* (2022).

6. Cechova, M. et al. Dynamic evolution of great ape Y chromosomes. *PNAS* **117**, 26273-26280 (2020).

7. Koerich, L.B., Wang, X., Clark, A.G. & Carvalho, A.B. Low conservation of gene content in the *Drosophila* Y chromosome. *Nature* **456**, 949-951 (2008).

8. Tobler, R., Nolte, V. & Schlötterer, C. High rate of translocation-based gene birth on the *Drosophila* Y chromosome. *Proceedings of the National Academy of Sciences* **114**, 11721-11726 (2017).

9. Smeds, L. et al. Evolutionary analysis of the female-specific avian W chromosome. *Nature Communications* **6**, 7330 (2015).

10. Peona, V. et al. The avian W chromosome is a refugium for endogenous retroviruses with likely effects on female-biased mutational load and genetic incompatibilities. *PHILOSOPHICAL TRANSACTIONSOF THE ROYAL SOCIETY B* **376**, 20200186 (2021).

11. Blackmon, H., Ross, L. & Bachtrog, D. Sex determination, sex chromosomes, and karyotype evolution in insects. *Journal of Heredity* **108**, 78-93 (2016).

12. Traut, W. & Marec, F. Sex chromatin in Lepidoptera. *The Quarterly Review of Biology* **2**, 239-259 (1996).

13. Fraïsse, C., Picard, M.A.L. & Vicoso, B. The deep conservation of the Lepidoptera Z chromosome suggests a non-canonical origin of the W. *Nature Communications* **8**, 1486 (2017).

14. Sahara, K., Yoshido, A. & Traut, W. Sex chromosome evolution in moths and butterflies. *Chromosome Research* **20**, 83-94 (2012).

15. Lewis, J.J. et al. The *Dryas iulia* genome supports multiple gains of a W chromosome from a B chromosome in butterflies. *Genome Biology and Evolution* **13**, evab128 (2021).

16. Wan, F. et al. A chromosome-level genome assembly of *Cydia pomonella* provides insights into chemical ecology and insecticide resistance. *Nature Communications* **10**, 4237 (2019).

17. Zhao, X. et al. A chromosome−level genome assembly of rice leaffolder, *Cnaphalocrocis medinalis*. *Molecular Ecology Resources* **21**, 561-572 (2021).

18. Zhang, F., Zhang, J., Yang, Y. & Wu, Y. A chromosome-level genome assembly for the beet armyworm (*Spodoptera exigua*) using PacBio and Hi-C sequencing. *bioRxiv* (2020).

19. Xiao, H. et al. The genetic adaptations of fall armyworm *Spodoptera frugiperda* facilitated its rapid global dispersal and invasion. *Molecular Ecology Resources* **20**, 1050-1068 (2020).

20. Chen, W. et al. A high-quality chromosome−level genome assembly of a generalist herbivore, *Trichoplusia ni*. *Molecular Ecology Resources* **19**, 485-496 (2019).

21. Fu, Y. et al. The genome of the Hi5 germ cell line from *Trichoplusia ni*, an agricultural pest and novel model for small RNA biology. *eLife* **7**, e31628 (2018).

22.   MarecKen, F., Sahara, S. & Traut, W. Rise and fall of the W chromosome in Lepidoptera. in
*Molecular biology and genetics of the Lepidoptera* (eds. MR, G. & F, M.) 49−63 (CRC, Boca Raton,
2010).

23.   Cheng, H., Concepcion, G.T., Feng, X., Zhang, H. & Li, H. Haplotype-resolved de novo assembly
using phased assembly graphs with hifiasm. *Nature Methods* **18**, 170-175 (2021).

24.   Roach, M.J., Schmidt, S.A. & Borneman, A.R. Purge Haplotigs: allelic contig reassignment for
third-gen diploid genome assemblies. *BMC Bioinformatics* **19**, 460 (2018).

25.   Rao, S.S.P. et al. A 3D map of the human genome at kilobase resolution reveals principles of
chromatin looping. *Cell* **162**, 687-688 (2015).

26.   Servant, N. et al. HiC-Pro: an optimized and flexible pipeline for Hi-C data processing. *Genome*
*Biology* **16**, 259 (2015).

27.   Burton, J.N. et al. Chromosome-scale scaffolding of de novo genome assemblies based on
chromatin interactions. *Nature Biotechnology* **31**, 1119-1125 (2013).

28.   Flynn, J.M. et al. RepeatModeler2 for automated genomic discovery of transposable element
families. *PNAS* **117**, 9451-9457 (2020).

29.   Bao, Z. & Eddy, S.R. Automated de novo identification of repeat sequence families in sequenced
genomes. *Genome Res* **12**, 1269-76 (2002).

30.   Price, A.L., Jones, N.C. & Pevzner, P.A. De novo identification of repeat families in large genomes.
*Bioinformatics* **21**, i351-i358 (2005).

31.   J, J. et al. Repbase Update, a database of eukaryotic repetitive elements. *Cytogenetic and Genome*
*Research* **110**, 462-467 (2005).

32.   Neumann, P., Novák, P., Hoštáková, N. & Macas, J. Systematic survey of plant LTR-
retrotransposons elucidates phylogenetic relationships of their polyprotein domains and provides a
reference for element classification. *Mobile DNA* **10**, 1 (2019).

33.   Wheeler, T.J. et al. Dfam: a database of repetitive DNA based on profile hidden Markov models.
*Nucleic Acids Research* **41**, D70-D82 (2012).

34.   Ou, S. & Jiang, N. LTR_retriever: a highly accurate and sensitive program for identification of long
terminal repeat retrotransposons. *Plant Physiology* **176**, 1410-1422 (2018).

35.   Ellinghaus, D., Kurtz, S. & Willhoeft, U. LTRharvest, an efficient and flexible software for de novo
detection of LTR retrotransposons. *BMC Bioinformatics* **9**, 18 (2008).

36.   Xu, Z. & Wang, H. LTR_FINDER: an efficient tool for the prediction of full-length LTR
retrotransposons. *Nucleic Acids Research* **35**, W265-W268 (2007).

37.   Tarailo-Graovac, M. & Chen, N. Using RepeatMasker to identify repetitive elements in genomic
sequences. *Curr. Protoc. Bioinform* **25**, 4.10.1-4.10.14 (2009).

38.   Benson, G. Tandem repeats finder: a program to analyze DNA sequences. *Nucleic Acids Research*
**27**, 573−580 (1999).

39.   Beier, S., Thiel, T., Munch, T., Scholz, U. & Mascher, M. MISA-web: a web server for
microsatellite prediction. *Bioinformatics* **33**, 2583-2585 (2017).

40.   Stanke, M., Diekhans, M., Baertsch, R. & Haussler, D. Using native and syntenically mapped
cDNA alignments to improve de novo gene finding. *Bioinformatics* **24**, 637-644 (2008).

41.   Korf, I. Gene finding in novel genomes. *BMC Bioinformatics* **5**, 59 (2004).

42.   Keilwagen, J. et al. Using intron position conservation for homology-based gene prediction. *Nucleic*
*Acids Research* **44**, e89-e89 (2016).

43.   Kim, D., Langmead, B. & Salzberg, S.L. HISAT: a fast spliced aligner with low memory

requirements. *Nature Methods* **12**, 357-360 (2015).

44. Pertea, M. et al. StringTie enables improved reconstruction of a transcriptome from RNA-seq reads.

*Nature Biotechnology* **33**, 290-295 (2015).

45. Tang, S., Lomsadze, A. & Borodovsky, M. Identification of protein coding regions in RNA

transcripts. *Nucleic Acids Research* **43**, e78-e78 (2015).

46. Haas, B.J. Improving the *Arabidopsis* genome annotation using maximal transcript alignment

assemblies. *Nucleic Acids Research* **31**, 5654-5666 (2003).

47. Grabherr, M.G. et al. Full-length transcriptome assembly from RNA-Seq data without a reference

genome. *Nature Biotechnology* **29**, 644-652 (2011).

48. Haas, B.J. et al. Automated eukaryotic gene structure annotation using EVidenceModeler and the

program to assemble spliced alignments. *Genome Biol* **9**, R7 (2008).

49. Finn, R.D. Pfam: clans, web tools and services. *Nucleic Acids Research* **34**, D247-D251 (2006).

50. Boeckmann, B. The SWISS-PROT protein knowledgebase and its supplement TrEMBL in 2003.

*Nucleic Acids Research* **31**, 365-370 (2003).

51. Lowe, T.M. & Eddy, S.R. tRNAscan-SE: a program for improved detection of transfer RNA genes

in genomic sequence. *Nucleic Acids Research* **25**, 955−964 (1997).

52. Griffiths-Jones, S. Rfam: annotating non-coding RNAs in complete genomes. *Nucleic Acids*

*Research* **33**, D121-D124 (2004).

53. Griffiths-Jones, S. miRBase: microRNA sequences, targets and gene nomenclature. *Nucleic Acids*

*Research* **34**, D140-D144 (2006).

54. Nawrocki, E.P. & Eddy, S.R. Infernal 1.1: 100-fold faster RNA homology searches. *Bioinformatics*

**29**, 2933-2935 (2013).

55. She, R., Chu, J.S.C., Wang, K., Pei, J. & Chen, N. genBlastA: Enabling BLAST to identify

homologous gene sequences. *Genome Research* **19**, 143-149 (2009).

56. Birney, E., Clamp, M. & Durbin, R. GeneWise and Genomewise. *Genome Research* **14**, 988-995

(2004).

57. Emms, D.M. & Kelly, S. OrthoFinder: phylogenetic orthology inference for comparative genomics.

*Genome Biology* **20**, 238 (2019).

58. Katoh, K. & Standley, D.M. MAFFT multiple sequence alignment software version 7:

improvements in performance and usability. *Molecular Biology and Evolution* **30**, 772-780 (2013).

59. Castresana, J. Selection of conserved blocks from multiple alignments for their use in phylogenetic

analysis. *Mol Biol Evol* **17**, 540-52 (2000).

60. Nguyen, L., Schmidt, H.A., von Haeseler, A. & Minh, B.Q. IQ-TREE: a fast and effective stochastic

algorithm for estimating maximum-likelihood phylogenies. *Molecular biology and evolution* **32**, 268-

274 (2015).

61. Yang, Z. PAML 4: phylogenetic analysis by maximum likelihood. *Molecular Biology and*

*Evolution* **24**, 1586-1591 (2007).

62. Mendes, F.K., Vanderpool, D., Fulton, B. & Hahn, M.W. CAFE 5 models variation in evolutionary

rates among gene families. *Bioinformatics* **36**, 5516-5518 (2020).

63. Mongue, A.J., Nguyen, P., Voleníková, A. & Walters, J.R. Neo-sex chromosomes in the monarch

butterfly, *Danaus plexippus. G3 Genes|Genomes|Genetics* **7**, 3281−3294 (2017).

64. Ranwez, V., Harispe, S., Delsuc, F., Douzery, E.J.P. & Murphy, W.J. MACSE: multiple alignment

of coding sequences accounting for frameshifts and stop codons. *PloS one* **6**, e22594-e22594 (2011).

65. Mank, J.E., Axelsson, E. & Ellegren, H. Fast-X on the Z: Rapid evolution of sex-linked genes in

birds. *Genome Research* **17**, 618-624 (2007).

66. Ashburner, M. et al. Gene Ontology: tool for the unification of biology. *Nature Genetics* **25**, 25−29 (2000).

67. Carbon, S. et al. The Gene Ontology resource: enriching a GOld mine. *Nucleic Acids Research* **49**, D325-D334 (2021).

68. Chen, S., Zhou, Y., Chen, Y. & Gu, J. fastp: an ultra-fast all-in-one FASTQ preprocessor. *Bioinformatics* **34**, i884−i890 (2018).

69. Mueller, J.C. et al. Micro germline-restricted chromosome in blue tits: evidence for meiotic functions. *Molecular Biology and Evolution* **40**, msad096 (2023).

70. Cheetham, S.W., Faulkner, G.J. & Dinger, M.E. Overcoming challenges and dogmas to understand the functions of pseudogenes. *Nature Reviews Genetics* **21**, 191−201 (2020).

71. Rogers, T.F., Pizzari, T. & Wright, A.E. Multi-copy gene family evolution on the avian W chromosome. *Journal of Heredity* **112**, 250-259 (2021).

72. H, A., K, M., Y, Y., T, O. & T, S. Retrotransposable elements on the W chromosome of the silkworm, *Bombyx mori*. *Cytogenetic and Genome Research* **110**, 144-151 (2015).

73. Pritham, E.J., Putliwala, T. & Feschotte, C. Mavericks, a novel class of giant transposable elements widespread in eukaryotes and related to DNA viruses. *Gene* **390**, 3-17 (2007).

74. Vítková, M., Fuková, I., Kubíčková, S. & Marec, F. Molecular divergence of the W chromosomes in pyralid moths (Lepidoptera). *Chromosome Research* **15**, 917-930 (2007).

75. Lukhtanov, V.A. Sex chromatin and sex chromosome systems in nonditrysian Lepidoptera (Insecta). *Journal of zoological systematics and evolutionary research* **38**, 73-79 (2000).

76. Bachtrog, D. et al. Sex determination: why so many ways of doing it. *PLoS biology* **12**, e1001899 (2014).