Zebrafish Rif1 Impacts Zygotic Genome Activation, Replication Timing, and Sex Determination

Emily A. Masser[1,2], Tyler D. Noble[1,2], Joseph C. Siefert[1,2], Duane Goins[2], Courtney G. Sansam[2], Christopher L. Sansam[1,2]

[1] University of Oklahoma Health Sciences Center, Department of Cell Biology, Oklahoma City, OK
[2] Oklahoma Medical Research Foundation, Cell Cycle and Cancer Biology Research Program, Oklahoma City, OK

ORCID IDs
Emily A. Masser: https://orcid.org/0000-0002-7934-7175
Tyler D. Noble: https://orcid.org/0000-0002-6073-7691
Joseph C. Siefert: https://orcid.org/0000-0001-5584-8672
Duane Goins: https://orcid.org/0000-0003-3154-9007
Courtney G. Sansam: https://orcid.org/0000-0002-0253-8643
Christopher L. Sansam: https://orcid.org/0000-0002-0545-2693

ABSTRACT

Deregulated DNA replication causes human developmental disorders and cancer, but we know little about how DNA replication is coordinated with changes in transcription and chromatin structure. The initiation of replication forks follows a spatiotemporal pattern called the replication timing program. We have developed the zebrafish into a model system to study the mechanisms by which the replication timing program changes during the extensive changes in the cell cycle, transcription, chromatin organization, and nuclear structure that occur during development. Our previous studies identified changes in DNA replication timing patterns occurring from the onset of zygotic transcription through gastrulation in zebrafish embryos. Rif1 is required for DNA replication timing in a wide range of eukaryotes. The broader role of Rif1 in establishing the replication timing program and chromatin structure during early vertebrate development remains unknown. We have generated Rif1 mutant zebrafish and have performed RNA sequencing and whole-genome replication timing analyses on multiple developmental stages. Rif1 mutants were viable but had a defect in female sex determination. Surprisingly, Rif1 loss predominantly affected DNA replication timing after gastrulation, while its impacts on transcription were more substantial during zygotic genome activation. Our results indicate that Rif1 has distinct roles in DNA replication and transcription control that manifest at different stages of development.

INTRODUCTION

The initiation of DNA replication follows a pattern where different genomic regions replicate at distinct times throughout S-phase. This temporal pattern is known as the replication-timing program [5]. Replication timing (RT) correlates with chromatin organization and gene activity [6]. In multicellular organisms, early replication often correlates with active transcription; whereas, late replication frequently coincides with transcriptionally silent genes [7-10]. In mammals, pluripotent cells exhibit a replication-timing pattern that changes with cellular differentiation; in addition, the switch in RT is related to changes in epigenetic modifications, transcriptional activity and the three-dimensional (3D) organization of chromatin [5-7, 11, 12]. The causal relationships between DNA replication and developmental changes, such as transcription, chromatin structure and 3D chromatin organization have been unclear.

The Rif1 (Rap1-interacting factor 1) protein is of particular interest because it has been implicated in the regulation of transcription, chromatin structure, 3D chromatin organization, DNA repair, and DNA replication timing. In fact, Rif1 is the single factor known to influence RT in all eukaryotes [5, 13-16]. Through conserved motifs, Rif1 has been shown to interact physically with the PP1 phosphatase [15, 17-20]. Rif1 controls RT by bringing PP1 to late-replicating regions of the genome, thereby causing de-phosphorylation of DNA replication factors during early S-phase and delaying initiation [15-19, 21, 22].

Our understanding of how Rif1 affects RT during embryonic development is incomplete. RT is developmentally regulated through Rif1 in *D. melanogaster* [23]. In early fly embryos, Rif1-activation initiates late replication of satellite repeats and causes the first lengthening of the cell cycle [23]. In Xenopus laevis, Rif1 knockdown accelerates early embryonic cell cycles and disrupts RT patterns in retinal progenitor cells [24]. Loss of Rif1 in mice is embryonic lethal with partial penetrance [25-28]. Cell lines derived from Rif1 mutant mice display altered RT profiles, but the extent to which RT is disrupted in the developing embryo has not been shown. Although Rif1 loss causes RT changes in cell lines from a wide range of species, it is unknown whether

Rif1 plays roles in the establishment or change of the RT program during vertebrate

embryogenesis.

We have developed the zebrafish into a model system to study how modifications in RT

coincide with extensive changes in cell cycle and transcription that occur in the developing

embryo. The earliest stages of zebrafish development have been extensively studied, so it is

possible to collect large numbers of synchronously developing embryos at developmental time

points when cells are known to be pluripotent, specified or committed to particular cell fates [29].

As a result, it is feasible to study how changes in RT accompany developmentally regulated

changes during cellular differentiation in vertebrates. We previously mapped RT at various

stages of early wild-type zebrafish development [30]. We revealed that a weak RT pattern

present at the onset of zygotic transcription becomes progressively more defined as the fish

develop. In addition, during gastrulation, large genomic domains switch from early-to-late

replication in the embryo. Finally, we observed that changes in epigenetic modifications involved

in enhancer activity correlate with developmental changes in RT [30].

To reveal the function of Rif1 in gene expression and RT during vertebrate development,

we have generated Rif1 mutant zebrafish, and we have performed RNA sequencing and whole-

genome RT analyses before the first major wave of zygotic transcription and before and after

gastrulation. We found that Rif1 is not essential for embryonic development, but its loss impairs

female sex determination. Although the Rif1 mutants had altered RT at each stage, the effects

of Rif1 loss on RT were minor compared to changes that naturally occur during development.

DNA replication timing was most affected by Rif1 loss after gastrulation, while transcription was

most affected early in development. Rif1 was required for normal expression of genes during

the first wave of zygotic genome activation. Our results indicate that in zebrafish, Rif1 has

distinct roles in DNA replication and transcription control that manifest at different stages of

development.

RESULTS

Generation of a Rif1 loss zebrafish model

The zebrafish genome has a single copy of the *rif1* gene on chromosome 9 (GRCz11

22,780,901-22,802,172). Similar to other vertebrates, the *zrif1* gene encodes a 2,347 amino

acid protein with heat repeats and PP1 phosphatase interaction sequences (Fig. 1B)[31]. To

determine the effects of Rif1 loss-of-function in the zebrafish, we mutated the gene using a

transcription activator-like effector nuclease targeting exon 8 (TALEN; Fig. 1A). Frameshift

mutations in exon eight would likely induce nonsense-mediated decay (NMD) of the mutant

mRNA. If expressed, the mutant allele would produce a protein ending within the sequence

encoding the heat repeats (Fig. 1A). We selected a stable line (*rif1*^{omf201}) with a seven base-pair

frameshift mutation for further analysis (Fig. 1A).

Fig. 1. A TALEN-induced mutation abrogates *rif1* expression in zebrafish. (A) Schematic illustration of the TALEN mutation in exon 8 of the zebrafish *rif1* gene. The DNA and protein sequences at the TALEN site are shown for the wild-type and a 7 bp deletion mutant. The deletion causes a frame shift and introduces a premature termination codon. The affected amino acids are in red. (B) Diagram of the *rif1* protein. The 7 bp frame shift mutation occurs in the conserved *rif1* N-terminal domain (Rif1-N), causing truncation of the Heat repeats and loss of the nuclear localization signal (NLS) and the PP1 phosphatase SILK and RVSF interaction motifs of the C-terminal domain (CTD). (C) Expression levels of *rif1* measured by quantitative RT-PCR analysis at pre-MBT (2.5 hours), dome (4.3 hours), shield (6 hours) and bud (10 hours) in wild-type (+/+), heterozygous (+/-) and homozygous (-/-) *rif1* zebrafish. Error bars represent standard deviation of the mean. (D) Rif1 and Smc3 protein expression levels were analyzed by immunoblotting at dome and shield in wild-type (+/+) and homozygous (-/-) *rif1* zebrafish.

We first tested whether zygotic *rif1^omf201^* mutants developed normally and survived to

sexual maturity. Most offspring from *rif1^omf201/+^* crosses hatched and began free feeding by five

days post-fertilization (dpf; data not shown). Therefore, we generated clutches of *rif1^+/+^*,

*rif1^omf201/+^*, or *rif1^omf201/omf201^* embryos spawning F2 mutant and wild-type fish. We then collected

embryos of each genotype at 2.5, 4.3, 6, and 10 hours post-fertilization (hpf) and measured *rif1*

mRNA by quantitative RT-PCR (Fig. 1C). These developmental time points span the start of

zygotic transcription through gastrulation[32]. Our data revealed that *rif1* expression in wild-type

zebrafish embryos is developmentally regulated, with the highest expression level at 2.5 hpf and

then a precipitous drop at all stages after that (Fig. 1C). The significance of this change in

mRNA expression for development is unknown, but we also found that Rif1 protein also

decreases in the embryo over these stages (data not shown). The expression level of *rif1*

mRNA in heterozygous and homozygous mutant fish was lower than wild-types at all stages

(Fig. 1C). In maternal-zygotic homozygous mutants, *rif1* mRNA was nearly undetectable (Fig.

1C). Therefore, we conclude the mutation causes a substantial decrease in *rif1* mRNA.

To determine if Rif1 protein levels were also affected by the mutation, we generated a

polyclonal antibody against the N-terminus of the protein. The antibody's antigen included the

portion of the truncated protein that the mutant allele would express (Fig. 1A, B). We did not

detect any residual full-length Rif1 protein on immunoblots of nuclear lysates from maternal-

zygotic homozygous mutants (Fig. 1D). Furthermore, we have not found evidence on the

immunoblots that the *rif1^omf201^* mutants express truncated Rif1 protein (Fig. 1D). Altogether,

these data suggest that the *rif1^omf201^* mutation eliminates Rif1 protein expression in the embryo.

Rif1 is not essential in zebrafish but its loss affects early development and sex

determination

Given that Rif1 loss in mice caused incompletely penetrant embryonic lethality, we

decided to track the development and survival of a large cohort of *rif1^omf201^* mutants [28]. We

Fig. 2. Loss of *rif1* affects zebrafish sex determination, epiboly, and primitive erythropoiesis. (A) Schematic of the heterozygous *rif1* mutant crosses used to test the essentiality of *rif1.* 766 fish were raised to sexual maturity (3 months) and then genotyped. (B) Quantitation of the genotypes from the heterozygous crosses shows the expected Mendelian ratios. (C) Counts of males and females from the heterozygous crosses shows fewer than expected *rif1* mutant female fish. (D) Time-lapse imaging of wild-type and maternal-zygotic *rif1* -/- early development shows a delay in epiboly completion in the mutants. (E) Expression of *gata1* and *B-globin* at 24 hpf marks primitive erythroid cells. In situ hybridization shows that *gata1* and *B-globin* expression at 24 hpf was decreased in rif1 homozygous mutants, indicating loss of primitive erythroid cells.

raised 766 offspring from *rif1$^{omf201/+}$* incrosses and counted fish of each genotype when they

reached sexual maturity (Fig. 2A, B). Zebrafish sex is determined by poorly defined genetic and

environmental factors, so we also counted fish of each gender [33]. Although all genotypes were

at their expected mendelian proportions, the number of female heterozygous and homozygous

mutants was reduced (Fig. 2B, C). We conclude that although the *rif1^{omf201}* mutation is not

zygotic lethal, it does perturb sex determination.

Our qRT-PCR analysis suggested that zebrafish have substantial *rif1* mRNA at pre-MBT

(midblastula transition), before full zygotic genome activation (Fig. 1A). Maternally supplied *rif1*

mRNA or protein might mask critical functions during early development in zygotic mutants.

Therefore, we carefully analyzed the maternal-zygotic *rif1$^{omf201/omf201}$* mutants, which lack

maternally supplied *rif1*, for any overt morphological defects during development. Mutant fish

development was delayed by approximately 1 hour at the end of gastrulation (10 hpf; Fig. 2D).

Moreover, *rif1^{omf201/omf201}* mutants had fewer circulating blood cells relative to wild types at one

dpf. Consistent with the blood cell deficiency, *rif1^{omf201/omf201}* mutants expressed lower levels of

hematopoiesis markers (*β-globin* and *gata-1*; Fig. 2E). Despite the delay in early development

and the deficiency in primitive hematopoiesis, maternal-zygotic *rif1^{omf201/omf201}* mutants recover

and develop to adulthood.

Rif1 loss impairs the establishment of the RT program genome-wide during zebrafish

development

Given that late replication and S-phase lengthening requires Rif1 in Drosophila, we

hypothesized that RT program establishment and change in vertebrates would also depend on

Fig. 3. Rif1 loss reduces the general sharpening of RT patterns that occur during normal zebrafish development. (A) Heat plot and dendograms showing hierarchical clustering of samples based on Euclidean distances calculated with RT values. (B) Principle component analysis reveals that 50% and 27% of variance in RT between samples is due to developmental stage (PC1) and genotype (PC2), respectively. (C) Autocorrelation analysis of RT values shows that RT profiles gradually gain structure during zebrafish development, and Rif1 loss delays that process. (D) Whole chromosome 1 RT profiles are shown for preMBT (top left), shield (top right), bud (bottom left) and 24 hpf (bottom right) stages. The RT profiles gradually develop definition, but Rif1 loss reduces the amplitude of the peaks and valleys. (E) Density plots of genome-wide RT values show a reduced dispersion of values from 0 in the *rif1* mutants compared to wild-types. Vertical black lines within density plots represent 25th and 75th percentiles.

Rif1 [23]. We tested this hypothesis by profiling RT in wild-type and $rif1^{omf201/omf201}$ mutant

zebrafish embryos at multiple stages. We focused on early development because we had

previously described changes in RT at multiple timepoints from the 256-cell through gastrulation

in zebrafish [30]. We showed that the 256-cell and shield stages in zebrafish are when the RT

pattern shifts from being relatively weak to defined [30]. Furthermore, the major wave of zygotic

genome activation occurs around the 256-cell stage. Therefore, 256-to-shield is analogous to

the stages in Drosophila development when Rif1 promotes the onset of late replication of

satellite repeats [23]. We also previously observed that the shield and bud stages are when

numerous broad RT changes occur [30]. Therefore, we used whole-genome sequencing to

generate RT maps for wild-type and Rif1 mutant zebrafish embryos at 128-cell/pre-MBT (2.25

hours), shield (6 hours), bud (10 hours), and 24 hpf.

We first assessed overall differences in RT between all genotypes and stages with

hierarchical clustering on Euclidean distances between samples (Fig. 3A). Using this approach,

we found that samples clustered first by developmental stage and then genotype, demonstrating

that the RT changes that occur during zebrafish development are more significant than those

caused by Rif1 loss. The Euclidean distances between wild-type and $rif1^{omf201/omf201}$ samples

were minor at the pre-MBT stage and increased progressively through development, reaching

their highest levels at 24 hpf. Importantly, biological replicates of the same developmental stage

and genotype were most similar, demonstrating the reproducibility of our RT measurements.

The conclusions drawn from the PCA were the same as Euclidean distance analysis

(Fig. 3B). All biological replicates clustered together on the PC2 vs. PC1 plot, showing that the

data is reproducible. Principal components 1 and 2 described RT differences caused by

developmental stage and genotype, respectively. The first principal component, which explained

50% of the RT variance, separated samples by developmental stage; the second component,

which explained 27% of the variance, separated samples by genotype. Therefore, the

developmental stage affected RT more strongly than the genotype. There was no difference

between wild-type and *rif1^omf201/omf201* samples in the first two principal components at the pre-

MBT stage (Fig. 3B; blue points), supporting that Rif1-loss had little effect on RT early in

development. As development proceeded, wild-type and *rif1^omf201/omf201* mutant samples

progressively separated on the second principal component, supporting that RT changes

caused by Rif1 loss were more pronounced later in development.

We previously showed that the RT pattern in zebrafish embryos gradually gained the

hallmark peak-and-valley RT profile structure as development proceeded [30]. We quantitatively

assessed the structure of the timing program in wild-type and *rif1^omf201/omf201* embryos throughout

development by calculating the autocorrelation of timing values along the lengths of the

chromosomes at each time point (Fig. 3C). As we described previously, RT value

autocorrelation increased between each stage in wild-type embryos [30]. The autocorrelation of

rif1^omf201/omf201 RT values also increased during development, but the change lagged behind wild-

type. For example, the autocorrelation of *rif1^omf201/omf201* RT values at the bud stage was like that

of wild-type at shield. The autocorrelation analysis indicated that Rif1 loss delayed the gradual

gain in RT profile structure during development.

Broad differences in RT were visually evident from plots of normalized (log2 S/G1) RT

values at all stages after the MBT (Fig. 3D). At the 128-cell stage (before the MBT), both

rif1^omf201/omf201 and wild-type RT profiles were flat compared to those at later stages (Fig. 3D).

This was consistent with replication origin firing synchrony at the 128-cell stage in both the wild-

type and *rif1* mutants. After the 128-cell stage, the RT profiles of *rif1^omf201/omf201* and wild-type

embryos developed peak-and-valley structures, but the *rif1^omf201/omf201* profiles were less defined

(Fig. 3D; compare black lines to colored lines). At each stage after 128-cell, the amplitudes of

peaks and valleys appeared lower in the *rif1^omf201/omf201* profiles (Fig. 3D). Overall, the RT values

(log2 S/G1) in *rif1^omf201/omf201* mutants at shield, bud, or 24 hpf were less dispersed from the

median than the RT values of wild-type embryos at the same stage (Fig. 3E). Altogether our

data show that Rif1 sharpens the genome-wide RT profile during zebrafish development.

Fig. 4. Rif1 loss does not prevent major changes in RT that occur during early zebrafish development. (A) Scatterplot of changes in RT of 2,498 putative enhancers that are H3K27 acetylated during gastrulation (DARs). A positive shield-to-24 hpf change indicates a switch to earlier replication during gastrulation or segmentation. The majority of acetylated regions switch to earlier replication in both wild-type and *rif1* mutants. (B) RT plotted across three regions with representative DARs. The positions of individual DARs are marked with triangles. (C-F) RT calculated for all of chr4:28,400,000-70,500,000 (C-D) or chr22:5,700,000-10,100,000 (E-F) at each stage and genotype. The broad early-to-late switches in RT during gastrulation occur in the *rif1* mutants.

Rif1 is not required for major changes in RT that occur during zebrafish development

We previously identified two significant changes in RT that occur during early zebrafish

development [30]. First, genomic domains spanning hundreds of kilobases underwent late-to-

early changes in RT. Both coincided with H3K27 acetylation or DNA demethylation at putative

enhancers within the domains. Second, ~40 million base-pairs of zebrafish chromosome 4q and

a large chromosomal domain on chromosome 22 (5,749,848-10,129,684) both underwent early-

to-late RT switches during gastrulation. The mechanisms underlying those changes are

unknown.

To show whether RT changes that coincide with enhancer activation require Rif1, we

calculated RT values at putative enhancers that become acetylated during early zebrafish

development (Fig. 3A-B)[34]. The majority (1,768/2,498) of differentially acetylated regions

(DARs) that become acetylated between 4.3–24 hpf changed to earlier replicating between the

shield (6 hpf) and 24 hpf stages in both wild-type and *rif1*[omf201/omf201] mutant embryos. Moreover,

the shield-to-24 hpf changes in RT for wild-type and *rif1*[omf201/omf201] DARs were strongly

correlated (rs=0.76, p<2.2e-16; Fig. 3A). Therefore, we conclude that Rif1-loss does not

generally affect the changes in RT that coincide with enhancer activation during early zebrafish

development.

To determine whether the early-to-late switches in RT that occur on chromosomes 4 and

22 require Rif1, we calculated RT values for Chr4q or Chr22:5,749,848-10,129,684 for both

wild-type and *rif1*[omf201/omf201] embryos at 128-cell, shield, bud, and 24 hpf stages (Fig. 4C-F).

Unexpectedly, Chr4q was replicated earlier during S-phase in the *rif1*[omf201/omf201] embryos than in

wild-type embryos at the 128-cell stage, but that difference was minor (Fig. 4C, D). As we

previously described, early-to-late RT switches on Chr4q (ANOVA p=7.27e-05) and Chr22

(ANOVA p=7.27e-05) occurred between the shield and 24 hpf stages in wild-type embryos (Fig.

4C-E). The early-to-late switches at Chr4q (ANOVA p=0.803) and Chr22 (ANOVA p=0.803) also

occurred in the *rif1*[omf201/omf201] mutant embryos (Fig. 4C-E). Although both Chr4q and

Chr22:5,749,848-10,129,684 underwent early-to-late switches in the *rif1*[omf201/omf201] mutant

embryos, both were earlier replicating in the mutant compared to wild-types at 24 hpf (Fig. 4C-

F). These differences might indicate that those specific timing switches were delayed or

incomplete, or it could be due to the genome-wide flattening of the RT profiles. Indeed, when we

compared all wild-type and *rif1*[omf201/omf201] mutant RT values at 24 hpf, we observed that most

late-replicating regions were earlier replicating in *rif1*[omf201/omf201] mutants (Fig. 3D, E). These data

have led us to conclude that although Rif1 promotes the establishment of the RT profile

genome-wide, it does not generally affect specific RT switches during zebrafish development.

237   Rif1 affects transcription at the earliest stages of zebrafish development

238   Mouse studies have revealed roles for Rif1 in regulating gene expression and epigenetic

239   marks in embryonic stem cells and cancer cell lines [21, 35, 36]. It has been suggested that Rif1

Fig. 5. Rif1 loss affects gene expression preferentially during the earliest stages of zebrafish development. (A) Matrix with colors and dendrograms representing Euclidean distances between 3' mRNA-Seq samples. The order of samples is based on hierarchical clustering and is the same for rows and columns. Each sample's genotype and developmental stage is marked by a color to the left of its row. (B) Principal component analysis (PCA) plot showing clusters of samples based on similarities of 3' mRNA-Seq data. The first two principal components are plotted, which represent 29.1% and 14.8% of the variance. Both the Euclidean distance and PCA analyses show that biological replicates are most similar, followed by samples from the same genotype and then stage. Rif1 mutant and wild-type gene expression differs most at dome and shield stages. (C) Fold-change in expression between *rif1* -/- and wild-type embryos versus normalized mean 3' mRNA-Seq read counts. Black data points represent differentially expressed genes (DEGs) with statistically significant (adjusted Wald test p-value < 0.05; Log2FC > 0.5) changes in expression in the rif1 mutants. (D) Counts of DEGs from C with lower (Down) or higher (Up) expression in *rif1* mutants.

affects transcription indirectly, possibly through its RT function [36]. We were particularly

interested in whether Rif1 loss would cause transcriptional changes later in development when

the RT program is established or in the early embryo, when there is a minimal RT pattern.

Therefore, we used 3' end RNA sequencing on maternal-zygotic *rif1*^{omf201/omf201} mutant and wild-

type zebrafish embryos at 1-cell (0.2 hours), 256-cell (2.5 hours), dome (4.3 hours), shield (6

hours), bud (10 hours) and day 1 (28 hours) (Fig. 5).

Fig. S5. Most DEGs are at normal levels in maternally-contributed mRNA. Heat plots showing the Log2 fold change in mRNA expression of *rif1* mutants vs wild types for all Down-DEGs (A) or Up-DEGs (B). Genes are displayed in rows and developmental stages are in columns. Rows were split by k-means clustering. Colors in the row annotations to the right of each heat plot denote the stage(s) at which each DEG was identified. The majority of 1-cell DEGs are expressed normally at other stages, and the majority of DEGs at other stages are expressed normally at 1-cell.

We determined the overall effects of Rif1 loss on the embryonic transcriptome by

analyzing the RNAseq data using principle component analysis (PCA) and hierarchical

clustering based on Euclidean distances (Fig. 5A, B). We found that biological replicates were

highly similar, and as expected, the transcriptome changed significantly between developmental

stages. Rif1 loss impacted the transcriptome at every stage evaluated but not all stages were

equally affected, with the most significant effects on gene expression at the dome and shield

stages (Fig. 5B). From this point forward, we will refer to differentially expressed genes (DEGs)

with higher or lower expression in the maternal-zygotic $rif1^{omf201/omf201}$ mutants as Up-DEGs or

Down-DEGs, respectively (adjusted Wald test p value < 0.05 and Log2-fold change > 0.5 or < -

0.5). The number of Up-DEGs peaked at the dome stage, while the number of Down-DEGs

peaked at the shield stage (Fig. 5C-D).

Zebrafish contribute most zygotic transcripts maternally, so the differences in transcript

abundance in the $rif1^{omf201/omf201}$ mutants could be due to changes in the supply or stability of

maternal mRNA (Harvey, 2013). We used k-means clustering to categorize all DEGs based on

their log-fold change in expression across all stages (Fig. S5A, B). The majority of one-cell

DEGs are expressed normally at other stages, and the majority of DEGs at other stages are

expressed normally at one-cell (Fig. S5A, B). Therefore, the differential expression of mRNAs at

later stages in the $rif1^{omf201/omf201}$ mutants is not due to differential maternal contribution.

Rif1 broadly suppresses transcription during Zygotic Genome Activation

To learn more about the genes affected by Rif1 loss, we looked at when zebrafish

embryos normally express the DEGs. White et al. profiled the zebrafish transcriptome at 18

developmental timepoints [1]. We used this dataset to categorize dome DEGs based on when

their peak expression occurs in wild-type embryos (Fig. 6A-D). We focused on dome DEGs

because they were the most abundant. Dome Up-DEGs are enriched in the Blastula/Gastrula

expression cluster (Fig. 6A, C). Since mRNAs that peak at the Blastula and Gastrula stages are

likely to be transcribed during the major wave of zygotic genome activation at the MBT, our data

suggests Rif1 loss stimulates the transcription of mRNAs at the first major wave of Zygotic

Genome Activation (ZGA). Dome Down-DEGs are enriched in the pre-ZGA cluster, which

suggests that Rif1 loss also enhances the destruction of maternal mRNAs (Fig. 6B, D). The

degradation of many maternal mRNAs in zebrafish is triggered by transcription of the *mir-430*

microRNA cluster before the major wave of zygotic transcription [37].

Using complementary approaches, Lee et al. and Heyn et al. defined "first wave" genes

that are transcribed immediately upon zygotic genome activation [2, 3]. We used gene set

enrichment analysis to determine the extent to which the $rif1^{omf201/omf201}$ mutants overexpress

first-wave genes (Fig. 6G). At the dome stage, 272 out of 383 first wave genes had a Log2FC > 0 in mutants compared to wild-types. Furthermore, the first wave gene sets were located towards the top of the list of all genes ranked by the dome stage Log2FC (Fig. 4G). Our data indicate that Rif1 loss generally increases the transcription of genes at the ZGA, but by including genes with substantial maternal contribution in our analysis, we might have underestimated that effect. Therefore, we used additional gene sets identified by Harvey et al. as being strictly zygotic or maternal (Fig. 6H)[4]. Like the "first wave" genes, the majority of genes in the zygotic set were expressed

Fig. 6. ZGA transcribed or degraded mRNAs are deregulated in *rif1* mutants. (A-B) Heat plots showing the expression (tags per million scaled to maximum) of *rif1* mutant Dome Up (A) or Down (B) DEGs (rows) across 18 developmental timepoints (columns) in wild type embryos. Expression data is from White, et al. [1]. The legend across the top of each heat plot denotes six developmental phases. (C-D) Log2 odds ratios from Fisher's exact tests of associations between dome Up (C) or Down (D) DEGs and the developmental phases in A and B. The total number of each DEG expressed in each phase is indicated by the size of the point, and point fills indicate statistical significance. Dome Up or Down DEGs are overrepresented in the Blastula/Gastrula or pre-ZGA phases, respectively. (E-F) Scaled QuantSeq counts for dome Up (E) or Down (F) DEGs across all developmental stages. Overall, dome Up or Down DEGs are normally up- or down-regulated by shield stage, so normal expression of dome DEGs is generally restored by shield. (G) Gene set enrichment analysis showing that genes known to be expressed during the first waves of zebrafish transcription are overexpressed upon Rif1 loss. The genes are ranked by Log2FC in expression between dome stage wild type and $rif1^{omf201/omf201}$ embryos, as measured by QuantSeq. (H) Gene set enrichment analysis showing that genes, which are strictly zygotic or maternal, are often up- or down-regulated at dome stage in *rif1* mutants, respectively.

Fig. 7. Labeling of nascent mRNAs reveals that the role for Rif1 in regulating ZGA transcription is pervasive. (A, D) Volcano plots showing the Log2-fold change in T-to-C conversions from SLAM-seq for each transcript between wild type and *rif1* mutant embryos at preMBT (A) or dome (D) stage. P values were calculated using the beta-binomial test and were corrected using the Benjamini & Hochberg method. The colors represent the density of overlapping data points. (B, E). Barplots of the number of differentially labeled genes (DLGs; >1.5-fold change in T-to-C conversions and BH-adjusted p value <= 0.1) in A or D. (C, F) Boxplots of T-to-C conversion percentages at preMBT (C) or dome (F) stage for the following: all genes; DEGs upregulated in *rif1* mutants from this study (256 Cell Up, Dome Up, or Shield Up), genes transcribed during the first ZGA wave from Heyn et al. or Lee et al., or genes identified by Harvey et al. as maternal or zygotic [2-4].

at higher levels (72/98 Log2FC > 0) in mutant embryos at the dome stage compared to wild-type, and most of the zygotic genes were towards the top of the list of genes ranked by dome Log2FC (Fig. 6H top). In contrast, genes in the maternal set were often down in the mutants at dome and were found towards the bottom of the ranked gene list (Fig. 6H bottom). Altogether our data suggest that Rif1 loss enhances the destruction of maternal

mRNAs and stimulates the expression of genes during zygotic genome activation.

To directly assess transcription during ZGA, we labeled nascent mRNAs in early

embryos with 4-Thiouridine (S4U)(Herzog et al. 2017). By injecting the label into one-cell stage

embryos we were able to identify mRNAs synthesized by the pre-MBT (2.5 hours) or dome (4.3

hours) stages. The incorporated label was mapped to mRNAs using thiol(SH)-linked alkylation

for metabolic sequencing of RNA (SLAM-seq). SLAM-seq introduces T-to-C conversions at the

sites of the incorporated S4U. Thus, each gene's percentage of T-to-C conversions indicates

the amount of its mRNA synthesized during the labeling period (zygotic) relative to pre-existing

mRNA (maternal).

At the pre-MBT stage, there was very little difference in T-to-C conversion percentages

between wild-type and *rif1^omf201/omf201* mutant embryos, and there were only 102 differentially

labeled genes (DLGs) at preMBT (Fig. 7A-C; fold-change in T-to-C conversion > 1.5 and beta-

binomial test FDR > 0.1). Therefore, we conclude that zygotic genome activation is not

occurring earlier than the MBT for most genes in the *rif1^omf201/omf201* mutants. In contrast, there

were 1,872 DLGs with increased labeling in mutants at dome (Fig. 7D, E; fold-change in T-to-C

conversion > 1.5 and beta-binomial test FDR > 0.1). Furthermore, the T-to-C conversion rates at

the dome stage were higher in the *rif1^omf201/omf201* mutants for all gene categories analyzed (Fig.

7F). The differences in T-to-C conversion were greatest for the genes we previously identified

as being overexpressed in the mutants (256-Cell and dome Up-DEGs) and for the genes known

to be transcribed during ZGA (First Wave and Zygotic). Notably, we detected a significant

difference in mean T-to-C conversion calculated for all labeled genes. Overall, these data

demonstrate that Rif1 has pervasive effects on transcription during zygotic genome activation.

DISCUSSION

Previous studies of Rif1 in vertebrates have mainly been limited to cell culture models

and have not addressed how its loss impacts RT establishment and change in the early embryo.

We have developed a zebrafish model to study the developmental effects of Rif1 loss. Using

this model, we have determined the impacts of Rif1 knockout on zebrafish embryonic

development and survival, RT, and gene expression.

Our work adds to our understanding of the importance of Rif1 for vertebrate

development. The extent to which Rif1 is essential for embryogenesis has been unclear. In

Drosophila melanogaster, Rif1 knockdown by RNA interference caused embryonic lethality, but

its knockout by CRISPR-Cas9 caused only mild effects on hatching rates[19, 23, 38]. Rif1

knockout phenotypes in mice depend on genetic background and sex. Rif1 is essential in

female mice, which die between the blastula and early gastrula stages[27]. Male mice are less

affected by Rif1 loss. Although most Rif1 mutant male mice die around mid-gestation, a fraction

survives to sexual maturity[25-28]. We observed only mild *rif1* phenotypes during embryonic

development. Rif1 loss only slightly delayed epiboly completion. The reason for that delay is

unknown, but it may be due to the transcriptional effects of Rif1 loss at the ZGA. The *rif1*

mutants also showed a deficiency in primitive hematopoiesis. The first wave of hematopoiesis is

not essential for zebrafish development, however it is in mice. Thus, defects in hematopoiesis

may contribute to the poor survival of Rif1 mutant mice.

The most evident phenotype in the *rif1* mutant zebrafish is the substantial reduction in

the number of females. It is tempting to speculate that this phenotype is related to female-

specific lethality in mice, but we suspect the underlying causes differ. In part, Rif1 mutant female

mice die due to an X chromosome inactivation defect, but zebrafish lack sex chromosomes [27].

Furthermore, since we showed that expected numbers of *rif1* mutant fish survive to sexual

maturity, the skew in gender ratio must be due to the preferential development of male fish

rather than female lethality. A reduction in the proliferation or survival of primordial germ cells

(PGCs) causes transdifferentiation of zebrafish to males [39]. To understand how Rif1 promotes

female development, we must determine whether its loss affects PGC proliferation. Intriguingly,

mouse PGCs exhibit increased replication-transcription conflicts and are sensitive to mutations

in Fanconi anemia pathway genes required for replication stress survival[40, 41]. Zebrafish

PGCs exhibit high levels of intergenic transcription, which suggests that they may also have

increased replication-transcription conflicts[42]. Furthermore, mutations in the Fanconi anemia

DNA repair pathway disrupt female zebrafish development[43, 44]. Therefore, the female

development defect in the Rif1 mutant zebrafish could be due to deregulated replication fork

initiation or impaired replication fork stability in PGCs.

Studies with flies and mammalian cells have shown situations where Rif1 becomes less

critical for RT once constitutive heterochromatin forms. For example, *D. melanogaster* Rif1

delays the replication of specific satellite repeats before H3K9me3 and HP1a associate with

them, but once constitutive heterochromatin forms, Rif1 is unnecessary [23, 45]. In human

HCT116 cells, late replication of genomic domains with high levels of the heterochromatin mark

H3K9me3 does not depend on Rif1 [36]. In contrast, Rif1 knockout nearly completely ablates

the RT pattern in human embryonic stem cells, which have relatively little constitutive

heterochromatin[36, 46]. Pre-gastrula zebrafish embryos express high levels of Rif1 and

substantially lack constitutive heterochromatin [47]. Thus, we were surprised that Rif1 loss had

relatively little effect on RT in early embryos. Instead, we found that RT differences were more

pronounced later in development. Understanding this paradox will likely reveal mechanisms that

control Rif1 activity.

Although the effects of Rif1 loss on RT increased as zebrafish development proceeded,

the significant developmental changes that shaped the RT profile of late-stage embryos were

not affected. For example, *rif1* was not required for the early-to-late timing switches that occur

over large segments of chromosomes 4 and 22 during gastrulation, nor was it required for the

widespread late-to-early switches that coincided with enhancer activation. Much of the variance

in RT between wild-type and mutant embryos was due to a general flattening of the timing

profiles, and Rif1 loss did not disrupt the overall peak and valley profile structure. Therefore, our

data suggest that instead of causing developmental RT switches in zebrafish, Rif1 sharpens RT

caused by other unknown determinants.

If Rif1 is not required, what drives developmental changes in RT? Zebrafish embryos

establish constitutive heterochromatin by the start of gastrulation, so the early-to-late RT

switches that occur during and after that stage might be due to further maturation of

heterochromatin[47]. The activation of enhancers might drive late-to-early RT changes

throughout the zebrafish genome. Early replicating control elements (ERCEs), which have

active epigenetic marks associated with enhancers, cause early replication of large domains in

the mouse genome [48]. More work is needed to determine whether the formation of constitutive

heterochromatin and activation of ERCEs drive developmental changes in RT in zebrafish.

We were particularly interested in whether RT changes in the embryo would lead to

transcriptional changes. A long-standing hypothesis is that the RT of a gene can affect its

transcriptional competence[49-51]. Recently, Klein et al. used human cell lines in which Rif1

could be conditionally depleted to show that its loss caused widespread and coordinated RT

and epigenomic changes[36]. Furthermore, the epigenomic alterations caused by Rif1 loss were

often associated with gene expression changes[36]. Although this work strongly supports the

hypothesis that RT maintains the epigenome, the effects of Rif1 loss in a developing embryo

where the RT program and the epigenome are established have not been examined. We used

whole-genome RT and RNAseq analysis to determine whether Rif1 loss caused RT changes

that led to dysregulated gene expression. Unexpectedly, we found that when Rif1 had the most

significant effect on RT during later development, there were few changes in mRNA levels. One

limitation of the current work was that we did not directly measure alterations in the epigenome,

which may have occurred without transcriptional changes. A second limitation was that we

measured RT and gene expression for all cells in the embryo in bulk. Consequently, we were

unable to detect cell-to-cell variability in either RT or gene expression. Indeed, RT dysregulation

could cause epigenetic variability. Nonetheless, our data show that RT changes caused by Rif1

loss do not substantially change overall gene expression later in development.

The most pronounced effects of Rif1 loss on transcription were during the earliest stages

of development. We previously showed that RT patterns are detected around the MBT in

zebrafish embryos, but those patterns are relatively weak compared to later development. In

this work, we showed that there was almost no detectable RT pattern in wild-type or Rif1

mutants before the ZGA. Despite the lack of RT patterns, Rif1 loss affected the first major wave

of transcription. Most genes affected by Rif1 loss at that stage increased slightly, but a

significant fraction of all genes transcribed at the ZGA were upregulated. Thus, Rif1 loss had a

strong overall effect on transcription at the ZGA.

Notably, the role for Rif1 in regulating ZGA transcription is evolutionarily conserved. As

we show here for zebrafish, mouse pluripotent cells express Rif1 at high levels, and Rif1

expression decreases during cellular differentiation[52]. Multiple groups have reported that Rif1

loss in mouse embryonic stem cells (mESCs) causes increased expression of two-cell (2C)

zygote-specific genes [35, 53-56]. Given that ZGA occurs through different transcription factors

in zebrafish and mice, it is likely that Rif1 affects genes expressed at the ZGA in both species by

regulating a common fundamental process rather than the activities of specific transcription

factors. mESCs in the 2C state have increased histone marks associated with active

transcription, and Rif1 loss is sufficient to cause these chromatin changes [56, 57]. Mouse Rif1

co-immunoprecipitates with multiple repressive histone methyltransferases and with the non-

canonical polycomb repressive complex PRC1.6, and Rif1 promotes the recruitment of

chromatin regulators to several 2C genes in mESCs [57]. We do not know how Rif1 affects the

epigenome in the early zebrafish, and this is an important goal for future research. Given the

striking similarity in Rif1 function in early zebrafish and mouse embryos, we anticipate the role of

Rif1 in regulating transcription in the early embryo will be broadly conserved in vertebrates.

In conclusion, we have described multiple phenotypic consequences of Rif1 loss in

zebrafish. We found that Rif1 undergoes a functional switch during zebrafish development.

When it is highly abundant during early development, Rif1 regulates transcription but does not

affect RT. Later in development, Rif1 is important for RT but does not substantially affect gene

expression. The zebrafish will be useful for understanding how Rif1 switches between these

functional modes at different stages of development. Furthermore, we have identified a new

function for Rif1 in sex determination. Thus, the *rif1* mutant zebrafish described here will serve

as a valuable model for understanding PGC biology and sex determination.

MATERIALS AND METHODS

Animal care. All animals were handled in strict accordance with protocols approved by

the OMRF Institutional Animal Care and Use Committee. Adult breeding fish were housed in an

aquatic animal facility in tanks of ~30 fish per tank and maintained at 26.5°C with 10-hour light

and 14-hour dark cycles.

TALEN production and microinjection . The TALEN expression constructs targeting

the seventh exon of the *rif1* gene were assembled in pCS2TAL3-DD and pCS2TAL3-RR using

Golden Gate Assembly as in Cermak et al. [58]. pCS2TAL3-DD and pCS2TAL3-RR were gifts

from David Grunwald (Addgene plasmids # 37275 and 37276; RRID:Addgene_37275;

RRID:Addgene_37276). mRNA encoding the TALE nuclease subunits was synthesized using

the mMESSAGE mMACHINE SP6 Transcription Kit (AM1340; ThermoFisher Scientific,

Waltham, MA) and purified using RNeasy Mini Kit (74104; Qiagen). 125pg of each TALEN

mRNA was injected into 1-cell stage zebrafish embryos. TALEN-injected embryos were raised

to adulthood and assessed for transmission of insertions/deletions at the TALEN cut site using

high resolution melt analysis (HRMA). F1 fish from mutation transmitting F0 animals were raised

to adulthood and analyzed for mutation by HRMA, and the specific *rif1* mutation was identified

by Sanger sequencing of PCR amplicons spanning the mutation site. *rif1* mutant fish were

genotyped using HRMA, Sanger sequencing, and/or competitive allele-specific PCR (KASP

assay; LGC).

Zebrafish embryo collection and processing for gDNA, mRNA, or protein.

Zebrafish were spawned as described previously [59]. For QuantSeq, 2-cell embryos were

injected with 50 pmoles of 4-Thiouridine (S4U). Embryos were incubated at 28.5°C. Before

collection, clutches were visually inspected to ensure that all embryos were developing

synchronously. Unfertilized or morphologically abnormal embryos were removed before

collection. Whole embryo pools to be used for gDNA or mRNA were rapidly frozen and stored at

-80°C for later processing. Embryos were prepared for fluorescence activated cell sorting based

on DNA content as described in Siefert et al[59]. Briefly, bud stage or 24 hpf embryos were

dechorionated with Pronase, deyolked, and then triturated to a single cell suspension in

Phosphate Buffered Saline. Nuclei were prepared from the cells were incubated in DNA staining

solution (PBS; 1% Bovine Serum Albumin; 50 ug/ml Propidium iodide P4170-25mg Sigma-

Aldrich; 100 ug/ml RNAse A 12091039 Invitrogen) on ice for at least one hour before sorting.

The nuclei were then filtered through a 40μm mesh before analysis on a BD LSRII (shield and

bud), BD Biosciences FACSAria IIIu (28hpf) or BD Biosciences FACSCalibur. Genomic DNA

was prepared from whole embryo pellets or sorted nuclei using the Quick-DNA Miniprep Plus Kit

(D4069; Zymo Research). mRNA was prepared from whole embryos using the RNeasy Mini Kit

(74104; Qiagen). For protein lysates, whole embryos in pools were deyolked and disaggregated

into a single cell suspension as previously described [60], and dry cell pellets were stored at -

80°C for later use. To make crude nuclear extract, cell pellets were re-suspended in Buffer A

(10mM HEPES, 10mM KCl, 1.5mM MgCl$_2$, 0.34M sucrose, 10% glycerol) with 0.1% triton and

protease inhibitors followed by Buffer A plus inhibitors, 0.1% triton and 1uL/mL Benzonase

(E1014; Sigma).

***In situ* Hybridization.** The Thisse Lab protocol was followed for the ISH [61]. Briefly,

ISH probes were constructed by PCR from genomic DNA and isothermal assembly into HindIII-

BamHI digested pUC57-Amp plasmids flanked by T3 and T7 promoters for sense (coding RNA)

and anti-sense strand synthesis, respectively. Primers used to amplify probe sequences are as

follows: *β-globin* forward acc ctc act aaa ggg aaC GCG CAG CGA TTC AGA ACA T, *β-globin*

reverse acg act cac tat agg gcC TTC TGA GGG CTG ACA CAA CA, *gata1a* forward acc ctc act

aaa ggg aaA CTC CTC TGA GCC TTC TCG T, and *gata1a* reverse acg act cac tat agg gcT

GTG GAG AAG GGC CGT AAA C. Dioxygenin (DIG)-labelled RNA probes were synthesized

with T3 and T7 polymerase with reagents from Roche. Embryos were dechorionated with

forceps and fixed in 4% paraformaldehyde in 1X PBS overnight. Fixed embryos were

dehydrated in 100% methanol and stored at -20°C. The 24 hpf embryos were proteinase K

treated for 10 minutes. The protocol was programmed into an Intavis VSi machine to automate

the prehybridization and washing procedures. Embryos were removed from the machine and

incubated in DIG-labelled RNA probes overnight in a 70°C water bath. Embryos were then

returned to the machine for washing, pre-incubation with Bovine Serum Albumin, incubation with

anti-DIG antibody, and additional washing. Labelled embryos were imaged with a Nikon

SMZ1500 stereomicroscope with Andor Zyla sCMOS camera.

Antibodies used for immunoblotting. Anti-zebrafish Rif1 antibodies were made by

immunizing rabbits with bacterially expressed purified fragments of zebrafish Rif1

(NP_001074275.1 amino acids 1-150 or 2199-2347) and were affinity purified from whole serum

(Cocalico Biologicals, Inc.). Rabbit polyclonal anti-human Smc3 was a gift from Susannah

Rankin [62]. Peroxidase AffiniPure Goat Anti-Rabbit IgG (H+L) was purchased from Jackson

Immuno Research Laboratories (RRID: AB_2313567).

Quantitative RT-PCR. mRNA was converted to cDNA using the AccuScript High

Fidelity Reverse Transcriptase (600089; Agilent, Santa Clara, CA). qPCR was done on a

LightCycler 480 with the SYBR Green I Master (04707516001; Roche) master mix. Three

technical replicates were performed for each sample. The relative mRNA expression was

calculated using the $2^{-\Delta\Delta Cq}$ method. The following primer sequences were used: *rif1* forward

ACT GGC TGA TGA CAT TGA TCG T, reverse ATC CAG GGC TAC GGA GGT TA; *eef1a1l1*

forward CTG GAG GCC AGC TCA AAC AT, reverse ATC AAG AAG AGT AGT ACC GCT AGC

ATT AC.

Next generation library preparation and sequencing. For whole-genome

sequencing, 50ng - 2µg of gDNA was sheared to approximately 300 bp fragments using a

sonicator (Covaris). The gDNA libraries were prepared using the KAPA Hyper Prep Kit (Roche)

with SeqCap (Roche) or Illumina TruSeq (Illumina) adapters. Genomic DNA libraries were

sequenced in 150bp paired-end read fragments on an Illumina HiSeq 2500 or NovaSeq 6000.

Before library preparation, the SlamSeq mRNA was Iodoacetamide treated using the SLAMseq

Kinetics Kit – Anabolic Kinetics Module (Lexogen). The QuantSeq and SlamSeq libraries were

prepared using the QuantSeq 3' mRNA-Seq Library Prep Kit for Illumina (Lexogen). RNA

sequencing was performed using custom primers on an Illumina Nextseq 500 with High Output

chemistry and 75bp single-ended reads.

Replication timing analysis. RT data were generated based on copy-number

variations between filtered G1 and S-phase whole-genome sequencing read counts, as

described previously (Siefert, 2017). Pools of >100 whole embryos at preMBT, dome, or shield

stage were treated as S-phase samples, since a high percentage cells at these stages are in S

phase. Nuclei from bud or 24 hpf embryos were sorted based on DNA content to isolate G1 and

S-phase fractions. The data was processed using our RepliTimer pipeline

(https://github.com/SansamLab/RepliTimer/tree/v1.0.11; DOI 10.5281/zenodo.6426313.) [63].

Whole genome sequencing reads were trimmed with fastp, aligned to the zebrafish genome

(GRCz11) using BWA-MEM, and duplicate reads were marked with Picard MarkDuplicates.

Properly-paired and non-duplicated reads with MapQ scores of at least 20 were filtered with

Bamtools. Alignments were read into GenomicRanges for downstream analysis in R. Reads

were also removed if they were within a set of 100 bp windows with anomalously high read

counts (exceeding 1.5 times the interquartile range in at least 4 out of 65 whole genome

sequencing runs from this study and previously published work)[30]. Genomic windows for

calculating RT were defined as regions with a minimum of 400 reads in every sample. An S/G1

quotient was calculated for each window, and quotients were smoothed and Log2 transformed

[30].

RNA sequencing analysis (QuantSeq 3' mRNA-Seq and SLAM-Seq). The QuantSeq

data was analyzed using our QuantSeqPipeline pipeline (https://github.com/SansamLab-

Pipelines-Genomics/QuantSeqPipeline/releases/tag/v0.1.2; DOI 10.5281/zenodo.6622164).

Reads were trimmed with bbmap and aligned to GRCz11 with STAR. Using a custom R script,

probable internal priming sites at polyA genomic sequences were identified and reads mapping

to those sites were removed. Reads aligning to 3' ends of transcripts are then counted. The

4.3.2 transcript annotation by Lawson et al was used [64]. Differentially expressed genes were

identified using DESeq2, and plots were generated in R. The SLAM-Seq sequencing reads

were aligned and T-to-C conversion rates were calculated using the SLAM-DUNK package with

default settings [65].

ACKNOWLEDGEMENTS

This work was supported by National Institutes of Health National Institute of General Medical

Sciences grant 1R01GM121703 and by the Oklahoma Center for Adult Stem Cell Research.

COMPETING INTERESTS

No competing interests declared.

AUTHOR CONTRIBUTIONS

cls, eam, jcs, and tdn conceived and designed the experiments and analysis. jcs generated the

rif1 mutant. eam, tdn, and cgs performed the RT experiments. jcs performed the in situ

hybridizations. eam performed the RNASeq and SlamSeq experiments. eam performed the live

embryo imaging and qRT-PCR analysis. eam and dg performed the large cohort genotyping

experiments. eam, dg, cgs, and tdn raised breeding cohorts for the majority of experiments. cls,

eam, and tdn analyzed the RT, RNASeq, and SlamSeq data. cls, eam, and tdn wrote the paper.

REFERENCES

1.   White RJ, Collins JE, Sealy IM, Wali N, Dooley CM, Digby Z, et al. A high-resolution mRNA expression time course of embryonic development in zebrafish. Elife. 2017;6. Epub 20171116. doi: 10.7554/eLife.30860. PubMed PMID: 29144233; PubMed Central PMCID: PMCPMC5690287.

2.   Heyn P, Kircher M, Dahl A, Kelso J, Tomancak P, Kalinka AT, et al. The earliest transcribed zygotic genes are short, newly evolved, and different across species. Cell Rep. 2014;6(2):285-92. Epub 20140116. doi: 10.1016/j.celrep.2013.12.030. PubMed PMID: 24440719.

3.   Lee MT, Bonneau AR, Takacs CM, Bazzini AA, DiVito KR, Fleming ES, et al. Nanog, Pou5f1 and SoxB1 activate zygotic gene expression during the maternal-to-zygotic transition. Nature. 2013;503(7476):360-4. Epub 20130922. doi: 10.1038/nature12632. PubMed PMID: 24056933; PubMed Central PMCID: PMCPMC3925760.

4.   Harvey SA, Sealy I, Kettleborough R, Fenyes F, White R, Stemple D, et al. Identification of the zebrafish maternal and paternal transcriptomes. Development. 2013;140(13):2703-10. Epub 20130529. doi: 10.1242/dev.095091. PubMed PMID: 23720042; PubMed Central PMCID: PMCPMC3678340.

5.   Rhind N, Gilbert DM. DNA replication timing. Cold Spring Harb Perspect Biol. 2013;5(8):a010132. Epub 20130801. doi: 10.1101/cshperspect.a010132. PubMed PMID: 23838440; PubMed Central PMCID: PMCPMC3721284.

6.   Hiratani I, Ryba T, Itoh M, Rathjen J, Kulik M, Papp B, et al. Genome-wide dynamics of replication timing revealed by in vitro models of mouse embryogenesis. Genome Res. 2010;20(2):155-69. Epub 20091201. doi: 10.1101/gr.099796.109. PubMed PMID: 19952138; PubMed Central PMCID: PMCPMC2813472.

7.   Hiratani I, Ryba T, Itoh M, Yokochi T, Schwaiger M, Chang CW, et al. Global reorganization of replication domains during embryonic stem cell differentiation. PLoS Biol. 2008;6(10):e245. doi: 10.1371/journal.pbio.0060245. PubMed PMID: 18842067; PubMed Central PMCID: PMCPMC2561079.

8.   MacAlpine DM, Rodríguez HK, Bell SP. Coordination of replication and transcription along a Drosophila chromosome. Genes Dev. 2004;18(24):3094-105. doi: 10.1101/gad.1246404. PubMed PMID: 15601823; PubMed Central PMCID: PMCPMC535919.

9.   White EJ, Emanuelsson O, Scalzo D, Royce T, Kosak S, Oakeley EJ, et al. DNA replication-timing analysis of human chromosome 22 at high resolution and different developmental states. Proc Natl Acad Sci U S A. 2004;101(51):17771-6. Epub 20041210. doi: 10.1073/pnas.0408170101. PubMed PMID: 15591350; PubMed Central PMCID: PMCPMC539744.

10.   Schübeler D, Scalzo D, Kooperberg C, van Steensel B, Delrow J, Groudine M. Genome-wide DNA replication profile for Drosophila melanogaster: a link between transcription and replication timing. Nat Genet. 2002;32(3):438-42. Epub 20020930. doi: 10.1038/ng1005. PubMed PMID: 12355067.

11.   Dixon JR, Jung I, Selvaraj S, Shen Y, Antosiewicz-Bourget JE, Lee AY, et al. Chromatin architecture reorganization during stem cell differentiation. Nature. 2015;518(7539):331-6. doi: 10.1038/nature14222. PubMed PMID: 25693564; PubMed Central PMCID: PMCPMC4515363.

12.   Zhou J, Ermakova OV, Riblet R, Birshtein BK, Schildkraut CL. Replication and subnuclear location dynamics of the immunoglobulin heavy-chain locus in B-lineage cells. Mol Cell Biol. 2002;22(13):4876-89. doi: 10.1128/mcb.22.13.4876-4889.2002. PubMed PMID: 12052893; PubMed Central PMCID: PMCPMC133899.

13.   Cornacchia D, Dileep V, Quivy JP, Foti R, Tili F, Santarella-Mellwig R, et al. Mouse Rif1 is a key regulator of the replication-timing programme in mammalian cells. EMBO J.

2012;31(18):3678-90. doi: 10.1038/emboj.2012.214. PubMed PMID: 22850673; PubMed
Central PMCID: PMCPMC3442270.
14.   Hayano M, Kanoh Y, Matsumoto S, Renard-Guillet C, Shirahige K, Masai H. Rif1 is a
global regulator of timing of replication origin firing in fission yeast. Genes Dev. 2012;26(2):137-
50. Epub 2012/01/27. doi: 10.1101/gad.178491.111. PubMed PMID: 22279046; PubMed
Central PMCID: PMCPMC3273838.
15.   Hiraga S, Alvino GM, Chang F, Lian HY, Sridhar A, Kubota T, et al. Rif1 controls DNA
replication by directing Protein Phosphatase 1 to reverse Cdc7-mediated phosphorylation of the
MCM complex. Genes Dev. 2014;28(4):372-83. Epub 2014/02/18. doi:
10.1101/gad.231258.113. PubMed PMID: 24532715; PubMed Central PMCID:
PMCPMC3937515.
16.   Yamazaki S, Ishii A, Kanoh Y, Oda M, Nishito Y, Masai H. Rif1 regulates the replication
timing domains on the human genome. EMBO J. 2012;31(18):3667-77. doi:
10.1038/emboj.2012.180. PubMed PMID: 22850674; PubMed Central PMCID:
PMCPMC3442267.
17.   Davé A, Cooley C, Garg M, Bianchi A. Protein phosphatase 1 recruitment by Rif1
regulates DNA replication origin firing by counteracting DDK activity. Cell Rep. 2014;7(1):53-61.
Epub 2014/03/25. doi: 10.1016/j.celrep.2014.02.019. PubMed PMID: 24656819; PubMed
Central PMCID: PMCPMC3989773.
18.   Mattarocci S, Shyian M, Lemmens L, Damay P, Altintas DM, Shi T, et al. Rif1 controls
DNA replication timing in yeast through the PP1 phosphatase Glc7. Cell Rep. 2014;7(1):62-9.
Epub 2014/04/02. doi: 10.1016/j.celrep.2014.03.010. PubMed PMID: 24685139.
19.   Sreesankar E, Bharathi V, Mishra RK, Mishra K. Drosophila Rif1 is an essential gene
and controls late developmental events by direct interaction with PP1-87B. Sci Rep.
2015;5:10679. Epub 2015/05/30. doi: 10.1038/srep10679. PubMed PMID: 26022086; PubMed
Central PMCID: PMCPMC4448129.
20.   Sukackaite R, Cornacchia D, Jensen MR, Mas PJ, Blackledge M, Enervald E, et al.
Mouse Rif1 is a regulatory subunit of protein phosphatase 1 (PP1). Sci Rep. 2017;7(1):2119.
Epub 2017/05/20. doi: 10.1038/s41598-017-01910-1. PubMed PMID: 28522851; PubMed
Central PMCID: PMCPMC5437018.
21.   Foti R, Gnan S, Cornacchia D, Dileep V, Bulut-Karslioglu A, Diehl S, et al. Nuclear
Architecture Organized by Rif1 Underpins the Replication-Timing Program. Mol Cell.
2016;61(2):260-73. Epub 20151224. doi: 10.1016/j.molcel.2015.12.001. PubMed PMID:
26725008; PubMed Central PMCID: PMCPMC4724237.
22.   Hiraga SI, Ly T, Garzón J, Hořejší Z, Ohkubo YN, Endo A, et al. Human RIF1 and
protein phosphatase 1 stimulate DNA replication origin licensing but suppress origin activation.
EMBO Rep. 2017;18(3):403-19. Epub 2017/01/13. doi: 10.15252/embr.201641983. PubMed
PMID: 28077461; PubMed Central PMCID: PMCPMC5331243.
23.   Seller CA, O'Farrell PH. Rif1 prolongs the embryonic S phase at the Drosophila mid-
blastula transition. PLoS Biol. 2018;16(5):e2005687. Epub 2018/05/11. doi:
10.1371/journal.pbio.2005687. PubMed PMID: 29746464; PubMed Central PMCID:
PMCPMC5963817.
24.   Meléndez García R, Haccard O, Chesneau A, Narassimprakash H, Roger J, Perron M,
et al. A non-transcriptional function of Yap regulates the DNA replication program in Xenopus
laevis. eLife. 2022;11:e75741. doi: 10.7554/eLife.75741.
25.   Chapman JR, Barral P, Vannier JB, Borel V, Steger M, Tomas-Loba A, et al. RIF1 is
essential for 53BP1-dependent nonhomologous end joining and suppression of DNA double-
strand break resection. Mol Cell. 2013;49(5):858-71. Epub 2013/01/22. doi:
10.1016/j.molcel.2013.01.002. PubMed PMID: 23333305; PubMed Central PMCID:
PMCPMC3594748.

26.   Daxinger L, Harten SK, Oey H, Epp T, Isbel L, Huang E, et al. An ENU mutagenesis
screen identifies novel and known genes involved in epigenetic processes in the mouse.
Genome Biol. 2013;14(9):R96. Epub 2013/09/13. doi: 10.1186/gb-2013-14-9-r96. PubMed
PMID: 24025402; PubMed Central PMCID: PMCPMC4053835.
27.   Enervald E, Powell LM, Boteva L, Foti R, Blanes Ruiz N, Kibar G, et al. RIF1 and KAP1
differentially regulate the choice of inactive versus active X chromosomes. Embo j.
2021;40(24):e105862. Epub 20211117. doi: 10.15252/embj.2020105862. PubMed PMID:
34786738; PubMed Central PMCID: PMCPMC8672179.
28.   Buonomo SB, Wu Y, Ferguson D, de Lange T. Mammalian Rif1 contributes to replication
stress survival and homology-directed repair. J Cell Biol. 2009;187(3):385-98. doi:
10.1083/jcb.200902039. PubMed PMID: 19948482; PubMed Central PMCID:
PMCPMC2779251.
29.   Ho RK, Kimmel CB. Commitment of cell fate in the early zebrafish embryo. Science.
1993;261(5117):109-11. doi: 10.1126/science.8316841. PubMed PMID: 8316841.
30.   Siefert JC, Georgescu C, Wren JD, Koren A, Sansam CL. DNA replication timing during
development anticipates transcriptional programs and parallels enhancer activation. Genome
Res. 2017;27(8):1406-16. doi: 10.1101/gr.218602.116. PubMed PMID: 28512193; PubMed
Central PMCID: PMCPMC5538556.
31.   Alavi S, Ghadiri H, Dabirmanesh B, Moriyama K, Khajeh K, Masai H. G-quadruplex
binding protein Rif1, a key regulator of replication timing. J Biochem. 2021;169(1):1-14. doi:
10.1093/jb/mvaa128. PubMed PMID: 33169133.
32.   Kimmel CB, Ballard WW, Kimmel SR, Ullmann B, Schilling TF. Stages of embryonic
development of the zebrafish. Dev Dyn. 1995;203(3):253-310. doi: 10.1002/aja.1002030302.
PubMed PMID: 8589427.
33.   Kossack ME, Draper BW. Genetic regulation of sex determination and maintenance in
zebrafish (Danio rerio). Curr Top Dev Biol. 2019;134:119-49. Epub 20190321. doi:
10.1016/bs.ctdb.2019.02.004. PubMed PMID: 30999973; PubMed Central PMCID:
PMCPMC6894417.
34.   Bogdanovic O, Fernandez-Miñán A, Tena JJ, de la Calle-Mustienes E, Hidalgo C, van
Kruysbergen I, et al. Dynamics of enhancer chromatin signatures mark the transition from
pluripotency to cell specification during embryogenesis. Genome Res. 2012;22(10):2043-53.
Epub 20120516. doi: 10.1101/gr.134833.111. PubMed PMID: 22593555; PubMed Central
PMCID: PMCPMC3460198.
35.   Dan J, Liu Y, Liu N, Chiourea M, Okuka M, Wu T, et al. Rif1 maintains telomere length
homeostasis of ESCs by mediating heterochromatin silencing. Developmental cell.
2014;29(1):7-19.
36.   Klein KN, Zhao PA, Lyu X, Sasaki T, Bartlett DA, Singh AM, et al. Replication timing
maintains the global epigenetic state in human cells. Science. 2021;372(6540):371-8. Epub
2021/04/24. doi: 10.1126/science.aba5545. PubMed PMID: 33888635; PubMed Central PMCID:
PMCPMC8173839.
37.   Giraldez AJ, Mishima Y, Rihel J, Grocock RJ, Van Dongen S, Inoue K, et al. Zebrafish
MiR-430 promotes deadenylation and clearance of maternal mRNAs. Science.
2006;312(5770):75-9. Epub 20060216. doi: 10.1126/science.1122689. PubMed PMID:
16484454.
38.   Armstrong RL, Das S, Hill CA, Duronio RJ, Nordman JT. Rif1 Functions in a Tissue-
Specific Manner To Control Replication Timing Through Its PP1-Binding Motif. Genetics.
2020;215(1):75-87. Epub 20200306. doi: 10.1534/genetics.120.303155. PubMed PMID:
32144132; PubMed Central PMCID: PMCPMC7198277.
39.   Tzung K-W, Goto R, Saju JM, Sreenivasan R, Saito T, Arai K, et al. Early depletion of
primordial germ cells in zebrafish promotes testis formation. Stem cell reports. 2015;4(1):61-73.

40.   Luo Y, Hartford SA, Zeng R, Southard TL, Shima N, Schimenti JC. Hypersensitivity of
primordial germ cells to compromised replication-associated DNA repair involves ATM-p53-p21
signaling. PLoS Genet. 2014;10(7):e1004471. Epub 20140710. doi:
10.1371/journal.pgen.1004471. PubMed PMID: 25010009; PubMed Central PMCID:
PMCPMC4091704.
41.   Yang Y, Xu W, Gao F, Wen C, Zhao S, Yu Y, et al. Transcription-replication conflicts in
primordial germ cells necessitate the Fanconi anemia pathway to safeguard genome stability.
Proc Natl Acad Sci U S A. 2022;119(34):e2203208119. Epub 20220815. doi:
10.1073/pnas.2203208119. PubMed PMID: 35969748; PubMed Central PMCID:
PMCPMC9407672.
42.   Redl S, de Jesus Domingues AM, Caspani E, Möckel S, Salvenmoser W, Mendez-Lago
M, et al. Extensive nuclear gyration and pervasive non-genic transcription during primordial
germ cell development in zebrafish. Development. 2021;148(2). Epub 20210120. doi:
10.1242/dev.193060. PubMed PMID: 33298460; PubMed Central PMCID: PMCPMC7847270.
43.   Rodríguez-Marí A, Cañestro C, Bremiller RA, Nguyen-Johnson A, Asakawa K,
Kawakami K, et al. Sex reversal in zebrafish fancl mutants is caused by Tp53-mediated germ
cell apoptosis. PLoS Genet. 2010;6(7):e1001034. Epub 20100722. doi:
10.1371/journal.pgen.1001034. PubMed PMID: 20661450; PubMed Central PMCID:
PMCPMC2908690.
44.   Ramanagoudr-Bhojappa R, Carrington B, Ramaswami M, Bishop K, Robbins GM, Jones
M, et al. Multiplexed CRISPR/Cas9-mediated knockout of 19 Fanconi anemia pathway genes in
zebrafish revealed their roles in growth, sexual development and fertility. PLoS Genet.
2018;14(12):e1007821. Epub 20181212. doi: 10.1371/journal.pgen.1007821. PubMed PMID:
30540754; PubMed Central PMCID: PMCPMC6328202.
45.   Yuan K, O'Farrell PH. TALE-light imaging reveals maternally guided, H3K9me2/3-
independent emergence of functional heterochromatin in Drosophila embryos. Genes Dev.
2016;30(5):579-93. Epub 20160225. doi: 10.1101/gad.272237.115. PubMed PMID: 26915820;
PubMed Central PMCID: PMCPMC4782051.
46.   Hawkins RD, Hon GC, Lee LK, Ngo Q, Lister R, Pelizzola M, et al. Distinct epigenomic
landscapes of pluripotent and lineage-committed human cells. Cell Stem Cell. 2010;6(5):479-91.
doi: 10.1016/j.stem.2010.03.018. PubMed PMID: 20452322; PubMed Central PMCID:
PMCPMC2867844.
47.   Laue K, Rajshekar S, Courtney AJ, Lewis ZA, Goll MG. The maternal to zygotic
transition regulates genome-wide heterochromatin establishment in the zebrafish embryo. Nat
Commun. 2019;10(1):1551. Epub 2019/04/06. doi: 10.1038/s41467-019-09582-3. PubMed
PMID: 30948728; PubMed Central PMCID: PMCPMC6449393.
48.   Sima J, Chakraborty A, Dileep V, Michalski M, Klein KN, Holcomb NP, et al. Identifying
cis Elements for Spatiotemporal Control of Mammalian DNA Replication. Cell. 2019;176(4):816-
30.e18. Epub 20181227. doi: 10.1016/j.cell.2018.11.036. PubMed PMID: 30595451; PubMed
Central PMCID: PMCPMC6546437.
49.   Gottesfeld J, Bloomer LS. Assembly of transcriptionally active 5S RNA gene chromatin
in vitro. Cell. 1982;28(4):781-91. Epub 1982/04/01. doi: 10.1016/0092-8674(82)90057-5.
PubMed PMID: 7201351.
50.   Lande-Diner L, Zhang J, Cedar H. Shifts in replication timing actively affect histone
acetylation during nucleosome reassembly. Mol Cell. 2009;34(6):767-74. doi:
10.1016/j.molcel.2009.05.027. PubMed PMID: 19560427; PubMed Central PMCID:
PMCPMC2717743.
51.   Zhang J, Xu F, Hashimshony T, Keshet I, Cedar H. Establishment of transcriptional
competence in early and late S phase. Nature. 2002;420(6912):198-202. Epub 2002/11/15. doi:
10.1038/nature01150. PubMed PMID: 12432398.

52. Adams IR, McLaren A. Identification and characterisation of mRif1: A mouse telomere-
associated protein highly expressed in germ cells and embryo-derived pluripotent stem cells.
Developmental Dynamics: An Official Publication of the American Association of Anatomists.
2004;229(4):733-44.
53. Li P, Wang L, Bennett BD, Wang J, Li J, Qin Y, et al. Rif1 promotes a repressive
chromatin state to safeguard against endogenous retrovirus activation. Nucleic acids research.
2017;45(22):12723-38.
54. Li L, Li P, Chen J, Li L, Shen Y, Zhu Y, et al. Rif1 interacts with non-canonical polycomb
repressive complex PRC1.6 to regulate mouse embryonic stem cells fate potential. Cell Regen.
2022;11(1):25. Epub 20220802. doi: 10.1186/s13619-022-00124-9. PubMed PMID: 35915272;
PubMed Central PMCID: PMCPMC9343540.
55. Zhang W, Yao C, Luo Y, Li Q, Zhao Q, Zhao Y, et al. Rif1 and Hmgn3 regulate the
conversion of murine trophoblast stem cells. Cell Rep. 2022;38(13):110570. doi:
10.1016/j.celrep.2022.110570. PubMed PMID: 35354046.
56. Yoshizawa-Sugata N, Yamazaki S, Mita-Yoshida K, Ono T, Nishito Y, Masai H. Loss of
full-length DNA replication regulator Rif1 in two-cell embryos is associated with zygotic
transcriptional activation. Journal of Biological Chemistry. 2021;297(6):101367. doi:
https://doi.org/10.1016/j.jbc.2021.101367.
57. Li P, Wang L, Bennett BD, Wang J, Li J, Qin Y, et al. Rif1 promotes a repressive
chromatin state to safeguard against endogenous retrovirus activation. Nucleic Acids Res.
2017;45(22):12723-38. Epub 2017/10/19. doi: 10.1093/nar/gkx884. PubMed PMID: 29040764;
PubMed Central PMCID: PMCPMC5727408.
58. Cermak T, Doyle EL, Christian M, Wang L, Zhang Y, Schmidt C, et al. Efficient design
and assembly of custom TALEN and other TAL effector-based constructs for DNA targeting.
Nucleic Acids Res. 2011;39(12):e82. Epub 2011/04/16. doi: 10.1093/nar/gkr218. PubMed
PMID: 21493687; PubMed Central PMCID: PMCPMC3130291.
59. Siefert JC, Clowdus EA, Goins D, Koren A, Sansam CL. Profiling DNA Replication
Timing Using Zebrafish as an In Vivo Model System. J Vis Exp. 2018;(134). Epub 2018/05/15.
doi: 10.3791/57146. PubMed PMID: 29757277; PubMed Central PMCID: PMCPMC6101039.
60. Link V, Carvalho L, Castanon I, Stockinger P, Shevchenko A, Heisenberg CP.
Identification of regulators of germ layer morphogenesis using proteomics in zebrafish. J Cell
Sci. 2006;119(Pt 10):2073-83. Epub 20060425. doi: 10.1242/jcs.02928. PubMed PMID:
16638810.
61. Thisse C, Thisse B. High-resolution in situ hybridization to whole-mount zebrafish
embryos. Nature Protocols. 2008;3(1):59-69. doi: 10.1038/nprot.2007.514.
62. Lafont AL, Song J, Rankin S. Sororin cooperates with the acetyltransferase Eco2 to
ensure DNA replication-dependent sister chromatid cohesion. Proceedings of the National
Academy of Sciences. 2010;107(47):20364-9. doi: doi:10.1073/pnas.1011069107.
63. Sansam CL. RepliTimer (v1.0.8). 2022.
64. Lawson ND, Li R, Shin M, Grosse A, Yukselen O, Stone OA, et al. An improved
zebrafish transcriptome annotation for sensitive and comprehensive detection of cell type-
specific genes. eLife. 2020;9:e55792. doi: 10.7554/eLife.55792.
65. Neumann T, Herzog VA, Muhar M, von Haeseler A, Zuber J, Ameres SL, et al.
Quantification of experimentally induced nucleotide conversions in high-throughput sequencing
datasets. BMC Bioinformatics. 2019;20(1):258. doi: 10.1186/s12859-019-2849-7.
827