Evidence for an Intricate Relationship Between Express Visuomotor Responses, Postural Control and Rapid Step Initiation in the Lower Limbs

Lucas S. Billen[1], Brian D. Corneil[2,3], Vivian Weerdesteyn[1,4]

[1]Department of Rehabilitation - Donders Institute for Brain, Cognition & Behavior, Radboud University Medical Center, Nijmegen, NL

[2]Department of Physiology & Pharmacology, Western University, London, CA

[3]Robarts Research Institute, London, CA

[4]Sint Maartenskliniek Research, Nijmegen, NL

Corresponding author:

Lucas Billen

Email: lucas.billen@radboudumc.nl

Key Points

• Express visuomotor responses (EVRs) are directionally tuned bursts of muscle activity that aid the

rapid initiation of a goal-directed movement. They are thought to be relayed to the motor

periphery along a rapid subcortical pathway involving the superior colliculus.

• While EVRs have predominantly been studied in reaching, it is unclear whether EVRs extend to

the lower extremities and if so, whether increasing the postural demands of a stepping task

interfere with lower-limb EVR expression.

• We found that when postural demands were low, strong EVRs in the hip abductor muscle gluteus

medius facilitated a rapid stepping response. Conversely, when postural demands were high,

EVRs hindered a fast stepping response, as they necessitated larger, compensatory postural

adjustments prior to step onset.

• These results help us better understand the interaction between ultra-rapid visuomotor

transformations in the EVR network, the postural demands of a given stepping task, and

subsequent step initiation.

Abstract

Recent work has described express visuomotor responses (EVRs) on the upper limb. EVRs are

directionally-tuned bursts of muscle activity that occur within 100 ms of visual stimulus appearance,

facilitating rapid reaching. Rapid stepping responses are also important in daily life, and while there is

evidence of EVR expression on lower limbs, it is unknown whether lower-limb EVRs are influenced by

increased postural demands. Here, we investigate the interaction between stepping-related EVRs and

anticipatory postural adjustments (APAs) that typically precede step initiation.

16 healthy young subjects rapidly stepped towards visual targets presented in front of the left

or right foot. We recorded bilateral surface EMG of gluteus medius (GM), a muscle involved in both

APAs and stepping, and bilateral ground reaction forces. Two conditions were introduced: a lateral or

medial stepping condition with reduced or increased postural demands, respectively. In the lateral

stepping condition, EVRs were robustly and strongly present in stance-side GM, and ground reaction

forces revealed the absence of APAs. Larger EVRs preceded shorter RTs, consistent with EVRs

facilitating step initiation. In contrast, in the medial stepping condition, EVRs were largely absent, and

ground reaction forces revealed the consistent presence of APAs. When occasionally present, EVRs in

the medial stepping condition preceded larger APAs and longer RTs. Thus, while EVRs in lower limbs

can facilitate rapid stepping, their expression is normally suppressed when postural stability is low.

Failing to appropriately suppress EVRs in such situations disrupts postural stability, necessitating larger

compensatory APAs and leading to longer stepping RTs.

36   **Introduction**

In everyday life, extremely fast reaction times are frequently required to adequately respond to

visual stimuli, for example when catching a ball that is suddenly thrown at us. Previous research has

shown that our fast visuomotor system allows for reaction times of ~120 ms in situations where mid-

flight reaching adjustments were required (Day & Brown, 2001; Day & Lyon, 2000; Fautrelle, Ballay,

et al., 2010; Fautrelle, Prablanc, et al., 2010; Soechting & Lacquaniti, 1983). A novel method that has

been proposed to study the fast visuomotor system from a static starting position is through

electromyographic (EMG) measurement of so-called *express visuomotor responses* (EVRs; formerly

called 'visual responses' or 'Stimulus-locked responses'; Corneil et al., 2004; Pruszynski et al., 2010).

These are defined as short-latency bursts of muscle activity that occur in a time-locked window ≈100

ms after stimulus presentation and precede the larger volley of muscle activity associated with

movement initiation. There are compelling similarities between the response properties of EVRs and

the response properties of mid-flight corrections. For example, like on-line corrections (Day & Lyon,

2000), EVRs are directionally tuned toward the visual stimulus, even when the intended movement is

in the opposite direction or temporarily supparessed (Atsma et al., 2018; Gu et al., 2016; Wood et al.,

2015). Further, earlier and larger-magnitude EVRs are provoked by stimuli of high contrast or low

spatial frequency (Kozak et al., 2019; Wood et al., 2015), similar to the response properties of on-line

corrections (Kozak et al., 2019; Veerman et al., 2008). Evidence suggests that EVRs are relayed to the

motor periphery along a subcortical tecto-reticulo-spinal pathway (Contemori et al., 2021b; Corneil &

Munoz, 2014; Glover & Baker, 2019; Gu et al., 2017; Kozak et al., 2020; Pruszynski et al., 2010). This

conclusion is based on the short latency of the EVR, its temporal separation from the larger wave of

muscle recruitment associated with the voluntary movement, and the similarity in stimulus

preferences with those seen in the visual responses in the intermediate superior colliculus (Chen et

al., 2018; Marino et al., 2012; Rezvani & Corneil, 2008).

60   The vast majority of studies have focused on fast visuomotor responses of the upper limb.

Fast stepping responses are arguably as important as fast reaching movements, for example when

rapidly adjusting one's stepping behavior while walking on uneven terrain or when intercepting a ball

while playing soccer. Indeed, there is strong evidence that the fast visuomotor network can also

recruit express responses in lower limb muscles, for instance during on-line pointing adjustments in

an upright standing posture (Fautrelle, Prablanc, et al., 2010), or while making on-line stepping

adjustments in response to an obstacle or a target shift (Nonnekes et al., 2010; Reynolds & Day,

2005). These on-line adjustments can be initiated substantially faster than voluntary stepping

adjustments, with reaction times ranging from ≈105-150 ms (Marigold et al., 2007; Reynolds & Day,

2005; Weerdesteyn et al., 2004). Importantly, postural demands are substantially higher in stepping

compared to reaching, because of the concurrent involvement of our legs in balancing the Centre of

Mass (CoM). This raises the question whether and how higher postural demands are integrated in

the express visuomotor response in the lower extremities. Intriguingly, a few studies have shown

that on-line step adjustments occurred at similar latencies irrespective of the postural demands of

the task (Nonnekes et al., 2010; Reynolds & Day, 2005). These findings suggest that a sudden visual

stimulus triggers a rapid goal-directed adjustment of the stepping movement without integration of

the postural demands, potentially destabilizing the body.

Here, we investigated the interaction between the ultra-rapid EVR response in the lower

extremities during goal-directed stepping and postural control in the form of *anticipatory postural*

adjustments (APAs). APAs typically precede step initiation from standstill and are closely tied to the

size and direction of the ensuing step (Bancroft & Day, 2016; Inaba et al., 2020). To trigger EVRs, we

used a recently developed emerging target paradigm that involves the sudden, but temporally

predictable, appearance of a moving visual target below an occluder (see Figure 1). This paradigm

has been shown to consistently evoke robust EVRs in upper limb muscles while making reaching

movements from a static position (Contemori et al., 2021a; Kozak et al., 2020; Kozak & Corneil,

2021). It is thought that the use of implied motion behind a barrier, which has been shown to

produce strong signals in motion-related areas in dorsal visual stream (Krekelberg et al., 2005),

combined with the high certainty of the time of target appearance, results in a strong visual transient

(Contemori et al., 2021a; Kozak et al., 2020). These properties have been shown to result in earlier,

stronger, and more prevalent EVRs compared to other paradigms. In the current study, we modified

this paradigm into a stepping task.

We recorded EMG from two muscles, tibialis anterior (TA) and gluteus medius (GM),

reasoning that they could be involved in both postural control and fast visuomotor responses. TA is

commonly used to characterize APAs involved in step initiation. It activates bilaterally (yet more

strongly in the stepping leg) during APAs to generate the initial forces that propel the CoM forward.

GM is involved in both the APA and the ensuing stepping movement. APA-related GM activity is

recruited on the *stepping* side to shift the CoM towards the stance side, followed by stepping-related

activity in the *stance* leg to propel the CoM forward and towards the stepping side. Thus, GM in

particular provides the unique opportunity to distinguish APA-related activity from stepping-related

activity prior to step onset.

In order to investigate whether EVRs correspond to APA-related activity (i.e. ipsilateral GM

activity in the stepping limb) or to stepping-related activity (i.e. contralateral GM activity in the

stance limb), we introduced contrasting balance demands, across different blocks of trials. This was

achieved by presenting the stepping targets in front of the participant either somewhat medially or

somewhat laterally, and by varying initial stance width. Based on previous research demonstrating

that stepping direction influences APA expression (Inaba et al., 2020), we expected that these task

manipulations would necessitate strong APAs when subjects step forward and medially from a wider

stance width (high postural demand), but would yield decreased APAs when subjects step forward

and laterally from a narrow stance (low postural demand) (Bancroft & Day, 2016; Inaba et al., 2020).

If, as in the upper limb, EVRs promote movement toward the target, then we expect EVRs to be

generated in the *stance* side GM and possibly TA, as these muscles rapidly propel the body forward

and toward the stepping side. Whereas such stance-leg EVRs may be beneficial for fast lateral

stepping, strong stance-leg EVRs are expected to compromise postural stability in the medial

stepping condition, because the narrow-based end position in the latter condition is less effective in

'catching' lateral CoM movements. Overall, we found that stance-leg EVRs were readily expressed

when subjects stepped laterally (i.e., towards a stable end position), but (when present) preceded

and necessitated larger APAs when subjects stepped medially towards a more unstable posture.

118   **Materials and Methods**

Subjects

16 healthy young subjects (4 males, 12 females) participated in this study. Ages ranged from 19 to 28

years (*M* = 23.35, *SD* = 2.37). Only participants with a BMI under 25 kg/m^2 were included in the study

to minimize the coverage of muscles by adipose tissue, which could compromise the quality of

surface EMG recordings, particularly of GM. None of the participants had any visual, neurological, or

motor-related disorders that could influence their performance in the study. The study protocol was

reviewed by the medical ethical committee (CMO Arnhem-Nijmegen, 2021-13269) and the study was

conducted in accordance with the latest version of the Declaration of Helsinki. All participants

provided written informed consent prior to participation and were free to withdraw from the study

at any time.

Experimental design

The experiment was performed using a Gait Real-time Analysis Interactive Lab (GRAIL, Motek

Medical, The Netherlands). The experimental setup included an M-gait dual-belt treadmill with two

embedded force plates (GRAIL, Motek Medical, The Netherlands) to measure ground reaction forces,

a Wave Wireless electromyography system (Wave Wireless EMG, Cometa, Italy) to record muscle

activity, and a projector (Optoma, UK) to project all visual stimuli. Participants stood on the

stationary M-Gait with each foot placed on a separate force plate. The stepping task was projected

on the treadmill in front of the participant (Figure 1). Previous research has demonstrated the

importance of high-contrast stimuli in eliciting strong EVRs (Kozak & Corneil, 2021; Wood et al.,

2015). We therefore covered the treadmill with a white vinyl mat and darkened the room.

Additionally, all participants wore a cap that shielded their eyes from the light of the projector, which

further increased the relative brightness of the projected stimuli. The stimuli had a luminance of 7

cd/m² against a black background of 0.23 cd/m² (contrast ratio: 30:1).

Each trial started with the appearance of a stationary visual target (solid white circle) 130 cm

in front of the participant, presented on a black background. This distance was chosen based on

previous research demonstrating that during walking, humans tend to fixate their eyes on the ground

approximately two steps ahead (Patla & Vickers, 2003). After being visible for 1000 ms, the target

started moving towards the participant at a constant speed of 0.44 m/s (with a retinal speed of

~11.0°/s for an average sized participant of 169 cm). The target then disappeared behind an occluder

(a light blue rectangle) and followed a straight track downwards to the base of the occluder. It

remained invisible to the participants for a fixed interval of 750 ms. Because the target continued to

move at a constant speed behind the occluder, participants could anticipate the timing of target

reappearance; in the upper limb, such certainty of the time of target reappearance increases EVR

prevalence, increases EVR magnitude, and decreases EVR latency (Contemori et al., 2021a). Once the

invisible target had reached the base of the occluder, the target reappeared randomly in front of the

left or right foot of the participant. Because this is the first study using this paradigm to study EVRs in

the lower extremities, we varied target appearance to investigate its influence on EVR properties.

The targets reappeared underneath the occluder as either (1) a target moving towards the lateral

edges of the treadmill at a constant speed of 0.23 m/s, or (2) a flashed target (one single flash with a

duration of 48 ms (i.e. three frames)). In the present article, only results for the flashed condition will

be reported. We realized in retrospect that moving targets introduce a confound between reaction

time and stepping length (since moving targets move away from the subject). Moreover, we aim to

use this paradigm for comparing EVRs between neurological patients and healthy controls. The

interpretation of between-group comparisons is more straight-forward when the behavioral

response (i.e. step length) is kept constant, which is why we prefer the flashed condition. We found

only minor differences in EVR expression between the flashed and the moving condition. These do

not influence our interpretation of the results (see Supplementary Materials).

Participants were instructed to divide their weight equally between both legs prior to step

onset and to avoid shifting the CoM forward in anticipation of the reappearing target. This was

visually checked during the experiment based on real-time force plate data. Participants were

instructed to perform a full stepping movement upon reappearance of the target, using the leg on

the side of target appearance (i.e. step with the left leg when the target appeared on the left side

and vice versa for the right leg). After having stepped onto the target with the stepping leg, the

stance leg had to be placed next to the stepping leg to complete the stepping movement. It was

emphasized to the participants that speed was the most important parameter in the present study

and that the step had to be initiated as rapidly as possible. After having completed the trial, the

participant returned to the starting position and the subsequent trial was initiated.

Trials were started manually via the D-flow software (Motekforce Link, The Netherlands) by

the experimenter. To account for small variable delays in target presentation, a photodiode

(TSL250R-LF, TAOS, USA) was used to measure the exact moment of target appearance. This was

achieved by placing the diode over a secondary peripheral target presented at the same time as the

actual stepping target. This secondary target was presented outside of the participant's field of view.

All reported measures (i.e. EMG and force plate measures) were aligned to the moment of stimulus

presentation detected by the photodiode.

In order to investigate the interaction between postural control in the form of APAs and

EVRs, targets were presented in front of the stepping foot, either somewhat laterally or somewhat

medially. In the lateral target condition, participants started at a narrow stance width (feet 23 cm

apart) and stepped forward and outward towards a lateral target presented 29 cm from the middle

line of the treadmill (retinal eccentricity ~10°; stepping eccentricity ~11.6°). In the medial target

condition, participants started at a wide stance width (feet 34 cm apart) and stepped forward and

inward towards a medial target presented 9 cm from the middle line (retinal eccentricity of ~3°;

stepping eccentricity ~8.1°). The variation in stance width between the medial and lateral stepping

condition was used to increase the relative contrast with regard to stepping direction and with

regard to balance control: stepping medially from a wide stance width increases the balance

demands, and consequently the need to make an APA, while a narrow stance width on lateral targets

has the opposite effect. The initial stance position was indicated by the projection of small circles at

the desired foot location.

Participants completed 4 blocks of 150 trials (600 in total). Each block consisted of either only

lateral targets or medial targets and the order of the blocks was counterbalanced. Participants were

informed about the condition before each block. Target appearance (moving/flashed) and target side

(left/right) were randomized on each trial. Participants started with a few practice trials to become

familiar with the task.

Figure 1 *- Experimental setup of the emerging target paradigm (not drawn to scale). The paradigm was projected on the floor in front of the participants. The visual target moved down towards the participants, disappeared behind the occluder, and then reappeared in front of right foot of the participant. Participant stepped onto the target upon reappearance, requiring either a lateral (left) or medial (right) stepping response*

Data collection

We recorded muscle activity from bilateral TA and GM using Ag/AgCl surface electrodes placed approximately 2 cm apart and longitudinally on the belly of the muscle (Wave Wireless EMG, Cometa, Italy). Skin preparation and electrode placement were performed in accordance with the SENIAM guidelines (Hermens & Merletti, 1999). The quality of the signal was checked before starting the recording session. EMG and force plate data were sampled at 2000 Hz.

Data processing and analysis

Incorrect trials were excluded from the analysis. Incorrect trials were defined as trials in which participants stepped towards the wrong direction or initiated stepping movement with the contralateral foot. Data analysis was performed using custom-written MATLAB scripts (version 2019a).

Reaction time

Stepping RT was defined as the time from visual target appearance, as measured by the photodiode, to the foot-off moment of the stepping foot. In line with previous research, foot-off was defined as the first sample at which the vertical ground reaction force component (Fz) was lower than one percent of the participants body weight (Rajachandrakumar et al., 2017).

EVR presence and latency

The raw EMG signals were first band-pass filtered between 20 and 450 Hz and subsequently rectified and low-passed filtered at 150 Hz. Second-order Butterworth filters were used. To determine the presence and latency of EVRs, we used a time-series receiver-operating characteristic (ROC) analysis,

as described previously (Gu et al., 2016; Kozak et al., 2020). EMG data were grouped based on target

location (lateral vs medial), and target side (left vs right). EMG activity of GM and TA was then

compared between leftward and rightward steps within any condition. For every sample between

100 ms before and 500 ms after visual stimulus appearance, an ROC analysis was performed and the

area under the ROC curve (AUC) was calculated. This metric indicates the probability that an ideal

observer can discriminate between the sides of stimulus location based solely on EMG activity. The

AUC values range between 0 and 1, where a value of 0.5 indicates chance discrimination and values

of 1 or 0 indicate perfectly correct or incorrect discrimination, respectively. Differences in muscle

recruitment on leftward vs rightward steps is thus crucial to ensure robust EVR detection. In line with

previous research, we set the discrimination threshold to 0.6 (Gu et al., 2016). The time of earliest

discrimination was defined as the time at which the AUC surpassed the discrimination threshold and

remained above the threshold for 16 out of 20 consecutive samples within the pre-defined EVR

epoch of 100-140 ms after stimulus presentation. Compared to reaching studies, which typically used

EVR windows of 80-120 ms (e.g. Gu et al., 2016; Kozak et al., 2020), the epoch used in the current

study was adjusted based on the use of lower contrast stimuli (Kozak & Corneil, 2021), the

conduction velocity of reticulospinal neurons and the additional distance to the lower extremities

(Buford, 2009).

Response magnitude in EVR window

The response magnitude in the EVR window was calculated for each condition within each

participant, regardless of whether an EVR was detected. On a single trial basis, the average EMG

activity during the EVR epoch (100-140 ms) was calculated and normalized against the median peak

EMG activity (in the interval from 140 ms to foot-off) within the moving target condition during

lateral stepping of the respective subject. EMG magnitudes of all trials were then averaged per

condition.

Statistical analysis

Statistical analyses were performed using IBM SPSS statistics software (version 27). The level of

significance was set to $p < .05$ for all analyses. We used paired t-tests to investigate differences in

EVR discrimination times between left and right muscles. Further t-tests were performed to study

whether EVR magnitude and subsequent stepping RT differed between lateral and medial stepping.

To further investigate the relationship between EVR magnitude and subsequent stepping RT, we

determined Spearman's rank correlation coefficients on the single trial data of all participants.

APA magnitude was defined as the maximum vertical ground reaction force component (Fz)

underneath the stepping leg in the interval from 140 ms after target appearance (i.e., the end of the

EVR window) and foot off, normalized to percent total body weight. To establish APA onset, a sliding

one-sample t-test was performed to test for any given time-point if the mean force plate data

(grouped by condition (lateral/medial) and stepping side (stepping/stance side)) significantly deviates

from the respective baseline value that was chosen to be at 100 ms after stimulus presentation. The

first of at least 10 consecutive statistically significant samples was defined as APA onset.

In order to further characterize the relationship between EVR presence and subsequent

stepping RT, trials were split based on median RT, thereby creating a "fast RT" and "slow RT" subset.

Subsequently, time-series ROC analyses were performed on each subset separately to determine EVR

presence within the two subsets. The same criteria for EVR presence as described above were

applied.

Results

Error Rates

Overall error rates were low; 1.06% of all steps involved errors in the medial condition and 0.43% of

all steps involved errors in the lateral condition. Error rates differed greatly between subjects. Two

subjects made only a single error out of 600 trials (error rate: 0.16%). In contrast, the most error

prone subject made 58 errors in total (error rate: 9.7%).

Visual Inspection of EMG Data Indicates the Presence of EVRs

Figure 2 shows muscle recordings of an exemplar participant aligned to visual stimulus onset. The

EMG signal of this participant showed one of the highest signal-to-noise ratios and exemplifies key

features of the recruitment patterns of interest (see supplementary materials for data from other

subjects). The first column shows the mean EMG activity of left and right GM (top rows) or TA

(bottom rows) on lateral and medial steps respectively. In addition, the time-series ROC curve is

plotted above the EMG activity; the times at which this curve goes above/below the 0.6/0.4

thresholds indicates consistent differences in muscle recruitment when the associated leg acts as the

stance or step limb.

The first two rows of Figure 2A, which demonstrate GM activation on lateral steps, indicate

that GM is primarily recruited on the stance limb side (e.g. left GM is active when stepping

rightward). There are two distinct bursts of activity; the first peaks at around 110 ms and the second

one peaks at around 190 ms after stimulus onset. These activation patterns are also visible in the

trial-by-trial representations of recruitment, for which the associated muscle is on the stance

(column 2) or stepping (column 3) limb. As can be inferred from the trial-by-trial activity in column 2,

there is a clear burst of activity ~110 ms after target presentation in almost every trial on the stance

limb. This initial burst of activity is followed at a variable interval by a second longer-lasting burst that

persists until shortly before foot-off (white dots). In contrast, column 3 shows that GM activity on the

stepping limb is suppressed at ~120 ms after target presentation and essentially remains silent

throughout the trial. The initial burst of activity observed on the stance side is the EVR, as it is time-

independent of the subsequent stepping reaction time. The ROC curve (column 1) supports this

notion as the 0.6 threshold is crossed at 102 ms and 110 ms for left and right GM, respectively.

The kinetic consequences of these recruitment patterns are shown in the force plate data in

column 4. From this, it is clear that ground reaction forces increase and decrease on the stance and

stepping side, respectively, soon after the EVR (~140 ms after target appearance). This presentation

is important because it indicates that APAs were not generated in the lateral stepping condition,

given the absence of an initial increase in vertical forces on the stepping side.

The activation patterns on GM and the force plate data are distinctly different in the medial

stepping condition. As expected, APAs were produced in this condition, as shown by the initial

increase in vertical forces on the stepping side at ~150 ms, which induced a CoM shift towards the

stance side before foot lift off (see column 4).

Further, the underlying pattern of GM activation in the medial condition exhibited a tri-

phasic recruitment of stance- and stepping-leg GM. Mean EMG traces show that GM on the stance

limb side was very briefly activated ~100 ms after stimulus presentation, but was then immediately

silenced. Subsequently, stepping-leg GM (e.g. left GM when stepping leftward) became active at

approximately 130-300 ms after stimulus presentation, which is the timing and patterning expected

of an APA. After 250 ms, GM on the stance limb became highly active and remained so through foot-

off.

Figure 2B shows TA activity of the same exemplar subject. Overall, and in contrast to what

was observed on GM, the initial activation pattern for both the lateral and the medial stepping

conditions looks rather similar. TA is symmetrically activated on both the stance and the stepping

side starting at around 120 ms after stimulus presentation. As can be inferred from the ROC curve

(column 1), a discrimination time within the pre-defined EVR window is absent in TA, due to this

symmetrical activation. On the stance side, TA activation is maintained through foot-off, whereas on

the stepping side, this initial activation is subsequently inhibited until shortly before foot-off. While

at first glance it seems that TA on the stepping limb shows EVR-like activity, given that the initial

recruitment is more time-locked to stimulus rather than movement onset, note that TA on the stance

limb is being recruited at the same time. Thus, unlike GM in the lateral stepping condition, the

recruitment of TA lacks the lateralization of recruitment to one limb or another that is a defining

characteristic of an EVR.

380 ***Figure 2.*** *GM (A) and TA (B) muscle activity, time-series ROC analysis and force plate data of an exemplar participant. Data*
is separated based on muscle side (left/right) and stepping condition (lateral/medial). Each condition is presented on a
381 *separate row. All data are aligned to visual stimulus onset (green line). **Column 1:** shows mean EMG activity for the stance-*
side (dark blue) and the stepping side (light blue) of the respective muscle. The time-series ROC curve is shown in gray.
382 *Discrimination times within the SLR epoch (100-140 ms) are indicated by the black vertical line. **Columns 2 and 3:** Trial-by-*
trial EMG activity of the stance side (column 2) and the stepping side (column 3). Intensity of color conveys the magnitude of
383 *EMG activity Each row represents a different trial. Trials are sorted by RT (white dots). **Column 4:** Mean vertical force (F_z)*
exerted by the stance (dark blue) and stepping leg (light blue).

Robust and postural-dependent expression of EVRs on GM but not TA

As is shown in Table 1, EVRs were robustly detected on stance-side GM in the lateral condition for

both left (16/16 participants) and right GM (15/16), whereas only a small number of participants

exhibited EVRs in the medial condition (1/16 for left GM, 3/16 for right GM). Discrimination times

(see Table 1) did not differ significantly between left and right GM during lateral stepping ($p > .1$).

Discrimination times on medial steps were on average later compared to lateral steps (Table 1).

Statistical analysis was, however, not possible due to the low number of participants who expressed

EVRs in the medial condition. Because the comparisons of both discrimination times and EMG

magnitudes in the EVR window between right and left GM yielded similar results, and for reasons of

conciseness, we chose to only report results for left GM for the remainder of the present paper.

Although an increase in TA activity was commonly observed in the EVR time window, initial

TA activation was often symmetrical, thus not yielding a discrimination time in the EVR window. As a

result, we detected EVRs in TA much less consistently compared to GM. The discrimination times for

TA were also considerably longer than for GM, and we suspect that they result from the lateralization

of muscle activity that blended into the later part of the EVR time window. TA is therefore not a

suitable candidate for studying EVRs in the context of this study, hence remaining analyses will focus

on GM. Note that this does not exclude possible EVR detection in other tasks or different initial

postures that might lead to a lateralization of TA activity.

Participant	Left GM		Right GM		Left TA		Right TA	
	Lateral	Medial	Lateral	Medial	Lateral	Medial	Lateral	Medial
1	107	-	100	-	130	-	119	134
2	101	-	102	-	127	138	-	-
3	105	-	107	-	-	117	-	-
4	102	-	110	-	136	-	-	-
5	103	-	111	-	-	-	-	-
6	106	-	100	118	-	-	-	-
7	106	-	-	-	133	134	130	-
8	115	-	112	-	130	-	140	-
9	103	-	102	-	-	-	139	121
10	115	-	113	-	-	-	-	-
11	101	-	106	107	-	-	-	-
12	122	-	113	-	129	122	-	-
13	110	-	105	111	121	-	-	-
14	109	113	103	-	-	-	136	104
15	106	-	103	-	-	-	138	-
16	119	-	107	-	-	-	131	-
Mean	108	113	106	112	129	127	135	120
SD	6.67	-	4.96	5.57	4.72	9.88	4.61	15.04

Table 1 Discrimination times for GM and TA across participants in lateral vs medial stepping (in ms). Empty cells indicate no discrimination time in that condition.

EVR magnitude is higher in lateral stepping compared to medial stepping

Consistent with the observations from visual inspection of the EMG data (Figure 2) and the results from the ROC analysis indicating higher EVR prevalence in the lateral condition, EVR magnitude was significantly higher in the lateral condition (M = 0.12 AU, SD = 0.05) compared to the medial condition (M = 0.05 AU, SD = 0.02; $t(15)$ = 6.12, $p < .001$, Hedges' g = 1.50).

Stronger EVRs precede shorter stepping reaction times

We investigated whether the contrasting postural demands of lateral and medial stepping had an impact on stepping reaction times. Indeed, stepping reaction times were significantly shorter in the lateral condition (M = 314 ms, SD = 61 ms) compared to the medial condition (M = 443 ms, SD = 54 ms; $t(15)$ = 26.23, $p < .001$, *Hedges' g* = 4.58; see Figure 3B). As previous work demonstrated that

stronger EVRs in a reaching paradigm precede short reach RTs, we evaluated whether this also

applied to our stepping paradigm. Indeed, in lateral stepping there was a strong negative correlation

between EVR magnitude and subsequent stepping RT (r = - .582, p < .01), indicating that stronger

EVRs preceded faster stepping RTs. This correlation was absent in the medial stepping condition (r = -

.040).

APAs are present in medial stepping, but not in lateral stepping

Based on previous studies investigating the relationship between APAs and stepping eccentricity

(Bancroft & Day, 2016; Inaba et al., 2020), we hypothesized that APAs would be present in the lateral

stepping condition, albeit with a strongly decreased magnitude compared to medial stepping.

However, the rapid average stepping reaction times of 314 ms in the lateral stepping condition leave

very little time to perform a complete APA. In addition, the representative subject described above

suggests that APAs may be absent altogether in the lateral stepping condition. We therefore

investigated the presence of APAs in the lateral and medial condition. Figure 3A shows the mean

vertical forces across all participants in the lateral and medial stepping conditions when stepping

towards the right side. As can be inferred from the magenta lines (i.e. medial steps) in Figure 3A, and

consistent with the force plate data shown for the representative subject, the small peak in vertical

force underneath the stepping leg and the dip in vertical force underneath the stance leg indicate the

presence of APAs. We found that the vertical force underneath the stepping leg started to exceed

the baseline force at 172 ms after stimulus presentation. In the lateral condition (green lines in Fig 3),

vertical forces immediately started to decrease underneath the stepping leg and increase

underneath the stance leg, indicating that, strikingly, APAs were not only decreased in magnitude,

but generally absent in this condition. The mean force started to significantly deviate from baseline

at 150 ms in both stance and stepping leg, consistent with a shift of body weight towards the stance

leg and unloading of body weight over the stepping leg.

Figure 3. A: *Mean APA activity based on force plate data in lateral and medial stepping underneath the stepping leg (left) and the stance leg (right). Shaded error bars indicate +/- 1 SD dispersion. Vertical lines indicate when the force plate data starts to significantly deviate from baseline, indicating APA onset. Arrows indicate mean stepping reaction times. **B:** Histogram of the trial-by-trial stepping reaction times of all subjects in the lateral (left) and medial (right) stepping condition.*

Relationship between EVRs, APAs and subsequent stepping RT

EVRs precede slow-RT steps in the medial condition

As shown above, APAs are generated in the medial condition. Interestingly, as shown in Fig. 2, EVRs

on the stance side preceded activity associated with APAs on the stepping side. These findings led us

to wonder if stance-side EVRs, if they were they produced in the medial stepping condition, would

influence APAs and overall subsequent stepping behavior. Although EVRs were indeed mostly absent

in the medial condition when performing the time-series ROC analysis for the whole set of trials, we

frequently observed EVR-like activity on the slower half of trials upon visually inspecting the single

trial data of all subjects. We therefore aimed to systematically investigate if EVRs indeed precede

slower stepping RTs by performing separate time-series ROC analyses on the fast and slow RT

subsets of trials, respectively. As shown in Table 2, EVRs were detected in no participants on the fast

subset of trials (2nd column in Table 2). In contrast, 9/16 participants exhibited EVRs on the slow half

of trials (3rd column in Table 2). Thus, EVRs regularly preceded steps with a subsequent slow stepping

reaction time, suggesting that EVR expression in this condition potentially compromised the

subsequent stepping behavior.

If, indeed, stance-side EVRs interfered with step initiation in situations with high postural

demands, then the disruptive consequences of stance-side EVR expression might have to be

compensated for by larger stepping-side APAs. We therefore investigated the relationship between

trial-by-trial EVR magnitude on the slow half of trials in the medial stepping condition and

subsequent APA magnitude (measured by the maximum vertical force underneath the stepping leg).

Indeed, we overall observed positive correlations between EVR and APA magnitude, which were

significant in eight participants, with stronger EVRs generally being followed by larger APAs.

Participant	Medial Stepping		Correlation EVR x APA	
	Fast half	Slow half	*r*	*p*
1	-	-	.112	.534
2	-	100	.320	.061
3	-	108	.474*	.002
4	-	-	.286	.054
5	-	116	.585**	< .001
6	-	-	.408*	.018
7	-	113	.386*	.020
8	-	113	.038	.823
9	-	107	.318	.062
10	-	-	.587**	< .001
11	-	-	.447*	.0003
12	-	102	.360*	.024
13	-	-	.250	.120
14	-	113	.229	.185
15	-	114	.114	.522
16	-	-	.463*	.008

Table 2. *Discrimination times indicating EVR presence across participants in the medial stepping condition for the fast and*

the slow half of trials (in ms). Empty cells indicate no discrimination time in that condition. The last two columns show

correlation coefficients and level of significance between trial-by-trial EVR magnitudes and subsequent APA magnitudes

*(based on force plate data) on the slow half of trials in the medial stepping condition. * Significant at the .05 level. ***

Significant at the .001 level.

Discussion

In this study, we investigated the relationship between postural control and express visuomotor

responses in tibialis anterior and gluteus medius using an emerging target paradigm. Participants

performed fast stepping movements towards projected targets on the floor. These targets were

presented anterolaterally or anteromedially relative to either a narrow or wide stance, respectively.

In line with our hypotheses, we found that the emerging target paradigm could robustly evoke EVRs

in GM. The EVRs corresponded to the stepping movement, meaning stance-side GM was active

within the EVR window, while stepping-side GM activity was virtually absent. In the lateral stepping

condition, this pattern of EVR expression facilitated the rapid execution of the step to propel the

body forward and laterally, as stronger EVRs were followed by shorter stepping reaction times. In

contrast, we found a substantial suppression of EVRs in the medial condition. Together, our findings

point towards an intricate relationship between EVRs, APAs, and step initiation. EVRs precede APAs,

and can be suppressed in a posturally-dependent fashion. Whenever this suppression temporarily

lapsed in a posturally-unstable condition, the EVR on the stance leg perturbed balance, necessitating

a stronger, longer-lasting, and presumably compensatory APA on the stepping leg, which ultimately

lead to longer reaction times.

EVR characteristics are in line with previous studies

EVR Prevalence

We consistently observed early target-directed activity in the GM abductor muscle across all

participants starting at a latency of ≈100ms after stimulus presentation. The short latency and time-

locked nature of the observed GM activity are in line with the definition of an EVR as proposed by

previous reaching studies (e.g. Contemori et al., 2021a; Glover & Baker, 2019; Gu et al., 2017; Kozak

et al., 2019; Pruszynski et al., 2010). Thus, we here demonstrated that the emerging target paradigm

that so efficiently introduced the cognitive factors necessary for EVR expression (Kozak et al., 2020),

can be adapted to a stepping paradigm leading to equally robust EVR expression in the lower limbs.

EVR Latencies

The average EVR latency was 107ms (ranging from 100-122 ms across subjects) and was similar

across all participants and conditions, regardless of stepping reaction time, which underlines the

time-locked nature of the EVRs. Compared to EVRs on upper limb muscles, the reported latencies are

consistent with neural signals needing more time to travel to lower limb. Indeed, the latencies found

in this study fit with the earliest EVR discrimination time in the pectoralis major muscle of 80 ms (Gu

et al., 2016; Kozak et al., 2020; Contemori et al., 2021a; Kozak & Corneil, 2021) plus an additional

average neural signal traveling time to the lower extremity of 20-25 ms via the reticulospinal tract

(Buford, 2009). EVR latencies also depend on stimulus contrast, with high-contrast stimuli promoting

earlier EVRs (Kozak & Corneil, 2021; Wood et al., 2015). This use of lower contrast stimuli in the

current study compared to previous reaching studies may also have led to the longer EVR latencies

reported here. In general, the EVRs in our study preceded the burst of muscle activity associated with

voluntary movement, except for some cases in which the EVR fused with the voluntary stepping

activity, especially on trials with shorter RT. Importantly, EVR latencies also preceded the onset of

APA-related muscle activity on the stepping side by ~60 ms as observed during medial stepping.

EVR Magnitudes

Another defining feature of the EVR is how its magnitude is inversely proportional to movement

reaction time (Corneil et al., 2004; Pruszynski et al., 2010), consistent with the EVR imparting

behaviorally-relevant forces (Gu et al., 2016). Typically, the r-value of such correlations range

between -0.3 and -0.4, meaning that the strength of the upper limb EVR predicts ~10-15% of the

variance of reach reaction time (Gu et al., 2016; Kozak & Corneil, 2021). We observed an even

stronger negative correlation in the lower limb (average r-value of -0.58), but we note that this

relationship is between the EVR magnitude on the *stance* leg and the RT of lift off of the *stepping* leg.

Our force plate data in the lateral stepping condition shows that the likely kinetic consequence of

stance-side GM recruitment and stepping-side GM suppression during the EVR is a rapid increase in

ground reaction force generated by the stance leg and unloading of the stepping leg within less than

150 ms of target presentation, which propels the CoM towards the direction of the target. This is

remarkably rapid and clearly preceded the APAs as generated in the medial stepping condition at

~170 ms post stimulus presentation.

Muscle considerations

The current study underlines the importance of lateralized muscle activity in order to guarantee

robust EVR detection through time-series ROC analysis. Target-selective muscle activity is a key

feature of the EVR, as this provides evidence that the brain accounts for the target location in its

visuomotor transformations, as opposed to a stereotyped response elicited, for example, by startling

stimuli. While TA did show early muscle onsets (i.e. generally within the EVR window), it did so

bilaterally, thus not meeting this criterion. TA was therefore unsuitable to study lower limb EVRs in

the current study context. In contrast, the experimental task required lateralized activity of GM for

both the APA and the stepping movement, but in an opposite way and with different timing, which

allowed us to study their interaction.

As EVR prevalence in the lower limbs is now established, future studies should look into EVR

expression in other lower-limb muscles. For example, it is plausible to assume that EVRs would also

be present in muscles on the stepping side, where they could facilitate step execution (e.g. by

expediting the lifting of the leg). A suitable candidates might be tensor fascia latae, which contributes

to both hip flexion and abduction and previously demonstrated short-latency recruitment (98 ms)

during on-line step adjustments in response to target shifts (Reynolds & Day, 2005).

EVRs facilitate rapid stepping movements in the lateral stepping condition, but are detrimental in

the medial stepping condition

We demonstrated that target location (lateral vs medial) had a major influence on the postural

demands of the stepping task. As emphasized before, step initiation usually starts with an APA phase

that ensures that balance demands are met prior to the subsequent stepping movement (Honeine et

al., 2016). Previous work by Inaba and colleagues (2020) showed that APA size generally decreases as

the stepping direction becomes more lateral (with stepping directions of up to 90° relative to a

forward step). In comparison, the stepping direction in the lateral stepping condition in the current

paradigm only deviated 11° from a forward step. Strikingly, we discovered that APAs were not only

decreased, but generally absent in lateral stepping. APAs were strongly present in medial stepping at

a latency of ~150ms after target appearance.

In conjunction with this finding, we found that the preceding EVR activity in GM

corresponded to stepping-related activity and not APA-related activity. From a biomechanical point

of view, this indicates that EVRs were favorable during lateral stepping and detrimental during

medial stepping. Because APAs apparently were not needed for maintaining balance in the lateral

stepping condition, the observed EVR activity in stance side GM allowed for an extremely rapid step

initiation, by directly propelling the CoM towards the stepping side.

In contrast, APAs were essential prior to step initiation in the medial condition to account for

the increased balance demands. Thus, initially GM needed to be activated on the ipsilateral side to

shift the CoM towards the stance limb. The occasionally observed stepping-related EVR activity in

stance-side GM *prior to the APA* was therefore detrimental to postural stability. Thus, following EVR

expression in this high-postural demand condition, larger APAs were required to compensate for the

EVRs. Due to the present study design, participants knew in advance whether a lateral or a medial

step needed to be performed. Higher-order cortical areas could therefore account for the increased

balance demands in the medial-target condition by a-priori downregulating the EVRs, as they would

otherwise negatively affect step initiation. However, on trials where this cortical inhibition

momentarily lapsed, the subsequently ensuing strong EVR hindered a fast stepping response towards

the target, as postural demands needed to be met first.

These findings complement an earlier study from Nonnekes et al. (2010), which investigated

on-line stepping adjustments in response to either lateral or medial target jumps in stroke patients.

Results showed that the correction speed in medial mid-flight adjustments was significantly slower

compared to lateral stepping adjustments, suggesting that the increased balance demands in the

medial condition compelled the stroke patients to suppress medial stepping adjustments in order to

safely complete the step. A similar comparison can be drawn to studies investigating the fast

visuomotor network during either pro-reaches (towards the target) or anti-reaches (away from the

target) (Gu et al., 2016; Kozak et al., 2020). In those studies, stronger EVRs preceded faster RTs in

pro-reaches, but slower RTs in anti-reaches. Taken together, these results support the notion that

the fast visuomotor system prioritizes the rapid goal-directed movement *towards* the target, while

the experimental context (i.e. reaching away from the target during reaching or increasing postural

demands during stepping) can dampen EVR expression. In situations where the dampening of EVR

expression fails, subsequent phases of muscle recruitment have to be increased in order to correct

for the kinetic consequences of the inappropriate EVR.

Neural correlates of EVRs and postural control

Although the present paradigm taps into a subcortical fast-latency pathway evoking the EVR, the

influence of higher-order cortical areas on the subcortical pathway via top-down processes should

not be underestimated. In fact, specific features of the emerging target paradigm proposed by Kozak

and colleagues (2020) engage higher-order processes, such as implied motion (Krekelberg et al.,

2005) and temporal predictability (Contemori et al., 2021a) that help potentiate EVRs. Multiple

studies have shown that top-down processes can significantly influence the properties of the EVRs.

For example, EVR characteristics are influenced by expectations or cueing of target location, the

immediacy of movement, hand-path barriers, or instructions to move away from the presented

target (Contemori et al., 2021a, 2021b, 2022; Gu et al., 2016, 2017; Kozak et al., 2020; Wood et al.,

2015). These findings are consistent with the hypothesis that EVRs are regulated via the midbrain

superior colliculus and its downstream connections with the brainstem reticular formation (Boehnke

& Munoz, 2008; Corneil et al., 2004; Corneil & Munoz, 2014; Glover & Baker, 2019). Here we

demonstrate that the expectation of postural instability can suppress EVRs in the lower limb. APAs

are also thought to be regulated via a complex cortical-subcortical interplay involving primary

sensorimotor cortex, supplementary motor area (Jacobs et al., 2009; Richard et al., 2017), basal

ganglia and brainstem (Takakusaki, 2017). In addition, evidence suggests that the reticulospinal tract

plays an important role in postural control and particularly the execution of APAs (Drew et al., 2004;

Massion, 1992; Takakusaki et al., 2016). The signals are then projected to interneurons and

motoneurons via the reticulospinal tract (Schepens & Drew, 2004; Takakusaki et al., 2016). Thus, it is

plausible to assume that both postural control and EVRs innervate GM via the reticulospinal tract.

However, APA onset latencies were longer compared to the hyper-direct subcortical EVR responses

(see Results). Suppression of the EVR response in high postural-demand conditions was therefore

crucial in order to maintain balance. This suppression potentially takes place at the level of the

reticular formation in the brainstem. Future studies should explore this further.

Limitations and future research

Participants knew in advance about the postural demands of the task. The decrease in EVR

prevalence and magnitude in the medial condition could therefore be caused by higher-order brain

areas suppressing the EVR network on a block-by-block basis. Future research could further

investigate this cortical-subcortical interaction by presenting lateral and medial targets in an

intermixed way rather than separated by block. If the decrease in EVR prevalence and magnitude in

medial conditions indeed depends on prior knowledge of the increased balance demands of the task,

multiple scenarios are possible in an intermixed paradigm: on the one hand, participants may

prioritize balance over speed in which case the EVRs will be minimized in both the medial and lateral

stepping conditions. On the other hand, the initial response may be to step as quickly as possible

until cortical areas intervene to produce an APA when stepping medially, in which case the EVRs will

appear strong in both medial and lateral stepping conditions, but as a consequence will negatively

impact stepping RTs in the medial condition.

Our study presented medial or lateral targets at a retinal eccentricity of 2.8° or 9.9°,

respectively, relative to the line of sight. We believe that this difference in retinal eccentricity did not

cause the general absence of EVRs to medial targets. Evidence from the upper limb suggests that the

EVR is tuned to target position relative to the hand, not relative to the current line of sight (Gu et al

2018). Indeed, robust EVRs on upper limb muscles (Gu et al 2018; Kearsley et al 2022) or other

related profiles of muscle recruitment (Cross et al. 2019) can be initiated by parafoveal visual events,

providing the eye and hand are not initially aligned. Nevertheless, future studies should adjust the

paradigm such that targets are presented at the same retinal eccentricity, or incorporate a range of

potential target locations for both posturally stable or unstable starting locations. Holding on to a

handrail could also provide another means of manipulating postural demands during medial

stepping.

Conclusions

Our results suggest an intricate relationship between ultra-rapid, albeit posturally-dependent

expression of EVRs, online control of the APA, and subsequent step initiation. The emerging target

paradigm evoked strong and reliable EVRs in the hip abductor muscle gluteus medius on the stepping

side in a low postural-demand condition, which facilitated rapid step initiation. Such strong EVRs

coincided with an absence of APAs. However, as postural demands were increased, APAs became

essential in order to maintain postural stability during step initiation. In this condition, stance-limb

EVRs were largely absent. On trials where EVR suppression temporarily lapsed, EVRs disrupted

postural stability, which necessitated larger APAs and longer stepping reaction times. We suggest

that regulation of the putative subcortical EVR network is mediated by higher-order cortical areas.

This top-down modulation is dependent on the expectation of postural (in)stability of the upcoming

step. The successful adaption of the emerging target paradigm into a stepping task greatly increases

the potential to further investigate this interaction between various neural networks in future

studies across different disease states and across the lifespan.

References

Atsma, J., Maij, F., Gu, C., Medendorp, W. P., & Corneil, B. D. (2018). Active Braking of Whole-Arm

Reaching Movements Provides Single-Trial Neuromuscular Measures of Movement

Cancellation. *Journal of Neuroscience*, *38*(18), 4367–4382.

https://doi.org/10.1523/JNEUROSCI.1745-17.2018

Bancroft, M. J., & Day, B. L. (2016). The Throw-and-Catch Model of Human Gait: Evidence from

Coupling of Pre-Step Postural Activity and Step Location. *Frontiers in Human Neuroscience*,

10. https://doi.org/10.3389/fnhum.2016.00635

Boehnke, S. E., & Munoz, D. P. (2008). On the importance of the transient visual response in the

superior colliculus. *Current Opinion in Neurobiology*, *18*(6), 544–551.

https://doi.org/10.1016/j.conb.2008.11.004

Buford, J. A. (2009). Reticulospinal System. In *Encyclopedia of Neuroscience* (pp. 151–158).

https://doi.org/10.1016/B978-008045046-9.01337-1

Chen, C.-Y., Sonnenberg, L., Weller, S., Witschel, T., & Hafed, Z. M. (2018). Spatial frequency

sensitivity in macaque midbrain. *Nature Communications*, *9*(1), 2852.

https://doi.org/10.1038/s41467-018-05302-5

Contemori, S., Loeb, G. E., Corneil, B. D., Wallis, G., & Carroll, T. J. (2021a). The influence of temporal

predictability on express visuomotor responses. *Journal of Neurophysiology*, *125*(3), 731–

747. https://doi.org/10.1152/jn.00521.2020

Contemori, S., Loeb, G. E., Corneil, B. D., Wallis, G., & Carroll, T. J. (2021b). Trial-by-trial modulation of

express visuomotor responses induced by symbolic or barely detectable cues. *Journal of*

Neurophysiology. https://doi.org/10.1152/jn.00053.2021

Contemori, S., Loeb, G. E., Corneil, B. D., Wallis, G., & Carroll, T. J. (2022). Symbolic cues enhance

express visuomotor responses in human arm muscles at the motor planning rather than the

visuospatial processing stage. *Journal of Neurophysiology*.

https://doi.org/10.1152/jn.00136.2022

Corneil, B. D., & Munoz, D. P. (2014). Overt Responses during Covert Orienting. *Neuron*, *82*(6), 1230–

1243. https://doi.org/10.1016/j.neuron.2014.05.040

Corneil, B. D., Olivier, E., & Munoz, D. P. (2004). Visual responses on neck muscles reveal selective

gating that prevents express saccades. *Neuron*, *42*(5), 831–841.

https://doi.org/10.1016/s0896-6273(04)00267-3

Day, B. L., & Brown, P. (2001). Evidence for subcortical involvement in the visual control of human

reaching. *Brain*, *124*(9), 1832–1840. https://doi.org/10.1093/brain/124.9.1832

Day, B. L., & Lyon, I. N. (2000). Voluntary modification of automatic arm movements evoked by
motion of a visual target. *Experimental Brain Research*, *130*(2), 159–168.
https://doi.org/10.1007/s002219900218

Drew, T., Prentice, S., & Schepens, B. (2004). Cortical and brainstem control of locomotion. *Progress*
in Brain Research, *143*, 251–261. https://doi.org/10.1016/S0079-6123(03)43025-2

Fautrelle, L., Ballay, Y., & Bonnetblanc, F. (2010). Muscular synergies during motor corrections:
Investigation of the latencies of muscle activities. *Behavioural Brain Research*, *214*(2), 428–
436. https://doi.org/10.1016/j.bbr.2010.06.015

Fautrelle, L., Prablanc, C., Berret, B., Ballay, Y., & Bonnetblanc, F. (2010). Pointing to double-step
visual stimuli from a standing position: Very short latency (express) corrections are observed
in upper and lower limbs and may not require cortical involvement. *Neuroscience*, *169*(2),
697–705. https://doi.org/10.1016/j.neuroscience.2010.05.014

Glover, I. S., & Baker, S. N. (2019). Multimodal stimuli modulate rapid visual responses during
reaching. *Journal of Neurophysiology*, *122*(5), 1894–1908.
https://doi.org/10.1152/jn.00158.2019

Gu, C., Pruszynski, J. A., Gribble, P. L., & Corneil, B. D. (2017). Done in 100 ms: Path-dependent
visuomotor transformation in the human upper limb. *Journal of Neurophysiology*, *119*(4),
1319–1328. https://doi.org/10.1152/jn.00839.2017

Gu, C., Wood, D. K., Gribble, P. L., & Corneil, B. D. (2016). A Trial-by-Trial Window into Sensorimotor
Transformations in the Human Motor Periphery. *The Journal of Neuroscience*, *36*(31), 8273–
8282. https://doi.org/10.1523/JNEUROSCI.0899-16.2016

Hermens, H. J., & Merletti, R., Freriks, Bart. (1999). *SENIAM: European recommendations for surface*
electromyography. Roessingh Research and Development.

Honeine, J.-L., Schieppati, M., Crisafulli, O., & Do, M.-C. (2016). The Neuro-Mechanical Processes That
Underlie Goal-Directed Medio-Lateral APA during Gait Initiation. *Frontiers in Human*
Neuroscience, *10*. https://doi.org/10.3389/fnhum.2016.00445

Inaba, Y., Suzuki, T., Yoshioka, S., & Fukashiro, S. (2020). Directional Control Mechanisms in
Multidirectional Step Initiating Tasks. *Frontiers in Human Neuroscience*, *14*, 178.
https://doi.org/10.3389/fnhum.2020.00178

Jacobs, J. V., Lou, J. S., Kraakevik, J. A., & Horak, F. B. (2009). The supplementary motor area
contributes to the timing of the anticipatory postural adjustment during step initiation in
participants with and without Parkinson's disease. *Neuroscience*, *164*(2), 877–885.
https://doi.org/10.1016/j.neuroscience.2009.08.002

Kozak, R. A., Cecala, A. L., & Corneil, B. D. (2020). An Emerging Target Paradigm to Evoke Fast

Visuomotor Responses on Human Upper Limb Muscles. *JoVE (Journal of Visualized*

Experiments), *162*, e61428. https://doi.org/10.3791/61428

Kozak, R. A., & Corneil, B. D. (2021). High-contrast, moving targets in an emerging target paradigm

promote fast visuomotor responses during visually guided reaching. *Journal of*

Neurophysiology, *126*(1), 68–81. https://doi.org/10.1152/jn.00057.2021

Kozak, R. A., Kreyenmeier, P., Gu, C., Johnston, K., & Corneil, B. D. (2019). Stimulus-Locked Responses

on Human Upper Limb Muscles and Corrective Reaches Are Preferentially Evoked by Low

Spatial Frequencies. *ENeuro*, *6*(5). https://doi.org/10.1523/ENEURO.0301-19.2019

Krekelberg, B., Vatakis, A., & Kourtzi, Z. (2005). Implied motion from form in the human visual cortex.

Journal of Neurophysiology, *94*(6), 4373–4386. https://doi.org/10.1152/jn.00690.2005

Marigold, D. S., Weerdesteyn, V., Patla, A. E., & Duysens, J. (2007). Keep looking ahead? Re-direction

of visual fixation does not always occur during an unpredictable obstacle avoidance task.

Experimental Brain Research, *176*(1), 32–42. https://doi.org/10.1007/s00221-006-0598-0

Marino, R. A., Levy, R., Boehnke, S., White, B. J., Itti, L., & Munoz, D. P. (2012). Linking visual response

properties in the superior colliculus to saccade behavior. *European Journal of Neuroscience*,

35(11), 1738–1752. https://doi.org/10.1111/j.1460-9568.2012.08079.x

Massion, J. (1992). Movement, posture and equilibrium: Interaction and coordination. *Progress in*

Neurobiology, *38*(1), 35–56. https://doi.org/10.1016/0301-0082(92)90034-c

Nonnekes, J. H., Talelli, P., de Niet, M., Reynolds, R. F., Weerdesteyn, V., & Day, B. L. (2010). Deficits

Underlying Impaired Visually Triggered Step Adjustments in Mildly Affected Stroke Patients.

Neurorehabilitation and Neural Repair, *24*(4), 393–400.

https://doi.org/10.1177/1545968309348317

Patla, A., & Vickers, J. (2003). How far ahead do we look when required to step on specific locations

in the travel path during locomotion? *Experimental Brain Research. Experimentelle*

Hirnforschung. Expérimentation Cérébrale, *148*, 133–138. https://doi.org/10.1007/s00221-

002-1246-y

Pruszynski, J. A., King, G. L., Boisse, L., Scott, S. H., Flanagan, J. R., & Munoz, D. P. (2010). Stimulus-

locked responses on human arm muscles reveal a rapid neural pathway linking visual input to

arm motor output. *European Journal of Neuroscience*, *32*(6), 1049–1057.

https://doi.org/10.1111/j.1460-9568.2010.07380.x

Rajachandrakumar, R., Fraser, J. E., Schinkel-Ivy, A., Inness, E. L., Biasin, L., Brunton, K., McIlroy, W. E.,

& Mansfield, A. (2017). Atypical anticipatory postural adjustments during gait initiation

among individuals with subacute stroke. *Gait & Posture*, *52*(1), 325–331.

https://doi.org/10.1016/j.gaitpost.2016.12.020

Reynolds, R. F., & Day, B. L. (2005). Rapid visuo-motor processes drive the leg regardless of balance

828   constraints. *Current Biology*, *15*(2), R48–R49. https://doi.org/10.1016/j.cub.2004.12.051

829 Rezvani, S., & Corneil, B. D. (2008). Recruitment of a head-turning synergy by low-frequency activity

830   in the primate superior colliculus. *Journal of Neurophysiology*, *100*(1), 397–411.

831   https://doi.org/10.1152/jn.90223.2008

832 Richard, A., Van Hamme, A., Drevelle, X., Golmard, J.-L., Meunier, S., & Welter, M.-L. (2017).

833   Contribution of the supplementary motor area and the cerebellum to the anticipatory

834   postural adjustments and execution phases of human gait initiation. *Neuroscience*, *358*, 181–

835   189. https://doi.org/10.1016/j.neuroscience.2017.06.047

836 Schepens, B., & Drew, T. (2004). Independent and convergent signals from the pontomedullary

837   reticular formation contribute to the control of posture and movement during reaching in

838   the cat. *Journal of Neurophysiology*, *92*(4), 2217–2238.

839   https://doi.org/10.1152/jn.01189.2003

840 Soechting, J. F., & Lacquaniti, F. (1983). Modification of trajectory of a pointing movement in

841   response to a change in target location. *Journal of Neurophysiology*, *49*(2), 548–564.

842   https://doi.org/10.1152/jn.1983.49.2.548

843 Takakusaki, K. (2017). Functional Neuroanatomy for Posture and Gait Control. *Journal of Movement*

844   *Disorders*, *10*(1), 1–17. https://doi.org/10.14802/jmd.16062

845 Takakusaki, K., Chiba, R., Nozu, T., & Okumura, T. (2016). Brainstem control of locomotion and

846   muscle tone with special reference to the role of the mesopontine tegmentum and

847   medullary reticulospinal systems. *Journal of Neural Transmission (Vienna, Austria: 1996)*,

848   *123*(7), 695–729. https://doi.org/10.1007/s00702-015-1475-4

849 Veerman, M. M., Brenner, E., & Smeets, J. B. J. (2008). The latency for correcting a movement

850   depends on the visual attribute that defines the target. *Experimental Brain Research*, *187*(2),

851   219–228. https://doi.org/10.1007/s00221-008-1296-x

852 Weerdesteyn, V., Nienhuis, B., Hampsink, B., & Duysens, J. (2004). Gait adjustments in response to an

853   obstacle are faster than voluntary reactions. *Human Movement Science*, *23*(3), 351–363.

854   https://doi.org/10.1016/j.humov.2004.08.011

855 Wood, D. K., Gu, C., Corneil, B. D., Gribble, P. L., & Goodale, M. A. (2015). Transient visual responses

856   reset the phase of low-frequency oscillations in the skeletomotor periphery. *European*

857   *Journal of Neuroscience*, *42*(3), 1919–1932. https://doi.org/10.1111/ejn.12976

Additional Information

Data availability statement

The authors confirm that the data supporting the findings of this study are available within the article

and its supplementary materials.

Competing interests

The authors declare that they have no competing interests.

Author Contributions

Lucas Billen: Conceptualization, Methodology, Software, Formal analysis, Investigation, Data

curation, Writing – Original Draft, Review & Editing, Visualization; **Brian Corneil:** Conceptualization,

Methodology, Software, Resources, Writing - Review & Editing, Supervision, Funding acquisition;

Vivian Weerdesteyn: Conceptualization, Methodology, Resources, Writing - Review & Editing,

Supervision, Project administration, Funding acquisition

Funding

This work was supported by a Donders Centre for Medical Neuroscience (DCMN) grant to BDC and

VW.

Acknowledgements

We thank Drs. Tim Carroll, Gerald Loeb, and Guy Wallis, as well as members of the Corneil and Carroll

lab, for comments on an earlier version of this manuscript. Thank you also to Maartje Meijer for

helping in the early stages of the project.

896   **Supplementary material**

897   **The influence of target appearance on EVR characteristics**

898   Similar to the flashed target condition, EVR prevalence in the moving target condition was high for

899   lateral stepping (16/16 participants) and low for medial stepping (4/16 participants).

900   We investigated the effect of target appearance (flashed vs moving) on EVR latency. Since

901   the medial target condition did not yield discrimination times in every participant, we restricted the

902   analysis to the lateral target condition. EVR latency was significantly longer in the flashed condition

903   (M = 112 ms, SD = 6.83) compared to the moving condition (M = 108 ms, SD = 7.21; $t(15)$ = 2.59, p =

904   .020, Hedges' g = 0.57).

905   A second t-test was performed to investigate the effect of target appearance on EVR

906   magnitude. Here we also included the medial condition. In the lateral conditions, moving targets

907   resulted in significantly higher EVR magnitudes (M = .127, SD = 0.06) compared to flashed targets (M

908   = .120, SD = 0.05; $t(15)$ = 2.32, p < .035, Hedges' g = 0.57). In the medial conditions, there were no

909   significant EVR magnitude differences between flashed and moving targets (p > .05).

910   Thus, our findings indicate that moving targets lead to faster and stronger EVRs compared to

flashed targets. This contradicts the findings of Contemori et al. (2021), who reported lower EVR

latency and higher EVR magnitude in reaches towards transient flashed targets compared to

sustained moving targets. However, several methodological deviations may be responsible for these

contradictory findings, such as the duration of the flashed target (8 ms in the Contemori study,

versus 48 ms here) and the direction of the moving target (downwards in the Contemori study,

sidewards here). The minor difference in EVR latency and EVR magnitude between stepping towards

flashed targets and moving targets might also be the result of a slightly different stepping direction

between the conditions. Since the moving target moved towards the edges of the treadmill,

participants had to make more lateral steps towards moving targets compared to flashed targets.

Summary figures for all participants. For figure description, see Figure 2.