1 **Tmem117 in AVP neurons regulates the counterregulatory response to hypoglycemia.**

4 Sevasti Gaspari, Gwenaël Labouèbe, Alexandre Picard, Xavier Berney, Ana Rodriguez

5 Sanchez-Archidona, Bernard Thorens[*]

7 Center for integrative Genomics, University of Lausanne, Lausanne, Switzerland

15 [*]Correspondence: Bernard Thorens, Center for Integrative Genomics, University of Lausanne CH-

16 1015 Lausanne, Switzerland. Phone: +41 21 2692 3981, Fax: +41 21 692 3985, e-mail:

17 bernard.thorens@unil.ch

Abstract

The counterregulatory response to hypoglycemia (CRR), which ensures a sufficient glucose supply to the brain, is an essential survival function. It is orchestrated by incompletely characterized glucose-sensing neurons, which trigger a coordinated autonomous and hormonal response that restores normoglycemia. Here, we investigated the role of hypothalamic *Tmem117*, identified in a genetic screen as a regulator of CRR. We show that *Tmem117* is expressed in vasopressin magnocellular neurons of the hypothalamus. *Tmem117* inactivation in these neurons increases hypoglycemia-induced vasopressin secretion leading to higher glucagon secretion, an estrus cycle phase-dependent effect in female mice. *Ex vivo* electrophysiological analysis, in-situ hybridization and *in vivo* calcium imaging reveal that *Tmem117* inactivation does not affect the glucose-sensing properties of vasopressin neurons but increases ER-stress, ROS production and intracellular calcium levels accompanied by increased AVP production and secretion. Thus, *Tmem117* in vasopressin neurons is a physiological regulator of glucagon secretion and highlight the role of these neurons in the coordinated response to hypoglycemia.

Introduction

The brain relies almost exclusively on glucose for metabolic energy production. Therefore, maintaining blood glucose levels no lower than ~5 mM is essential for survival (Marty, Dallaporta and Thorens, 2007). When glycemic levels drop below this threshold, a brain-orchestrated neuroendocrine reflex triggers a counterregulatory response (CRR). This is characterized by the secretion of multiple hormones that act on peripheral target organs to stimulate endogenous glucose production, suppress insulin secretion and minimize glucose utilization to restore normoglycemia (Tesfaye and Seaquist, 2010). A key aspect of the CRR is the induction of glucagon (GCG) secretion from pancreatic alpha cells, which trigger enhanced glucose production from the liver (Ramnanan *et al.*, 2011; Thorens, 2022). Patients with type 1 or advanced type 2 diabetes display loss of hypoglycemia-induced GCG secretion (Cryer, 2012; Siafarikas *et al.*, 2012; Bisgaard Bengtsen and Møller, 2021), but the underlying mechanism is poorly understood. Evidence suggests that defects in the central nervous system (CNS) control of GCG secretion contribute to this blunted response (Beall, Ashford and McCrimmon, 2012; Stanley, Moheet and Seaquist, 2019).

In the CNS, changes in glucose concentrations are detected by neurons located in several regions but have been studied most extensively in the hypothalamus and brainstem. Several hypothalamic areas are equipped with glucose-sensing neurons, including the ventromedial (VMH), dorsomedial (DMH), lateral, arcuate (ARC), paraventricular (PVN) and supraoptic (SON) nuclei (Stanley, Moheet and Seaquist, 2019). Glucose responsive neurons are activated by either a rise (glucose excited / GE neurons) or a fall (glucose inhibited / GI neurons) in glucose levels. The response to hypoglycemia involves activation by GI neurons of both the parasympathetic and sympathetic branches of the autonomic nervous system, which stimulate GCG secretion and regulates hepatic glucose production, as well as the activation of the hypothalamus-pituitary-adrenal (HPA) axis to stimulate epinephrine and glucocorticoid secretion (Tesfaye and Seaquist, 2010).

Multiple studies have provided evidence in support for the role of hypothalamic and

brainstem glucose sensing neurons in CRR. For instance, cell-specific activation of VMH neuronal

subpopulations expressing glucokinase (Gck) (Meek *et al.*, 2016) or steroidogenic-factor 1 (Sf1)

(Stanley *et al.*, 2016) increases GCG secretion and glycemia, while their inhibition blocks CRR.

Interestingly, genetic inactivation of *Gck* in Sf1 neurons impairs GCG secretion in a sex-

dependent manner (Steinbusch *et al.*, 2016). Glucose responsive neurons located in other nuclei

have also been shown to control GCG secretion. This is the case of cholecystokinin-expressing

neurons of the parabrachial nucleus that control VMH neurons (Garfield *et al.*, 2014); of fibroblast

growth factor 15-expressing neurons of the DMH (Picard *et al.*, 2016; Picard *et al.*, 2021); and of

Glut2 GI neurons of the nucleus tractus solitarius, which activate vagal nerve firing to stimulate

GCG secretion (Lamy *et al.*, 2014).

To identify in an unbiased manner novel hypothalamic regulators of CRR induced by

insulin-induced hypoglycemia, we performed a new genetic screen using a panel of BXD

recombinant inbred mice (Picard *et al.*, 2022). This led to the identification of *Agpat5*, encoding a

lipid biosynthesis enzyme, and the characterization of its essential role in the activation by

hypoglycemia of AgRP neurons and GCG secretion (Strembitska *et al.*, 2022). We further

identified *Tmem117*, located in a QTL on chromosome 15, as another candidate regulator of GCG

secretion.

Here, we show that *Tmem117* is expressed in vasopressin (AVP) magnocellular neurons,

that AVP neurons are in large part GI neurons, and that hypoglycemia induces copeptin (CPP; an

AVP surrogate) and GCG secretion. Genetic inactivation of *Tmem117* in AVP neurons increases,

in a sex-dependent manner, hypoglycemia-induced CPP and GCG secretion. *Tmem117*

inactivation does not affect the glucose sensing properties of AVP neurons, but induces ER-

stress and increases intracellular ROS and Ca^{++} levels as well as AVP mRNA expression.

Results

***Tmem117* is expressed in AVP magnocellular neurons.**

A genetic screen of a panel of 36 recombinant inbred BXD mice for hypoglycemia-induced GCG secretion identified a cQTL on chromosome 8 and one on chromosome 15 (Picard *et al.*, 2022). The candidate gene on chromosome 8 is the lipid-modifying enzyme Agpat5, which is required for hypoglycemia sensing by AgRP neurons, vagal nerve activation and GCG secretion (Strembitska *et al.*, 2022). On chromosome 15 two genes showed strong negative correlation with the GCG trait, *Irak4* and *Tmem117* (Fig S1A-D). The role of *Irak4* in controlling hypothalamic Il-1β signaling and GCG secretion has been reported (Picard *et al.*, 2022). Here, we performed an expression QTL analysis (eQTL) for the level of *Tmem117* mRNA in the hypothalamus of the BXD mouse lines. We identified an eQTL on chromosome 15 at the same position of the cQTL (Fig S1E). Thus indicating that this genomic locus controls both GCG secretion and *Tmem117* expression.

To establish the sites of *Tmem117* expression in the hypothalamus we performed immunofluorescence microscopy analysis. We found strong immunolabeling for Tmem117 in the SON and PVN (Fig 1A) and in the posterior pituitary (Fig 1B). Co-staining for neuropeptides revealed that *Tmem117* was expressed in AVP magnocellular cells (Fig 1E-G). The specificity of the immunostaining was confirmed by the lack of signal in brains of mice with *Tmem117* gene inactivation in AVP neurons (*Tmem117^{fl/fl}*;*AVP-IRES-Cre-D^{tg/+}* mice) (Fig 1C,D). Quantification in three mice and two consecutive hypothalamic sections for each mouse (bregma -0.7 and -0.8) showed that ~90 percent of AVP neurons in the PVN and ~97 percent in the SON were Tmem117 positive (Fig 1H,I). Super-resolution microscopy revealed a punctuated intracellular distribution of Tmem117, which colocalized in part with AVP granules in both the soma and axons of magnocellular neurons (Fig 1J).

Figure S1. Hypothalamic *Tmem117* is a candidate gene for the control of insulin-induced glucagon secretion.

(A) Clinical quantitative trait locus (cQTL) analysis revealed two genomic regions controlling insulin-induced GCG secretion, on chromosome 8 and 15.

(B) The peak on chromosome 15, between markers c15.loc87 and rs13482723, contains 42 genes.

(C) The genes showing the highest correlation between their hypothalamic mRNA expression and plasma GCG are *Irak4* and *Tmem117*.

(D) Correlation between the level of *Tmem117* expression in the hypothalamus of the 36 BXD mouse lines and insulin-induced plasma GCG levels.

(E) Expression quantitative trait locus (eQTL) analysis for *Tmem117* expression in the hypothalamus of the 36 BXD mouse strains. A cis eQTL was identified on chromosome 15 with a LRS of 43.3.

Panels A-C are adapted from *Picard et al., 2022*.

LRS: likelihood ratio statistic, TPM: transcripts per million.

Figure 1. *Tmem117* is expressed in AVP magnocellular neurons.

Immunofluorescence microscopy detection of Tmem117 in the mouse hypothalamus and pituitary:

(A, B) Positive immunostaining was observed in the PVN, the SON and the posterior pituitary.

(C, D) No immunostaining was detected in the hypothalamus and pituitary of mice with constitutive

inactivation of *Tmem117* in AVP neurons.

(E) Co-staining for Tmem117 and AVP in the PVN.

(F) Co-staining for Tmem117 and AVP in the SON.

(G) Co-staining for Tmem117 and AVP in the posterior pituitary.

(H, I) Quantification of AVP$^+$/Tmem117$^+$ cells in the PVN and SON, respectively.

(J) Higher resolution depiction of an AVP$^+$/Tmem117$^+$ neuron in the PVN captured with structured

illumination microscopy. The white arrows highlight representative positions of signal co-localization.

3V: 3rd ventricle, AP: Anterior pituitary, MZ: Medial zone, opt: optic tract, PP: Posterior pituitary, PVN:

Paraventricular nucleus, SCh: Suprachiasmatic nucleus, SON: Supraoptic nucleus. Scale bar=100μm for

panels A-G, =10μm for panel J.

Tmem117 inactivation in AVP neurons increases insulin-induced copeptin and glucagon

secretion.

To inactivate *Tmem117* specifically in AVP neurons, we generated *Tmem117^{fl}* mice that

allowed Cre-dependent excision of exon 3, which encodes the third transmembrane domain

(amino acids 93-136) of the predicted structure of Tmem117 (Jumper *et al.*, 2021; Varadi *et al.*,

2022) (Fig S2A,B). To induce *Tmem117^{fl}* recombination in AVP neurons, we used a viral-mediated

approach for expression of Cre recombinase. This was preferred over the use of *AVP-IRES-Cre-*

D$^{tg/+}$ mice because these display reduced endogenous AVP levels (Cheng, Fung and Cheng,

2019). We, thus, constructed an AAV plasmid containing the full-length AVP promoter placed

upstream of a codon-improved Cre recombinase (iCre) sequence (Ponzio *et al.*, 2012) (Fig S2C).

The resulting plasmid was packaged in serotype 6 adeno-associated viruses (AAV6-AVP-iCre),

which infect magnocellular neurons with high efficacy (Ponzio *et al.*, 2012) and have a high

retrograde transport capacity (Salegio *et al.*, 2013). Injection of AAV6-AVP-iCre in the posterior

pituitary of adult *Tmem117$^{fl/fl}$* mice triggered efficient recombination in the PVN and SON as

detected by PCR analysis of genomic DNA (Fig S2D) and by the expression of tdTomato when

the AAV6-AVP-iCre was injected in the posterior pituitary of *Rosa26^{tdTomato}* mice (Fig S2E).

Figure S2. *Tmem117^{fl}* mice and viral construct generation.

(A) Predicted structure and transmembrane localization of the mouse Tmem117 (adapted from AlphaFold).

The pink aminoacids highlighted by the pink arrows are the first (aa 93) and the last (aa 136) encoded by

the exon 3 sequence that corresponds to the whole 3rd transmembrane helix.

(B) Schematic representation of the *Tmem117* wild-type locus (+), the *Tmem117^{fl}* locus (fl), and the

recombined *Tmem117* locus (KO). Primer binding sites and expected amplicon sizes are indicated.

(C) Generation of AAV-AVP-iCre plasmid. The pCDH-CB-iCre plasmid was used as a donor for the iCre
sequence located between the NotI and AgeI restriction sites. The p2.0VPI.EGFP plasmid contained the
full-length 2kb AVP promoter in an AAV vector. The GFP-containing NotI-AgeI fragment of this plasmid was
replaced with the iCre sequence to generate the AAV-AVP-iCre plasmid.
(D) PCR analysis of *Tmem117* recombination. DNA extracted from microdissected PVN and SON of mice
injected in the posterior pituitary with the AAV6-AVP-iCre showed a 225 bp fragment indicative of
Tmem117^{fl} recombination. SON DNA from *Tmem117^{fl}* mice crossed with *AVP-IRES2-cre-D$^{tg/+}$* mice was
used as positive control.
(E) Injection of AAV6-AVP-iCre in the posterior pituitary of *Rosa26$^{tdTomato/+}$* mice led to efficient expression
of tdTomato in AVP magnocellular neurons of the PVN and SON.
AVP: Vasopressin, PP: Posterior pituitary, PVN: Paraventricular nucleus, SON: Supraoptic nucleus. Scale
bar=100μm.
207

To assess the role of *Tmem117* in GCG secretion, *Tmem117$^{fl/fl}$* and *Tmem117$^{+/+}$* mice

were injected with the AAV6-AVP-icre in the posterior pituitary to generate AVP^{TM117KO} and

AVP^{TM117WT} mice, respectively (Fig 2A). Two weeks later the mice received an intraperitoneal (i.p.)

injection of saline and blood was collected one hour later for plasma CPP and GCG

measurements. The same experiment was repeated one week later with injection of insulin

instead of saline to induce hypoglycemia. Insulin induced the same hypoglycemic levels in

AVP^{TM117WT} and AVP^{TM117KO} male mice (Fig 2B); CPP secretion was increased in both groups of

mice, but significantly more in AVP^{TM117KO} mice than in AVP^{TM117WT} mice (Fig 2C); the same pattern

was observed for the secretion of GCG (Fig 2D).

Female AVP^{TM117WT} mice and *AVP^{TM117KO}* mice were then tested in the same conditions.

Insulin induced the same level of hypoglycemia (Fig 2E) and triggered comparable secretion of

CPP and GCG in both groups of mice (Fig 2F,G). However, when the hormone levels were

analyzed separately for each phase of the estrus cycle, we found that while the insulin-induced

hypoglycemic levels were comparable across the estrus cycle between genotypes (Fig 2H), the

level of secreted CPP and GCG were significantly higher in *AVP^{TM117KO}* mice as compared to

AVP^{TM117WT} mice during the proestrus phase (Fig 2I,J).

Figure 2. *Tmem117* inactivation in AVP neurons enhances hypoglycemia-induced copeptin and glucagon secretion

(A) Experimental scheme. AAV6-AVP-iCre was injected in the posterior pituitary of Tmem117[fl/fl] (AVP[TM117KO]) or Tmem117[+/+] (AVP[TM117WT]) mice.

(B) Glycemia one hour after saline (black) or insulin (red) i.p. injection [n=15-16 per group].

(C) CPP plasma levels one hour after saline or insulin injection [n=13 per group].

(D) GCG plasma levels one hour after saline or insulin injection [n=15-16 per group].

(E) Glycemia one hour after saline (black) or insulin (red) i.p. injection [n=18-22 per group].

(F) CPP plasma levels one hour after saline or insulin injection [n=18-19 per group].

(G) GCG plasma levels one hour after saline or insulin injection [n=17-20 per group].

(H-J) Analysis of glycemic levels and of CPP and GCG secretion following insulin injection in female mice

at each stage of the estrus cycle [n=3-7 per group].

(H) Glycemic levels

(I) CPP plasma levels

(J) GCG plasma levels.

d: day, di: diestrus, est: estrus, INS: insulin, met: metestrus, PP: posterior pituitary, pro: proestrus, SAL:

saline. For panels H-J, lines correspond to the mean value per group and error bars represent ± SEM. B-

G: 2-way ANOVA RM with Bonferroni post-hoc test; H-J: unpaired t-test; *p<0.05, **p<0.01, ***p<0.001.

Glucose responsiveness of SON AVP neurons is not affected by *Tmem117* inactivation.

To assess whether AVP neurons were activated by insulin-induced hypoglycemia

C57Bl/6N male mice were injected with a saline solution or with insulin and their brains were

collected two hours later. Immunofluorescence microscopy analysis revealed that hypoglycemia

(<3.9 mmol/L) robustly increased c-Fos expression in both PVN and SON (Fig 3B,E) with a ~3-

fold increase in the number of c-Fos$^+$ AVP neurons (Fig 3C,F) compared to the saline injected

controls (Fig 3A,D).

To determine whether *Tmem117* inactivation would modify insulin-induced c-Fos

expression in AVP neurons, we prepared AVP^{TM117KO} mice and AVP^{TM117WT} control mice by

injection of AAV6-AVP-iCre in the posterior pituitary of *Tmem117$^{fl/fl}$* and *Tmem117$^{+/+}$* male mice

(Fig 4A). Two weeks later the mice were treated as described above for the C57Bl/6N. Induction

of c-Fos expression in AVP neurons of the SON was comparable between the two mouse groups

(Fig 4B,C).

To determine whether AVP neurons were directly sensitive to hypoglycemia and whether

Tmem117 would modify this sensitivity, we performed patch clamp recordings of AVP neurons in

the presence of 2.5 mM and 0.1 mM glucose (Labouèbe, Thorens and Lamy, 2018). *Tmem117$^{+/+}$*

and *Tmem117$^{fl/fl}$* adult male mice were co-injected in the SON with AAV6-AVP-iCre and an AAV8-

DIO-mCherry to allow fluorescence visualization of the AVP neurons (Fig 4D). SON was preferred

over PVN due to its homogeneous magnocellular AVP population. Direct injection was used

instead of retrograde labeling by injection in the posterior pituitary due to its higher labeling

efficiency. One to two weeks after surgery, we performed patch clamp analysis of AVP neurons.

In brain slices of AVP^{TM117WT} mice, we found that approximately half of the recorded AVP neurons

were activated by low glucose (GI neurons) and that the other half were glucose non-responder

(NR) neurons; no glucose excited (GE) neurons were observed (Fig 4E-H; black). When the same

measurements were performed in cells with *Tmem117* gene inactivation, the proportion of GI and

NR cells and their membrane potential responses were similar to that of AVP^{TM117WT} mice (Fig 4E-

H; red). Thus, AVP neurons were in large part GI neurons and *Tmem117* inactivation did not alter

their glucose responsiveness.

***Tmem117* inactivation increases ER stress, ROS production, intracellular Ca^{++} levels, and**

AVP mRNA expression

Tmem117 has been identified in a siRNA screen of the HTC116 colon cancer cell line as

a negative regulator of ER stress and ROS production (Tamaki *et al.*, 2017). To verify its

contribution to ER stress we used the colon adenocarcinoma cell line SW480, which

endogenously expresses *Tmem117*. SW480 cells were transfected with a siRNA targeting

Tmem117 or a siRNA control and 48 hours later the cells were collected for RNA and protein

extraction. A small reduction in *Tmem117* mRNA levels (log$_2$fold ~ -0.5) robustly increased

expression of the ER stress markers *sXbp1* and *BiP* (Fig 3SA) and calnexin (Fig 3SB). *Tmem117*

silencing in this cell line did not increase ROS production (Fig 3SC) but *Tmem117* overexpression

significantly reduced ROS production (Fig 3SD). These data confirm a link between *Tmem117*

expression and ER stress and ROS production in carcinoma cell lines.

Figure 3. AVP magnocellular neurons are activated by insulin-induced hypoglycemia.

C57BL6/N male mice were injected i.p. with saline or insulin and c-Fos expression (green) in AVP (magenta) neurons was quantitated by immunofluorescence microscopy 2 hours later.

(A) C-Fos expression in the PVN after saline injection.

(B) C-Fos expression in the PVN after insulin injection.

(C) Quantitation of c-Fos positive cells in the PVN after saline and insulin injections [n=4 per group].

(D) C-Fos expression in the SON after saline injection.

(E) C-Fos expression in the SON after insulin injection.

(F) Quantitation of c-Fos positive cells in the SON after saline and insulin injections [n=4 per group].

3V: 3rd ventricle, AVP: Vasopressin, INS: insulin, opt: optic tract, PVN: Paraventricular nucleus, SAL: saline, SON: Supraoptic nucleus. Scale bar=100μm. Lines correspond to the mean value per group and error bars represent ± SEM. Unpaired t-test; *p<0.05, **p<0.01.

Figure 4. Glucose responsiveness of *Tmem117* KO AVP neurons.

(A) Experimental scheme. AAV6-AVP-iCre was injected in the posterior pituitary of Tmem117[fl/fl] (AVP[TM117KO]) or Tmem117[+/+] (AVP[TM117WT]) mice.

(B) C-Fos expression in the SON

(C) Quantitation of c-Fos positive cells in the SON after saline and insulin injections [n=4-5 per group].

(D) Experimental scheme. AAV6-AVP-iCre was co-injected with AAV8-DIO-mcherry in the SON of Tmem117[fl/fl] (AVP[TM117KO]) or Tmem117[+/+] (AVP[TM117WT]) mice.

(E) Proportion of GI and NR AVP neurons in the SON [n=16-18 per group].

(F) Representative traces of GI and NR AVP neurons in each group.

(G) Membrane potential was increased comparably in response to 0.1 mM glucose in both groups.

(H) No change in membrane potential in NR neurons of both genotypes [n=7-9 per group].

AVP: Vasopressin, d: day, GI: glucose inhibited, INS: insulin, NR: non-responding, opt: optic tract, SAL: saline, w: week. Scale bar=100µm. Lines correspond to the mean value per group and error bars represent

 ± SEM. C: 2-way ANOVA with Tukey post-hoc test; E: Fisher's exact test; G-H: paired t-test for all graphs

314 except Δ membrane potential (unpaired); nsp>0.05, *p<0.05, ***p>0.001.

Figure S3. Tmem117 is modulating ER stress and ROS production in SW480 cells.

(A) Expression levels of ER stress markers in SW480 cells. qRT-PCR quantification [n=8-9 per group].

(B) Protein levels of calnexin in SW480 cells upon *Tmem117* silencing. Western blot quantification [[n=6 per group].

(C) Intracellular ROS measurement in SW480 cells. DHE fluorescent signal in SW480 cells [n=16 per group].

(D) Intracellular ROS measurement in SW480 cells upon overexpression of *Tmem117*. DHE fluorescent signal in SW480 cells [n=16 per group].

DHE: dihydroethidium, RFU: relative

fluorescent units. Bars correspond to the mean value per group and error bars represent ± SEM. Unpaired

t-test; nsp>0.05, **p<0.01, ***p>0.001.

To test whether inactivation of *Tmem117* in AVP magnocellular neurons *in vivo* would also

affect ER stress and ROS production, we prepared *AVP^{TM117KO}* and *AVP^{TM117WT}* mice by injecting

AAV6-AVP-iCre in the posterior pituitary of *Tmem117^{fl/fl}* and *Tmem117^{+/+}* mice. Two weeks later

we collected their brains and measured the level of *Bip* mRNA by in situ hybridization

(RNAScope). Quantitative analysis at the single-cell level revealed higher expression of *Bip*

mRNA in AVP neurons of *AVP^{TM117KO}* mice as compared to those of *AVP^{TM117WT}* mice (Fig 5A,D);

non-AVP cells in the SON showed no difference in *Bip* mRNA expression between the two groups

of mice (Fig 5B,D). Furthermore, we found markedly elevated levels of AVP mRNA in AVP^{TM117KO}

mice as compared to control mice (Fig 5C,D). AVP transcription has been reported to be induced

by ROS (St-Louis et al., 2012, 2014). We, thus, assessed the ROS levels in the SON, by injecting

i.p. the fluorescent ROS indicator dihydroethidium (DHE) in $AVP^{TM117KO}$ and $AVP^{TM117WT}$ mice

twenty-four hours before tissue collection (Fig 5E). Quantification of the DHE fluorescence

intensity revealed higher ROS production in the SON of $AVP^{TM117KO}$ mice (Fig 5F,G).

Both ER stress and ROS can increase intracellular Ca^{++} concentrations (Giorgi *et al.*,

2018), which would enhance neurosecretion. We, thus, measured the intracellular Ca^{++} levels in

AVP neuronal terminals by *in vivo* fiber photometry. $AVP^{TM117KO}$ and $AVP^{TM117WT}$ mice were

prepared by co-injection of the AAV6-AVP-iCre virus and an AAV9-hsyn-FLEX-jGCaMP7 in the

posterior pituitary of $Tmem117^{fl/fl}$ and $Tmem117^{+/+}$ mice. During the same surgery an optic fiber

cannula was implanted in the posterior pituitary for fluorescence monitoring (Fig 5H). Two weeks

later the baseline jGCaMP7 fluorescence signal was recorded in both control and knock-out mice.

This showed that inactivation of *Tmem117* induced a markedly higher baseline Ca^{++} signal (Fig

5I). We then performed Ca^{++} recordings after insulin-induced hypoglycemia. The first measures

were performed at d14-16 after viral injection. A second session of recording was performed over

d19-24 in $AVP^{TM117WT}$ mice injected with saline and these data were used as control for the first

set of experiments. For both sessions, the Ca^{++} signal was recorded during a 30-minute baseline

period and for one hour following i.p. insulin or saline injections (Fig 5J). Each signal was

normalized to the mean intensity of the baseline (corresponding to the last 10 minutes preceding

the i.p. injection) and the area under the curve (AUC) was quantified for the mean trace of each

group over 10-minute periods (Fig 5K). In $AVP^{TM117WT}$ mice insulin injection increased the Ca^{++}

signal as compared to saline injection, a difference that became significant 20 minutes after the

injection. In $AVP^{TM117KO}$ mice insulin induced a significantly higher Ca^{++} signal than in $AVP^{TM117WT}$

mice already 10 minutes after injection (Fig 5J,K). The correct placement of the optic fiber (Fig

5L) was assessed at the end of each experiment and the expression of jGCaMP7 in AVP

magnocellular terminals was verified by fluorescence microscopy (Fig 5M).

Figure 5. *Tmem117* inactivation in AVP neurons increases AVP and BiP mRNA, ROS levels and intracellular calcium in vivo.

(A) Quantification of BiP mRNA per cell in AVP-positive cells of the SON [n=483-704 per group].

(B) Quantification of BiP mRNA per cell in AVP-negative cells of the SON [n=347-584 per group].

(C) Quantification of AVP mRNA per cell [n=483-704 per group].

(D) Fluorescence microscopy detection of AVP mRNA (green) and BiP mRNA (magenta) in the SON.

(E) Experimental scheme. AAV6-AVP-iCre was injected in the posterior pituitary of Tmem117[fl/fl]

(AVP[TM117KO]) or Tmem117[+/+] (AVP[TM117WT]) mice. DHE was injected i.p. (50mg/kg) 24 hours before tissue

collection.

(F) Fluorescence microscopy detection of DHE-derived fluorescence (red) in the SON.

(G) Quantification of DHE-derived fluorescence in the SON [n=8 per group].

(H) Experimental scheme. AAV6-AVP-iCre was injected in the posterior pituitary of Tmem117[fl/fl]

(AVP[TM117KO]) or Tmem117[+/+] (AVP[TM117WT]) mice. An optic fiber was implanted in the same area for signal

analysis in freely moving mice.

(I) Basal calcium levels measured in AVP magnocellular terminals of AVP[TM117WT] and AVP[TM117KO] mice. The

gray dashed line represents the time at which the optic fiber cable was plugged onto the implanted cannula

[n=5-6 per group].

(J) Calcium recordings in AVP magnocellular terminals 10 minutes before and one hour after the i.p injection

of insulin (red and orange) or saline (black). The vertical arrow corresponds to the i.p. injection (time=0).

(K) Quantification of mean signal intensity in 10-minute time bins. [n=6-8 per group].

(L) Stereoscope image of the pituitary gland depicting the optic fiber tract in the posterior pituitary.

(M) Fluorescence microscopy detection of GCaMP (green) in posterior pituitary. Immunofluorescent

labelling was used to localize the AVP terminals (magenta).

AUC: area under the curve, AVP: Vasopressin, d: day, DHE: dihydroethidium, INS: insulin, opt: optic tract,

PP: Posterior pituitary, SAL: saline, SON: Supraoptic nucleus,. Scale bar=20µm. For panels G and K, data

are represented as mean ± SEM. For the violin plots of panels A-C, dashed lines correspond to the median

value and dotted lines to the quartile values. A-C,G: unpaired t-test; I: 2-way ANOVA RM with Bonferroni

post-hoc test; K: 1-way ANOVA with Tukey's post-hoc test; *p<0.05, ***p>0.001.

Together, the above experiments indicate that *Tmem117* inactivation in AVP

magnocellular neurons triggered ER stress, increased ROS production, intracellular Ca^{++} levels

and AVP mRNA levels.

***Tmem117* inactivation progressively triggers cell death.**

Permanently elevated ER stress and increased levels of intracellular ROS and Ca^{++} may

trigger cell death (Zhivotovsky and Orrenius, 2011; Iurlaro and Muñoz-Pinedo, 2016; Giorgi *et al.*,

2018). Therefore, we investigated the fate of AVP neurons one month after *Tmem117* inactivation

in *Tmem117*^{fl/fl} mice co-injected in the SON with AAV6-AVP-iCre and AAV9-hsyn-DIO-EGFP

(*AVPT*^{M117KO} mice). Two groups of control mice were prepared. One consisted of *Tmem117*^{+/+}

mice injected with both viruses (*AVP*^{TM117WT} mice); these allowed to assess by

immunofluorescence microscopy the total number of Tmem117+ cells and the total number of

infected, EGFP-expressing cells. The second group consisted of *Tmem117*^{fl/fl} mice injected only

with AAV9-hsyn-DIO-EGFP (*AVP*^{TM117FL} mice), which were used as control for the effect of the

genotype on the total number of Tmem117+ cells in the SON (Fig S4A). Fluorescence microscopy

images of EGFP and Tmem117 expression and of DAPI staining for the three groups are

presented in Figure S4B.

Quantitative analysis of the immunofluorescence data showed that the total number of

Tmem117+ cells was the same in the two control groups but was reduced by ≥ 50% in *AVP*^{TM117KO}

mice as compared to the control mice (Fig S4C). When considering only the EGFP+ and

Tmem117+ double labelled cells, the reduction was even more striking, reaching ~75% (Fig S4D)

indicating that most of the Cre-infected AVP neurons had disappeared during the post-infection

period. This was also reflected by a reduction in the total number of neurons in the SON, as

measured by DAPI staining (Fig S4E).

To determine whether the loss of AVP neurons would suppress the oversecretion

phenotype, we injected AAV6-AVP-iCre in the posterior pituitary of *Tmem117*^{fl/fl} (AVP^{TM117KO}) and

Tmem117^{+/+} (AVP^{TM117WT}) male mice and, one month later, we assessed hormonal secretion upon

insulin-induced hypoglycemia and collected the brains for histological analysis (Fig 6A).

Immunofluorescence microscopy analysis revealed a significant reduction in the number of AVP

neurons in the PVN and SON of *AVP*^{TM117KO} mice as compared to *AVP*^{TM117WT} mice (Fig 6B-D).

Insulin-induced hypoglycemia reached the same level in both groups (Fig 6E), and induced CPP

and GCG secretion to the same extent (Fig 6F,G). Thus, inactivation of *Tmem117* in AVP neurons

led, over time, to cell death and the disappearance of the oversecretion phenotype.

Figure S4. *Tmem117* inactivation in AVP neurons of the SON overtime leads to cell death.

(A) Experimental scheme. AAV6-AVP-iCre was co-injected with AAV9-DIO-EGFP in the SON of

Tmem117^{fl/fl} (AVP^{TM117KO}) or Tmem117^{+/+} (AVP^{TM117WT}) mice. AAV9-DIO-EGFP alone was injected in the

SON of Tmem117^{fl/fl} (AVP^{TM117FL}) to control for any genotype-derived effect.

(B) Tmem117 (magenta), EGFP (green) and DAPI nuclear staining (white) in the SON. The dashed line is

delimiting the SON area.

(C) Quantification of Tmem117 expressing cells in the SON [n=3-4 per group].

(D) Quantitation of Tmem117 and EGFP double positive cells in the SON [n=3 per group].

(E) Quantitation of the number of cells (DAPI positive) in the SON [n=3-4 per group].

opt: optic tract, SON: Supraoptic nucleus, w: week. Scale bar=100μm. Data are represented as mean ±

SEM. C,E: 1-way ANOVA with Tukey post-hoc test; D: unpaired t-test; *p<0.05, **p<0.01, ***p<0.001.

Figure 6. *Tmem117* inactivation in AVP neurons over time triggers neuronal death.

(A) Experimental timeline. AAV6-AVP-iCre was injected in the posterior pituitary of Tmem117[fl/fl]

(AVP[TM117KO]) or Tmem117[+/+] (AVP[TM117WT]) mice.

(B) Immunofluorescence microscopy detection of AVP in the PVN and the SON.

(C) Quantitation of AVP positive cells in the PVN [n=3-4 per group].

(D) Quantitation of AVP positive cells in the SON [n=3-4 per group].

(E) Glycemia one hour after saline (black) or insulin (red) i.p. injections in AVP[TM117WT] and AVP[TM117KO] male

mice [n=8-9 per group].

(F) CPP plasma levels one hour after saline or insulin injection in AVP[TM117WT] and AVP[TM117KO] male mice

[n=8 per group].

(G) GCG plasma levels one hour after saline or insulin injection in AVP[TM117WT] and AVP[TM117KO] male mice

[n=8-9 per group].

3V: 3[rd] ventricle, AVP: Vasopressin, d: day, INS: insulin, KO: AVP[TM117KO], SAL: saline, opt: optic tract, PVN:

paraventricular nucleus, SON: Supraoptic nucleus, WT: AVP[TM117WT]. Scale bar=100μm. For panels C and

D lines correspond to the mean value per group and error bars represent ± SEM. C,D: unpaired t-test; E-

G: 2-way ANOVA RM with Bonferroni post-hoc test; *p<0.05, **p<0.01, ***p<0.001.

Discussion

In this study we characterized the site of expression and functional role of *Tmem117*, a

genetically controlled regulator of insulin-induced GCG secretion. We found that Tmem117 is

expressed in AVP magnocellular neurons and *Tmem117* inactivation increases hypoglycemia-

induced CPP and GCG secretion. This phenotype is observed in male mice and only in the

proestrus phase in female mice. C-Fos immunodetection showed that AVP neurons are activated

by hypoglycemia and patch clamp analysis revealed that about half of them are GI neurons.

Inactivation of *Tmem117* did not affect the glucose responsiveness of AVP neurons, but instead

increased ER stress, elevated intracellular ROS and Ca^{++} concentrations, leading to increased

AVP production and secretion. These results highlight the physiological role of AVP neurons in

the control of GCG secretion and identify *Tmem117* as a novel regulator of their function.

Tmem117 is an eight transmembrane domain-containing protein (Bürgi *et al.*, 2016). We

found it to be present in AVP neurons of the PVN and SON, with a widespread intracellular

localization, from the soma to the axons and in their terminals located in the posterior pituitary,

the site of AVP secretion in the bloodstream. The level of *Tmem117* mRNA expression in the

hypothalamus of the BXD mouse lines used for the genetic screen showed strong and negative

correlation with hypoglycemia-induced GCG secretion (Picard *et al.*, 2022) (Fig S1D). Our

physiological studies indicated that AVP neurons are activated by insulin-induced hypoglycemia,

leading to CPP secretion, a response that was augmented by *Tmem117* inactivation and was

accompanied by increased GCG secretion. These observations are in agreement with previous

studies showing increased secretion of AVP in response to hypoglycemia (Baylis and Robertson,

1980; Baylis, Zerbe and Robertson, 1981; Chiodera *et al.*, 1992) and that AVP stimulates GCG

secretion through its binding to the AVP V1b receptor of pancreatic α cells (Dunning, Moltz and

Fawcett, 1984; Spruce *et al.*, 1985; Gao, Gérard and Henquin, 1992; Yibchok-anun *et al.*, 2004;

Kim *et al.*, 2021; Liu *et al.*, 2021). On the other hand, these results are also in line with our genetic

screen that showed a negative correlation between *Tmem117* expression and insulin-induced

GCG secretion.

492   Interestingly, the effect of *Tmem117* inactivation on GCG secretion is sex-dependent; it is

present in male mice, but only during the proestrus phase in female mice. Circulating levels of

estradiol peak during proestrus and have been linked to adaptations in AVP magnocellular

neurons and their afferent connections (Sladek and Somponpun, 2008), as well as to decreased

GCG secretion from pancreatic alpha cells (Godsland, 2005; Mårtensson *et al.*, 2009). Thus, our

data suggest a possible contribution of estradiol to the AVP-stimulated GCG release in conditions

of hypoglycemia. Preceding studies have also reported that the CRR displays sex-dimorphism

(Steinbusch *et al.*, 2016; Briski *et al.*, 2017), although the precise mechanisms involved still need

to be fully characterized.

501   In order to investigate the mechanisms by which *Tmem117* inactivation regulates CPP

secretion in hypoglycemic conditions, we explored the glucose responsiveness of the AVP

neurons. First, c-Fos immunofluorescence analysis showed that insulin-induced hypoglycemia

led to a ~3 fold increase in the activation of AVP neurons in the SON, suggesting that they are

glucose responsive. This was confirmed by patch clamp analysis, which revealed that ~50% of

them are GI neurons activated by low extracellular glucose concentrations. Thus, AVP neurons

can directly respond to hypoglycemia. Interestingly, a recent study reported that hypoglycemia

can also activate AVP neurons indirectly, through afferent connections arising from GI neurons of

the basolateral medulla (Kim *et al.*, 2021). Thus, AVP neurons are part of a brainstem-

hypothalamus neuronal circuit where hypoglycemia can be sensed by neurons located at multiple

sites to activate the secretion of AVP leading to increased secretion of GCG.

Patch clamp and c-Fos immunofluorescence analysis revealed that inactivation of

Tmem117 does not affect the glucose responsiveness of AVP neurons. Instead, it induces ER

stress and increases intracellular ROS and Ca^{++} levels, leading to enhanced AVP production and

secretion. These results extend previous observations made in the HCT116 cancer cells showing

that *Tmem117* silencing increases ER stress and ROS production (Tamaki *et al.*, 2017) and are

in line with the reports that increased intracellular ROS stimulate AVP transcription in

magnocellular neurons (St-Louis *et al.*, 2012, 2014). It is striking that a small reduction in

Tmem117 expression in the SW480 cell line (~30%) led to a significantly increased expression of

ER stress markers. This suggests that *Tmem117* expression level has an important regulatory

role. This is compatible with the genetic screen that showed strong correlation across the BXD

lines between the level of *Tmem117* expression and hypoglycemia-induced GCG secretion. Thus,

fine tuning of *Tmem117* expression appears as a physiological mean of controlling AVP neuron

secretory capacity. In contrast, genetic inactivation of *Tmem117*, which leads to permanent

increase in ER stress, ROS production and elevated intracellular Ca^{++} levels, is not compatible

with the long-term survival of these neurons. Indeed, one month after induction of *Tmem117*

recombination we observe neuronal death, probably as a result of apoptosis, as reported in the

HCT116 cells with silencing of *Tmem117* (Tamaki *et al.*, 2017).

Collectively, the present study highlights the physiological role of the AVP neurons in the

CRR and identifies *Tmem117* as a genetic determinant of AVP and GCG secretion. Inactivation

of *Tmem117*, did not affect the glucose sensing properties of AVP GI neurons, but led to increased CPP secretion associated with increased ER stress, intracellular ROS and Ca^{++} levels, and increased AVP mRNA expression. Thus, *Tmem117* emerges as a so far uncharacterized regulator of AVP neuron secretory capacity, by increasing AVP mRNA expression and secretion in response to hypoglycemia. Defining how Tmem117 controls these intracellular processes at the molecular level will, however, require future studies. Finally, this study further demonstrates that the central response to hypoglycemia is highly complex, integrating not only brainstem and hypothalamic glucose responsive neurons that activate the HPA axis and both branches of the autonomic nervous system, but also the secretion in the blood of AVP by magnocellular neurons.

Materials & Methods

RESOURCE AVAILABILITY

Lead contact

Further information and requests for resources and reagents should be directed to and will be fulfilled by the Lead Contact, Bernard Thorens (bernard.thorens@unil.ch).

Materials availability

Free of charge for noncommercial purposes

Data and code availability

Data reported in this paper will be shared by the lead contact upon request.

This paper does not report original code.

Any additional information required to reanalyze the data reported in this paper is available from the lead contact upon request.

EXPERIMENTAL MODELS AND SUBJECT DETAILS

Mice

All animal care and experimental procedures were in accordance with the Swiss National

Institutional Guidelines of Animal Experimentation (OExA; 455.163) with license-approval

(VD3363, VD3674) issued by the Vétérinaire Cantonal (Vaud, Switzerland). Mice were housed

up to 5 per cage in a temperature-controlled room with a 12hr light/dark cycle and ad libitum

access to water and standard laboratory chow (Kliba Nafag). Eight- to ten-week-old mice were

used for all the experiments.

<u>Generation of Tmem117^{fl} mice.</u>

Tmem117^{fl} mice were generated by homologous recombination in embryonic stem cells

(genOway, Lyon, France). A reporter cassette for targeting exon 3, preceded by a splice acceptor

was inserted in anti-sense into the respective intron. Both the cassette and the targeted exon

were flanked by loxP sites and mutated loxP sites to enable deletion monitoring using the FLEx

approach (Schnütgen *et al.*, 2003) (FigS1A). The derived mouse line was developed onto a

C57BL/6N background.

<u>Tmem117^{fl}; AVP-IRES2-cre-D mice.</u>

Tmem117$^{fl/fl}$; AVP-IRES2-cre-D$^{tg/+}$ mice were generated by crossing the Tmem117^{fl} line with the

AVP-IRES2-cre-D (Jackson Laboratory code: 023530) for verifying the specificity of the Tmem117

antibody (Fig 1C,D).

<u>Rosa26^{tdTomato} (Ai14) mice.</u>

Ai14 mice (Jackson Laboratory code: 007914) were used for verifying the specificity of the newly

generated AAV6-AVP-icre.

Cell lines

<u>SW480 cells</u>

The human colorectal cancer cell line SW480 (ATCC CCL-228) was cultured in Leibovitz's L-15

Medium, supplemented with 10% fetal bovine serum (FBS). Cells were incubated at 37°C in a

0.3% CO_2 atmosphere and passaged twice per week up to 30 passages.

METHOD DETAILS

In vivo

Stereotaxic surgery / Optic fiber cannula implantation.

Mice were anesthetized with a ketamine and xylazine solution (100mg/kg ketamine, 5mg/kg xylazine injected i.p.) and were placed on a stereotaxic frame (Stoelting, Chicago, USA). A small incision was made in the skin to reveal the skull. A small opening, to allow the passage of a 33 gauge needle, was made and the viral solution was injected at a rate of 100nl/min. The stereotaxic coordinates (relative to the bregma) and total volume injected for each area were: posterior pituitary: AP: -3, ML: 0, DV: -5.8, 0° angle / 400nl; SON: AP: -0.46, ML: ±1.15, DV: -5.45, 0° angle / 140nl per hemisphere. After the viral injection, the incision was closed by suturing and an analgesic solution was provided (0.1mg/kg buprenorphine s.c.). Ophthalmic ointment was used throughout the procedure to prevent eye dryness. A heat pad was used to maintain the proper body temperature.

For fiber photometry recordings an optic fiber (Doric Lenses B280-2408-6.4) was implanted in the posterior pituitary after the viral injection and the incision was closed with dental cement.

For insulin-induced hypoglycemia experiments 400nl of AA6-AVP-icre [8.1×10^{12} (viral genomes) vg/ml] were injected in the posterior pituitary. For electrophysiological recordings (Fig 4) and viability assessment (Fig S4) 140nl of a 1:1 mix of AAV6-AVP-icre (8.1×10^{12} vg/ml) with AAV8-syn-DIO-mCherry (2.2×10^{13} vg/ml) or AAV9-syn-DIO-EGFP (2.4×10^{13} vg/ml) were injected per hemisphere in SON. For fiber photometry 400nl of a 1:1 mix of AAV6-AVP-icre (8.1×10^{12} vg/ml) with AAV9-syn-FLEX-jGaMP7s (2.5×10^{13} vg/ml) were injected in the posterior pituitary.

Insulin-induced hypoglycemia; blood collection.

Mice were food deprived for 6 hours (8am-2pm). At 12pm, they were placed in individual cages and glycemia was measured. At 1pm, glycemia was measured again and the mice were injected i.p. with either saline (baseline control) or insulin (0.8U/kg). At 2p.m. glycemia was measured

again and 100-120μl of blood were collected under isoflurane-induced general anesthesia by

submandibular vein incision.

Fiber photometry.

Starting 5 days after surgery, mice were habituated to the recording cage and placement of the

optic fiber in 30min sessions at least twice per week for two weeks. On the day of recording, the

mice were placed in the recording cage for 30min and signal recording was started. The optic

fiber was then connected to capture the baseline and stimulated signal intensity. The fiber

photometry system was from Doric Lenses using LED light sources for GCaMP (465nm) and

isosbestic (405nm) measurements; signal was recorded and analyzed using the Doric

Neuroscience Studio software. The GCaMP7 signal was first normalized over the isosbestic

signal ($i=i_{465}-i_{405}$) and then over the mean signal for the whole trace (ΔF/F0). Each point was then

normalized to the baseline signal to obtain the corresponding z-score calculated using the

formula: $z_{(i)}=[\Delta F/F0_{(i)} - median(\Delta F/F0_{baseline})/ MAD_{baseline}]$. For baseline intensity traces z-score

generation was calculated based on an 1 second recording period before the cable was plugged

onto the optic cannula. For insulin-induced hypoglycemia traces z-score generation was

calculated based on the baseline signal recorded over 10 minutes before the i.p. injection.

Estrus cycle monitoring.

The estrus cycle phase was identified based on vaginal cytology samples after violet crystal

staining (McLean *et al.*, 2012).

Tissue collection.

For immunofluorescence microscopy analysis mice were transcardially perfused with 10ml ice-

cold phosphate buffered saline (PBS: 137mM NaCl, 2.7mM KCl, 10mM Na2HPO4, 1.8mM

KH2PO4) followed by 40ml ice-cold paraformaldehyde (PFA, 4%) in PBS. Brain and pituitary were

postfixed in 4% PFA O/N at 4°C and then kept for 24 hours in 30% sucrose in PBS at 4°C. Tissues

were frozen and stored at -80°C.

For SON and PVN DNA extraction, fresh brains were collected and kept in RNAlater solution

(Thermo Fisher Scientific) at 4°C for one week. Then, 250μm thick vibratome sections were

prepared and PVN and SON nuclei were dissected under a stereomicroscope. Microdissected

tissue was frozen with dry ice and stored at -80°C, before DNA extraction.

Ex vivo

<u>Electrophysiological recordings</u>

Tmem117$^{fl/fl}$ and Tmem117$^{+/+}$ mice were injected in the SON with an AAV6-AVP-iCre and an

AAV8-DIO-mCherry. One to two weeks later, they were deeply anesthetized with isoflurane,

decapitated, and their brain taken out and immediately placed in an ice-cold high glucose artificial

cerebrospinal fluid (ACSF) containing (in mM): 125 NaCl, 2.5 KCl, 1.25 NaH2PO4, 1 MgCl2, 2

CaCl2, 26 NaHCO3, and 10 glucose (300 ± 5 mOsm) equilibrated with 95% O2 / 5% CO2. 250

μm coronal sections containing SON were prepared using a vibratome (VT1000S, Leica) and

immediately transferred to an oxygenated ACSF solution containing 2.5mM glucose and

maintained at 32°C for at least 1 h before starting the recordings. Experiments were performed

using an upright epifluorescence microscope (BX51WI, Olympus, Japan) mounted on a motorized

stage coupled to a micromanipulator (MPC-325, Sutter Instrument, USA) and equipped with a

mercury lamp and an Evolve EMCCD camera (Teledyne Photometrics Technology, USA) and

appropriate filters allowing the visualization of mCherry-expressing neurons.

Glucose responsiveness of AVP neurons was then assessed in the whole-cell configuration using

a MultiClamp 700B amplifier associated with a 1440A Digidata digitizer (Molecular Devices).

Borosilicate glass pipettes (resistance: 2-5 MΩ.; GC150F-7.5, Harvard Apparatus, USA) were

prepared with a P-97 horizontal micropipette puller (Sutter Instrument, USA). The patch pipettes

were filled with an intrapipette solution containing (in mM): 130 K-gluconate, 5 NaCl, 1 MgCl2, 10

NaPhosphocreatinine, 10 HEPES, 0.2 EGTA, 4 MgATP, 0.5 Na2GTP (pH 7.2-7.4; 280 ± 5

mOsm). Membrane potential and membrane resistance were monitored in current-clamp mode

in the presence of 2.5 mM or 0.1 mM glucose after a 10–15 min baseline. Neurons with an access

resistance >25 MΩ or changing by >20% during the recording were excluded from the analysis.

Signals were filtered at 2 kHz, digitized at 10 kHz, and collected online with a pClamp 10 data

acquisition system (Molecular Devices).

Cell culture

SW480 cells were plated in 6-well tissue culture plate at a density of 300.000 cells/well. After 48

hours cells were transfected with siRNA against *Tmem117* or siRNA control (30pmol/well) with

lipofectamine RNAiMAX (Thermo Fisher Scientific, Cat# 13778100). 48 hours after transfection

cells were trypsinized and pelleted by mild centrifugation (1200rpm for 3min). Supernatant was

removed and cell pellet was used for RNA and protein extraction using QIAGEN RNeasy Mini kit.

Sample preparation / analysis

Immunofluorescence microscopy.

25 μm-thick tissue sections were washed with PBS for 5min at RT, incubated with blocking

solution (2% normal goat serum + 0.3% Triton x-100 in PBS) for 1 hour at RT and then incubated

O/N with primary antibodies (Guinea Pig anti-(Arg8)-Vasopressin, BMA BIOMEDICALS, Cat# T-

5048; Rabbit anti-Tmem117, Novus, Cat# NBP1-94078) in blocking solution at 4°C. Next, sections

were washed with PBS three times for 10min, incubated with secondary antibodies (Goat anti-

Rabbit Alexa Fluor 488, Molecular Probes, Cat# A-11078; Goat anti-Guinea pig Cy5, Abcam,

Cat# ab102372) in blocking solution for 2 hours at RT, washed again with PBS three times for

10min, incubated with DAPI (1:10000 in PBS) for 20min at RT, washed with PBS three times for

10min, allowed to dry and finally mounted using fluoromount (Sigma F4680).

For c-Fos immunodetection tissue sections were washed with PBS for 5min at RT, incubated with

blocking solution (4% normal goat serum + 0.3% Triton x-100 in PBS) for 1 hour at RT and then

incubated O/N with primary antibody for c-Fos (Rabbit anti-cFos, Cell Signaling, Cat# 2250) in

blocking solution at RT. Next, sections were washed with PBS three times for 10min, incubated

with secondary antibody in blocking solution for 3 hours at RT, washed again with PBS three

times for 10min and incubated O/N with primary antibody for AVP in blocking solution at 4°C. Next,

sections were washed with PBS three times for 10min, incubated with secondary antibody in

blocking solution for 3 hours at RT, washed again with PBS three times for 10min, incubated with

DAPI (1:10000 in PBS) for 20min at RT, washed with PBS three times for 10min, allowed to dry

and finally mounted using fluoromount.

For all secondary antibodies the final concentration in the working solution was 1:500. For primary

antibodies the concentrations used were: anti-Tmem117 1:250, anti-AVP 1:250, anti-cFos

1:1000.

<u>In-situ hybridization (RNAscope)</u>

25 μm-thick tissue sections containing the SON were co-stained by in situ hybridization for AVP

mRNA (Cat# 401391) and BiP mRNA (Cat# 438831-C3) using RNAscope probes and RNAscope

Fluorescent Multiplex Detection Reagents (Advanced Cell Diagnostics, Newark, CA, USA)

following manufacturer's instructions.

<u>DHE fluorescent labelling</u>

Tissue sections were washed with PBS for 5min at RT, allowed to dry and mounted using

fluoromount (Sigma F4680).

<u>Quantitative microscopy analysis</u>

Fluorescence images were acquired on a ZEISS Axio Imager.M2 microscope, equipped with

ApoTome.2 and a Camera Axiocam 702 mono (Zeiss, Germany). Specific filter cubes were used

for the visualization of green (Filter set 38 HE eGFP shift free (E) EX BP 470/40, BS FT 495, EM

BP 525/50), red (Filter set 43 HE Cy 3 shift free (E) EX BP 550/25, BS FT 570, EM BP 605/70),

blue (Filter set 49 DAPI shift free (E) EX G 365, BS FT 395, EM BP 445/50) fluorescence and far

red (Filter set 50 Cy 5 shift free [E] EX BP 640/30, BS FT 660, EM BP 690/50). Different

magnifications were selected using a Zeiss x20 objective (Objective Plan-Apochromat 20x/0.8

M27, FWD=0.55mm) and a x40 oil-immersion objective (Objective C Plan-Apochromat ×40/1.4

Oil DIC M27 [FWD = 0.13 mm]). For super-resolution microscopy the ZEISS ELYRA7 SIM[2]

system was used with an x63 oil-immersion objective.

Cell quantification in PVN and SON was performed on images derived from two consecutive

hypothalamic sections for each mouse (bregma -0.7 and -0.8). Fluorescence intensity

quantification was performed using the QuPath-0.3.2 software on images derived from two

consecutive hypothalamic sections for each mouse (bregma -0.7 and -0.8). Neuroanatomical

landmarks (3rd ventricle, optic tract, fornix) and DAPI staining were used to define the region of

interest and cell area was marked manually by a blinded experimenter.

<u>Western blot.</u>

The protein content of the samples was quantified by Bradford assay. 25µg of total protein per

sample were prepared in SDS-PAGE sample buffer (2% SDS, 10% glycerol, 50mM Tris, 0.1%

bromophenol blue, 5% β-mercaptoethanol) and separated on a 10% SDS-PAGE gel. Proteins

were transferred onto a nitrocellulose membrane, incubated with blocking solution [3% milk in

PBS-T (PBS + 0.1% Tween-20)] for 1 hour at RT, washed with PBS-T 3 times for 10min and

incubated O/N at 4°C with primary antibody (Rabbit anti-Calnexin, Abcam, Cat# ab133615) at a

dilution of 1:1000 in PBS-T. Then membranes were washed with PBS-T 3 times for 10min,

incubated for 1 hour at RT with horseradish peroxidase (HRP) conjugated secondary antibody

(Anti-Rabbit HRP, Amersham, Cat# NA934) at a dilution of 1:10000 in blocking solution, washed

again with PBS-T 3 times for 10min, incubated for 1min with enhanced chemiluminescence (ECL,

Amersham) buffer and the HRP mediated chemiluminescence signal was detected/imaged using

the Fusion FX6 Spectra imaging platform (VILBER). The derived images were analyzed for band

intensity using imageJ. The calnexin signal of each sample was normalized to the total protein

(Pierce Reversible Protein Stain Kit). Data are presented as percent of the control.

<u>DNA extraction / PCR.</u>

DNA was extracted from microdissected tissue using the Arcturus Picopure DNA extraction kit

(Applied biosystems, Ref #11815-00). The extracted DNA was then subjected to PCR for

detection of recombination using the following primers: 5'-CTTTCTTCATAAAAAGCCGGAA

GGCATTAC-3' (212), 5'-GCCTGAAATATAAATATCGCAAGTGAGTGTGC-3' (213), 5'-

CAACTGACCTTGGGCAAGAACATAAAGTG-3' (216) (graphical representation Fig S1A).

RNA extraction / Real time-PCR

Total RNA was extracted from SW480 cell pellets using the QIAGEN RNeasy Mini kit based on

manufacturer's instructions. RNA was then reverse transcribed, and the derived cDNA was

quantified using specific primers against *Tmem117* (FW:5'-TGTGATGCAGGACTGGGAAT-3';

RV:5'-TTGAACTGCATGTGAGGCGT-3'), *Xbp1* (FW:5'- TGGCCGGGTCTGCTGAGTCCG -3';

RV:5'-ATCCATGGGGAGATGTTCTGG -3'), *usXbp1* (FW:5'-CAGCACTCAGACTACGTGCA-3';

RV:5'- ATCCATGGGGAGATGTTCTGG-3'), *sXbp1* (FW:5'-CTGAGTCCGAATCAGGTGCAG-3';

RV:5'-ATCCATGGGGAGATGTTCTGG-3'), *BiP* (FW:5'-TGTTCAACCAATTATCAGCAAACTC-

3'; RV:5'-TAGGTGGTCCCCAAGTCGAT-3'). Expression data were normalized to the

housekeeping gene *Gusb* (FW: 5'-CCACCAGGGACCATCCAAT-3'; RV: 5'-

AGTCAAAATATGTGTTCTGGACAAAGTAA-3').

Quantification of circulating hormones

CPP and GCG levels in the plasma were quantified by Copeptin ELISA (MyBioSource, Cat#

MBS2020621) and Glucagon ELISA (Mercodia, Cat# 10-1281-01), respectively, based on the

manufacturer's instructions.

Constructs / Cloning / AAV packaging

The p2.0VPI.icre (AVP-icre) plasmid was generated by subcloning the iCre sequence (AgeI

2732nt, NotI 3813nt) from pCDH-CB-iCre plasmid (Addgene # 72257) in the p2.0VPI.EGFP

backbone (AgeI 6237nt, NotI 6972nt; Addgene # 40868) (see Fig S1). The packaging of the

plasmid in the AAV6 was performed by the University of North Carolina (Chapel Hill, NC) vector

core.

The TM117-GFP plasmid was provided by the Van der Goot group (EPFL, Lausanne; Bürgi *et*

al., 2016).

eQTL analysis

eQTL mapping was performed using the R package R/qtl (Broman *et al.*, 2003) with a genotype

map from the BXD panel composed of GeneNetwork genotypes (www.genenetwork.org) merged

with available RNA sequencing data from whole hypothalamic tissue (Picard *et al.*, 2016). eQTL

interval mapping was calculated using the expected-maximization algorithm, a 5% genotyping

error rate, and pseudomarkers were generated every cM. eQTL location was obtained by 6.915

likelihood ratio statistics (LRS) support intervals. Significant eQTLs were determined for the trait

using a 5% false discovery rate threshold estimated from 1000 permutations.

QUANTIFICATION AND STATISTICAL ANALYSIS

All graphs and statistical analysis were generated using Prism software (GraphPad Prism 9),

further details regarding sample size and the statistical analysis used in each case can be found

in the corresponding figure legends. In brief, for experiments concerning comparisons between

two groups on a single independent variable (Fig 2H-J; Fig 3C,F; Fig 4G$^\Delta$; Fig 5A,B,D-F,I; Fig

6C,DFig S3D,E) we used unpaired two-tailed t-test. For experiments concerning comparisons

between two dependent measurements (Fig 4G,H) we used paired two-tailed t-test. For those

concerning comparisons among three groups on a single independent variable (Fig 5M; Fig

S3C,F) we used 1-way ANOVA with Tukey post-hoc test. For two-factor designs concerning

repeated measures (Fig 2B-G; Fig 5K; Fig 6E-G) we used 2-way ANOVA RM with Bonferroni

post-hoc test. For two-factor designs concerning non repeated measures (Fig 4C) we used 2-

way ANOVA with Tukey post-hoc test. For the comparison of neuronal subpopulation

proportion (Fig 4E) we used Fisher's exact test. In all graphs error bars are depicting ±SEM.

Acknowledgements

We thank Wanda Dolci for the excellent technical support. We thank Prof. G Van der Goot for

kindly providing the TM117-GFP plasmid. We thank ZEISS and the EPFL Bioimaging and Optics

Platform for the access to the ELYRA7 SIM^2 super-resolution microscopy apparatus. This work

was supported by a European Research Council Advanced Grant (INTEGRATE, No. 694798)

and a Swiss National Science Foundation grant (310030-182496) to BT, and has received funding

from the Innovative Medicines Initiative 2 Joint Undertaking (JU) under grant agreement No

777460 (HypoRESOLVE). The JU receives support from the European Union's Horizon 2020

research and innovation program and EFPIA and T1D Exchange, JDRF, International Diabetes

Federation (IDF), The Leona M. and Harry B. Helmsley Charitable Trust.

Author Contributions

SG: conceptualization, methodology, investigation, formal analysis, visualization, project

administration, writing-original draft, editing manuscript. **GL:** methodology, investigation, formal

analysis, visualization, editing manuscript. **AP:** formal analysis, editing manuscript. **XB:**

methodology, formal analysis. **ARSA:** software, visualization, editing manuscript. **BT:**

conceptualization, supervision, project administration, resources, visualization, writing-original

draft, editing manuscript, funding acquisition.

Declaration of Interest

The authors have no conflict of interest to declare.

References

Baylis, P.H. and Robertson, G.L. (1980) 'Rat vasopressin response to insulin-induced
hypoglycemia', *Endocrinology*, 107(6), pp. 1975–1979. Available at:
https://doi.org/10.1210/endo-107-6-1975.

Baylis, P.H., Zerbe, R.L. and Robertson, G.L. (1981) 'Arginine vasopressin response to insulin-
induced hypoglycemia in man', *The Journal of Clinical Endocrinology and Metabolism*, 53(5),
pp. 935–940. Available at: https://doi.org/10.1210/jcem-53-5-935.

Beall, C., Ashford, M.L. and McCrimmon, R.J. (2012) 'The physiology and pathophysiology of
the neural control of the counterregulatory response', *American Journal of Physiology-*

Regulatory, Integrative and Comparative Physiology, 302(2), pp. R215–R223. Available at:
https://doi.org/10.1152/ajpregu.00531.2011.

Bisgaard Bengtsen, M. and Møller, N. (2021) 'Mini-review: Glucagon responses in type 1
diabetes – a matter of complexity', *Physiological Reports*, 9(16), p. e15009. Available at:
https://doi.org/10.14814/phy2.15009.

Briski, K.P. *et al.* (2017) 'Sex Differences and Role of Estradiol in Hypoglycemia-Associated
Counter-Regulation', in F. Mauvais-Jarvis (ed.) *Sex and Gender Factors Affecting Metabolic*
Homeostasis, Diabetes and Obesity. Cham: Springer International Publishing (Advances in
Experimental Medicine and Biology), pp. 359–383. Available at: https://doi.org/10.1007/978-3-
319-70178-3_17.

Broman, K.W. *et al.* (2003) 'R/qtl: QTL mapping in experimental crosses', *Bioinformatics*, 19(7),
pp. 889–890. Available at: https://doi.org/10.1093/bioinformatics/btg112.

Bürgi, J. *et al.* (2016) 'Intrinsic Disorder in Transmembrane Proteins: Roles in Signaling and
Topology Prediction', *PloS One*, 11(7), p. e0158594. Available at:
https://doi.org/10.1371/journal.pone.0158594.

Cheng, A.H., Fung, S.W. and Cheng, H.-Y.M. (2019) 'Limitations of the Avp-IRES2-Cre (JAX
#023530) and Vip-IRES-Cre (JAX #010908) Models for Chronobiological Investigations',
Journal of Biological Rhythms, 34(6), pp. 634–644. Available at:
https://doi.org/10.1177/0748730419871184.

Chiodera, P. *et al.* (1992) 'Hypoglycemia-induced arginine vasopressin and oxytocin release is
mediated by glucoreceptors located inside the blood-brain barrier', *Neuroendocrinology*, 55(6),
pp. 655–659. Available at: https://doi.org/10.1159/000126185.

Cryer, P.E. (2012) 'Minireview: Glucagon in the Pathogenesis of Hypoglycemia and
Hyperglycemia in Diabetes', *Endocrinology*, 153(3), pp. 1039–1048. Available at:
https://doi.org/10.1210/en.2011-1499.

Dunning, B.E., Moltz, J.H. and Fawcett, C.P. (1984) 'Modulation of insulin and glucagon
secretion from the perfused rat pancreas by the neurohypophysial hormones and by desamino-
D-arginine vasopressin (DDAVP)', *Peptides*, 5(5), pp. 871–875. Available at:
https://doi.org/10.1016/0196-9781(84)90109-8.

Gao, Z.Y., Gérard, M. and Henquin, J.C. (1992) 'Glucose- and concentration-dependence of
vasopressin-induced hormone release by mouse pancreatic islets', *Regulatory Peptides*, 38(1),
pp. 89–98. Available at: https://doi.org/10.1016/0167-0115(92)90075-6.

Garfield, A.S. *et al.* (2014) 'A Parabrachial-Hypothalamic Cholecystokinin Neurocircuit Controls
Counterregulatory Responses to Hypoglycemia', *Cell Metabolism*, 20(6), pp. 1030–1037.
Available at: https://doi.org/10.1016/j.cmet.2014.11.006.

Giorgi, C. *et al.* (2018) 'Calcium Dynamics as a Machine for Decoding Signals', *Trends in Cell*
Biology, 28(4), pp. 258–273. Available at: https://doi.org/10.1016/j.tcb.2018.01.002.

Godsland, I.F. (2005) 'Oestrogens and insulin secretion', *Diabetologia*, 48(11), pp. 2213–2220.
Available at: https://doi.org/10.1007/s00125-005-1930-0.

Iurlaro, R. and Muñoz-Pinedo, C. (2016) 'Cell death induced by endoplasmic reticulum stress',
The FEBS Journal, 283(14), pp. 2640–2652. Available at: https://doi.org/10.1111/febs.13598.

Jumper, J. *et al.* (2021) 'Highly accurate protein structure prediction with AlphaFold', *Nature*,
596(7873), pp. 583–589. Available at: https://doi.org/10.1038/s41586-021-03819-2.

Kim, A. *et al.* (2021) 'Arginine-vasopressin mediates counter-regulatory glucagon release and is
diminished in type 1 diabetes', *eLife*, 10, p. e72919. Available at:
https://doi.org/10.7554/eLife.72919.

Labouèbe, G., Thorens, B. and Lamy, C. (2018) 'GLUT2-Expressing Neurons as Glucose
Sensors in the Brain: Electrophysiological Analysis', *Methods in Molecular Biology (Clifton,*
N.J.), 1713, pp. 255–267. Available at: https://doi.org/10.1007/978-1-4939-7507-5_19.

Lamy, C.M. *et al.* (2014) 'Hypoglycemia-activated GLUT2 neurons of the nucleus tractus
solitarius stimulate vagal activity and glucagon secretion', *Cell Metabolism*, 19(3), pp. 527–538.
Available at: https://doi.org/10.1016/j.cmet.2014.02.003.

Liu, L. *et al.* (2021) 'Gq signaling in α cells is critical for maintaining euglycemia', *JCI insight*,
6(24), p. e152852. Available at: https://doi.org/10.1172/jci.insight.152852.

Mårtensson, U.E.A. *et al.* (2009) 'Deletion of the G protein-coupled receptor 30 impairs glucose
tolerance, reduces bone growth, increases blood pressure, and eliminates estradiol-stimulated
insulin release in female mice', *Endocrinology*, 150(2), pp. 687–698. Available at:
https://doi.org/10.1210/en.2008-0623.

Marty, N., Dallaporta, M. and Thorens, B. (2007) 'Brain glucose sensing, counterregulation, and
energy homeostasis', *Physiology (Bethesda, Md.)*, 22, pp. 241–251. Available at:
https://doi.org/10.1152/physiol.00010.2007.

McLean, A.C. *et al.* (2012) 'Performing Vaginal Lavage, Crystal Violet Staining, and Vaginal
Cytological Evaluation for Mouse Estrous Cycle Staging Identification', *JoVE (Journal of*
Visualized Experiments), (67), p. e4389. Available at: https://doi.org/10.3791/4389.

Meek, T.H. *et al.* (2016) 'Functional identification of a neurocircuit regulating blood glucose',
Proceedings of the National Academy of Sciences, 113(14), pp. E2073–E2082. Available at:
https://doi.org/10.1073/pnas.1521160113.

Picard, A. *et al.* (2016) 'A Genetic Screen Identifies Hypothalamic Fgf15 as a Regulator of
Glucagon Secretion', *Cell Reports*, 17(7), pp. 1795–1806. Available at:
https://doi.org/10.1016/j.celrep.2016.10.041.

Picard, A. *et al.* (2021) 'Fgf15 Neurons of the Dorsomedial Hypothalamus Control Glucagon
Secretion and Hepatic Gluconeogenesis', *Diabetes*, 70(7), pp. 1443–1457. Available at:
https://doi.org/10.2337/db20-1121.

Picard, A. *et al.* (2022) 'Hypothalamic Irak4 is a genetically controlled regulator of hypoglycemia-
induced glucagon secretion', *Molecular Metabolism*, 61, p. 101479. Available at:
https://doi.org/10.1016/j.molmet.2022.101479.

Ponzio, T.A. *et al.* (2012) 'Cell-type specific expression of the vasopressin gene analyzed by
AAV mediated gene delivery of promoter deletion constructs into the rat SON in vivo', *PloS One*,
7(11), p. e48860. Available at: https://doi.org/10.1371/journal.pone.0048860.

Ramnanan, C.J. *et al.* (2011) 'Physiologic action of glucagon on liver glucose metabolism',
Diabetes, Obesity and Metabolism, 13(s1), pp. 118–125. Available at:
https://doi.org/10.1111/j.1463-1326.2011.01454.x.

Salegio, E.A. *et al.* (2013) 'Axonal transport of adeno-associated viral vectors is serotype-
dependent', *Gene Therapy*, 20(3), pp. 348–352. Available at: https://doi.org/10.1038/gt.2012.27.

Schnütgen, F. *et al.* (2003) 'A directional strategy for monitoring Cre-mediated recombination at
the cellular level in the mouse', *Nature Biotechnology*, 21(5), pp. 562–565. Available at:
https://doi.org/10.1038/nbt811.

Siafarikas, A. *et al.* (2012) 'Early loss of the glucagon response to hypoglycemia in adolescents
with type 1 diabetes', *Diabetes Care*, 35(8), pp. 1757–1762. Available at:
https://doi.org/10.2337/dc11-2010.

Sladek, C.D. and Somponpun, S.J. (2008) 'Estrogen Receptors: Their Roles in Regulation of
Vasopressin Release for Maintenance of Fluid and Electrolyte Homeostasis', *Frontiers in*
neuroendocrinology, 29(1), pp. 114–127. Available at:
https://doi.org/10.1016/j.yfrne.2007.08.005.

Spruce, B.A. *et al.* (1985) 'The effect of vasopressin infusion on glucose metabolism in man',
Clinical Endocrinology, 22(4), pp. 463–468. Available at: https://doi.org/10.1111/j.1365-
2265.1985.tb00145.x.

Stanley, S., Moheet, A. and Seaquist, E.R. (2019) 'Central Mechanisms of Glucose Sensing and
Counterregulation in Defense of Hypoglycemia', *Endocrine Reviews*, 40(3), pp. 768–788.
Available at: https://doi.org/10.1210/er.2018-00226.

Stanley, S.A. *et al.* (2016) 'Bidirectional electromagnetic control of the hypothalamus regulates
feeding and metabolism', *Nature*, 531(7596), pp. 647–650. Available at:
https://doi.org/10.1038/nature17183.

Steinbusch, L.K.M. *et al.* (2016) 'Sex-Specific Control of Fat Mass and Counterregulation by
Hypothalamic Glucokinase', *Diabetes*, 65(10), pp. 2920–2931. Available at:
https://doi.org/10.2337/db15-1514.

St-Louis, R. *et al.* (2012) 'Reactive oxygen species are required for the hypothalamic
osmoregulatory response', *Endocrinology*, 153(3), pp. 1317–1329. Available at:
https://doi.org/10.1210/en.2011-1350.

St-Louis, R. *et al.* (2014) 'Reactive oxygen species are physiological mediators of the
noradrenergic signaling pathway in the mouse supraoptic nucleus', *Free Radical Biology &*
Medicine, 71, pp. 231–239. Available at: https://doi.org/10.1016/j.freeradbiomed.2014.03.024.

Strembitska, A. *et al.* (2022) 'Lipid biosynthesis enzyme Agpat5 in AgRP-neurons is required for
insulin-induced hypoglycemia sensing and glucagon secretion', *Nature Communications*, 13(1),
p. 5761. Available at: https://doi.org/10.1038/s41467-022-33484-6.

Tamaki, T. *et al.* (2017) 'A novel transmembrane protein defines the endoplasmic reticulum
stress-induced cell death pathway', *Biochemical and Biophysical Research Communications*,
486(1), pp. 149–155. Available at: https://doi.org/10.1016/j.bbrc.2017.03.017.

Tesfaye, N. and Seaquist, E.R. (2010) 'Neuroendocrine Responses to Hypoglycemia', *Annals of
the New York Academy of Sciences*, 1212, pp. 12–28. Available at:
https://doi.org/10.1111/j.1749-6632.2010.05820.x.

Thorens, B. (2022) 'Neuronal regulation of glucagon secretion and gluconeogenesis', *Journal of
Diabetes Investigation*, n/a(n/a). Available at: https://doi.org/10.1111/jdi.13745.

Varadi, M. *et al.* (2022) 'AlphaFold Protein Structure Database: massively expanding the
structural coverage of protein-sequence space with high-accuracy models', *Nucleic Acids
Research*, 50(D1), pp. D439–D444. Available at: https://doi.org/10.1093/nar/gkab1061.

Yibchok-anun, S. *et al.* (2004) 'The role of arginine vasopressin in diabetes-associated increase
in glucagon secretion', *Regulatory Peptides*, 122(3), pp. 157–162. Available at:
https://doi.org/10.1016/j.regpep.2004.06.010.

Zhivotovsky, B. and Orrenius, S. (2011) 'Calcium and cell death mechanisms: A perspective
from the cell death community', *Cell Calcium*, 50(3), pp. 211–221. Available at:
https://doi.org/10.1016/j.ceca.2011.03.003.