Title: The NQR pathway regulates the immunomodulatory function of *Bacteroides thetaiotaomicron*

Morgan J. Engelhart[1,2,3], Robert W. P. Glowacki[1,3], Clifford V. Harding[4], Eric C. Martens[5], Philip P. Ahern[1,2,3*]

[1]Department of Cardiovascular and Metabolic Sciences, Lerner Research Institute, Cleveland Clinic, Cleveland, OH 44195, USA;
[2]Department of Molecular Medicine, Lerner College of Medicine, Case Western Reserve University, Cleveland, OH;
[3]Center for Microbiome and Human Health, Lerner Research Institute, Cleveland Clinic, Cleveland, OH 44195, USA;
[4]Department of Pathology, Case Western Reserve University/University Hospitals Cleveland Medical Center, Cleveland, Ohio, USA;
[5]Department of Microbiology and Immunology, University of Michigan Medical School, Ann Arbor, MI 48109, USA.
*Correspondence: ahernp@ccf.org

Abstract

The gut microbiome and intestinal immune system are engaged in a dynamic interplay that provides myriad benefits to host health. However, the microbiome also has the capacity to elicit damaging inflammatory responses, and thus the establishment of harmonious immune-microbiome interactions is essential to maintain homeostasis. Gut microbes actively coordinate the induction of anti-inflammatory immune responses that establish and sustain these mutualistic interactions. Despite this, the microbial pathways that govern this dialogue remain poorly understood. We investigated the mechanisms through which the gut symbiont *Bacteroides thetaiotaomicron* (*B. theta*), a known modulator of intestinal immune function, exerts its immunomodulatory functions. Our data reveal that *B. theta* stimulates production of the cytokine IL-10 in a Toll-like receptor 2 dependent manner, independent of its known capsular polysaccharides. This process depends on an intact NQR pathway, which regenerates NAD+ during respiration. Deletion of *nqr* genes significantly reduces the capacity of *B. theta* to induce IL-10 by impairing biogenesis of outer membrane vesicles. These data identify a microbial pathway that may be targeted to intentionally manipulate the immunomodulatory capacity of the microbiome.

Introduction

The gastrointestinal tract is densely colonized with a complex array of bacteria, fungi, archaea, and viruses collectively referred to as the gut microbiome[1-3]. The intestine is also home to a multifaceted immune network that is separated from the gut microbiome by an overlying epithelial and mucus layer[4,5]. Despite this physical barrier, the intestinal immune system is in continual contact with the microbiome. Given the important contribution of the microbiome to maximizing host health, it is essential that the host maintains harmonious relations with these microbial communities to sustain a mutualistic state, while remaining vigilant for potential infectious agents. To this end, the intestinal immune system is enriched with cell types and molecules that prevent unwarranted deleterious inflammatory responses, and limit the penetration of the microbiome into host tissues[6]. This includes a large population of CD4+ regulatory T cells (Tregs)[7-14], and expression of cytokines such as IL-10[8,15] and transforming growth factor-beta (TGF-β)[16], potent anti-inflammatory mediators. The essential nature of these responses is highlighted by the deleterious impact of defects in these pathways, which leads to the development of tissue-destructive, microbiome-targeted pro-inflammatory immune responses, as occurs in inflammatory bowel disease[17-27]. However, the molecular factors that drive the accumulation/production of these tolerance inducing factors, and the microbial pathways involved, remain poorly understood.

A large body of evidence has established that specific gut microbes actively coordinate the activation of host anti-inflammatory responses. These include, bacteria such as *Helicobacter hepaticus*[15,28,29], *Bacteroides fragilis*[8,30-33], *Bacteroides thetaiotaomicron* (*B. theta* hereafter) [7,34,35], *Bifidobacterium bifidum*[10], members of the altered Schaedler flora (ASF)[13] and collections of *Clostridium* species[9,14,36] among others, that have been

shown to induce the accumulation of colonic Tregs and/or production of the anti-inflammatory cytokine IL-10. Immunomodulatory microbiome-derived products identified to date have fallen into two main categories (i) products of microbial metabolism like short-chain fatty acids or microbial-modified bile acids[36-41] and (ii) cellular components like capsular or cell-surface polysaccharides[8,10,15,31,32] that are sensed by host pattern recognition receptors (PRRs)[10,15,32,33,42]. Although our knowledge of the precise pathways through which the host is exposed to these molecules is limited, outer membrane vesicles (OMVs), spherical structures derived from the outer membrane of many Gram-negative bacteria[33,43-47], are posited to be packaged with proteins, lipids, and polysaccharides, and delivered to the intestinal immune system to shape the local immunological milieu. Indeed, members of the *Bacteroides*, a predominant genera within the gut microbiome[48-50], including *B. fragilis*[33] and *B. theta*[43,51], produce OMVs that contain immunomodulatory factors that can invoke beneficial immune response that limit inflammation, *e.g.* OMVs from *B. fragilis* contain polysaccharide A (PSA) that elicits IL-10 production[33].

Given their enormous potential to shape immune responses, gut microbes and/or their derived products have the potential to act as novel therapeutics for inflammatory diseases by favoring the induction of anti-inflammatory responses. Despite this promise, and notwithstanding some key advances, our understanding of these molecules, and how they are delivered to the host, remains limited. As our knowledge of the host-microbiome dialogue that promotes mutualism grows, there is an increasing need to understand the microbial pathways that govern the production and secretion of these factors. Defining such pathways will provide an opportunity to potentially expand production of molecules of interest, or heterologously reconstruct these pathways in synthetic probiotics to maximize beneficial effects[52].

We have leveraged the genetically tractable gut bacterium, *B. theta*, a prominent microbiome member that we and others[7,34,35,41] have shown to induce anti-inflammatory responses, to understand the microbial pathways that impact gut bacterial interaction with the immune system. Our data have identified that *B. theta* induces the anti-inflammatory cytokine IL-10 in a TLR2 dependent but capsular polysaccharide independent manner. In addition, we have identified that the Na[+]-translocating <u>N</u>ADH:ubi<u>q</u>uinone oxido<u>r</u>eductase (*nqr*) locus[53-55]—encoded by genes *BT1155-BT1160*—is required for the optimal induction of IL-10. We show that disruption of the *nqr* operon impairs OMV biogenesis and that this altered OMV generation is responsible for the reduced IL-10 induction. Our results provide a formal demonstration that couples OMV production with the induction of IL-10 by members of the microbiome, and offer new insights into the pathways that regulate the generation of OMVs. Thus, the NQR complex represents a potential microbiome target that could alter host-microbe interactions and favor the promotion of an anti-inflammatory intestinal environment.

Results

B. thetaiotaomicron secretes immunomodulatory factors that are sensed via TLR2 in a capsular-polysaccharide independent manner.

B. thetaiotaomicron (*B. theta*) has been established as a potent immunomodulatory microbiome member[7,34,35,43,51]. We sought to define the pathways that underlie the capacity of *B. theta* to mediate such effects. Several bacterial members of the microbiome have been shown to stimulate the production of IL-10 via secreted factors[10,15,31,33]. To determine if *B. theta* also stimulates immune responses via secreted factors, we treated unfractionated splenocytes from wild-type mice with either heat-inactivated *B. theta* (strain VPI-5482) cell pellets (OD_{600}=1.0 per 5×10^5 splenocytes), conditioned media (sterile-filtered TYG media conditioned by growth of *B. theta*; 1% v/v), or sterile TYG bacterial culture media for two days, and assessed the production of the cytokine IL-10 in culture supernatants by ELISA. Both bacterial pellets and conditioned media stimulated significant production of IL-10, demonstrating that *B. theta* drives production of IL-10 and that this can be mediated by factors that are secreted/released from the cell (Figure 1A). Furthermore, these findings were replicated using murine bone-marrow derived macrophages (BMDMs) suggesting that this response is not limited to splenocytes (Supplemental Figure 1A). As with many microbes/microbial products, both cell pellet and conditioned media also induced significant production of IL-6 (Supplemental Figure 1B), often associated with inflammation[56], but notably, IL-6 is also required for the development of bacterial-reactive anti-

inflammatory Tregs in the intestine[57,58]. Thus, IL-6 also forms part of a loop that sustains mutualism. As *B. theta* elicits Treg development and does not elicit inflammation in wild-type mice[34,43], induction of IL-10 and IL-6 likely both contribute to its anti-inflammatory properties, and thus we focused our remaining analyses on its ability to induce IL-10. Outer membrane vesicles (OMVs) are spherical vesicles that are released by many Gram-negative bacteria. *B. theta* produces OMVs that have been posited to represent the dominant process through which it interacts with the host[43]. Moreover, OMVs from the related bacterium, *B. fragilis*, have been shown to induce IL-10[33]. To determine if OMVs secreted into the conditioned media by *B. theta* can stimulate IL-10, we isolated OMVs from stationary phase cultures via ultracentrifugation, resuspended them in a volume of PBS equivalent to that from which they were prepared, and used them to stimulate splenocytes for two days (1% v/v). These OMVs also induced significant production of IL-10 relative to TYG culture media control which had been passed through the same process, although notably, the level of induction was not equivalent to that seen with unfractionated conditioned media, suggesting that additional factors are present that can prime IL-10 (Figure 1B).

B. fragilis produces a capsular polysaccharide (CPS), polysaccharide A (PSA), that can be packaged into OMVs and drives IL-10 induction in a TLR2-dependent manner[33]. To determine if there were parallels with how *B. theta* induced IL-10, we first assessed the role of TLR2 in responsiveness to *B. theta*. To do so, we stimulated wild-type or TLR2-/- splenocytes with conditioned media and assessed IL-10 in culture supernatant after two days. As with *B. fragilis*, TLR2 was required for responsiveness to *B. theta* as TLR2-/- splenocytes produced significantly less IL-10 than wild-type splenocytes in response to *B. theta* conditioned media (Figure 1C), despite mounting a robust response to stimulation with LPS (data not shown). Stimulation of BMDMs generated from the bone-marrow of wild-type or TLR2-/- mice, also showed a role for TLR2 in producing IL-10 in response to *B. theta* (Supplemental Figure 1C), confirming that the requirement for TLR2 was not limited to the splenocyte assay. Conditioned media from an additional isolate of *B. theta* (strain 0940-1) also induced IL-10 in a TLR2 dependent manner (Supplemental Figure 1D), suggesting that TLR2 is a sensor for *B. theta* as a species, in keeping with a recent report showing that a large panel of distinct *Bacteroides* are sensed by this PRR[59]. TLR2 can coordinate activation of the immune system in concert with the PRR Dectin-1, and the dual engagement of TLR2 and Dectin-1 by *B. fragilis* derived PSA drives IL-10 production[30,60]. However, we found that the ability of secreted factors produced by *B. theta* to induce IL-10 is not dependent on Dectin-1 (Supplemental Figure 1E). *B. theta* also synthesizes CPSs that are encoded by 8 distinct *cps* loci. Stimulation of splenocytes with an acapsular mutant (Δ*tdk*Δ*cps*1-8) that lacks the ability to synthesize any of these capsular polysaccharides (CPSs)[61], induced equivalent IL-10 to the parent strain (note, the parent strain is represented by *B. theta* 5482 on the Δ*tdk* background that is used to construct mutants) (Figure 1D), in keeping with a report that *B. theta* CPSs act to mask it from the immune system rather than stimulate immune responses[62]. Thus, *B. theta*, unlike *B. fragilis*, does not stimulate IL-10 induction via any of its known CPSs. These data suggest the existence of a novel factor in *B. theta* that induces IL-10.

Gene *BT1160* is required for *B. thetaiotaomicron* mediated IL-10 induction.

To further investigate how *B. theta* induces the production of IL-10 we aimed to identify genes that that were required for its IL-10 inducing capacity. To do so, we used an arrayed transposon mutagenesis library of 2,009 *B. theta* transposon mutants, made on the acapsular (*B. theta* Δ*tdk*Δ*cps*1-8) genetic background[35], allowing the assessment of individual mutations on IL-10 inductive capacity. Splenocytes were stimulated with the conditioned media from individually-grown *B. theta* Δ*tdk*Δ*cps*1-8 transposon mutants (1% v/v per well) for two days (note, to facilitate testing of a large batch of mutants, the conditioned media was centrifuged to remove cellular debris but was not sterile filtered). Pam3CSK4, a defined TLR2 agonist[63-65], was used to stimulate splenocytes on each plate, to allow for normalization of data and minimize the effect of plate-to-plate variation. We then assessed IL-10 induction in culture supernatants by ELISA, and normalized the values for each transposon mutant to the positive control, Pam3CSK4. In addition, we performed growth assessment of the individual mutants and 27 mutants that did not exhibit detectable growth were removed from our analysis. Several mutants displayed impaired IL-10 induction as evidenced by a lower IL-10/Pam3CSK4 ratio (Figure 2A). We chose one mutant for follow-up (mutant from well H4 of plate 19; 19_H4), and validated its impaired ability to

induce IL-10 by generating an independent batch of sterile-filtered conditioned media which confirmed its impaired ability to induce IL-10 (Figure 2B). Next we mapped the location of the transposon insertion, and found that transposon mutant 19_H4 had a disrupted, out-of-frame insertion in gene *BT1160*, which is predicted based on gene homology to encode subunit A (*nqrA*) of the Na+-transporting NADH:ubiquinone oxidoreductase (NQR complex). The NQR complex is comprised of six subunits encoded by a six gene cluster (*nqr* locus, genes *BT1155-BT1160*) (Figure 2C). The NQR complex has been characterized in other bacteria, including *B. fragilis*, and is found within the inner membrane, responsible for converting NADH to NAD$^+$ and shuttling electrons down the electron transport chain, to generate ATP, while pumping sodium from the cytosol into the periplasm to create an ion gradient and converting fumarate to succinate[53-55]. To validate the role of this gene we sought to create engineered in-frame deletions of gene *BT1160*. First, we generated a truncated knock-out of gene *BT1160* (ΔBT1160_trunc) on both the acapsular (Δ*tdk*Δ*cps*1-8) and wild-type (Δ*tdk*) backgrounds. We engineered this truncated deletion of gene *BT1160* with the goal of limiting modifications to the downstream genes of the *nqr* locus as the predicted promotor for the *nqr* locus is directly upstream of gene *BT1160*. The truncated deletion left the first 8 nucleotide bases of gene *BT1160*—including the start codon of *BT1160*—intact. However, the same reading frame encodes a stop codon prior to gene *BT1159*, therefore, downstream genes are not shifted out of frame. This deletion of gene *BT1160* resulted in significant impairment in IL-10 induction by *B. theta* compared to the parent strain, on both the wild-type and acapsular backgrounds (Figure 2D). Although we observed statistically significant impacts on growth following deletion of gene *BT1160*, the magnitude of these effects was minimal, and the mutant reached equivalent levels as wild-type in stationary phase (Supplemental Figure 2A,B; note, the ΔBT1160_trunc mutant grew significantly more slowly than parent strain directly from frozen stocks as has been reported for mutants in this pathway in *B. fragilis*[53], however, parent strain-levels of growth were observed when sub-cultured, and thus in all assays, the cultures used were subcultures from initial overnight cultures that were grown to stationary phase). The ability of *B. theta* conditioned media to induce IL-10 from BMDMs was also shown to be dependent on gene *BT1160* (Supplemental Figure 2C). Thus, our screen identified gene *BT1160* as a *bona fide* regulator of IL-10 induction in *B. theta*.

B. theta has been reported to stimulate IL-10 production from human peripheral blood mononuclear cells (PBMCs)[51]. To determine if *BT1160* played a role in the immunomodulatory function of *B. theta* on human cells, we stimulated PBMCs from two different healthy donors with the conditioned media from wild-type (Δ*tdk*) *B. theta* or the truncated *BT1160* deletion mutant and found that as with murine cells, the induction of IL-10 in human PBMCs also required *BT1160* (Figure 2E). Complementation of *BT1160 in trans* by introduction of the full-length *BT1160* gene and the promotor region into the *att2* (tRNA^{ser}) site, partially restored the ability of *B. theta* to induce IL-10, compared to wild-type (Figure 2F), with notable variance between experiments (Supplemental Figure 3A). Next, we generated a complete in-frame deletion of gene *BT1160* on the Δ*tdk* background, that also significantly diminished the ability of *B. theta* to induce IL-10 compared to the parent strain, but did not impair IL-10 induction as greatly as the truncated genetic mutant (Figure 2G). These data show that disruption of gene *BT1160* significantly impairs *B. theta* mediated IL-10 induction. However, given the differences between the two deletion mutants and the partial impact of complementation, the effects observed in the *BT1160* deletion mutants may be attributable to other genes that are impacted in these mutants, likely the other genes in the *nqr* locus.

The NQR complex coordinates IL-10 induction by *B. thetaiotaomicron*.

The NQR complex is encoded by a bacterial locus (*nqr*) made up of six genes, genes *BT1155-BT1160* (Figure 3A). Genes *BT1155-BT1159* code for subunits B-F (*nqrB-F*) in the NQR complex and are directly downstream of *BT1160* (Figure 3A). To determine if deletion of gene *BT1160* impacted expression of these downstream genes, we performed RT-qPCR to assess gene expression at mid-log phase in both the truncated and full genetic deletion mutants. These data showed that both *BT1160* mutants displayed a reduction in expression of downstream genes *BT1156-BT1159* within the *nqr* locus compared to wild-type *B. theta* (Δ*tdk*) (Figure 3B), suggesting that the reduced IL-10 induction by these mutants may be attributable, in part, to altered expression of these genes (as expected, expression of gene *BT1160* was not detectable in either mutant; not shown). To

determine if alterations in expression of other *nqr* genes could also impact IL-10 induction, we engineered *B. theta* mutants lacking either the entire *nqr* locus (ΔBT1155-BT1160) or the entire locus except *BT1160* (ΔBT1155-BT1159). Both mutants were significantly impaired in their capacity to induce IL-10 from splenocytes across a range of doses (0.5%, 1%, and 3% v/v) (Figure 3C, D and Supplemental Figure 3B). Furthermore, the induction of IL-10 by *B. theta* conditioned media, and its dependence on *nqr* genes, was observed in cells from both male and female mice, suggesting this pathway is not limited to one biological sex (Supplemental Figure 4A). Importantly, the expression of genes *BT1154* and *BT1161*, which lie directly downstream and upstream of the *nqr* locus, was unaltered by the deletion of *BT1160* and a control deletion mutant of gene *BT1161* has no discernible impact on IL-10 induction (Figure 3B, E and Supplemental Figure 3B). These data suggest that (i) disruption of the *nqr* locus impairs the ability of *B. theta* to induce IL-10, but that deletion of a gene directly upstream of the locus does not, and that (ii) *BT1160* is not sufficient for induction of IL-10, and thus the impact of *BT1160* deletion may be mediated by impacts on other genes within the *nqr* locus.

The NQR complex influences *B. thetaiotaomicron* OMV biogenesis and composition.

We next aimed to uncover how the NQR complex could impact induction of IL-10. As OMVs have been shown to be key mediators of the anti-inflammatory effect of other members of the *Bacteroides*[33], and a central mechanism through which *B. theta* interacts with the host[43], we reasoned that the loss of NQR could alter the production of the molecule(s) packaged into OMVs that drive IL-10 induction, or impair the production/secretion of OMVs into the conditioned media. To first determine the impact of gene *BT1160* on production of immunomodulatory OMVs we prepared OMVs from the truncated *BT1160* mutant and the parental strain, and assessed the ability of these preparations to stimulate IL-10. In these assays, OMVs were resuspended in an equivalent volume of PBS as that of the conditioned media from which they were isolated, *i.e.* if 1 ml of media was used to isolate OMVs, the pelleted vesicles were resuspended in 1 ml of PBS and not normalized based on OMV quantity. These experiments recapitulated what we observed with conditioned media, with OMV preparations from the *BT1160* deficient strain (*B. theta*$^{\Delta tdk \Delta BT1160\_trunc}$) showing significant impairment in IL-10 induction when compared to the parent strain (*B. theta*$^{\Delta tdk}$) (Figure 4A). OMVs isolated from the partial *nqr* locus deletion mutant (*BT1155-BT1159*) that retains *BT1160* (Figure 4B) or the entire deletion mutant (*BT1155-BT1160*), were also impaired in IL-10 induction ability compared to OMVs from the parent strain (Figure 4C). However, OMVs isolated from a mutant harboring a deletion of a gene outside the *nqr* locus (*BT1161*), did not alter the ability of *B. theta* OMVs to induce IL-10 compared to the parent strain (Figure 4D). This pattern was observed using male and female mice (Supplemental Figure 4B), suggesting it is not specific to a single biological sex. These data suggest that gene *BT1160* and the *nqr* locus influence the ability of *B. theta* to produce immunomodulatory OMVs.

To determine if loss of *BT1160* impacted the specific proteins packaged into OMVs, including lipoproteins which are abundant in *B. theta* OMVs and known TLR2 agonists, we performed proteomic assessment of OMVs isolated from the *BT1160* truncated mutant (*B. theta*$^{\Delta tdk \Delta BT1160\_trunc}$) (the truncated mutant was chosen as it was a less profound modification of the *B. theta* genome yet had large effects on IL-10 induction) or the parent strain (*B. theta*$^{\Delta tdk}$) using LC-MS (detected proteins were identified in the reference *B. theta*$^{VPI-5482}$ UniProtKB protein database using the program Andomeda). We performed two independent proteomic assessments with three replicates of wild-type and mutant in each analysis. A total of 187 total proteins were detected in each of the two experiments (i.e. they were detected in both experiments), the majority of which (87.7%) remained unchanged between the parent and mutant strain (Figure 5A).The proteins that were (i) detected, (ii) found to have a fold change ≥2 in abundance, and (iii) whose P-value for differential abundance was <0.05, were considered to represent those proteins that are robustly regulated by *BT1160*. Of the 187 proteins detected, 6.42% were found to be higher in abundance in the parent strain compared to the mutant, while 5.88% were found to be higher in abundance in the *B. theta* mutant lacking *BT1160* (Figure 5A). Those proteins whose abundance was significantly impacted by deletion of *BT1160* from the first and second proteomics assessments are shown in Figure 5B and 5C respectively. 14 of the 23 significantly altered proteins identified were also found to have significantly changed abundance between the OMVs isolated from the acapsular *B. theta* mutant lacking gene *BT1160* and the acapsular parent strain (data not shown). The

relatively low number of altered proteins prompted us to search for alternative explanations for the impaired induction of IL-10.

Interestingly, during the preparation of OMVs for proteomic analysis, we noted that the *BT1160* deficient strain produced smaller pellets than the parent strain from equivalent amounts of conditioned media, suggesting impairment in OMV biogenesis. Quantification of the OMVs by assessment of the protein concentration in OMV preparations revealed that deletion of gene *BT1160* leads to a reduction in the concentration of OMVs produced compared to the parent strain (Figure 6A). Note, due to variability in the raw OMV values we quantified OMVs from batches of wild-type and *B. theta* mutants that were processed at the same time, and normalized the value of the mutant to that of the wild-type which was set to 100%. The data presented show the pooled results of multiple individual biological replicate batches. Similar to the induction of IL-10 by conditioned media, complementation of gene *BT1160 in trans* restored OMV biogenesis to varying degrees (Figure 6A). As the specific method used for OMV quantitation can impact conclusions[66], we sought an alternative approach for assessment of OMV concentration. To this end we used a particle analyzer, ZetaView, to quantify OMVs. These data also showed impaired OMV production from the truncated *B. theta*$^{\Delta tdk \Delta BT1160\_trunc}$ mutant compared to the parent strain (Supplemental Figure 5A), validating the Qubit-based quantification data, and suggesting that the lower protein levels reflected fewer OMVs rather than a reduced protein level within the OMVs. Deletion of the remaining *nqr* locus genes except for *BT1160 (BT1155-BT1159)*, and the entire *nqr* locus (*BT1155-BT1160*) also led to a reduction in the concentration of OMVs compared to the parent strain (Figure 6B), suggesting that this pathway is impacted by the NQR complex rather than by gene *BT1160* in isolation. Importantly, the inability of *B. theta* lacking gene *BT1160* to induce IL-10 is not from a total loss of OMV biogenesis. ZetaView analysis and transmission electron microscopy (TEM) both revealed that *B. theta* still produces OMVs when gene *BT1160* is deleted (Supplemental Figure 5B), highlighting that these data reflect a quantitative reduction in OMVs rather than a complete loss of OMV production. Collectively, these data suggest that the NQR complex plays an important role in OMV biogenesis in *B. theta*.

NQR-dependent OMV production is directly linked to the ability of *B. thetaiotaomicron* to induce IL-10.

Our proteomic and OMV quantification data suggests that impaired NQR function leads to slight alterations in the specific proteins packaged in OMVs as well as a reduced capacity to generate OMVs. Thus, the impaired IL-10 induction could result from a reduction in the TLR2 agonist within the OMVs that drives the IL-10 production, a reduction in OMV quantity without changes to the specific TLR2 agonist that drives IL-10 production, or a combination of these two possibilities acting synergistically. We reasoned that if the phenotype was driven primarily by impaired OMV biogenesis, then the use of equivalent OMV concentrations from wild-type and *BT1160* mutant *B. theta* should lead to a similar amount of IL-10 induction. Alternatively, if there was impaired packaging of the specific TLR2 agonist that drives IL-10 production in OMV from the *BT1160* mutant (*B. theta*$^{\Delta tdk \Delta BT1160\_trunc}$), then normalizing OMV concentrations to that of the parent strain would not restore IL-10 induction by the *BT1160* mutant strain. To test this, we stimulated splenocytes with OMV preparations from the *BT1160* mutant and parental strain as above, or OMVs from the *BT1160* deletion mutant such that they were the same concentration as the OMVs from the parent strain. These data show that normalizing the OMV concentration between the mutant and parent strain largely restored the ability of *B. theta*$^{\Delta tdk \Delta BT1160\_trunc}$ to induce IL-10 (Figure 6C). This result held true whether OMVs were normalized using Qubit protein concentration or ZetaView particles/mL (data not shown). These data suggest that gene *BT1160* plays an important role in OMV biogenesis in *B. theta* and explains, in large part, how this gene—and the *nqr* locus as a whole—impacts the induction of innate immune responses by *B. theta*. Interestingly, these data also suggest that IL-10 induction by the immune system is dependent on the quantity of OMVs carrying immunomodulatory factors, and that deletion of NQR complex genes disrupts canonical OMV biogenesis. Thus, our data establishes that not only are OMVs sufficient to elicit IL-10 production, but their optimal production is required for the immunomodulatory function of *B. theta*.

Discussion

The basis through which the immune system maintains immune tolerance has fascinated immunologists for decades, but until relatively recently, this interest has focused on the induction and maintenance of tolerance to self-antigens encoded within the host genome, or non-self as represented by organs/tissues during transplantation[6,67,68]. However, the intestine is laden with another form of non-self, the microbiome, which also necessitates the development of immune tolerance due to the myriad benefits it provides to the host[6]. A failure in immune tolerance of the microbiome can lead to development of chronic diseases like inflammatory bowel disease (IBD) that severely impact host health[18,21-23,67-69]. The intestinal immune system is therefore faced with the challenge of differentiating between potential pathogens and resident microbiome members, despite both types of microorganisms producing agonists of host-PRRs such as Toll-like receptor 4 (TLR4), TLR2, Dectin-1, etc[59,60,63].

While the "Danger Theory" provides a conceptual framework to understand how this is achieved, the molecular processes that underlie this decision remain enigmatic[70]. Several studies have established that the gut microbiome can coordinate the induction of anti-inflammatory immune responses that limit inflammation and favor mutualism via the production of a variety of metabolites and cellular products[8-11,13-16,28,30,31,33,36-41,43,51,57-59,62,71,72]. Despite this, there remains a dearth of knowledge regarding the nature of these microbiome-derived molecules and the microbial pathways that govern their production and secretion. Here, we leveraged the genetic tractability of the gut symbiont *B. theta*, a potent modulator of immune function, to identify pathways that contributed its ability to promote production of the anti-inflammatory molecule IL-10. Our data revealed that secreted factors from *B. theta* promoted IL-10 production via a TLR2-dependent but capsular polysaccharide independent mechanism. Furthermore, we uncovered that the gene, *BT1160*, and the *nqr* locus in which it resides (predicted to encode a the Na+-transporting NADH:ubiquinone oxidoreductase (NQR) complex)[53-55], played a central role in the ability of *B. theta*-derived conditioned media to drive the production of IL-10. We further show that the impairment in IL-10 induction was linked to the ability to produce outer membrane vesicles (OMVs), reinforcing the notion that OMVs play important roles as mediators of symbiosis.

The existence of a microbiome-TLR2-IL-10 axis has emerged due to the work of several groups. This is exemplified by *B.* fragilis which produces the capsular polysaccharide, polysaccharide A (PSA), which is packaged into OMVs and drives IL-10 production via TLR2[8,30-33,42]. Mirroring these findings, *Bifidobacterium bifidum*[10] and *Helicobacter hepaticus*[15] produce cellular polysaccharides that stimulate IL-10 production via TLR2, and members of the Bacteroidetes as a whole have recently been shown to promote IL-10 induction through a TLR2-MyD88 axis[59]. While our data reveal that *B. theta* secreted factors also stimulated IL-10 production through a TLR2-dependent mechanism, we found no requirement for any of the known capsular polysaccharides produced by *B. theta* in this process. Although it is possible that additional, as yet unindentified, cellular polysaccharides are involved in this process, this suggests that *B. theta* shapes immune function by a pathway distinct from *B. fragilis*. Notably, it has recently emerged that chemically synthesized PSA lacks IL-10 inductive capacity, by contrast with PSA isolated from *B. fragilis*, and instead, a small lipid covalently attached to PSA is critical for its immunomodulatory function[30], in accordance with structural studies of TLR2-based recognition of synthetic agonists that show that TLR2 directly binds lipids[65,73]. In keeping with the lack of a role for capsular polysaccharides, we did not detect a role for Dectin-1 in promoting IL-10 production, which contrasts with *B. fragilis* whereby dual engagement of TLR2 by lipid and Dectin-1 by PSA stimulate IL-10[30]. Neither *H. hepaticus* nor *B. bifidum* require sensing by Dectin-1 for their immunomodulatory capacity[10,15]. It is therefore unclear if there are as yet unidentified lipids/lipid moieties that are required for the immunomodulatory function of *H. hepaticus* or *B. bifidum* derived polysaccharides, and further investigation is required to identify the specific factor from *B. theta* that stimulates IL-10 induction. Given the lack of molecularly defined factors that drive IL-10, it is as yet unknown if TLR2 represents a common host sensor for a wide variety of microbiome members through recognition of distinct molecules, or if there is a common structurally similar lipid produced across a range of microbes that promotes IL-10 induction. Moreover, whether specific chemical features of gut microbe-derived molecules are sensed to allow induction of tolerance, or other facets of the initial recognition event are responsible for the observed responses is unknown and warrants intensive study.

A number of studies have shown OMVs to be sufficient to mediate the immunologic function of gut bacteria[33,35,43,46,51,74]. Our data now provide evidence that OMV production is directly linked to immunomodulation and that impaired OMV production can reduce the capacity to shape immune function. Our data identified the existence of a pathway, namely the NQR pathway, which is required for optimal production of OMVs, and demonstrated that this OMV biogenesis could be directly linked to immunomodulatory capacity. Despite the attention that has been focused on OMVs[47,75,76], there is still a dearth of information regarding the molecular pathways that coordinate their biogenesis. Seminal studies on *B. theta* OMVs have revealed that they are composed of phospholipids, lipooligosaccharide (LOS), outer membrane and periplasmic proteins, sphingolipids, and lipoproteins[74,77]. These OMVs contain distinct protein profiles from the *B. theta* outer membrane and differ from other microbes in that they are highly enriched with lipoproteins[74,78-80]. The OMV-enriched lipoproteins contain a lipoprotein export sequence believed to coordinate packaging into OMVs, suggesting the existence of an ordered process[74]. Our studies extend on these and identify a locus that sustains the optimal production of OMVs. The *nqr* locus supports energy production during respiration, and in doing so, shuttles Na+ across the membrane to create Na+ gradients[53-55]. While we currently do not understand which of these functions, if either, are involved in OMV biogenesis, it is tempting to speculate that energetic requirements of OMV generation necessitate optimal ATP generation, and hence a functional NQR pathway is essential.

Collectively, our studies have identified a novel regulator of the immunomodulatory function of *B. theta*. The capacity to modulate OMV biogenesis has applicability to both promote tolerance and limit pathogenicity by increasing or decreasing OMV production in mutualists or pathogens respectively, but this will require detailed understanding of how NQR contributes to this process. Notably, antimicrobial compounds such as natural quinone-like inhibitors—HQNO and korormicin—target the NQR complex by binding to NqrB and preventing Na+-dependent reduction of quinones[81]. These compounds effectively target and kill pathogenic species, like *Vibrio cholerae* and *Pseudomonas aeruginosa*, yet gut commensals like *B. fragilis* and *B. theta* remain unharmed[81]. Thus, the selective inhibition or stimulation of the NQR pathway may allow rational manipulation of microbiome function to enhance anti-inflammatory function, or boost the delivery of molecules of interest via OMVs. Further insight into how biogenesis is coordinated, and defining the specific immunomodulatory factors they harbor is essential to gain a more complete understanding of how gut microbes like *B. theta* coordinate host immune responses, and will provide an opportunity to identify pathways that can be manipulated to shape intestinal immune responses.

Acknowledgments

We are grateful to Anagha Kadam, Jessica Till, Orion Brock, Adeline Hajjar, and Jan Claesen for their intellectual input throughout these studies, and their advice during preparation of the manuscript; Meng Wu for her assistance with data analysis from the transposon mutagenesis screen; Belinda Willard for proteomic analysis; and William Massey for his assistance with generation of bone-marrow derived macrophages. In addition, we would like to thank the following for their provision of mice used in these studies: Nancy Nagy, Case Western Reserve University (TLR2$^{-/-}$), and Luis Ramirez and Tanya Freedman, University of Minnesota (wild-type and Dectin-1$^{-/-}$). OMV proteomics was made possible by the Fusion Lumos LC-MS instrument purchased with a National Institutes of Health (NIH) Shared Instrument Grant: 1S10OD023436-01. This work is supported by R01DK126772 from the National Institute of Diabetes and Digestive and Kidney Diseases, NIH. C.V.H. acknowledges support by R01AI034343 from the National Institute of Allergy and Infectious Diseases, NIH. E.C.M. acknowledges support by R01DK118024 from the National Institute of Diabetes and Digestive and Kidney Diseases, NIH.

Author Contributions

M.J.E. designed and executed experiments to test the role of various forms of *B. theta* to induce IL-10, identified transposon mutants with impaired IL-10 induction capacity, generated mutant forms of *B. theta*,

analyzed data and helped write the manuscript. R.W.P.G. generated and grew the transposon mutagenesis library used in this study, generated mutant forms of *B. theta*, and designed experiments to test the role of the *nqr* locus, C.V.H. provided TLR2$^{-/-}$ mice and guidance on their use in these studies. E.C.M. supervised the development of the transposon mutagenesis library, and provided the library and acpasular mutant of *B. theta* for use in our studies, in addition to intellectual input for data interpretations. P.P.A. designed experiments described within, supervised the study, and helped write the manuscript.

Declaration of Interests

P.P.A. serves as a consultant to Novome Biotechnologies, a company that genetically engineers microbes to act as therapeutic agents.

Figure 1. *B. theta* secretes immunomodulatory factors that are sensed via TLR2 in a CPS independent manner.

Figure 1. *B. theta* secretes immunomodulatory factors that are sensed via TLR2 in a CPS independent manner.

(A) Unfractionated WT splenocytes were stimulated with *B. theta*[VPI-5482] heat-inactivated pellet (OD =1.0) or *B. theta*[VPI-5482] conditioned media (1% v/v), or appropriate media control for 2 days and IL-10 was assessed in the supernatant by ELISA.
(B) WT splenocytes were stimulated with *B. theta*[VPI-5482] untreated conditioned media or TYG media (1% v/v), or isolated OMVs from the conditioned media or TYG control media that underwent the same process as that for OMV isolation and was resuspended in PBS (1% v/v). Samples were incubated for 2 days and IL-10 was assessed in the supernatant by ELISA.
(C) WT and TLR2[-/-] splenocytes were stimulated with *B. theta*[VPI-5482] conditioned media or TYG media control (1% v/v) for 2 days and IL-10 was assessed in the supernatant by ELISA.
(D) WT splenocytes were stimulated with varying doses of *B. theta* (Δ*tdk*) or acapsular *B. theta* (Δ*tdk*Δ*cps*1-8) conditioned media or TYG control media for 2 days and IL-10 was assessed in the supernatant by ELISA.

Data points represent an independent technical replicate and horizontal lines show the mean **(A-C)**, or data points represent the mean and error bars represent the standard deviation of three technical replicates **(D)**. Graphs are representative of 3 experiments **(A-C)**, wild-type versus acapsular conditioned media dose curve represents a single experiment, while the 1% v/v dose is representative of 4 experiments **(D)**. Statistical significance was determined using Student's T test: **(A, B)**, Two-way ANOVA with Sidak's post hoc test, comparisons to WT **(C)**, and Student's T test on the area under the curve: **(D)**. ns (not significant) P≥0.05; **P≤0.01; ***P≤0.001, ****P≤0.0001.

Figure 2. Gene BT1160 is required for *B. theta* mediated induction of IL-10.

Figure 2. Gene *BT1160* is required for *B. theta* mediated induction of IL-10.

(A) WT splenocytes were stimulated with conditioned media from a library of 2,009 individual *B. theta*$^{\Delta tdk \Delta cps1\text{-}8}$ transposon mutants (1% v/v) or Pam3CSK4 for 2 days and IL-10 levels in culture supernatants were measured by ELISA. For each plate, IL-10 levels were normalized to the IL-10 level induced by the positive control Pam3CSK4. Data shows the Log2 transformed value normalized to the positive control Pam3CSK4.

(B) WT splenocytes were stimulated with conditioned media from *B. theta*$^{\Delta tdk \Delta cps1\text{-}8}$ or transposon mutant 19_H4 (1% v/v) for 2 days and IL-10 was assessed by ELISA.

(C) The transposon insertion in transposon mutant 19_H4 was mapped to gene *BT1160* that is predicted to encode the first subunit of the NQR complex.

(D) WT splenocytes were stimulated with conditioned media (1% v/v) from *B. theta* harboring a truncated deletion of gene *BT1160* (acapsular background, left; wild-type background, right) for 2 days, and IL-10 levels were measured by ELISA.

(E) Human PBMCs from two healthy donors were stimulated with conditioned media (3% v/v) from *B. theta* harboring a truncated deletion of gene *BT1160* in the wild-type background, parent strain (Δtdk), or TYG media for 2 days, and IL-10 levels were measured by ELISA.

(F) WT splenocytes were stimulated with conditioned media (1% v/v) from *B. theta*$^{\Delta tdk}$, *B. theta*$^{\Delta tdk \Delta BT1160\_trunc}$, or *B. theta*$^{\Delta tdk \Delta BT1160\_trunc}$ complemented with *BT1160* (*B. theta*$^{\Delta tdk \Delta BT1160\_trunc::BT1160}$) for 2 days, and IL-10 levels were measured by ELISA.

(G) WT splenocytes were stimulated with conditioned media (1% v/v) from *B. theta* harboring a truncated deletion of gene *BT1160* (red), a full deletion of gene *BT1160* (grey) (Δtdk background), or the parent strain *B. theta*$^{\Delta tdk}$ (blue) for 2 days, and IL-10 levels were measured by ELISA.

Individual data points show mean values **(A)** or individual technical replicates **(B, D-G)**, and horizontal lines represent the mean. Graphs are representative of 4 individual experiments **(B, D, F, and G)** and 1 individual experiment **(A, E)**. Statistical significance was determined by Student's T-test: **(B, D)**, and One-way ANOVA with Dunnett's post-hoc test, comparisons to *B. theta*$^{\Delta tdk}$ or between all **(E-G)**. ns (not significant) $P \geq 0.05$, *$P < 0.05$, **$P \leq 0.01$, ***$P \leq 0.001$, ****$P \leq 0.0001$.

Figure 3. *B. theta* NQR complex promotes IL-10 induction.

Figure 3. *B. theta* NQR complex promotes IL-10 induction.

(A) Schematic of the gene locus that encodes the NQR complex and flanking genes.

(B) The expression ($2^{\Delta/\Delta \text{ Ct values}}$) of *nqr* locus genes *BT1155-BT1159* and control flanking genes (*BT1154* and *BT1161*). *B. theta*$^{\Delta tdk}$, *B. theta*$^{\Delta tdk\Delta BT1160\_\text{trunc}}$, and *B. theta*$^{\Delta tdk\Delta BT1160\_\text{full}}$ were grown *in vitro* in TYG media and cells were harvested at mid-log growth, and RNA expression profiles were assessed by RT-qPCR with *B. theta* 16S rRNA as the housekeeping gene. *BT1160* was below the limit of detection in both mutants.

(C-E) WT splenocytes were stimulated with different doses of conditioned media (0.5%, 1%, 3% v/v) from *B. theta* harboring deletions of genes *BT1155-BT1159* **(C)**, or genes *BT1155-BT1160* **(D)**, and conditioned media (1% v/v) from *B. theta* harboring a deletion in gene *BT1161* **(E)**, on the *B. theta*$^{\Delta tdk}$ background for 2 days, and IL-10 levels were measured by ELISA.

Data points represent the mean of technical replicates of independent biological replicates **(B)** or independent technical replicates **(E)**, or the mean of three technical replicates **(C, D)** and horizontal bars represent the mean **(B, E)**, and error bars shown the standard deviation **(C, D)**. Graphs are representative of a single experiment **(B)** or 4 experiments **(C-E)**. Statistical significance was determined using One-way ANOVA with Dunnett's post hoc test and comparisons between all groups: **(B)**, and Student's T test **(E)** on area under the curve: **(C-D)**. ns (not significant) P≥0.05, *P<0.05, **P≤0.01, ***P≤0.001, ****P≤0.0001.

Figure 4. The *nqr* locus influences *B. theta* OMV biogenesis.

Figure 4. The *nqr* locus influences B. theta OMV biogenesis.

(A) WT splenocytes were stimulated with conditioned media (1% v/v) or OMVs isolated from the conditioned media (1% v/v of OMVs resuspended in PBS) from *B. theta*$^{\Delta tdk}$ or *B. theta*$^{\Delta tdk \Delta BT1160}$_trunc for 2 days, and IL-10 levels were measured by ELISA.

(B-D) WT splenocytes were stimulated with OMVs isolated from the conditioned media (1% v/v of OMVs resuspended in PBS) from *B. theta*$^{\Delta tdk}$ or *B. theta*$^{\Delta tdk \Delta BT1155-BT1159}$ **(B)**, *B. theta*$^{\Delta tdk}$ or *B. theta*$^{\Delta tdk \Delta BT1155-BT1160}$ **(C)**, or *B. theta*$^{\Delta tdk}$ or *B. theta*$^{\Delta tdk \Delta BT1161}$ **(D)** for 2 days, and IL-10 levels were measured by ELISA.

Data points represent independent technical replicates and horizontal bars represent the mean **(A-D)**. Graphs are representative of 4 experiments. Statistical significance was determined using Student's T test: **(A-D)**. ns (not significant) P≥0.05, *P<0.05, **P≤0.01, ***P≤0.001, ****P≤0.0001.

Figure 5. Gene *BT1160* impacts OMV protein composition.

Figure 5. Gene *BT1160* impacts OMV protein composition.

(A) OMVs were isolated from the conditioned media of *B. theta*$^{\Delta tdk}$ and *B. theta*$^{\Delta tdk \Delta BT1160\_trunc}$ and were subjected to proteomic assessment via LC-MS. The percentage of proteins that were detected in two separate OMV proteomic analyses between *B. theta*$^{\Delta tdk}$ and *B. theta*$^{\Delta tdk \Delta BT1160\_trunc}$ and identified as having no change in abundance or statistically significant (P-value ≤0.05) ≥2-fold difference in abundance between the parent and mutant strain in both analyses are shown.

(B, C) Proteins with significant (P-value ≤0.05) differential abundance and with fold changes ≥2, identified in two separate OMV proteomics analyses between *B. theta*$^{\Delta tdk}$ and *B. theta*$^{\Delta tdk \Delta BT1160\_trunc}$ are shown. Proteins identified are represented by the gene locus tag. Total of 6 replicates per group (3 per proteomic assessment), first batch (B), and second batch (C). Bar color corresponds to gene locus tag between graphs and bars represents the mean difference in abundance in OMVs from *B. theta*$^{\Delta tdk}$ relative to *B. theta*$^{\Delta tdk \Delta BT1160\_trunc}$.

Proteins shown were selected based on the ratio (Δ*tdk*/Δ*tdk*Δ*BT1160*_trunc) of their LFQ intensities (the averaged normalized protein quantities) and their P-values (determined using Student's T-test). Ns (not significant) P>0.05, significant P≤0.05.

Figure 6. OMV production is directly linked to the ability of *B. theta* to induce IL-10.

Figure 6. OMV production is directly linked to the ability of *B. theta* to induce IL-10.

(A, B) Protein concentration of biological batches of OMVs isolated from the conditioned media of *B. theta*$^{\Delta tdk}$, *B. theta*$^{\Delta tdk\Delta BT1160\_trunc}$, or *B. theta*$^{\Delta tdk\Delta BT1160\_trunc::BT1160}$ (A), or *B. theta*$^{\Delta tdk}$, *B. theta*$^{\Delta tdk\Delta BT1155-BT1159}$, or *B. theta*$^{\Delta TDK\Delta BT1155-BT1160}$ (B), as measured by Qubit. Protein concentration was normalized to the value from a biological batch prepared from the parent strain (*B. theta*$^{\Delta tdk}$) at the same time which was set to 100. **(C)** WT splenocytes were incubated with OMVs isolated from the conditioned media of *B. theta*$^{\Delta tdk}$ or *B. theta*$^{\Delta tdk\Delta BT1160\_trunc}$ (1% v/v). Non-normalized OMVs were resuspended in an equivalent volume of PBS to the original volume of conditioned media they were isolated from (blue and red) and normalized OMVs were concentrated to normalize the protein concentration between the mutant and the parent strain (pink). Samples were incubated for 2 days, and IL-10 was measured in culture supernatants by ELISA.

Data points represent an independent biological **(A, B)** or technical replicate **(C)** where the horizontal bars represent the median **(A, B)**, or mean **(C)**. Graphs represent 5 pooled experiments **(A)**, 3 pooled experiments **(B)**, or are representative of 4 experiments **(C)**. Statistical significance was determined using Kruskal-Wallis test with Dunn's post-hoc test, comparisons between all: **(A, B)**, or One-way ANOVA with Dunnett's post-hoc test, comparisons between all: **(C)**. ns (not significant) $P \geq 0.05$, $*P < 0.05$, $**P \leq 0.01$, $***P \leq 0.001$, $****P \leq 0.0001$.

Methods:

Bacterial strains, culturing conditions, and molecular genetics.

Bacteroides thetaiotaomicron ATCC 29148 (VPI-5482) and its genetic variants were grown in tryptone-yeast extract-glucose (TYG)) broth medium supplemented with 0.005% w/v hemin (see Supplemental Table #1 for recipe) or on brain heart infusion (BHI) agar (BD, 211059) supplemented with 10% v/v defibrinated horse blood (Quad Five, Cat. # 210-500) at 37°C in an anaerobic chamber (Compressed gas: 5% CO_2, 5% H_2, 90% N_2, chamber set to~2.5% H_2, Coy Manufacturing, Grass Lake, MI). Genetic deletions were performed by counter-selectable allelic exchange as previously described[82], and all bacterial strains, plasmids, and primers are referenced in Supplemental Table #2. Briefly, deletion of genes encoding the NQR complex was done using a modified form of *B. theta* strain VPI-5482 with a deletion of the thymidine kinase gene *tdk* (*BT2275*) to facilitate counter-selection. In addition, deletion of genes encoding the NQR complex were also made as indicated using *B. theta*$^{\Delta tdk\Delta cps1-8}$ as the parent strain[61]. Deletion of genes or bacterial gene clusters *BT1160, BT1161, BT1155-BT1159,* and *BT1155-BT1160,* was completed by ligating PCR-amplified 750bp fragments flanking the gene of interest into the suicide vector (pExchange-*tdk*) using Sal1 (NEB, Cat. # R0138L) and Xbal (NEB, Cat. # R0145L) cut sites. This vector contains a cloned copy of *tdk* (provides resistance to the toxic nucleotide analog 5-fluoro-2'-deoxyuridine (FUdR))*, bla* (ampicillin resistance)*,* and *ermGb* (erythromycin resistance), to facilitate selection and counter-selection. The resulting construct was conjugated into either *B. theta*$^{\Delta tdk}$ or *B. theta*$^{\Delta tdk\Delta cps1-8}$. Individual-recombinant merodiploid colonies, selected on erythromycin (25 μg/mL), were pooled and plated on BHI-blood agar containing FUdR (200 μg/mL) to select for recombinants. Candidate gene deletions were screened by PCR and DNA sequencing to identify isolates that had the desired gene deletions.

Complementation of gene *BT1160* was accomplished as previously described in[83]. Briefly, wild-type alleles encoding *BT1160* and promoter regions were amplified using the primers listed in Supplemental Table #2. Complementing alleles were ligated into the *pNBU2-ermGb* vector using Sal1 and BamH1 (NEB, Cat. # R0136L) cut sites and the resulting construct was inserted into *att2* (tRNA^{ser}) in *B. theta*$^{\Delta tdk\Delta BT1160\_trunc}$. Individual recombinant colonies, selected on erythromycin (25 μg/mL) were screened by PCR and DNA sequencing to identify successfully complemented isolates.

B. theta heat-inactivated pellets and conditioned media.

B. theta was grown anaerobically in TYG growth medium overnight to stationary phase (~16 hours) and the OD$_{600}$ measured (ranging from ~2.0-3.5). When comparing the effects of multiple *B. theta* strains or mutants on cytokine induction, cultures were normalized by OD using TYG. Normalized cultures were spun down at 7000 x g for 5 minutes at 4°C, and the supernatant was separated from the pellet by pouring off into a fresh tube. The supernatant was filtered through a 0.22 μm PVDF filter (Millex, Cat. # SLGVR33RS) to make 1-~1.5 mL aliquots of cell-free supernatant, referred to as conditioned media hereafter. Conditioned media was frozen at -20°C until further use. The bacterial pellet was washed by resuspending in sterile PBS, spun down as before, and resuspended in a volume of sterile PBS equivalent to the volume of media from which the pellet was obtained, the OD$_{600}$ was measured again, and additional dilutions were made in PBS such that equivalent CFU (based on OD$_{600}$) of different strains or mutants were used to stimulate cells in downstream assays. Aliquots of the washed and OD$_{600}$ normalized pellets (1 mL) were transferred to sterile Eppendorf tubes and placed at 70°C for 30 minutes to heat-inactivate them. Heat-inactivated pellets were stored at -20°C until use. The effectiveness of sterile filtering the conditioned media was confirmed by plating on BHI blood agar plates and incubated at 37°C anaerobically for >3 days (data not shown).

Outer Membrane Vesicle (OMV) isolation.

OMVs were isolated from conditioned media using an ultracentrifugation based approach as previously described[84]. A set volume of conditioned media was put into 1.5 mL ultracentrifugation tubes (ThermoScientific, Cat. # 314352H01) and spun at 100,000 x g for 2 hours at 4°C. Following the spin, the supernatant was removed and the pellet (containing the OMVs) was resuspended and washed in 1 mL of sterile PBS (all PBS was first double filtered through a 0.1 μm PES filter (Thermo Scientific, Cat. #5670010) before use). This resuspension was spun down again at 100,000 x g for 2 hours, the supernatant was removed, and pellet containing the washed OMVs, were resuspended in 0.1 μm filtered PBS. OMVs were either concentrated (for Qubit concentration analysis, see below) or resuspended in a volume of PBS equivalent to the volume of conditioned media from which they were extracted (for IL-10 induction assay, see below). Isolated OMVs were stored at -20°C until use.

In vitro IL-10 assay.

The ability of the heat-inactivated pellets and conditioned media from *B. theta* strains or mutants to induce IL-10 was tested with an *in vitro* assay using bulk unfractionated murine splenocytes. Splenocytes were isolated from the spleen of wild-type, TLR2$^{-/-}$, or Dectin-1$^{-/-}$ mice by forcing through a 70 μm cell-strainer using sterile PBS containing 0.1% w/v Bovine Serum Albumin (BSA; Sigma). Cells were pipetted up and down to generate a single cell suspension and collected in a 15 mL conical tube, and pelleted by centrifugation at 454 x g for 5 minutes. The supernatant was removed, the pellet was resuspended by flicking the tube and 750 μL (per spleen) of room temperature ACK lysis buffer (Gibco, Cat. # A10492) was added, followed by immediate vortexing. The cells were incubated for 3 minutes at room temperature, and ~10 mL of PBS/0.1% BSA was then added to the cells to halt lysis. The cell suspension was then passed through a 70 μm cell-strainer and the flow-through collected into a new 15 mL conical tube. Cells were pelleted as before, the supernatant removed, and the pellet was resuspended in 5 mL complete RPMI (RPMI 1640 supplemented with L-glutamine, 20 mM final concentration HEPES (Gibco, Cat # 15630-080), 10% v/v final concentration heat-inactivated fetal bovine serum (FBS), 100 units/mL Penicillin and 100 μg/mL Streptomycin) per spleen. Live cells were counted by hemocytometer (stained with trypan blue) and diluted with complete RPMI to 5x10^6 cells/mL.

Splenocytes were plated in a flat-bottomed sterile 96-well tissue culture treated plate (Gibco, Cat. # 353072) by adding 100 μL to each well for a final concentration of 5x10^5 cells/well. Splenocytes were then incubated with heat-inactivated bacterial pellets (OD$_{600}$ 0.05-1.0), conditioned media (0.1-7% v/v), or OMVs (0.1-5% v/v) and complete RPMI was added to each well to bring the final volume to 200 μL. Cells and samples were incubated at 37°C in 5% CO$_2$ for 48 hours, following which the cells were pelleted by centrifugation at 454 x g for 5 minutes and the supernatants collected. Supernatants were frozen at -20°C until further processing, and IL-10 was measured by ELISA (Biolegend, Cat. # 431411).

The *in vitro* IL-10 assay was also run on BMDMs from wild-type or TLR2$^{-/-}$ mice. Briefly, bone marrow was isolated from the femur and tibia using a tube-in-tube centrifugation method, where a hole was poked in the bottom of a sterile PCR tube using a sterile 18G needle and then placed in a sterile Eppendorf tube. The ends of the femur and tibia were removed and the bones were placed open side down in the PCR tube. The tubes were centrifuged briefly to remove the bone marrow and lysed with ACK lysis buffer (as described above; 100uL per tube). Lysis was halted with PBS/0.1% BSA and all the bone marrow was combined and the cells were pelleted at 454 x g for 5 minutes before being resuspended in BMDM media (DMEM-low glucose with L-glutamine and sodium pyruvate, 20% v/v FBS, 100 units/mL Penicillin, 100 μg/mL Streptomycin, 30% v/v L929-cell media, 20mM HEPES). Note, the bone marrow was differentiated into BMDMs using macrophage colony-stimulating factor (M-CSF) provided by the L929-cell media (LCM)[85]. LCM was generated by harvesting and sterile filtering the supernatant from confluent cell culturing of L929 cells (NCPC L929, ATCC # CCL-1) in DMEM media (DMEM-low glucose with L-glutamine and sodium pyruvate, 100 units/mL Penicillin and 100 μg/mL Streptomycin, 10% v/v FBS). ACK lysed bone marrow was plated at 2x10^6 cells per non-tissue culture treated petri dish in 10 mL BMDM media. The media was changed every three days, and the macrophages were harvested using versene (Gibco, Cat. #15040-066) after day 10 when the cells were adherent and elongated. BMDMs were plated the same as the splenocytes (as described above), at a final concentration of 5x10^5 cells/well, and the supernatant was collected after 2 days for detection of IL-10. All spleens/bone marrow were isolated from animals whose use was approved by the Institutional Animal Care and Use Committee (IACUC) of the Cleveland Clinic (wild-type [Jackson Labs, strain #: 000664] and TLR2-/- [Jackson Labs, strain #: 004650]), the IACUC of Case Western Reserve University (TLR2-/- [generously provided to C.V.H. by Shizuo Akira, Research Institute for Microbial Disease, Osaka University, Osaka, Japan, and back-crossed to C57BL/6J mice a minimum of eight times as per [86]]) or the IACUC of the University of Minnesota (Jackson Labs, wild-type [strain #: 000664] and Dectin-1-/- [Jackson Labs, strain #: 012337]). Dectin-1-/- mice are reported by Jackson Labs to be on a mixed C57BL/6N;C57BL/6J genetic background. All mice used were male unless otherwise indicated.

In vitro IL-10 assays were also performed using human Peripheral Blood Mononuclear Cells (PBMCs) (STEMCELL Technologies, Cat. # 70025.1). The assay was performed as described above, and PBMCs were thawed and handled according to the manufacturer instructions. Briefly, the outside of the frozen vial of cells was wiped with 70% ethanol, vented in a biosafety cabinet by quickly uncapping the tube to release built up

pressure, and then recapped. The cells were thawed at 37°C in a water bath by gently shaking the vial. The vial was removed from the water bath while a small frozen cell pellet remained. The outside of the vial was cleaned with 70% ethanol again. Cells were transferred to a 50 mL conical and 1 mL of complete RPMI was added to the cells dropwise. Cells were washed by slowly adding 15-20 mL complete RPMI dropwise while swirling the tube. Cells were spun down at 454 x g for 10 minutes at room temperature and the supernatant was carefully removed. Cells were resuspended by gently flicking the tube in the remaining volume. 100 μg/mL of DNase 1 (STEMCELL Technologies, Cat. # 07900) was added to the cell suspension for 15 minutes at room temperature to prevent cell clumping. Cells were washed with 15-20 mL complete RPMI and spun as described above. The supernatant was removed and the pellet was resuspended by gently flicking the tube. The volume was measured and live cells were counted by hemocytometer (stained with trypan blue) and diluted with complete RPMI to $5x10^6$ cells/mL. PBMCs were then plated as described above and human IL-10 was measured in the cell supernatants by ELISA (Biolegend, Cat. # 430604).

IL-10 induction by transposon mutagenesis library.

The ability of *B. theta* transposon mutants to induce IL-10 was tested using an arrayed *B. theta*$^{\Delta tdk\Delta cps1-8}$ transposon mutagenesis library. The library was previously generated using the *mariner* transposon[35]. Each 96-well plate of mutants was grown anaerobically at 37°C in TYG media, to stationary phase, overnight, and frozen at -80°C until use. The plates were thawed and spun at 454 x g for 5 minutes to separate the bacterial pellet from the TYG media supernatant. Splenocytes were stimulated as described above using a dose of 1% v/v conditioned media or Pam3CSK4 (each plate had wells stimulated with Pam3CSK4 as a positive control), and IL-10 induction was measured in the supernatant after 2 days by ELISA. Each mutant was tested in duplicate and normalized to the Pam3CSK4 positive control value to account for plate to plate variation. The data was processed with custom R script. Briefly, the normalized IL-10 expression levels were converted to log2 scale for normalization, and then any transposon mutant with an IL-10/Pam3CSK4 ratio data point below 2 standard deviation were filtered, and z-score and p-value were computed for each using normal distribution statistics. Adjusted p-values were calculated by the false discovery rate (FDR) to correct for multiple hypothesis testing. Transposon mutant 19_H4 and the acapsular parent strain were sent for whole Illumina whole genome sequencing (2 X 151 bp) to identify the insertion site of the transposon.

Nqr locus gene expression

RNA was isolated from three 5 mL cultures of *B. theta*$^{\Delta tdk\Delta BT1160}$_trunc, *B. theta*$^{\Delta tdk\Delta BT1160}$_full, and their parent strains (*B. theta*$^{\Delta tdk}$) grown to mid-log phase (OD 0.5-0.8) in TYG. The 5 mL cultures were transferred to a 15 mL conical tube and spun down at 7000 x g for 5 minutes at room temperature. The supernatant was removed (note, a 1 mL aliquot was sterile filtered and kept to test conditioned media for IL-10 induction (as above)) and replaced with 5 mL of RNA-protect (Qiagen, Cat. # 1018390), mixed by pipetting, and allowed to sit for 5 minutes at room temperature. Samples were spun for 10 minutes at 7000 x g at room temperature and the supernatant was decanted, and all additional liquid was removed by gently taping the tube on a paper towel. Samples were stored at -80°C until RNA was isolated.

RNA was isolated following manufactures instructions provided by RNeasy kit (Qiagen, Cat. # 74004). Briefly, pellets were brought to room temperature and resuspended in 200 μL of TE buffer containing 1 mg/mL lysozyme (Sigma, Cat. #L4919) by pipetting, and allowed to incubate for 5 minutes at room temperature, pipetting for 10 seconds every 2 minutes. 700 μL of RLT buffer (containing β-mercaptoethanol (BME) 10 μL/mL) was added to the solution and mixed by pipetting, followed by 500 μL of 100% ethanol and mixed by pipetting. The solution was transferred to an RNeasy Mini Spin column (Econospin Cat. # 1940-250) and spun for 15 seconds at 18,111 x g and the flow-through was discarded. 700 μL of Buffer RW1 was added to the column and spun again, as above, to wash the membrane. The flow through was discarded and 500 μL of Buffer RPE was added to the column and spun again. The flow-through was discard and this step was repeated followed by a 2 minute spin. The flow-through was discarded and the tube was spun again for 1 minute to remove any residual ethanol. The column was placed in an RNase-free 1.5 mL collection tube (Ambion, Cat. # AM12400) and 30 μL of nuclease free water was added to the column membrane, let to sit for 5 minutes at room temperature and then spun at 8476 x g for 1 minute. This step was repeated for a total of 60μL of elution volume. RNA concentration was determined by nanodrop and DNase 1 treated (NEB, Cat. # M0303L). DNase 1 (4 units) was added to each sample and let to incubate at 37°C for 30 minutes. RNA was re-purified by adding 350 μL of RLT buffer containing BME (10 μL/mL), followed by 250 μL of 100% ethanol and mixed by pipetting. The solution was transferred to an RNeasy Mini Spin column and washed and eluted

as above. The concentration of each sample was determined by nanodrop. RNA was confirmed free of DNA contamination by running a qPCR for the *BT1155* gene of *B. theta* (see Supplemental Table #2) using an in-house qPCR mix (see below) on the purified RNA prior to cDNA synthesis, and were considered DNA-free when a Ct value greater than our background value of 30 was obtained[87,88].

Reverse transcription was performed using an adapted SuperScript III First-Strand Synthesis System (Invitrogen, Cat. # 18080-051) to generate cDNA. In brief, 1 μg of RNA was incubated with 10 mM dNTP mix (1 μL/sample) and random hexamers (200 ng/sample) in a final volume of 13 μL at 65°C for 5 minutes followed by 5 minutes on ice. Each sample was spun down and first strand buffer (1X, 4 μL/sample), 0.1M DTT (1 μL/sample), SuperScript III enzyme (200 units/sample), and water (1 μL/sample) were added to a final volume of 20 μL. The sample was mixed by pipetting gently and the tube was spun down prior to incubating for 1 hour at 60°C. The reaction was halted by inactivating the enzyme at 70°C for 15 minutes. A 1-to-1 conversion of RNA to cDNA was assumed and each sample was diluted with water to 5 ng/μL and stored at -20°C until use.

The abundance of each target transcript (*BT1154-BT1161*) was quantified using an in-house qPCR mix as described in[87,88] (see Supplemental Table #2 for qPCR primer sequences). In brief, each 20 μL reaction contained 1X SYBR Green 1 (Lonza, Cat. # 50513), 1X Thermopol Reaction Buffer (NEB, Cat. # B9004S), 125 μM dNTPs, 2.5 mM MgSO4, 500 nM of gene specific primers, and 0.5 units Hot Start *Taq* Polymerase (NEB, Cat. # M0495L), and 10 ng of template cDNA. 16S rRNA was used as a housekeeping gene for each sample, and a water control was run for each primer condition. A Ct value greater than our background value of 30 was considered non-detectable[87,88]. Each sample was run in duplicate and the Ct values were averaged. The ΔCt values were calculated by normalizing the gene of interest Ct value to the 16S rRNA housekeeping gene value for each sample. Next, the Δ/ΔCt value was calculated by normalizing the ΔCt value for *BT1160* deletion samples to the ΔCt value of the parent strain (Δ*tdk*). Fold change was calculating using the $2^{\Delta/\Delta Ct}$ method.

OMV comparative proteomics using liquid chromatography with tandem mass spectrometry (LC-MS/MS).

Outer membrane vesicles were isolated from *B. theta*$^{\Delta tdk\Delta BT1160\_trunc}$, *B. theta*$^{\Delta tdk\Delta cps1\text{-}8\Delta BT1160\_trunc}$, and their corresponding parent strains (*B. theta*$^{\Delta tdk}$ and *B. theta*$^{\Delta tdk\Delta cps1\text{-}8}$) grown to stationary phase in TYG. Conditioned media from three 250 mL cultures of each mutant was made as described above. Each 250 mL batch of conditioned media was divided between six 39 mL QuickSeal Ultracentrifugation tubes (Beckman Coulter, Cat. # 342414) and OMVs isolated by ultracentrifugation at 100,000 $x\ g$ (as describe above). The remaining conditioned media was kept to test its ability to induce IL-10 using the *in vitro* splenocyte assay (describe above) to confirm that the conditioned media used in these measurements displayed the expected immune-stimulatory capacity. OMV pellets were resuspended and combined in a total of 1 mL sterile PBS to wash, and put through another round of ultracentrifugation. Washed OMV pellets were resuspended in 300 μL sterile milli-Q water and frozen at -80°C. An aliquot was taken to test their ability to induce IL-10 to confirm that the OMV used in these measurements displayed the expected immune-stimulatory capacity, and were further assessed by ZetaView (as describe below) to ensure OMV were successfully isolated.

Prior to LC-MS/MS, samples were lyophilized and then lysed in 200 μL Radio-Immunoprecipitation Assay (RIPA) buffer using a probe sonicater, and protein concentration was measured by BCA assay. A 100 μg aliquot of each sample was precipitated with acetone overnight, and the pellets dried and reconstituted in 6M urea/Tris buffer. Samples were then reduced with 1 μL of 200 mM dithiothreitol, alkylated by 4 μL of 200 mM iodoacetamide, then digested with 4 μL of a trypsin/Lys-C Mix (0.5 μg/uL) (Promega # V5071) using a two-step in-solution digestion overnight at 37°C. Digestion was terminated by the addition of trifluoracetic acid to a final concentration of 0.5-1% v/v. The samples were then desalted using Sep-Pak C18 cc Vac Cartridge (WAT054955) and reconstituted in 30 μL 0.1% v/v formic acid.

Prepared samples were run on a LC-MS Dionex Ultimate 3000 nano-flow high performance liquid chromatography (HPLC) interfacing with a ThermoScientific Fusion Lumos mass spectrometer system. The HPLC system used an Acclaim PepMap 100 precolum (75 μm x 2 cm, C18, 3 μm, 100 A) followed by an Acclaim PepMap RSLC analytical column (75 μm x 15 cm, C18, 2 μm, 100 A). 5 μL volumes of the extract were injected and the peptides eluted from the column by an acetonitrile/0.1% v/v formic acid gradient at a flow rate of 0.3 μL/min were introduced into the source of the mass spectrometer online. The micro-electrospray ion source is operated at 2.5 kV. The digest was analyzed using the data dependent multitask capability of the

instrument acquiring full scan mass spectra to determine peptide molecular weights and product ion spectra to determine amino acid sequence in successive instrument scans. The data were searched against the *B. theta* (VPI-5482) UniProtKB protein database with the program Andomeda (4782 sequences). LC-MS spectra intensities for identified proteins were normalized and expressed as label free quantity (LFQ) intensities calculated in the program MaxQuant version 2.0.1.0. The data matrix was uploaded into Perseus and proteins that were not found in at least 2 samples were discarded from the analyses. Zero intensity values were imputed by replacing intensities with a normal distribution. The Δ*tdk*/Δ*tdk*Δ*BT1160*_trunc LFQ ratio (fold change) was calculated on averaged LFQ intensities and the p-value calculated using Student's T-test.

OMV quantification and normalization.
The concentration of OMVs isolated from *B. theta* mutants lacking *BT1160, BT1155-BT1159,* and *BT1155-BT1160* and their parent strain were determined using Qubit (measures protein concentration) or ZetaView (a nanoparticle analyzer with Particle Metrix software)[66]. OMVs were isolated from 1 mL aliquots of conditioned media from *B. theta*$^{\Delta tdk}$ and mutant strains as described above. The concentration of isolated OMVs were measured by Qubit and Zetaview. For Qubit (Invitrogen, Cat. # Q33211), the OMVs from 1 mL of conditioned media were concentrated 50X by resuspending in 20 μL of sterile filtered PBS. The concentration was measured in μg /mL and then the 1X original concentration of OMVs was calculated. For ZetaView, OMVs isolated from 1 mL of conditioned media were resuspended in a volume of sterile filtered PBS equivalent to the original volume from which they were obtained (as described above) and 1 mL was injected into the ZetaView. Once the concentration was determined for both the mutants and parent strain, OMVs were isolated from the same batch of conditioned media and resuspended in PBS at either the original volume (for *B. theta*$^{\Delta tdk}$) or, for mutants whose OMV concentration was lower (example: *B. theta*$^{\Delta tdk\Delta BT1160}$), in a volume that resulted in the same concentration of OMVs as *B. theta*$^{\Delta tdk}$. Concentrated OMVs from the mutants and OMVs from the parent strain were then tested for their ability to induce IL-10 using the *in vitro* IL-10 assay described above. The same percentage of isolated OMVs, from both the mutant and parent strain, were added to the splenocytes.

Transmission electron microscopy (TEM) of OMVs.
50 mL cultures of *B. theta*$^{\Delta tdk}$ and *B. theta*$^{\Delta tdk\Delta BT1160}$_trunc were grown in TYG anaerobically overnight to stationary phase at 37°C and the OD$_{600}$ was measured. Mutants were normalized to a final OD of 2.95 using TYG and the conditioned media was isolated as described above. Briefly, cultures were spun at 7000 $x\ g$ for 5 minutes and the supernatant was filtered through a 0.2 μm PES filter (MilliQ Steriflip, Cat. #SCGP00525). 1 mL aliquots of conditioned media were removed and frozen at -20°C to test the mutant and parent strains ability to induce IL-10 using the assay described above to confirm that those OMV that were imaged showed the expected immune-stimulatory characteristics. The conditioned media from each mutant was then concentrated from 50 mL to 1 mL using 100 kDa size cut off filters (Vivaspin20 Cat. # 28932363). OMVs remain in the larger than 100 kDa top fraction and were then subject to OMV isolation using the ultracentrifugation steps described above. The washed pellet containing the OMVs was resuspended in 100 μL of PBS, the solution was turbid, and frozen at -20°C until use.
OMVs were prepared for TEM by negative staining. 5 μL of each OMV sample was added to the carbon-coated copper 400 mesh grid and incubated at room temperature for 10 minutes. The samples were washed with 5 drops of 5 mM Tris buffer (pH 7.1) and then with 5 drops of distilled water. The specimen was stained with 5 drops of 2% uranyl acetate. The grid was allowed to dry completely for several hours or overnight. The prepared specimens were visualized with an electron microscope operated at 80 kV.

Statistical analysis
Statistical analyses were performed in GraphPad Prism version 9.0.0. Details of the specific statistical tests used is included in the figure legends where appropriate.

Supplemental Figure 1. Cytokine induction by *B. theta* strains on WT, TLR2$^{-/-}$, and Dectin-1$^{-/-}$ splenocytes and BMDMs, related to Figure 1.

(A) WT BMDMs were stimulated with *B. theta*$^{VPI-5482}$ heat-inactivated pellet (OD$_{600}$=1.0) or RPMI media, or *B. theta*$^{VPI-5482}$ conditioned media (1% v/v) or TYG media (1% v/v) control for 2 days and IL-10 was assessed in the supernatant by ELISA.
(B) WT splenocytes were stimulated with *B. theta*$^{VPI-5482}$ heat-inactivated pellet (OD$_{600}$=1.0) or RPMI media, or *B. theta*$^{VPI-5482}$ conditioned media (1% v/v) or TYG media control for 2 days and IL-6 was assessed in the supernatant by ELISA.
(C) WT and TLR2$^{-/-}$ BMDMs were stimulated with *B. theta*$^{VPI-5482}$ conditioned media (1% v/v) or TYG media control for 2 days and IL-10 was assessed in the supernatant by ELISA.
(D) WT and TLR2$^{-/-}$ splenocytes were stimulated with conditioned media (1% v/v) from *B. theta* (strain 0940-1) or TYG media control for 2 days and IL-10 was assessed in the supernatant by ELISA.
(E) WT and Dectin-1$^{-/-}$ splenocytes were stimulated with *B. theta*$^{VPI-5482}$ conditioned media (1% v/v) or TYG media control for 2 days and IL-10 was assessed in the supernatant by ELISA.

Data points represent an independent technical replicate where the horizontal bars represent the mean **(A-E)**. Graphs are representative of at least 2 experiments. Statistical significance was determined Student's T test: **(A, B)** or Two-way ANOVA with Sidak's post hock test, comparisons to WT **(C-E)**. ns P≥0.05, **P≤0.01, ****P≤0.0001.

Supplemental Figure 2.

Supplemental Figure 2. *B. theta* gene *BT1160* mutants on acapsular and wild-type backgrounds, or parent strain, growth curves and IL-10 induction on BMDMs, related to Figure 2.

(A, B) *B. theta* wild-type (Δtdk) and acapsular *B. theta* ($\Delta tdk \Delta cps1$-8) and *B. theta* with the truncated deletion of gene *BT1160* on each background were grown in TYG for growth curve analysis. Growth curves were performed in 96 well plates with 12 technical replicates per mutant. Growth (OD_{600}) was measured continuously over 48 hours with readings every 10 minutes. Tables represent the average and standard deviation of the Specific Rate, Total growth, and Lag Time of all 12 technical replicates for each mutant. Statistical significance calculated comparing the mutant to the appropriate parent strain and reported in the tables under the mutant.

(C) WT BMDMs were stimulated with conditioned media (1% v/v) from *B. theta* wild-type (Δtdk) and the *B. theta* with the truncated deletion of gene *BT1160* for 2 days and IL-10 was assessed in the supernatant by ELISA.

Data points represent the average of the 12 technical replicates for each time point **(A, B)**, or individual technical replicates of a single experiment and horizontal bars represent the mean **(C)**. Statistical significance was determined by Student's T test: **(A-C)**. **P≤0.01.

Supplemental Figure 3.

Supplemental Figure 3. Additional biological replicates of *B. theta BT1160* complementation and *nqr* locus mutants, related to Figure 3.

(A) WT splenocytes were stimulated with *B. theta*$^{\Delta tdk}$, *B. theta*$^{\Delta tdk \Delta BT1160\_\text{trunc}}$, or *B. theta*$^{\Delta tdk \Delta BT1160\_\text{trunc}::BT1160}$ conditioned media (1% v/v) for 2 days and IL-10 was assessed in the supernatant by ELISA. Each graph shows the results from an independent experiment. **(B)** WT splenocytes were stimulated with *B. theta*$^{\Delta tdk}$, *B. theta*$^{\Delta tdk \Delta BT1160\_\text{trunc}}$, *B. theta*$^{\Delta tdk \Delta BT1155\text{-}BT1159}$, *B. theta*$^{\Delta tdk \Delta BT1155\text{-}BT1160}$, or *B. theta*$^{\Delta tdk \Delta BT1161}$ conditioned media or TYG media control (1% v/v) for 2 days and IL-10 was assessed in the supernatant by ELISA. Each graph shows the results from an independent experiment.

Data points on each graph represent an independent technical replicate where the horizontal bars represent the mean **(A, B)**. Statistical significance was determined using One-way ANOVA with Dunnett's post-hoc test, comparisons between all: **(A, B)**. ns P≥0.05, *P<0.05, **P≤0.01, ***P≤0.001, ****P≤0.0001.

Supplemental Figure 4.

Supplemental Figure 4. IL-10 induction by *B. theta nqr* locus mutants on WT splenocytes isolated from male and female mice, related to Figure 3 and 4.

(A) WT splenocytes, isolated from male and female mice, were stimulated with *B. theta*$^{\Delta tdk}$, *B. theta*$^{\Delta tdk \Delta BT1160}$_trunc, or *B. theta*$^{\Delta tdk \Delta BT1155-BT1160}$ conditioned media, or TYG control media (1% v/v) for 2 days and IL-10 was assessed in the supernatant by ELISA.
(B) WT splenocytes, isolated from male and female mice, were stimulated with *B. theta*$^{\Delta tdk}$, *B. theta*$^{\Delta tdk \Delta BT1160}$_trunc, or *B. theta*$^{\Delta tdk \Delta BT1155-BT1160}$ OMVs isolated from conditioned media or TYG control media (1% v/v; resuspended in PBS) for 2 days and IL-10 was assessed in the supernatant by ELISA.

Data points represent an independent technical replicate where the horizontal bars represent the mean **(A, B)**. Statistical significance was determined using One-way ANOVA with Dunnett's post-hoc test, comparisons to *B. theta*$^{\Delta tdk}$: **(A, B)**. ****P≤0.0001.

Supplemental Figure 5.

Supplemental Figure 5. OMV images and concentration measurement, related to Figure 4, 5.

(A) Qubit and ZetaView were used to measure the concentration of OMVs isolated from the same batch of conditioned media of *B. theta*$^{\Delta tdk}$ and *B. theta*$^{\Delta tdk\Delta BT1160}$_trunc. Each data point represents an independent biological replicate where the horiziontal bars represent the median.
(B) Representative TEM images of concentrated OMVs isolated from the conditioned media of *B. theta*$^{\Delta tdk}$ and *B. theta*$^{\Delta tdk\Delta BT1160}$_trunc.

References:

1. Ley, R.E., Lozupone, C.A., Hamady, M., Knight, R., and Gordon, J.I. (2008). Worlds within worlds: evolution of the vertebrate gut microbiota. Nat Rev Microbiol *6*, 776-788. 10.1038/nrmicro1978.
2. Savage, D.C. (1977). Microbial ecology of the gastrointestinal tract. Annu Rev Microbiol *31*, 107-133. 10.1146/annurev.mi.31.100177.000543.
3. Walter, J., and Ley, R. (2011). The human gut microbiome: ecology and recent evolutionary changes. Annu Rev Microbiol *65*, 411-429. 10.1146/annurev-micro-090110-102830.
4. Marchiando, A.M., Graham, W.V., and Turner, J.R. (2010). Epithelial barriers in homeostasis and disease. Annu Rev Pathol *5*, 119-144. 10.1146/annurev.pathol.4.110807.092135.
5. Turner, J.R. (2009). Intestinal mucosal barrier function in health and disease. Nat Rev Immunol *9*, 799-809. 10.1038/nri2653.
6. Mowat, A.M. (2003). Anatomical basis of tolerance and immunity to intestinal antigens. Nat Rev Immunol *3*, 331-341. 10.1038/nri1057.
7. Faith, J.J., Ahern, P.P., Ridaura, V.K., Cheng, J., and Gordon, J.I. (2014). Identifying gut microbe-host phenotype relationships using combinatorial communities in gnotobiotic mice. Sci Transl Med *6*, 220ra211. 10.1126/scitranslmed.3008051.
8. Round, J.L., and Mazmanian, S.K. (2010). Inducible Foxp3+ regulatory T-cell development by a commensal bacterium of the intestinal microbiota. Proc Natl Acad Sci U S A *107*, 12204-12209. 10.1073/pnas.0909122107.
9. Atarashi, K., Tanoue, T., Shima, T., Imaoka, A., Kuwahara, T., Momose, Y., Cheng, G., Yamasaki, S., Saito, T., Ohba, Y., et al. (2011). Induction of colonic regulatory T cells by indigenous Clostridium species. Science *331*, 337-341. 10.1126/science.1198469.
10. Verma, R., Lee, C., Jeun, E.J., Yi, J., Kim, K.S., Ghosh, A., Byun, S., Lee, C.G., Kang, H.J., Kim, G.C., et al. (2018). Cell surface polysaccharides of Bifidobacterium bifidum induce the generation of Foxp3(+) regulatory T cells. Sci Immunol *3*. 10.1126/sciimmunol.aat6975.
11. Read, S., Malmström, V., and Powrie, F. (2000). Cytotoxic T lymphocyte-associated antigen 4 plays an essential role in the function of CD25(+)CD4(+) regulatory cells that control intestinal inflammation. J Exp Med *192*, 295-302. 10.1084/jem.192.2.295.
12. Maloy, K.J., Antonelli, L.R., Lefevre, M., and Powrie, F. (2005). Cure of innate intestinal immune pathology by CD4+CD25+ regulatory T cells. Immunol Lett *97*, 189-192. 10.1016/j.imlet.2005.01.004.
13. Geuking, M.B., Cahenzli, J., Lawson, M.A., Ng, D.C., Slack, E., Hapfelmeier, S., McCoy, K.D., and Macpherson, A.J. (2011). Intestinal bacterial colonization induces mutualistic regulatory T cell responses. Immunity *34*, 794-806. 10.1016/j.immuni.2011.03.021.
14. Atarashi, K., Tanoue, T., Oshima, K., Suda, W., Nagano, Y., Nishikawa, H., Fukuda, S., Saito, T., Narushima, S., Hase, K., et al. (2013). Treg induction by a rationally selected mixture of Clostridia strains from the human microbiota. Nature *500*, 232-236. 10.1038/nature12331.
15. Danne, C., Ryzhakov, G., Martinez-Lopez, M., Ilott, N.E., Franchini, F., Cuskin, F., Lowe, E.C., Bullers, S.J., Arthur, J.S.C., and Powrie, F. (2017). A Large Polysaccharide Produced by Helicobacter hepaticus Induces an Anti-inflammatory Gene Signature in Macrophages. Cell Host Microbe *22*, 733-745 e735. 10.1016/j.chom.2017.11.002.
16. Martin-Gallausiaux, C., Béguet-Crespel, F., Marinelli, L., Jamet, A., Ledue, F., Blottière, H.M., and Lapaque, N. (2018). Butyrate produced by gut commensal bacteria activates TGF-beta1 expression through the transcription factor SP1 in human intestinal epithelial cells. Sci Rep *8*, 9742. 10.1038/s41598-018-28048-y.
17. Garrett, W.S., Gordon, J.I., and Glimcher, L.H. (2010). Homeostasis and inflammation in the intestine. Cell *140*, 859-870. 10.1016/j.cell.2010.01.023.
18. Bennett, C.L., Christie, J., Ramsdell, F., Brunkow, M.E., Ferguson, P.J., Whitesell, L., Kelly, T.E., Saulsbury, F.T., Chance, P.F., and Ochs, H.D. (2001). The immune dysregulation, polyendocrinopathy, enteropathy, X-linked syndrome (IPEX) is caused by mutations of FOXP3. Nat Genet *27*, 20-21. 10.1038/83713.
19. Asseman, C., Mauze, S., Leach, M.W., Coffman, R.L., and Powrie, F. (1999). An essential role for interleukin 10 in the function of regulatory T cells that inhibit intestinal inflammation. J Exp Med *190*, 995-1004. 10.1084/jem.190.7.995.
20. Asseman, C., Read, S., and Powrie, F. (2003). Colitogenic Th1 cells are present in the antigen-experienced T cell pool in normal mice: control by CD4+ regulatory T cells and IL-10. J Immunol *171*, 971-978. 10.4049/jimmunol.171.2.971.

21. Kühn, R., Löhler, J., Rennick, D., Rajewsky, K., and Müller, W. (1993). Interleukin-10-deficient mice develop chronic enterocolitis. Cell *75*, 263-274. 10.1016/0092-8674(93)80068-p.
22. Glocker, E.O., Kotlarz, D., Boztug, K., Gertz, E.M., Schäffer, A.A., Noyan, F., Perro, M., Diestelhorst, J., Allroth, A., Murugan, D., et al. (2009). Inflammatory bowel disease and mutations affecting the interleukin-10 receptor. N Engl J Med *361*, 2033-2045. 10.1056/NEJMoa0907206.
23. Wildin, R.S., Ramsdell, F., Peake, J., Faravelli, F., Casanova, J.L., Buist, N., Levy-Lahad, E., Mazzella, M., Goulet, O., Perroni, L., et al. (2001). X-linked neonatal diabetes mellitus, enteropathy and endocrinopathy syndrome is the human equivalent of mouse scurfy. Nat Genet *27*, 18-20. 10.1038/83707.
24. Tedde, A., Laura Putignano, A., Bagnoli, S., Congregati, C., Milla, M., Sorbi, S., Genuardi, M., and Papi, L. (2008). Interleukin-10 promoter polymorphisms influence susceptibility to ulcerative colitis in a gender-specific manner. Scand J Gastroenterol *43*, 712-718. 10.1080/00365520701885507.
25. Powrie, F., Carlino, J., Leach, M.W., Mauze, S., and Coffman, R.L. (1996). A critical role for transforming growth factor-beta but not interleukin 4 in the suppression of T helper type 1-mediated colitis by CD45RB(low) CD4+ T cells. J Exp Med *183*, 2669-2674. 10.1084/jem.183.6.2669.
26. Shull, M.M., Ormsby, I., Kier, A.B., Pawlowski, S., Diebold, R.J., Yin, M., Allen, R., Sidman, C., Proetzel, G., Calvin, D., and et al. (1992). Targeted disruption of the mouse transforming growth factor-beta 1 gene results in multifocal inflammatory disease. Nature *359*, 693-699. 10.1038/359693a0.
27. Tamizifar, B., Lankarani, K.B., Naeimi, S., Rismankar Zadeh, M., Taghavi, A., and Ghaderi, A. (2008). Promoter polymorphism of transforming growth factor-beta1 gene and ulcerative colitis. World J Gastroenterol *14*, 243-247. 10.3748/wjg.14.243.
28. Kullberg, M.C., Jankovic, D., Gorelick, P.L., Caspar, P., Letterio, J.J., Cheever, A.W., and Sher, A. (2002). Bacteria-triggered CD4(+) T regulatory cells suppress Helicobacter hepaticus-induced colitis. J Exp Med *196*, 505-515. 10.1084/jem.20020556.
29. Xu, M., Pokrovskii, M., Ding, Y., Yi, R., Au, C., Harrison, O.J., Galan, C., Belkaid, Y., Bonneau, R., and Littman, D.R. (2018). c-MAF-dependent regulatory T cells mediate immunological tolerance to a gut pathobiont. Nature *554*, 373-377. 10.1038/nature25500.
30. Erturk-Hasdemir, D., Oh, S.F., Okan, N.A., Stefanetti, G., Gazzaniga, F.S., Seeberger, P.H., Plevy, S.E., and Kasper, D.L. (2019). Symbionts exploit complex signaling to educate the immune system. Proc Natl Acad Sci U S A *116*, 26157-26166. 10.1073/pnas.1915978116.
31. Mazmanian, S.K., Round, J.L., and Kasper, D.L. (2008). A microbial symbiosis factor prevents intestinal inflammatory disease. Nature *453*, 620-625. 10.1038/nature07008.
32. Round, J.L., Lee, S.M., Li, J., Tran, G., Jabri, B., Chatila, T.A., and Mazmanian, S.K. (2011). The Toll-like receptor 2 pathway establishes colonization by a commensal of the human microbiota. Science *332*, 974-977. 10.1126/science.1206095.
33. Shen, Y., Giardino Torchia, M.L., Lawson, G.W., Karp, C.L., Ashwell, J.D., and Mazmanian, S.K. (2012). Outer membrane vesicles of a human commensal mediate immune regulation and disease protection. Cell Host Microbe *12*, 509-520. 10.1016/j.chom.2012.08.004.
34. Sefik, E., Geva-Zatorsky, N., Oh, S., Konnikova, L., Zemmour, D., McGuire, A.M., Burzyn, D., Ortiz-Lopez, A., Lobera, M., Yang, J., et al. (2015). MUCOSAL IMMUNOLOGY. Individual intestinal symbionts induce a distinct population of RORγ$^+$ regulatory T cells. Science *349*, 993-997. 10.1126/science.aaa9420.
35. Wegorzewska, M.M., Glowacki, R.W.P., Hsieh, S.A., Donermeyer, D.L., Hickey, C.A., Horvath, S.C., Martens, E.C., Stappenbeck, T.S., and Allen, P.M. (2019). Diet modulates colonic T cell responses by regulating the expression of a Bacteroides thetaiotaomicron antigen. Sci Immunol *4*. 10.1126/sciimmunol.aau9079.
36. Furusawa, Y., Obata, Y., Fukuda, S., Endo, T.A., Nakato, G., Takahashi, D., Nakanishi, Y., Uetake, C., Kato, K., Kato, T., et al. (2013). Commensal microbe-derived butyrate induces the differentiation of colonic regulatory T cells. Nature *504*, 446-450. 10.1038/nature12721.
37. Smith, P.M., Howitt, M.R., Panikov, N., Michaud, M., Gallini, C.A., Bohlooly, Y.M., Glickman, J.N., and Garrett, W.S. (2013). The microbial metabolites, short-chain fatty acids, regulate colonic Treg cell homeostasis. Science *341*, 569-573. 10.1126/science.1241165.
38. Arpaia, N., Campbell, C., Fan, X., Dikiy, S., van der Veeken, J., deRoos, P., Liu, H., Cross, J.R., Pfeffer, K., Coffer, P.J., and Rudensky, A.Y. (2013). Metabolites produced by commensal bacteria promote peripheral regulatory T-cell generation. Nature *504*, 451-455. 10.1038/nature12726.

39. Li, W., Hang, S., Fang, Y., Bae, S., Zhang, Y., Zhang, M., Wang, G., McCurry, M.D., Bae, M., Paik, D., et al. (2021). A bacterial bile acid metabolite modulates Treg activity through the nuclear hormone receptor NR4A1. Cell Host Microbe *29*, 1366-1377 e1369. 10.1016/j.chom.2021.07.013.

40. Paik, D., Yao, L., Zhang, Y., Bae, S., D'Agostino, G.D., Zhang, M., Kim, E., Franzosa, E.A., Avila-Pacheco, J., Bisanz, J.E., et al. (2022). Human gut bacteria produce TauEta17-modulating bile acid metabolites. Nature *603*, 907-912. 10.1038/s41586-022-04480-z.

41. Song, X., Sun, X., Oh, S.F., Wu, M., Zhang, Y., Zheng, W., Geva-Zatorsky, N., Jupp, R., Mathis, D., Benoist, C., and Kasper, D.L. (2020). Microbial bile acid metabolites modulate gut RORγ(+) regulatory T cell homeostasis. Nature *577*, 410-415. 10.1038/s41586-019-1865-0.

42. Wang, Q., McLoughlin, R.M., Cobb, B.A., Charrel-Dennis, M., Zaleski, K.J., Golenbock, D., Tzianabos, A.O., and Kasper, D.L. (2006). A bacterial carbohydrate links innate and adaptive responses through Toll-like receptor 2. J Exp Med *203*, 2853-2863. 10.1084/jem.20062008.

43. Hickey, C.A., Kuhn, K.A., Donermeyer, D.L., Porter, N.T., Jin, C., Cameron, E.A., Jung, H., Kaiko, G.E., Wegorzewska, M., Malvin, N.P., et al. (2015). Colitogenic Bacteroides thetaiotaomicron Antigens Access Host Immune Cells in a Sulfatase-Dependent Manner via Outer Membrane Vesicles. Cell Host Microbe *17*, 672-680. 10.1016/j.chom.2015.04.002.

44. Prados-Rosales, R., Baena, A., Martinez, L.R., Luque-Garcia, J., Kalscheuer, R., Veeraraghavan, U., Camara, C., Nosanchuk, J.D., Besra, G.S., Chen, B., et al. (2011). Mycobacteria release active membrane vesicles that modulate immune responses in a TLR2-dependent manner in mice. J Clin Invest *121*, 1471-1483. 10.1172/jci44261.

45. Jan, A.T. (2017). Outer Membrane Vesicles (OMVs) of Gram-negative Bacteria: A Perspective Update. Front Microbiol *8*, 1053. 10.3389/fmicb.2017.01053.

46. Roier, S., Blume, T., Klug, L., Wagner, G.E., Elhenawy, W., Zangger, K., Prassl, R., Reidl, J., Daum, G., Feldman, M.F., and Schild, S. (2015). A basis for vaccine development: Comparative characterization of Haemophilus influenzae outer membrane vesicles. Int J Med Microbiol *305*, 298-309. 10.1016/j.ijmm.2014.12.005.

47. Roier, S., Zingl, F.G., Cakar, F., Durakovic, S., Kohl, P., Eichmann, T.O., Klug, L., Gadermaier, B., Weinzerl, K., Prassl, R., et al. (2016). A novel mechanism for the biogenesis of outer membrane vesicles in Gram-negative bacteria. Nat Commun *7*, 10515. 10.1038/ncomms10515.

48. Hayashi, H., Sakamoto, M., and Benno, Y. (2002). Phylogenetic analysis of the human gut microbiota using 16S rDNA clone libraries and strictly anaerobic culture-based methods. Microbiol Immunol *46*, 535-548. 10.1111/j.1348-0421.2002.tb02731.x.

49. Hold, G.L., Pryde, S.E., Russell, V.J., Furrie, E., and Flint, H.J. (2002). Assessment of microbial diversity in human colonic samples by 16S rDNA sequence analysis. FEMS Microbiol Ecol *39*, 33-39. 10.1111/j.1574-6941.2002.tb00904.x.

50. Moore, W.E., and Holdeman, L.V. (1974). Human fecal flora: the normal flora of 20 Japanese-Hawaiians. Appl Microbiol *27*, 961-979. 10.1128/am.27.5.961-979.1974.

51. Durant, L., Stentz, R., Noble, A., Brooks, J., Gicheva, N., Reddi, D., O'Connor, M.J., Hoyles, L., McCartney, A.L., Man, R., et al. (2020). Bacteroides thetaiotaomicron-derived outer membrane vesicles promote regulatory dendritic cell responses in health but not in inflammatory bowel disease. Microbiome *8*, 88. 10.1186/s40168-020-00868-z.

52. Chua, K.J., Kwok, W.C., Aggarwal, N., Sun, T., and Chang, M.W. (2017). Designer probiotics for the prevention and treatment of human diseases. Curr Opin Chem Biol *40*, 8-16. 10.1016/j.cbpa.2017.04.011.

53. Ito, T., Gallegos, R., Matano, L.M., Butler, N.L., Hantman, N., Kaili, M., Coyne, M.J., Comstock, L.E., Malamy, M.H., and Barquera, B. (2020). Genetic and Biochemical Analysis of Anaerobic Respiration in Bacteroides fragilis and Its Importance In Vivo. mBio *11*. 10.1128/mBio.03238-19.

54. Reyes-Prieto, A., Barquera, B., and Juárez, O. (2014). Origin and evolution of the sodium -pumping NADH: ubiquinone oxidoreductase. PLoS One *9*, e96696. 10.1371/journal.pone.0096696.

55. Barquera, B. (2014). The sodium pumping NADH:quinone oxidoreductase (Na^+-NQR), a unique redox-driven ion pump. J Bioenerg Biomembr *46*, 289-298. 10.1007/s10863-014-9565-9.

56. Kishimoto, T., and Kang, S. (2022). IL-6 Revisited: From Rheumatoid Arthritis to CAR T Cell Therapy and COVID-19. Annu Rev Immunol *40*, 323-348. 10.1146/annurev-immunol-101220-023458.

57. Ohnmacht, C., Park, J.H., Cording, S., Wing, J.B., Atarashi, K., Obata, Y., Gaboriau-Routhiau, V., Marques, R., Dulauroy, S., Fedoseeva, M., et al. (2015). MUCOSAL IMMUNOLOGY. The microbiota regulates type 2 immunity through RORγt+ T cells. Science *349*, 989-993. 10.1126/science.aac4263.

58. Yissachar, N., Zhou, Y., Ung, L., Lai, N.Y., Mohan, J.F., Ehrlicher, A., Weitz, D.A., Kasper, D.L., Chiu, I.M., Mathis, D., and Benoist, C. (2017). An Intestinal Organ Culture System Uncovers a Role for the Nervous System in Microbe-Immune Crosstalk. Cell *168*, 1135-1148.e1112. 10.1016/j.cell.2017.02.009.

59. Spindler, M.P., Siu, S.S., Mogno, I., Li, Z., Yang, C., Mehandru, S., Britton, G.J., and Faith, J.J. (2022). Human gut microbiota stimulate defined innate immune responses that vary from phylum to strain. Cell Host Microbe. 10.1016/j.chom.2022.08.009.

60. Goodridge, H.S., and Underhill, D.M. (2008). Fungal Recognition by TLR2 and Dectin-1. Handb Exp Pharmacol, 87-109. 10.1007/978-3-540-72167-3_5.

61. Rogers, T.E., Pudlo, N.A., Koropatkin, N.M., Bell, J.S., Moya Balasch, M., Jasker, K., and Martens, E.C. (2013). Dynamic responses of Bacteroides thetaiotaomicron during growth on glycan mixtures. Mol Microbiol *88*, 876-890. 10.1111/mmi.12228.

62. Hsieh, S., Porter, N.T., Donermeyer, D.L., Horvath, S., Strout, G., Saunders, B.T., Zhang, N., Zinselmeyer, B., Martens, E.C., Stappenbeck, T.S., and Allen, P.M. (2020). Polysaccharide Capsules Equip the Human Symbiont Bacteroides thetaiotaomicron to Modulate Immune Responses to a Dominant Antigen in the Intestine. J Immunol *204*, 1035-1046. 10.4049/jimmunol.1901206.

63. Takeda, K., Kaisho, T., and Akira, S. (2003). Toll-like receptors. Annu Rev Immunol *21*, 335-376. 10.1146/annurev.immunol.21.120601.141126.

64. Gambhir, V., Yildiz, C., Mulder, R., Siddiqui, S., Guzzo, C., Szewczuk, M., Gee, K., and Basta, S. (2012). The TLR2 agonists lipoteichoic acid and Pam3CSK4 induce greater pro-inflammatory responses than inactivated Mycobacterium butyricum. Cell Immunol *280*, 101-107. 10.1016/j.cellimm.2012.12.001.

65. Jin, M.S., Kim, S.E., Heo, J.Y., Lee, M.E., Kim, H.M., Paik, S.G., Lee, H., and Lee, J.O. (2007). Crystal structure of the TLR1-TLR2 heterodimer induced by binding of a tri-acylated lipopeptide. Cell *130*, 1071-1082. 10.1016/j.cell.2007.09.008.

66. Bitto, N.J., Zavan, L., Johnston, E.L., Stinear, T.P., Hill, A.F., and Kaparakis-Liaskos, M. (2021). Considerations for the Analysis of Bacterial Membrane Vesicles: Methods of Vesicle Production and Quantification Can Influence Biological and Experimental Outcomes. Microbiol Spectr *9*, e0127321. 10.1128/Spectrum.01273-21.

67. Charlton, B., Auchincloss, H., Jr., and Fathman, C.G. (1994). Mechanisms of transplantation tolerance. Annu Rev Immunol *12*, 707-734. 10.1146/annurev.iy.12.040194.003423.

68. Nossal, G.J. (1983). Cellular mechanisms of immunologic tolerance. Annu Rev Immunol *1*, 33-62. 10.1146/annurev.iy.01.040183.000341.

69. Landers, C.J., Cohavy, O., Misra, R., Yang, H., Lin, Y.C., Braun, J., and Targan, S.R. (2002). Selected loss of tolerance evidenced by Crohn's disease-associated immune responses to auto- and microbial antigens. Gastroenterology *123*, 689-699. 10.1053/gast.2002.35379.

70. Matzinger, P. (1994). Tolerance, danger, and the extended family. Annu Rev Immunol *12*, 991-1045. 10.1146/annurev.iy.12.040194.005015.

71. Erttmann, S.F., Swacha, P., Aung, K.M., Brindefalk, B., Jiang, H., Härtlova, A., Uhlin, B.E., Wai, S.N., and Gekara, N.O. (2022). The gut microbiota prime systemic antiviral immunity via the cGAS-STING-IFN-I axis. Immunity *55*, 847-861.e810. 10.1016/j.immuni.2022.04.006.

72. Jiang, T.T., Shao, T.Y., Ang, W.X.G., Kinder, J.M., Turner, L.H., Pham, G., Whitt, J., Alenghat, T., and Way, S.S. (2017). Commensal Fungi Recapitulate the Protective Benefits of Intestinal Bacteria. Cell Host Microbe *22*, 809-816.e804. 10.1016/j.chom.2017.10.013.

73. Botos, I., Segal, D.M., and Davies, D.R. (2011). The structural biology of Toll-like receptors. Structure *19*, 447-459. 10.1016/j.str.2011.02.004.

74. Valguarnera, E., Scott, N.E., Azimzadeh, P., and Feldman, M.F. (2018). Surface Exposure and Packing of Lipoproteins into Outer Membrane Vesicles Are Coupled Processes in Bacteroides. mSphere *3*. 10.1128/mSphere.00559-18.

75. Roier, S., Zingl, F.G., Cakar, F., and Schild, S. (2016). Bacterial outer membrane vesicle biogenesis: a new mechanism and its implications. Microb Cell *3*, 257-259. 10.15698/mic2016.06.508.

76. Sartorio, M.G., Pardue, E.J., Feldman, M.F., and Haurat, M.F. (2021). Bacterial Outer Membrane Vesicles: From Discovery to Applications. Annu Rev Microbiol *75*, 609-630. 10.1146/annurev-micro-052821-031444.

77. Sartorio, M.G., Valguarnera, E., Hsu, F.F., and Feldman, M.F. (2022). Lipidomics Analysis of Outer Membrane Vesicles and Elucidation of the Inositol Phosphoceramide Biosynthetic Pathway in Bacteroides thetaiotaomicron. Microbiol Spectr *10*, e0063421. 10.1128/spectrum.00634-21.

78. Wensink, J., and Witholt, B. (1981). Outer-membrane vesicles released by normally growing Escherichia coli contain very little lipoprotein. Eur J Biochem *116*, 331-335. 10.1111/j.1432-1033.1981.tb05338.x.

79. Hoekstra, D., van der Laan, J.W., de Leij, L., and Witholt, B. (1976). Release of outer membrane fragments from normally growing Escherichia coli. Biochim Biophys Acta *455*, 889-899. 10.1016/0005-2736(76)90058-4.

80. Elhenawy, W., Debelyy, M.O., and Feldman, M.F. (2014). Preferential packing of acidic glycosidases and proteases into Bacteroides outer membrane vesicles. mBio *5*, e00909-00914. 10.1128/mBio.00909-14.

81. Maynard, A., Butler, N.L., Ito, T., da Silva, A.J., Murai, M., Chen, T., Koffas, M.A.G., Miyoshi, H., and Barquera, B. (2019). Antibiotic Korormicin A Kills Bacteria by Producing Reactive Oxygen Species. J Bacteriol *201*. 10.1128/jb.00718-18.

82. Koropatkin, N.M., Martens, E.C., Gordon, J.I., and Smith, T.J. (2008). Starch catabolism by a prominent human gut symbiont is directed by the recognition of amylose helices. Structure *16*, 1105-1115. 10.1016/j.str.2008.03.017.

83. Martens, E.C., Chiang, H.C., and Gordon, J.I. (2008). Mucosal glycan foraging enhances fitness and transmission of a saccharolytic human gut bacterial symbiont. Cell Host Microbe *4*, 447-457. 10.1016/j.chom.2008.09.007.

84. Chutkan, H., Macdonald, I., Manning, A., and Kuehn, M.J. (2013). Quantitative and qualitative preparations of bacterial outer membrane vesicles. Methods Mol Biol *966*, 259-272. 10.1007/978-1-62703-245-2_16.

85. Englen, M.D., Valdez, Y.E., Lehnert, N.M., and Lehnert, B.E. (1995). Granulocyte/macrophage colony-stimulating factor is expressed and secreted in cultures of murine L929 cells. J Immunol Methods *184*, 281-283. 10.1016/0022-1759(95)00136-x.

86. Drage, M.G., Pecora, N.D., Hise, A.G., Febbraio, M., Silverstein, R.L., Golenbock, D.T., Boom, W.H., and Harding, C.V. (2009). TLR2 and its co-receptors determine responses of macrophages and dendritic cells to lipoproteins of Mycobacterium tuberculosis. Cell Immunol *258*, 29-37. 10.1016/j.cellimm.2009.03.008.

87. Speer, M.A. (2013). Development of a Genetically Modified Silage Inoculant for the Biological Pretreatment of Lignocellulosic Biomass. Doctor of Philosophy (The Pennsylvania State University).

88. Glowacki, R.W.P., Pudlo, N.A., Tuncil, Y., Luis, A.S., Sajjakulnukit, P., Terekhov, A.I., Lyssiotis, C.A., Hamaker, B.R., and Martens, E.C. (2020). A Ribose-Scavenging System Confers Colonization Fitness on the Human Gut Symbiont Bacteroides thetaiotaomicron in a Diet-Specific Manner. Cell Host Microbe *27*, 79-92.e79. 10.1016/j.chom.2019.11.009.