Transcriptomic responses to multigenerational environmental warming in a cold-adapted

salmonid

Running title: Lake trout transcriptomic responses to warming

Chantelle M. Penney[1], Gary Burness[2], and Chris C. Wilson[3]

[1] Environmental and Life Sciences Graduate Program, Trent University, Peterborough, Ontario,

K9J 7B8

[2] Department of Biology, Trent University, Peterborough, Ontario, K9L 0G2

[3] Aquatic Research and Development Section, Ontario Ministry of Natural Resources and

Forestry, Trent University, Peterborough, Ontario, K9L 0G2

Author for correspondence: Chantelle Penney, Environmental and Life Sciences Graduate

Program, Trent University, Peterborough, Ontario, K9L 0G2, chantellepenney@trentu.ca

Authors' contributions: CMP, GB and CCW conceived the ideas and designed methodology;

CMP collected and analysed the data, and led the writing of the manuscript. All authors

contributed critically to manuscript drafts and gave final approval for publication.

Conflict of interest: The authors have no conflict of interest to declare.

Keywords: acclimation, climate change, RNA sequencing, *Salvelinus namaycush*,

transgenerational plasticity, transcriptomics.

Abstract

Cold-adapted species are particularly threatened by climate change as rates of environmental

warming outpace the ability of many populations adapt. Recent evidence suggest that

transgenerational thermal plasticity may play a role in the response of cold-adapted organisms to

long-term changes in temperature. Using RNA sequencing, we explored differential gene

expression of lake trout (*Salvelinus namaycush*), a cold-adapted species, to examine the

molecular processes that respond to elevated temperatures under conditions of within-generation

(offspring) and transgenerational (parental) warm-acclimation. We hypothesized that genes

associated with metabolism, growth and thermal stress/tolerance would be differentially

expressed in juvenile lake trout offspring depending on their own acclimation temperature and

that of their parents. We found enriched pathways for thermal stress, signaling processes,

immune function, and transcription regulation, although the specific genes and direction of

differential expression (up- or downregulation) depended on the combination of offspring and

parental thermal rearing conditions. While parental warm-acclimation did have a

transgenerational effect on gene expression in their offspring, within-generation warm-

acclimation had a larger effect on differential expression, enriching more pathways. We provide

evidence of the transgenerational response to warming at the transcriptional level in lake trout,

which should be useful for future studies of transcriptomics and plasticity in these and other

cold-adapted species.

Keywords: acclimation, climate change, RNA sequencing, *Salvelinus namaycush*,

transgenerational plasticity, transcriptomics.

Introduction

Temperate and Arctic regions are warming rapidly; surface temperatures in Canada have

increased by ~1.7°C in under a century with even greater warming happening in the Arctic

(~2.3°C; Zhang et al. 2019). As a result, high-latitude, cold-adapted organisms are experiencing,

and will continue to experience, rapid temperature increases that may threaten their survival

(Beitinger and Bennett 2000; Burkhead 2012; IPCC 2021). Range expansion to track suitable

conditions is particularly challenging for species and populations with limited dispersal options,

such as many freshwater fishes (Guzzo and Blanchfield 2017; Smith et al. 2020). Warming rates

may outpace adaptive responses (Visser 2008; Crozier and Hutchings 2014; Comte and Olden

2017), particularly for those with low standing genetic variation (Willi et al. 2006; Wilson 2017).

In the absence of standing genetic variation for coordinated adaptive responses throughout core

metabolic pathways, population responses would be limited to plastic thermal acclimation, but

the low physiological plasticity typical of many cold-adapted species means they likely will be

particularly vulnerable to rapidly changing environments (Somero 2010; Kelly et al. 2014).

Individuals that can withstand warming may have to rely heavily on energetic resources that

would normally support growth or reproduction, thus having potentially negative downstream

effects on the population (Somero 2010; Rosenfeld et al. 2020). A lack of adaptive genetic

variation, low capacity for temperature acclimation, and rapid environmental change are

predicted to result in widespread species extirpations (Bennett et al. 2019; Morash et al. 2021).

Some species may be able to buffer the negative impact of environmental warming

through transgenerational acclimation, also known as transgenerational plasticity (TGP). As the

name suggests, TGP refers to the phenotypic changes that occur over the course of multiple

generations (Jablonka et al. 1992; Bell and Hellmann 2019; Bonduriansky 2021). This involves

non-genetic inheritance of environmentally-driven changes, including parental effects, where the

experiences of one generation are reflected in the morphology, physiology or behaviour of a

subsequent generation, often F1 or F2 (Hellman et al. 2020; Lee et al. 2020; Bonduriansky

2021).

TGP occurs through modulation of gene expression through non-genetic inheritance,

which includes parental effects and the transfer of epigenetic factors such as patterns of DNA

methylation and histone modification, and microRNA activity (Dai et al. 2020; Spadafora 2020;

Venney et al. 2020). TGP can potentially be beneficial when a parent's environment coincides

with what their offspring are expected to experience (Bernatchez 2016; Ashe et al. 2021). For

example, TGP could be beneficial in a population that experiences environmental warming over

multiple generations whereby the offspring are phenotypically "primed" for a warmer

environment (Donelan et al. 2020; McCaw et al. 2020; Venney et al. 2022). In the context of

climate change, this phenotypic priming could buffer the effects of rapid warming, allowing

additional time for evolutionary processes to occur (Bernatchez 2016; Ashe et al. 2021), though

the extent to which TGP can be considered adaptive is arguable (Uller et al. 2013; Sánchez-Tójar

et al. 2020). Nevertheless, TGP with temperature has been identified in a number of species

(Greenspoon and Spencer 2018; Yin et al. 2019). For example, the coral reef fish

Acanthochromis polyacanthus exhibits TGP for increased temperatures by improving its aerobic

scope in a warm environment by increasing maximum metabolic rate or decreasing resting

metabolic rate (Donelson et al. 2012; Bernal et al. 2018). Many of the studies on thermal TGP in

aquatic organisms have used eurythermal, temperate or tropical model species; by contrast, cold-

adapted aquatic vertebrates are relatively understudied but could stand to benefit from TGP

given their vulnerability to environmental warming due to climate change.

Acute and chronic temperature acclimation has obvious impacts on the metabolism of

ectotherms, at both the whole organism and molecular level (Pérez-Ruzafa et al. 2018; Petitjean

et al. 2019; Morash et al. 2021). Relatively few studies to date report the effect of

transgenerational warming on phenotypic variation in tandem with gene expression, or with

follow-up experiments testing the differential gene expression underlying transgenerational

effects (Shama et al. 2016; Bernal et al. 2018; Veilleux et al. 2018). Examining differential gene

expression associated with changes in metabolism with thermal experiences can be informative

(Oomen and Hutchings 2017). Shama et al. (2016), for example, found that genes involved in

metabolism, mitochondrial protein synthesis, hemostasis and apoptosis were differentially

expressed in offspring depending on the multigenerational thermal experiences down the

maternal line. These findings underscore the relationship between the molecular mechanisms

underlying physiological plasticity and parental thermal experiences.

We sought to explore differential gene expression in lake trout (*Salvelinus namaycush*) to

examine the molecular processes potentially underlying transgenerational responses to elevated

temperatures. The lake trout is a long-lived, cold-adapted salmonid that is largely limited to cold,

oligotrophic lakes in formerly glaciated regions of North America (Riley et al. 2021; Wilson and

Mandrak 2021) and is under significant threat from climate change (Casselman 2008; Guzzo and

Blanchfield 2017). Being stenothermal, lake trout behaviourally thermoregulate by migrating to

deeper, cooler water when lakes thermally stratify during the summer (Martin and Olver 1980;

Guzzo and Blanchfield 2017). Low genetic variation for some lake trout populations (Perrier et

al. 2017), and little within- and among-population variation for thermal acclimation capacity

(McDermid et al. 2013; Kelly et al. 2014) may limit the adaptive potential of lake trout

populations for warming temperate and arctic habitats. A previous study investigated the capacity

for transgenerational thermal plasticity in lake trout as a means of coping with warming

associated with climate change and found that they exhibit only limited transgenerational

plasticity at the whole organism (phenotypic) level (Penney et al. 2021). Unexpectedly, cold-

acclimated offspring of parents from elevated temperatures exhibited higher resting metabolic

rate than those with one or both cold acclimated parents (Penney et al. 2021). This study

confirmed that the parental environment can influence offspring phenotypic expression, but did

not expand the investigation into the mechanisms underlying this transgenerational response.

For this study, we assessed gene expression in previously studied offspring from factorial

mating crosses between cold- and warm-acclimated lake trout adults (Penney et al. 2021) to

assess transcriptional responses to within- and transgenerational thermal acclimation.

Acclimation of the parents (transgenerational) to one of two environmental temperatures (10 or

17°C) was combined with subsequent acclimation of the offspring (within-generation) to cool

(11°C) or warm (15°C) temperatures (Fig. 1). Evidence exists to support both maternal and

paternal contributions to transgenerational effects (Marshall 2015; Shama et al. 2016; Penney et

al. 2021). Our experimental design allowed us to examine the transgenerational effect of

temperature acclimation of each parent, and both parents together, in combination with the

offspring's (within-generation) temperature acclimation on gene expression in juvenile lake trout.

We hypothesized that genes associated with metabolism, growth and thermal

stress/tolerance would be differentially expressed in juvenile lake trout depending on both the

acclimation temperature of the offspring and the acclimation temperature of their parents. Based

on studies of within-generation and transgenerational warming on differential gene expression in

fish (Quinn et al. 2011; Veilleux et al. 2015; Akbarzadeh and Leder 2016; Shama et al. 2016), we

predicted that 1) genes involved in growth, heat shock and hypoxia responses, and metabolic

pathways would be upregulated in warm-acclimated offspring, and 2) differential expression of

these genes would vary depending on whether one or both parents were warm-acclimated,

although the size and direction of the effect is difficult to predict given that parental influence on

offspring gene expression has been shown to be sex-specific; depending on the sex of the

offspring and that of the parent (Best et al. 2018; Baustista et al. 2020).

Methods

Hatchery-raised adult lake trout from the Ontario provincial Seneca Lake broodstock

were used as parents for the experiment. This hatchery strain originated from Seneca Lake, one

of the Finger Lakes in central New York State (42°41' N, 76°54' W). The Ontario Ministry of

Natural Resources and Forestry (OMNRF) hatchery system has maintained this lake trout strain

for over five generations, using rotational line crossing (Kincaid 1977) to minimize inbreeding

and maintain the founding genetic diversity of the strain (OMNRF Fish Culture Stocks

Catalogue 2005).

The tissue samples used for these experiments were collected from juvenile (young of

year) lake trout that were previously used in a separate study on metabolic rate with

transgenerational warming described in Penney et al. (2021). In this previous study, adult and

juvenile offspring lake trout were acclimated to two different temperatures and the offspring

were subjected to an acute thermal challenge of $+2°C·h^{-1}$ to determine the effect of within- and

transgenerational temperature acclimation on upper thermal tolerance and metabolic rate. Details

of the prior experiment (Penney et al. 2021) that are relevant to the current study are summarized

below.

All experiments adhered to the Canadian Council on Animal Care guidelines and were

approved by the Trent University Animal Care Committee (Protocol # 24794) and the OMNRF

Aquatic Animal Care Committee (Protocol # FACC_136).

Experimental design and rearing conditions

Two groups (n = 8 and 9) of mature lake trout (age 8; 2.3 - 4.2 kg) were each held in

6,000 L tanks (1 x 1 x 6 m), with males and females kept together, at the OMNRF White Lake

Fish Culture Station (Sharbot Lake, Ontario, Canada) with water being supplied from a nearby

lake. As the source waterbody (White Lake) began to stratify in early June, 2015, we began to

increase tank water temperature by 1°C per day starting from approximately 9°C until one tank

reached $10 \pm 0.5°C$ and the other $17 \pm 0.5°C$. These temperatures were chosen because 10°C

represents the thermal requirements for lake trout spawning while 17°C was intended to induce

physiological thermal stress without preventing reproduction (Casselman 2008; Penney et al.

2021) at the higher temperature. Temperatures were controlled and maintained by drawing in and

mixing water from above and below the lake's thermocline as it flowed into the holding tanks.

Fish were acclimated to these temperatures for approximately 3 months, mirroring thermal

stratification in the source waterbody. Starting in mid-September, the temperature in the warm-

water treatment was allowed to gradually cool by holding the proportional inflows from above

and below the thermocline constant as the lake's surface water cooled. Temperatures in the two

tanks converged at 10°C after fall turnover (late September) in the lake, and gradually cooled to

7.8°C by the end of the breeding interval (mid-November).

Adult lake trout were reproductive by the end of October with the warm-acclimated fish

first spawning on October 30[th]. The cold-acclimated adults were first ready to spawn on

November 5[th] and all mating crosses were completed by November 19th. Experimental offspring

families were produced by a full factorial mating cross using two males and two females from

each of the two temperature treatments (4 x 4, 8 adults in total) resulting in a total of 16 offspring

families; four families from each of four parental treatment groups (W♀xW♂, W♀xC♂, C♀xW♂

and C♀xC♂, where W = warm-acclimated, and C = cold-acclimated; Figure 1). Fertilized eggs

were transported to the OMNRF Codrington Fish Research Facility (Codrington, Ontario,

Canada) where they were transferred into 200 L tanks receiving freshwater at ambient

temperature (5-6 °C) and natural photoperiod under dim light with each family separately

contained in steel mesh boxes (9 x 9 x 7.5 cm, one family per box).

In March (2016), when hatched offspring were ready to begin exogenous feeding, we

randomly selected 14 individuals from each of the 16 families to transfer to 200 L tanks for

temperature acclimation. Seven fry from each family were acclimated to a cold temperature

(11°C) and the other 7 from each family were acclimated to a warm temperature (15°C). The

offspring's lower acclimation temperature differs from that of their parents because of reported

differences in optimum temperature for both life stages (11 vs 10°C; Edsall 2000, Casselman

2008). The upper acclimation temperature for the offspring represents the anticipated increase in

temperature associated with climate change by year 2100 (Hayhoe et al. 2010; IPCC 2021),

however, for the parents the upper acclimation temperature was two degrees higher to ensure that

temperature was high enough to elicit a physiological thermal response without compromising

reproduction. To begin acclimating the offspring, water temperature was increased at a rate of

1°C per day until target temperatures were reached (11 and 15°C). Due to space constraints,

tanks were subdivided by adding steel inserts with perforated bottoms. Each insert was divided

into 4 sections (24 x 25 x 28 cm per section) with two paternal half-sibling families (n = 7 each;

shared male parent) housed per section. Offspring from pooled families were later individually

identified to family using microsatellite genotyping (Penney et al. 2021).

After the 3-4 week acclimation period, offspring were subjected to an acute temperature

increase of $+2°C·h^{-1}$ from their acclimation temperature until loss of equilibrium was observed.

Eight fish from the same acclimation temperature (11°C or 15°C) were simultaneously tested per

day, with each fish held in its own experimental chamber (Penney et al. 2021). Following loss of

equilibrium, each fish was quickly transferred to a bath of 0.3 g l−1 of buffered tricaine

methanesulfonate (MS-222; Aqua Life, Syndel Laboratories Ltd, BC, Canada) for euthanasia.

Whole liver was rapidly dissected from each fish, blotted on a lab wipe and preserved in

RNAlater (Invitrogen, Thermo Fisher Scientific) as per manufacturer's instructions. After 24-48

hours in a 4°C refrigerator, the RNAlater was pipetted from the tissues and the samples were

stored at -80°C until RNA extraction. We chose to study gene expression in the liver because it

is a metabolically active tissue involved in a number of physiological processes that respond to

warming (Quinn et al. 2011; Akbarzadeh and Leder, 2016; Dammark et al. 2018). Additionally,

transcriptional responses to multigenerational warming have been observed in the livers of

zebrafish: Luu et al. (2021). Therefore, the liver ought to provide insight into the metabolic

pathways that are affected by transgenerational temperature acclimation in lake trout. Liver

tissue was collected between June 28 and August 9 in 2016.

RNA isolation and sequencing

RNA was extracted from the preserved liver tissues using a phenol-chloroform extraction

method (Chomczynski and Sacchi 2006) after the tissues had been individually homogenized via

a FastPrep-24 BeadBeater (MP Biomedicals) with 2 mL Lysing Matrix D tubes (MP

Biomedicals) and 1 mL of Trizol reagent (Invitrogen, Thermo Fisher Scientific). RNA was

precipitated with RNA precipitation solution (Sambrook and Russel 2001) and isopropanol, and

washed with 75% ethanol. RNA samples were resuspended in nuclease-free water (Thermo

Fisher Scientific). The purity and concentration of the RNA was quantified using a NanoDrop-

8000 spectrophotometer, and RNA quality was assessed by gel electrophoresis of glyoxylated

RNA on a 1.5% agarose gel (Sambrook and Russel 2001).

Two lots of RNA samples were sent for high-throughput sequencing over a two-year

span. In 2018, liver RNA samples from 24 individuals were sent to The Centre for Applied

Genomics (Sick Kids Hospital, Toronto, Ontario, Canada), and in 2020 another 30 liver samples

(individuals) were sent to Genome Quebec (Montreal, Quebec, Canada). Altogether, the 54

samples included tissue from six individuals from each of the seven experimental treatment

groups, however, the control group (11°C-acclimated offspring from cold-acclimated (CxC)

parents) had a total of 12 individuals sequenced: 6 individuals in 2018 and an additional 6 in

2020, so that each treatment group was compared against the control group sequenced in the

same year to eliminate batch effects in the analysis of differential expression (Table 1).

Individuals from each family were represented in their respective treatment group (there were 4

families per offspring-parent treatment group) and because each sequenced group comprised of 6

individuals, each group had 2 representatives from 2 of the 4 families. Overall, this did not lead

to a bias toward any one family being overrepresented in the samples, except for the 2020

control group which had 3 individuals from one family, but this family is not overrepresented in

any of the other groups. Subsequent genetic testing for males and females (Yano et al. 2013)

confirmed that each group also contained members of both sexes (Table 1). Both facilities

assessed the RNA quality via a Bioanalyzer (Agilent Technologies) and all samples passed

quality control (RNA integrity number: ≥ 7.5). cDNA libraries were constructed by enriching

the poly(A) tails of mRNA with oligo dT-beads using NEBNext Ultra II Directional polyA

mRNA Library Prep. In 2018, barcoded libraries were distributed among two and a half lanes

and were sequenced on the Illumina HiSeq2500 instrument producing an average of 28 million

reads per sample (n = 24, paired-end, 2x126 bp). In 2020, libraries were sequenced on the

Illumina HiSeq4000 with barcoded libraries distributed among three lanes producing 36 million

reads per sample (n = 30, paired-end, 2x126 bp).

De novo assembly, annotation and analysis

For our study, we opted to use a de novo transcriptome assembly for the reads rather than

a genome-guided approach. For non-model species with complex genomes, a de novo approach

can often produce better assemblies than a genome-guided approach (DeWoody et al. 2013), and

the de novo approach using Trinity has been successfully used for transcriptome assembly for

salmonids (Carruthers et al. 2018; Narum and Campbell 2015; Nazari et al. 2021). We did not

use the recently published lake trout genome as a reference because the reads generated from this

experiment were also used to develop the lake trout genome (Smith et al. 2022).

RNA reads from both years were assembled and annotated by the Canadian Centre for

Computational Genomics (C3G, Quebec) in 2020. Adapters and reads with a Phred quality score

<30 were trimmed using Trimmomatic (v.0.36), and normalization and *de novo* assembly was

performed using Trinity assembly software (v.2.0.4) following Haas et al. (2013) and Grabherr et

al. (2011). Pre- and post-trimming read counts and Trinity statistics can be found in the

supplementary file (Tables S1 and S2). Functional annotation was performed with the Trinotate

tool (v.2.0.2) and using the TransDecoder pipeline (v.2.0.1) to identify candidate coding regions

and open reading frames. Candidate peptides were searched against known proteins in the Swiss-

Prot (UniProtKB/Swiss-Prot) database via BLASTp (v.2.3.0) and the pfam database. The pfam

database was also used with HMMER (v.3.1b2) to identify protein domains. Signal peptides,

transmembrane regions and ribosomal RNA genes were predicted using the programs signalP

(v.4.1), tmHMM (v.2.0c) and RNAMMER (v.1.2), respectively.

Seven comparisons were made where offspring treatment groups were each compared to

the control group (cold-acclimated offspring with cold-acclimated parents) to observe the effect

that within- and transgenerational plasticity (acclimation) had on gene expression in juvenile lake

trout (Table 1). Transcript abundance was estimated using 'RNA-Seq by Estimation

Maximization' (RSEM; Li and Dewey, 2011) in Trinity with the

'align_and_estimate_abundance.pl' utility. Differential gene expression was analyzed between

the groups of lake trout offspring in R using the limma package (v.3.40.6; Ritchie et al., 2015).

The log2fold change (LFC) was calculated for each gene, representing the magnitude of up- or

downregulation; negative LFC are downregulated genes. A principle component analysis was

performed to explore the DEGs for possible batch effects between the years given the differences

in sequencing depth (28 vs. 36 million reads per sample), and we tested the correlation between

DEGs for each of the pairwise comparisons to check whether application of a batch correction,

using the twelve control group individuals as a shared baseline, changed the level of expression

of DEGs.

From the list of annotated genes, we applied a LFC cutoff of $\leq$ -1 to $\geq$ 1 and used this

limited list of DEGs for an enrichment analysis performed using the Database for Annotation,

Visualization and Integrated Discovery (DAVID) tool (v.6.8) (Huang et al. 2009A; Huang et al.

2009B). The DAVID tool required that a model species be specified for our gene list: we opted

to use *Salmo salar* as DAVID does not yet include *S. namaycush* despite the recent publication

of the *S. namaycush* reference genome (Smith et al. 2022), and *S. salar* had the best coverage for

annotated proteins among orthologous species available in the Swiss-Prot database (determined

via NCBI Eukaryotic genome annotation pipeline, v.8.5). The output from the DAVID tool

mapped genes to functional categories (GO) and metabolic pathways (Kyoto Encyclopedia of

Genes and Genomes: KEGG); we ran our list of DEGs against backgrounds for fish species in

the database (*Oncorhynchus mykiss*, *S. salar*, *Danio rerio*, *Oryzias latipes*), using the DAVID

tool default criterion of reporting functions that are enriched by a minimum count of 2 genes and

p<0.1. In our study, we only discuss DEGs that mapped to GO or KEGG in DAVID tool because

the function of these DEGs have been confirmed in fish species. Tables outlining the enriched

functions are included in the supplemental material. DEGs that were not included in the

enrichment analysis are only discussed in the terms of the overall size of the effect (i.e. total

number of DEGs, annotated and not annotated) on gene expression under conditions of within-

and.or transgenerational warm-acclimation.

Results

Transcriptome sequencing yielded 52-57 million reads per sample sequenced in 2018,

and 55-104 millions reads per sample sequenced in 2020 (Table S1). Trinity assembled 524,988

transcripts with 261,620 genes for all groups sequenced in 2018 and 413,029 transcripts with

269,149 genes for all groups sequenced in 2020 (Table S2). The transcriptome data showed both

transgenerational and within-generation environmental influences on gene expression. Offspring

within-generation warm-acclimation resulted in 3,853 DEGs with a LFC of $\leq$ -1 to $\geq$ 1 (Fig. 2).

By comparison, transgenerational warm-acclimation resulted in fewer differentially expressed

genes (maternal: 1,958; paternal: 1,777; both parents: 2,244; Fig. 3). The combined effect of

offspring warm-acclimation with that of either or both parents often resulted in more DEGs than

offspring within- or transgenerational warm-acclimation in isolation (offspring and maternal:

4,478; offspring and paternal: 3,638; offspring and both parents: 4,712; Fig. 4). Among all

groups, 1,005 of these DEGs were recognized by the DAVID tool and used for enrichment

analysis. The level of differential expression of these genes are displayed in a heatmap (Fig. 2).

The DEGs in the heatmap are sorted by LFC according to gene expression level in warm-

acclimated offspring to show that any given gene up- or downregulated by offspring warm

acclimation could be expressed differently with the influence of maternal and/or paternal warm

acclimation. The list of genes along with their LFC values and adjusted *p*-value can be found in

the supplementary information (Suppl. Table 17).

Principal component analysis may have detected a batch effect between the two years

with 11% of the variation apparently due to offspring acclimation temperature and 4-6% of the

variation potentially due to different sequencing years (Fig. S1). An analysis of the log2 fold-

changes before and after correction showed that there was a strong correlation across all

comparisons ($r \geq 0.91$), indicating that the batch correction provided essentially the same results

(Fig. S2). Although we must still acknowledge the potential for a batch effect occurring in our

results, we are confident in the trends observed in both years. For example, the within-generation

(offspring) warm-acclimation group had a higher number of DEGs than did the groups that

received either maternal or paternal transgenerational warm-acclimation. These latter two

(parent/transgenerational) groups were sequenced in 2020, meaning that they received the higher

sequencing depth yet still had fewer differentially expressed genes compared to the within-

generation group which was sequenced at a lower depth in 2018.

Within-generation acclimation: offspring warm-acclimation

As predicted, warm-acclimation of lake trout offspring resulted in differential gene

expression compared to cold-acclimated offspring, providing support for our hypothesis that

warm acclimation would upregulate genes associated with metabolism, growth and thermal

tolerance. Our enrichment analysis determined that 29 of these upregulated DEGs were involved

in signalling processes, 14 related to cytoplasm functions and 9 related to immunoglobulin

(Suppl. Table 3). There were 4 genes associated with methyltransferase activity, though this

function was not significantly enriched (p=0.09; Suppl. Table 3). Upregulated DEGs that

mapped to metabolic (KEGG) pathways included those involved in degradation (*HMGCLL1*,

CNOT1) such as peroxisome activity and RNA and ketone degradation (Suppl. Table 4). Genes

involved in the heat shock response were also upregulated, specifically *hsp90aa1.1* (Suppl. Table

4). Others included cell adhesion (*CD99L2, CTNNA2*) and signalling (*JAK1, PYCARD, CHD8*;

Suppl. Table 4). There were 19 downregulated DEGs related to metal- and heme-binding and 3

DEGs associated with cyclin activity (Suppl. Table 3). Contrary to our predictions,

downregulated DEGs involved in KEGG pathways included glycolysis/gluconeogenesis

(*GAPDH*) and metabolism of fructose and mannose (*ALDOB*; Suppl. Table 4). Other DEGs

involved in peroxidase activity were also downregulated with offspring warm-acclimation

(*MPV17L2*; Suppl. Table 4).

Transgenerational acclimation: parental warm-acclimation

Parental warm-acclimation influenced gene expression in cold-acclimated offspring,

however, transgenerational warm-acclimation (either or both parents) affected fewer genes than

did within-generation warm-acclimation (Fig. 3). The direction of differential expression (up- or

down-regulation) in the offspring depended on which parent was warm-acclimated and

sometimes expression of a gene due to maternal or paternal warm-acclimation contrasted with

the expression level when only the offspring were warm-acclimated (positive vs. negative LFC;

Fig. 2). Nevertheless, parental warm-acclimation influenced differential gene expression in the

lake trout offspring, lending support to our prediction that differential expression of genes would

vary in offspring depending on parental temperature acclimation.

Maternal warm-acclimation had the effect of upregulating processes related to muscle

function in their offspring, specifically 2 DEGs involved in calcium-binding and 3 DEGs in

troponin synthesis (Suppl. Table 5). There were 13 upregulated DEGs associated with

transferases and 3 DEGs related to glycolysis (Suppl. Table 5). Upregulated genes involved in

KEGG pathways included DNA damage/repair and fatty acid metabolism (*DDIT4*, *hsd17b12a*;

Suppl. Table 6). There were 14 downregulated DEGs involved in cell signalling and 10 involved

in hydrolase/protease activity (Suppl. Table 5). The KEGG pathway analysis revealed that

maternal warm-acclimation downregulated *GAPDH* involved in glycolysis/gluconeogenesis, and

functions related to RNA transport and degradation (*CNOT1* and *EIF3C*; Suppl. Table 6).

Paternal warm-acclimation also had an effect on offspring gene expression, providing

further support for our hypothesis regarding a transgenerational of warming on gene expression,

although fewer genes were differentially expressed with paternal warming compared to maternal

warming (Fig. 3). There were 10 upregulated DEGs related to transferase activity along with 3

DEGs involved in glycolytic processes (Suppl. Table 7). Upregulated KEGG pathways included

the mTOR signalling pathway (*DDIT4*) and protein processing (*ERO1A*; Suppl. Table 8).

Downregulated DEGs included 16 DEGs involved in cell-signalling and 7 DEGs associated with

protease activity (Suppl. Table 7). The KEGG pathway analysis revealed there were two

transcripts (one upregulated and the other downregulated), that were identified as *GAPDH* and

mapped to glycolysis (Suppl. Table 8). Comparing the adjusted *p*-value of both transcripts,

where a lower *p*-value means it is less likely a false positive (i.e. type I error), suggests that

GAPDH was actually downregulated ($p = 0.88$). Up- and downregulation was also observed with

CNOT1 (Suppl. Table 8), but because the $p = 1$ in both cases, we cannot conclude whether this

DEG was up- or downregulated.

As expected, transgenerational effects were also observed in offspring gene expression

when both mothers and fathers were warm-acclimated. The combined effect of both parents

resulted in more differentially expressed genes in the offspring compared to the transgenerational

effect of warm acclimation of either parent in isolation (Fig. 3). We observed 16 upregulated

DEGs that were associated with cell signalling processes, 9 DEGs related to

hydrolase/proteases/peptidases function and 2 DEGs associated with peroxisome activity (Suppl.

Table 9). Upregulated KEGG pathways included the mTOR pathway (*DDIT4*), RNA

degradation (*CNOT1*) and amino acid transferase activity (*BHMT*, *GATM*, *GAMT*; Suppl. Table

10). Downregulated functions involved 4 DEGs included in glycosyltransferase activity, 3 DEGs

related to lectin and 2 DEGs associated with tRNA function (Suppl. Table 9). There were 7

downregulated DEGs associated with translation/ribosomal proteins, but this function was not

significantly enriched ($p=0.09$; Suppl. Table 9). Downregulated KEGG pathways included

peroxisome activity (*HMGCLL1*) and apoptotic signaling (*pycard*) and Notch signaling pathways

(*APH1B*; Suppl. Table 10). Again, *GAPDH* appeared as both up- and downregulated in the

KEGG analysis.

Multigenerational acclimation: offspring and parental warm-acclimation

As hypothesized, genes associated with metabolism, growth and thermal stress/tolerance

were differentially expressed with the combined effect of offspring and parent warm acclimation.

The combination of offspring and parent (both together) warm-acclimation resulted in more

DEGs (e.g. >200) being up- or downregulated relative to the offspring group that experienced

only within-generation warm-acclimation (Fig. 4). We must acknowledge here that the groups

that represent the combined effect of offspring and maternal warm-acclimation, and the

combined effect of offspring and paternal warm-acclimation were sequenced at a higher read

depth than the within-generation only group (36 vs 28 million reads, respectively), meaning that

the differences in differential expression here may be partly due to an experimental artefact of

sequencing depth. For example, a total of 4,478 transcripts were differentially expressed in the

offspring that received the combined treatment of within- and maternal transgenerational warm-

acclimation compared to the 3,853 transcripts of the group that received the within-generation

warm-acclimation only (Fig. 4).

The combined effect of offspring and maternal warm-acclimation resulted in the

upregulation of 20 DEGs related transferase function, 4 DEGs involved in

metalloprotease/endopeptidase activity and 4 associated with the PDZ domain (Suppl. Table 11).

There were also 3 DEGs associated with protein folding, 3 DEGs associated with DNA repair,

and 3 associated with muscle contraction/ZASP (Suppl. Table 11). Genes that encode heat shock

proteins (*HSP70*, *HSPB1*, *HSP7C*, *HSP30*) and proteins involved in the respiratory chain (*COX2*,

COX1) were also upregulated, though these functions were not significantly enriched (p=0.07;

Suppl. Table 11). Similarly, upregulated KEGG pathways included heat shock responses and

RNA degradation (Suppl. Table 12). Downregulated functions included 21 DEGs with

transferase functions, 20 with function related to the cell nucleus and 19 DEGs involved in ATP

binding (Suppl. Table 11). Additionally, there were 12 DEGs associated with kinases, 5 that had

cyclin functions and 5 associated with DNA replication (Suppl. Table 11). Downregulated

KEGG pathways included glycolysis/gluconeogenesis (*ALDOB*, *GAPDH*; Suppl. Table 12).

Offspring and paternal warm-acclimation combined to influence the differential

expression of similar numbers and sorts of pathways as did maternal warm-acclimation, again

supporting our hypothesis. This combination of offspring and paternal warm-acclimation had the

effect of upregulating 4 DEGs that have proteases/peptidases functions and 2 DEGs involved in

glycolysis (Suppl. Table 13). There were 2 upregulated DEGs involved in complex IV of the

respiratory chain, however, this function was not significantly enriched (p=0.07; Suppl. Table

13). Similar to the combined effect of offspring and maternal warm-acclimation, offspring and

paternal warm-acclimation upregulated KEGG pathways involved in cell signaling/inflammation

(*mapk14a*), but offspring and paternal warm-acclimation uniquely upregulated KEGG pathways

associated with protein folding (*HYOU1*) and insulin regulation (*foxo1a*; Suppl. Table 14).

Enriched downregulated functions included ATP binding (14 DEGs), hydrolase/protease activity

(13 DEGs), DNA replication (4 DEGs) and cyclin function (4 DEGs; Suppl. Table 13). There

were also 2 DEGs associated with tRNA ligase activity and 2 DEGs related to peroxisome

function (Suppl. Table 13). Downregulated KEGG pathways included glycolysis (*ALDOB*,

GAPDH), amino acid metabolism (*hpda*, *ALDH4A1, RRM1, RRM2, tdo2a*), fatty acid

metabolism (*ELOVL6*; Suppl. Table 14).

Of all groups compared so far, the combined multigenerational effect of offspring,

maternal and paternal warm-acclimation had the greatest effect on differential gene expression in

the offspring (Fig. 4). There were 41 of upregulated DEGs associated with signalling function

(Suppl. Table 15). Immunoglobulin and protease/peptidase activity were also enriched (9 and 5

DEGs, respectively; Suppl. Table 15). There were 2 DEGs involved in the respiratory chain

complex IV but complex IV was not significantly enriched (p=0.07; Suppl. Table 15). Some of

the affected KEGG pathways included the heat shock response (*hsp90aa1.1*), signalling

pathways associated with inflammation (*mapk14a*) and insulin regulation (*foxo1a*; Suppl. Table

16). Downregulated functions included transferase activity (24 DEGs), DNA replication (6

DEGs) and cyclin functions (4 DEGs; Suppl. Table 15). Binding functions were downregulated,

such as those related to magnesium, nucleic acid and iron binding (5 DEGs each; Suppl. Table

15). Downregulated DEGs involved in KEGG pathways included peroxisome activity

(*MPV17L2*), glycolysis/gluconeogenesis (*ALDOB*), and response to DNA damage (*DDIT4*;

Suppl. Table 16).

Discussion

Our hypothesis that genes associated with metabolism, growth and thermal

stress/tolerance would be differentially expressed in juvenile lake trout depending on the

acclimation temperatures of the offspring as well as their parents was supported by the sequence

data. Genes involved in growth, heat shock and hypoxia responses, and metabolic pathways were

differentially expressed with within-generation, transgenerational and multigenerational warm-

acclimation. The degree to which each gene was affected depended on whether it was the

offspring or parent that had been warm-acclimated, with the offspring thermal experience often

having a stronger effect compared to the parental thermal experience. Maternal and paternal

warm-acclimation sometimes had opposing effects on the direction of differential expression of

certain genes in their offspring with one parent stimulating gene upregulation and the other

parent stimulating downregulation.

We identified five noteworthy trends when considering the experimental results all

together: 1) Within-generation (offspring) warm-acclimation had a larger effect on the overall

number of mapped DEGs compared to transgenerational contributions from warm-acclimated

parents (mother, father or both). Unsurprisingly, the combination of offspring, maternal and

paternal warm-acclimation had the greatest effect on the level of differential expression in the

offspring with an overall 471 mapped DEGs. 2) Cell signalling processes were upregulated (29

DEGs) with offspring (within-generation) warm-acclimation, but maternal or paternal

(transgenerational) warm acclimation had the effect of downregulating cell-signalling functions

(14-16 DEGs). When both parents were warm-acclimated, cell signalling was upregulated (16

DEGs). Cell signalling was most enriched (41 DEGs) with multigenerational (offspring,

maternal and paternal) warm-acclimation. 3) Transferase functions appeared to be regulated

mainly by transgenerational (parental) warm acclimation rather than with within-generation

(offspring) warm-acclimation, however, the degree of enrichment (i.e. gene count) and the

direction of the effect (up- or downregulation) appeared to depend upon whether one or both

parents were warm-acclimated. 4) Regulation of cyclin functions appeared to be primarily under

the control of the offspring because this process was only downregulated when offspring were

warm-acclimated, thus not influenced by transgenerational (either parent or both) warm-

acclimation. 5) Mothers seemed to have an influence over muscle functions (calcium binding,

troponin synthesis, ZASP) in their offspring, upregulating these functions when mothers

experienced a warmer environment, whereas there was no evidence of paternal influence on

offspring muscle functions.

General effect of warming on differential gene expression

Several DEGs that were associated with thermal stress were apparent across all

comparisons. Among these were upregulated DEGs associated with molecular chaperone and co-

chaperone functions (*hsp90aa1.1, hsp70, hsp30, hsp7c, hspB1, hspB7, HYOU1, FKBP5* and

FKBP8). Upregulation of heat shock proteins and FK506-binding proteins (HSP90a, HSP70,

FKBP10) has been observed in other salmonids experiencing warm acclimation and with an

528 acute temperature increase (Akbarzadeh et al. 2018; Houde et al. 2019; Shi et al. 2019). Another

529 common indicator of heat stress in salmonids is hsp47, also known as serpinh1 (Tomalty et al.

530 2015; Akbarzadeh et al. 2018). We did not see differential expression of serpinh1 in our study,

531 however, upregulation of this gene does not always occur with heat stress (Mackey et al. 2021).

532 *HYOU1* in particular was upregulated with within- or multigenerational warm-acclimation, but

533 not with transgenerational warm-acclimation in isolation (Suppl. Table 17). *HYOU1* (also known

534 as *GRP170* and *HSP12A*) is part of the heat shock protein 70 family and its expression is

535 primarily induced with hypoxia. Although, upregulation of this gene has previously been

536 observed in response to heat stress in fish (Huang et al. 2018; Bilyk et al. 2021), its expression

537 may be linked to a hypoxia response or may potentially reflect the increased demands on aerobic

538 metabolism/oxidative stress (Olsvik et al. 2013; Huang et al. 2018). The temperature-controlled

539 water in the source tank supplying the respirometer setup was supplemented with compressed

540 oxygen at higher temperatures in an effort to maintain saturation levels and minimize exposure to

541 hypoxia. However, we must acknowledge the possibility that some hypoxic responses may still

542 have occurred.

543   In general, within- and transgenerational warm-acclimation resulted in upregulated

544 mitogen-activated protein kinases (MAPK): *MAPK3*, *MAPK14a* and *MAPK14b* (also known as

545 *ERK1*, *p38a* and *p38b*, respectively). MAPKs are important in the regulation of many processes,

546 including mitosis during growth and development. They are also upregulated in response to

547 various stressors, such as osmotic, hypoxic and thermal stress (Cowan and Storey 2003; Tian et

548 al. 2019), for example, upregulation of these proteins has been observed in Atlantic salmon and

549 *Oncorhynchus* spp. with thermal stress (Akbarzadeh et al. 2018; Shi et al. 2019). MAPKs are

550 associated with the induction of HSPs, for example, *MAPK3* has been shown to induce

551 expression of *HSP70* in zebrafish under thermal stress (Keller et al. 2008). Both *MAPK3* and

552 *HSP70* were upregulated in warm-acclimated lake trout offspring with warm-acclimated

553 mothers, suggesting that these fish relied on MAPK signaling cascades for resilience to warming.

554 This is perhaps not surprising given that MAPK cascades and the induction of HSPs are

555 evolutionarily conserved across numerous taxa (Krens et al. 2006; Kostenko et al. 2011), and

556 that upregulation of these genes has been observed previously in other taxa following heat stress

557 (Keller et al. 2008; Liu et al. 2021). Other affected pathways indicating upper thermal stress

558 included peroxisome activity (*HMGCLL1*), DNA damage (*DDIT4*) and apoptotic signaling

(*pycard*), though whether these DEGs were up- or downregulated depended on whether it was

the parental or offspring generation that had experienced warming. Taken together, these DEGs

indicate increasing temperatures caused a stress response in lake trout, which we would expect

given that each offspring was subjected to an acute temperature increase (+2°C/h) and these

DEGs were likely upregulated to respond to cellular/protein damage occurring at their upper

critical thermal limit (Penney et al. 2021).

There was a general enrichment of pathways associated with metabolism, including genes

such as *COX1*, *COX2, GAPDH* and *ALDOB*. In general, *GAPDH* and *ALDOB* were

downregulated with within- or transgenerational warm acclimation, however, *COX1* and *COX2*

were upregulated in warm-acclimated offspring with warm-acclimated mothers following an

acute temperature challenge. The effect of acute warming on differential expression of

cytochrome c oxidase subunits is equivocal in fish: for example, acute exposure to warmer

temperatures resulted in downregulation of *COX1* and *COX2* in the liver of yellow croaker (Qian

and Xue 2016) but upregulation of *COX1* in the liver of juvenile lake whitefish (Zak and Manzon

2019). For salmonids in particular, upregulation *COX2* also occurred with heat stress in chinook

salmon (*O. tshawytscha*) likely in response to the increased metabolic demands associated with

warming (Tomalty et al. 2015). Upregulation COX6B1 has been shown to occur with heat stress

in Pacific salmon (Akbarzadeh et al. 2018), though we did not detect expression of this subunit

in our study. In our experiment, the matching of offspring and maternal thermal experience

increased COX gene expression to a greater extent compared to other groups, potentially

allowing offspring to meet the energetic demands associated with acute warming. We did not see

upregulation of any other COX subunits, but it seems that expression of COX subunits does not

follow stoichiometric expectations (Bremer and Moyes 2014) which may explain why we

observed upregulation of only two of the 14 subunits that make up the COX holoenzyme.

We found some indication that warm-acclimation and upper thermal stress affected the

growth of the lake trout offspring. We expected to see upregulation of some growth-related

genes given that the warm-acclimated offspring were heavier than the cold-acclimated offspring

at the time of sampling (Penney et al. 2021). Common molecular indicators of growth in fish

include genes encoding growth hormone (*GH*) and insulin-like growth factor-1 (*IGF-1*), and

genes related to thyroid function (Hevrøy et al. 2015; Li et al. 2021). We did not observe

differential expression of *IGF-1* in the liver of our lake trout, but the thyroid receptors *TRα* and

TRβ were downregulated in offspring that experienced warm-acclimation. Thyroid hormones

have important roles in growth and development (reviewed by Deal and Volkoff 2020), however,

it is curious that *TRα* and *TRβ* were downregulated given we showed previously that warm-

acclimated lake trout grew larger than the cold-acclimated group (Penney et al. 2021). Indicators

of growth, measured as expression of *GH* and *IGF-1*, increased in *Labeo rohita* at slightly

warmer acclimation temperatures but was suppressed at higher acclimation temperatures

(Shahjahan et al. 2021). We collected liver samples at each fish's CTMax, so it is possible that

fish were responding to the stress of the acute temperature increase and may have suppressed

growth at this point in favour of a heat stress response. Lastly, there is evidence suggesting that

growth hormone signaling pathways in the muscle are more sensitive to elevated temperature

than in the liver of Atlantic salmon and rainbow trout (Hevrøy et al. 2015), thus it is possible that

we may have seen a stronger transcriptional response for growth pathways in the muscle

compared to the liver. Confirmation of this would require further experimentation.

Transcriptomic responses to warm-acclimation across generations

We found that an offspring's thermal acclimation often had a greater influence on the

level (LFC) of gene expression than did the thermal experience of its parents and in some cases

multigenerational (offspring and parental) warm acclimation had an additive effect. For example,

cold-acclimated offspring downregulated the molecular chaperone *HYOU1* when either parent

was warm-acclimated (LFC maternal: -1.21, paternal: -0.94; Suppl. Table 17), however, a

combination of offspring and parental warm-acclimation upregulated *HYOU1* (LFC offspring &

maternal: 0.72, offspring & paternal 0.80, offspring & both parents 1.14), though to a lesser

extent than when offspring were warm-acclimated in isolation (LFC 1.29; Suppl. Table 17). This

trend suggests that within- and transgenerational warm acclimation had an additive effect on the

expression level of *HYOU1*. The gene *MAPK14a* was upregulated whenever the offspring were

warm-acclimated, regardless of parental thermal experience (LFC ranged from 1.19 to 2.06) and

seemed to override the isolated effect of transgenerational (parental) warm-acclimation (LFC

ranged from -0.67 to 0.52; Suppl. Table 17). The idea of WGP overriding TGP has been

presented before (Shama 2017), and our experiment provides evidence of this phenomenon at the

transcriptomics level. It has been suggested that transgenerational effects are strongest in early-

juvenile life stages (Yin et al. 2019) and it would be interesting to see if this is reflected at the

transcriptomics level by looking at gene expression at different life stages following within- and

transgenerational warm-acclimation.

Compared to within-generation (offspring) warm-acclimation, transgenerational warm-

acclimation (parents only) had a smaller effect on differential gene expression in lake trout

offspring, supporting our previous study of whole animal metabolic responses to acute warming

(Penney et al. 2021). Maternal and paternal warm-acclimation separately elevated DEGs

functioning in the DNA damage response (*DDIT4*) and heat shock proteins (maternal only:

HSPB1, *HSPB7*). These transgenerationally elevated pathways associated with DNA damage and

HSPs support the idea of an adaptive response whereby parents may improve offspring fitness

when they correctly anticipate their offspring's environment (Bateson et al. 2014; Bernatchez

2016; Ashe et al. 2021).

Our prediction that transgenerational warm acclimation would upregulate metabolic

processes in the lake trout offspring was partially supported. We observed differential expression

of the genes GAPDH, ALDOB and ENOA which relate to the glycolytic enzymes

glyceraldehyde-3-phosphate dehydrogenase, aldolase and enolase, respectively. These three

transcripts tended to be downregulated in all groups except for the two groups that included the

combined effect of offspring cold acclimation and paternal warm acclimation which indicates a

paternal transgenerational effect of increasing offspring metabolism (Suppl. Table 17). Despite

this evidence, we cannot be fully confident that glycolysis was upregulated in these offspring

because, while the differential expression of GAPDH, ALDOB and ENOA suggests an effect of

warm acclimation on the rate of glycolysis in the offspring, these three transcripts did not

significantly enrich the 'glycolysis' pathway in our analysis. While the expression of these genes

may vary depending on species, tissue type and treatment, the enzymes these genes encode are

not considered rate-limiting enzymes, and these genes have even been suggested as

housekeeping genes for experiments exploring differential gene expression (Hori et al. 2012;

Purohit et al. 2016; Shekh et al. 2017).

Although transgenerational warming can result in larger offspring (Salinas et al. 2012),

we did not find support for a transgenerational effect of warm-acclimation on growth in lake

trout at the transcriptome level, consistent with a lack of response at the whole animal level

(Penney et al. 2021). Lastly, despite some genes being transgenerationally expressed in the same

direction by either the maternal or paternal warm-acclimation, none of these genes appeared to

have an additive or synergistic effect on expression level when both parents were simultaneously

warm-acclimated (Suppl. Table 14).

The combination of offspring, maternal and paternal warm-acclimation had the greatest

effect on the overall number genes that were differentially expressed and, for some genes, the

greatest effect on up- or downregulation. This adds to a growing literature suggesting that

transgenerational plasticity occurs when offspring and parental thermal experiences are

coincident (Shama et al. 2014; Donelson et al. 2017; Yin et al. 2019). Among all of the groups

compared, the combination of offspring and parent warm-acclimation had the strongest effect on

cytochrome c oxidase subunit upregulation. Differential expression of COX-related genes has

elsewhere been recorded with transgenerational acclimation (Shama et al. 2016; Bernal et al.

2018). Also, the expression level of *COX1* and *COX2* subunits (among 7 of 14 subunits tested)

correlated with cytochrome c oxidase activity in muscle of fish (Bremer and Moyes 2014). Our

result could be indicative of a higher metabolic rate in warm-acclimated offspring with warm-

acclimated parents, although neither resting or peak metabolic rates differed between warm-

acclimated offspring from warm- or cold-acclimated parents (Penney et al. 2021). It is worth

noting, however, that gene expression does not necessarily confer a protein to function given that

post-transcriptional and post-translational activity could also tag a protein for degradation.

There were a few instances where some genes such as *GAPDH* and *CNOT1* appeared to

be simultaneously upregulated and downregulated within treatments, though why this occurred is

not immediately clear. One possible explanation is that the direction of the responses were sex-

specific (Best et al. 2018; Bell and Hellman 2019; Bautista et al. 2020). For example, in

stickleback, mothers can influence the direction (up- or downregulation) of differentially

expressed genes depending on whether the offspring is male or female (Metzger and Schulte

2016). It is possible that these genes were up- and downregulated because we had representatives

of both sexes in each of our groups.

The offspring exhibited a stronger within-generation affect on gene expression

transgenerational effects, with more DEGs in the offspring warm-acclimation treatments than

from parental warm-acclimation. Some evidence suggests that transgenerational warming has a

stronger effect on offspring differential gene expression compared to developmental warming

(Veillleux et al. 2015; Shama et al. 2016). Prior to initiating warm-acclimation, all offspring in

our experiment were reared in the same environment and temperature regime to minimize

differential developmental effects. The influence of thermal acclimation during early life stages

on differential gene expression at subsequent life stages compared to within- and

transgenerational acclimation may be a fruitful area for study.

686   Venney et al. (2022) used a similar breeding design as this study to assess temperature-

related DNA methylation in adult brook trout and their offspring. In contrast to our findings,

Venney et al. (2022) observed a stronger transgenerational effect from adults held at higher

temperatures during gonad development compared with methylation from offspring rearing

temperatures. Despite the apparent contrasts between the results of Venney et al. (2022) and our

study, the findings from the two studies are complementary rather than contradictory. Some

studies on salmonids have shown that changes in global methylation do not appear to influence

differential gene expression (Christensen et al. 2021; Leitwein et al. 2022). DNA methylation

can influence gene expression in an individual and subsequent generations, but gene expression

is also a direct response to physiological, metabolic, and bioenergetic requirements including

acclimation. The greater methylation observed from reproductive adults by Venney at al. (2022)

may reflect the lower temperature tolerances and scope of reproductive adults compared with

other free-swimming life stages in many fish species (Dahlke et al. 2020). The lesser extent of

DNA methylation observed in juvenile brook trout by Venney et al. may also reflect the ability

of subadult brook trout to utilize warmer temperatures than adults (Smith and Ridgway 2019).

By contrast, our study examined levels of gene expression by transcriptome sequencing, and

showed a greater effect of within-generation acclimation on gene expression, paralleling results

from a previous study looking at whole-organism respirometry in lake trout (Penney at al. 2021).

Conclusions

706   Our study confirmed that lake trout exhibit both transgenerational (both maternal and

paternal) and within-generation thermal acclimation, with both levels of acclimation influencing

the strength and direction of differential expression of genes associated with metabolism and

thermal stress/tolerance. Despite this, the limited transgenerational effect on DEGs indicates that

adaptive transgenerational acclimation is unlikely to contribute sufficiently substantive benefits

to enable lake trout to cope with rapidly changing environmental conditions related to global

warming (Penney et al. 2021). Instead, population-level responses to temperature-related stress

will likely be limited to within-generation acclimation (Kelly et al. 2014) and longer-term

adaptive responses. Beyond acclimation, persistence of cold-water species and populations may

be reliant on existing genetic resources to cope with chronic warming (Comte and Olden 2017;

Crozier and Hutchings 2014; Guzzo and Blanchfield 2017), which is particularly daunting for

long-lived species (Willi et al. 2006). It therefore seems likely that lake trout and similarly

vulnerable cold-water species may require increased management and conservation efforts to

ensure their future in a rapidly warming world.

Acknowledgements

The Ontario Ministry of Natural Resources and Forestry (OMNRF) Fish Culture Section

provided adult lake trout, rearing space and logistical support at the White Lake Fish Culture

Station. Bill Sloan, Scott Ferguson, Anne McCarthy (OMNRF) and John Dewart provided

assistance with breeding adult fish; Bill Sloan and Scott Ferguson also provided invaluable

logistical support for all aspects of fish husbandry from care of the juvenile trout to assisting with

experiments at the OMNRF Codrington Fisheries Research Facility. Caleigh Smith (OMNRF)

genotyped adult and juvenile fish for parentage assignment and family identification of juveniles.

Paul Craig and Heather Ikert (University of Waterloo) kindly provided advice and guidance on

the enrichment analysis. Madison Wikston provided advice for figure presentation. Paul Craig

and Graham Scott provided feedback on early drafts of the manuscript. The RNA sequencing

was conducteddone by The Centre for Applied Genomics (Sick Kids Hospital, Toronto, Ontario,

Canada) and Genome Quebec (Montreal, Quebec, Canada). The bioinformatic analyses was

performed by Canadian Center for Computational Genomics (Montreal, Quebec, Canada). The

Canadian Center for Computational Genomics (C3G) is a Genomics Technology Platform (GTP)

supported by the Canadian Government through Genome Canada.

Research funding was provided by the Canada-Ontario Great Lakes Agreement (COA) to

CCW, by the Ontario Federation of Anglers and Hunters (OFAH) to CMP, and a Natural

Sciences and Engineering Research Council (NSERC) Discovery Grant to GB.

Data archiving statement: Data for this study are available at: *to be completed after manuscript*

is accepted for publication.

Literature Cited

Akbarzadeh, A., Leder, E.H. (2016). Acclimation of killifish to thermal extremes of hot spring: transcription of gonadal and liver heat shock genes. *Comparative Biochemistry and Physiology Part A: Molecular & Integrative Physiology*, 191, 89-97.

Akbarzadeh A., Günther O.P., Houde A.L.S., Ming T. J., Jeffries K.M., Hinch S.G., Miller, K.M. (2018). Developing specific molecular biomarkers for thermal stress in salmonids. *BMC Genomics*, 19, 1-28.

Ashe, A., Colot, V., Oldroyd, B.P. (2021). How does epigenetics influence the course of evolution? *Philosophical Transactions of the Royal Society B: Biological Sciences*, 376(1826), 20200111.

Bateson, P., Gluckman, P., Hanson, M. (2014). The biology of developmental plasticity and the Predictive Adaptive Response hypothesis. *The Journal of Physiology*, 592, 2357-2368.

Bautista, N.M., Burggren, W.W. (2019). Parental stressor exposure simultaneously conveys both adaptive and maladaptive larval phenotypes through epigenetic inheritance in the zebrafish (*Danio rerio*). *Journal of Experimental Biology*, 222, jeb208918.

Bautista, N.M., Crespel, A., Crossley, J., Padilla, P., Burggren, W. (2020). Parental transgenerational epigenetic inheritance related to dietary crude oil exposure in *Danio rerio*. *Journal of Experimental Biology*, 223, jeb222224.

Beitinger, T.L., Bennett, W.A. (2000). Quantification of the role of acclimation temperature in temperature tolerance of fishes. *Environmental Biology of Fishes*, 58, 277-288.

Bell, A.M., Hellmann, J. (2019). An integrative framework for understanding the mechanisms and multigenerational consequences of transgenerational plasticity. *Annual Review of Ecology, Evolution and Systemics*, 50, 97–118.

Bennett, S., Duarte, C.M., Marbà, N., Wernberg, T. (2019). Integrating within-species variation

in thermal physiology into climate change ecology. *Philosophical Transactions of the Royal*

Society B: Biological Sciences, 374, 20180550.

Bernal, M.A., Donelson, J.M., Veilleux, H.D., Ryu, T., Munday, P.L., Ravasi, T. (2018).

Phenotypic and molecular consequences of stepwise temperature increase across generations in a

coral reef fish. *Molecular Ecology*, 27, 4516-4528.

Bernatchez, L. (2016). On the maintenance of genetic variation and adaptation to environmental

change: considerations from population genomics in fishes. *Journal of Fish Biology*, 89, 2519-

2556.

Best, C., Ikert, H., Kostyniuk, D. J., Craig, P. M., Navarro-Martin, L., Marandel, L., Mennigen,

J. A. (2018). Epigenetics in teleost fish: from molecular mechanisms to physiological

phenotypes. *Comparative Biochemistry and Physiology Part B: Biochemistry and Molecular*

Biology, 224, 210-244.

Bilyk, K.T., Zhuang, X., Vargas-Chacoff, L., Cheng, C.C. (2021). Evolution of chaperome gene

expression and regulatory elements in the antarctic notothenioid fishes. *Heredity*, 126, 424-441.

Bonduriansky R. (2021). Plasticity across generations. In: D.W. Pfennig (Ed.). Phenotypic

Plasticity & Evolution: Causes, Consequences, Controversies (pp. 327-348). Boca Raton: CRC

Press/Taylor & Francis Group.

Bremer, K., Moyes, C.D. (2014). mRNA degradation: an underestimated factor in steady-state

transcript levels of cytochrome c oxidase subunits? *Journal of Experimental Biology,* 217, 2212-

2220.

Burkhead, N.M. (2012). Extinction rates in North American freshwater fishes, 1900-

2010. *BioScience*, 62: 798-808.

Carruthers, M., Yurchenko, A.A., Augley, J.J., Adams, C.E., Herzyk, P., Elmer, K.R. (2018). De

novo transcriptome assembly, annotation and comparison of four ecological and evolutionary

model salmonid fish species. *BMC genomics*, 19, 1-17.

Casselman, J.M . (2008). Effects of climate and climate change on lake trout populations and

fisheries. In: K.H. Mills, M. Dyck, & L.A. Harwood (Eds.). Proceedings of the Second North

American Lake Trout Symposium (pp. 103-113). Yellowknife, NT, Canada: Canadian Technical

Report of Fisheries and Aquatic Sciences.

Chomczynski, P., Sacchi, N. (2006). The single-step method of RNA isolation by acid

guanidinium thiocyanate–phenol–chloroform extraction: twenty-something years on. *Nature*

Protocols, 1, 581-585.

Christensen, K.A., Le Luyer, J., Chan, M.T., Rondeau, E.B., Koop, B.F., Bernatchez, L., Devlin,

R.H. (2021). Assessing the effects of genotype-by-environment interaction on epigenetic,

transcriptomic, and phenotypic response in a Pacific salmon. *G3 Genes|Genomes|Genetics*, 11,

jkab021.

Comte, L., Olden, J.D. (2017). Climatic vulnerability of the world's freshwater and marine

fishes. *Nature Climate Change*, 7, 718–722.

Cowan, K.J. Storey, K.B. (2003). Mitogen-activated protein kinases: new signaling pathways

functioning in cellular responses to environmental stress. *Journal of Experimental Biology*, 206,

1107-1115.

Crozier, L.G. J.A. Hutchings. (2014). Plastic and evolutionary responses to climate change in

fish. *Evolutionary Applications*, 7, 68-87.

Dai, Z., Ramesh, V., Locasale, J.W. (2020). The evolving metabolic landscape of chromatin

biology and epigenetics. *Nature Reviews Genetics*, 21, 737-753.

Dahlke, F.T., Wohlrab, S., Butzin, M., Pörtner, H.O. (2020). Thermal bottlenecks in the life
cycle define climate vulnerability of fish. *Science*, *369*, 65-70.

Dammark, K. B., Ferchaud, A.L., Hansen, M.M., Sørensen, J.G. (2018). Heat tolerance and gene
expression responses to heat stress in threespine sticklebacks from ecologically divergent
environments. *Journal of Thermal Biology*, 75, 88-96.

Deal, C.K., Volkoff, H. (2020). The role of the thyroid axis in fish. *Frontiers in*
Endocrinology, 11, 596585.

DeWoody, J.A., Abts, K.C., Fahey, A.L., Ji, Y., Kimble, S.J., Marra, N.J., Ayawardena, B.K.,
Willoughby, J.R. (2013). Of contigs and quagmires: next-generation sequencing pitfalls
associated with transcriptomic studies. *Molecular Ecology Resources*, 13, 551-558.

Donelan, S.C, Hellman, J.K., Bell, A.M., Luttbeg, B., Orrock, J.L., Sheriff, M.J., Sih, A. (2020).
Transgenerational plasticity in human-altered environments. *Trends in Ecology & Evolution*, 35,
115-124.

Donelson, J.M., Munday, P.L., McCormick, M.I., Pritcher, C.R. (2012). Rapid transgenerational
acclimation of a tropical reef fish to climate change. *Nature Climate Change*, 2, 30–32.

Donelson, J.M., Sunday, J.M., Figueira, W.F., Gaitán-Espitia, J.D., Hobday, A.J., Johnson, C.R.,
Leis, J.M., Ling, S.D., Marshall, D., Pandolfi, J.M., Pecl, G., Rodgers, G.G., Booth, D.J.,
Munday, P.L. (2019). Understanding interactions between plasticity, adaptation and range shifts
in response to marine environmental change. *Philosophical Transactions of the Royal Society B:*
Biological Sciences, 374, 20180186.

Edsall, T.A., Cleland, J. (2000). Optimum temperature for growth and preferred temperatures of
age-0 lake trout. *North American Journal of Fisheries Management*, 20, 804-809.

Grabherr, M.G., Haas, B.J., Yassour, M., Levin, J.Z., Thompson, D.A., Amit, I., Adiconis, X.,

Fan, L., Raychowdhury, R., Zeng, Q., Chen, Z., Mauceli, E., Hacohen, N., Gnirke, A., Rhind, N.,

di Palma, F., Birren, B.W., Nusbaum, C., Lindblad-Toh, K., Friedman, N., Regev, A. (2011).

Full-length transcriptome assembly from RNA-Seq data without a reference genome. *Nature*

Biotechnology, 29, 644-652.

Greenspoon, P.B., Spencer, H.G. (2018). The evolution of epigenetically mediated adaptive

transgenerational plasticity in a subdivided population. *Evolution*, 72, 2773-2780.

Guzzo, M.M., Blanchfield, P.J. (2017). Climate change alters the quantity and phenology of

habitat for lake trout (*Salvelinus namaycush*) in small Boreal Shield lakes. *Canadian Journal of*

Fisheries and Aquatic Sciences, 74, 871-884.

Haas, B.J., Papanicolaou, A., Yassour, M., Grabherr, M., Blood, P.D., Bowden, J., Couger, M.

B., Eccles, D., Li, B., Lieber, M., MacManes, M.D., Ott, M., Orvis, J., Pochet, N., Strozzi, F.,

Weeks, N., Westerman, R., William, T., Dewey, C.N., Henschel, R., LeDuc, R.D. Friedman, N.,

Regev, A. (2013). De novo transcript sequence reconstruction from RNA-seq using the Trinity

platform for reference generation and analysis. *Nature Protocols*, 8, 1494-1512.

Hayhoe, K., VanDorn, J., Croley II, T., Schlegal, N., Nwuebbles, D. (2010). Regional climate

change projections for Chicago and the US Great Lakes. *Journal of Great Lakes Research*, 36

(Suppl 2), 7-21.

Hellmann, J.K., Carlson, E.R., Bell, A.M. (2020). Sex-specific plasticity across generations II:

Grandpaternal effects are lineage specific and sex specific. *Journal of Animal Ecology*, 89, 2800-

2812.

Hevrøy, E.M., Tipsmark, C.K., Remø, S.C., Hansen, T., Fukuda, M., Torgersen, T., Vikeså, V.,

Olsvik, P.A., Waagbø, R., Shimizu, M. (2015). Role of the GH-IGF-1 system in Atlantic salmon

and rainbow trout postsmolts at elevated water temperature. *Comparative Biochemistry and*

Physiology Part A: Molecular & Integrative Physiology, 188, 127-138.

Hori, T. S., Gamperl, A. K., Afonso, L. O., Johnson, S. C., Hubert, S., Kimball, J., Bowman, S.,

Rise, M. L. (2010). Heat-shock responsive genes identified and validated in Atlantic cod (Gadus

morhua) liver, head kidney and skeletal muscle using genomic techniques. *BMC Genomics,* 11,

1-22.

Houde, A.L.S., Akbarzadeh, A., Günther, O.P., Li, S., Patterson, D.A., Farrell, A.P., Hinch, S.G.,

Miller, K. M. (2019). Salmonid gene expression biomarkers indicative of physiological

responses to changes in salinity and temperature, but not dissolved oxygen. *Journal of*

Experimental Biology, 222, jeb198036.

Huang, D.W., Sherman, B.T., Lempicki, R.A. (2009A). Systematic and integrative analysis of

large gene lists using DAVID Bioinformatics Resources. *Nature Protocols*, 4, 44-57.

Huang, D.W., Sherman B.T., Lempicki, R.A. (2009B). Bioinformatics enrichment tools: paths

toward the comprehensive functional analysis of large gene lists. *Nucleic Acids Research*, 37, 1-

13.

Huang, J., Li, Y., Liu, Z., Kang, Y., Wang, J. (2018). Transcriptomic responses to heat stress in

rainbow trout *Oncorhynchus mykiss* head kidney. *Fish & Shellfish Immunology*, 82, 32-40.

IPCC (2021). Summary for Policymakers. In: V. Masson-Delmotte, P. Zhai, A. Pirani, S.L.

Connors, C. Péan, S. Berger, N. Caud, Y. Chen, L. Goldfarb, M.I. Gomis, M. Huang, K. Leitzell,

E. Lonnoy, J.B.R. Matthews, T.K. Maycock, T. Waterfield, O. Yelekçi, R. Yu, & B. Zhou

(Eds.), Climate Change 2021: The Physical Science Basis. Contribution of Working Group I to

the Sixth Assessment Report of the Intergovernmental Panel on Climate Change (pp. 3−32).

Cambridge, United Kingdom and New York, NY, USA:Cambridge University Press.

Jablonka, E., Lachmann, M., Lamb, M.J. (1992). Evidence, mechanisms and models for the

inheritance of acquired characters. *Journal of Theoretical Biology*, 158, 245-268.

Keller, J.M., Escara-Wilke, J.F., Keller, E.T. (2008). Heat stress-induced heat shock protein 70

expression is dependent on ERK activation in zebrafish (*Danio rerio*) cells. *Comparative*

Biochemistry and Physiology Part A: Molecular & Integrative Physiology, 150, 307-314.

Kelly, N.I., Burness, G., McDermid, J.L., Wilson, C.C. (2014). Ice age fish in a warming world:

minimal variation in thermal acclimation capacity among lake trout (*Salvelinus namaycush*)

populations. *Conservation Physiology*, 2, doi:10.1093/conphys/cou025.

Kincaid, H.L. (1977). Rotational line crossing: an approach to the reduction of inbreeding

accumulation in trout brood stocks. *The Progressive Fish-Culturist*, 39, 179-181.

Kostenko, S., Dumitriu, G., Lægreid, K.J., Moens, U. (2011). Physiological roles of mitogen-

activated-protein-kinase-activated p38-regulated/activated protein kinase. *World Journal of*

Biological Chemistry, 2, 73-89.

Krens, S.G., He, S., Spaink, H.P., Snaar-Jagalska, B.E. (2006). Characterization and expression

patterns of the MAPK family in zebrafish. *Gene Expression Patterns*, 6, 1019-1026.

Lee, W.S., Salinas, S., Lee, Y.R., Siskidis, J.A., Mangel, M., Munch, S.B. (2020). Thermal

transgenerational effects remain after two generations. *Ecology and Evolution*, 10, 11296-11303.

Leitwein, M., Wellband, K., Cayuela, H., Le Luyer, J., Mohns, K., Withler, R., Bernatchez, L.

(2022). Strong parallel differential gene expression induced by hatchery rearing weakly

associated with methylation signals in adult Coho Salmon (O. kisutch). *Genome Biology and*

Evolution, 14, evac036.

Li, B., & Dewey, C.N. (2011). RSEM: accurate transcript quantification from RNA-Seq data

with or without a reference genome. *BMC Bioinformatics*, 12, 1-16.

Li, S., Guo, H., Chen, Z., Jiang, Y., Shen, J., Pang, X., Li, Y. (2021). Effects of acclimation

temperature regime on the thermal tolerance, growth performance and gene expression of a cold-

water fish, *Schizothorax prenanti*. *Journal of Thermal Biology*, 98, 102918.

Liu, Z., Huang, X., Yang, Z., Peng, C., Yu, H., Cui, C., Hu, Y., Wang, X., Xing, Q., Hu, J., Bao,

Z. (2021). Identification, characterization, and expression analysis reveal diverse regulated roles

of three MAPK genes in *Chlamys farreri* under heat stress. *Frontiers in Physiology*, 12, 688626.

Luu, I., Ikert, H., Craig, P.M. (2021). Chronic exposure to anthropogenic and climate related

stressors alters transcriptional responses in the liver of zebrafish (*Danio rerio*) across multiple

generations. *Comparative Biochemistry and Physiology Part C: Toxicology &*

Pharmacology, 240, 108918.

Mackey, T.E., Hasler, C.T., Durhack, T., Jeffrey, J.D., Macnaughton, C.J., Ta, K., Enders, E.C.,

Jeffries, K. M. (2021). Molecular and physiological responses predict acclimation limits in

juvenile brook trout (*Salvelinus fontinalis*). *Journal of Experimental Biology*, *224*, jeb241885.

Marshall, D.J. (2015). Environmentally induced (co)variance in sperm and offspring phenotypes

as a source of epigenetic effects. *Journal of Experimental Biology*, 218, 107–113.

Martin N.V., Olver C.H. (1980). The lake charr, Salvelinus namaycush. In: E.K. Balon (Ed.),

Charrs: Salmonid Fishes of the Genus Salvelinus, Ed 1st (pp. 205–277). Netherlands: Springer.

McCaw, B.A., T.J. Stevenson, L.T. Lancaster. (2020). Epigenetic responses to temperature and

climate. *Integrative and Comparative Biology*, 60, 1469-1480.

McDermid, J.L.,Wilson, C.C., Sloan, W.N., Shuter, B.J. (2013). Intraspecific differences in

thermal biology of inland lake trout populations. *Transactions of the American Fisheries Society*,

142, 756–766.

Metzger, D.C., Schulte, P.M. (2016). Maternal stress has divergent effects on gene expression
patterns in the brains of male and female threespine stickleback. *Proceedings of the Royal*
Society B: Biological Sciences, 283, 20161734.
990
991 Morash, A.J., Speers-Roesch, B., Andrew, S., Currie, S. (2021). The physiological ups and
992 downs of thermal variability in temperate freshwater ecosystems. *Journal of Fish Biology*, 98,
993 1524-1535.
994
995 Narum, S.R., Campbell, N.R. (2015). Transcriptomic response to heat stress among ecologically
996 divergent populations of redband trout. *BMC Genomics*, 16, 1-12.
997
998 Nazari, S., Pourkazemi, M., Paknejad, H., Kazemi, E., Ghaderi, M., Eslamloo, K. (2021).
999 Transcriptome profiling of farmed rainbow trout (Oncorhynchus mykiss) liver from different
1000 sources of dietary zinc. *Aquaculture*, 543, 737017.
1001
1002 Olsvik, P.A., Vikeså, V., Lie, K.K., Hevrøy, E.M. (2013). Transcriptional responses to
1003 temperature and low oxygen stress in Atlantic salmon studied with next-generation sequencing
1004 technology. *BMC Genomics,* 14, 1-21.
1005
1006 OMNR Fish Culture Stocks Catalogue. (2005). Fish Culture Section. Peterborough, ON: Ontario
1007 Ministry of Natural Resources.
1008
1009 Oomen, R. A., Hutchings, J. A. (2017). Transcriptomic responses to environmental change in
1010 fishes: Insights from RNA sequencing. *Facets*, *2*, 610-641.
1011
1012 Penney, C.M., Burness, G., Tabh, J.K.R., Wilson, C.C. (2021). Limited transgenerational effects
1013 of environmental temperatures on thermal performance of a cold-adapted salmonid.
1014 *Conservation Physiology*, 9, coab021.
1015

1016 Pérez-Ruzafa, A., Pérez-Marcos, M., Marcos, C. (2018). From fish physiology to ecosystems

1017 management: Keys for moving through biological levels of organization in detecting

1018 environmental changes and anticipate their consequences. *Ecological Indicators*, 90, 334-345.

1019

1020 Perrier, C., Ferchaud, A.L., Sirois, P., Thibault, I., Bernatchez, L. (2017). Do genetic drift and

1021 accumulation of deleterious mutations preclude adaptation? Empirical investigation using

1022 RADseq in a northern lacustrine fish. *Molecular Ecology*, 26, 6317–6335.

1023

1024 Petitjean, Q., Jean, S., Gandar, A., Côte, J., Laffaille, P., Jacquin, L. (2019). Stress responses in

1025 fish: From molecular to evolutionary processes. *Science of the Total Environment*, 684, 371-380.

1026

1027 Purohit, G. K., Mahanty, A., Mohanty, B. P., Mohanty, S. (2016). Evaluation of housekeeping

1028 genes as references for quantitative real-time PCR analysis of gene expression in the murrel

1029 *Channa striatus* under high-temperature stress. *Fish Physiology and Biochemistry*, 42, 125-135.

1030

1031 Qian, B., Xue, L. (2016). Liver transcriptome sequencing and de novo annotation of the large

1032 yellow croaker (*Larimichthy crocea*) under heat and cold stress. *Marine Genomics*, 25, 95-102.

1033

1034 Quinn, N.L., McGowan, C.R., Cooper, G.A., Koop, B.F., Davidson, W.S. (2011). Identification

1035 of genes associated with heat tolerance in Arctic charr exposed to acute thermal

1036 stress. *Physiological Genomics*, 43, 685-696.

1037

1038 Riley, S.C., Hansen, M.J., Krueger, C.C., Noakes, D.L.G., Muir, A.M. (2021). Introduction. The

1039 Lake Charr: Biology, Ecology, Distribution, and Management. In: A.M. Muir, M.J. Hansen, C.C.

1040 Krueger, & S.C. Riley (Eds.), The Lake Charr *Salvelinus namaycush*: Biology, Ecology,

1041 Distribution, and Management, Fish & Fisheries Series 39 (pp. 1-12). Switzerland: Springer

1042 Nature.

1043

1044 Ritchie, M.E., Phipson, B., Wu, D.I., Hu, Y., Law, C.W., Shi, W., Smyth, G.K. (2015). limma

1045 powers differential expression analyses for RNA-sequencing and microarray studies. *Nucleic

1046 Acids Research*, 43, e47-e47.

1047

1048 Rosenfeld, J., Richards, J., Allen, D., Van Leeuwen, T., Monnet, G. (2020). Adaptive trade-offs

1049 in fish energetics and physiology: insights from adaptive differentiation among juvenile

1050 salmonids. *Canadian Journal of Fisheries and Aquatic Sciences*, 77, 1243-1255.

1051

1052 Salinas, S., Munch, S.B. (2012). Thermal legacies: transgenerational effects of temperature on

1053 growth in a vertebrate. *Ecology Letters*, 15, 159-163.

1054

1055 Sambrook, J., Russell, D.W. (2001). Molecular Cloning: a laboratory manual. Cold Spring

1056 Harbor: Cold Spring Harbor Laboratory Press.

1057

1058 Sánchez-Tójar, A., Lagisz, M., Moran, N.P., Nakagawa, S., Noble, D.W., Reinhold, K. (2020).

1059 The jury is still out regarding the generality of adaptive 'transgenerational' effects. *Ecology*

1060 *Letters*, 23, 1715–1718.

1061

1062 Shahjahan, M., Zahangir, M.M., Islam, S.M.M., Ashaf-Ud-Doulah, M., Ando, H. (2021). Higher

1063 acclimation temperature affects growth of rohu (*Labeo rohita*) through suppression of GH and

1064 IGFs genes expression actuating stress response. *Journal of Thermal Biology*, 100, 103032.

1065

1066 Shama, L.N., Strobel, A., Mark, F.C., Wegner, K.M. (2014). Transgenerational plasticity in

1067 marine sticklebacks: maternal effects mediate impacts of a warming ocean. *Functional Ecology*,

1068 28, 1482–1493.

1069

1070 Shama, L.N., Mark, F.C., Strobel, A., Lokmer, A., John, U., Mathias Wegner, K. (2016).

1071 Transgenerational effects persist down the maternal line in marine sticklebacks: gene expression

1072 matches physiology in a warming ocean. *Evolutionary Applications*, 9, 1096–1111.

1073

1074 Shama, L.N. (2017). The mean and variance of climate change in the oceans: hidden

1075 evolutionary potential under stochastic environmental variability in marine

1076 sticklebacks. *Scientific Reports*, 7, 1-14.

1077

1078 Shekh, K., Tang, S., Niyogi, S., Hecker, M. (2017). Expression stability and selection of optimal

1079 reference genes for gene expression normalization in early life stage rainbow trout exposed to

1080 cadmium and copper. *Aquatic Toxicology,* 190, 217-227.

1081

1082 Shi, K.P., Dong, S.L., Zhou, Y.G., Li, Y., Gao, Q.F., Sun, D.J. (2019). RNA-seq reveals temporal

1083 differences in the transcriptome response to acute heat stress in the Atlantic salmon (*Salmo*

1084 *salar*). *Comparative Biochemistry and Physiology Part D: Genomics and Proteomics*, 30, 169-

1085 178.

1086

1087 Smith, D. A., Ridgway, M. S. (2019). Temperature selection in brook charr: lab experiments,

1088 field studies, and matching the Fry curve. *Hydrobiologia*, *840*, 143-156.

1089

1090 Smith, D.A., Jackson, D.A., Ridgway, M.S. (2020). Thermal habitat of brook trout in lakes of

1091 different size. *Freshwater Science*, 39, 56-69.

1092

1093 Somero, G.N. (2010). The physiology of climate change: how potentials for acclimatization and

1094 genetic adaptation will determine 'winners' and 'losers'. *Journal of Experimental Biology*, 213,

1095 912–920.

1096

1097 Smith, S.R., Normandeau, E., Djambazian, H., Nawarathna, P.M., Berube, P., Muir, A.M.,

1098 Ragoussis, J., Penney, C.M., Scribner, K.T., Luikart, G., Wilson, C.C., Bernatchez, L. (2022). A

1099 chromosome-anchored genome assembly for Lake Trout (*Salvelinus namaycush*). *Molecular*

1100 *Ecology Resources*, 22, 679-694.

1101

1102 Spadafora, C. (2020). Transgenerational epigenetic reprogramming of early embryos: a

1103 mechanistic model. *Environmetal Epigenetics*, 6, dvaa009.

1104

1105 Tian, Y., Wen, H., Qi, X., Zhang, X., Li, Y. (2019). Identification of mapk gene family in

1106 *Lateolabrax maculatus* and their expression profiles in response to hypoxia and salinity

1107 challenges. *Gene*, 684, 20-29.

1108

Tomalty, K.M., Meek, M.H., Stephens, M.R., Rincón, G., Fangue, N.A., May, B.P., Baerwald, M.R. (2015). Transcriptional response to acute thermal exposure in juvenile Chinook salmon determined by RNAseq. *G3: Genes, Genomes, Genetics*, *5*, 1335-1349.

Uller, T., Nakagawa, S., English, S. (2013). Weak evidence for anticipatory parental effects in plants and animals. *Journal of Evolutionary Biology*, 26, 2161-2170.

Veilleux, H.D., Ryu, T., Donelson, J.M., Van Herwerden, L., Seridi, L., Ghosheh, Y., Berumen, M.L., Leggat, W., Ravasi, T., Munday, P. L. (2015). Molecular processes of transgenerational acclimation to a warming ocean. *Nature Climate Change*, 5, 1074-1078.

Veilleux, H.D., Donelson, J.M., Munday, P.L. (2018). Reproductive gene expression in a coral reef fish exposed to increasing temperature across generations. *Conservation Physiology*, 6, doi: 10.1093/conphys/ cox077.

Venney, C.J., Love, O.P., Drown, E.J., Heath, D.D. (2020). DNA methylation profiles suggest intergenerational transfer of maternal effects. *Molecular Biology and Evolution*, 37, 540-548.

Venney, C.J., Wellband, K.W., Normandeau, E., Houle, C., Garant, D., Audet, C., Bernatchez, L. (2022). Thermal regime during parental sexual maturation, but not during offspring rearing, modulates DNA methylation in brook charr (*Salvelinus fontinalis*). *Proceedings of the Royal Society B: Biological Sciences*, 289, 20220670.

Visser, M.E. (2008). Keeping up with a warming world; assessing the rate of adaptation to climate change. *Proceedings of the Royal Society B: Biological Sciences*, 275, 649–659.

Willi, Y., Van Buskirk, J., Hoffmann, A.A. (2006). Limits to the adaptive potential of small populations. *Annual Review of Ecology, Evolution and Systematics*, 37, 433–458.

Wilson, C. (2017). Genetic aspects of climate change influences on inland fishes and fisheries. *Fisheries*, 42, 125–126.

1140

1141 Wilson, C.C., Mandrak, N.E. (2021). Paleoecology. In: A.M. Muir, M.J. Hansen, C.C. Krueger,

1142 & S.C. Riley (Eds.), The Lake Charr *Salvelinus namaycush*: Biology, Ecology, Distribution, and

1143 Management, Fish & Fisheries Series 39 (pp. 41-68). Switzerland: Springer Nature.

1144

1145 Yano, A., Nicol, B., Jouanno, E., Quillet, E., Fostier, A., Guyomard, R., Guiguen, Y. (2013). The

1146 sexually dimorphic on the Y-chromosome gene (sdY) is a conserved male-specific Y-

1147 chromosome sequence in many salmonids. Evolutionary applications, 6, 486-496.

1148

1149 Yin, J., Zhou, M., Lin, Z., Li, Q.Q., Zhang, Y.Y. (2019). Transgenerational effects benefit

1150 offspring across diverse environments: a meta-analysis in plants and animals. *Ecology Letters*,

1151 22, 1976–1986.

1152

1153 Zak, M.A., Manzon, R.G. (2019). Expression and activity of lipid and oxidative metabolism

1154 enzymes following elevated temperature exposure and thyroid hormone manipulation in juvenile

1155 lake whitefish (*Coregonus clupeaformis*). *General and Comparative Endocrinology*, 275, 51-64.

1156

1157 Zhang, X., Flato, G., Kirchmeier-Young, M., Vincent, L., Wan, H., Wang, X., Rong, R., Fyfe, J.,

1158 Li, G., Kharin, V.V. (2019). Changes in Temperature and Precipitation Across Canada. In: E.

1159 Bush & D.S. Lemmen (Eds.) Canada's Changing Climate Report (pp. 112-193). Ottawa, Ontario:

1160 Government of Canada.

1161

1162

1163

1164 **Tables**

1165

1166 **Table 1**: Summary of the comparison treatments used for assessing differential gene expression in juvenile lake trout following

1167 within- and transgenerational thermal acclimation, showing treatment groups (offspring and parental treatments), and measured

1168 effects.

	Treatment group (parent treatment)	Sequence year (sample size)	Offspring sex ratio (♂,♀)	Effect measured
Control group	Cold-acclimated offspring (C♀xC♂)	2018 (n=6) & 2020 (n=6)	2,4 5,1	*No effect (all other groups are compared to the control group)*
Within-generation warm-acclimation	Warm-acclimated offspring (C♀xC♂)	2018 (n=6)	3,3	Offspring warm-acclimation
Transgenerational warm-acclimation	Cold-acclimated offspring (W♀xC♂)	2020 (n=6)	4,2	Maternal warm-acclimation
	Cold-acclimated offspring (C♀xW♂)	2020 (n=6)	4,2	Paternal warm-acclimation
	Cold-acclimated offspring (W♀xW♂)	2018 (n=6)	3,3	Maternal & paternal warm-acclimation
Within- and transgenerational warm-acclimation	Warm-acclimated offspring (W♀xC♂)	2020 (n=6)	2,4	Offspring & maternal warm-acclimation
	Warm-acclimated offspring (C♀xW♂)	2020 (n=6)	3,3	Offspring & paternal warm-acclimation

| Warm-acclimated offspring (W♀xW♂) | 2018 (n=6) | 5,1 | Offspring, maternal & paternal warm-acclimation |

1169

1170 **Figures**

1177
1178

1179 **Figure 1**: Experimental treatment groups of lake trout (*Salvelinus namaycush*). Two males and
1180 two females each from cold- and warm-acclimation tanks were bred in a full factorial cross,
1181 resulting in four families per parental treatment group (C♀xC♂, W♀xC♂, C♀xW♂, W♀xW♂; one
1182 each depicted here for simplicity). Offspring from each family was divided and acclimated to
1183 either a cold (11°C) or warm (15°C) temperature.

1184
1185

1186
Figure 2: Heatmap comparing the effect of parental and offspring thermal acclimation on
differential gene expression in juvenile lake trout, grouped by combined parental and offspring
warm acclimation (transgenerational warming; top) and offspring acclimation treatment (within-
generation; bottom). Each column represents a treatment group (n = 6 individuals) compared
against the control group (11°C C♀xC♂; n = 6; not shown). Gene expression levels are
represented as log2-fold change (LFC; blue = downregulate, red = upregulated). LFC is sorted
according to gene expression level in warm-acclimated offspring to show that any given gene up-
or downregulated by offspring warm acclimation could be expressed differently with the

influence of maternal and/or paternal warm acclimation. The list of genes along with their LFC
values and FDR (adjusted p-value) can be found in the supplementary information (Suppl. Table
3).
1198

1199
1200
1201
1202

1203

Figure 3: Number of transcripts that were differentially expressed in response to warm-acclimation of juvenile lake trout offspring (n = 6) when compared against the control group (11°C C♀xC♂; n = 12). Genes were expressed at log2-fold change (LFC) of ≤ -1 or ≥ +1 and are presented to show the total differentially expressed transcripts (top) which were then separated to show the upregulated (bottom, left) and downregulated (bottom, right) transcripts.

Total transcripts

Upregulated

Downregulated

Warm acclimation of offspring & both parents

Offspring & maternal warm acclimation

Offspring & paternal warm acclimation

Offspring warm acclimation

Figure 4: Number of transcripts that were differentially expressed in response to
transgenerational (parental) warm-acclimation of juvenile lake trout offspring when the mothers
(A) or the fathers (B) were warm-acclimated, and when both parents (C) were warm-acclimated.
Treatment groups (n = 6 each) were compared against control group (11°C C♀xC♂; n = 12).
Genes were expressed at log2-fold change (LFC) of $\leq$ -1 or $\geq$ +1 and are presented to show the
total differentially expressed transcripts (top) which were then separated to show the upregulated
(bottom, left) and downregulated (bottom, right) transcripts.