Transcriptomic responses to multigenerational environmental warming in a cold-adapted

salmonid using *de novo* RNA sequencing

Running title: Lake trout transcriptomic responses to warming

Chantelle M. Penney[1], Gary Burness[2], and Chris C. Wilson[3]

[1] Environmental and Life Sciences Graduate Program, Trent University, Peterborough, Ontario,

K9J 7B8

[2] Department of Biology, Trent University, Peterborough, Ontario, K9L 0G2

[3] Aquatic Research and Development Section, Ontario Ministry of Natural Resources and

Forestry, Trent University, Peterborough, Ontario, K9L 0G2

Author for correspondence: Chantelle Penney, Environmental and Life Sciences Graduate

Program, Trent University, Peterborough, Ontario, K9L 0G2, chantellepenney@trentu.ca

Authors' contributions: CMP, GB and CCW conceived the ideas and designed methodology;

CMP collected and analysed the data, and led the writing of the manuscript. All authors

contributed critically to manuscript drafts and gave final approval for publication.

Conflict of interest: The authors have no conflict of interest to declare.

Abstract

Cold-adapted species are particularly threatened by climate change as rates of environmental warming outpace the ability of many populations adapt. Recent evidence suggest that transgenerational thermal plasticity may play a role in the response of cold-adapted organisms to long-term changes in temperature. Using RNA sequencing, we explored differential gene expression of lake trout (*Salvelinus namaycush*), a cold-adapted species, to examine the molecular processes that respond to elevated temperatures under conditions of within-generation (offspring) and transgenerational (parental) warm-acclimation. We hypothesized that genes associated with metabolism, growth and thermal stress/tolerance would be differentially expressed in juvenile lake trout offspring depending on their own acclimation temperature and that of their parents. We found enriched pathways for thermal stress, signaling processes, immune function, and transcription regulation, although the specific genes and direction of differential expression (up- or downregulation) depended on the combination of offspring and parental thermal rearing conditions. While parental warm-acclimation did have a transgenerational effect on gene expression in their offspring, within-generation warm-acclimation had a larger effect on differential expression, enriching more pathways. We provide evidence of the transgenerational response to warming at the transcriptional level in lake trout, which should be useful for future studies of transcriptomics and plasticity in these and other cold-adapted species.

Keywords: acclimation, climate change, RNA sequencing, *Salvelinus namaycush*, transgenerational plasticity, transcriptomics.

Introduction

Temperate and Arctic regions are warming rapidly; surface temperatures in Canada have

increased by ~1.7°C in under a century with even greater warming happening in the Arctic

(~2.3°C; Zhang et al. 2019). As a result, high-latitude, cold-adapted organisms are experiencing,

and will continue to experience, rapid temperature increases that may threaten their survival

(Beitinger and Bennett 2000; Burkhead 2012; IPCC 2021). Range expansion to track suitable

conditions is particularly challenging for species and populations with limited dispersal options,

such as many freshwater fishes (Guzzo and Blanchfield 2017; Smith et al. 2020). Warming rates

may outpace adaptive responses (Visser 2008; Crozier and Hutchings 2014; Comte and Olden

2017), particularly for those with low standing genetic variation (Willi et al. 2006; Wilson 2017).

In the absence of standing genetic variation for coordinated adaptive responses throughout core

metabolic pathways, population responses would be limited to plastic thermal acclimation, but

the low physiological plasticity typical of many cold-adapted species means they likely will be

particularly vulnerable to rapidly changing environments (Somero 2010; Kelly et al. 2014).

Individuals that can withstand warming may have to rely heavily on energetic resources that

would normally support growth or reproduction, thus having potentially negative downstream

effects on the population (Somero 2010; Rosenfeld et al. 2020). A lack of adaptive genetic

variation, low capacity for temperature acclimation, and rapid environmental change are

predicted to result in widespread species extirpations (Bennett et al. 2019; Morash et al. 2021).

81   Some species may be able to buffer the negative impact of environmental warming

through transgenerational acclimation, also known as transgenerational plasticity (TGP). As the

name suggests, TGP refers to the phenotypic changes that occur over the course of multiple

generations (Jablonka et al. 1992; Bell and Hellmann 2019; Bonduriansky 2021). This involves

non-genetic inheritance of environmentally-driven changes, including parental effects, where the

experiences of one generation are reflected in the morphology, physiology or behaviour of a

subsequent generation, often F1 or F2 (Hellman et al. 2020; Lee et al. 2020; Bonduriansky

2021).

89   TGP occurs through modulation of gene expression through non-genetic inheritance,

which includes parental effects and the transfer of epigenetic factors such as patterns of DNA

methylation and histone modification, and microRNA activity (Dai et al. 2020; Spadafora 2020;

Venney et al. 2020). TGP can potentially be beneficial when a parent's environment coincides

with what their offspring are expected to experience (Bernatchez 2016; Ashe et al. 2021). For

example, TGP could be beneficial in a population that experiences environmental warming over

multiple generations whereby the offspring are phenotypically "primed" for a warmer

environment (Donelan et al. 2020; McCaw et al. 2020; Venney et al. 2022). In the context of

climate change, this phenotypic priming could buffer the effects of rapid warming, allowing

additional time for evolutionary processes to occur (Bernatchez 2016; Ashe et al. 2021), though

the extent to which TGP can be considered adaptive is arguable (Uller et al. 2013; Sánchez-Tójar

et al. 2020). Nevertheless, TGP with temperature has been identified in a number of species

(Greenspoon and Spencer 2018; Yin et al. 2019). For example, the coral reef

fish *Acanthochromis polyacanthus* exhibits TGP for increased temperatures by improving its

aerobic scope in a warm environment by increasing maximum metabolic rate or decreasing

resting metabolic rate (Donelson et al. 2012; Bernal et al. 2018). Many of the studies on thermal

TGP in aquatic organisms have used eurythermal, temperate or tropical model species; by

contrast, cold-adapted aquatic vertebrates are relatively understudied but could stand to benefit

from TGP given their vulnerability to environmental warming due to climate change.

Acute and chronic temperature acclimation has obvious impacts on the metabolism of

ectotherms, at both the whole organism and molecular level (Pérez-Ruzafa et al. 2018; Petitjean

et al. 2019; Morash et al. 2021). Relatively few studies to date report the effect of

transgenerational warming on phenotypic variation in tandem with gene expression, or with

follow-up experiments testing the differential gene expression underlying transgenerational

effects (Shama et al. 2016; Bernal et al. 2018; Veilleux et al. 2018). Examining differential gene

expression associated with changes in metabolism with thermal experiences can be informative.

Shama et al. (2016), for example, found that genes involved in metabolism, mitochondrial

protein synthesis, hemostasis and apoptosis were differentially expressed in offspring depending

on the multigenerational thermal experiences down the maternal line. These findings underscore

the relationship between the molecular mechanisms underlying physiological plasticity and

parental thermal experiences.

We sought to explore differential gene expression in lake trout (*Salvelinus namaycush*) to

examine the molecular processes potentially underlying transgenerational responses to elevated

temperatures. The lake trout is a long-lived, cold-adapted salmonid that is largely limited to cold,

oligotrophic lakes in formerly glaciated regions of North America (Riley et al. 2021; Wilson and

Mandrak 2021) and is under significant threat from climate change (Casselman 2008; Guzzo and

Blanchfield 2017). Being stenothermal, lake trout behaviourally thermoregulate by migrating to

deeper, cooler water when lakes thermally stratify during the summer (Martin and Olver 1980;

Guzzo and Blanchfield 2017). Low genetic variation for some lake trout populations (Perrier et

al. 2017), and little within- and among-population variation for thermal acclimation capacity

(McDermid et al. 2013; Kelly et al. 2014) may limit the adaptive potential of lake trout

populations for warming temperate and arctic habitats. A previous study investigated the capacity

for transgenerational thermal plasticity in lake trout as a means of coping with warming

associated with climate change and found that they exhibit only limited transgenerational

plasticity (Penney et al. 2021). Unexpectedly, cold-acclimated offspring of parents from elevated

temperatures exhibited higher resting metabolic rate than those with one or both cold acclimated

parents (Penney et al. 2021).

For this study, we assessed gene expression in previously studied offspring from factorial

mating crosses between cold- and warm-acclimated lake trout adults (Penney et al. 2021) to

assess transcriptional responses to within- and transgenerational thermal acclimation.

Acclimation of the parents (transgenerational) to one of two environmental temperatures (10 or

17°C) was combined with subsequent acclimation of the offspring (within-generation) to cool

(11°C) or warm (15°C) temperatures (Fig. 1). Evidence exists to support both maternal and

paternal contributions to transgenerational effects (Marshall 2015; Shama et al. 2016; Penney et

al. 2021). Our experimental design allowed us to examine the transgenerational effect of

temperature acclimation of each parent, and both parents together, in combination with the

offspring's (within-generation) temperature acclimation on gene expression in juvenile lake trout.

We hypothesized that genes associated with metabolism, growth and thermal

stress/tolerance would be differentially expressed in juvenile lake trout depending on both the

acclimation temperature of the offspring and the acclimation temperature of their parents. Based

on studies of within-generation and transgenerational warming on differential gene expression in

fish (Quinn et al. 2011; Veilleux et al. 2015; Akbarzadeh and Leder 2016; Shama et al. 2016), we

predicted that 1) genes involved in growth, heat shock and hypoxia responses, and metabolic

pathways would be upregulated in warm-acclimated offspring, and 2) differential expression of

these genes would vary depending on whether one or both parents were warm-acclimated,

although the size and direction of the effect is difficult to predict given that parental influence on

offspring gene expression has been shown to be sex-specific; depending on the sex of the

offspring and that of the parent (Best et al. 2018; Baustista et al. 2020).

Methods

Hatchery-raised adult lake trout from the Ontario provincial Seneca Lake broodstock

were used as parents for the experiment. This hatchery strain originated from Seneca Lake, one

of the Finger Lakes in central New York State (42°41' N, 76°54' W). The Ontario Ministry of

Natural Resources and Forestry (OMNRF) hatchery system has maintained this lake trout strain

for over five generations, using rotational line crossing (Kincaid 1977) to minimize inbreeding

and maintain the founding genetic diversity of the strain (OMNRF Fish Culture Stocks

Catalogue 2005).

The tissue samples used for these experiments were collected from juvenile (young of

year) lake trout that were previously used in a separate study on metabolic rate with

transgenerational warming described in Penney et al. (2021). In this previous study, adult and

juvenile offspring lake trout were acclimated to two different temperatures and the offspring

were subjected to an acute thermal challenge of $+2°C·h^{-1}$ to determine the effect of within- and

transgenerational temperature acclimation on upper thermal tolerance and metabolic rate. Details

of the prior experiment (Penney et al. 2021) that are relevant to the current study are summarized

below.

All experiments adhered to the Canadian Council on Animal Care guidelines and were

approved by the Trent University Animal Care Committee (Protocol # 24794) and the OMNRF

Aquatic Animal Care Committee (Protocol # FACC_136).

Experimental design and rearing conditions

Two groups (n = 8 and 9) of mature lake trout (age 8; 2.3 - 4.2 kg) were each held in

6,000 L tanks (1 x 1 x 6 m), with males and females kept together, at the OMNRF White Lake

Fish Culture Station (Sharbot Lake, Ontario, Canada) with water being supplied from a nearby

lake. As the source waterbody (White Lake) began to stratify in early June, 2015, we began to

increase tank water temperature by 1°C per day starting from approximately 9°C until one tank

reached 10 ± 0.5°C and the other 17 ± 0.5°C. These temperatures were chosen because 10°C

represents the thermal requirements for lake trout spawning while 17°C is meant to push the fish

to their physiological limits without compromising reproduction (Casselman 2008; Penney et al.

2021) at the higher temperature. Temperatures were controlled and maintained by drawing in and

mixing water from above and below the lake's thermocline as it flowed into the holding tanks.

Fish were acclimated to these temperatures for approximately 3 months, mirroring thermal

stratification in the source waterbody. Starting in mid-September, the temperature in the warm-

water treatment was allowed to gradually cool by holding the proportional inflows from above

and below the thermocline constant as the lake's surface water cooled. Temperatures in the two

tanks converged at 10°C after fall turnover (late September) in the lake, and gradually cooled to

7.8°C by the end of the breeding interval (mid-November).

Adult lake trout were reproductive by the end of October with the warm-acclimated fish

first spawning on October 30[th]. The cold-acclimated adults were first ready to spawn on

November 5[th] and all mating crosses were completed by November 19th. Experimental offspring

families were produced by a full factorial mating cross using two males and two females from

each of the two temperature treatments (4 x 4, 8 adults in total) resulting in a total of 16 offspring

families; four families from each of four parental treatment groups (W♀xW♂, W♀xC♂, C♀xW♂

and C♀xC♂, where W = warm-acclimated, and C = cold-acclimated; Figure 1). Fertilized eggs

were transported to the OMNRF Codrington Fish Research Facility (Codrington, Ontario,

Canada) where they were transferred into 200 L tanks receiving freshwater at ambient

temperature (5-6 °C) and natural photoperiod under dim light with each family separately

contained in steel mesh boxes (9 x 9 x 7.5 cm, one family per box).

In March (2016), when hatched offspring were ready to begin exogenous feeding, we

randomly selected 14 individuals from each of the 16 families to transfer to 200 L tanks for

temperature acclimation. Seven fry from each family were acclimated to a cold temperature

(11°C) and the other 7 from each family were acclimated to a warm temperature (15°C). The

offspring's lower acclimation temperature differs from that of their parents because of reported

differences in optimum temperature for both life stages (11 vs 10°C; Edsall 2000, Casselman

2008). The upper acclimation temperature for the offspring represents the anticipated increase in

temperature associated with climate change by year 2100 (Hayhoe et al. 2010; IPCC 2021),

however, for the parents the upper acclimation temperature was two degrees higher to ensure that

temperature was high enough to elicit a physiological thermal response without compromising

reproduction. To begin acclimating the offspring, water temperature was increased at a rate of

1°C per day until target temperatures were reached (11 and 15°C). Due to space constraints,

tanks were subdivided by adding steel inserts with perforated bottoms. Each insert was divided

into 4 sections (24 x 25 x 28 cm per section) with two paternal half-sibling families (n = 7 each;

shared male parent) housed per section. Offspring from pooled families were later individually

identified to family using microsatellite genotyping (Penney et al. 2021).

After the 3-4 week acclimation period, offspring were subjected to an acute temperature

increase of $+2°C·h^{-1}$ from their acclimation temperature until loss of equilibrium was observed,

and then quickly transferred to a bath of 0.3 g l−1 of buffered tricaine methanesulfonate (MS-

222; Aqua Life, Syndel Laboratories Ltd, BC, Canada). Following euthanasia, the whole liver

was rapidly dissected from each fish, blotted on a lab wipe and preserved in RNAlater

(Invitrogen, Thermo Fisher Scientific) as per manufacturer's instructions. After 24-48 hours in a

4°C refrigerator, the RNAlater was pipetted from the tissues and the samples were stored at -

80°C until RNA extraction. We chose to study gene expression in the liver because it is a

metabolically active tissue involved in a number of physiological processes that respond to

warming (Quinn et al. 2011; Akbarzadeh and Leder, 2016; Dammark et al. 2018). Additionally,

transcriptional responses to multigenerational warming have been observed in the livers of

zebrafish: Luu et al. (2021). Therefore, the liver ought to provide insight into the metabolic

pathways that are affected by transgenerational temperature acclimation in lake trout. Liver

tissue was collected between June 28 and August 9 in 2016.

RNA isolation and sequencing

RNA was extracted from the preserved liver tissues using a phenol-chloroform extraction

method (Chomczynski and Sacchi 2006) after the tissues had been individually homogenized via

a FastPrep-24 BeadBeater (MP Biomedicals) with 2 mL Lysing Matrix D tubes (MP

Biomedicals) and 1 mL of Trizol reagent (Invitrogen, Thermo Fisher Scientific). RNA was

precipitated with RNA precipitation solution (Sambrook and Russel 2001) and isopropanol, and

washed with 75% ethanol. RNA samples were resuspended in nuclease-free water (Thermo

Fisher Scientific). The purity and concentration of the RNA was quantified using a NanoDrop-

8000 spectrophotometer, and RNA quality was assessed by gel electrophoresis of glyoxylated

RNA on a 1.5% agarose gel (Sambrook and Russel 2001).

Two lots of RNA samples were sent for high-throughput sequencing over a two-year

span. In 2018, liver RNA samples from 24 individuals were sent to The Centre for Applied

Genomics (Sick Kids Hospital, Toronto, Ontario, Canada), and in 2020 another 30 liver samples

(individuals) were sent to Genome Quebec (Montreal, Quebec, Canada). Altogether, the 54

samples included tissue from six individuals from each of the seven experimental treatment

groups, however, the control group (11°C-acclimated offspring from cold-acclimated (CxC)

parents) had a total of 12 individuals sequenced: 6 individuals in 2018 and an additional 6 in

2020, so that each treatment group was compared against the control group sequenced in the

same year to eliminate batch effects in the analysis of differential expression (Table 1). Each

group included at least one representative from each family. Both facilities assessed the RNA

quality via a Bioanalyzer (Agilent Technologies) and all samples passed quality control (RNA

integrity number: ≥ 7.5). cDNA libraries were constructed by enriching the poly(A) tails of

mRNA with oligo dT-beads using NEBNext Ultra II Directional polyA mRNA Library Prep. In

2018, barcoded libraries were distributed among two and a half lanes and were sequenced on the

Illumina HiSeq2500 instrument producing an average of 28 million reads per sample (n = 24,

paired-end, 2x126 bp). In 2020, libraries were sequenced on the Illumina HiSeq4000 with

barcoded libraries distributed among three lanes producing 36 million reads per sample (n = 30,

paired-end, 2x126 bp).

De novo assembly, annotation and analysis

RNA reads from both years were assembled and annotated by the Canadian Centre for

Computational Genomics (C3G, Quebec) in 2020. Adapters and reads with a Phred quality score

<30 were trimmed using Trimmomatic (v.0.36), and normalization and *de novo* assembly was

performed using Trinity assembly software (v.2.0.4) following Haas et al. (2013) and Grabherr et

al. (2011). Pre- and post-trimming read counts and Trinity statistics can be found in the

supplementary file (Tables S1 and S2). Functional annotation was performed with the Trinotate

tool (v.2.0.2) and using the TransDecoder pipeline (v.2.0.1) to identify candidate coding regions

and open reading frames. Candidate peptides were searched against known proteins in the Swiss-

Prot (UniProtKB/Swiss-Prot) database via BLASTp (v.2.3.0) and the pfam database. The pfam

database was also used with HMMER (v.3.1b2) to identify protein domains. Signal peptides,

transmembrane regions and ribosomal RNA genes were predicted using the programs signalP

(v.4.1), tmHMM (v.2.0c) and RNAMMER (v.1.2), respectively.

279   Seven comparisons were made where offspring treatment groups were each compared to

the control group (cold-acclimated offspring with cold-acclimated parents) to observe the effect

that within- and transgenerational plasticity (acclimation) had on gene expression in juvenile lake

trout (Table 1). Transcript abundance was estimated using 'RNA-Seq by Estimation

Maximization' (RSEM; Li and Dewey, 2011) in Trinity with the

'align_and_estimate_abundance.pl' utility. Differential gene expression was analyzed between

the groups of lake trout offspring in R using the limma package (v.3.40.6; Ritchie et al., 2015)

with nominal p-values corrected for multiple testing using the Benjamini-Hochberg method with

the false discovery rate (FDR) of 0.1. The log2fold change (LFC) was also calculated for DEGs,

representing the magnitude of up- or downregulation; negative LFC are downregulated genes.

289   For each comparison, an enrichment analysis was performed on DEGs using the goseq

package (v.1.28.0) in R which assigned Gene Ontology (GO) terms to transcripts and categorized

them based on function (CC: cellular components, BP: biological processes, MF: molecular

function). This initial analysis returned results that ran DEGs against databases for multiple

species (e.g., human, mice, etc.). Using a limited list of DEGs (LFC of <-1 to >1), we validated

the function of these genes for lake trout with the Database for Annotation, Visualization and

Integrated Discovery (DAVID) tool (v.6.8) (Huang et al. 2009A; Huang et al. 2009B). The

DAVID tool identified the genes as official gene symbols and required that a model species be

specified for our gene list, so we opted to use *Salmo salar* because *S. namaycush* was not an

available option and, among orthologous species, *S. salar* had the best coverage for annotated

proteins available in the Swiss-Prot database (determined via NCBI Eukaryotic genome

annotation pipeline, v.8.5). The output from the DAVID tool mapped genes to functional

categories (GO) and metabolic pathways (Kyoto Encyclopedia of Genes and Genomes: KEGG),

and we limited the list of gene annotations to backgrounds belonging to fish species

(*Oncorhynchus mykiss*, *S. salar*, *Danio rerio*, *Oryzias latipes*). Tables outlining the enriched

functions are included in the supplemental material. We opted to keep the DAVID tool default

criteria of reporting functions that are enriched by a minimum of 2 genes (count) and *p*<0.1. We

discuss significantly (*p*≤0.05) enriched functions and pathways in the following sections with

gene IDs provided in parentheses.

Results

The transcriptome data showed both transgenerational and within-generation

environmental influences on gene expression (Fig. 2). Among all groups, there were 1,005

unique DEGs with a LFC of ≤ -1 to ≥ 1 that were recognized by the DAVID tool and used for

enrichment analysis. The level of differential expression of these genes are displayed in a

heatmap (Fig. 2). The first column represents the effect of offspring (within-generation) warm-

acclimation on differential gene expression and sets the order of the DEG list, sorted from

highest to lowest LFC. The list of genes along with their LFC values and adjusted *p*-value can be

found in the supplementary information (Suppl. Table 17).

Within-generation acclimation: offspring warm-acclimation

As predicted, warm-acclimation of lake trout offspring resulted in differential gene

expression compared to cold-acclimated offspring, providing support for our hypothesis that

warm acclimation would upregulate genes associated with metabolism, growth and thermal

tolerance. There were 193 upregulated DEGs mapped to GO terms or metabolic (KEGG)

pathways. These 193 genes are displayed in Fig. 3 which shows the number of genes that had

overlapping KEGG and GO functions. For example, there were 13 upregulated genes that had

overlapping functions relating to KEGG pathways, biological processes, molecular function and

cellular components whereas there were 27 genes that mapped uniquely the molecular function

(Fig. 3). Among these genes were 29 upregulated DEGs involved in signalling processes, 14

related to cytoplasm functions and 9 related to immunoglobulin (Suppl. Table 3). There were 4

genes associated with methyltransferase activity, though this function was not significantly

enriched (*p*=0.09; Suppl. Table 3). Upregulated DEGs that mapped to metabolic (KEGG)

pathways included those involved in degradation (*HMGCLL1*, *CNOT1*) such as peroxisome

activity and RNA and ketone degradation (Suppl. Table 4). Genes involved in the heat shock

response were also upregulated, specifically *hsp90aa1.1* (Suppl. Table 4). Others included cell

adhesion (*CD99L2, CTNNA2*) and signalling (*JAK1, PYCARD, CHD8*; Suppl. Table 4). Warm-

acclimated offspring had 180 downregulated DEGs that mapped to GO terms or KEGG

pathways (Fig. 3, right). There were 19 downregulated DEGs related with metal- and heme-

binding and 3 DEGs associated with cyclin activity (Suppl. Table 3). Contrary to our predictions,

downregulated DEGs involved in KEGG pathways included glycolysis/gluconeogenesis

(*GAPDH*) and metabolism of fructose and mannose (*ALDOB*; Suppl. Table 4). Other DEGs

involved in peroxidase activity were also downregulated with offspring warm-acclimation

(*MPV17L2*; Suppl. Table 4).

Transgenerational acclimation: parental warm-acclimation

345   Parental warm-acclimation influenced gene expression in cold-acclimated offspring. The

direction of differential expression (up- or down-regulation) in the offspring depended on which

parent was warm-acclimated and sometimes expression of a gene due to maternal or paternal

warm-acclimation contrasted with the expression level when only the offspring were warm-

acclimated (positive vs. negative LFC; Fig. 2). Nevertheless, parental warm-acclimation

influenced differential gene expression in the lake trout offspring, lending support to our

prediction that differential expression of genes would vary in offspring depending on parental

temperature acclimation.

353   With maternal warm-acclimation, 107 upregulated DEGs mapped to GO terms or KEGG

pathways (Fig. 4A). Maternal warm-acclimation had the effect of upregulating processes related

to muscle function in their offspring, specifically 2 DEGs involved in calcium-binding and 3

DEGs in troponin synthesis (Suppl. Table 5). There were 13 upregulated DEGs associated with

transferases and 3 DEGs related to glycolysis (Suppl. Table 5). Upregulated genes involved in

KEGG pathways included DNA damage/repair and fatty acid metabolism (*DDIT4*, *hsd17b12a*;

Suppl. Table 6). Maternal warm-acclimation resulted in 61 downregulated offspring DEGs

mapped to GO terms or KEGG pathways (Fig. 4A). There were 14 downregulated DEGs

involved in cell signalling and 10 involved in hydrolase/protease activity (Suppl. Table 5). The

KEGG pathway analysis revealed that maternal warm-acclimation downregulated *GAPDH*

involved in glycolysis/gluconeogenesis, and functions related to RNA transport and degradation

(*CNOT1* and *EIF3C*; Suppl. Table 6).

365   Paternal warm-acclimation also had an effect on offspring gene expression, providing

further support for our hypothesis regarding a transgenerational of warming on gene expression,

although fewer genes were differentially upregulated with paternal warming compared to

maternal warming. There were 69 upregulated DEGs mapped to GO terms or KEGG pathways

(Fig. 4B) which included 10 DEGs related to transferase activity along with 3 DEGs involved in

glycolytic processes (Suppl. Table 7). Upregulated KEGG pathways included the mTOR

signalling pathway (*DDIT4*) and protein processing (*ERO1A*; Suppl. Table 8). Slightly more (n =

75) DEGs were downregulated (Fig. 4B), including 16 DEGs involved in cell-signalling and 7

DEGs associated with protease activity (Suppl. Table 7). The KEGG pathway analysis revealed

there were two transcripts (one upregulated and the other downregulated), that were identified as

GAPDH and mapped to glycolysis (Suppl. Table 8). Comparing the adjusted *p*-value of both

transcripts, where a lower *p*-value means it is less likely a false positive (i.e. type I error),

suggests that *GAPDH* was actually downregulated ($p = 0.88$). Up- and downregulation was also

observed with *CNOT1* (Suppl. Table 8), but because the $p = 1$ in both cases, we cannot conclude

whether this DEG was up- or downregulated.

As expected, transgenerational effects were also observed in offspring gene expression

when both mothers and fathers were warm-acclimated. A total of 101 upregulated DEGs mapped

to GO terms or KEGG pathways (Fig. 4C). Among these were 16 DEGs associated with cell

signalling processes, 9 DEGs related to hydrolase/proteases/peptidases function and 2 DEGs

associated with peroxisome activity (Suppl. Table 9). Upregulated KEGG pathways included the

mTOR pathway (*DDIT4*), RNA degradation (*CNOT1*) and amino acid transferase activity

(*BHMT*, *GATM*, *GAMT*; Suppl. Table 10). A similar number of downregulated genes (n = 101)

mapped to GO terms or KEGG pathways (Fig. 4C) which included 4 DEGs involved in

glycosyltransferase activity, 3 DEGs related to lectin and 2 DEGs associated with tRNA function

(Suppl. Table 9). There were 7 downregulated DEGs associated with translation/ribosomal

proteins, but this function was not significantly enriched ($p=0.09$; Suppl. Table 9).

Downregulated KEGG pathways included peroxisome activity (*HMGCLL1*) and apoptotic

signaling (*pycard*) and Notch signaling pathways (*APH1B*; Suppl. Table 10). Again, *GAPDH*

appeared as both up- and downregulated in the KEGG analysis.

Multigenerational acclimation: offspring and parental warm-acclimation

As hypothesized, genes associated with metabolism, growth and thermal stress/tolerance

were differentially expressed with the combined effect of offspring and parent warm acclimation.

In some cases, a combination of offspring and parent warm-acclimation resulted in more DEGs

(e.g. >200) being up- or downregulated relative to when the within-generation warm-acclimation

and transgenerational warm-acclimation were considered in isolation.

With offspring and maternal warm-acclimation, 193 upregulated genes mapped to GO

terms or KEGG pathways (Fig. 5A). These included 20 DEGs related transferase function, 4

DEGs involved in metalloprotease/endopeptidase activity and 4 associated with the PDZ domain

(Suppl. Table 11). There were also 3 DEGs associated with protein folding, 3 DEGs associated

with DNA repair, and 3 associated with muscle contraction/ZASP (Suppl. Table 11). Genes that

encode heat shock proteins (*HSP70*, *HSPB1*, *HSP7C*, *HSP30*) and proteins involved in the

respiratory chain (*COX2*, *COX1*) were also upregulated, though these functions were not

significantly enriched (p=0.07; Suppl. Table 11). Similarly, upregulated KEGG pathways

included heat shock responses and RNA degradation (Suppl. Table 12). There were 236

downregulated genes which mapped to GO terms or KEGG pathways (Fig. 5A). Among these

were 21 DEGs with transferase functions, 20 with function related to the cell nucleus and 19

DEGs involved in ATP binding (Suppl. Table 11). Additionally, there were 12 DEGs associated

with kinases, 5 that had cyclin functions and 5 associated with DNA replication (Suppl. Table

11). Downregulated KEGG pathways included glycolysis/gluconeogenesis (*ALDOB*, *GAPDH*;

Suppl. Table 12).

Offspring and paternal warm-acclimation combined to influence the differential

expression of similar numbers and sorts of pathways as did maternal warm-acclimation, again

supporting our hypothesis. There were 187 upregulated DEGs mapped to GO terms or KEGG

pathways (Fig. 5B). The combination of offspring and paternal warm-acclimation had the effect

of upregulating 4 DEGs that have proteases/peptidases functions and 2 DEGs involved in

glycolysis (Suppl. Table 13). There were 2 upregulated DEGs involved in complex IV of the

respiratory chain, however, this function was not significantly enriched (p=0.07; Suppl. Table

13). Similar to the combined effect of offspring and maternal warm-acclimation, offspring and

paternal warm-acclimation upregulated KEGG pathways involved in cell signaling/inflammation

(*mapk14a*), but offspring and paternal warm-acclimation uniquely upregulated KEGG pathways

associated with protein folding (*HYOU1*) and insulin regulation (*foxo1a*; Suppl. Table 14). A

similar number (n = 180) of DEGs mapped to downregulated GO terms or KEGG pathways (Fig.

5B). Enriched downregulated functions included ATP binding (14 DEGs), hydrolase/protease

activity (13 DEGs), DNA replication (4 DEGs) and cyclin function (4 DEGs; Suppl. Table 13).

There were also 2 DEGs associated with tRNA ligase activity and 2 DEGs related to peroxisome

function (Suppl. Table 13). Downregulated KEGG pathways included glycolysis (*ALDOB*,

GAPDH), amino acid metabolism (*hpda*, *ALDH4A1, RRM1, RRM2, tdo2a*), fatty acid

metabolism (*ELOVL6*; Suppl. Table 14).

Of all groups compared so far, the combined multigenerational effect of offspring,

maternal and paternal warm-acclimation had the greatest effect on differential gene expression in

the offspring with a total of 471 identified DEGs (Fig. 5C). There were 230 upregulated DEGs

mapped to GO terms or KEGG pathways (Fig. 5C) with 41 of these upregulated DEGs

associated with signalling function (Suppl. Table 15). Immunoglobulin and protease/peptidase

activity were also enriched (9 and 5 DEGs, respectively; Suppl. Table 15). There were 2 DEGs

involved in the respiratory chain complex IV but complex IV was not significantly enriched

(*p*=0.07; Suppl. Table 15). Some of the affected KEGG pathways included the heat shock

response (*hsp90aa1.1*), signalling pathways associated with inflammation (*mapk14a*) and insulin

regulation (*foxo1a*; Suppl. Table 16). There were 241 downregulated DEGs mapped to GO terms

or KEGG pathways (Fig. 5C). Enriched functions included transferase activity (24 DEGs), DNA

replication (6 DEGs) and cyclin functions (4 DEGs; Suppl. Table 15). Binding functions were

downregulated, such as those related to magnesium, nucleic acid and iron binding (5 DEGs each;

Suppl. Table 15). Downregulated DEGs involved in KEGG pathways included peroxisome

activity (*MPV17L2*), glycolysis/gluconeogenesis (*ALDOB*), and response to DNA damage

(*DDIT4*; Suppl. Table 16).

Summary of general trends

We have identified five noteworthy trends when considering the experimental results all

together: 1) Within-generation (offspring) warm-acclimation had a larger effect on the overall

number of mapped DEGs compared to transgenerational contributions from warm-acclimated

parents (mother, father or both). Unsurprisingly, the combination of offspring, maternal and

paternal warm-acclimation had the greatest effect on the level of differential expression in the

offspring with an overall 471 mapped DEGs. 2) Cell signalling processes were upregulated (29

DEGs) with offspring (within-generation) warm-acclimation, but maternal or paternal

(transgenerational) warm acclimation had the effect of downregulating cell-signalling functions

(14-16 DEGs). When both parents were warm-acclimated, cell signalling was upregulated (16

DEGs). Cell signalling was most enriched (41 DEGs) with multigenerational (offspring,

maternal and paternal) warm-acclimation. 3) Transferase functions appeared to be regulated

mainly by transgenerational (parental) warm acclimation rather than with within-generation

(offspring) warm-acclimation, however, the degree of enrichment (i.e. gene count) and the

direction of the effect (up- or downregulation) appeared to depend upon whether one or both

parents were warm-acclimated. 4) Regulation of cyclin functions appeared to be primarily under

the control of the offspring because this process was only downregulated when offspring were

warm-acclimated, thus not influenced by transgenerational (either parent or both) warm-

acclimation. 5) Mothers seemed to have an influence over muscle functions (calcium binding,

troponin synthesis, ZASP) in their offspring, upregulating these functions when mothers

experienced a warmer environment, whereas there was no evidence of paternal influence on

offspring muscle functions.

Discussion

There was evidence to support our hypothesis that genes associated with metabolism,

growth and thermal stress/tolerance would be differentially expressed in juvenile lake trout,

depending on the acclimation temperatures of the offspring as well as their parents. Genes

involved in growth, heat shock and hypoxia responses, and metabolic pathways were

differentially expressed with within-generation, transgenerational and multigenerational warm-

acclimation. The degree to which each gene was affected depended on whether it was the

offspring or parent that had been warm-acclimated, with the offspring thermal experience often

having a stronger effect compared to the parental thermal experience. Maternal and paternal

warm-acclimation sometimes had opposing effects on the direction of differential expression of

certain genes in their offspring with one parent stimulating gene upregulation and the other

parent stimulating downregulation.

The general effect of warming on differential gene expression

Across all comparisons there were several DEGs that were indicative of thermal stress.

Among these were upregulated DEGs associated with molecular chaperone and co-chaperone

functions (*hsp90aa1.1, hsp70, hsp30, hsp7c, hspB1, hspB7, HYOU1, FKBP5* and *FKBP8*).

HYOU1 in particular was upregulated with within- or multigenerational warm-acclimation, but

not with transgenerational warm-acclimation in isolation (Suppl. Table 17). *HYOU1* (also known

as *GRP170* and *HSP12A*) is part of the heat shock protein 70 family and its expression is

primarily induced with hypoxia. Although, upregulation of this gene has previously been

observed in response to heat stress in fish (Huang et al. 2018; Bilyk et al. 2021), its expression

may be linked to a hypoxia response or may potentially reflect the increased demands on aerobic

metabolism/oxidative stress (Olsvik et al. 2013; Huang et al. 2018). The temperature-controlled

water in the source tank supplying the respirometer setup was supplemented with compressed

oxygen at higher temperatures in an effort to maintain saturation levels and minimize exposure to

hypoxia. However, we must acknowledge the possibility that some hypoxic responses may still

have occurred.

In general, within- and transgenerational warm-acclimation resulted in upregulated

mitogen-activated protein kinases (MAPK): *MAPK3*, *MAPK14a* and *MAPK14b* (also known as

ERK1, *p38a* and *p38b*, respectively). MAPKs are important in the regulation of many processes,

including mitosis during growth and development. They are also upregulated in response to

various stressors, such as osmotic, hypoxic and thermal stress (Cowan and Storey 2003; Tian et

al. 2019). MAPKs are associated with the induction of HSPs, for example, *MAPK3* has been

shown to induce expression of *HSP70* in zebrafish under thermal stress (Keller et al. 2008). Both

MAPK3 and *HSP70* were upregulated in warm-acclimated lake trout offspring with warm-

acclimated mothers, suggesting that these fish relied on MAPK signaling cascades for resilience

to warming. This is perhaps not surprising given that MAPK cascades and the induction of HSPs

are evolutionarily conserved across numerous taxa (Krens et al. 2006; Kostenko et al. 2011), and

that upregulation of these genes has been observed previously in other taxa following heat stress

(Keller et al. 2008; Liu et al. 2021). Other affected pathways indicating upper thermal stress

included peroxisome activity (*HMGCLL1*), DNA damage (*DDIT4*) and apoptotic signaling

(*pycard*), though whether these DEGs were up- or downregulated depended on whether it was

the parental or offspring generation that had experienced warming. Taken together, these DEGs

indicate increasing temperatures caused a stress response in lake trout, which we would expect

given that each offspring was subjected to an acute temperature increase (+2°C/h) and these

DEGs were likely upregulated to respond to cellular/protein damage occurring at their upper

critical thermal limit (Penney et al. 2021).

There was a general enrichment of pathways associated with metabolism, including genes

such as *COX1*, *COX2, GAPDH* and *ALDOB*. In general, *GAPDH* and *ALDOB* were

downregulated with within- or transgenerational warm acclimation, however, *COX1* and *COX2*

were upregulated in warm-acclimated offspring with warm-acclimated mothers following an

acute temperature challenge. The effect of acute warming on differential expression of

cytochrome c oxidase subunits is equivocal in fish: for example, acute exposure to warmer

temperatures resulted in downregulation of *COX1* and *COX2* in the liver of yellow croaker (Qian

and Xue 2016) but upregulation of *COX1* in the liver of juvenile lake whitefish (Zak and Manzon

2019). In our experiment, the matching of offspring and maternal thermal experience increased

COX gene expression to a greater extent compared to other groups, potentially allowing

offspring to meet the energetic demands associated with acute warming. We did not see

upregulation of any other COX subunits, but it seems that expression of COX subunits does not

follow stoichiometric expectations (Bremer and Moyes 2014) which may explain why we

observed upregulation of only two of the 14 subunits that make up the COX holoenzyme.

We found some indication that warm-acclimation and upper thermal stress affected the

growth of the lake trout offspring. We expected to see upregulation of some growth-related

genes given that the warm-acclimated offspring were heavier than the cold-acclimated offspring

at the time of sampling (Penney et al. 2021). Common molecular indicators of growth in fish

include genes encoding growth hormone (*GH*) and insulin-like growth factor-1 (*IGF-1*), and

genes related to thyroid function (Hevrøy et al. 2015; Li et al. 2021). We did not observe

differential expression with *GH* or *IGF-1* in our experiment, but the thyroid receptors *TRα* and

TRβ were downregulated in offspring that experienced warm-acclimation. Thyroid hormones

have important roles in growth and development (reviewed by Deal and Volkoff 2020), however,

it is curious that *TRα* and *TRβ* were downregulated given we showed previously that warm-

acclimated lake trout grew larger than the cold-acclimated group (Penney et al. 2021). Indicators

of growth, measured as expression of *GH* and *IGF-1*, increased in *Labeo rohita* at slightly

warmer acclimation temperatures but was suppressed at higher acclimation temperatures

(Shahjahan et al. 2021). We collected liver samples at each fish's CTMax, so it is possible that

fish were responding to the stress of the acute temperature increase and may have suppressed

growth at this point in favour of a heat stress response. Lastly, there is evidence suggesting that

growth hormone signaling pathways in the muscle are more sensitive to elevated temperature

than in the liver of Atlantic salmon and rainbow trout (Hevrøy et al. 2015), thus it is possible that

we may have seen a stronger transcriptional response for growth pathways in the muscle

compared to the liver. Confirmation of this would require further experimentation.

Transcriptomic responses to warm-acclimation across generations

We found that an offspring's thermal acclimation often had a greater influence on the

level (LFC) of gene expression than did the thermal experience of its parents and in some cases

multigenerational (offspring and parental) warm acclimation had an additive effect. For example,

cold-acclimated offspring downregulated the molecular chaperone *HYOU1* when either parent

was warm-acclimated (LFC maternal: -1.21, paternal: -0.94; Suppl. Table 17), however, a

combination of offspring and parental warm-acclimation upregulated *HYOU1* (LFC offspring &

maternal: 0.72, offspring & paternal 0.80, offspring & both parents 1.14), though to a lesser

extent than when offspring were warm-acclimated in isolation (LFC 1.29; Suppl. Table 17). This

trend suggests that within- and transgenerational warm acclimation had an additive effect on the

expression level of *HYOU1*. The gene *MAPK14a* was upregulated whenever the offspring were

warm-acclimated, regardless of parental thermal experience (LFC ranged from 1.19 to 2.06) and

seemed to override the isolated effect of transgenerational (parental) warm-acclimation (LFC

ranged from -0.67 to 0.52; Suppl. Table 17). The idea of WGP overriding TGP has been

presented before (Shama 2017), and our experiment provides evidence of this phenomenon at the

transcriptomics level. It has been suggested that transgenerational effects are strongest in early-

juvenile life stages (Yin et al. 2019) and it would be interesting to see if this is reflected at the

transcriptomics level by looking at gene expression at different life stages following within- and

transgenerational warm-acclimation.

Compared to within-generation (offspring) warm-acclimation, transgenerational warm-

acclimation (parents only) had only a modest effect on differential gene expression in lake trout

offspring, supporting our previous study of whole animal metabolic responses to acute warming

(Penney et al. 2021). Transgenerational effects can include parental effects (e.g. egg provisioning

and transfer of sperm cell cytoplasmic components) and epigenetics (Dai et al. 2020; Spadafora

2020; Venney et al. 2020). For example, brook trout (*S. fontinalis*) offspring incurred more

differentially methylated regions due to transgenerational warm-acclimation than with within-

generation warm-acclimation (Venney et al. 2022). In the current study, maternal and paternal

warm-acclimation separately elevated DEGs functioning in the DNA damage response (*DDIT4*)

and heat shock proteins (maternal only: *HSPB1*, *HSPB7*). These transgenerationally elevated

pathways associated with DNA damage and HSPs supports the idea of an adaptive response

whereby parents may improve offspring fitness when they correctly anticipate their offspring's

environment (Bateson et al. 2014; Bernatchez 2016; Ashe et al. 2021).

Our prediction that transgenerational warm acclimation would upregulate metabolic

processes in the lake trout offspring was partially supported. We observed differential expression

of the genes GAPDH, ALDOB and ENOA which relate to the glycolytic enzymes

glyceraldehyde-3-phosphate dehydrogenase, aldolase and enolase, respectively. These three

transcripts tended to be downregulated in all groups except for the two groups that included the

combined effect of offspring cold acclimation and paternal warm acclimation which indicates a

paternal transgenerational effect of increasing offspring metabolism (Suppl. Table 17). Despite

this evidence, we cannot be fully confident that glycolysis was upregulated in these offspring

because, while the differential expression of GAPDH, ALDOB and ENOA suggests an effect of

warm acclimation on the rate of glycolysis in the offspring, these three transcripts did not

significantly enrich the 'glycolysis' pathway in our analysis. Also, these genes may be

constitutively expressed and have even been proposed as housekeeping genes for experiments

exploring differential gene expression, however, the expression level may vary depending on

species, tissue type and treatment (Hori et al. 2012; Purohit et al. 2016; Shekh et al. 2017).

Although transgenerational warming can result in larger offspring (Salinas et al. 2012),

we did not find support for a transgenerational effect of warm-acclimation on growth in lake

trout at the transcriptomics level, consistent with a lack of response at that whole animal level

(Penney et al. 2021). Lastly, despite some genes being transgenerationally expressed in the same

direction by either the maternal or paternal warm-acclimation, none of these genes appeared to

have an additive effect on expression level when both parents were simultaneously warm-

acclimated (Suppl. Table 14).

The combination of offspring, maternal and paternal warm-acclimation had the greatest

effect on the overall number genes that were differentially expressed and, for some genes, the

greatest effect on up- or downregulation. This adds to a growing literature suggesting that

transgenerational plasticity occurs when offspring and parental thermal experiences are

coincident (Shama et al. 2014; Donelson et al. 2017; Yin et al. 2019). Among all of the groups

compared, the combination of offspring and parent warm-acclimation had the strongest effect on

cytochrome c oxidase subunit upregulation. Differential expression of COX-related genes has

elsewhere been recorded with transgenerational acclimation (Shama et al. 2016; Bernal et al.

2018). Also, the expression level of *COX1* and *COX2* subunits (among 7 of 14 subunits tested)

correlated with cytochrome c oxidase activity in muscle of fish (Bremer and Moyes 2014). Our

result could be indicative of a higher metabolic rate in warm-acclimated offspring with warm-

acclimated parents, although neither resting or peak metabolic rates differed between warm-

acclimated offspring from warm- or cold-acclimated parents (Penney et al. 2021). It is worth

noting, however, that gene expression does not necessarily confer a protein to function given that

post-transcriptional and post-translational activity could also tag a protein for degradation.

There were a few instances where some genes such as *GAPDH* and *CNOT1* appeared to

be simultaneously upregulated and downregulated within treatments, though why this occurred is

not immediately clear. One possible explanation is that the direction of the responses were sex-

specific (Best et al. 2018; Bell and Hellman 2019; Bautista et al. 2020). For example, in

stickleback, mothers can influence the direction (up- or downregulation) of differentially

expressed genes depending on whether the offspring is male or female (Metzger and Schulte

2016). We did not test for the sex of the offspring so we cannot confirm this for the lake trout.

Conclusions

We found that transgenerational thermal acclimation in isolation (either or both parents)

or in combination with offspring (within-generation) thermal acclimation can influence the

strength and direction of differential expression of genes associated with metabolism and thermal

stress/tolerance in lake trout experiencing acute thermal stress. In general, transgenerational

warm acclimation had a smaller effect on differential gene expression compared to the strength

of within-generation warm acclimation and, for some genes, transgenerational acclimation had

opposing effects on offspring gene expression (mothers vs. fathers, or parents vs. offspring). The

influence of transgenerational acclimation on offspring responses to environmental changes is

complex, may not always be apparent in the phenotype or may have weak effects (Uller et al.

2013; Bautista and Burggren 2019; Lee et al. 2020; Sánchez-Tójar et al. 2020). However, the

patterns of gene expression we observed here could present itself in the offspring's phenotype

and we have previously shown that transgenerational environmental mismatching can have

maladaptive effects in lake trout (Penney et al. 2021).

Without sufficient time or standing genetic variation necessary for adaptation (Willi et al.

2006, Comte and Olden 2017, Wilson 2017) populations of many cold-adapted species will need

to rely on plastic thermal acclimation to withstand chronic increases in environmental

temperature. This may be particularly important for stenothermal coldwater species like lake

trout, and for landlocked or isolated populations with finite adaptive resources. It is therefore

critically important that we improve our understanding of the transcriptomic and phenotypic

responses of organisms that are vulnerable to climate change by exploring the mechanisms

driving phenotypic changes, and how those mechanisms act to mediate stresses associated with

chronic warming.

Acknowledgements

The Ontario Ministry of Natural Resources and Forestry (OMNRF) Fish Culture Section

provided adult lake trout, rearing space and logistical support at the White Lake Fish Culture

Station. Bill Sloan, Scott Ferguson, Anne McCarthy (OMNRF) and John Dewart provided

assistance with breeding adult fish; Bill Sloan and Scott Ferguson also provided invaluable

logistical support for all aspects of fish husbandry from care of the juvenile trout to assisting with

experiments at the OMNRF Codrington Fisheries Research Facility. Caleigh Smith (OMNRF)

genotyped adult and juvenile fish for parentage assignment and family identification of juveniles.

Paul Craig and Heather Ikert (University of Waterloo) kindly provided advice and guidance on

the enrichment analysis. Madison Wikston provided advice for figure presentation. Paul Craig

and Graham Scott provided feedback on early drafts of the manuscript.

Research funding was provided by the Canada-Ontario Great Lakes Agreement (COA) to

CCW, by the Ontario Federation of Anglers and Hunters (OFAH) to CMP, and a Natural

Sciences and Engineering Research Council (NSERC) Discovery Grant to GB.

Data archiving statement: Data for this study are available at: *to be completed after manuscript*

is accepted for publication.

Literature Cited

Akbarzadeh, A., Leder, E.H. (2016). Acclimation of killifish to thermal extremes of hot spring: transcription of gonadal and liver heat shock genes. *Comparative Biochemistry and Physiology Part A: Molecular & Integrative Physiology*, 191, 89-97.

Ashe, A., Colot, V., Oldroyd, B.P. (2021). How does epigenetics influence the course of evolution? *Philosophical Transactions of the Royal Society B: Biological Sciences*, 376(1826), 20200111.

Bateson, P., Gluckman, P., Hanson, M. (2014). The biology of developmental plasticity and the Predictive Adaptive Response hypothesis. *The Journal of Physiology*, 592, 2357-2368.

Bautista, N.M., Burggren, W.W. (2019). Parental stressor exposure simultaneously conveys both adaptive and maladaptive larval phenotypes through epigenetic inheritance in the zebrafish (*Danio rerio*). *Journal of Experimental Biology*, 222, jeb208918.

Bautista, N.M., Crespel, A., Crossley, J., Padilla, P., Burggren, W. (2020). Parental transgenerational epigenetic inheritance related to dietary crude oil exposure in *Danio rerio*. *Journal of Experimental Biology*, 223, jeb222224.

Beitinger, T.L., Bennett, W.A. (2000). Quantification of the role of acclimation temperature in temperature tolerance of fishes. *Environmental Biology of Fishes*, 58, 277-288.

Bell, A.M., Hellmann, J. (2019). An integrative framework for understanding the mechanisms and multigenerational consequences of transgenerational plasticity. *Annual Review of Ecology, Evolution and Systemics*, 50, 97–118.

Bennett, S., Duarte, C.M., Marbà, N., Wernberg, T. (2019). Integrating within-species variation in thermal physiology into climate change ecology. *Philosophical Transactions of the Royal Society B: Biological Sciences*, 374, 20180550.

Bernal, M.A., Donelson, J.M., Veilleux, H.D., Ryu, T., Munday, P.L., Ravasi, T. (2018).

Phenotypic and molecular consequences of stepwise temperature increase across generations in a

coral reef fish. *Molecular Ecology*, 27, 4516-4528.

Bernatchez, L. (2016). On the maintenance of genetic variation and adaptation to environmental

change: considerations from population genomics in fishes. *Journal of Fish Biology*, 89, 2519-

2556.

Best, C., Ikert, H., Kostyniuk, D. J., Craig, P. M., Navarro-Martin, L., Marandel, L., Mennigen,

J. A. (2018). Epigenetics in teleost fish: from molecular mechanisms to physiological

phenotypes. *Comparative Biochemistry and Physiology Part B: Biochemistry and Molecular*

Biology, 224, 210-244.

Bilyk, K.T., Zhuang, X., Vargas-Chacoff, L., Cheng, C.C. (2021). Evolution of chaperome gene

expression and regulatory elements in the antarctic notothenioid fishes. *Heredity*, 126, 424-441.

Bonduriansky R. (2021). Plasticity across generations. In: D.W. Pfennig (Ed.). Phenotypic

Plasticity & Evolution: Causes, Consequences, Controversies (pp. 327-348). Boca Raton: CRC

Press/Taylor & Francis Group.

Bremer, K., Moyes, C.D. (2014). mRNA degradation: an underestimated factor in steady-state

transcript levels of cytochrome c oxidase subunits? *Journal of Experimental Biology,* 217, 2212-

2220.

Burkhead, N.M. (2012). Extinction rates in North American freshwater fishes, 1900-

2010. *BioScience*, 62: 798-808.

Casselman, J.M . (2008). Effects of climate and climate change on lake trout populations and

fisheries. In: K.H. Mills, M. Dyck, & L.A. Harwood (Eds.). Proceedings of the Second North

American Lake Trout Symposium (pp. 103-113). Yellowknife, NT, Canada: Canadian Technical

Report of Fisheries and Aquatic Sciences.

Chomczynski, P., Sacchi, N. (2006). The single-step method of RNA isolation by acid

guanidinium thiocyanate–phenol–chloroform extraction: twenty-something years on. *Nature*

Protocols, 1, 581-585.

Comte, L., Olden, J.D. (2017). Climatic vulnerability of the world's freshwater and marine

fishes. *Nature Climate Change*, 7, 718–722.

Cowan, K.J. Storey, K.B. (2003). Mitogen-activated protein kinases: new signaling pathways

functioning in cellular responses to environmental stress. *Journal of Experimental Biology*, 206,

1107-1115.

Crozier, L.G. J.A. Hutchings. (2014). Plastic and evolutionary responses to climate change in

fish. *Evolutionary Applications*, 7, 68-87.

Dai, Z., Ramesh, V., Locasale, J.W. (2020). The evolving metabolic landscape of chromatin

biology and epigenetics. *Nature Reviews Genetics*, 21, 737-753.

Dammark, K. B., Ferchaud, A.L., Hansen, M.M., Sørensen, J.G. (2018). Heat tolerance and gene

expression responses to heat stress in threespine sticklebacks from ecologically divergent

environments. *Journal of Thermal Biology*, 75, 88-96.

Deal, C.K., Volkoff, H. (2020). The role of the thyroid axis in fish. *Frontiers in*

Endocrinology, 11, 596585.

Donelan, S.C, Hellman, J.K., Bell, A.M., Luttbeg, B., Orrock, J.L., Sheriff, M.J., Sih, A. (2020).

Transgenerational plasticity in human-altered environments. *Trends in Ecology & Evolution*, 35,

115-124.

Donelson, J.M., Munday, P.L., McCormick, M.I., Pritcher, C.R. (2012). Rapid transgenerational

acclimation of a tropical reef fish to climate change. *Nature Climate Change*, 2, 30–32.

Donelson, J.M., Sunday, J.M., Figueira, W.F., Gaitán-Espitia, J.D., Hobday, A.J., Johnson, C.R.,

Leis, J.M., Ling, S.D., Marshall, D., Pandolfi, J.M., Pecl, G., Rodgers, G.G., Booth, D.J.,

Munday, P.L. (2019). Understanding interactions between plasticity, adaptation and range shifts

in response to marine environmental change. *Philosophical Transactions of the Royal Society B:*

Biological Sciences, 374, 20180186.

Edsall, T.A., Cleland, J. (2000). Optimum temperature for growth and preferred temperatures of

age-0 lake trout. *North American Journal of Fisheries Management*, 20, 804-809.

Grabherr, M.G., Haas, B.J., Yassour, M., Levin, J.Z., Thompson, D.A., Amit, I., Adiconis, X.,

Fan, L., Raychowdhury, R., Zeng, Q., Chen, Z., Mauceli, E., Hacohen, N., Gnirke, A., Rhind, N.,

di Palma, F., Birren, B.W., Nusbaum, C., Lindblad-Toh, K., Friedman, N., Regev, A. (2011).

Full-length transcriptome assembly from RNA-Seq data without a reference genome. *Nature*

Biotechnology, 29, 644-652.

Greenspoon, P.B., Spencer, H.G. (2018). The evolution of epigenetically mediated adaptive

transgenerational plasticity in a subdivided population. *Evolution*, 72, 2773-2780.

Guzzo, M.M., Blanchfield, P.J. (2017). Climate change alters the quantity and phenology of

habitat for lake trout (*Salvelinus namaycush*) in small Boreal Shield lakes. *Canadian Journal of*

Fisheries and Aquatic Sciences, 74, 871-884.

Haas, B.J., Papanicolaou, A., Yassour, M., Grabherr, M., Blood, P.D., Bowden, J., Couger, M.

B., Eccles, D., Li, B., Lieber, M., MacManes, M.D., Ott, M., Orvis, J., Pochet, N., Strozzi, F.,

Weeks, N., Westerman, R., William, T., Dewey, C.N., Henschel, R., LeDuc, R.D. Friedman, N.,

Regev, A. (2013). De novo transcript sequence reconstruction from RNA-seq using the Trinity

platform for reference generation and analysis. *Nature Protocols*, 8, 1494-1512.

Hayhoe, K., VanDorn, J., Croley II, T., Schlegal, N., Nwuebbles, D. (2010). Regional climate

change projections for Chicago and the US Great Lakes. *Journal of Great Lakes Research*, 36

(Suppl 2), 7-21.

Hellmann, J.K., Carlson, E.R., Bell, A.M. (2020). Sex-specific plasticity across generations II:

Grandpaternal effects are lineage specific and sex specific. *Journal of Animal Ecology*, 89, 2800-

2812.

Hevrøy, E.M., Tipsmark, C.K., Remø, S.C., Hansen, T., Fukuda, M., Torgersen, T., Vikeså, V.,

Olsvik, P.A., Waagbø, R., Shimizu, M. (2015). Role of the GH-IGF-1 system in Atlantic salmon

and rainbow trout postsmolts at elevated water temperature. *Comparative Biochemistry and*

Physiology Part A: Molecular & Integrative Physiology, 188, 127-138.

Hori, T. S., Gamperl, A. K., Afonso, L. O., Johnson, S. C., Hubert, S., Kimball, J., Bowman, S.,

Rise, M. L. (2010). Heat-shock responsive genes identified and validated in Atlantic cod (Gadus

morhua) liver, head kidney and skeletal muscle using genomic techniques. *BMC Genomics,* 11,

1-22.

Huang, D.W., Sherman, B.T., Lempicki, R.A. (2009A). Systematic and integrative analysis of

large gene lists using DAVID Bioinformatics Resources. *Nature Protocols*, 4, 44-57.

Huang, D.W., Sherman B.T., Lempicki, R.A. (2009B). Bioinformatics enrichment tools: paths

toward the comprehensive functional analysis of large gene lists. *Nucleic Acids Research*, 37, 1-

13.

Huang, J., Li, Y., Liu, Z., Kang, Y., Wang, J. (2018). Transcriptomic responses to heat stress in

rainbow trout *Oncorhynchus mykiss* head kidney. *Fish & Shellfish Immunology*, 82, 32-40.

IPCC (2021). Summary for Policymakers. In: V. Masson-Delmotte, P. Zhai, A. Pirani, S.L.

Connors, C. Péan, S. Berger, N. Caud, Y. Chen, L. Goldfarb, M.I. Gomis, M. Huang, K. Leitzell,

E. Lonnoy, J.B.R. Matthews, T.K. Maycock, T. Waterfield, O. Yelekçi, R. Yu, & B. Zhou

(Eds.), Climate Change 2021: The Physical Science Basis. Contribution of Working Group I to

the Sixth Assessment Report of the Intergovernmental Panel on Climate Change (pp. 3−32).

Cambridge, United Kingdom and New York, NY, USA:Cambridge University Press.

Jablonka, E., Lachmann, M., Lamb, M.J. (1992). Evidence, mechanisms and models for the

inheritance of acquired characters. *Journal of Theoretical Biology*, 158, 245-268.

Keller, J.M., Escara-Wilke, J.F., Keller, E.T. (2008). Heat stress-induced heat shock protein 70

expression is dependent on ERK activation in zebrafish (*Danio rerio*) cells. *Comparative*

Biochemistry and Physiology Part A: Molecular & Integrative Physiology, 150, 307-314.

Kelly, N.I., Burness, G., McDermid, J.L., Wilson, C.C. (2014). Ice age fish in a warming world:

minimal variation in thermal acclimation capacity among lake trout (*Salvelinus namaycush*)

populations. *Conservation Physiology*, 2, doi:10.1093/conphys/cou025.

Kincaid, H.L. (1977). Rotational line crossing: an approach to the reduction of inbreeding

accumulation in trout brood stocks. *The Progressive Fish-Culturist*, 39, 179-181.

Kostenko, S., Dumitriu, G., Lægreid, K.J., Moens, U. (2011). Physiological roles of mitogen-

activated-protein-kinase-activated p38-regulated/activated protein kinase. *World Journal of*

Biological Chemistry, 2, 73-89.

Krens, S.G., He, S., Spaink, H.P., Snaar-Jagalska, B.E. (2006). Characterization and expression

patterns of the MAPK family in zebrafish. *Gene Expression Patterns*, 6, 1019-1026.

Lee, W.S., Salinas, S., Lee, Y.R., Siskidis, J.A., Mangel, M., Munch, S.B. (2020). Thermal

transgenerational effects remain after two generations. *Ecology and Evolution*, 10, 11296-11303.

Li, B., & Dewey, C.N. (2011). RSEM: accurate transcript quantification from RNA-Seq data

with or without a reference genome. *BMC Bioinformatics*, 12, 1-16.

Li, S., Guo, H., Chen, Z., Jiang, Y., Shen, J., Pang, X., Li, Y. (2021). Effects of acclimation

temperature regime on the thermal tolerance, growth performance and gene expression of a cold-

water fish, *Schizothorax prenanti*. *Journal of Thermal Biology*, 98, 102918.

Liu, Z., Huang, X., Yang, Z., Peng, C., Yu, H., Cui, C., Hu, Y., Wang, X., Xing, Q., Hu, J., Bao,

Z. (2021). Identification, characterization, and expression analysis reveal diverse regulated roles

of three MAPK genes in *Chlamys farreri* under heat stress. *Frontiers in Physiology*, 12, 688626.

Luu, I., Ikert, H., Craig, P.M. (2021). Chronic exposure to anthropogenic and climate related

stressors alters transcriptional responses in the liver of zebrafish (*Danio rerio*) across multiple

generations. *Comparative Biochemistry and Physiology Part C: Toxicology &*

Pharmacology, 240, 108918.

Marshall, D.J. (2015). Environmentally induced (co)variance in sperm and offspring phenotypes

as a source of epigenetic effects. *Journal of Experimental Biology*, 218, 107–113.

Martin N.V., Olver C.H. (1980). The lake charr, Salvelinus namaycush. In: E.K. Balon (Ed.),

Charrs: Salmonid Fishes of the Genus Salvelinus, Ed 1st (pp. 205–277). Netherlands: Springer.

McCaw, B.A., T.J. Stevenson, L.T. Lancaster. (2020). Epigenetic responses to temperature and

climate. *Integrative and Comparative Biology*, 60, 1469-1480.

McDermid, J.L.,Wilson, C.C., Sloan, W.N., Shuter, B.J. (2013). Intraspecific differences in

thermal biology of inland lake trout populations. *Transactions of the American Fisheries Society*,

142, 756–766.

Metzger, D.C., Schulte, P.M. (2016). Maternal stress has divergent effects on gene expression

patterns in the brains of male and female threespine stickleback. *Proceedings of the Royal*

Society B: Biological Sciences, 283, 20161734.

Morash, A.J., Speers-Roesch, B., Andrew, S., Currie, S. (2021). The physiological ups and
downs of thermal variability in temperate freshwater ecosystems. *Journal of Fish Biology*, 98,
1524-1535.
891

892   Olsvik, P.A., Vikeså, V., Lie, K.K., Hevrøy, E.M. (2013). Transcriptional responses to
893   temperature and low oxygen stress in Atlantic salmon studied with next-generation sequencing
894   technology. *BMC Genomics,* 14, 1-21.
895

896   OMNR Fish Culture Stocks Catalogue. (2005). Fish Culture Section. Peterborough, ON: Ontario
897   Ministry of Natural Resources.
898

899   Penney, C.M., Burness, G., Tabh, J.K.R., Wilson, C.C. (2021). Limited transgenerational effects
900   of environmental temperatures on thermal performance of a cold-adapted salmonid.
901   *Conservation Physiology*, 9, coab021.
902

903   Pérez-Ruzafa, A., Pérez-Marcos, M., Marcos, C. (2018). From fish physiology to ecosystems
904   management: Keys for moving through biological levels of organization in detecting
905   environmental changes and anticipate their consequences. *Ecological Indicators*, 90, 334-345.
906

907   Perrier, C., Ferchaud, A.L., Sirois, P., Thibault, I., Bernatchez, L. (2017). Do genetic drift and
908   accumulation of deleterious mutations preclude adaptation? Empirical investigation using
909   RADseq in a northern lacustrine fish. *Molecular Ecology*, 26, 6317–6335.
910

911   Petitjean, Q., Jean, S., Gandar, A., Côte, J., Laffaille, P., Jacquin, L. (2019). Stress responses in
912   fish: From molecular to evolutionary processes. *Science of the Total Environment*, 684, 371-380.
913

914   Purohit, G. K., Mahanty, A., Mohanty, B. P., Mohanty, S. (2016). Evaluation of housekeeping
915   genes as references for quantitative real-time PCR analysis of gene expression in the murrel
916   *Channa striatus* under high-temperature stress. *Fish Physiology and Biochemistry*, 42, 125-135.
917

Qian, B., Xue, L. (2016). Liver transcriptome sequencing and de novo annotation of the large

yellow croaker (*Larimichthy crocea*) under heat and cold stress. *Marine Genomics*, 25, 95-102.

Quinn, N.L., McGowan, C.R., Cooper, G.A., Koop, B.F., Davidson, W.S. (2011). Identification

of genes associated with heat tolerance in Arctic charr exposed to acute thermal

stress. *Physiological Genomics*, 43, 685-696.

Riley, S.C., Hansen, M.J., Krueger, C.C., Noakes, D.L.G., Muir, A.M. (2021). Introduction. The

Lake Charr: Biology, Ecology, Distribution, and Management. In: A.M. Muir, M.J. Hansen, C.C.

Krueger, & S.C. Riley (Eds.), The Lake Charr *Salvelinus namaycush*: Biology, Ecology,

Distribution, and Management, Fish & Fisheries Series 39 (pp. 1-12). Switzerland: Springer

Nature.

Ritchie, M.E., Phipson, B., Wu, D.I., Hu, Y., Law, C.W., Shi, W., Smyth, G.K. (2015). limma

powers differential expression analyses for RNA-sequencing and microarray studies. *Nucleic*

Acids Research, 43, e47-e47.

Rosenfeld, J., Richards, J., Allen, D., Van Leeuwen, T., Monnet, G. (2020). Adaptive trade-offs

in fish energetics and physiology: insights from adaptive differentiation among juvenile

salmonids. *Canadian Journal of Fisheries and Aquatic Sciences*, 77, 1243-1255.

Salinas, S., Munch, S.B. (2012). Thermal legacies: transgenerational effects of temperature on

growth in a vertebrate. *Ecology Letters*, 15, 159-163.

Sambrook, J., Russell, D.W. (2001). Molecular Cloning: a laboratory manual. Cold Spring

Harbor: Cold Spring Harbor Laboratory Press.

Sánchez-Tójar, A., Lagisz, M., Moran, N.P., Nakagawa, S., Noble, D.W., Reinhold, K. (2020).

The jury is still out regarding the generality of adaptive 'transgenerational' effects. *Ecology*

Letters, 23, 1715–1718.

Shahjahan, M., Zahangir, M.M., Islam, S.M.M., Ashaf-Ud-Doulah, M., Ando, H. (2021). Higher
acclimation temperature affects growth of rohu (*Labeo rohita*) through suppression of GH and
IGFs genes expression actuating stress response. *Journal of Thermal Biology*, 100, 103032.
952
953 Shama, L.N., Strobel, A., Mark, F.C., Wegner, K.M. (2014). Transgenerational plasticity in
954 marine sticklebacks: maternal effects mediate impacts of a warming ocean. *Functional Ecology*,
955 28, 1482–1493.
956
957 Shama, L.N., Mark, F.C., Strobel, A., Lokmer, A., John, U., Mathias Wegner, K. (2016).
958 Transgenerational effects persist down the maternal line in marine sticklebacks: gene expression
959 matches physiology in a warming ocean. *Evolutionary Applications*, 9, 1096–1111.
960
961 Shama, L.N. (2017). The mean and variance of climate change in the oceans: hidden
962 evolutionary potential under stochastic environmental variability in marine
963 sticklebacks. *Scientific Reports*, 7, 1-14.
964
965 Shekh, K., Tang, S., Niyogi, S., Hecker, M. (2017). Expression stability and selection of optimal
966 reference genes for gene expression normalization in early life stage rainbow trout exposed to
967 cadmium and copper. *Aquatic Toxicology,* 190, 217-227.
968
969 Smith, D.A., Jackson, D.A., Ridgway, M.S. (2020). Thermal habitat of brook trout in lakes of
970 different size. *Freshwater Science*, 39, 56-69.
971
972 Somero, G.N. (2010). The physiology of climate change: how potentials for acclimatization and
973 genetic adaptation will determine 'winners' and 'losers'. *Journal of Experimental Biology*, 213,
974 912–920.
975
976 Spadafora, C. (2020). Transgenerational epigenetic reprogramming of early embryos: a
977 mechanistic model. *Environmetal Epigenetics*, 6, dvaa009.
978

Tian, Y., Wen, H., Qi, X., Zhang, X., Li, Y. (2019). Identification of mapk gene family in
Lateolabrax maculatus and their expression profiles in response to hypoxia and salinity
challenges. *Gene*, 684, 20-29.

Uller, T., Nakagawa, S., English, S. (2013). Weak evidence for anticipatory parental effects in
plants and animals. *Journal of Evolutionary Biology*, 26, 2161-2170.

Veilleux, H.D., Ryu, T., Donelson, J.M., Van Herwerden, L., Seridi, L., Ghosheh, Y., Berumen,
M.L., Leggat, W., Ravasi, T., Munday, P. L. (2015). Molecular processes of transgenerational
acclimation to a warming ocean. *Nature Climate Change*, 5, 1074-1078.

Veilleux, H.D., Donelson, J.M., Munday, P.L. (2018). Reproductive gene expression in a coral
reef fish exposed to increasing temperature across generations. *Conservation Physiology*, 6, doi:
10.1093/conphys/ cox077.

Venney, C.J., Love, O.P., Drown, E.J., Heath, D.D. (2020). DNA methylation profiles suggest
intergenerational transfer of maternal effects. *Molecular Biology and Evolution*, 37, 540-548.

Venney, C.J., Wellband, K.W., Normandeau, E., Houle, C., Garant, D., Audet, C., Bernatchez, L.
(2022). Thermal regime during parental sexual maturation, but not during offspring rearing,
modulates DNA methylation in brook charr (*Salvelinus fontinalis*). *Proceedings of the Royal*
Society B: Biological Sciences, 289, 20220670.

1001

1002 Visser, M.E. (2008). Keeping up with a warming world; assessing the rate of adaptation to
1003 climate change. *Proceedings of the Royal Society B: Biological Sciences*, 275, 649–659.

1004

1005 Willi, Y., Van Buskirk, J., Hoffmann, A.A. (2006). Limits to the adaptive potential of small
1006 populations. *Annual Review of Ecology, Evolution and Systematics*, 37, 433–458.

1007

1008 Wilson, C. (2017). Genetic aspects of climate change influences on inland fishes and fisheries.
1009 *Fisheries*, 42, 125–126.

1010

1011   Wilson, C.C., Mandrak, N.E. (2021). Paleoecology. In: A.M. Muir, M.J. Hansen, C.C. Krueger,

1012   & S.C. Riley (Eds.), The Lake Charr *Salvelinus namaycush*: Biology, Ecology, Distribution, and

1013   Management, Fish & Fisheries Series 39 (pp. 41-68). Switzerland: Springer Nature.

1014

1015   Yin, J., Zhou, M., Lin, Z., Li, Q.Q., Zhang, Y.Y. (2019). Transgenerational effects benefit

1016   offspring across diverse environments: a meta-analysis in plants and animals. *Ecology Letters*,

1017   22, 1976–1986.

1018

1019   Zak, M.A., Manzon, R.G. (2019). Expression and activity of lipid and oxidative metabolism

1020   enzymes following elevated temperature exposure and thyroid hormone manipulation in juvenile

1021   lake whitefish (*Coregonus clupeaformis*). *General and Comparative Endocrinology*, 275, 51-64.

1022

1023   Zhang, X., Flato, G., Kirchmeier-Young, M., Vincent, L., Wan, H., Wang, X., Rong, R., Fyfe, J.,

1024   Li, G., Kharin, V.V. (2019). Changes in Temperature and Precipitation Across Canada. In: E.

1025   Bush & D.S. Lemmen (Eds.) Canada's Changing Climate Report (pp. 112-193). Ottawa, Ontario:

1026   Government of Canada.

1027

1028

1029

1030

1031

1032

1033

1034

1035

1036

1037

1038

1039

1040

1041 **Tables**

1042

1043 **Table 1**: Summary of the seven comparisons made to measure differential gene expression in juvenile lake trout following within- and
1044 transgenerational thermal acclimation.

	Control group	Treatment group	Effect measured
Within-generation warm-acclimation	Cold-acclimated offspring (C♀xC♂)	Warm-acclimated offspring (C♀xC♂)	Offspring warm-acclimation
		Cold-acclimated offspring (W♀xC♂)	Maternal warm-acclimation
Transgenerational warm-acclimation	Cold-acclimated offspring (C♀xC♂)	Cold-acclimated offspring (C♀xW♂)	Paternal warm-acclimation
		Cold-acclimated offspring (W♀xW♂)	Maternal & paternal warm-acclimation
		Warm-acclimated offspring (W♀xC♂)	Offspring & maternal warm-acclimation
Within- and transgenerational warm-acclimation	Cold-acclimated offspring (C♀xC♂)	Warm-acclimated offspring (C♀xW♂)	Offspring & paternal warm-acclimation
		Warm-acclimated offspring (W♀xW♂)	Offspring, maternal & paternal warm-acclimation

1045 Sample sizes were n=12 for the control group and n=6 for each treatment group. Parental crosses are given in parentheses where C =
1046 cold and W = warm, with mothers listed first (♀ x ♂).

1047

1048

1049 **Figures**
1050
1051

1052
1053

1054 **Figure 1**: Experimental treatment groups of lake trout (*Salvelinus namaycush*). Two males and
1055 two females each from cold- and warm-acclimation tanks were bred in a full factorial cross,
1056 resulting in four families per parental treatment group (C♀xC♂, W♀xC♂, C♀xW♂, W♀xW♂; one
1057 each depicted here for simplicity). Offspring from each family was divided and acclimated to
1058 either a cold (11°C) or warm (15°C) temperature.

Figure 2: Heatmap comparing the effect of warm-acclimation on offspring (within-generation) and parental (transgenerational) gene expression in juvenile lake trout. Changes in expression are represented as log2-fold change (LFC; blue = downregulate, red = upregulated) in offspring (n = 6) for each treatment compared against the control group (11°C C♀xC♂; n = 12) where both parents and offspring were cold-acclimated. Labels for each column indicate whether the offspring, parents or both were acclimated to higher ("warm") temperature. Parental treatment (acclimation) is given in parentheses where "C" refers to a cold-acclimated parent and "W" refers to a warm-acclimated parent, mothers are listed first. The list of genes along with their

1084   LFC values and FDR (adjusted p-value) can be found in the supplementary information (Suppl.
1085   Table 3).
1086
1087
1088

Figure 3: Enrichment of pathways using genes that were differentially expressed in response to warm-acclimation of juvenile lake trout offspring (n = 6) when compared against the control group (11°C C♀xC♂; n = 12) where both parents and offspring were cold-acclimated. Genes were expressed at log2-fold change (LFC) of ≤ -1 or ≥ +1 and were mapped to GO categories (BP: biological processes, CC: cellular components, MF: molecular function) and KEGG pathways. Treatment groups include representatives from each family.

1104

Figure 4: Enrichment of pathways using genes that were differentially expressed in response to transgenerational (parental) warm-acclimation of juvenile lake trout offspring when the mothers (A) or the fathers (B) were warm-acclimated, and when both parents (C) were warm-acclimated. Treatment groups (n = 6 each) were compared against control group (11°C C♀xC♂; n = 12) where both parents and offspring were cold-acclimated. Genes were expressed at log2-fold change (LFC) of ≤ -1 or ≥ +1 and were mapped to GO categories (BP: biological processes, CC: cellular components, MF: molecular function) and KEGG pathways. Treatment groups include representatives from each family.

1113

Figure 5: Enrichment of pathways using genes that were differentially expressed in response to the combined effect of within- and transgenerational warm-acclimation in juvenile lake trout offspring. Offspring were warm-acclimated along with warm-acclimated mothers (A), fathers (B) or both parents (C). Treatment groups (n = 6 individuals per group) were compared against a control group (11°C C♀xC♂; n = 12) where both parents and offspring were cold-acclimated. Genes were expressed at log2-fold change (LFC) of ≤ -1 or ≥ +1 and were mapped to GO categories (BP: biological processes, CC: cellular components, MF: molecular function) and KEGG pathways. Treatment groups include representatives from each family.