Transcriptomic responses to within- and transgenerational environmental warming in a cold-adapted salmonid

Running title: Lake trout transcriptomic responses to warming

Chantelle M. Penney[1], Gary Burness[2], Gerardo Zapata[3], François Lefebvre[3] and Chris C. Wilson[4]

[1] Environmental and Life Sciences Graduate Program, Trent University, Peterborough, Ontario, K9J 7B8

[2] Department of Biology, Trent University, Peterborough, Ontario, K9L 0G2

[3] Canadian Centre for Computational Genomics, McGill University, Montréal, Québec, H3A 1A3

[4] Aquatic Research and Development Section, Ontario Ministry of Natural Resources, Trent University, Peterborough, Ontario, K9L 0G2

Author for correspondence: Chantelle Penney, Environmental and Life Sciences Graduate Program, Trent University, Peterborough, Ontario, K9L 0G2, chantellepenney@trentu.ca

Authors' contributions: CMP, GB and CCW conceived the ideas and designed methodology; CMP collected and analysed the data, and led the writing of the manuscript. FL and GZ completed the bioinformatics. All authors gave final approval for publication.

Conflict of interest: The authors have no conflict of interest to declare.

Keywords: acclimation, climate change, RNA sequencing, *Salvelinus namaycush*, transgenerational plasticity, transcriptomics.

Abstract

Cold-adapted species are particularly threatened by climate change as rates of environmental warming outpace the ability of many populations adapt. Recent evidence suggests that transgenerational thermal plasticity may play a role in the response of cold-adapted organisms to long-term changes in temperature. Using RNA sequencing, we explored differential gene expression of lake trout (*Salvelinus namaycush*), a cold-adapted species, to examine the molecular processes that respond to elevated temperatures under conditions of within-generation (offspring) and transgenerational (parental) warm acclimation. We hypothesized that genes associated with metabolism, growth and thermal stress/tolerance would be differentially expressed in juvenile lake trout offspring depending on their own acclimation temperature and that of their parents. While parental warm acclimation did have a transgenerational effect on gene expression in their offspring, within-generation (offspring) warm acclimation had a larger effect on the number of differentially expressed genes. Differentially expressed genes enriched pathways for thermal stress, signaling processes, immune function, and transcription regulation and depended on the acclimation temperature of the offspring in isolation or in combination with parental warm acclimation. We provide evidence of the transgenerational response to warming at the transcriptional level in lake trout, which should be useful for future studies of transcriptomics and plasticity in this and other cold-adapted species.

Introduction

Temperate and Arctic regions are warming rapidly; surface temperatures in Canada have

increased by ~1.7°C in under a century with even greater warming happening in the Arctic

(~2.3°C; Zhang et al. 2019). As a result, high-latitude and cold-adapted organisms are

experiencing, and will continue to experience, rapid temperature increases that may threaten their

survival (Beitinger and Bennett 2000; Burkhead 2012; IPCC 2021). Range expansion to track

suitable conditions is particularly challenging for species and populations with limited dispersal

options, such as many freshwater fishes (Guzzo and Blanchfield 2017; Smith et al. 2020).

Warming rates may outpace adaptive responses (Visser 2008; Crozier and Hutchings 2014;

Comte and Olden 2017), particularly for those with low standing genetic variation (Willi et al.

2006; Wilson 2017). In the absence of standing genetic variation for coordinated adaptive

responses throughout core metabolic pathways, population responses would be limited to plastic

thermal acclimation, but the low physiological plasticity typical of many cold-adapted species

means they likely will be particularly vulnerable to rapidly changing environments (Somero

2010; Kelly et al. 2014; Kelly et al. 2018). Individuals that can withstand warming may have to

rely heavily on energetic resources that would normally support growth or reproduction, thus

having potentially negative downstream effects on the population (Somero 2010; Rosenfeld et al.

2020). A lack of adaptive genetic variation, low capacity for temperature acclimation, and rapid

environmental change are predicted to result in widespread species extirpations (Bennett et al.

2019; Morash et al. 2021).

Species may be able to buffer the negative impact of environmental warming to some

extent through transgenerational acclimation, also known as transgenerational plasticity (TGP).

As the name suggests, TGP refers to the phenotypic changes that occur over the course of

multiple generations (Jablonka et al. 1992; Bell and Hellmann 2019; Bonduriansky 2021). This

involves non-genetic inheritance of environmentally-driven changes, including parental effects,

where the experiences of one generation are reflected in the morphology, physiology or

behaviour of a subsequent generation (Hellman et al. 2020; Lee et al. 2020; Bonduriansky 2021).

TGP occurs through modulation of gene expression through non-genetic inheritance,

which includes parental effects and the transfer of epigenetic factors such as patterns of DNA

methylation and histone modification, and microRNA activity (Dai et al. 2020; Spadafora 2020;

Venney et al. 2020). TGP can potentially be beneficial when a parent's environment coincides

with what their offspring are expected to experience (Bernatchez 2016; Ashe et al. 2021). For

example, TGP could be beneficial in a population that experiences environmental warming over

multiple generations whereby the offspring are phenotypically "primed" for a warmer

environment (Donelan et al. 2020; McCaw et al. 2020; Venney et al. 2022). In the context of

climate change, this phenotypic priming could buffer the effects of rapid warming, allowing

additional time for evolutionary processes to occur (Bernatchez 2016; Ashe et al. 2021), though

the extent to which TGP can be considered adaptive is arguable (Uller et al. 2013; Sánchez-Tójar

et al. 2020). Nevertheless, TGP with temperature has been identified in a number of species

(Greenspoon and Spencer 2018; Yin et al. 2019). For example, the coral reef fish

Acanthochromis polyacanthus exhibits TGP for increased temperatures by improving its aerobic

scope in a warm environment by increasing maximum metabolic rate or decreasing resting

metabolic rate (Donelson et al. 2012; Bernal et al. 2018). Many of the studies on thermal TGP in

aquatic organisms have used eurythermal, temperate or tropical model species; by contrast, cold-

adapted aquatic vertebrates are relatively understudied but could stand to benefit from TGP

given their vulnerability to environmental warming due to climate change.

Acute and chronic temperature acclimation has obvious impacts on the metabolism of

ectotherms, at both the whole organism and molecular level (Pérez-Ruzafa et al. 2018; Petitjean

et al. 2019; Morash et al. 2021). Examining the differential gene expression associated with these

impacts on metabolism and thermal experiences in general can be informative (Oomen and

Hutchings 2017), and some studies have reported the differential gene expression underlying

previously observed transgenerational effects (Shama et al. 2016; Bernal et al. 2018; Veilleux et

al. 2018). In marine sticklebacks, for example, genes involved in metabolism, mitochondrial

protein synthesis, hemostasis and apoptosis were differentially expressed in offspring depending

on the multigenerational thermal experiences down the maternal line (Shama et al. 2016). These

findings underscore the relationship between the molecular mechanisms underlying

physiological plasticity and parental thermal experiences.

We sought to explore differential gene expression in lake trout (*Salvelinus namaycush*) to

examine the molecular processes potentially underlying transgenerational responses to elevated

temperatures. The lake trout is a long-lived, cold-adapted salmonid that is largely limited to cold,

oligotrophic lakes in formerly glaciated regions of North America (Riley et al. 2021; Wilson and

Mandrak 2021) and is under significant threat from climate change (Casselman 2008; Guzzo and

Blanchfield 2017). Being stenothermal, lake trout behaviourally thermoregulate by migrating to

deeper, cooler water when lakes thermally stratify during the summer (Martin and Olver 1980;

Guzzo and Blanchfield 2017). Low genetic variation for some lake trout populations (Perrier et

al. 2017), and little within- and among-population variation for thermal acclimation capacity

(McDermid et al. 2013; Kelly et al. 2014) may limit the adaptive potential of lake trout

populations for warming temperate and arctic habitats. A previous study investigated the capacity

for transgenerational thermal plasticity in lake trout as a means of coping with warming

associated with climate change and found that they exhibit only limited transgenerational

plasticity at the whole organism (phenotypic) level (Penney et al. 2021). Unexpectedly, cold-

acclimated offspring of warm-acclimated parents exhibited a higher resting metabolic rate than

those with one or both cold-acclimated parents (Penney et al. 2021). This study confirmed that

the parental environment can influence offspring phenotypic expression, but did not assess

potential mechanisms underlying this transgenerational response.

In the current study, we assessed gene expression in previously studied offspring from

factorial mating crosses between cold- and warm-acclimated lake trout adults (Penney et al.

2021) to assess transcriptional responses to within- and transgenerational thermal acclimation.

Acclimation of the parents (transgenerational) to one of two environmental temperatures (10°C

or 17°C) was combined with subsequent acclimation of the offspring (within-generation) to cool

(11°C) or warm (15°C) temperatures (Fig. 1). Evidence exists to support both maternal and

paternal contributions to transgenerational effects (Marshall 2015; Shama et al. 2016; Penney et

al. 2021). Our experimental design allowed us to examine the transgenerational effect of

temperature acclimation of each parent, and both parents together, in combination with offspring

(within-generation) temperature acclimation on gene expression in juvenile lake trout.

We hypothesized that genes associated with metabolism, growth and thermal

stress/tolerance would be differentially expressed in juvenile lake trout depending on both the

acclimation temperature of the offspring and the acclimation temperature of their parents. Based

on studies of within-generation and transgenerational warming on differential gene expression in

fish (Quinn et al. 2011; Veilleux et al. 2015; Akbarzadeh and Leder 2016; Shama et al. 2016), we

predicted that 1) genes involved in growth, heat shock and hypoxia responses, and metabolic

pathways would be upregulated in warm-acclimated offspring, and 2) differential expression of

these genes would vary depending on whether one or both parents were warm-acclimated,

although the size and direction of the effect is difficult to predict given that parental influence on

offspring gene expression has been shown to be sex-specific; depending on the sex of the

offspring and that of the parent (Best et al. 2018; Baustista et al. 2020).

Methods

Hatchery-raised adult lake trout from the Ontario provincial Seneca Lake broodstock

were used as parents for the experiment. This hatchery strain originated from Seneca Lake, one

of the Finger Lakes in central New York State (42°41' N, 76°54' W). The Ontario Ministry of

Natural Resources (OMNR) hatchery system has maintained this lake trout strain for over five

generations, using rotational line crossing (Kincaid 1977) to minimize inbreeding and maintain

the founding genetic diversity of the strain (OMNR Fish Culture Stocks Catalogue 2005).

The juvenile (young of year) lake trout used for this experiment were also previously

used in a separate, earlier study (Penney et al. 2021). The details of this earlier experiment that

are relevant to the current study are summarized in the next section below.

All experiments adhered to the Canadian Council on Animal Care guidelines and were

approved by the Trent University Animal Care Committee (Protocol # 24794) and the OMNRF

Aquatic Animal Care Committee (Protocol # FACC_136).

Experimental design and rearing conditions

Mature lake trout (age 8; 2.3 - 4.2 kg) were split into two groups with both sexes kept

together (n=8: 4 males and 4 females, and n=9: 4 males and 5 females), and each group was held

in a 6,000 L flow-through tank (1 x 1 x 6 m) at the OMNR White Lake Fish Culture Station

(Sharbot Lake, Ontario, Canada) with water being supplied from a nearby lake. As the source

waterbody (White Lake) began to stratify in early June, 2015, tank water temperatures were

increased by 1°C per day starting from approximately 9°C until one tank reached 10 ± 0.5°C and

the other 17 ± 0.5°C for the cold-acclimated and warm-acclimated adult lake trout groups,

respectively. These temperatures (10°C and 17°C) were chosen because 10°C represents the

thermal requirements for lake trout spawning while 17°C was intended to induce physiological

thermal stress without preventing reproduction at the higher temperature (Casselman 2008;

Penney et al. 2021). Temperatures were controlled and maintained by drawing in and mixing

water from above and below the lake's thermocline as it flowed into the holding tanks. Fish were

acclimated to these temperatures for approximately 3 months, mirroring thermal stratification in

the source waterbody. Starting in mid-September, the temperature in the warm-water treatment

was allowed to gradually cool by holding the proportional inflows from above and below the

thermocline constant as the lake's surface water cooled. Temperatures in the two tanks converged

at 10°C after fall turnover (late September) in the lake, and gradually cooled to 7.8°C by the end

of the breeding interval (mid-November).

Adult lake trout were reproductive by the end of October with the warm-acclimated fish

first spawning on October 30[th]. The cold-acclimated adults were first ready to spawn on

November 5[th] and all mating crosses were completed by November 19th. Experimental offspring

families were produced by a full factorial mating cross using two males and two females from

each of the two temperature treatments (4 x 4, 8 adults in total) resulting in a total of 16 offspring

families; four families from each of four parental treatment groups (W♀xW♂, W♀xC♂, C♀xW♂

and C♀xC♂, where W = warm-acclimated, and C = cold-acclimated; Fig. 1). Fertilized eggs were

transported to the OMNR Codrington Fish Research Facility (Codrington, Ontario, Canada)

where they were transferred into 200 L tanks receiving freshwater at ambient temperature (5-

6°C) and natural photoperiod under dim light with each family separately contained in steel mesh

boxes (9 x 9 x 7.5 cm, one family per box).

In March 2016, when hatched offspring were ready to begin exogenous feeding, we

randomly selected 14 individuals from each of the 16 families to transfer to 200 L tanks for

temperature acclimation. Seven fry from each family were acclimated to a cold temperature

(11°C) and the other 7 from each family were acclimated to a warm temperature (15°C). The

offspring's lower acclimation temperature differs from that of their parents because of reported

differences in optimum temperature for both life stages (11 vs 10°C; Edsall 2000, Casselman

2008). The upper acclimation temperature for the offspring represents the anticipated increase in

temperature associated with climate change by year 2100 (Hayhoe et al. 2010; IPCC 2021),

however, for the parents the upper acclimation temperature was two degrees higher to ensure that

temperature was high enough to elicit a physiological thermal response without compromising

reproduction. To begin acclimating the offspring, water temperature was increased at a rate of

1°C per day until target temperatures were reached (11°C and 15°C). Due to space constraints,

tanks were subdivided by adding steel inserts with perforated bottoms. Each insert was divided

into 4 sections (24 x 25 x 28 cm per section) with two paternal half-sibling families (n = 7 each;

shared male parent) housed per section. Offspring from pooled families were later individually

identified to family using microsatellite genotyping (Penney et al. 2021).

After the 3-4 week acclimation period, offspring were subjected to an acute temperature

increase of $+2°C·h^{-1}$ from their acclimation temperature until loss of equilibrium was observed.

Eight fish from the same acclimation temperature (11°C or 15°C) were simultaneously tested per

day, with each fish held in its own experimental chamber (Penney et al. 2021). Following loss of

equilibrium, each fish was quickly transferred to a bath of 0.3 g l−1 of buffered tricaine

methanesulfonate (MS-222; Aqua Life, Syndel Laboratories Ltd, BC, Canada) for euthanasia.

Whole liver was rapidly dissected from each fish, blotted on a lab wipe and preserved in

RNAlater (Invitrogen, Thermo Fisher Scientific) as per manufacturer's instructions. After 24-48

hours in a 4°C refrigerator, the RNAlater was pipetted from the tissues and the samples were

stored at -80°C until RNA extraction. Liver tissue was collected between June 28 and August 9,

2016. We chose to study gene expression in the liver because it is a metabolically active tissue

involved in a number of physiological processes that respond to warming (Quinn et al. 2011;

Akbarzadeh and Leder 2016; Dammark et al. 2018), and transcriptional responses to

multigenerational warming have previously been observed in the livers of zebrafish (Luu et al.

2021).

RNA isolation and sequencing

RNA was extracted from the preserved liver tissues using a phenol-chloroform extraction

method (Chomczynski and Sacchi 2006) after the tissues had been individually homogenized via

a FastPrep-24 BeadBeater (MP Biomedicals) with 2 mL Lysing Matrix D tubes (MP

Biomedicals) and 1 mL of Trizol reagent (Invitrogen, Thermo Fisher Scientific). RNA was

precipitated with RNA precipitation solution (Sambrook and Russel 2001) and isopropanol, and

washed with 75% ethanol. RNA samples were resuspended in nuclease-free water (Thermo

Fisher Scientific). The purity and concentration of the RNA was quantified using a NanoDrop-

8000 spectrophotometer, and RNA quality was assessed by gel electrophoresis of glyoxylated

RNA on a 1.5% agarose gel (Sambrook and Russel 2001).

Two lots of RNA samples were sent for high-throughput sequencing over a two-year

span. In 2018, liver RNA samples from 24 randomly chosen individuals were sent to The Centre

for Applied Genomics (Sick Kids Hospital, Toronto, Ontario, Canada), and in 2020 another 30

liver samples (individuals) were sent to Genome Quebec (Montreal, Quebec, Canada).

Altogether, the 54 samples included tissue from six individuals from each of the seven

experimental treatment groups, however, the control group (11°C-acclimated offspring from

cold-acclimated (CxC) parents) had a total of 12 individuals sequenced: 6 individuals in 2018

and an additional 6 in 2020, so that each treatment group was compared against the control group

sequenced in the same year to eliminate batch effects in the analysis of differential expression

(Table 1). Individuals from each family were represented in their respective treatment group

(there were 4 families per offspring-parent treatment group) and because each sequenced group

comprised of 6 individuals, each group had 2 representatives from 2 of the 4 families. Overall,

this did not lead to a bias toward any one family being overrepresented in the samples, except for

the 2020 control group which had 3 individuals from one family, but this family is not

overrepresented in any of the other groups. Subsequent genetic testing for males and females

using the salmonid *sdY* sex maker (Yano et al. 2013) confirmed that each group also contained

members of both sexes (Table 1). Both facilities assessed the RNA quality via a Bioanalyzer

(Agilent Technologies) and all samples passed quality control (RNA integrity number: ≥ 7.5).

cDNA libraries were constructed by enriching the poly(A) tails of mRNA with oligo dT-beads

using NEBNext Ultra II Directional polyA mRNA Library Prep. In 2018, barcoded libraries

were distributed among two and a half lanes and were sequenced on the Illumina HiSeq2500

instrument producing an average of 28 million reads per sample (n = 24, paired-end, 2x100 bp).

In 2020, libraries were sequenced on the Illumina HiSeq4000 with barcoded libraries distributed

among three lanes producing 36 million reads per sample (n = 30, paired-end, 2x126 bp).

Data processing and analysis

To analyze reads we used GenPipes, the main in-house framework of the Canadian

Centre for Computational Genomics (C3G), to perform major processing steps (Bourgey et al.

2019). Adapters and reads with a Phred quality score of <30 were trimmed using Trimmomatic

(v.0.36; Bolger, Lohse and Usadel 2014). The trimmed reads were aligned to the SaNama_1.0,

GCF_016432855.1 lake trout (*Salvelinus namaycush*) reference genome (Smith et al. 2022)

using STAR (Dobin et al. 2012) and read counts were obtained using HTSeq (Anders et al.

2015).

To measure the effect that offspring (within-generation) and parent (transgenerational,

one or both parents) warm acclimation had on gene expression in juvenile lake trout, we

compared the differential gene expression of each treatment group to that of the control group.

We had a total of seven treatment groups and one control group, providing us with a total of

seven comparisons which allowed us to test the effect of offspring (within-generation) warm

acclimation in isolation, parent (transgenerational) warm acclimation (either parent in isolation,

and both parents together), and the combined effect of the warm acclimation of the offspring

with one or both parents. A summary of the comparisons and the effect measured is provided in

Table 1. Differential gene expression was determined for each experimental treatment group by

comparing it to the control group within a common year to attempt to reduce or eliminate a

potential batch effect due to sequencing year in the expression data. For example, experimental

groups that were sequenced in 2020 were compared against the control group also sequenced in

2020.

Differences in gene expression levels between the treatment groups were identified using

DESeq2 (Love et al. 2014) using negative binomial GLM fitting and Wald statistics

(nbiomWaldTest), and ashr (Stephens et al. 2017) was used to shrink log2 fold changes (LFC) in

the gene expression data. Significance testing was accomplished by obtaining adjusted *p*-values

via the Benjamini-Hochberg approach for detecting false discovery rate (FDR) and differentially

expressed genes were considered as those with an adjusted p-value (FDR) of 0.05. Significantly

expressed genes with a Log2 Fold Change -0.5 and 0.5 were used in a functional enrichment

analysis with Gene Ontology (GO) in clusterProfiler (Yu et al. 2012), using the entire list of

expressed genes as background and using the Benjamini-Hochberg method to test for FDR of the

enriched terms. We plotted the top five most significantly enriched (i.e. lowest adjusted p-value)

GO terms for each category (molecular function: MF; cellular component: CC; and biological

process: BP) for each of the seven offspring comparison groups.

A principal component analysis (PCA) was performed using the rlog counts of all genes

to explore the variance across all samples and to double check for possible batch effects between

the 2018 and 2020 datasets. Seven samples appeared as outliers on the PCA plot. Of these seven

samples, two were from the control group (one from 2018 and one from 2020), and a sample

each from the following experimental groups: warm-acclimated offspring (C♀xC♂), cold-

acclimated offspring (C♀xW♂), cold-acclimated (W♀xW♂), warm-acclimated offspring

(W♀xC♂), and warm-acclimated offspring (W♀xW♂). To account for the batch effect, we simply

matched the samples to the corresponding sequencing group as well as the experimental groups

into the differential expression design formula allowing us to account for both batch effect and

treatment group to identify differentially expressed genes. We applied the batch correction to the

samples using the twelve control group individuals (i.e. n=6 from 2018 and n=6 from 2020

control groups combined) as a shared baseline. One other sample, from a warm-acclimated

offspring (W♀xW♂) experimental treatment group, had a substantial number of reads align to

Homo sapiens genome indicating potential contamination had occurred during RNA isolation.

This sample was excluded from the final dataset.

Results

The total number of paired-end reads for each of the 53 lake trout liver samples ranged

from 26,150,786 to 52,743,330. Between 88.0% and 93.2% of these reads successfully aligned

with the lake trout reference genome (Smith et al. 2022).

The PCA of offspring gene expression data did not show clear segregation among sample

groups or a significant batch effect between the two sets of sequence data (Fig. 2). Seven

samples appeared as outliers on the PCA plot: of these seven samples, two were from the control

group (one from 2018 and one from 2020), and single outliers were detected from the warm-

acclimated offspring (C♀xC♂), cold-acclimated offspring (C♀xW♂), cold-acclimated (W♀xW♂),

warm-acclimated offspring (W♀xC♂), and warm-acclimated offspring (W♀xW♂) experimental

groups. Before the batch correction was applied, principal components (PC) 1 and 2 explained

26% and 9% of the variance across all groups (Fig. 2A). The variance explained by PC1 and PC2

changed to 11% and 11% respectively after the batch correction was applied (Fig. 2B),

suggesting that a batch effect was originally present in our dataset. Other principal components

explained 9% or less of the variation (Fig. 2C). While the correction method we used was

effective at mitigating the batch effect in our dataset, we are aware that some of the batch effect

may still be present to a lesser extent and may influence the interpretation of the results.

Within-generation acclimation: offspring warm acclimation

Offspring within-generation warm acclimation resulted in 821 DEGs with a LFC of ≤ -

0.5 to ≥ 0.5 (Fig. 2 and 3). Of these genes, 318 were uniquely differentially expressed under the

isolated effect of within-generation warm acclimation and an additional 106 DEGs were

differentially expressed whenever offspring were warm-acclimated (Fig. 3).

The isolated effect of offspring within-generation warm acclimation impacted functions

in each GO category (molecular function: MF; cellular component: CC; and biological process:

BP) related to oxygen binding and gas transport (GO:0005344, 0020037, 0005833, 0031838,

0072562, 0015669; Fig. 4). Also affected were biological processes and molecular functions

related to immune function and cell signaling (GO:0048018, 0030546), such as cellular

responses to biotic stimulus/lipopolysaccharide/molecules of bacterial origin (GO:0071222,

0071219, 0071216), and biological processes related to dendrite membranes (GO:0032590) and

toll-like receptor signalling (GO:0002224; Fig. 4). Offspring warm acclimation also affected

aggresome activity (GO:0016235) and cytokine activity (GO:0005125; Fig. 4).

Transgenerational acclimation: parental warm acclimation

An effect of transgenerational (parental) warm acclimation on offspring differential gene

expression was evident, however, parental warm acclimation resulted in fewer DEGs compared

to the isolated effect of offspring (within-generation) warm acclimation (Fig. 2 and 3). Paternal

warm acclimation had the smallest effect, resulting in just 4 DEGs in their offspring (third row

down, Fig. 3). The combined effect of warm acclimation of both parents resulted in fewer

differentially expressed genes in the offspring compared to when offspring were under the

influence of maternal warm acclimation alone (both parents: 13 DEGs vs. maternal: 63 DEGs;

sixth and fourth row down, Fig. 3), suggesting that paternal warming can dampen some of the

effect of maternal warm acclimation on offspring gene expression. Of these DEGS, 49 were

uniquely expressed (i.e. did not overlap with other transgenerational warm acclimation treatment

groups) when mothers were warm-acclimated, 3 DEGs were uniquely expressed with paternal

warm acclimation and only 8 DEGs were uniquely expressed when both parents were warm-

acclimated (Fig. 3).

Maternal warm acclimation impacted the regulation of cellular components and

biological processes related to muscle function, specifically cardiac muscle repolarization, in

their offspring (GO:0030016, 0043292, 0031674, 0031430, 0008016, 0060307, 0099625,

0099622, 0099623; Fig. 5). Other enriched GO terms included cellular components associated

with the apical plasma membrane (GO:0016324), and molecular functions associated with

transmembrane transporter and ion channel activity (GO:0015106, 0015276, 0022834, 0015103,

0008509; Fig. 5). With so few DEGs, no GO output was obtained for the transgenerational effect

of paternal warm acclimation or warm acclimation of both parents.

Combined generational acclimation: offspring and parental warm acclimation

The combined effect of offspring warm acclimation with that of either or both parents

often resulted in a higher number of DEGs than seen with offspring within- or transgenerational

warm acclimation in isolation, except for the combined effect of offspring and paternal warm

acclimation which resulted in fewer DEGs than seen with the isolated effect of offspring warm

acclimation (offspring and maternal: 1,060; offspring and paternal: 492; offspring and both

parents: 1,149; Fig. 2 and 3). Unsurprisingly, the greatest number of uniquely expressed DEGs

occurred with the combined effect of offspring, maternal and paternal warm acclimation with

581 DEGs (Fig. 3). This was closely followed by the combined effect of offspring and maternal

warm acclimation with 560 DEGs, and the combined effect of the offspring and paternal warm

acclimation resulted in 149 DEGs (Fig. 3).

When both the offspring and the mothers were warm-acclimated, molecular functions

associated with an increased metabolism (e.g. acetyl-CoA C-acyltransferase; GO:0003988) and

protein folding chaperone activity (GO:0044183) were enriched (Fig. 6A). Also enriched were

molecular functions and biological processes related to DNA synthesis/replication (GO:0004748,

0061731, 0006260, 0006270), and cellular components involving the nuclear membrane activity

(GO:0034992, 0034993, 0106083, 0106094 ; Fig. 6A). Biological processes associated with

responses to monoamine catecholamine signaling (e.g. dopamine) were also affected by the

combined effect of offspring and maternal warm acclimation (GO:0007191, 0071868, 0071870;

Fig. 6A).

When both the offspring and fathers were warm-acclimated (Fig 6B), molecular functions

involving protein folding chaperone activity (GO:0044183) and phosphorylation (e.g. JUNKK

kinases activity; GO:00004706) were enriched (Fig. 6B). Other molecular functions included

membrane receptor and transporter activity for cell uptake of lipoproteins (GO:0019534,

0005041, 0030228; Fig. 6B). Cellular components included increased cellular respiration (e.g.

pyruvate dehydrogenase complex activity; GO:0005967, 0045254), DNA transcription (e.g.

PRC1; GO;0035102) and components related to the development of microvilli (GO:0005902)

and stress fibers (actin filaments; GO:0001725; Fig. 6B). Some enriched biological processes

were related to molecular function, such as protein refolding (GO:0042026). Lake trout offspring

showed an increased general response to heat (GO:0009408), and indicators of growth, such as

cartilage and genital growth (GO: 0060536, 0035261, 0060596; Fig. 6B).

When the offspring and both parents were warm-acclimated, most enriched GO terms

across all categories were related to DNA replication and repair (GO:0004748, 0061731,

0016728; 0006260, 0006261, 0044786, 0000727; Fig. 6C). Other molecular functions were

related to ferrous binding (GO:0008198) and protein chaperone activity (GO:0044183), while

other cellular components included functions involved in protein complexes (GO:0000796,

0031261, 0000811, 0000228), specifically with regards to post-translational modification of

protein (e.g. oligosaccharyltransferase complex; GO:0008250; Fig. 6C). Other biological

processes were related to apoptotic functions (GO:2000352; Fig. 6C).

Discussion

Our hypothesis that genes associated with metabolism, growth and thermal

stress/tolerance would be differentially expressed in juvenile lake trout depending on the

acclimation temperatures of the offspring as well as their parents was supported by the

transcriptome data. Genes involved in growth, heat stress responses and metabolic pathways

were differentially expressed with within-generation and transgenerational warm acclimation,

and with the combination of warm acclimation of both generations. The amount of differential

expression (total DEGs) and the type of enriched functions depended on whether it was the

offspring or parent(s) that had been warm-acclimated. Within-generation (offspring) warm

acclimation had a larger effect on the overall number of DEGs compared to transgenerational

contributions from warm-acclimated parents (mother, father or both). As might be expected,

however, the combination of offspring, maternal and paternal warm acclimation had the greatest

effect on the total number of DEGs.

We considered up- and downregulated differentially expressed genes together in order to

assess which pathways/processes are involved in the within- and transgenerational responses to

warm acclimation. Although, the both up- and downregulated genes were analyzed together, it

should be recognized that warm acclimation often results in disproportionately more upregulated

genes compared to downregulated genes (Healy et al. 2017).

General effect of warming on differential gene expression

In general, within- and transgenerational warm acclimation resulted in differential

expression of genes associated with heat tolerance and a heat stress response. In particular, our

enrichment analysis highlighted increased aggresome, apoptosis and protein chaperone activity

indicating heat-related injury to tissues. Some of these pathways included genes that encode heat

shock proteins (HSPs), which is not surprising given that induction of HSPs are evolutionarily

conserved across numerous taxa (Krens et al. 2006; Kostenko et al. 2011) and that increased

expression of these proteins have been observed previously in other fish taxa following heat

stress (Keller et al. 2008; Kelly et al. 2018; Liu et al. 2021). Warm-acclimated lake trout

offspring also showed an increase in immune system activity. The lake trout offspring in our

study demonstrated differential expression of genes associated with a toll-like receptor signalling

pathway, and similar results have been observed in other fish species. For example, upregulation

of the toll-like receptor 3 gene (*tlr3*) occurred in the liver of the flounder, *Paralichthys olivaceus*,

following heat shock (Lee et al. 2023). An increase in immune-related functions is common in

heat stressed fish and often appear to coincide with indicators of temperature-induced tissue

damage (Huang et al. 2018; Beemelmanns et al. 2021). Taken together, these pathways indicate

increasing temperatures elicited a stress response in lake trout, which we would expect given that

each offspring was subjected to an acute temperature increase ($+2°C/h$) and these pathways were

likely upregulated to respond to cellular/protein damage occurring at their upper critical thermal

limit (Penney et al. 2021).

With warm acclimation, there was a general enrichment of pathways associated with an

increased metabolic demand, including O_2 binding and gas transport. Upregulation of processes

related to O_2 transport, such as heme binding, have previously been observed in the liver of fish

with a history of heat stress (Duan et al. 2024) and could mean that the lake trout offspring were

attempting to increase their O_2 carrying capacity to meet increased metabolic demands (Lu et al.

2020). We also observed differential expression of metabolic enzymes such as acetyl-CoA C-

acyltransferase activity and pyruvate dehydrogenase. Both of these enzymes are used in the

production of acetyl CoA: a key coenzyme involved in several metabolic functions, including the

citric acid cycle (Shi and Tu 2015). Our transcriptomic findings here are indicative of an increase

in O_2 demand which aligns with the findings of Penney et al. 2021 in that offspring metabolic

rates, measured as rates of O_2 consumption (MO_2), were elevated at the time of tissue sampling.

We found some indication that warm acclimation and upper thermal stress affected the

growth of the lake trout offspring. We expected to see differential expression of some growth-

related functions given that the warm-acclimated offspring were heavier than the cold-acclimated

offspring at the time of sampling (Penney et al. 2021). These functions included oxygen

transport, cartilage growth and synthesis of cellular membrane and cytoskeletal components.

Other studies have observed upregulation of similar pathways involved in growth in fish

(Danzmann et al. 2016; Lu et al. 2020; Vergauwen et al. 2010). For example, we saw an

enrichment of acetyl-CoA C-acyltransferase activity which could indicate a higher demand for

acetyl-CoA production for growth, just as a transcriptome analysis of the liver of *Salmo salar*

revealed an increased demand for acetyl-CoA in faster growing groups of fish (Caballero-Solares

et al. 2018).

It is curious that enrichment of more growth-related processes was not discovered in the

heavier, warm-acclimated lake trout offspring in our study. For this reason, we cannot be entirely

confident that the enriched pathways we observed resulted primarily from growth of the

offspring. We collected liver samples at each fish's critical thermal maximum, so it is possible

that fish were responding to the stress of the acute temperature increase and may have

suppressed growth during the acute temperature challenge in favour of a heat stress response.

Suppression of growth at high temperatures is not unusual. For example, increased expression of

the genes *GH* and *IGF-1* (indicators of growth) occurred in *Labeo rohita* at slightly warmer

acclimation temperatures but was suppressed at higher acclimation temperatures (Shahjahan et

al. 2021). Alternatively, there is evidence to suggest that growth hormone signaling pathways are

more sensitive to elevated temperature in the muscle compared to other tissues (Hevrøy et al.

2015). For example, Houde et al. (2019) did not see differential expression of insulin-like growth

factor binding proteins *IGFBP-1* and *IGFBP-10-v1* in the gills of *Onchorhynchus tshawytscha*.

Thus, it is possible that we may have seen a stronger transcriptional response for growth-related

pathways in the muscle rather than in the liver of the lake trout used in our study, though

confirmation of this would require further experimentation.

Transcriptomic responses to transgenerational warm acclimation

We found that an offspring's thermal acclimation often had a greater influence on the

number of DEGs than did the thermal experience of its parents, suggesting that transgenerational

transcriptomic responses to warming are weaker than those that occur within a generation. This

result agrees with a previous study that saw that the effect of transgenerational warm acclimation

on offspring metabolic rate (rate of O_2 consumption) was limited (Penney et al. 2021). In

contrast to our findings, some previously published evidence suggests that transgenerational

warming has a stronger effect on offspring differential gene expression compared to

developmental warming (Veilleux et al. 2015; Shama et al. 2016). It has also been suggested that

transgenerational effects are strongest in early-juvenile life stages (Yin et al. 2019) and it would

be interesting to see if this is reflected at the transcriptomics level by looking at gene expression

at different life stages following within- and transgenerational warm acclimation in lake trout.

Related to this, it may worth noting that prior to warm acclimation all offspring in our

experiment were reared in the same environment and temperature regime to minimize

differential developmental effects. How developmental and transgenerational warming interact in

these fish is not fully understood, but the influence of thermal acclimation on differential gene

expression during early, developmental life stages compared to later life stages may be a fruitful

area for study. Lastly, the combination of offspring, maternal and paternal warm acclimation had

the greatest effect on the overall number genes that were differentially expressed. This adds to

the growing literature suggesting that the strength of transgenerational plasticity is greatest when

offspring and parental thermal experiences coincided (Shama et al. 2014; Donelson et al. 2017;

Yin et al. 2019).

Paternal transgenerational warm acclimation appeared to dampen the effect of maternal

warm acclimation on offspring gene expression in that fewer genes were differentially expressed

in the offspring when both parents were warm-acclimated compared to when offspring were

under the influence of maternal warm acclimation only. Earlier discussions of transgenerational

plasticity speculated that fathers could influence maternal investment in their offspring through

non-genetic means, including imprinting and components included in glandular secretions (Uller

2008; Bonduriansky and Day 2009). Investigations of paternal effects are less common than that

of maternal effects and there have been calls for more studies on paternally mediated non-genetic

inheritance on offspring phenotypes (Rutkowska et al. 2020). Recent studies have shown that a

stronger paternal contribution is possible. In brook trout (*S. fontinalis*), paternal transgenerational

warming can contribute more to offspring thermal responses and survival compared to the

maternal contribution (Penney et al. 2022; Houle et al. 2023). In contrast, we did not previously

see a stronger paternal effect in the thermal response of the lake trout offspring used for this

experiment (Penney et al. 2021). Whether this was due to suppression of offspring gene

expression is not entirely clear but further investigation into this effect could be insightful,

potentially revealing a mechanism by which fathers can control their offspring's developmental

rate or resource use in competition or in concert with maternal effects.

533   An effect of parental (transgenerational) warm acclimation on the offspring's heat shock

response was evident, providing support for our hypothesis. The combination of offspring and

maternal warm acclimation led to the differential expression of genes associated with cellular

responses to catecholamines (e.g. dopamine). Modification of cellular sensitivity to

catecholamines can influence responses to warmer temperatures. For example, sensitivity to

dopamine can regulate thermal preference behaviour (Ueno et al. 2012), and blocking of the β-

adrenergic pathway can lead to a decreased cellular heat shock response in rainbow trout

(Templeman et al. 2014). Additionally, biological processes related to protein refolding and a

general response to heat were differentially expressed under conditions of offspring and paternal

warm acclimation. Our results suggest that perhaps thermal sensitivity is regulated when

offspring and parental warm acclimation coincide which is supported by the idea that

transgenerational plasticity can be adaptive when environmental changes trend in the same

direction across generations (Shama et al. 2014; Donelson et al. 2017; Yin et al. 2019).

546   Our prediction that transgenerational warm acclimation would upregulate metabolic

processes in lake trout offspring was partially supported, and is indicative of a higher metabolic

rate in warm-acclimated offspring with warm-acclimated parents. Maternal transgenerational

warm acclimation in isolation led to the enrichment of processes related to cellular

transmembrane ion transport, including the transfer of bicarbonate ions. The increase in higher

metabolic rates of ectotherms with warming can result in metabolic (lactic) acidosis and the

release of bicarbonate ions into the blood counteracts the shift in pH (Stewart et al. 2019). The

combined effect of warm-acclimated mothers and offspring led to differential expression of

genes associated with acetyl-CoA C-acyltransferase activity, and the combined effect of warm-

acclimation of the fathers and offspring led to differential expression of genes associated with the

pyruvate dehydrogenase complex. These two enzymes are involved in regulating metabolism

through their involvement in the production of acetyl CoA (Shi and Tu 2015), however, previous

work showed that neither resting or peak metabolic rates differed between warm-acclimated

offspring from warm- or cold-acclimated parents (Penney et al. 2021). It is worth noting that

gene expression does not necessarily confer a protein to function given that post-transcriptional

and post-translational activity could also tag a protein for degradation; the enrichment analysis

may suggest an upregulation of metabolic processes, but these proteins may have been rendered

inactive, or perhaps were not expressed highly enough to be detected at the level of the

phenotype in the lake trout offspring.

We found some support for a transgenerational effect of warm acclimation on growth in

lake trout at the transcriptome level. Molecular functions including DNA synthesis (e.g.

ribonucleoside-diphosphate reductase activity), biological processes related to DNA replication,

and cellular components involved in stabilizing the nuclear membrane were influenced by the

combined effect of maternal and offspring warm acclimation. An upregulation in cellular

structural components has previously been observed with warm acclimation and growth in fish

(Vergauwen et al. 2010; Lu et al. 2020). Protein folding chaperone activity also appeared in the

molecular function GO category for each of the combined warm acclimation groups: warm-

acclimated offspring with warm-acclimated mothers, fathers, or both parents and may be

upregulated for the formation of new proteins. An alternative explanation could be that these

functions were involved in other processes related to cell repair/replacement as cells and proteins

sustain heat damage, especially considering that apoptosis appeared in the enrichment analysis

when offspring and both parents were warm-acclimated. Finally, biological processes related to

genitalia and cartilage differentiation were differentially expressed when fathers and offspring

were warm-acclimated and may be indicative of growth of reproductive organs as the offspring

develop towards sexual maturity. Transgenerational warming can result in larger offspring

(Salinas et al. 2012), however, our transcriptome findings are not consistent with the findings of

Penney et al. (2021) where transgenerational warm acclimation had no effect on the mass of the

offspring. Nevertheless, our transcriptome results suggest that there is some transgenerational

plasticity in growth-related functions when both parents and offspring experience warming.

Strangely, warm-acclimated mothers seemed to have an influence over processes related

to muscle function in their offspring. These included cellular components and biological

processes associated with contractile fibers and cardiac muscle cell membrane function. These

same functions were not apparent with the combined effect of maternal and offspring warm
acclimation. It is curious that GO terms specifying muscle functions appeared in our results
given that the focal tissue was liver. However, a transgenerational effect of thermal acclimation
on muscle function has been observed in liver samples fish. For example, transgenerational
warm acclimation resulted in differentially methylated regions of genes associated with cardiac
muscle function, heart rate and cardiac myofibril assembly in liver tissue of *Acanthochromis*
polyacanthus (Ryu et al. 2020). Future studies could test the tissue-specific transcriptomic
responses to transgenerational warm acclimation in lake trout or other salmonids to better
understand the functions under the influence of transgenerational plasticity.

Venney et al. (2022) used a similar breeding design as this study to assess temperature-
related DNA methylation in adult brook trout and their offspring. In contrast to our findings,
Venney et al. (2022) observed a stronger transgenerational effect from adults held at higher
temperatures during gonad development compared with methylation from offspring rearing
temperatures. Despite the apparent contrasts between our results and those of Venney et al.
(2022), the findings from the two studies are in fact complementary rather than contradictory.
Some studies on salmonids have shown that changes in global methylation do not appear to
influence differential gene expression (Christensen et al. 2021; Leitwein et al. 2022). DNA
methylation can influence gene expression in an individual and subsequent generations, but gene
expression is also a direct response to physiological, metabolic, and bioenergetic requirements
including acclimation. The greater methylation observed from reproductive adults by Venney et
al. (2022) may reflect the lower temperature tolerances of reproductive adults compared with
other free-swimming life stages (Dahlke et al. 2020). The lesser extent of DNA methylation
observed in juvenile brook trout by Venney et al. may also reflect the ability of subadult brook
trout to use warmer temperatures compared to adults (Smith and Ridgway 2019), as well as the
greater metabolic scope of juvenile fish compared with reproductive adults (Dahlke et al. 2020).
By contrast, rather than studying methylation, our study examined levels of gene expression and
showed a greater effect of within-generation acclimation. Our current results parallel those from
a previous study looking at whole-organism respirometry in lake trout (Penney at al. 2021).

Conclusions

Our study has shown that lake trout exhibit both transgenerational (both maternal and

paternal) and within-generation thermal acclimation, with both levels of acclimation influencing

the strength of differential expression of genes associated with thermal stress/tolerance,

metabolism and growth. Despite this, the limited transgenerational effect on gene expression

indicates that adaptive transgenerational acclimation is unlikely to contribute sufficiently

substantive benefits to enable lake trout to cope with rapidly changing environmental conditions

related to global warming (Penney et al. 2021). Instead, population-level responses to

temperature-related stress will likely be restricted to within-generation acclimation (Kelly et al.

2014) and longer-term adaptive responses. Beyond acclimation, persistence of cold-water species

and populations may be reliant on existing genetic resources to cope with chronic warming

(Comte and Olden 2017; Crozier and Hutchings 2014; Guzzo and Blanchfield 2017), which is

particularly daunting for long-lived species (Willi et al. 2006). It therefore seems likely that lake

trout and similarly vulnerable cold-water species may require increased management and

conservation efforts to ensure their future in a rapidly warming world.

Acknowledgements

The Ontario Ministry of Natural Resources (OMNR) Fish Culture Section provided adult

lake trout, rearing space and logistical support at the White Lake Fish Culture Station. Bill Sloan,

Scott Ferguson, Anne McCarthy (OMNR) and John Dewart provided assistance with breeding

adult fish; Bill Sloan and Scott Ferguson also provided invaluable logistical support for all

aspects of fish husbandry from care of the juvenile trout to assisting with experiments at the

OMNR Codrington Fisheries Research Facility. Caleigh Smith (OMNRF) genotyped adult and

juvenile fish for parentage assignment, family identification of juveniles and sex marker typing.

Paul Craig and Heather Ikert (University of Waterloo) kindly provided advice and guidance on

the enrichment analysis. Paul Craig, Graham Scott, Anne Dalziel and Aaron Shafer provided

feedback on early drafts of the manuscript. The RNA sequencing was conducted by The Centre

for Applied Genomics (Sick Kids Hospital, Toronto, Ontario, Canada) and Genome Quebec

(Montreal, Quebec, Canada). The bioinformatic analyses was performed by Canadian Center for

Computational Genomics (Montreal, Quebec, Canada). The Canadian Center for Computational

Genomics (C3G) is a Genomics Technology Platform (GTP) supported by the Canadian

Government through Genome Canada.

Research funding was provided by the Canada-Ontario Great Lakes Agreement (COA) to

CCW, by the Ontario Federation of Anglers and Hunters (OFAH) to CMP, and a Natural

Sciences and Engineering Research Council (NSERC) Discovery Grant to GB.

Data archiving statement: Data for this study are available at: *to be completed after manuscript*

is accepted for publication.

Literature Cited

Akbarzadeh, A., Leder, E.H. (2016). Acclimation of killifish to thermal extremes of hot spring: transcription of gonadal and liver heat shock genes. *Comparative Biochemistry and Physiology Part A: Molecular & Integrative Physiology*, 191, 89-97.

Anders, S., Pyl, P.T., Huber, W. (2015). HTSeq—a Python framework to work with high-throughput sequencing data. *Bioinformatics*, 31, 166-169.

Ashe, A., Colot, V., Oldroyd, B.P. (2021). How does epigenetics influence the course of evolution? *Philosophical Transactions of the Royal Society B: Biological Sciences*, 376(1826), 20200111.

Beemelmanns, A., Zanuzzo, F.S., Sandrelli, R.M., Rise, M.L., Gamperl, A.K. (2021). The Atlantic salmon's stress-and immune-related transcriptional responses to moderate hypoxia, an incremental temperature increase, and these challenges combined. *G3*, 11, jkab102.

Beitinger, T.L., Bennett, W.A. (2000). Quantification of the role of acclimation temperature in temperature tolerance of fishes. *Environmental Biology of Fishes*, 58, 277-288.

Bell, A.M., Hellmann, J. (2019). An integrative framework for understanding the mechanisms and multigenerational consequences of transgenerational plasticity. *Annual Review of Ecology, Evolution and Systemics*, 50, 97–118.

Bennett, S., Duarte, C.M., Marbà, N., Wernberg, T. (2019). Integrating within-species variation in thermal physiology into climate change ecology. *Philosophical Transactions of the Royal Society B: Biological Sciences*, 374, 20180550.

Bernal, M.A., Donelson, J.M., Veilleux, H.D., Ryu, T., Munday, P.L., Ravasi, T. (2018). Phenotypic and molecular consequences of stepwise temperature increase across generations in a coral reef fish. *Molecular Ecology*, 27, 4516-4528.

Bernatchez, L. (2016). On the maintenance of genetic variation and adaptation to environmental
change: considerations from population genomics in fishes. *Journal of Fish Biology*, 89, 2519-
2556.

Best, C., Ikert, H., Kostyniuk, D. J., Craig, P. M., Navarro-Martin, L., Marandel, L., Mennigen,
J. A. (2018). Epigenetics in teleost fish: from molecular mechanisms to physiological
phenotypes. *Comparative Biochemistry and Physiology Part B: Biochemistry and Molecular*
Biology, 224, 210-244.

Bolger, A.M., Lohse, M., Usadel, B. (2014). Trimmomatic: a flexible trimmer for Illumina
sequence data. *Bioinformatics*, 30, 2114-2120.

Bonduriansky R. (2021). Plasticity across generations. In: D.W. Pfennig (Ed.). Phenotypic
Plasticity & Evolution: Causes, Consequences, Controversies (pp. 327-348). Boca Raton: CRC
Press/Taylor & Francis Group.

Bourgey, M., Dali, R., Eveleigh, R., Chen, K.C., Letourneau, L., Fillon, J., Michaud, M., Caron,
M., Sandoval, J., Lefebvre, F., Leveque, G., Mercier, E., Bujold, D., Marquis, P., Van, P.T.,
Anderson de Lima Morais, D., Tremblay, J., Shao, X., Henrion, E., Gonzalez, E., Quirion, P.,
Caron, B., Bourque, G. (2019). GenPipes: an open-source framework for distributed and scalable
genomic analyses. *GigaScience*, 8, giz037.

Burkhead, N.M. (2012). Extinction rates in North American freshwater fishes, 1900-
2010. *BioScience*, 62: 798-808.

Caballero-Solares, A., Xue, X., Parrish, C. C., Foroutani, M. B., Taylor, R. G., Rise, M. L.
(2018). Changes in the liver transcriptome of farmed Atlantic salmon (*Salmo salar*) fed
experimental diets based on terrestrial alternatives to fish meal and fish oil. *BMC Genomics*, 19,
1-26.

Casselman, J.M . (2008). Effects of climate and climate change on lake trout populations and
fisheries. In: K.H. Mills, M. Dyck, & L.A. Harwood (Eds.). Proceedings of the Second North
American Lake Trout Symposium (pp. 103-113). Yellowknife, NT, Canada: Canadian Technical
Report of Fisheries and Aquatic Sciences.
720
721 Chomczynski, P., Sacchi, N. (2006). The single-step method of RNA isolation by acid
722 guanidinium thiocyanate–phenol–chloroform extraction: twenty-something years on. *Nature*
723 *Protocols*, 1, 581-585.
724
725 Christensen, K.A., Le Luyer, J., Chan, M.T., Rondeau, E.B., Koop, B.F., Bernatchez, L., Devlin,
726 R.H. (2021). Assessing the effects of genotype-by-environment interaction on epigenetic,
727 transcriptomic, and phenotypic response in a Pacific salmon. *G3 Genes|Genomes|Genetics*, 11,
728 jkab021.
729
730 Comte, L., Olden, J.D. (2017). Climatic vulnerability of the world's freshwater and marine
731 fishes. *Nature Climate Change*, 7, 718–722.
732
733 Crozier, L.G. J.A. Hutchings. (2014). Plastic and evolutionary responses to climate change in
734 fish. *Evolutionary Applications*, 7, 68-87.
735
736 Curley, J.P., Mashoodh, R., Champagne, F.A. (2011). Epigenetics and the origins of paternal
737 effects. *Hormones and behavior*, 59, 306-314.
738
739 Dai, Z., Ramesh, V., Locasale, J.W. (2020). The evolving metabolic landscape of chromatin
740 biology and epigenetics. *Nature Reviews Genetics*, 21, 737-753.
741
742 Dahlke, F.T., Wohlrab, S., Butzin, M., Pörtner, H.O. (2020). Thermal bottlenecks in the life
743 cycle define climate vulnerability of fish. *Science*, *369*, 65-70.
744

Dammark, K. B., Ferchaud, A.L., Hansen, M.M., Sørensen, J.G. (2018). Heat tolerance and gene
expression responses to heat stress in threespine sticklebacks from ecologically divergent
environments. *Journal of Thermal Biology*, 75, 88-96.

Danzmann, R.G., Kocmarek, A.L., Norman, J.D., Rexroad, C.E., Palti, Y. (2016). Transcriptome
profiling in fast versus slow-growing rainbow trout across seasonal gradients. *BMC Genomics*,
17, 1-18.

Dobin, A., Davis, C.A., Schlesinger, F., Drenkow, J., Zaleski, C., Jha, S., Batut, P., Chaisson, M.
Gingeras, T.R. (2013). STAR: ultrafast universal RNA-seq aligner. *Bioinformatics*, 29, 15-21.

Donelan, S.C, Hellman, J.K., Bell, A.M., Luttbeg, B., Orrock, J.L., Sheriff, M.J., Sih, A. (2020).
Transgenerational plasticity in human-altered environments. *Trends in Ecology & Evolution*, 35,
115-124.

Donelson, J.M., Munday, P.L., McCormick, M.I., Pritcher, C.R. (2012). Rapid transgenerational
acclimation of a tropical reef fish to climate change. *Nature Climate Change*, 2, 30–32.

Donelson, J.M., Sunday, J.M., Figueira, W.F., Gaitán-Espitia, J.D., Hobday, A.J., Johnson, C.R.,
Leis, J.M., Ling, S.D., Marshall, D., Pandolfi, J.M., Pecl, G., Rodgers, G.G., Booth, D.J.,
Munday, P.L. (2019). Understanding interactions between plasticity, adaptation and range shifts
in response to marine environmental change. *Philosophical Transactions of the Royal Society B:*
Biological Sciences, 374, 20180186.

Duan, C., Tian, C., Guan, Y., Xu, H., Yang, L., Chen, Y., Liu, Y., Shen, Y., Zhang, Y., Cao, S.,
Huang, Y., Li, G. (2024). Long-term thermal stress induces hepatic injury and alters the
thermotolerance response in Hong Kong catfish (*Clarias fuscus*). *Aquaculture*, 590, 741041.

Edsall, T.A., Cleland, J. (2000). Optimum temperature for growth and preferred temperatures of
age-0 lake trout. *North American Journal of Fisheries Management*, 20, 804-809.

Greenspoon, P.B., Spencer, H.G. (2018). The evolution of epigenetically mediated adaptive
transgenerational plasticity in a subdivided population. *Evolution*, 72, 2773-2780.

Guzzo, M.M., Blanchfield, P.J. (2017). Climate change alters the quantity and phenology of
habitat for lake trout (*Salvelinus namaycush*) in small Boreal Shield lakes. *Canadian Journal of*
Fisheries and Aquatic Sciences, 74, 871-884.

Hayhoe, K., VanDorn, J., Croley II, T., Schlegal, N., Nwuebbles, D. (2010). Regional climate
change projections for Chicago and the US Great Lakes. *Journal of Great Lakes Research*, 36
(Suppl 2), 7-21.

Healy, T.M., Bryant, H.J., Schulte, P.M. (2017). Mitochondrial genotype and phenotypic
plasticity of gene expression in response to cold acclimation in killifish. *Molecular Ecology*, 26,
814-830.

Hellmann, J.K., Carlson, E.R., Bell, A.M. (2020). Sex-specific plasticity across generations II:
Grandpaternal effects are lineage specific and sex specific. *Journal of Animal Ecology*, 89, 2800-
2812.

Hevrøy, E.M., Tipsmark, C.K., Remø, S.C., Hansen, T., Fukuda, M., Torgersen, T., Vikeså, V.,
Olsvik, P.A., Waagbø, R., Shimizu, M. (2015). Role of the GH-IGF-1 system in Atlantic salmon
and rainbow trout postsmolts at elevated water temperature. *Comparative Biochemistry and*
Physiology Part A: Molecular & Integrative Physiology, 188, 127-138.

Houde, A.L.S., Akbarzadeh, A., Günther, O.P., Li, S., Patterson, D.A., Farrell, A.P., Hinch, S.G.,
Miller, K. M. (2019). Salmonid gene expression biomarkers indicative of physiological
responses to changes in salinity and temperature, but not dissolved oxygen. *Journal of*
Experimental Biology, 222, jeb198036.

Houle, C., Gossieaux, P., Bernatchez, L., Audet, C., Garant, D. (2023). Transgenerational effects
on body size and survival in brook charr (*Salvelinus fontinalis*). *Evolutionary Applications*, 16,
1061-1070.

Huang, J., Li, Y., Liu, Z., Kang, Y., Wang, J. (2018). Transcriptomic responses to heat stress in
rainbow trout *Oncorhynchus mykiss* head kidney. *Fish & Shellfish Immunology*, 82, 32-40.

IPCC (2021). Summary for Policymakers. In: V. Masson-Delmotte, P. Zhai, A. Pirani, S.L.
Connors, C. Péan, S. Berger, N. Caud, Y. Chen, L. Goldfarb, M.I. Gomis, M. Huang, K. Leitzell,
E. Lonnoy, J.B.R. Matthews, T.K. Maycock, T. Waterfield, O. Yelekçi, R. Yu, & B. Zhou
(Eds.), Climate Change 2021: The Physical Science Basis. Contribution of Working Group I to
the Sixth Assessment Report of the Intergovernmental Panel on Climate Change (pp. 3−32).
Cambridge, United Kingdom and New York, NY, USA:Cambridge University Press.

Jablonka, E., Lachmann, M., Lamb, M.J. (1992). Evidence, mechanisms and models for the
inheritance of acquired characters. *Journal of Theoretical Biology*, 158, 245-268.

Keller, J.M., Escara-Wilke, J.F., Keller, E.T. (2008). Heat stress-induced heat shock protein 70
expression is dependent on ERK activation in zebrafish (*Danio rerio*) cells. *Comparative*
Biochemistry and Physiology Part A: Molecular & Integrative Physiology, 150, 307-314.

Kelly, N.I., Burness, G., McDermid, J.L., Wilson, C.C. (2014). Ice age fish in a warming world:
minimal variation in thermal acclimation capacity among lake trout (*Salvelinus namaycush*)
populations. *Conservation Physiology*, 2, doi:10.1093/conphys/cou025.

Kelly, N.I., Wilson, C.C., Currie, S., Burness, G. (2018). Acclimation capacity of the cardiac
HSP70 and HSP90 response to thermal stress in lake trout (*Salvelinus namaycush*), a
stenothermal ice-age relict. *Comparative Biochemistry and Physiology Part B: Biochemistry and*
Molecular Biology, 224, 53-60.

Kincaid, H.L. (1977). Rotational line crossing: an approach to the reduction of inbreeding

accumulation in trout brood stocks. *The Progressive Fish-Culturist*, 39, 179-181.

Kostenko, S., Dumitriu, G., Lægreid, K.J., Moens, U. (2011). Physiological roles of mitogen-

activated-protein-kinase-activated p38-regulated/activated protein kinase. *World Journal of*

Biological Chemistry, 2, 73-89.

Krens, S.G., He, S., Spaink, H.P., Snaar-Jagalska, B.E. (2006). Characterization and expression

patterns of the MAPK family in zebrafish. *Gene Expression Patterns*, 6, 1019-1026.

Lee, W.S., Salinas, S., Lee, Y.R., Siskidis, J.A., Mangel, M., Munch, S.B. (2020). Thermal

transgenerational effects remain after two generations. *Ecology and Evolution*, 10, 11296-11303.

Lee, D., Kim, K.H., Park, J.W., Lee, J.H., Kim, J.H. (2023). High water temperature-mediated

immune gene expression of olive flounder, *Paralichthys olivaceus* according to pre-stimulation

at high temperatures. *Environmental Toxicology and Pharmacology*, 101, 104159.

Leitwein, M., Wellband, K., Cayuela, H., Le Luyer, J., Mohns, K., Withler, R., Bernatchez, L.

(2022). Strong parallel differential gene expression induced by hatchery rearing weakly

associated with methylation signals in adult Coho Salmon (O. kisutch). *Genome Biology and*

Evolution, 14, evac036.

Liu, Z., Huang, X., Yang, Z., Peng, C., Yu, H., Cui, C., Hu, Y., Wang, X., Xing, Q., Hu, J., Bao,

Z. (2021). Identification, characterization, and expression analysis reveal diverse regulated roles

of three MAPK genes in *Chlamys farreri* under heat stress. *Frontiers in Physiology*, 12, 688626.

Love, M.I., Huber, W., Anders, S. (2014). Moderated estimation of fold change and dispersion

for RNA-seq data with DESeq2. *Genome Biology*, 15, 1-21.

Lu, X., Chen, H.M., Qian, X.Q., Gui, J.F. (2020). Transcriptome analysis of grass carp

(*Ctenopharyngodon idella*) between fast-and slow-growing fish. *Comparative Biochemistry and*

Physiology Part D: Genomics and Proteomics, 35, 100688.

Luu, I., Ikert, H., Craig, P.M. (2021). Chronic exposure to anthropogenic and climate related

stressors alters transcriptional responses in the liver of zebrafish (*Danio rerio*) across multiple

generations. *Comparative Biochemistry and Physiology Part C: Toxicology &*

Pharmacology, 240, 108918.

Marshall, D.J. (2015). Environmentally induced (co)variance in sperm and offspring phenotypes

as a source of epigenetic effects. *Journal of Experimental Biology*, 218, 107–113.

Martin N.V., Olver C.H. (1980). The lake charr, *Salvelinus namaycush*. In: E.K. Balon (Ed.),

Charrs: Salmonid Fishes of the Genus Salvelinus, Ed 1st (pp. 205–277). Netherlands: Springer.

McCaw, B.A., T.J. Stevenson, L.T. Lancaster. (2020). Epigenetic responses to temperature and

climate. *Integrative and Comparative Biology*, 60, 1469-1480.

McDermid, J.L.,Wilson, C.C., Sloan, W.N., Shuter, B.J. (2013). Intraspecific differences in

thermal biology of inland lake trout populations. *Transactions of the American Fisheries Society*,

142, 756–766.

Morash, A.J., Speers-Roesch, B., Andrew, S., Currie, S. (2021). The physiological ups and

downs of thermal variability in temperate freshwater ecosystems. *Journal of Fish Biology*, 98,

1524-1535.

OMNR Fish Culture Stocks Catalogue. (2005). Fish Culture Section. Peterborough, ON: Ontario

Ministry of Natural Resources.

Oomen, R. A., Hutchings, J. A. (2017). Transcriptomic responses to environmental change in

fishes: Insights from RNA sequencing. *Facets*, *2*, 610-641.

Penney, C.M., Burness, G., Tabh, J.K.R., Wilson, C.C. (2021). Limited transgenerational effects

of environmental temperatures on thermal performance of a cold-adapted salmonid.

Conservation Physiology, 9, coab021.

Penney, C.M., Tabh, J.K., Wilson, C.C., Burness, G. (2022). Within-generation and

transgenerational plasticity of a temperate salmonid in response to thermal acclimation and acute

temperature stress. *Physiological and Biochemical Zoology*, 95, 484-499.

Pérez-Ruzafa, A., Pérez-Marcos, M., Marcos, C. (2018). From fish physiology to ecosystems

management: Keys for moving through biological levels of organization in detecting

environmental changes and anticipate their consequences. *Ecological Indicators*, 90, 334-345.

Perrier, C., Ferchaud, A.L., Sirois, P., Thibault, I., Bernatchez, L. (2017). Do genetic drift and

accumulation of deleterious mutations preclude adaptation? Empirical investigation using

RADseq in a northern lacustrine fish. *Molecular Ecology*, 26, 6317–6335.

Petitjean, Q., Jean, S., Gandar, A., Côte, J., Laffaille, P., Jacquin, L. (2019). Stress responses in

fish: From molecular to evolutionary processes. *Science of the Total Environment*, 684, 371-380.

Quinn, N.L., McGowan, C.R., Cooper, G.A., Koop, B.F., Davidson, W.S. (2011). Identification

of genes associated with heat tolerance in Arctic charr exposed to acute thermal

stress. *Physiological Genomics*, 43, 685-696.

Riley, S.C., Hansen, M.J., Krueger, C.C., Noakes, D.L.G., Muir, A.M. (2021). Introduction. The

Lake Charr: Biology, Ecology, Distribution, and Management. In: A.M. Muir, M.J. Hansen, C.C.

Krueger, & S.C. Riley (Eds.), The Lake Charr *Salvelinus namaycush*: Biology, Ecology,

Distribution, and Management, Fish & Fisheries Series 39 (pp. 1-12). Switzerland: Springer

Nature.

Rosenfeld, J., Richards, J., Allen, D., Van Leeuwen, T., Monnet, G. (2020). Adaptive trade-offs
in fish energetics and physiology: insights from adaptive differentiation among juvenile
salmonids. *Canadian Journal of Fisheries and Aquatic Sciences*, 77, 1243-1255.
928

929 Rutkowska, J., Lagisz, M., Bonduriansky, R., Nakagawa, S. (2020). Mapping the past, present
930 and future research landscape of paternal effects. *BMC Biology*, 18, 1-24.
931

932 Ryu, T., Veilleux, H.D., Munday, P.L., Jung, I., Donelson, J.M., Ravasi, T. (2020). An
933 epigenetic signature for within-generational plasticity of a reef fish to ocean warming. *Frontiers
934 in Marine Science*, 7, 284.
935

936 Salinas, S., Munch, S.B. (2012). Thermal legacies: transgenerational effects of temperature on
937 growth in a vertebrate. *Ecology Letters*, 15, 159-163.
938

939 Sambrook, J., Russell, D.W. (2001). Molecular Cloning: a laboratory manual. Cold Spring
940 Harbor: Cold Spring Harbor Laboratory Press.
941

942 Sánchez-Tójar, A., Lagisz, M., Moran, N.P., Nakagawa, S., Noble, D.W., Reinhold, K. (2020).
943 The jury is still out regarding the generality of adaptive 'transgenerational' effects. *Ecology
944 Letters*, 23, 1715–1718.
945

946 Shahjahan, M., Zahangir, M.M., Islam, S.M.M., Ashaf-Ud-Doulah, M., Ando, H. (2021). Higher
947 acclimation temperature affects growth of rohu (*Labeo rohita*) through suppression of GH and
948 IGFs genes expression actuating stress response. *Journal of Thermal Biology*, 100, 103032.
949

950 Shama, L.N., Strobel, A., Mark, F.C., Wegner, K.M. (2014). Transgenerational plasticity in
951 marine sticklebacks: maternal effects mediate impacts of a warming ocean. *Functional Ecology*,
952 28, 1482–1493.
953

Shama, L.N., Mark, F.C., Strobel, A., Lokmer, A., John, U., Mathias Wegner, K. (2016).

Transgenerational effects persist down the maternal line in marine sticklebacks: gene expression

matches physiology in a warming ocean. *Evolutionary Applications*, 9, 1096–1111.

Shi, L., Tu, B.P. (2015). Acetyl-CoA and the regulation of metabolism: mechanisms and

consequences. *Current opinion in cell biology*, 33, 125-131.

Smith, D. A., Ridgway, M. S. (2019). Temperature selection in brook charr: lab experiments,

field studies, and matching the Fry curve. *Hydrobiologia*, *840*, 143-156.

Smith, D.A., Jackson, D.A., Ridgway, M.S. (2020). Thermal habitat of brook trout in lakes of

different size. *Freshwater Science*, 39, 56-69.

Smith, S.R., Normandeau, E., Djambazian, H., Nawarathna, P.M., Berube, P., Muir, A.M.,

Ragoussis, J., Penney, C.M., Scribner, K.T., Luikart, G., Wilson, C.C., Bernatchez, L. (2022). A

chromosome-anchored genome assembly for Lake Trout (*Salvelinus namaycush*). *Molecular

Ecology Resources*, 22, 679-694.

Somero, G.N. (2010). The physiology of climate change: how potentials for acclimatization and

genetic adaptation will determine 'winners' and 'losers'. *Journal of Experimental Biology*, 213,

912–920.

Spadafora, C. (2020). Transgenerational epigenetic reprogramming of early embryos: a

mechanistic model. *Environmental Epigenetics*, 6, dvaa009.

Stephens, M. (2017). False discovery rates: a new deal. *Biostatistics*, 18, 275-294.

Stewart, H.A., Aboagye, D.L., Ramee, S.W., Allen, P.J. (2019). Effects of acute thermal stress on

acid–base regulation, haematology, ion-osmoregulation and aerobic metabolism in Channel

Catfish (*Ictalurus punctatus*). *Aquaculture Research*, 50, 2133-2141.

Templeman, N.M., LeBlanc, S., Perry, S.F., Currie, S. (2014). Linking physiological and cellular
responses to thermal stress: β-adrenergic blockade reduces the heat shock response in fish.
Journal of Comparative Physiology B, 184, 719-728.
988
989 Ueno, T., Tomita, J., Kume, S., Kume, K. (2012). Dopamine modulates metabolic rate and
990 temperature sensitivity in *Drosophila melanogaster. PloS one*, 7, e31513.
991
992 Uller, T., Nakagawa, S., English, S. (2013). Weak evidence for anticipatory parental effects in
993 plants and animals. *Journal of Evolutionary Biology*, 26, 2161-2170.
994
995 Veilleux, H.D., Ryu, T., Donelson, J.M., Van Herwerden, L., Seridi, L., Ghosheh, Y., Berumen,
996 M.L., Leggat, W., Ravasi, T., Munday, P. L. (2015). Molecular processes of transgenerational
997 acclimation to a warming ocean. *Nature Climate Change*, 5, 1074-1078.
998
999 Veilleux, H.D., Donelson, J.M., Munday, P.L. (2018). Reproductive gene expression in a coral
1000 reef fish exposed to increasing temperature across generations. *Conservation Physiology*, 6, doi:
1001 10.1093/conphys/ cox077.
1002
1003 Venney, C.J., Love, O.P., Drown, E.J., Heath, D.D. (2020). DNA methylation profiles suggest
1004 intergenerational transfer of maternal effects. *Molecular Biology and Evolution*, 37, 540-548.
1005
1006 Venney, C.J., Wellband, K.W., Normandeau, E., Houle, C., Garant, D., Audet, C., Bernatchez, L.
1007 (2022). Thermal regime during parental sexual maturation, but not during offspring rearing,
1008 modulates DNA methylation in brook charr (*Salvelinus fontinalis*). *Proceedings of the Royal
1009 Society B: Biological Sciences*, 289, 20220670.
1010
1011 Vergauwen, L., Benoot, D., Blust, R., Knapen, D. (2010). Long-term warm or cold acclimation
1012 elicits a specific transcriptional response and affects energy metabolism in zebrafish.
1013 *Comparative Biochemistry and Physiology Part A: Molecular & Integrative Physiology*, 157,
1014 149-157.
1015

Visser, M.E. (2008). Keeping up with a warming world; assessing the rate of adaptation to climate change. *Proceedings of the Royal Society B: Biological Sciences*, 275, 649–659.

Willi, Y., Van Buskirk, J., Hoffmann, A.A. (2006). Limits to the adaptive potential of small populations. *Annual Review of Ecology, Evolution and Systematics*, 37, 433–458.

Wilson, C. (2017). Genetic aspects of climate change influences on inland fishes and fisheries. *Fisheries*, 42, 125–126.

Wilson, C.C., Mandrak, N.E. (2021). Paleoecology. In: A.M. Muir, M.J. Hansen, C.C. Krueger, & S.C. Riley (Eds.), The Lake Charr *Salvelinus namaycush*: Biology, Ecology, Distribution, and Management, Fish & Fisheries Series 39 (pp. 41-68). Switzerland: Springer Nature.

Yano, A., Nicol, B., Jouanno, E., Quillet, E., Fostier, A., Guyomard, R., Guiguen, Y. (2013). The sexually dimorphic on the Y-chromosome gene (sdY) is a conserved male-specific Y-chromosome sequence in many salmonids. Evolutionary applications, 6, 486-496.

Yin, J., Zhou, M., Lin, Z., Li, Q.Q., Zhang, Y.Y. (2019). Transgenerational effects benefit offspring across diverse environments: a meta-analysis in plants and animals. *Ecology Letters*, 22, 1976–1986.

Yu, G., Wang, L., Han, Y., He, Q. (2012). clusterProfiler: an R package for comparing biological themes among gene clusters. *OMICS: A Journal of Integrative Biology*, 16, 284-287.

Zhang, X., Flato, G., Kirchmeier-Young, M., Vincent, L., Wan, H., Wang, X., Rong, R., Fyfe, J., Li, G., Kharin, V.V. (2019). Changes in Temperature and Precipitation Across Canada. In: E. Bush & D.S. Lemmen (Eds.) Canada's Changing Climate Report (pp. 112-193). Ottawa, Ontario: Government of Canada.

1044 **Table 1**: Details for the lake trout offspring groups used to assess differential gene expression following within- and transgenerational
1045 thermal acclimation as shown in Figure 1. The measured effect is determined by comparing each treatment group against the control
1046 group.

	Treatment group (parent treatment)	Sequence year (sample size)	Offspring sex ratio (♂,♀)	Measured effect
Control group	Cold-acclimated offspring (C♀xC♂)	2018 (n=6) & 2020 (n=6)	2,4 5,1	*No effect (all other groups are compared to the control group)*
Within-generation warm acclimation	Warm-acclimated offspring (C♀xC♂)	2018 (n=6)	3,3	Offspring warm acclimation
Transgenerational warm acclimation	Cold-acclimated offspring (W♀xC♂)	2020 (n=6)	4,2	Maternal warm acclimation
	Cold-acclimated offspring (C♀xW♂)	2020 (n=6)	4,2	Paternal warm acclimation
	Cold-acclimated offspring (W♀xW♂)	2018 (n=6)	3,3	Maternal & paternal warm acclimation
Within- and transgenerational warm acclimation	Warm-acclimated offspring (W♀xC♂)	2020 (n=6)	2,4	Offspring & maternal warm acclimation
	Warm-acclimated offspring (C♀xW♂)	2020 (n=6)	3,3	Offspring & paternal warm acclimation
	Warm-acclimated offspring (W♀xW♂)	2018 (n=6)	5,1	Offspring, maternal & paternal warm acclimation

1047

1048   Treatment group describes the offspring's thermal acclimation as cold or warm, and the parent treatment is described in brackets,

1049   mother listed first (C=cold, W=warm).

1050 **Figures**

1051

1052

Control group:
Offspring C-A
Parental C-A
(C♀ x C♂)

WGP:
Offspring W-A
Parental C-A
(C♀ x C♂)

TGP:
Offspring C-A
Maternal W-A
(W♀ x C♂)

WGP + TGP:
Offspring W-A
Maternal W-A
(W♀ x C♂)

TGP:
Offspring C-A
Paternal W-A
(C♀ x W♂)

WGP + TGP:
Offspring W-A
Paternal W-A
(C♀ x W♂)

TGP:
Offspring C-A
2 Parents W-A
(W♀ x W♂)

WGP + TGP:
Offspring W-A
2 Parents W-A
(W♀ x W♂)

1053

1054

Figure 1: Experimental treatment groups of lake trout (*Salvelinus namaycush*). Two males and
two females each from cold- and warm acclimation (C-A and W-A, respectively) tanks were
bred in a full factorial cross, resulting in four families per parental treatment group (C♀xC♂,
W♀xC♂, C♀xW♂, W♀xW♂; one each depicted here for simplicity). Offspring from each family
was divided and acclimated to either a cold (control; blue: 11°C) or warm (pink: 15°C)
temperature.

1061

Figure 2: Principal component analysis to explore variance and correct the batch effect between sequencing runs in the gene expression data of the lake trout offspring groups. The PCA plots show ordination of all samples A) before and B) after batch correction. Panel C shows a scree plot of all principal components that cumulatively explain approximately 90% of the variance in the dataset.

1069

1070

Figure 3: Distribution of up- and downregulated differentially expressed genes in juvenile lake
trout with within-generation warm acclimation (WGP), transgenerational warm acclimation
(TGP) and the combined effect of both conditions (WGP + TGP) compared against the control
group (11°C C♀xC♂; not shown). Each comparison represents a treatment group (n = 6
individuals). Offspring acclimation temperature (11 or 15°C) and parental treatment (C=cold,
W=warm) are provided in parentheses. Gene expression levels are represented as log2-fold
change with negative values representing downregulated genes, and vice versa.

1078

1079

Figure 4: The number of differentially expressed genes that were commonly expressed (or
overlapped) between the juvenile lake trout offspring treatment groups (n=6 per group)
compared against the control group (11°C C♀xC♂). The number of differentially expressed genes
is listed above each vertical bar. The dots under each bar denote the groups that share those
genes (i.e. overlapped genes between groups). A single dot beneath a bar means that these genes
are uniquely expressed in that one group. Treatments included within-generation warm
acclimation (WGP), transgenerational warm acclimation (TGP) and the combined effect of both
conditions (WGP + TGP). Offspring acclimation temperature (11°C or 15°C) and parental
treatment (C=cold, W=warm) are provided in parentheses.

1089

Figure 5: Enriched GO (Gene Ontology) terms for differentially expressed genes in juvenile lake trout offspring (n=6) that were warm-acclimated (within-generation). The GO plot lists the top five most enriched GO terms for each GO category: molecular function (MF), cellular components (CC) and biological processes (BP). The number of differentially expressed genes is provided at the end of each bar. The -log$_e$ of the adjusted p-values is given on the x-axis and the threshold (p≤0.05) is indicated by the red vertical line at 2.99.

1101

Figure 6: Enriched GO (Gene Ontology) terms for differentially expressed genes in juvenile lake
trout offspring (n=6) that received maternal transgenerational warm acclimation. There were no
enriched GO terms for paternal transgenerational warm acclimation, or for transgenerational
warm acclimation of both parents. The GO plot lists the top five most enriched GO terms for
each GO category: molecular function (MF), cellular components (CC) and biological processes
(BP). The number of differentially expressed genes is provided at the end of each bar. . The -log$_e$
of the adjusted p-values is given on the x-axis and the threshold (p≤0.05) is indicated by the red
vertical line at 2.99.

1110

1111

Figure 7: Enriched GO (Gene Ontology) terms for differentially expressed genes in juvenile lake
trout offspring (n=6 per group) that were warm-acclimated (within-generation) in combination
with transgenerational (parental) warm acclimation of A) the mothers, B) the fathers and C) both
parents. The GO plot lists the top five most enriched GO terms for each GO category: molecular
function (MF), cellular components (CC) and biological processes (BP). The number of

1117   differentially expressed genes is provided at the end of each bar. The -log$_e$ of the adjusted p-
1118   values is given on the x-axis and the threshold (p≤0.05) is indicated by the red vertical line at
1119   2.99.