1
2
Styxl2 regulates *de novo* sarcomere assembly by binding to non-muscle myosin IIs and promoting their degradation

Xianwei Chen[1#], Yanfeng Li[1#], Jin Xu[1], Yong Cui[2], Qian Wu[3], Haidi Yin[3], Yuying Li[4], Chuan Gao[1], Liwen Jiang[2], Huating Wang[4], Zilong Wen[1], Zhongping Yao[3], Zhenguo Wu[1*]

1. Division of Life Science, the Hong Kong University of Science & Technology, Hong Kong, China; 2. School of Life Sciences, the Chinese University of Hong Kong, Hong Kong, China; 3. Department of Applied Biology and Chemical Technology, the Hong Kong Polytechnic University, Hong Kong, China; 4. Department of Orthopaedics and Traumatology, Li Ka Shing Institute of Health Sciences, the Chinese University of Hong Kong, Hong Kong, China

#equal contribution

*Corresponding author:

Division of Life Science,
the Hong Kong University of Science & Technology,
Clearwater Bay, Kowloon
Hong Kong
China
Tel: (852)23588704
Fax: (852)23581552
Email: bczgwu@ust.hk

Running title: Styxl2 promotes sarcomere assembly

Key words:

Styxl2; pseudophosphatase; sarcomere; non-muscle myosin II; autophagy

Abstract

Styxl2, a poorly characterized pseudophosphatase, was identified as a transcriptional target of the Jak1-Stat1 pathway during myoblast differentiation in culture. Styxl2 is specifically expressed in vertebrate striated muscles. By gene knockdown or genetic knockout, we found that Styxl2 plays an essential role in maintaining sarcomere integrity in developing muscles of both zebrafish and mice. To further reveal the functions of Styxl2 in adult muscles, we generated two inducible knockout mouse models: one with *Styxl2* being deleted in mature myofibers to assess its role in sarcomere maintenance, and the other in adult muscle satellite cells (MuSCs) to assess its role in *de novo* sarcomere assembly. We find that Styxl2 is not required for sarcomere maintenance but functions in *de novo* sarcomere assembly during injury-induced muscle regeneration. Mechanistically, Styxl2 interacts with non-muscle myosin IIs, enhances their ubiquitination, and targets them for autophagy-dependent degradation. Without Styxl2, the degradation of non-muscle myosin IIs is delayed, which leads to defective sarcomere assembly and force generation. Thus, Styxl2 promotes *de novo* sarcomere assembly by interacting with non-muscle myosin IIs and facilitating their autophagic degradation.

Introduction

Striated muscles, including skeletal and cardiac muscles of vertebrates and some muscles of invertebrates, are unique in that they contain highly-organized, multi-protein contractile structures called myofibrils in the cytosol of myofibers or cardiomyocytes (Clark et al., 2002; Mukund & Subramaniam, 2020; Sparrow & Schöck, 2009). Sarcomeres are the smallest repeating structural units of myofibrils responsible for muscle contraction (Avellaneda et al., 2021; Squire et al., 2005). A sarcomere is flanked by Z-discs (or Z lines) with multiple actin filaments (or thin filaments) anchored to the Z-discs in parallel and the interdigitating bipolar myosin bundles (or thick filaments) anchored to the M-line in the middle of the sarcomere (Squire et al., 2005). As a key component of the Z-disc, an anti-parallel α-actinin dimer directly binds to two actin filaments in opposite directions (Squire et al., 2005). A sarcomere can be further divided into the lighter I (isotropic)-band and the darker A (anisotropic)-band based on their microscopic appearances with the I-band containing actin filaments only and the A-band containing both the myosin bundles and partially overlapping actin filaments (Craig & Padrón, 2004; Squire et al., 2005). Mutations in sarcomeric proteins are linked to congenital human cardiac and skeletal muscle diseases (Bönnemann & Laing, 2004; Parker et al., 2020; Parker & Peckham, 2020). Although the overall structures of sarcomeres are well-established and the protein components of sarcomeres are largely known, it remains controversial how sarcomeres correctly assemble *in vivo*. Several different but not mutually exclusive models including the template model, the stitching model, and the premyofibril model have been proposed (Martin & Kirk, 2020; Rui et al., 2010; Sanger et al., 2006). The template model proposes that actin stress fibers serve as an early template to recruit other sarcomeric components including Z-disc proteins and the thin and thick filaments (Dlugosz et al., 1984). The stitching model or the titin model proposes that the actin filaments held by the Z-disc (i.e., the I-Z-I complexes) and the myosin filaments first assemble independently, and they are then stitched together to form intact sarcomeres with the help of titin as titin is known to simultaneously interact with Z-disc proteins via its N-terminal region and M-line proteins via its C-terminal region (Holtzer et al., 1997; Lu et al., 1992). The premyofibril model states that mini-sarcomeres consisting of actin filaments, α-actinin-containing Z bodies, and non-muscle myosin IIs first assemble near cell periphery. Then, non-muscle myosin IIs are replaced by muscle myosin II during the transition from premyofibrils to nascent myofibrils and finally to mature myofibrils (Rhee et al., 1994; Sanger et al., 2006). Each of these models is supported by some experimental

observations but may be incompatible with others (Chopra et al., 2018; Fenix et al., 2018;

Rui et al., 2010). In addition, mechanical tension generated by attachment of myofibers to

tendon, myofiber bundling, and formation of mini-sarcomeres is also found to promote

sarcomere assembly in both fly and mammalian muscle cells (Chopra et al., 2018; Loison et

al., 2018; Luis & Schnorrer, 2021; Mao et al., 2022; Weitkunat et al., 2014).

Protein phosphatases are a family of enzymes that catalyze the removal of the

phosphate group from covalently-linked substrates including proteins, lipids, nucleic acids,

and polysaccharides (Alonso et al., 2004; Tonks, 2006; Weinfeld et al., 2011; Worby et al.,

2006). They antagonize the actions of kinases and regulate various biological processes

including tissue development, cell proliferation, differentiation, and cell migration.

Interestingly, some of the protein phosphatases are found to be catalytically inactive due to

the absence of essential amino acid residues such as the catalytic cysteine in an otherwise

well-conserved phosphatase domain (M. J. Chen et al., 2017; Reiterer et al., 2020). These

confirmed or putative catalytically inactive phosphatases are also called pseudophosphatases.

Although catalytically inactive, several pseudophosphatases are found to play important roles

in various organisms (Azzedine et al., 2003; Cheng et al., 2009; Hinton, 2020; Reiterer et al.,

2017). For example, Styx and Sbf1, both members of the pseudophosphatase family, were

found to regulate sperm development. Loss of Styx or Sbf1 resulted in male infertility in mice

(Firestein et al., 2002; Wishart & Dixon, 2002).

In our current study, we identified Styxl2 (previously named as Dusp27), a poorly

characterized pseudophosphatase, as a new regulator involved in sarcomere assembly in both

zebrafish and mice. The original name of the gene and its protein product (i.e., Dusp27) has

caused confusion in the literature due to the fact that the same name was once used for Dupd1

(now re-named as Dusp29), a catalytically active phosphatase (Friedberg et al., 2007).

Recently, it was proposed that Dusp27 be re-named as Serine/threonine/tyrosine-interacting-

like 2 (Styxl2) (Cooper & Waddell, 2020). To avoid further confusion, we will use Styxl2

hereafter. Styxl2 is specifically expressed in vertebrate striated muscles. Our work in both

zebrafish and mouse models shows that it plays an important role in *de novo* sarcomere

assembly instead of sarcomere maintenance. Mechanistically, we found that Styxl2 directly

interacts with non-muscle myosin IIs that are involved in the early stage of sarcomere

assembly and subsequently replaced by muscle myosin IIs in mature sarcomeres (Wang et al.,

2018). Styxl2 promotes ubiquitination and autophagy-dependent degradation of non-muscle

myosin IIs, which ultimately facilitates sarcomere assembly *in vivo*.

Results

***Styxl2* encodes a striated muscle-specific pseudophosphatase and is a transcriptional target of the Jak1-Stat1 pathway in myoblasts**

We previously reported that the Jak1-Stat1 pathway plays an important role in myogenic differentiation in cultured myoblasts (Sun et al., 2007). However, it remains unclear what genes are targeted by the pathway. To examine the changes in transcriptomes, we conducted a microarray analysis in C2C12 myoblasts: cells were first transfected with either a control siRNA or specific siRNAs targeting *Jak1* or *Stat1* and the cells were harvested at two different time points: (1) after 24 hours of growth in the growth medium (GM 24h, the proliferation stage); (2) after 12 hours in the differentiation medium following 24 hours of growth in the GM (DM 12h, the early differentiation stage). The total mRNA was extracted and subjected to microarray analysis. Among differentially expressed genes (DEGs) in cells transfected with *Jak1*-siRNA and *Stat1*-siRNA, many were shared, which was expected as both Jak1 and Stat1 function in the same pathway. Some representative targets shared by Jak1 and Stat1 were shown (Table 1). Consistent with our previous findings (Sun et al., 2007), *myogenin* gene was negatively regulated by both Jak1 and Stat1. Among the genes targeted by both Jak1 and Stat1, *serine/threonine/tyrosine-interacting-like 2* (*Styxl2*), previously also named as *dual-specificity phosphatase 27 (putative)* (*Dusp27*), attracted our attention: it is a poorly characterized gene with only one published report in zebrafish so far (Fero et al., 2014). It encodes a member of the dual-specificity phosphatase family with a conserved dual-specificity phosphatase catalytic (DSPc) domain located at its N-terminus (Figure 1A). However, Styxl2 is predicted to be catalytically inactive due to substitution of a highly-conserved catalytic cysteine by serine in its DSPc domain (Figure 1-figure supplement 1A), thus making it a member of the pseudophosphatase subfamily (Hinton, 2019; Kharitidi et al., 2014). By both RT-qPCR and Western blot, we confirmed that both the mRNA and protein levels of *Styxl2* were induced by siRNAs against *Jak1* or *Stat1* in C2C12 myoblasts, with the extent of induction being more obvious during early differentiation (Figure 1-figure supplement 1B, C). Consistently, treatment of C2C12 myoblasts with leukemia inhibitory factor (LIF) or oncostatin M (OSM), two cytokines that are known to potently activate the Jak1-Stat1 pathway in myoblasts (Sun et al., 2007; Xiao et al., 2011), indeed led to a gradual decrease of Styxl2 protein (Figure 1-figure supplement 1D, E). By Western blot, we found that *Styxl2* has very limited tissue distribution pattern: it is specifically expressed in striated muscles including the heart and skeletal muscles (Figure 1B). Consistently, the expression of

Styxl2 mRNA is confined to the heart and somites of early mouse embryos (Figure 1-figure

supplement 1F). In C2C12 myoblasts, Styxl2 protein was already present in proliferating

myoblasts but was further induced during myogenic differentiation (Figure 1C). We also

examined the subcellular localization of Styxl2 in myoblasts. As our home-made polyclonal

antibody is not suitable for immunostaining of the endogenous Styxl2, we examined

subcellular localization of Flag- or HA-tagged Styxl2 instead in C2C12 cells by

immunostaining. We found that both Flag- and HA-tagged Styxl2 were predominantly

localized in the cytoplasm (Figure 1-figure supplement 1G). Collectively, Styxl2 is a

cytoplasmic protein that is specifically expressed in striated muscles.

Knockdown of *Styxl2* in zebrafish leads to severe defects in sarcomere structure

An earlier study by Fero *et al.* in zebrafish showed that Styxl2 regulates the integrity of

sarcomeres (Fero et al., 2014). We also confirmed this finding in zebrafish using a

morpholino oligo (MO) that blocks the translation initiation of *Styxl2*. The effectiveness of

the MO was verified using a transgene encoding green fluorescence protein (GFP) fused in-

frame with the same sequence of *Styxl2* (i.e., the first 25 nucleotides of the coding region)

targeted by the MO: when zebrafish zygotes were microinjected with the transgenes with or

without the MO, the expression of GFP was suppressed only when the MO was co-injected

(Figure 1-figure supplement 1H). We then injected zebrafish zygotes with or without the

Styxl2-MO and examined the structures of skeletal muscles by immunostaining at 24 hours

after fertilization (hpf). Consistent with the report by Fero *et al.* (Fero et al., 2014), we found

that sarcomeres in both fast and slow muscles were severely disrupted when *Styxl2* was

knocked down with the MO (Figure 1E). Moreover, electron microscopy analysis further

revealed the disrupted sarcomere structures in zebrafish muscles injected with the MO

(Figure 1D).

Conditional deletion of *Styxl2* leads to neonatal lethality in mutant mice

To further explore the role of Styxl2 in mammals, we generated *Styxl2* floxed mice with two

LoxP sites flanking exons 3~5 of the *Styxl2* gene (Figure 2-figure supplement 1A). Exons 4

and 5 encode a part of the DSPc domain of Styxl2. We first generated germline *Styxl2*

knockout (KO) mice using EIIA-Cre mice (Lakso et al., 1996). Western blot analysis showed

that Styxl2 protein was totally absent in KO mice (Figure 2A). In multiple litters, no live

homozygous KO pups were found after birth (Figure 2-figure supplement 1B). By examining

developing embryos of different ages, we found that *Styxl2* KO embryos died between E13.5

and E15.5 (Figure 2-figure supplement 1C). In some dead *Styxl2* KO foetuses, edema was
found (Figure 2B), suggesting potential problems with the heart (Conway et al., 2003). This
is consistent with the fact that Styxl2 is also highly expressed in cardiac muscles as well. The
electron microscopy analysis of the heart muscles from E14.5 embryos revealed that the
sarcomeric structures of cardiac muscles from *Styxl2* KO embryos were severely disrupted
(Figure 2C). To focus on the functions of Styxl2 in skeletal muscles, we generated
conditional *Styxl2* KO (CKO) mice by crossing *Styxl2$^{f/f}$* mice with *Pax7$^{Cre/+}$* mice (Keller et
al., 2004) (Figure 2-figure supplement 1D). We found that *Styxl2* CKO pups were born at the
expected Mendelian ratio but all the homozygous mutant pups died in a few hours after birth
(Figure 2D, E and Figure 2-figure supplement 1D). As Pax7 is also expressed in other non-
muscle cell types including neurons during embryo development, to be certain that
embryonic lethality of *Pax7$^{Cre/+}$; Styxl2$^{f/f}$* mice was indeed caused by severe defects in
skeletal muscles, we also specifically deleted *Styxl2* in skeletal muscles using a *Myf5^{Cre}* line
in which the Cre gene was inserted in the *Myf5* locus (Tallquist et al., 2000). Consistently, all
skeletal muscle-specific *Styxl2* KO pups died within one day after birth (Figure 2-figure
supplement 1D).
As the skin of *Styxl2* CKO pups appeared cyanotic, we speculated that the new-born
mutant mice had lung defects that led to respiratory distress and perinatal death. The lungs of
the CKO pups were examined and we found that the pulmonary alveoli indeed failed to open
(Figure 2F), which was likely caused by defective diaphragm muscle as defects in a number
of myogenic genes exhibited similar perinatal lethality (Hasty et al., 1993; Rudnicki et al.,
1993). Electron microscopy analysis confirmed that the sarcomere structure was severely
disrupted in diaphragm muscle of the mutant mice (Figure 2G). Consistently, electron
microscopy revealed that the sarcomere structure of the limb muscle was also severely
disrupted (Figure 2-figure supplement 1E). Notably, some sarcomeres were still present in the
skeletal muscles of the CKO mice (Figure 2-figure supplement 1E), suggesting that Styxl2 is
important but not absolutely required for the formation of the sarcomeres. Therefore, Styxl2
plays an important role in the development of striated muscles, including the skeletal muscle
and the heart muscle.
220

Styxl2 is dispensable for the maintenance of mature sarcomeres but critical for *de novo*
sarcomere assembly in adult skeletal muscles
Although our data above indicated that Styxl2 plays an important role in regulating
sarcomere formation during embryonic skeletal muscle development, it remained unclear

whether it was necessary for the maintenance of the sarcomere structure in adult skeletal

muscles. To address this question, we generated a tamoxifen (TMX) inducible myofiber-

specific *Styxl2* knockout mouse model (also named *Styxl2* MF-iKO hereafter) using *Tg (HSA-*

MerCreMer) (McCarthy et al., 2012). In this model, *Styxl2* was efficiently deleted in mature

skeletal muscles of adult mice (Figure 3A, B). We directly assessed the contractile functions

of Styxl2-null muscles. Three weeks after the last TMX injection (Figure 3A), the

gastrocnemius muscle was isolated and the twitch and tetanic force generated by the muscle

were measured (Guo et al., 2015). No obvious difference in twitch or tetanic force was

observed between the control and *Styxl2* MF-iKO muscles (Figure 3C, D), suggesting that

Styxl2 is dispensable for the maintenance of existing sarcomeres in adult skeletal muscles.

Next, we assessed whether Styxl2 played a role in the *de novo* formation of nascent

sarcomeres in regenerating muscles of adult mice. To do so, we generated another tamoxifen

inducible, MuSC-specific *Styxl2* KO mouse model using *Pax7^{CreERT2/CreERT2}; Styxl2^{f/f}* (also

named *Styxl2* SC-iKO mice hereafter) (Murphy et al., 2011). *Styxl2* was first deleted in

muscle satellite cells (MuSCs) of adult mice according to the scheme in Figure 3E, followed

by acute injury to the hindlimb skeletal muscles with Cardiotoxin (CTX) (Figure 3E). As

injured muscles are repaired by MuSCs, the nascent sarcomeres in regenerating myofibers of

Styxl2 SC-iKO mice would be formed in the absence of Styxl2 (Figure 3F). At 30 days post

injury, a time point when injured muscles are known to be fully regenerated (Dumont et al.,

2015), the gastrocnemius muscles were isolated and functionally tested *in vitro*. Both the

twitch and tetanic force were measured separately. Compared to muscles from the control

mice, the twitch force generated by muscles from *Styxl2* SC-iKO mice was decreased by ~30%

(Figure 3G), while the tetanic force generated by muscles from the mutant mice was not

affected (Figure 3H). This suggested that sarcomeres formed *de novo* in regenerating

myofibers from Styxl2-null MuSCs were partially defective. When we examined sarcomeres

by electron microscopy, we noticed that the length of individual sarcomeres from *Styxl2* SC-

iKO mice was shorter in regenerated muscles (Figure 3I, J). Thus, our data showed that

Styxl2 functions in *de novo* sarcomere assembly in regenerating muscles from adult mice.

Styxl2 directly binds to non-muscle myosin II (NM II) in myoblasts

To understand how Styxl2 functions to regulate sarcomere formation, we employed BioID, a

proximity-dependent protein labelling method (Roux et al., 2012, 2018), to search for binding

partners of Styxl2. The full-length Styxl2 was fused in-frame with BirA*, a mutant form of

biotin ligase BirA (Figure 4A). The fused gene encoding Styxl2-BirA*-HA was stably

expressed in C2C12 myoblasts using the Tet-on system (Das et al., 2016) (Figure 4-figure

supplement 1A). As Styxl2 is expected to function during late myogenic differentiation, we

followed the scheme in Figure 4B and added biotin into the culture medium with or without

doxycycline (Dox) for 24 hours before harvest. By Western blot, we showed that Styxl2-

BirA*-HA was strongly induced by Dox as expected (Figure 4-figure supplement 1A, middle

panel). Importantly, more cellular proteins were biotinylated in Styxl2-BirA*-HA-expressing

cells induced by Dox (Figure 4-figure supplement 1A, top panel). These biotinylated proteins

were isolated and enriched by streptavidin beads and analysed by iTRAQ-based mass

spectrometry (Wiese et al., 2007). The biotinylated proteins enriched in Styxl2-BirA*-HA-

expressing cells (Enrichment score > 1) were further analysed by gene ontology (GO)

analysis. Notably, a number of the enriched proteins were related to "myofibril", "sarcomere",

and "A band" (Figure 4C). This supports our hypothesis that Styxl2 functions to regulate

sarcomere assembly. To help identify direct interaction partners of Styxl2, the $\log_2$

Enrichment scores (i.e., the peak intensity ratios of the reporter ions) were plotted against the

minus $\log_{10}$(p.values). Those enriched proteins with the Enrichment score > 1.2 and $p < 0.05$

were highlighted in red on the plot (Figure 4-figure supplement 1B), and some of the

enriched proteins (Figure 4-figure supplement 1C) were selected for further confirmation by

co-immunoprecipitation (co-IP) assays using lysates from differentiated C2C12 myotubes.

Among them, Myh9 and Myh10, the members of the non-muscle myosin IIs (NM IIs)

(Newell-Litwa et al., 2015), were found to specifically interact with the endogenous Styxl2

(Figure 4D). Moreover, by protein truncation analysis in combination with the co-IP assays,

we showed that the head domains of Myh9 and Myh10 were responsible for their interaction

with Styxl2 (Figure 4E, F).

Styxl2 promotes sarcomere assembly by down-regulating NM IIs

Our data above indicated that Styxl2 regulates sarcomere assembly. NM IIs were reported to

participate in sarcomere assembly in several organisms (Fenix et al., 2018; Sanger et al., 2009;

Tullio et al., 1997). However, NM IIs are not detected in mature sarcomeres and their

replacement by muscle-specific myosin is a marker of skeletal muscle maturation. Therefore,

we examined protein levels of Myh9 and Myh10 in both differentiating C2C12 myotubes and

postnatal mouse skeletal muscles over time. In differentiating C2C12 cells, we found that the

protein levels of Myh9 and Myh10 slightly increased at the beginning of myogenic

differentiation, but started to decrease ~four days after myogenic differentiation, which

coincided with the time when the sarcomeric myosin heavy chain started to increase (Figure

5-figure supplement 1A). In postnatal mouse skeletal muscles, the protein levels of Myh9 and

Myh10 started to drop at P8 and became non-detectable by P30 (Figure 5A), which was

consistent with previous reports (Fenix et al., 2018; Sanger et al., 2009). The protein levels of

Styxl2 remained constant and abundant in the first month after birth but began to drop to a

lower level by P65 (Figure 5A). When we examined the protein levels of Myh9 and Myh10

in skeletal muscles of P1 pups, we noticed an obvious increase in protein levels of Myh9 and

Myh10 in *Styxl2* CKO mice (Figure 5B, C). We also examined protein levels of Myh9 and

Myh10 in regenerating adult muscles. Using SC-iKO mice and *Pax7*^CreERT2/CreERT2^ control

mice as described in Figure 3E, we first induced *Styxl2* deletion by tamoxifen in adult MuSCs

followed by BaCl$_2$-induced muscle injury. At 7 dpi, the levels of Myh9 and Myh10 were

comparable between control and *Styxl2* SC-iKO mice (Figure 5D). Interestingly, at 10 dpi,

Myh9 and Myh10 were nearly undetectable in muscles from control mice, but they remained

at a higher level in muscles from *Styxl2* SC-iKO mice (Figure 5D). Our data above suggested

that loss of Styxl2 results in delayed degradation of NM II during sarcomere assembly. To

further prove this is the case, we turned to the zebrafish model. It was reported that *Styxl2*^-/-^

zebrafish exhibited severe motility defects in response to needle touch (Fero et al., 2014).

Consistently, we also found that *Styxl2* morphants displayed similar phenotypes: most of the

Styxl2 morphants did not respond to the needle touch (Figure 5E), which was caused by

defective sarcomere structures in skeletal muscles (Figure 5F). Knowing that NM IIs are

involved in the early phase of sarcomere assembly followed by their replacement with the

sarcomeric myosin (Fenix et al., 2018; White et al., 2014), we suspected that delayed

degradation of NM IIs was responsible for defective sarcomere structure in *Styxl2* morphants.

Therefore, it was likely that simultaneous knockdown of both NM IIs and *Styxl2* may rescue

the sarcomeric defects in *Styxl2* morphants. It was reported that both z*Myh9b* and z*Myh10* are

expressed in skeletal muscles of zebrafish (Gutzman et al., 2015). When we knocked down

z*Myh9b* with a MO in *Styxl2* morphants, the embryos died very early before skeletal muscles

were formed (our unpublished data). When zebrafish zygotes were co-injected with the

Styxl2-MO and a MO targeting z*Myh10* (Figure 5-figure supplement 1B), the percentage of

fishes responding to the needle touch was obviously increased (Figure 5E). Consistently, the

disrupted sarcomere structure caused by *Styxl2*-MO was indeed partially restored in the

Styxl2/Myh10 double-knockdown fishes (Figure 5F).

Styxl2 induces the degradation of NM II via its C-terminal domain

Our data above showed that loss of Styxl2 caused abnormal accumulation of NM IIs (Figure
5), which suggested that Styxl2 normally promote degradation of NM IIs during nascent
sarcomere assembly. To test whether this is the case, we co-expressed HA-Myh9 with either
Mst1 (as a negative control) or Styxl2 in C2C12 myoblasts. The protein levels of HA-Myh9
were indeed reduced by co-transfected Styxl2 (Figure 6A). We further found that Styxl2-
mediated reduction in Myh9 protein levels was not caused by reduced *Myh9* mRNA (Figure
6-figure supplement 1A). To determine which part of Styxl2 promotes degradation of NM IIs,
we truncated Styxl2 into two fragments (Figure 6B, upper panel): an N-terminal fragment
containing the DSPc domain (i.e., aa 1-513, also termed Styxl2N513 hereafter) and a C-
terminal fragment without the DSPc domain (i.e., aa 510-1138, also named Styxl2ΔN509
hereafter). When we co-transfected 293T cells with a construct encoding HA-Myh9 together
with another construct encoding Mst1, Styxl2N513, or Styxl2ΔN509, we found that only
Styxl2ΔN509 promoted degradation of Myh9 (Figure 6B, lane 3). As we previously showed
that the head domains of NM IIs were involved in interacting with Styxl2 (Figure 4E, F), we
then tested whether the protein levels of the head domain of Myh9 were also regulated by
Styxl2. Consistently, only Styxl2ΔN509 promoted degradation of the head domain of Myh9
(Figure 6B, lane 6). To further verify this finding *in vivo*, we microinjected the mRNA
encoding the full-length or the C-terminus of zStyxl2, along with the MO for z*Styxl2*, into the
zebrafish one-cell embryos. As expected, the defective sarcomere structure caused by *zStyxl2*
knockdown was efficiently rescued by overexpressing the mRNA encoding the full-length
zStyxl2, as revealed by the α-actinin staining (Figure 6C). Importantly, the mRNA encoding
the C-terminus of zStyxl2 was nearly as effective as that encoding the full-length zStyxl2
(Figure 6C). Similar results were also obtained when we subjected fish to myosin staining
(Figure 6-figure supplement 1B). Thus, the C-terminus of Styxl2, without the DSPc domain,
is both necessary and sufficient to promote degradation of NM IIs during nascent sarcomere
assembly.
352

Styxl2 promotes the degradation of NM IIs via the autophagy-lysosome pathway
In order to understand how Styxl2 promotes the degradation of NM IIs, we first turned to the
ubiquitination/proteasome pathway (Nandi et al., 2006). MG132, a specific proteasome
inhibitor, was added to C2C12 cells transfected with Styxl2. No abnormal accumulation of
NM IIs was found in the presence of MG132 (Figure 7-figure supplement 1), suggesting that
the proteasome is not involved in Styxl2-mediated degradation of NM IIs. Next, we turned to

the autophagy-lysosome-dependent protein degradation system. In C2C12 myoblasts, we co-
expressed Myh9 with either Mst1 (control) or Styxl2 together with an siRNA against GFP
(control) or ATG5, a key component of the autophagy pathway (Mizushima et al., 2010). In
the presence of the control siRNA, Styxl2 efficiently promoted degradation of Myh9 as
expected (Figure 7A, lane 2). However, in the presence of ATG5 siRNA, the protein levels of
Myh9 were restored to levels comparable to that of the control even in the presence of the co-
expressed Styxl2 (Figure 7A, lane 3). This result supported the involvement of the autophagy
pathway in Styxl2-mediated degradation of NM IIs. Furthermore, we also examined the
degradation of Myh9 induced by Styxl2Δ509 with or without several different autophagy
inhibitors (Figure 7B). In the absence of any autophagy inhibitors, Styxl2ΔN509 induced
degradation of Myh9 as expected (Figure 7B, lanes 3 and 4). In the presence of LY294002
(an inhibitor of PI3K), chloroquine (an inhibitor that blocks the binding of autophagosomes
to lysosomes), bafilomycin A1 (BafA1, an inhibitor of V-ATPase that can block autophagic
flux), and NH_4Cl (an inhibitor of the degradative enzymes inside lysosomes), the protein
levels of Myh9 were largely restored to the levels comparable to that in the controls (Figure
7B, lanes 1 and 2). Thus, we concluded that the autophagy-lysosome pathway is involved in
the degradation of NMIIs induced by Styxl2.

It is known that some ubiquitinated proteins (such as sarcomeric titin) are degraded by
selective autophagy (Bogomolovas et al., 2021; R.-H. Chen et al., 2019; Kirkin et al., 2009;
Pankiv et al., 2007; Shaid et al., 2013; Zaffagnini & Martens, 2016). A recent report showed
that Myh9 could also undergo Nek9-mediated selective autophagy (Yamamoto et al., 2021),
suggesting that Myh9 is ubiquitinated. To find out whether Styxl2 promotes ubiquitination of
NM IIs, we co-transfected HEK 293T cells with plasmids encoding Myh9 and ubiquitin
together with a plasmid encoding either Mst1 or Styxl2. Compared to the Mst1 control, we
found that Styxl2 indeed promoted the ubiquitination of Myh9 (Figure 7C).

Discussion

Styxl2 is a pseudophosphatase that functions independently of its conserved phosphatase domain. Pseudophosphatases are present in multicellular organisms ranging from *C. elegans* to human (Hinton, 2019; Reiterer et al., 2020). Moreover, pseudophosphatases have also been identified in plants (Bellec et al., 2002). Recently, 26 pseudophosphatase domains, including the phosphatase domain of Styxl2, were bioinformatically identified in the human genome (M. J. Chen et al., 2017). Due to the absence of one or several conserved amino acids including the catalytic cysteine, pseudophosphatases are not expected to possess the phosphatase activity. STYX, the first molecularly characterized pseudophosphatase, was identified in 1995 as a phosphoserine/threonine/tyrosine-binding protein (Wishart et al., 1995). The substitution of the catalytic cysteine by glycine in the active site motif leads to the loss of its catalytic activity (Reiterer et al., 2013; Wishart et al., 1995). In addition to STYX, the functions of several other pseudophosphatases from *C. elegans* to human have also been characterized (Cheng et al., 2009; Hinton et al., 2010; Maruyama et al., 2007; Nakhro et al., 2013; Tomar et al., 2019). These reports unequivocally demonstrate that pseudophosphatases are not simply evolution relics but play essential roles in multicellular organisms.

Pseudophosphatases exert their functions via diverse modes including direct competition with other proteins by binding to a phospho-residue-containing motif, modulation of the phosphatase activity by binding to an active phosphatase, sequestration of target proteins in specific subcellular locations, and integration of signals by simultaneous binding with multiple proteins (Hinton, 2019; Reiterer et al., 2020). The original report on STYX showed that mutation of the glycine in the conserved HGX$_5$R motif back to cysteine restored the phosphatase activity (Wishart et al., 1995), suggesting that the three-dimensional structure of the phosphatase domain in STYX is still folded properly. Thus, some pseudophosphatases are thought to function by binding to specific phosphorylated substrates, which is the case for pseudophosphatases EGG-4/5 and Pas2 (Bellec et al., 2002; Cheng et al., 2009; Da Costa et al., 2006). In these examples, pseudophosphatases bind the phosphorylated sites of selected target proteins through their inactive phosphatase domains and prevent them from binding other interacting proteins or active phosphatases. However, this is not the only way pseudophosphatases function. MTMR13, a pseudophosphatase, functions by binding to MTMR2, an active lipid phosphatase, to regulate its phosphatase activity. Mutations in either *MTMR13* or *MTMR2* are causally linked to human Type 4B Charcot-Marie-Tooth disease, an

inherited disorder of the peripheral nervous system with abnormal nerve myelination (Berger

et al., 2006; Robinson & Dixon, 2005).

Unlike several other characterized pseudophosphatases that function via their

conserved phosphatase domains, Styxl2 appears to function independently of its phosphatase

domain as the phosphatase domain of Styxl2 is dispensable for its ability to induce

degradation of NM IIs and to rescue the sarcomere defects in the *Styxl2* morphant zebrafish

(Figure 6B-C). Such a phosphatase domain-independent mode of action is also observed in

His Domain Protein Tyrosine Phosphatase (HD-PTP or PTPN23), a pseudophosphatase

known to facilitate endosomal cargo sorting and the degradation of multiple membrane

receptors including epidermal growth factor receptor, platelet-derived growth factor receptor,

MHC class I and Integrin α5β1 (Doyotte et al., 2008; Kharitidi et al., 2015; H. Ma et al., 2015;

Parkinson et al., 2015). Interestingly, a truncated fragment of HD-PTP without its

phosphatase domain is still capable of promoting endosomal trafficking of cargo proteins like

epidermal growth factor (EGF) (Doyotte et al., 2008).

Styxl2 facilitates *de novo* sarcomere assembly by promoting autophagy-dependent

degradation of non-muscle myosin IIs. Although absent in invertebrates, Styxl2 is

expressed and conserved in vertebrates. The specific involvement of Styxl2 in sarcomere

assembly is supported by its restricted expression in striated muscles. Interestingly, mutation

in the coding region of *STYXL2* and abnormal CpG methylation at the *STYXL2* locus were

detected in some patients with congenital heart defects (CHD), suggesting a possible link of

STYXL2 to CHD (Arrington et al., 2012; Radhakrishna et al., 2016). Moreover, its role in

sarcomere assembly is also conserved in vertebrates as interference of the gene expression in

zebrafish by either a transposon integration or morpholino oligos leads to severe defects in

sarcomere assembly (Fero et al., 2014) (Figure 1D, E). However, Styxl2 is not essential for

sarcomere assembly as sarcomeres in *Drosophila* flight muscles can normally assemble in the

absence of a Styxl2 homolog. Furthermore, we could still find residual sarcomeres in skeletal

muscles of conditional *Styxl2* KO mice (driven by either *Pax7^Cre^* or *Myf5^Cre^*) (Figure 2-figure

supplement 1E).

Although the highly-ordered sarcomere structure has been known for many years, it

remains controversial how sarcomeres correctly assemble. Several different but not mutually

exclusive models have been proposed (Rui et al., 2010; Sanger et al., 2006, 2017). Studies in

Drosophila, *C. elegans*, and mice underscore an essential role for integrin in initiating the

assembly of sarcomeres (Rui et al., 2010; Sparrow & Schöck, 2009). For example, β1

integrin was shown to be essential for sarcomere assembly in mice (Schwander et al., 2003).

Among the existing sarcomere assembly models, the prevailing model of sarcomere assembly

is called premyofibril model (Rhee et al., 1994; Sanger et al., 2006, 2017). In this model, α-

actinin, skeletal muscle actin and non-muscle myosin IIs (NM IIs) first assemble to form

mini-sarcomeres or premyofibrils at the cell periphery. Then, additional sarcomeric proteins

like titin, nebulin, troponin, and myomesin will be incorporated and NM IIs will be

eventually replaced by skeletal muscle myosin in mature sarcomeres (White et al., 2014,

2018). Consistent with this model, germline knockout of NM IIB (i.e., Myh10) in mice led to

early lethality between E14.5 and birth and sarcomere structural defects in embryonic

cardiomyocytes (Tullio et al., 1997). Similarly, defective sarcomeres were also observed in

embryonic muscles of stage-17 mutant fly null for *zipper* (*zip*), the *Drosophila* gene encoding

the heavy chain of NM II (Bloor & Kiehart, 2001; Loison et al., 2018). Moreover, a recent

paper using human iPSC-derived cardiomyocytes demonstrated that nascent sarcomeres

assemble directly from muscle actin stress fibers (Fenix et al., 2018). Both MYH9 and

MYH10 are present in muscle actin stress fibers with MYH10 being essential for sarcomere

assembly. Consistent with the premyofibril model, NM IIs are found to be replaced by

muscle myosin IIs in mature sarcomeres (Fenix et al., 2018). Nevertheless, the view of the

involvement of NM IIs in early sarcomere assembly is also challenged by several studies.

Unlike germline knockout of NM IIB, cardiac-specific knockout of NM IIB in adult mice did

not generate obvious defects in sarcomere structures (X. Ma et al., 2009). Although NM IIA

is not detected in embryonic hearts, loss of cardiac NM IIB led to increased expression of

NM IIA, suggesting that NM IIA may compensate for the loss of NM IIB in NM IIB-null

embryonic cardiomyocytes (Fenix et al., 2018; Tullio et al., 1997). Moreover, it was also

suspected that the cardiomyocyte-specific deletion of NM IIB occurs after the initial

sarcomere assembly due to the expression timing of the Cre recombinase under the control of

the cardiac α-MHC promoter (Fenix et al., 2018; X. Ma et al., 2009). Furthermore, a recent

study using CRISPR/Cas9 to knock out *MYH9* and *MYH10* genes either individually or in

combination in human iPSC-derived cardiomyocytes showed that neither gene is required for

sarcomere assembly in their cell culture models (Chopra et al., 2018). It is puzzling at present

why the results on the involvement of NM IIs in *de novo* sarcomere assembly differ in the

two studies by Fenix *et al.* and Chopra *et al.* even though both studies used human iPSC-

derived cardiomyocytes to assay *de novo* sarcomere assembly in cell culture.

It also remains unclear how NM IIs are replaced by muscle myosin during *de novo*
sarcomere assembly in vertebrates. Our current study showed that Styxl2 interacts with NM
IIs and induces their degradation via the autophagy-lysosome pathway. Autophagy is known
to regulate sarcomere integrity (Masiero et al., 2009; Nakai et al., 2007). In particular,
selective autophagy is known to degrade multiple ubiquitinated sarcomeric proteins including
titin and filamin (Arndt et al., 2010; Bogomolovas et al., 2021). p62/SQSTM1 and NBR1
have been shown to act as adaptors linking ubiquitinated substrates to Atg8/LC3 that is
localized on the membrane of autophagosome (Kirkin et al., 2009; Pankiv et al., 2007).
Interestingly, a recent study on ciliogenesis showed that MYH9 is also degraded by selective
autophagy in an NEK9-mediated manner (Yamamoto et al., 2021). Consistently, we found
that Styxl2 promotes the ubiquitination of Myh9 (Figure 7C). It remains to be established
whether Nek9 is involved in Styxl2-mediated autophagic degradation of NM IIs and whether
ubiquitination of NM IIs is dependent on Nek9-mediated phosphorylation during
myofibrillogenesis.

Our data showed that Styxl2 plays indispensable role during embryonic myogenesis
as conditional deletion of *Styxl2* during mouse embryo development using either *Pax7^{Cre}* or
Myf5^{Cre} resulted in severe disruption of sarcomere structures in skeletal muscles and perinatal
lethality (Figure 2 and Figure 2-figure supplement 1). In contrast, inducible deletion of *Styxl2*
in adult muscles only led to mild defects in sarcomere structures (Figure 3F-J). It is likely that
additional unknown molecules can compensate for the loss of Styxl2 in adult regenerating
muscles. Such distinct roles of Styxl2 during development are also supported by differential
expression patterns of Styxl2 and NM IIs: both are abundantly expressed in new-born and
juvenile mice. While the protein levels of Myh9 and Myh10 start to drop one week after birth
and become undetectable by three weeks after birth, that of Styxl2 is relatively stable in the
first month after birth but starts to drop sharply by two months after birth (Figure 5A). This is
also consistent with our findings that Styxl2 is not required for the maintenance of the
sarcomeres in adult muscles (Figure 3A-D).

In summary, our current study shows that Styxl2 critically regulates *de novo*
sarcomere assembly by binding to NM IIs, enhancing their ubiquitination, and promoting
their autophagic degradation, which ultimately facilitates sarcomere maturation *in vivo*
(Figure 7D).

Materials and Methods

Animals

ABSR wild-type zebrafish were maintained following standard procedures and the zygotes were collected for microinjection experiments. $Pax7^{CreERT2/+}$ mice (Stock No: 017763) were purchased from the Jackson Laboratory (Bar Harbor, ME, USA). $Myf5^{Cre/+}$, $Pax7^{Cre/+}$, and Tg $(HSA-MerCreMer)$ mice were generous gifts from Drs. Michael Rudnicki (Ottawa Hospital Research Institute, ON, Canada), Mario Capecchi (University of Utah, UT, USA), and Karyn Esser (University of Florida, FL, USA), respectively. $Styxl2^{flox/flox}$ mice were generated by the Model Animal Research Centre (Nanjing University, Nanjing, China). To delete $Styxl2$ in adult mice, tamoxifen dissolved in corn oil (15 mg/ml) was injected into adult mice intraperitoneally at 5 µl per gram of body weight for 5 consecutive days followed by another 5 doses of tamoxifen injection every 3-4 days. All animals were maintained and handled following the protocols approved by the Animal Ethics Committee of the Hong Kong University of Science and Technology.

Cell culture and DNA constructs

C2C12 cells and HEK 293T cells were purchased from ATCC (Manassas, VA, USA). C2C12 cells were maintained in growth medium (GM) (Dulbecco's modified Eagle's medium supplemented with 20% fetal bovine serum, 100 Units/ml penicillin, and 100 µg/ml streptomycin) and differentiation was induced with differentiation medium (DM) (Dulbecco's modified Eagle's medium supplemented with 2% horse serum, 100 Units/ml penicillin, and 100 µg/ml streptomycin). HEK 293T cells were maintained in Dulbecco's modified Eagle's medium supplemented with 10% fetal bovine serum, 100 Units/ml penicillin, and 100 µg/ml streptomycin. Mouse Styxl2, Styxl2ΔN509 and Styxl2N513 were amplified from a mouse muscle cDNA library and inserted into a modified pcDNA3.0 vector containing a Flag-expressing cassette to generate constructs encoding Flag-Styxl2, Flag-Styxl2ΔN509 and Flag-Styxl2N513. BirA* was amplified from pcDNA3.1 mycBioID (addgene, #35700) and an HA tag was added to its C-terminus. The cDNA encoding BirA*-HA was fused in-frame with that encoding mouse Styxl2 and cloned into the pRetroX-Tight-Pur vector (kindly provided by Dr. Yusong Guo, the Hong Kong University of Science and Technology) to get pTight-Styxl2-BirA*-HA. Mouse Myh9, Myh9-head and Myh9-tail were amplified from a plasmid encoding Myosin-IIA-GFP (addgene, #38297) and inserted into a modified pcDNA3.0 vector containing an HA-expressing cassette to generate constructs

encoding HA-Myh9, HA-Myh9-head and HA-Myh9-tail. Mouse Myh10-head and Myh10-

tail were amplified from a mouse C2C12 cell cDNA library and inserted into a modified

pcDNA3.0 vector containing an HA-expressing cassette to generate constructs encoding HA-

Myh10-head and HA-Myh10-tail. Fish Styxl2 and Styxl2ΔN493 were amplified from a

zebrafish cDNA library and inserted into either pCS2+ vector or pCS2+-GFP vector to

generate pCS2+-zStyxl2 and pCS2+-zStyxl2ΔN493 or pCS2+-GFP-zStyxl2. And the

construct coding Myc-Ub was kindly provided by Dr. Shengcai Lin (Xiamen University).

RNA extraction and RT-qPCR

Total RNA was extracted from C2C12 cells using the TRIzol Reagent (Invitrogen) following

the manufacturer's instructions. The reverse transcription was performed using an oligo-dT

primer and the ImRrom-II Reverse Transcription System (Promega). The ABI 7700 Sequence

Detection System (PE Applied-Biosystems) was used for qPCR and the relative fold change

of genes of interest was analysed based on the $2^{-\Delta\Delta Ct}$ method (Livak & Schmittgen, 2001).

The qPCR primers used were listed below:

Styxl2-F: 5'- GCCCATCCACCTCTCCTC-3'

Styxl2-R: 5'- GCTCCTGTCATCCATCTTCTC-3'

Myh9-F: 5'-GGCCCTGCTAGATGAGGAGT-3'

Myh9-R: 5'-CTTGGGCTTCTGGAACTTGG-3'

Myh10-F: 5'-GGAATCCTTTGGAAATGCGAAGA-3'

Myh10-R: 5'-GCCCCAACAATATAGCCAGTTAC-3'

Gapdh-F: 5'-CCCACTCTTCCACCTTCG-3'

Gapdh-R: 5'-TCCTTGGAGGCCATGTAG-3'

siRNA transfection

The following siRNAs were used to knock down specific genes. They were transfected into

C2C12 cells using the Lipofectamine RNAiMax reagent (Thermo Fisher Scientific) following

manufacturer's instructions.

GFP (negative control): 5'-GCTGACCCTGAAGTTCATC-3'

Jak1: 5'-GCCTGAGAGTGGAGGTAAC-3'

Stat1: 5'-GGATCAAGTCATGTGCATA-3'

ATG5: 5'-GGCTCCTGGATTATGTCAT-3'

Microarray

Total mRNA were extracted from duplicated samples and subjected to microarray analysis
using the Affymetrix GeneChip® Mouse Genome 430 2.0 Array. The data were analysed by
the GeneSpring software.

Plasmid transfection
The Lipofectamine Transfection Reagent (Thermo Fisher Scientific) and the PLUS Reagent
(Thermo Fisher Scientific) were used to transfect plasmids into C2C12 myoblasts following
manufacturer's instructions. Polyethylenimine (PEI, Polysciences) was used to transfect
plasmids into HEK 293T cells.

Western blot
For cultured cells, Western blot was performed according to the standard procedures. For
mouse tissues, the samples were homogenized using a T-25 digital IKA®ULTRA-TURRAX
disperser in a cell lysis buffer (25 mM Tris-HCl pH7.4, 150 mM NaCl, 1 mM EDTA, 1%
NP-40, 5% glycerol) with a cocktail of protease inhibitors (20 mM p-nitrophenylphosphate,
20 mM β-glycerolphosphate, 0.5 mM phenylmethylsulfonyl fluoride, 2 μg/ml aprotinin, 0.5
μg/ml leupeptin, 0.7 μg/ml pepstatin, 50 μM sodium vanadate). Then the soluble lysates were
used for Western blot. The primary antibodies used include anti-Jak1 (Upstate, 06-272), anti-
Stat1 (Upstate, 06-501), anti-β-Tubulin (Sigma, T4026), anti-sarcomere myosin heavy chain
(MHC) (DSHB, MF20), anti-GAPDH (Ambion, AM4300), anti-Myh9 (Cell Signaling, 3403),
anti-Myh10 (Santa Cruz, sc-376942), anti-Flag (Sigma, F3165), and anti-HA (Roche,
12CA5). The Styxl2 antiserum was generated in-house by injecting rabbits with recombinant
proteins spanning aa 2-142 of Styxl2 fused in-frame with glutathione S-transferase. The
antiserum was purified with protein-G beads. The streptavidin-HRP was purchased from GE
Healthcare (#RPN1231).

Immunostaining
C2C12 myoblasts were fixed with 4% paraformaldehyde (PFA) for 10 min at room
temperature, and then washed with 0.1% PBST. Then samples were permeabilized with 0.5%
PBST for 20 min. Zebrafishes were fixed with PFA for 2-4 h at room temperature and
washed with 0.1% PBST. Then the fishes were permeabilized with cold acetone for 8 min at -
20°C. The following procedures are the same for two types of samples. After
permeabilization, the samples were washed with 0.1% PBST and blocked with 4% IgG-free

BSA for 1 h and then incubated with the primary antibody at 4°C overnight. Then samples

were washed with 0.1% PBST and incubated with the secondary antibody at room

temperature for 1 h. The DNA was stained with 100 ng/ml 4′,6‐diamidino‐2‐phenylindole

(DAPI). For fish slow muscles, the nucleus was stained with an anti-Prox1 antibody. The

primary antibodies used include anti-Flag (Sigma, F7425), anti-HA (Santa Cruz, sc-805),

anti-α-Actinin (Sigma, A7732), anti-Prox1 (Angiobio, 11-002P), anti-slow myosin heavy

chain (DSHB, F59), anti-myosin heavy chain (DSHB, A4.1025), and anti-sarcomere myosin

heavy chain (MHC) (DSHB, MF20).

Microinjection of zebrafish zygotes

The zygotes of zebrafish were collected in the morning. 2 nl of either plasmid DNA- or

morpholino-containing solution was injected into fish zygotes as report before (Jin et al.,

2009). The concentration of pCS2+-GFP-zStyxl2 was 25 ng/µl. The concentration of

morpholinos used for microinjection was 0.3 ~ 0.5 mM. A control morpholino targeting a

non-related gene was used as negative control (Ctrl-MO). Morpholinos used were

synthesized by Gene Tools (USA) and their sequences were listed below:

Ctrl-MO: 5'-AGCACACAAAGGCGAAGGTCAACAT-3'

Styxl2-MO: 5'-GCTGATCCTCCACAGACGACGCCAT-3'

Myh10-MO: 5'-CTTCACAAATGTGGTCTTACCTTGA-3'

Atg5-MO:5'-CACATCCTTGTCATCTGCCATTATC-3'

The DNA constructs of pCS2+-zStyxl2 (with several synonymous mutations near the start

codon of zStyxl2 to make it MO-resistant) and pCS2+-zStyxl2ΔN493 were linearized and

used as templates for *in vitro* transcription. The mRNA was synthesized using the

mMESSAGE mMACHINE® SP6 Kit (Ambion, AM1340) followed by purification with the

MEGAclear™ Transcription Clean-Up Kit (Invitrogen, AM1908). The purified mRNA was

dissolved in RNase-free water and diluted to 150 ng/µl immediately before microinjection.

Transmission Electron Microscopy

The heart and skeletal muscles were processed following the published protocols (Graham &

Orenstein, 2007). Then the samples were examined by a Hitachi H7650 transmission electron

microscope (Hitachi High-Technologies). The sarcomere length was measured from electron

microscopic images with a magnification of 50,000 X. The length of 3-5 sarcomeres was

measured for each mouse and a total of six mice were examined. The results were presented

as mean ± SD. The unpaired Student's t-test was performed and p < 0.05 was considered

statistically significant.

Measurement of muscle force generation *In vitro*

The gastrocnemius muscle of left hindlimb was carefully dissected from mice together with

femur condyle and Achilles tendon. The twitch and tetanic force generated by the dissected

gastrocnemius muscle and the muscle cross-sectional area (MCSA) were measured following

published protocols (Guo et al., 2015). The force generated was recorded by the Dynamic

Muscle Control system (DMC v5.4; Aurora Scientific) and analysed by the Dynamic Muscle

Analysis system (DMA v3.2; Aurora Scientific). Both twitch and tetanic force were then

normalized to the MCSA.

BioID and mass spectrometry

C2C12 myoblasts were infected by a retrovirus carrying pRetroX-Tet-On Advanced (kindly

provided by Dr. Yusong Guo, the Hong Kong University of Science and Technology) and

selected by 1 mg/ml G418 to get the Tet-on cells. Then the Tet-on cells were infected by a

retrovirus carrying pTight-Styxl2-BirA*-HA and selected by 1.5 μg/ml puromycin to obtain

stable Tet-Styxl2-BirA*-HA cells. Thus, the expression of Styxl2-BirA*-HA is controlled by

the Tet-on system in these stable C2C12 myoblasts. These stable cells were seeded in GM

and cultured for 24 h before being induced to differentiate in DM for 48 h. The medium was

then replaced with DM containing 50 μM biotin (control) or 50 μM biotin plus 1 μg/ml

doxycycline. After another 24 h, the control and doxycycline-treated cells were processed for

mass spectrometry analysis following the published protocols (Roux et al., 2018). Briefly, the

biotinylated proteins were enriched by Dynabeads MyOne Streptavidin C1 (Invitrogen,

#65001). The proteins bound to the beads were reduced with 10 mM Tris(2-

carboxyethyl)phosphine hydrochloride (Sigma, C4706) at 40°C for 30 min and alkylated with

20 mM Iodoacetamide (Sigma, I1149) in the dark for 30 min to prevent re-formation of the

disulfide bond. The samples were digested with Trypsin Gold (Promega, V5280) overnight at

37°C on a Thermomixer at 700 rpm followed by centrifugation. The supernatant was

transferred to a new tube, acidified to pH2-3 by trifluoroacetic acid (Thermofisher, 85183),

and desalted with a Pierce C18 spin column (Thermo Fisher Scientific, 89873). The tryptic

peptides were labelled with iTRAQ reagents (AB Sciex, 4390812) following the

manufacturer's instructions. Briefly, peptides from control cells were labelled with iTRAQ

tag 114 and those from doxycycline-treated cells were labelled with tag 115. The labelled

samples were mixed and fractionated by an UltiMate 3000 RSLCnano system (Thermo

Fisher Scientific). A C18 Acclaim PepMap RSLC analytical column (75 μm × 250 mm, 2 μm,

100Å) with a C18 nano Viper trap-column (0.3 mm × 5 mm, Thermo Fisher Scientific) was

used for peptide elution and separation at a flow rate of 300 nl/min. The mobile phase

contained buffer A (0.1% formic acid, 2% acetonitrile) and B (0.1% formic acid, 98%

acetonitrile). The gradient was set as follows: 0 min, 3% B; 10 min, 3% B; 12 min, 7% B; 62

min, 20% B; 64 min, 30% B; 65 min, 80% B; 73 min, 80% B; 73.1 min, 3% B; and 90 min, 3%

B. MS data were then acquired using an Orbitrap Fusion Lumos mass spectrometer (Thermo

Fisher Scientific) in a positive mode, with the following settings: 1) MS1 scan: 400 to 1,600

m/z; resolution: 120,000; automatic gain control (AGC): 400,000; and maximum injection

time: 50 ms; 2) The collision energy was set at 35% and orbitrap was used for MS2 scan as

well; 3) MS2 scan starts from 100 m/z; resolution: 30,000; AGC: 50,000; and maximum

injection time: 250 ms; 4) Exclusion window was set for 40 s; 5) The intensity threshold was

set at 25,000. Data analysis was carried out using Mascot Deamon version 2.5.0 (Matrix

Science) with standard settings for each instrument type and the data were searched against

the *Mus musculus* database downloaded from SwissProt. A peptide tolerance of 10 ppm and

fragment ion tolerance of 20 ppm were used. Carbamidomethylation of cysteine was

specified as fixed modification, while oxidation of methionine and acetylation of protein N

termini were set as variable modifications. The false discovery rate was set at 0.01 for

proteins and peptide spectrum match. The 115/114 iTRAQ ratio of one protein was calculated

based on the peak intensity of reporter ions by the software, and defined as Enrichment score

for proteins in doxycycline-treated cells over that in control cells. The *p* values for each

protein generated by the software were used for gene ontology analysis.

708 Gene ontology enrichment analysis

709   All the biotinylated proteins enriched (with Enrichment score > 1) in cells expressing Styxl2-

710   BirA*-HA were imported and analysed at Geneontology (http://geneontology.org/). The

711   terms in the cellular component category were presented.

713 Immunoprecipitation

714   C2C12 cells or HEK293T cells were lysed in the lysis buffer with a cocktail of protease

715   inhibitors. The soluble whole cell lysates were first immunoprecipitated with an antibody

716   targeting Styxl2, MEF2A (negative control), or Flag (Roche, F3165). The proteins co-

precipitated were analysed by Western blot. In the ubiquitination assays, HEK 293T cells were lysed in the RIPA buffer (50 mM Tris-HCl, pH 7.4, 150 mM NaCl, 1 mM EDTA, 1% Triton X-100, 0.5% sodium deoxycholate, 0.1% SDS) with a cocktail of protease inhibitors and the soluble whole cell lysates were immunoprecipitated with an anti-Myc antibody (Santa Cruz, sc-40). The immunoprecipitated proteins were then analysed by Western blot.

Injury-induced muscle regeneration

Avertin was used to anaesthetize mice at 0.5 mg per gram of body weight by intraperitoneal injection. Then 120 µl (for females) or 150 µl (for males) of 10 µM Cardiotoxin was injected into muscles of each hind limb to induce acute injury. After >30 days, the muscles were dissected and tested. To examine protein levels at different time points, each tibialis anterior muscle was injected with 30 µl of 1.2% $BaCl_2$ to induce acute damage. The regenerating muscles were collected for Western blot at different time points.

Acknowledgements

The project was supported by research grants from the Hong Kong Research Grant Council (C6018-19G, AoE/M-604/16, T13-602/21-N, and T13-605/18-W), the State Key Laboratory of Molecular Neuroscience at HKUST, the Hong Kong Center for Neurodegenerative Diseases (HKCeND), and the Shenzhen Bay Laboratory (S201101002, Guangdong, China). We thank Drs. K Esser, M Rudnicki, and M Capecchi for providing mouse Cre lines used in this study. We also thank Dr. Y Guo for providing the BioID vector and Drs. Y Dai and S Zhao for their help with zebrafish work.

Competing interests

The authors declare no competing interests.

References:

Alonso, A., Sasin, J., Bottini, N., Friedberg, I., Friedberg, I., Osterman, A., Godzik, A., Hunter, T., Dixon, J., & Mustelin, T. (2004). Protein tyrosine phosphatases in the human genome. *Cell*, *117*(6), 699–711. https://doi.org/10.1016/j.cell.2004.05.018

Arndt, V., Dick, N., Tawo, R., Dreiseidler, M., Wenzel, D., Hesse, M., Fürst, D. O., Saftig, P., Saint, R., Fleischmann, B. K., Hoch, M., & Höhfeld, J. (2010). Chaperone-Assisted Selective Autophagy Is Essential for Muscle Maintenance. *Current Biology*, *20*(2), 143–148. https://doi.org/10.1016/j.cub.2009.11.022

Arrington, C. B., Bleyl, S. B., Matsunami, N., Bonnell, G. D., Otterud, B. E. M., Nielsen, D. C., Stevens, J., Levy, S., Leppert, M. F., & Bowles, N. E. (2012). Exome analysis of a family with pleiotropic congenital heart disease. *Circulation. Cardiovascular Genetics*, *5*(2), 175–182. https://doi.org/10.1161/CIRCGENETICS.111.961797

Avellaneda, J., Rodier, C., Daian, F., Brouilly, N., Rival, T., Luis, N. M., & Schnorrer, F. (2021). Myofibril and mitochondria morphogenesis are coordinated by a mechanical feedback mechanism in muscle. *Nature Communications*, *12*(1), Article 1. https://doi.org/10.1038/s41467-021-22058-7

Azzedine, H., Bolino, A., Taïeb, T., Birouk, N., Di Duca, M., Bouhouche, A., Benamou, S., Mrabet, A., Hammadouche, T., Chkili, T., Gouider, R., Ravazzolo, R., Brice, A., Laporte, J., & LeGuern, E. (2003). Mutations in MTMR13, a new pseudophosphatase homologue of MTMR2 and Sbf1, in two families with an autosomal recessive demyelinating form of Charcot-Marie-Tooth disease associated with early-onset glaucoma. *American Journal of Human Genetics*, *72*(5), 1141–1153. https://doi.org/10.1086/375034

Bellec, Y., Harrar, Y., Butaeye, C., Darnet, S., Bellini, C., & Faure, J.-D. (2002). Pasticcino2 is a protein tyrosine phosphatase-like involved in cell proliferation and differentiation

in Arabidopsis. *The Plant Journal*, *32*(5), 713–722. https://doi.org/10.1046/j.1365-

313X.2002.01456.x

Berger, P., Berger, I., Schaffitzel, C., Tersar, K., Volkmer, B., & Suter, U. (2006). Multi-level

regulation of myotubularin-related protein-2 phosphatase activity by myotubularin-

related protein-13/set-binding factor-2. *Human Molecular Genetics*, *15*(4), 569–579.

https://doi.org/10.1093/hmg/ddi473

Bloor, J. W., & Kiehart, D. P. (2001). Zipper Nonmuscle Myosin-II Functions Downstream

of PS2 Integrin in Drosophila Myogenesis and Is Necessary for Myofibril Formation.

Developmental Biology, *239*(2), 215–228. https://doi.org/10.1006/dbio.2001.0452

Bogomolovas, J., Fleming, J. R., Franke, B., Manso, B., Simon, B., Gasch, A., Markovic, M.,

Brunner, T., Knöll, R., Chen, J., Labeit, S., Scheffner, M., Peter, C., & Mayans, O.

(2021). Titin kinase ubiquitination aligns autophagy receptors with mechanical signals

in the sarcomere. *EMBO Reports*, *22*(10), e48018.

https://doi.org/10.15252/embr.201948018

Bönnemann, C. G., & Laing, N. G. (2004). Myopathies resulting from mutations in

sarcomeric proteins. *Current Opinion in Neurology*, *17*(5), 529–537.

Chen, M. J., Dixon, J. E., & Manning, G. (2017). Genomics and evolution of protein

phosphatases. *Science Signaling*, *10*(474), eaag1796.

https://doi.org/10.1126/scisignal.aag1796

Chen, R.-H., Chen, Y.-H., & Huang, T.-Y. (2019). Ubiquitin-mediated regulation of

autophagy. *Journal of Biomedical Science*, *26*(1), 80. https://doi.org/10.1186/s12929-

019-0569-y

Cheng, K. C.-C., Klancer, R., Singson, A., & Seydoux, G. (2009). Regulation of MBK-

2/DYRK by CDK-1 and the Pseudophosphatases EGG-4 and EGG-5 during the

Oocyte-to-Embryo Transition. *Cell*, *139*(3), 560–572.

https://doi.org/10.1016/j.cell.2009.08.047

Chopra, A., Kutys, M. L., Zhang, K., Polacheck, W. J., Sheng, C. C., Luu, R. J., Eyckmans, J.,

Hinson, J. T., Seidman, J. G., Seidman, C. E., & Chen, C. S. (2018). Force Generation

via β-Cardiac Myosin, Titin, and α-Actinin Drives Cardiac Sarcomere Assembly from

Cell-Matrix Adhesions. *Developmental Cell*, *44*(1), 87-96.e5.

https://doi.org/10.1016/j.devcel.2017.12.012

Clark, K. A., McElhinny, A. S., Beckerle, M. C., & Gregorio, C. C. (2002). Striated Muscle

Cytoarchitecture: An Intricate Web of Form and Function. *Annual Review of Cell and*

Developmental Biology, *18*(1), 637–706.

https://doi.org/10.1146/annurev.cellbio.18.012502.105840

Conway, S. J., Kruzynska□Frejtag, A., Kneer, P. L., Machnicki, M., & Koushik, S. V. (2003).

What cardiovascular defect does my prenatal mouse mutant have, and why? *Genesis*,

35(1), 1–21. https://doi.org/10.1002/gene.10152

Cooper, L. M., & Waddell, D. S. (2020). A tale of two DUSP27s: Proposed resolution for the

naming of distinct dual-specificity phosphatases. *American Journal of Physiology-*

Cell Physiology, *319*(1), C148–C150. https://doi.org/10.1152/ajpcell.00201.2020

Craig, R., & Padrón, R. (2004). Chapter 7 Molecular Structure of the Sarcomere. In *Myology*

(pp. 129–166).

Da Costa, M., Bach, L., Landrieu, I., Bellec, Y., Catrice, O., Brown, S., De Veylder, L.,

Lippens, G., Inzé, D., & Faure, J.-D. (2006). Arabidopsis PASTICCINO2 Is an

Antiphosphatase Involved in Regulation of Cyclin-Dependent Kinase A. *The Plant*

Cell, *18*(6), 1426–1437. https://doi.org/10.1105/tpc.105.040485

Das, A. T., Tenenbaum, L., & Berkhout, B. (2016). Tet-On Systems For Doxycycline-

inducible Gene Expression. *Current Gene Therapy*, *16*(3), 156–167.

https://doi.org/10.2174/1566523216666160524144041

Dlugosz, A. A., Antin, P. B., Nachmias, V. T., & Holtzer, H. (1984). The relationship

between stress fiber-like structures and nascent myofibrils in cultured cardiac

myocytes. *Journal of Cell Biology*, *99*(6), 2268–2278.

https://doi.org/10.1083/jcb.99.6.2268

Doyotte, A., Mironov, A., McKenzie, E., & Woodman, P. (2008). The Bro1-related protein

HD-PTP/PTPN23 is required for endosomal cargo sorting and multivesicular body

morphogenesis. *Proceedings of the National Academy of Sciences of the United States*

of America, *105*(17), 6308–6313. https://doi.org/10.1073/pnas.0707601105

Dumont, N., Bentzinger, C., Sincennes, M.-C., & Rudnicki, M. (2015). Satellite Cells and

Skeletal Muscle Regeneration. *Comprehensive Physiology*, *5*, 1027–1059.

https://doi.org/10.1002/cphy.c140068

Fenix, A. M., Neininger, A. C., Taneja, N., Hyde, K., Visetsouk, M. R., Garde, R. J., Liu, B.,

Nixon, B. R., Manalo, A. E., Becker, J. R., Crawley, S. W., Bader, D. M., Tyska, M.

J., Liu, Q., Gutzman, J. H., & Burnette, D. T. (2018). Muscle-specific stress fibers

give rise to sarcomeres in cardiomyocytes. *eLife*, *7*, e42144.

https://doi.org/10.7554/eLife.42144

Fero, K., Bergeron, S. A., Horstick, E. J., Codore, H., Li, G. H., Ono, F., Dowling, J. J., &

Burgess, H. A. (2014). Impaired embryonic motility in dusp27 mutants reveals a

developmental defect in myofibril structure. *Disease Models & Mechanisms*, *7*(2),

289–298. https://doi.org/10.1242/dmm.013235

Firestein, R., Nagy, P. L., Daly, M., Huie, P., Conti, M., & Cleary, M. L. (2002). Male

infertility, impaired spermatogenesis, and azoospermia in mice deficient for the

840   pseudophosphatase Sbf1. *The Journal of Clinical Investigation*, *109*(9), 1165–1172.

841   https://doi.org/10.1172/JCI12589

842 Friedberg, I., Nika, K., Tautz, L., Saito, K., Cerignoli, F., Friedberg, I., Godzik, A., &

843   Mustelin, T. (2007). Identification and characterization of DUSP27, a novel dual-

844   specific protein phosphatase. *FEBS Letters*, *581*(13), 2527–2533.

845   https://doi.org/10.1016/j.febslet.2007.04.059

846 Graham, L., & Orenstein, J. M. (2007). Processing tissue and cells for transmission electron

847   microscopy in diagnostic pathology and research. *Nature Protocols*, *2*(10), 2439–

848   2450. https://doi.org/10.1038/nprot.2007.304

849 Guo, A., Leung, K., Qin, J., Chow, S. K., & Cheung, W. (2015). Effect of Low-Magnitude,

850   High-Frequency Vibration Treatment on Retardation of Sarcopenia: Senescence-

851   Accelerated Mouse-P8 Model. *Rejuvenation Research*, *19*(4), 293–302.

852   https://doi.org/10.1089/rej.2015.1759

853 Gutzman, J. H., Sahu, S. U., & Kwas, C. (2015). Non-muscle myosin IIA and IIB

854   differentially regulate cell shape changes during zebrafish brain morphogenesis.

855   *Developmental Biology*, *397*(1), 103–115. https://doi.org/10.1016/j.ydbio.2014.10.017

856 Hasty, P., Bradley, A., Morris, J. H., Edmondson, D. G., Venuti, J. M., Olson, E. N., & Klein,

857   W. H. (1993). Muscle deficiency and neonatal death in mice with a targeted mutation

858   in the myogenin gene. *Nature*, *364*(6437), Article 6437.

859   https://doi.org/10.1038/364501a0

860 Hinton, S. D. (2019). The role of pseudophosphatases as signaling regulators. *Biochimica et*

861   *Biophysica Acta (BBA) - Molecular Cell Research*, *1866*(1), 167–174.

862   https://doi.org/10.1016/j.bbamcr.2018.07.021

Hinton, S. D. (2020). Pseudophosphatase MK-STYX: The atypical member of the MAP

kinase phosphatases. *The FEBS Journal*, *287*(19), 4221–4231.

https://doi.org/10.1111/febs.15426

Hinton, S. D., Myers, M. P., Roggero, V. R., Allison, L. A., & Tonks, N. K. (2010). The

pseudophosphatase MK-STYX interacts with G3BP and decreases stress granule

formation. *Biochemical Journal*, *427*(3), 349–357.

https://doi.org/10.1042/BJ20091383

Holtzer, H., Hijikata, T., Lin, Z. X., Zhang, Z. Q., Holtzer, S., Protasi, F., Franzini-Armstrong,

C., & Sweeney, H. L. (1997). Independent assembly of 1.6 microns long bipolar

MHC filaments and I-Z-I bodies. *Cell Structure and Function*, *22*(1), 83–93.

https://doi.org/10.1247/csf.22.83

Jin, H., Sood, R., Xu, J., Zhen, F., English, M. A., Liu, P. P., & Wen, Z. (2009). Definitive

hematopoietic stem/progenitor cells manifest distinct differentiation output in the

zebrafish VDA and PBI. *Development*, *136*(4), 647–654.

https://doi.org/10.1242/dev.029637

Keller, C., Hansen, M. S., Coffin, C. M., & Capecchi, M. R. (2004). Pax3:Fkhr interferes

with embryonic Pax3 and Pax7 function: Implications for alveolar

rhabdomyosarcoma cell of origin. *Genes & Development*, *18*(21), 2608–2613.

https://doi.org/10.1101/gad.1243904

Kharitidi, D., Apaja, P. M., Manteghi, S., Suzuki, K., Malitskaya, E., Roldan, A., Gingras,

M.-C., Takagi, J., Lukacs, G. L., & Pause, A. (2015). Interplay of Endosomal pH and

Ligand Occupancy in Integrin α5β1 Ubiquitination, Endocytic Sorting, and Cell

Migration. *Cell Reports*, *13*(3), 599–609. https://doi.org/10.1016/j.celrep.2015.09.024

Kharitidi, D., Manteghi, S., & Pause, A. (2014). Pseudophosphatases: Methods of analysis

and physiological functions. *Methods*, *65*(2), 207–218.

https://doi.org/10.1016/j.ymeth.2013.09.009

Kirkin, V., McEwan, D. G., Novak, I., & Dikic, I. (2009). A Role for Ubiquitin in Selective

Autophagy. *Molecular Cell*, *34*(3), 259–269.

https://doi.org/10.1016/j.molcel.2009.04.026

Lakso, M., Pichel, J. G., Gorman, J. R., Sauer, B., Okamoto, Y., Lee, E., Alt, F. W., &

Westphal, H. (1996). Efficient in vivo manipulation of mouse genomic sequences at

the zygote stage. *Proceedings of the National Academy of Sciences of the United*

States of America, *93*(12), 5860–5865. https://doi.org/10.1073/pnas.93.12.5860

Livak, K. J., & Schmittgen, T. D. (2001). Analysis of relative gene expression data using

real-time quantitative PCR and the 2(-Delta Delta C(T)) Method. *Methods (San Diego,*

Calif.), *25*(4), 402–408. https://doi.org/10.1006/meth.2001.1262

Loison, O., Weitkunat, M., Kaya-Çopur, A., Alves, C. N., Matzat, T., Spletter, M. L.,

Luschnig, S., Brasselet, S., Lenne, P.-F., & Schnorrer, F. (2018). Polarization-

resolved microscopy reveals a muscle myosin motor-independent mechanism of

molecular actin ordering during sarcomere maturation. *PLOS Biology*, *16*(4),

e2004718. https://doi.org/10.1371/journal.pbio.2004718

Lu, M. H., DiLullo, C., Schultheiss, T., Holtzer, S., Murray, J. M., Choi, J., Fischman, D. A.,

& Holtzer, H. (1992). The vinculin/sarcomeric-alpha-actinin/alpha-actin nexus in

cultured cardiac myocytes. *The Journal of Cell Biology*, *117*(5), 1007–1022.

https://doi.org/10.1083/jcb.117.5.1007

Luis, N. M., & Schnorrer, F. (2021). Mechanobiology of muscle and myofibril

morphogenesis. *Cells & Development*, *168*, 203760.

https://doi.org/10.1016/j.cdev.2021.203760

Ma, H., Wardega, P., Mazaud, D., Klosowska-Wardega, A., Jurek, A., Engström, U.,

Lennartsson, J., & Heldin, C.-H. (2015). Histidine-domain-containing protein tyrosine

phosphatase regulates platelet-derived growth factor receptor intracellular sorting and

degradation. *Cellular Signalling*, *27*(11), 2209–2219.

https://doi.org/10.1016/j.cellsig.2015.07.020

Ma, X., Takeda, K., Singh, A., Yu, Z.-X., Zerfas, P., Blount, A., Liu, C., Towbin, J. A.,

Schneider, M. D., Adelstein, R. S., & Wei, Q. (2009). Conditional Ablation of

Nonmuscle Myosin II-B Delineates Heart Defects in Adult Mice. *Circulation*

Research, *105*(11), 1102–1109. https://doi.org/10.1161/CIRCRESAHA.109.200303

Mao, Q., Acharya, A., Rodríguez-delaRosa, A., Marchiano, F., Dehapiot, B., Al Tanoury, Z.,

Rao, J., Díaz-Cuadros, M., Mansur, A., Wagner, E., Chardes, C., Gupta, V., Lenne,

P.-F., Habermann, B. H., Theodoly, O., Pourquié, O., & Schnorrer, F. (2022).

Tension-driven multi-scale self-organisation in human iPSC-derived muscle fibers.

eLife, *11*, e76649. https://doi.org/10.7554/eLife.76649

Martin, T. G., & Kirk, J. A. (2020). Under construction: The dynamic assembly, maintenance,

and degradation of the cardiac sarcomere. *Journal of Molecular and Cellular*

Cardiology, *148*, 89–102. https://doi.org/10.1016/j.yjmcc.2020.08.018

Maruyama, R., Velarde, N. V., Klancer, R., Gordon, S., Kadandale, P., Parry, J. M., Hang, J.

S., Rubin, J., Stewart-Michaelis, A., Schweinsberg, P., Grant, B. D., Piano, F.,

Sugimoto, A., & Singson, A. (2007). EGG-3 Regulates Cell-Surface and Cortex

Rearrangements during Egg Activation in Caenorhabditis elegans. *Current Biology*,

17(18), 1555–1560. https://doi.org/10.1016/j.cub.2007.08.011

Masiero, E., Agatea, L., Mammucari, C., Blaauw, B., Loro, E., Komatsu, M., Metzger, D.,

Reggiani, C., Schiaffino, S., & Sandri, M. (2009). Autophagy Is Required to Maintain

935   Muscle Mass. *Cell Metabolism*, *10*(6), 507–515.

936   https://doi.org/10.1016/j.cmet.2009.10.008

937 McCarthy, J. J., Srikuea, R., Kirby, T. J., Peterson, C. A., & Esser, K. A. (2012). Inducible

938   Cre transgenic mouse strain for skeletal muscle-specific gene targeting. *Skeletal*

939   *Muscle*, *2*(1), 8. https://doi.org/10.1186/2044-5040-2-8

940 Mizushima, N., Yoshimori, T., & Levine, B. (2010). Methods in Mammalian Autophagy

941   Research. *Cell*, *140*(3), 313–326. https://doi.org/10.1016/j.cell.2010.01.028

942 Mukund, K., & Subramaniam, S. (2020). Skeletal muscle: A review of molecular structure

943   and function, in health and disease. *Wiley Interdisciplinary Reviews. Systems Biology*

944   *and Medicine*, *12*(1), e1462. https://doi.org/10.1002/wsbm.1462

945 Murphy, M. M., Lawson, J. A., Mathew, S. J., Hutcheson, D. A., & Kardon, G. (2011).

946   Satellite cells, connective tissue fibroblasts and their interactions are crucial for

947   muscle regeneration. *Development*, *138*(17), 3625–3637.

948   https://doi.org/10.1242/dev.064162

949 Nakai, A., Yamaguchi, O., Takeda, T., Higuchi, Y., Hikoso, S., Taniike, M., Omiya, S.,

950   Mizote, I., Matsumura, Y., Asahi, M., Nishida, K., Hori, M., Mizushima, N., & Otsu,

951   K. (2007). The role of autophagy in cardiomyocytes in the basal state and in response

952   to hemodynamic stress. *Nature Medicine*, *13*(5), Article 5.

953   https://doi.org/10.1038/nm1574

954 Nakhro, K., Park, J.-M., Hong, Y. B., Park, J. H., Nam, S. H., Yoon, B. R., Yoo, J. H., Koo,

955   H., Jung, S.-C., Kim, H.-L., Kim, J. Y., Choi, K.-G., Choi, B.-O., & Chung, K. W.

956   (2013). SET binding factor 1 (SBF1) mutation causes Charcot-Marie-Tooth disease

957   type 4B3. *Neurology*, *81*(2), 165–173.

958   https://doi.org/10.1212/WNL.0b013e31829a3421

Nandi, D., Tahiliani, P., Kumar, A., & Chandu, D. (2006). The ubiquitin-proteasome system.

Journal of Biosciences, *31*(1), 137–155. https://doi.org/10.1007/BF02705243

Newell-Litwa, K. A., Horwitz, R., & Lamers, M. L. (2015). Non-muscle myosin II in disease:

Mechanisms and therapeutic opportunities. *Disease Models & Mechanisms*, *8*(12),

1495–1515. https://doi.org/10.1242/dmm.022103

Pankiv, S., Clausen, T. H., Lamark, T., Brech, A., Bruun, J.-A., Outzen, H., Øvervatn, A.,

Bjørkøy, G., & Johansen, T. (2007). P62/SQSTM1 Binds Directly to Atg8/LC3 to

Facilitate Degradation of Ubiquitinated Protein Aggregates by Autophagy*. *Journal

of Biological Chemistry*, *282*(33), 24131–24145.

https://doi.org/10.1074/jbc.M702824200

Parker, F., Baboolal, T. G., & Peckham, M. (2020). Actin Mutations and Their Role in

Disease. *International Journal of Molecular Sciences*, *21*(9), E3371.

https://doi.org/10.3390/ijms21093371

Parker, F., & Peckham, M. (2020). Disease mutations in striated muscle myosins. *Biophysical

Reviews*, *12*(4), 887–894. https://doi.org/10.1007/s12551-020-00721-5

Parkinson, M. D. J., Piper, S. C., Bright, N. A., Evans, J. L., Boname, J. M., Bowers, K.,

Lehner, P. J., & Luzio, J. P. (2015). A non-canonical ESCRT pathway, including

histidine domain phosphotyrosine phosphatase (HD-PTP), is used for down-

regulation of virally ubiquitinated MHC class I. *Biochemical Journal*, *471*(1), 79–88.

https://doi.org/10.1042/BJ20150336

Radhakrishna, U., Albayrak, S., Alpay-Savasan, Z., Zeb, A., Turkoglu, O., Sobolewski, P., &

Bahado-Singh, R. O. (2016). Genome-Wide DNA Methylation Analysis and

Epigenetic Variations Associated with Congenital Aortic Valve Stenosis (AVS).

PLOS ONE, *11*(5), e0154010. https://doi.org/10.1371/journal.pone.0154010

Reiterer, V., Fey, D., Kolch, W., Kholodenko, B. N., & Farhan, H. (2013).

Pseudophosphatase STYX modulates cell-fate decisions and cell migration by

spatiotemporal regulation of ERK1/2. *Proceedings of the National Academy of*

Sciences, *110*(31), E2934–E2943. https://doi.org/10.1073/pnas.1301985110

Reiterer, V., Pawłowski, K., Desrochers, G., Pause, A., Sharpe, H. J., & Farhan, H. (2020).

The dead phosphatases society: A review of the emerging roles of

pseudophosphatases. *The FEBS Journal*, *287*(19), 4198–4220.

https://doi.org/10.1111/febs.15431

Reiterer, V., Pawłowski, K., & Farhan, H. (2017). STYX: A versatile pseudophosphatase.

Biochemical Society Transactions, *45*(2), 449–456.

https://doi.org/10.1042/BST20160279

Rhee, D., Sanger, J. M., & Sanger, J. W. (1994). The premyofibril: Evidence for its role in

myofibrillogenesis. *Cell Motility*, *28*(1), 1–24. https://doi.org/10.1002/cm.970280102

Robinson, F. L., & Dixon, J. E. (2005). The phosphoinositide-3-phosphatase MTMR2

associates with MTMR13, a membrane-associated pseudophosphatase also mutated in

type 4B Charcot-Marie-Tooth disease. *The Journal of Biological Chemistry*, *280*(36),

31699–31707. https://doi.org/10.1074/jbc.M505159200

Roux, K. J., Kim, D. I., Burke, B., & May, D. G. (2018). BioID: A Screen for Protein-Protein

Interactions. *Current Protocols in Protein Science*, *91*(1), 19.23.1-19.23.15.

https://doi.org/10.1002/cpps.51

Roux, K. J., Kim, D. I., Raida, M., & Burke, B. (2012). A promiscuous biotin ligase fusion

protein identifies proximal and interacting proteins in mammalian cells. *Journal of*

Cell Biology, *196*(6), 801–810. https://doi.org/10.1083/jcb.201112098

Rudnicki, M. A., Schnegelsberg, P. N. J., Stead, R. H., Braun, T., Arnold, H.-H., & Jaenisch,

    R. (1993). MyoD or Myf-5 is required for the formation of skeletal muscle. *Cell*,

    75(7), 1351–1359. https://doi.org/10.1016/0092-8674(93)90621-V

Rui, Y., Bai, J., & Perrimon, N. (2010). Sarcomere Formation Occurs by the Assembly of

    Multiple Latent Protein Complexes. *PLOS Genetics*, *6*(11), e1001208.

    https://doi.org/10.1371/journal.pgen.1001208

Sanger, J. W., Kang, S., Siebrands, C. C., Freeman, N., Du, A., Wang, J., Stout, A. L., &

    Sanger, J. M. (2006). How to build a myofibril. *Journal of Muscle Research and Cell*

    Motility, *26*(6–8), 343–354. https://doi.org/10.1007/s10974-005-9016-7

Sanger, J. W., Wang, J., Fan, Y., White, J., Mi-Mi, L., Dube, D. K., Sanger, J. M., & Pruyne,

    D. (2017). Assembly and Maintenance of Myofibrils in Striated Muscle. *Handbook of*

    Experimental Pharmacology, *235*, 39–75. https://doi.org/10.1007/164_2016_53

Sanger, J. W., Wang, J., Holloway, B., Du, A., & Sanger, J. M. (2009). Myofibrillogenesis in

    skeletal muscle cells in zebrafish. *Cell Motility*, *66*(8), 556–566.

    https://doi.org/10.1002/cm.20365

Schwander, M., Leu, M., Stumm, M., Dorchies, O. M., Ruegg, U. T., Schittny, J., & Müller,

    U. (2003). B1 Integrins Regulate Myoblast Fusion and Sarcomere Assembly.

    Developmental Cell, *4*(5), 673–685. https://doi.org/10.1016/S1534-5807(03)00118-7

Shaid, S., Brandts, C. H., Serve, H., & Dikic, I. (2013). Ubiquitination and selective

    autophagy. *Cell Death & Differentiation*, *20*(1), Article 1.

    https://doi.org/10.1038/cdd.2012.72

Sparrow, J. C., & Schöck, F. (2009). The initial steps of myofibril assembly: Integrins pave

    the way. *Nature Reviews Molecular Cell Biology*, *10*(4), 293–298.

    https://doi.org/10.1038/nrm2634

Squire, J. M., Al□khayat, H. A., Knupp, C., & Luther, P. K. (2005). Molecular Architecture

in Muscle Contractile Assemblies. In *Advances in Protein Chemistry* (Vol. 71, pp.

17–87). Academic Press. https://doi.org/10.1016/S0065-3233(04)71002-5

Sun, L., Ma, K., Wang, H., Xiao, F., Gao, Y., Zhang, W., Wang, K., Gao, X., Ip, N., & Wu, Z.

(2007). JAK1–STAT1–STAT3, a key pathway promoting proliferation and

preventing premature differentiation of myoblasts. *The Journal of Cell Biology*,

179(1), 129–138. https://doi.org/10.1083/jcb.200703184

Tallquist, M. D., Weismann, K. E., Hellström, M., & Soriano, P. (2000). Early myotome

specification regulates PDGFA expression and axial skeleton development.

Development (Cambridge, England), *127*(23), 5059–5070.

https://doi.org/10.1242/dev.127.23.5059

Tomar, V. S., Baral, T. K., Nagavelu, K., & Somasundaram, K. (2019).

Serine/threonine/tyrosine-interacting-like protein 1 (STYXL1), a pseudo phosphatase,

promotes oncogenesis in glioma. *Biochemical and Biophysical Research*

Communications, *515*(1), 241–247. https://doi.org/10.1016/j.bbrc.2019.05.093

Tonks, N. K. (2006). Protein tyrosine phosphatases: From genes, to function, to disease.

Nature Reviews Molecular Cell Biology, *7*(11), Article 11.

https://doi.org/10.1038/nrm2039

Tullio, A. N., Accili, D., Ferrans, V. J., Yu, Z.-X., Takeda, K., Grinberg, A., Westphal, H.,

Preston, Y. A., & Adelstein, R. S. (1997). Nonmuscle myosin II-B is required for

normal development of the mouse heart. *Proceedings of the National Academy of*

Sciences, *94*(23), 12407–12412. https://doi.org/10.1073/pnas.94.23.12407

Wang, J., Fan, Y., Sanger, J. M., & Sanger, J. W. (2018). Nonmuscle myosin II in cardiac

and skeletal muscle cells. *Cytoskeleton*, *75*(8), 339–351.

https://doi.org/10.1002/cm.21454

Weinfeld, M., Mani, R. S., Abdou, I., Aceytuno, R. D., & Glover, J. N. M. (2011). Tidying

up loose ends: The role of polynucleotide kinase/phosphatase in DNA strand break

repair. *Trends in Biochemical Sciences*, *36*(5), 262–271.

https://doi.org/10.1016/j.tibs.2011.01.006

Weitkunat, M., Kaya-Çopur, A., Grill, S. W., & Schnorrer, F. (2014). Tension and Force-

Resistant Attachment Are Essential for Myofibrillogenesis in Drosophila Flight

Muscle. *Current Biology*, *24*(7), 705–716. https://doi.org/10.1016/j.cub.2014.02.032

White, J., Barro, M. V., Makarenkova, H. P., Sanger, J. W., & Sanger, J. M. (2014).

Localization of Sarcomeric Proteins During Myofibril Assembly in Cultured Mouse

Primary Skeletal Myotubes. *The Anatomical Record*, *297*(9), 1571–1584.

https://doi.org/10.1002/ar.22981

White, J., Wang, J., Fan, Y., Dube, D. K., Sanger, J. W., & Sanger, J. M. (2018). Myofibril

Assembly in Cultured Mouse Neonatal Cardiomyocytes. *The Anatomical Record*,

301(12), 2067–2079. https://doi.org/10.1002/ar.23961

Wiese, S., Reidegeld, K. A., Meyer, H. E., & Warscheid, B. (2007). Protein labeling by

iTRAQ: A new tool for quantitative mass spectrometry in proteome research.

PROTEOMICS, *7*(3), 340–350. https://doi.org/10.1002/pmic.200600422

Wishart, M. J., Denu, J. M., Williams, J. A., & Dixon, J. E. (1995). A Single Mutation

Converts a Novel Phosphotyrosine Binding Domain into a Dual-specificity

Phosphatase. *Journal of Biological Chemistry*, *270*(45), 26782–26785.

https://doi.org/10.1074/jbc.270.45.26782

Wishart, M. J., & Dixon, J. E. (2002). The archetype STYX/dead-phosphatase complexes

with a spermatid mRNA-binding protein and is essential for normal sperm production.

Proceedings of the National Academy of Sciences, *99*(4), 2112–2117.

https://doi.org/10.1073/pnas.251686198

Worby, C. A., Gentry, M. S., & Dixon, J. E. (2006). Laforin, a Dual Specificity Phosphatase

That Dephosphorylates Complex Carbohydrates *. *Journal of Biological Chemistry*,

281(41), 30412–30418. https://doi.org/10.1074/jbc.M606117200

Xiao, F., Wang, H., Fu, X., Li, Y., Ma, K., Sun, L., Gao, X., & Wu, Z. (2011). Oncostatin M

inhibits myoblast differentiation and regulates muscle regeneration. *Cell Research*,

21(2), Article 2. https://doi.org/10.1038/cr.2010.144

Yamamoto, Y., Chino, H., Tsukamoto, S., Ode, K. L., Ueda, H. R., & Mizushima, N. (2021).

NEK9 regulates primary cilia formation by acting as a selective autophagy adaptor for

MYH9/myosin IIA. *Nature Communications*, *12*(1), Article 1.

https://doi.org/10.1038/s41467-021-23599-7

Zaffagnini, G., & Martens, S. (2016). Mechanisms of Selective Autophagy. *Journal of*

Molecular Biology, *428*(9, Part A), 1714–1724.

https://doi.org/10.1016/j.jmb.2016.02.004

1093

Figure 1. Styxl2 regulates sarcomere integrity during zebrafish muscle development. (A) A schematic of mouse Styxl2 protein. DSPc (red): dual-specificity phosphatase, catalytic domain. The numbers denote the positions of various amino acids in mouse Styxl2 including Ser (S)-225. (B) The expression of Styxl2 protein in different mouse tissues. 100 µg of soluble mouse tissue lysates were subjected to Western blot analysis. GAPDH: glyceraldehyde-3-phosphate dehydrogenase. SM: skeletal muscles. (C) The expression of Styxl2 protein in C2C12 myoblasts before and after differentiation in culture. MHC: myosin heavy chain. (D) Zebrafish zygotes were either mock-injected or injected with a Styxl2-specific morpholino (MO). Sarcomeres in muscles at 48 hpf were revealed by transmission electron microscopy (TEM). Representative images were shown. Scale bar: 500 nm. (E) Sarcomeres in fast and slow muscles of zebrafish embryos at 24 hpf were revealed by immunostaining using specific antibodies as indicated. Ctrl: control (non-treated); MO: *Styxl2*-morpholino-treated. The nuclei of slow and fast muscle fibers were counter-stained with an anti-Prox1 antibody and DAPI, respectively. Scale bar: 10 µm.

Figure 2. Conditional deletion of mouse *Styxl2* leads to defective sarcomeres in striated muscles. (A) The expression of Styxl2 protein in skeletal muscles of wild-type (+/+), heterozygous (+/−) and *Styxl2* knockout (KO) (−/−) (driven by *EIIA*-Cre) mice. Soluble tissue lysates from limbs of E14.5 mouse embryos were subjected to Western blot analysis. N.S.: non-specific. (B) E14.5 embryos of control (Ctrl) and *EIIA*-Cre; *Styxl2^f/f* KO mice. *: edema was found in some *Styxl2* KO foetuses. (C) The sarcomere structures of cardiomyocytes in the heart of E14.5 embryos from control and *EIIA*-Cre; *Styxl2^f/f* KO mice were revealed by TEM. Scale bar: 1 μm. (D) The expression of Styxl2 protein in skeletal muscles of control (Ctrl) and *Styxl2* CKO P1 mice. (E) The new-born pups (P1) from control (Ctrl) and *Pax7*^Cre/+; *Styxl2^f/f* KO mice (CKO). (F) Lung sections of control and Styxl2 CKO P1 mice. Scale bar: 50 μm. (G) The sarcomere structure of diaphragm muscles from control (Ctrl) and *Styxl2* CKO P1 mice revealed by TEM. Scale bar: 1 μm.

Figure 3. Styxl2 is not required for sarcomere maintenance but involved in *de novo* sarcomere assembly in adult muscles. (A) The schematic to assess the role of Styxl2 in sarcomere maintenance in adult muscles. TMX: tamoxifen. (B) Styxl2 protein levels in Ctrl and MF-iKO mouse gastrocnemius muscles used for force test. The band intensity of Styxl2 was quantified by ImageJ. (C, D) The twitch and tetanic force generated by gastrocnemius muscles of three Ctrl and four MF-iKO mice were measured. In (A-D), Ctrl: *Styxl2^{f/f}* mice; MF-iKO: *Tg: HSA-MerCreMer; Styxl2^{f/f}* mice. (E) The schematic to assess the role of Styxl2 in *de novo* sarcomere assembly in adult muscles. CTX: cardiotoxin. (F) Styxl2 protein levels in regenerated gastrocnemius muscles used for force test. N.S.: non-specific. The band intensity of Styxl2 was quantified. (G, H) Twitch and tetanic force by regenerated gastrocnemius muscles were measured from four pairs of control and SC-iKO mice. (I) TEM images of the muscle used for force generation test. Scale bar: 500 nm. (J) Quantification of sarcomere length in (I). TEM images from three pairs of mice were examined and the length of 3-5 sarcomeres from each mouse was measured. In (E-J), Ctrl: *Pax7^{CreERT2/CreERT2}* mice; SC-iKO: *Pax7^{CreERT2/CreERT2}; Styxl2^{f/f}* mice. n.s.: not significant. *, p-value < 0.05. ***, p-value < 0.001.

Figure 4. Styxl2 binds to non-muscle myosin IIs. (A) The schematic of the fusion protein Styxl2-BirA*-HA. (B) The workflow of sample preparation for BioID. (C) Enriched (with Enrichment score >1) biotinylated proteins in cells expressing Styxl2-BirA*-HA relative to that in control cells were identified by mass spectrometry followed by Gene Ontology (GO) analysis. Myofibril-related terms were shown. (D) C2C12 cells were harvested after 2 days in DM and the soluble whole cell lysates were subjected to immunoprecipitation (IP) using either an anti-MEF2A (negative control) or an anti-Styxl2 antibody. The endogenous Myh9 or Myh10 was detected by Western blot. (E) The schematic showing the domains of non-muscle myosin IIs (NM II). (F) HEK 293T cells were transfected with Flag-Styxl2 (black arrow) together with HA-NM II-head (red arrowheads) or HA-NM II-tail (black arrowheads). The soluble whole cell lysates were subjected to IP with an anti-Flag antibody and the co-immunoprecipitated proteins were detected by Western blot.

Figure 5. Styxl2 protein levels inversely correlates with that of non-muscle myosin IIs. (A) Protein levels of Styxl2 and non-muscle myosin IIs in skeletal muscles of wild-type mice of different ages were determined by Western blot. (B) Protein levels of Styxl2 and non-muscle myosin IIs in skeletal muscles of control and *Styxl2* CKO P1 mice were determined by Western blot. (C) Relative ratios of non-muscle myosin II protein levels in control and *Styxl2* CKO P1 mice were quantified by ImageJ based on the data from four (for Myh9) or five (for Myh10) sets of Western blots. (D) Protein levels of Styxl2 and non-muscle myosin IIs in regenerating tibialis anterior muscles were determined by Western blot. The experiments were done for four times with similar results and a representative gel was shown. The protein levels of NM IIs in 10 dpi muscles revealed by Western blot were quantified in the right panel. Ctrl: *Pax7^{CreERT2/CreERT2}* mice; SC-iKO: *Pax7^{CreERT2/CreERT2}*; *Styxl2^{f/f}* mice; dpi: days post-injury. (E) Zebrafish zygotes were separately injected with the following morpholinos (MO): control (Ctrl-MO), *Styxl2*-MO, and mixed morpholinos targeting both *Styxl2* and *Myh10* (dMO). At 48 hpf, zebrafishes were poked by a fine needle and their responses were recorded. Quantification of responses of zebrafish to needle touch was shown. In each group, at least 8 zebrafishes were tested and the percentage of zebrafishes responding to needle touch was calculated. Each dot represents one group of experiment. (F) Zebrafishes from each condition in (E) were subjected to immunostaining for α-actinin to visualize sarcomere structures. Representative images were shown. The inset at the top right corner of each image shows the actinin intensity along the yellow line. Scale bar: 10 µm. *, p-value < 0.05. **, p-value < 0.01. ****, p-value < 0.0001.

Figure 6. The DSPc domain of Styxl2 is dispensable for degradation of non-muscle myosin IIs. (A) DNA constructs encoding Myh9, Styxl2 or Mst1 (negative control) were co-expressed in C2C12 cells. Soluble whole cell extracts were subjected to Western blot analysis. OE: overexpression. (B) The schematic shows the truncated fragments of Styxl2 protein. The red blocks indicate the DSPc domain of Styxl2. HEK 293T cells were transfected with different plasmids as indicated. 24 h after transfection, cells were harvested and soluble whole cell extracts were subjected to Western blot analysis. The arrow indicates the head domain of Myh9, and the arrowhead indicates Styxl2 missing the N-terminal 509 aa (Styxl2ΔN509). Mst1: negative control. N.S.: non-specific. (C) Zebrafish zygotes were injected with various morpholinos as indicated with or without the co-injected *Styxl2* mRNAs. z*Styxl2*FL: the full-length fish *Styxl2* mRNA; z*Styxl2*ΔN493: the truncated fish *Styxl2* mRNA missing the 5' region encoding the N-terminal 493 aa. Representative α-actinin immunofluorescent images were shown. The number in numerator at the top left corner of each image represents fish embryos showing normal α-actinin staining pattern, while that in denominator represents the total number of embryos examined. The inset at the top right corner of each image shows the actinin intensity along the yellow line. Scale bar: 10 μm.

Figure 7. The Styxl2-mediated degradation of non-muscle myosin IIs is autophagy dependent. (A) Plasmids encoding Myh9, Styxl2 or Mst1 (negative control) were co-expressed in C2C12 cells together with various siRNAs as indicated. Soluble whole cell extracts were subjected to Western blot analysis. OE, overexpression. N.S.: non-specific. The protein level of Myh9 was quantified at the bottom panel. *, p-value < 0.05. (B) HEK 293T cells were transfected with various constructs as indicated. Various inhibitors of the autophagy-lysosome pathway (50 μM LY294002, 100 μM Chloroquine, 100 nM BafA1, 20 mM NH₄Cl) were added to the cell culture 6 h before harvest. OE, overexpression. (C) HEK 293T cells were co-transfected with the plasmids as indicated. 18 h later, 100 nM of BafA1 was added to the culture medium. After another 6 h, the soluble whole cell lysates were subjected to immunoprecipitation (IP) with an anti-Myc antibody and the immunoprecipitated proteins were further analysed by Western blot (WB). The red arrow indicates Flag-Styxl2. (D) The schematic showing that Styxl2 promotes sarcomere assembly by binding to and targeting non-muscle myosin IIs for degradation, which facilitates their eventual replacement by muscle myosin II during sarcomere maturation.

| Probe set ID | Gene | Fold change | | | |
| | | P Stage | | D Stage | |
		siJak1	siStat1	siJak1	siStat1
1417889_at	apolipoprotein B editing complex 2	1.53	4.43	2.43	4.07
1419391_at	myogenin	4.29	9.99	2.62	4.26
1422088_at	v-myc myelocytomatosis viral oncogene homolog 1	2.12	3.9	1.88	3.68
1422606_at	C1q and tumor necrosis factor related protein 3	2.23	13.4	1.68	5.45
1426971_at	ubiquitin-activating enzyme E1-like	3.53	7.05	2.42	3.03
1427306_at	ryanodine receptor 1	2.11	6	2.71	4.52
1449178_at	PDZ and LIM domain 3	3.45	12.3	2.94	7
1451453_at	death-associated kinase 2	1.52	3.14	1.96	3.81
1452520_a_at	cholinergic receptor	6.76	15.5	2.6	4.13
1429223_a_at	hemochromatosis type 2 (juvenile) (human homolog)	2.16	28.8	2	7.66
1429459_at	sema domain	2.77	4.43	2.27	5.58
1435828_at	avian musculoaponeurotic fibrosarcoma (v-maf) AS42 oncogene homolog	1.96	3.38	1.63	3.27
1439658_at	leiomodin 3 (fetal)	1.91	17.9	2.83	18.8
1439746_at	dual specificity phosphatase 27 (putative) (dusp27)	1.82	4.78	1.73	2.36
1444494_at	kelch repeat and BTB (POZ) domain containing 10	2.46	9.63	2.79	6.38
1421426_at	Hedgehog-interacting protein	-2.04	-3.55	-1.62	-6.65
1435438_at	SRY-box containing gene 8 (sox8)	1.85	4.01	1.51	5.7

Table 1. Candidate genes regulated by both Jak1 and Stat1. Selected candidate genes targeted by both si-Jak1 and si-Stat1 were shown. The fold change was determined by the relative levels of a gene in cells treated with si-Jak1 or si-Stat1 over that in control cells. Genes with fold change > 1.5 for si-Jak1 and > 2 for si-Stat1 were listed in this table. Genes with positive fold change are up-regulated (positive value), while those with negative fold change are down-regulated. P Stage and D Stage denote the proliferation stage (GM 24h) and early differentiation stage (DM 12h), respectively.