Styxl2 regulates *de novo* sarcomere assembly by binding to non-muscle myosin IIs and promoting their degradation

Xianwei Chen[1#], Yanfeng Li[1#], Jin Xu[1], Yong Cui[2], Qian Wu[3], Haidi Yin[3], Yuying Li[4], Liwen Jiang[2], Huating Wang[4], Zilong Wen[1], Zhongping Yao[3], Zhenguo Wu[1*]

1. Division of Life Science, the Hong Kong University of Science & Technology, Hong Kong, China; 2. School of Life Sciences, the Chinese University of Hong Kong, Hong Kong, China; 3. Department of Applied Biology and Chemical Technology, the Hong Kong Polytechnic University, Hong Kong, China. 4. Department of Orthopaedics and Traumatology, Li Ka Shing Institute of Health Sciences, Chinese University of Hong Kong, Hong Kong, China

\# equal contribution

*Corresponding author:

Division of Life Science,
the Hong Kong University of Science & Technology,
Clearwater Bay, Kowloon
Hong Kong
China
Tel: (852)23588704
Fax: (852)23581552
Email: bczgwu@ust.hk

Running title: Styxl2 promotes sarcomere assembly

Key words:

Styxl2; pseudophosphatase; sarcomere; non-muscle myosin II; autophagy

Abstract

Styxl2, a poorly characterized pseudophosphatase, was identified as a transcriptional target of the Jak1-Stat1 pathway during myoblast differentiation in culture. Styxl2 is specifically expressed in vertebrate striated muscles. By morpholino-mediated gene knockdown or genetic knockout, we found that Styxl2 plays an essential role in maintaining sarcomere integrity in developing muscles of both zebrafish and mice. To further reveal the functions of Styxl2 in adult muscles, we employed two inducible knockout mouse models: one with *Styxl2* deleted in mature myofibers to assess its role in sarcomere maintenance, and the other in adult muscle satellite cells (MuSCs) to assess its role in *de novo* sarcomere assembly. We find that Styxl2 is not required for sarcomere maintenance but plays a role in *de novo* sarcomere assembly during injury-induced muscle regeneration. Mechanistically, Styxl2 interacts with non-muscle myosin IIs and targets them for autophagy-dependent degradation. Without Styxl2, the degradation of non-muscle myosin IIs is delayed, which leads to defective sarcomere assembly and force generation. Thus, Styxl2 promotes *de novo* sarcomere assembly by interacting with non-muscle myosin IIs and facilitating their degradation.

Introduction

Striated muscles, including skeletal and cardiac muscles of vertebrates and some muscles of invertebrates, are unique in that they contain highly-organized, multi-protein contractile structures called myofibrils in the cytosol of myofibers or cardiomyocytes (Clark et al., 2002; Mukund and Subramaniam, 2020; Sparrow and Schöck, 2009). Sarcomeres are the smallest repeating structural units of myofibrils responsible for muscle contraction (Avellaneda et al., 2021; Squire et al., 2005). A sarcomere is flanked by Z-discs (or Z lines) with multiple actin filaments (or thin filaments) anchored to the Z-discs in parallel and the interdigitating bipolar myosin bundles (or thick filaments) anchored to the M-line in the middle of the sarcomere (Squire et al., 2005). As a key component of the Z-disc, an anti-parallel α-actinin dimer directly binds to two actin filaments in opposite directions (Squire et al., 2005). A sarcomere can be further divided into the lighter I (isotropic)-band and the darker A (anisotropic)-band based on their microscopic appearances with the I-band containing actin filaments only and the A-band containing both the myosin bundles and partially overlapping actin filaments (Craig and Padrón, 2004; Squire et al., 2005). Mutations in sarcomeric proteins are linked to congenital human cardiac and skeletal muscle diseases (Bönnemann and Laing, 2004; Parker et al., 2020; Parker and Peckham, 2020). Although the overall structures of sarcomeres are well-established and the protein components of sarcomeres are largely known, it remains controversial how sarcomeres correctly assemble *in vivo*. Several different but not mutually exclusive models including the template model, the stitching model, and the premyofibril model have been proposed (Martin and Kirk, 2020; Rui et al., 2010; Sanger et al., 2006). The template model proposes that actin stress fibers serve as an early template to recruit other sarcomeric components including Z-disc proteins and the thin and thick filaments (Dlugosz et al., 1984). The stitching model or the titin model proposes that the actin filaments held by the Z-disc (i.e., the I-Z-I complexes) and the myosin filaments first assemble independently, and they are then stitched together to form intact sarcomeres with the help of titin as titin is known to simultaneously interact with Z-disc proteins via its N-terminal region and M-line proteins via its C-terminal region (Holtzer et al., 1997; Lu et al., 1992). The premyofibril model states that mini-sarcomeres consisting of actin filaments, α-actinin-containing Z bodies, and non-muscle myosin IIs first assemble near cell periphery. Then, non-muscle myosin IIs are replaced by muscle myosin II during the transition from premyofibrils to nascent myofibrils and finally to mature myofibrils (Rhee et al., 1994; Sanger et al., 2006). Each of these models is supported by some experimental

observations but may be incompatible with others (Chopra et al., 2018; Fenix et al., 2018;
Rui et al., 2010). In addition, mechanical tension generated by attachment of myofibers to
tendon or myofiber bundling is also found to promote sarcomere assembly (Luis and
Schnorrer, 2021).

Protein phosphatases are a family of enzymes that catalyze the removal of the
phosphate group from covalently-linked substrates including proteins, lipids, nucleic acids,
and polysaccharides (Alonso et al., 2004; Tonks, 2006; Weinfeld et al., 2011; Worby et al.,
2006). They antagonize the actions of kinases and regulate various biological processes
including tissue development, cell proliferation, differentiation, and cell migration.
Interestingly, some of the protein phosphatases are found to be catalytically inactive due to
the absence of essential amino acid residues such as the catalytic cysteine in an otherwise
well-conserved phosphatase domain (Chen et al., 2017; Reiterer et al., 2020). These
confirmed or putative catalytically inactive phosphatases are also called pseudophosphatases.
Although catalytically inactive, several pseudophosphatases are found to play important roles
in various organisms (Azzedine et al., 2003; Cheng et al., 2009; Hinton, 2020; Reiterer et al.,
2017). For example, Styx and Sbf1, both members of the pseudophosphatase family, were
found to regulate sperm development. Loss of Styx or Sbf1 resulted in male infertility in mice
(Firestein et al., 2002; Wishart and Dixon, 2002).

In our current study, we identified Styxl2 (also named Dusp27), a poorly-
characterized pseudophosphatase, as a new regulator involved in sarcomere assembly in both
zebrafish and mice. The original name of the gene and its protein product (i.e., Dusp27) has
caused confusion in the literature due to the fact that the same name was once used for
DUPD1 (now re-named as Dusp29), a catalytically active phosphatase (Friedberg et al.,
2007). Recently, it was suggested that Dusp27 be re-named as Serine/threonine/tyrosine-
interacting-like protein 2 (Styxl2) (Cooper and Waddell, 2020). To avoid any potential
confusion, we will use Styxl2 hereafter. Styxl2 is specifically expressed in vertebrate striated
muscles. Our work in both zebrafish and mouse models shows that it plays an important role
in *de novo* sarcomere assembly instead of sarcomere maintenance. Mechanistically, we found
that Styxl2 directly interacts with non-muscle myosin IIs that are involved in the early stage
of sarcomere assembly and subsequently replaced by muscle myosin IIs in mature sarcomeres
(Wang et al., 2018). Styxl2 promotes autophagy-dependent degradation of non-muscle
myosin IIs, which ultimately facilitates sarcomere assembly *in vivo*.

Results

Styxl2 encodes a sarcomeric muscle-specific pseudophosphatase and is a transcriptional target of the Jak1-Stat1 pathway in myoblasts

We previously reported that the Jak1-Stat1 pathway plays an important role in myogenic differentiation in cultured myoblasts (Sun et al., 2007). However, it remains unclear what genes are targeted by the pathway. To examine the changes in transcriptomes, we conducted a microarray analysis in C2C12 myoblasts: cells were first transfected with either a control siRNA or specific siRNAs targeting *Jak1* or *Stat1* and the cells were harvested at two different time points: (1): after 24 hours of growth in the growth medium (GM 24h) (the proliferation stage); (2): after 12 hours in the differentiation medium following 24 hours of growth in the GM (DM 12h, the early differentiation stage). The total mRNA was extracted and subjected to microarray analysis. Among differentially expressed genes (DEGs) in cells transfected with *Jak1*-siRNA and *Stat1*-siRNA, many were shared, which was expected as both Jak1 and Stat1 function in the same pathway. Some representative targets shared by Jak1 and Stat1 were shown (Table S1). Consistent with our previous findings (Sun et al., 2007), *myogenin* gene was negatively regulated by both Jak1 and Stat1. Among the genes targeted by both Jak1 and Stat1, *dual specificity phosphatase 27 (putative) (Dusp27 or Styxl2)* attracted our attention: it is a poorly characterized gene with only one published report in zebrafish so far (Fero et al., 2014). It encodes a member of the dual-specificity phosphatase family with a conserved dual-specificity phosphatase catalytic (DSPc) domain located at its N-terminus (Fig. 1A). However, Styxl2 is predicted to be catalytically inactive due to substitution of a highly-conserved catalytic cysteine by serine in its DSPc domain (Fig. S1A), thus making it a member of the pseudophosphatase subfamily (Hinton, 2019; Kharitidi et al., 2014). By both RT-qPCR and Western blot, we confirmed that both the mRNA and protein levels of *Styxl2* were induced by siRNAs against *Jak1* or *Stat1* in C2C12 myoblasts, with the extent of induction being more obvious during early differentiation (Fig. S1B, C). Consistently, treatment of C2C12 myoblasts with leukemia inhibitory factor (LIF) or oncostatin M (OSM), two cytokines that are known to potently activate the Jak1-Stat1 pathway in myoblasts (Sun et al., 2007; Xiao et al., 2011), indeed led to a decrease of Styxl2 protein (Fig. S1D, E). By Western blot, we found that *Styxl2* has very limited tissue distribution pattern: it is specifically expressed in sarcomeric muscles including the heart and skeletal muscles (Fig. 1B). Consistently, the expression of *Styxl2* mRNA is confined to the heart and somites of early mouse embryos (Fig. S1F). In C2C12 myoblasts, Styxl2 protein

was already present in proliferating myoblasts but was further induced during myogenic
differentiation (Fig. 1C). We also tried to determine the subcellular localization of Styxl2 in
myoblasts. As our home-made polyclonal antibody is not suitable for immunostaining of the
endogenous Styxl2, we examined subcellular localization of Flag- or HA-tagged Styxl2
instead in C2C12 cells by immunostaining. We found that both Flag- and HA-tagged Styxl2
were predominantly localized in the cytoplasm (Fig. S1G). Collectively, Styxl2 is a
cytoplasmic protein that is specifically expressed in sarcomeric muscles.
165
Knockdown of *Styxl2* in zebrafish leads to severe defects in sarcomere structure
The earlier study by *Fero et al.* in zebrafish showed that Styxl2 is required for the integrity of
sarcomeres (Fero et al., 2014). We also confirmed this finding in zebrafish using a
morpholino oligo (MO) that blocks the translation initiation of *Styxl2*. The effectiveness of
the MO was verified using a transgene encoding green fluorescence protein (GFP) fused in-
frame with the same sequence of *Styxl2* (i.e., the first 25 nucleotides of the coding region)
targeted by the MO: when the one-cell zebrafish embryos were microinjected with the
transgenes with or without the MO, the expression of GFP was suppressed only when the MO
was co-injected (Fig. S1H). We then injected one-cell zebrafish embryos with or without the
Styxl2-MO and examined the structures of skeletal muscles by immunostaining at 24 hours
after fertilization (hpf). Consistent with the report by *Fero et al.* (Fero et al., 2014), we found
that both fast and slow myofibers were severely disrupted when *Styxl2* was knocked down
with the MO (Fig. 1E). Moreover, electron microscopy analysis further revealed the disrupted
sarcomere structures in zebrafish muscles injected with the MO (Fig. 1D).
180
Conditional deletion of *Styxl2* leads to neonatal lethality in mutant mice
To explore the role of Styxl2 in mammals, we generated *Styxl2* floxed mice with two LoxP
sites flanking exons 3~5 of the *Styxl2* gene (Fig. S2A). Exons 4 and 5 encode a part of the
DSPc domain of Styxl2. We first aimed to generate germ line *Styxl2* knockout (KO) mice
using EIIA-Cre mice (Lakso et al., 1996). Western blot analysis showed that Styxl2 protein
was totally absent in KO mice (Fig. 2A). In multiple litters, no live homozygous KO pups
were found after birth (Fig. S2B). By examining developing embryos of different ages, we
found that *Styxl2* KO embryos died between E13.5 to E15.5 (Fig. S2C). In some dead *Styxl2*
KO foetuses, edema was found (Fig. 2B), suggesting potential problems with the heart
(Conway et al., 2003). This is consistent with the fact that Styxl2 is also highly expressed in
cardiac muscles as well. The electron microscopy analysis of the heart muscles from E14.5

embryos revealed that the sarcomeric structures of cardiac muscles from *Styxl2* KO embryos
were severely disrupted (Fig. 2C). To focus on the functions of Styxl2 in skeletal muscles, we
generated coditional *Styxl2* KO (CKO) mice by crossing *Styxl2$^{f/f}$* mice (Fig. S2A) with
Pax7$^{Cre/+}$ mice (Keller et al., 2004). We found that *Styxl2* CKO pups were born at the
expected Mendelian ratio but all the homozygous mutant pups died in a few hours after birth
(Fig. 2D, E and Fig. S2D). As Pax7 is also expressed in other non-muscle cell types including
neurons during embryo development, to be certain that embryonic lethality of *Pax7$^{Cre/+}$;*
Styxl2$^{f/f}$ mice was indeed caused by severe defects in skeletal muscles, we also specifically
deleted *Styxl2* in skeletal muscles using a *Myf5^{Cre}* line in which the Cre gene was inserted in
the *Myf5* locus (Tallquist et al., 2000). Consistently, all skeletal muscle-specific *Styxl2* KO
pups died within one day after birth (Fig. S2D).
As the skin of *Styxl2* CKO pups appeared cyanotic, we speculated that the new-born mutant
mice had lung defects that led to respiratory distress and perinatal death. The lungs of the
CKO pups were examined and we found that the pulmonary alveoli indeed failed to open
(Fig. 2F), which was likely caused by defective diaphragm muscle as defects in a number of
myogenic genes exhibited similar perinatal lethality (Hasty et al., 1993; Rudnicki et al., 1993).
Electron microscopy analysis confirmed that the sarcomere structure was severely disrupted
in diaphragm muscle of the mutant mice (Fig. 2G). Consistently, electron microscopy
revealed that the sarcomere structure of the limb muscle was also severely disrupted (Fig.
S2E). Notably, some sarcomeres were still present in the skeletal muscles of the CKO mice
(Fig. S2E), suggesting that Styxl2 is important but not absolutely required for the formation
of the sarcomere structure. Therefore, Styxl2 plays an important role in the development of
striated muscles, including the skeletal muscle and the heart muscle.
215

Styxl2 is dispensable for the maintenance of mature sarcomeres but critical for *de novo*
sarcomere assembly in adult skeletal muscles
Although our data above indicated that Styxl2 plays an important role in regulating
sarcomere formation during embryonic skeletal muscle development, it remained unclear
whether it was necessary for the maintenance of the sarcomere structure in adult skeletal
muscles. To address this question, we generated a tamoxifen (TMX) inducible myofiber-
specific *Styxl2* knockout mouse model (also named *Styxl2* MF-iKO hereafter) using *Tg (HSA-*
MerCreMer) (McCarthy et al., 2012). In this model, *Styxl2* was efficiently deleted in mature
skeletal muscles of adult mice (Fig. 3A, B). We directly assessed the contractile functions of
Styxl2-null muscles. Three weeks after the last TMX injection (Fig. 3A), the gastrocnemius

muscle was isolated and the twitch and tetanic force generated by the muscle were measured (Guo et al., 2015). No obvious difference in twitch or tetanic force was observed between the control and *Styxl2* MF-iKO muscles (Fig. 3C, D), suggesting that Styxl2 is dispensable for the maintenance of existing sarcomeres in adult skeletal muscles. Next, we assessed whether Styxl2 played a role in the *de novo* formation of nascent sarcomeres in regenerating muscles of adult mice. To do so, we generated another tamoxifen inducible, MuSC-specific *Styxl2* KO mouse model using *Pax7^{CreERT2/CreERT2}; Styxl2^{f/f}* (also named *Styxl2* SC-iKO mice hereafter) (Murphy et al., 2011). *Styxl2* was first deleted in muscle satellite cells (MuSCs) of adult mice according to the scheme in Fig. 3E, followed by acute injury to the hindlimb skeletal muscles with Cardiotoxin (CTX) (Fig. 3E). As injured muscles are repaired by MuSCs, the nascent sarcomeres in regenerating myofibers of *Styxl2* SC-iKO mice would be formed in the absence of Styxl2 (Fig. 3F). At 30 days post injury, a time point when injured muscles are known to be fully regenerated (Dumont et al., 2015), the gastrocnemius muscles were isolated and functionally tested *in vitro*. Both the twitch and tetanic force were measured separately. Compared to muscles from the control mice, the twitch force generated by muscles from *Styxl2* SC-iKO mice was decreased by ~30% (Fig. 3G), while the tetanic force generated by muscles from the mutant mice was not affected (Fig. 3H). This suggested that sarcomeres formed *de novo* in regenerating myofibers from Styxl2-null MuSCs were partially defective. When we examined sarcomeres by electron microscopy, we noticed that the length of individual sarcomeres from *Styxl2* SC-iKO mice was shorter in regenerated muscles (Fig. 3I, 3J). Thus, our data showed that Styxl2 functions in *de novo* sarcomere assembly in regenerating muscles from adult mice.

Styxl2 directly binds to non-muscle myosin II (NM II) in myoblasts

To understand how Styxl2 functions to regulate sarcomere formation, we employed BioID, a proximity-dependent protein labelling method (Roux et al., 2018, 2012), to search for binding partners of Styxl2. The full-length Styxl2 was fused in-frame with BirA*, a mutant form of biotin ligase BirA (Fig. 4A). The fused gene encoding Styxl2-BirA*-HA was stably expressed in C2C12 myoblasts using the Tet-on system (Das et al., 2016) (Fig. S3A). As Styxl2 is expected to function during late myogenic differentiation, we followed the scheme in Fig. 4B and added biotin into the culture medium with or without doxycycline (Dox) for 24 hours before harvest . By Western blot, we showed that Styxl2-BirA*-HA was strongly induced by Dox as expected (Fig. S3A, middle panel). Importantly, more cellular proteins were biotinylated in Styxl2-BirA*-HA-expressing cells induced by Dox (Fig. S3A, top panel).

These biotinylated proteins were isolated and enriched by streptavidin beads and analysed by

iTRAQ-based mass spectrometry (Wiese et al., 2007). The biotinylated proteins enriched in

Styxl2-BirA*-HA-expressing cells (Enrichment score > 1) were further analysed by gene

ontology (GO) analysis. Notably, a number of the enriched proteins were related to

"myofibril", "sarcomere", and "A band" (Fig. 4C). This supports our hypothesis that Styxl2

functions to regulate sarcomere assembly. To help identify direct interaction partners of

Styxl2, the $\log_2$ Enrichment scores (i.e., the peak intensity ratios of the reporter ions) were

plotted against the minus $\log_{10}$(p.values). Those enriched proteins with the Enrichment score >

1.2 and $p < 0.05$ were highlighted in red on the plot (Fig. S3B), and some of the enriched

proteins (Fig. S3C) were selected for further confirmation by co-immunoprecipitation (co-IP)

assays using lysates from differentiated C2C12 myotubes. Among them, Myh9 and Myh10,

the members of the non-muscle myosin IIs (NM IIs) (Newell-Litwa et al., 2015), were found

to specifically interact with the endogenous Styxl2 (Fig. 4D). In addition, we also examined

the subcellular localization of Styxl2 using a plasmid expressing zStyxl2 fused with green

fluorescent protein (GFP) in zebrafish. zStyxl2 was found to alternate with α-actinin in

myofibrils of zebrafishes at 48 hpf (Fig. S3D), which is consistent with the previous report

(Fero et al., 2014). Interestingly, it was reported that Myh9 is localized in the mini-A-band in

sarcomeres of zebrafish (Sanger et al., 2009), which supports our finding that Styxl2 interacts

with NM IIs during sarcomere assembly. Moreover, by protein truncation analysis in

combination with the co-IP assays, we showed that the head domains of Myh9 and Myh10

were responsible for their interaction with Styxl2 (Fig. 4E, F).

Styxl2 promotes sarcomere assembly by down-regulating NM IIs

Our data above indicated that Styxl2 regulates sarcomere assembly. NM IIs were reported to

participate in sarcomere assembly in several organisms (Fenix et al., 2018; Sanger et al., 2009;

Tullio et al., 1997). However, NM IIs are not detected in mature sarcomeres and their

replacement by muscle-specific myosin is a marker of skeletal muscle maturation. Therefore,

we examined protein levels of Myh9 and Myh10 in both differentiating C2C12 myotubes and

postnatal mouse skeletal muscles over time. In differentiating C2C12 cells, we found that the

protein levels of Myh9 and Myh10 slightly increased at the beginning of myogenic

differentiation, but started to decrease ~four days after myogenic differentiation, which

coincided with the time when the sarcomeric myosin heavy chain started to increase (Fig.

S4A). In postnatal mouse skeletal muscles, the protein levels of Myh9 and Myh10 started to

drop at P8 and became non-detectable by P30 (Fig. 5A), which was consistent with previous

reports (Fenix et al., 2018; Sanger et al., 2009). The protein levels of Styxl2 remained constant and abundant in the first month after birth but began to drop to a lower level by P65 (Fig. 5A). When we examined the protein levels of Myh9 and Myh10 in skeletal muscles of P1 pups, we noticed an obvious increase in protein levels of Myh9 and Myh10 in *Styxl2* CKO mice (Fig. 5B, C). We also examined protein levels of Myh9 and Myh10 in regenerating adult muscles. Using SC-iKO mice and *Pax7*$^{CreERT2/CreERT2}$ control mice as described in Fig. 3E, we first induced *Styxl2* deletion by tamoxifen in adult MuSCs followed by BaCl$_2$-induced muscle injury. At 7 dpi, the levels of Myh9 and Myh10 were comparable between control and *Styxl2* SC-iKO mice (Fig. 5D). Interestingly, at 10 dpi, Myh9 and Myh10 were nearly undetectable in muscles from control mice, but they remained at a higher level in muscles from *Styxl2* SC-iKO mice (Fig. 5D). Our data above suggested that loss of Styxl2 results in delayed degradation of NM II during sarcomere assembly. To further prove this is the case, we turned to the zebrafish model. It was reported that *Styxl2*$^{-/-}$ zebrafish exhibited severe motility defects in response to needle touch (Fero et al., 2014). Consistently, we also found that *Styxl2* morphants displayed similar phenotypes: most of the *Styxl2* morphants did not respond to the needle touch (Fig. 5E, F), which was caused by defective sarcomere structure in skeletal muscles (Fig. 5G). Knowing that NM IIs are involved in the early phase of sarcomere assembly followed by their replacement with the sarcomeric myosin (Fenix et al., 2018; White et al., 2014), we suspected that delayed degradation of NM IIs was responsible for defective sarcomere structure in *Styxl2* morphants. Therefore, it was likely that simultaneous knockdown of both NM IIs and *Styxl2* may rescue the defects in the sarcomere structure. It was reported that both z*Myh9b* and z*Myh10* are expressed in skeletal muscles of zebrafish (Gutzman et al., 2015). When we knocked down z*Myh9b* with a MO in *Styxl2* morphants, the embryos died very early before skeletal muscles were formed (our unpublished data). When zebrafish zygotes were co-injected with the *Styxl2*-MO and a MO targeting z*Myh10*, the ratio of fishes responding to the needle touch was obviously increased (Fig. 5F). Consistently, the disrupted sarcomere structure caused by *Styxl2*-MO was partially restored in the *Styxl2/Myh10* double-knockdown fishes (Fig. 5G).

Styxl2 induces the degradation of NM II via its C-terminal domain

Our data above showed that loss of Styxl2 caused abnormal accumulation of NM IIs (Fig. 5), which suggested that Styxl2 normally promote degradation of NM IIs during nascent sarcomere assembly. To test whether this is the case, we co-expressed HA-Myh9 with either a control protein or Styxl2 in C2C12 myoblasts. The protein levels of HA-Myh9 were indeed

reduced by co-transfected Styxl2 (Fig. 6A). We further found that Styxl2-mediated reduction

in Myh9 protein levels was not caused by reduced *Myh9* mRNA (data not shown). To

determine which part of Styxl2 promotes degradation of NM IIs, we truncated Styxl2 into

two fragments: an N-terminal fragment containing the DSPc domain (i.e., aa 1-513, also

termed Styxl2N513 hereafter) and a C-terminal fragment without the DSPc domain (i.e., aa

510-1138, also named Styxl2ΔN509 hereafter). When we co-transfected 293T cells with a

construct encoding HA-Myh9 together with another construct encoding Mst1 (as a negative

control), Styxl2N513, or Styxl2ΔN509, we found that only Styxl2ΔN509 promoted

degradation of Myh9 (Fig. 6C, lane 3). As we previously showed that the head domains of

NM IIs were involved in interacting with Styxl2 (Fig. 4E, F), we then tested whether the

protein levels of the head domain of Myh9 were also regulated by Styxl2. Consistently, only

Styxl2ΔN509 promoted degradation of the head domain of Myh9 (Fig. 6C, lane 6). To further

verify this finding *in vivo*, we microinjected the mRNA encoding the full-length or the C-

terminus of zStyxl2, along with the MO for z*Styxl2*, into the zebrafish one-cell embryos. As

expected, the defective sarcomere structure caused by *Styxl2* knockdown was efficiently

rescued by overexpressing the mRNA encoding the full-length zStyxl2. Importantly, the

mRNA encoding the C-terminus of zStyxl2 was nearly as effective as that encoding the full-

length zStyxl2 (Fig. 6B). Thus, the C-terminus of Styxl2, without the DSPc domain, is both

necessary and sufficient to promote degradation of NM IIs during nascent sarcomere

assembly.

Styxl2 promotes the degradation of NM IIs via the autophagy-lysosome pathway

In order to understand how Styxl2 promotes the degradation of NM IIs, we first turned to the

ubiquitination/proteasome pathway (Nandi et al., 2006). MG132, a specific inhibitor of the

proteasome-dependent protein degradation pathway, was added to C2C12 cells transfected

with Styxl2 (Fig. S4B). No abnormal accumulation of NM IIs was found in the presence of

MG132, suggesting that the proteasome-dependent system is less likely involved in Styxl2-

mediated degradation of NM IIs. Next, we turned to the autophagy-lysosome-dependent

protein degradation system. In C2C12 myoblasts, we co-expressed Myh9 with either a

control protein or Styxl2 together with an siRNA against GFP (control) or ATG5, a key

component of the autophagy pathway (Mizushima et al., 2010). In the presence of the control

siRNA, Styxl2 efficiently promoted degradation of Myh9 as expected (Fig. 7A, lane 2).

However, in the presence of ATG5 siRNA, the protein levels of Myh9 were restored to levels

comparable to that of the control (Fig. 7A, lane 1) even in the presence of the co-expressed

Styxl2 (Fig. 7A, lane 3). This result supported the involvement of the autophagy pathway in

Styxl2-mediated degradation of NM IIs. Furthermore, we also examined the degradation of

Myh9 induced by the C-terminal domain of Styxl2 with or without several different

autophagy inhibitors (Fig. 7B). In the absence of any autophagy inhibitors, Styxl2ΔN509

induced degradation of Myh9 as expected (Fig. 7B, lanes 3 and 4). In the presence of

LY294002 (the inhibitor for PI3K), chloroquine (the inhibitor of the binding of

autophagosomes to lysosomes), bafilomycin A1 (BafA1, the inhibitor of the fusion between

autophagosomes and lysosomes), and NH_4Cl (the inhibitor of the degradative enzyme

activation inside lysosomes), the protein levels of Myh9 were largely restored to the levels

comparable to that in the controls (Fig. 7B, lanes 1 and 2). Thus, we concluded that the

autophagy-lysosome pathway is involved in the degradation of NMIIs induced by Styxl2.

Discussion

Styxl2 is a pseudophosphatase that functions independently of its conserved phosphatase domain. Pseudophosphatases are present in multicellular organisms ranging from *C. elegans* to human (Hinton, 2019; Reiterer et al., 2020). Moreover, pseudophosphatases have also been identified in plants (Bellec et al., 2002). Recently, 26 pseudophosphatase domains, including the phosphatase domain of Styxl2, were bioinformatically identified in the human genome (Chen et al., 2017). Due to the absence of one or several conserved amino acids including the catalytic cysteine, pseudophosphatases are not expected to possess the phosphatase activity. STYX, the first molecularly characterized pseudophosphatase, was identified in 1995 as a phosphoserine/threonine/tyrosine-binding protein (Wishart et al., 1995). The substitution of the catalytic cysteine by glycine in the active site motif leads to the loss of its catalytic activity (Reiterer et al., 2013; Wishart et al., 1995). In addition to STYX, the functions of several other pseudophosphatases from *C. elegans* to human have also been characterized (Cheng et al., 2009; Hinton et al., 2010; Maruyama et al., 2007; Nakhro et al., 2013; Tomar et al., 2019). These reports unequivocally demonstrate that pseudophosphatases are not simply evolution relics but play essential roles in multicellular organisms.

Pseudophosphatases exert their functions via diverse modes including direct competition with other proteins by binding to a phospho-residue-containing motif, modulation of the phosphatase activity by binding to an active phosphatase, sequestration of target proteins in specific subcellular locations, and integration of signals by simultaneous binding with multiple proteins (Hinton, 2019; Reiterer et al., 2020). The original report on STYX showed that mutation of the glycine in the conserved HGX$_5$R motif back to cysteine restored the phosphatase activity (Wishart et al., 1995), suggesting that the three-dimensional structure of the phosphatase domain in STYX is still folded properly. Thus, some pseudophosphatases are thought to function by binding to specific phosphorylated substrates, which is the case for pseudophosphatases EGG-4/5 and Pas2 (Bellec et al., 2002; Cheng et al., 2009; Da Costa et al., 2006). In these examples, pseudophosphatases bind the phosphorylated sites of selected target proteins through their inactive phosphatase domains and prevent them from binding other interacting proteins or active phosphatases. However, this is not the only way pseudophosphatases function. MTMR13, a pseudophosphatase, functions by binding to MTMR2, an active lipid phosphatase, to regulate its phosphatase activity. Mutations in either *MTMR13* or *MTMR2* are causally linked to human Type 4B Charcot-Marie-Tooth disease, an

inherited disorder of the peripheral nervous system with abnormal nerve myelination (Berger

et al., 2006; Robinson and Dixon, 2005).

Unlike several other characterized pseudophosphatases that function via their

conserved phosphatase domains, Styxl2 appears to function independently of its phosphatase

domain as the phosphatase domain of Styxl2 is dispensable for its ability to induce

degradation of NM IIs and to rescue the sarcomere defects in the *Styxl2* morphant zebrafish

(Fig. 6B, C). Such a phosphatase domain-independent mode of action is also observed in His

Domain Protein Tyrosine Phosphatase (HD-PTP or PTPN23), a pseudophosphatase known to

facilitate endosomal cargo sorting and the degradation of multiple membrane receptors

including epidermal growth factor receptor, platelet-derived growth factor receptor, MHC

class I and Integrin α5β1 (Doyotte et al., 2008; Kharitidi et al., 2015; Ma et al., 2015;

Parkinson et al., 2015). Interestingly, a truncated fragment of HD-PTP without its

phosphatase domain is still capable of promoting endosomal trafficking of cargo proteins like

epidermal growth factor (EGF) (Doyotte et al., 2008).

Styxl2 facilitates *de novo* sarcomere assembly by promoting autophagy-dependent

degradation of non-muscle myosin IIs. Although absent in invertebrates, Styxl2 is

expressed and conserved in vertebrates. The specific involvement of Styxl2 in sarcomere

assembly is supported by its restricted expression in sarcomeric muscles. Interestingly,

mutation in the coding region of *STYXL2* and abnormal CpG methylation at the *STYXL2*

locus were detected in some patients with congenital heart defects (CHD), suggesting a

possible link of *STYXL2* to CHD (Arrington Cammon B. et al., 2012; Radhakrishna et al.,

2016). Moreover, its role in sarcomere assembly is also conserved in vertebrates as

interference of the gene expression in zebrafish by either a transposon integration or

morpholino oligos leads to severe defects in sarcomere assembly (Fero et al., 2014) (Fig. 1D,

E). However, Styxl2 is not absolutely essential for sarcomere assembly as sarcomeres in

Drosophila flight muscles can normally assemble in the absence of a Styxl2 homolog.

Furthermore, we could still find residual sarcomeres in skeletal muscles of conditional *Styxl2*

KO mice (driven by either Pax7[Cre] or Myf5[Cre]) (Fig. S2E).

Although the highly-ordered sarcomere structure has been known for many years, it

remains controversial how sarcomeres correctly assemble. Several different but not mutually

exclusive models have been proposed (Rui et al., 2010; Sanger et al., 2017, 2006). Studies in

Drosophila, *C. elegans*, and mice underscore an essential role for integrin in initiating the

assembly of sarcomeres (Rui et al., 2010; Sparrow and Schöck, 2009). For example, β1

integrin was shown to be essential for sarcomere assembly in mice (Schwander et al., 2003).
Among the existing sarcomere assembly models, the prevailing model of sarcomere assembly
is called premyofibril model (Rhee et al., 1994; Sanger et al., 2017, 2006). In this model, α-
actinin, skeletal muscle actin and non-muscle myosin IIs (NM IIs) first assemble to form
mini-sarcomeres or premyofibrils at the cell periphery. Then, additional sarcomeric proteins
like titin, nebulin, troponin, and myomesin will be incorporated and NM IIs will be
eventually replaced by skeletal muscle myosin in mature sarcomeres (White et al., 2018,
2014). Consistent with this model, germline knockout of NM IIB (i.e., Myh10) in mice led to
early lethality between E14.5 and birth and sarcomere structural defects in embryonic
cardiomyocytes (Tullio et al., 1997). Similarly, defective sarcomeres were also observed in
embryonic muscles of stage-17 mutant fly null for *zipper* (*zip*), the *Drosophila* gene encoding
the heavy chain of NM II (Bloor and Kiehart, 2001; Loison et al., 2018). Moreover, a recent
paper using human iPSC-derived cardiomyocytes demonstrated that nascent sarcomeres
assemble directly from muscle actin stress fibers (Fenix et al., 2018). Both MYH9 and
MYH10 are present in muscle actin stress fibers with MYH10 being essential for sarcomere
assembly. Consistent with the Premyofibril model, NM IIs are found to be replaced by
muscle myosin IIs in mature sarcomeres (Fenix et al., 2018). Nevertheless, the view of the
involvement of NM IIs in early sarcomere assembly is also challenged by several studies.
Unlike germline knockout of NM IIB, cardiac-specific knockout of NM IIB in adult mice did
not generate obvious defects in sarcomere structures (Ma et al., 2009). Although NM IIA is
not detected in embryonic hearts, loss of cardiac NM IIB led to increased expression of NM
IIA, suggesting that NM IIA may compensate for the loss of NM IIB in NM IIB-null
embryonic cardiomyocytes (Fenix et al., 2018; Tullio et al., 1997). Moreover, it was also
suspected that the cardiomyocyte-specific deletion of NM IIB occurs after the initial
sarcomere assembly due to the expression timing of the Cre recombinase under the control of
the cardiac α-MHC promoter (Fenix et al., 2018; Ma et al., 2009). Furthermore, a recent
study used CRISPR/Cas9 to knock out *MYH9* and *MYH10* genes either individually or in
combination in human iPSC-derived cardiomyocytes and showed that neither gene is
required for sarcomere assembly in their cell culture models (Chopra et al., 2018). It is
puzzling at present why the results on the involvement of NM IIs in *de novo* sarcomere
assembly differ in the two studies by Fenix *et al.* and Chopra *et al.* even though both studies
used human iPSC-derived cardiomyocytes to assay *de novo* sarcomere assembly in cell
culture.

It also remains unclear how NM IIs are replaced by muscle myosin during *de novo*
sarcomere assembly in vertebrates. Our current study showed that Styxl2 interacts with NM
IIs and induces their degradation via the autophagy-lysosome pathway. Autophagy is known
to regulate sarcomere integrity (Masiero et al., 2009; Nakai et al., 2007). In particular,
selective autophagy is known to degrade proteins, protein aggregates, and organelles
(Bogomolovas et al., 2021; Kirkin et al., 2009; Lange et al., 2005; Shaid et al., 2013;
Zaffagnini and Martens, 2016). Some ubiquitinated proteins including sarcomeric titin are
degraded by selective autophagy. p62/SQSTM1 and NBR1 have been shown to act as
adaptors linking ubiquitinated substrates to Atg8/LC3 that is localized on the membrane of
autophagosome (Kirkin et al., 2009; Pankiv et al., 2007). Interestingly, a recent study on
ciliogenesis showed that MYH9 could be degraded by selective autophagy in an NEK9-
mediated manner (Yamamoto et al., 2021). It remains to be tested whether NM IIs are
ubiquitinated during the late stage of sarcomere assembly before their autophagy-dependent
degradation and whether NEK9 is involved in this process. It also remains to be elucidated
how exactly Styxl2 promotes autophagy-dependent degradation of NM IIs.

Our data showed that Styxl2 plays indispensable role during embryonic myogenesis
as conditional deletion of *Styxl2* during mouse embryo development using either *Pax7^{Cre}* or
Myf5^{Cre} resulted in severe disruption of sarcomere structures in skeletal muscles and perinatal
lethality (Fig. 2 and Fig. S2). In contrast, inducible deletion of *Styxl2* in adult muscles only
led to mild defects in sarcomere structures (Fig. 3F-J). It is likely that additional unknown
molecules can compensate for the loss of Styxl2 in adult regenerating muscles. Such distinct
roles of Styxl2 during development are also supported by differential expression patterns of
Styxl2 and NM IIs: both are abundantly expressed in new-born and juvenile mice. While the
protein levels of Myh9 and Myh10 start to drop one week after birth and become
undetectable by three weeks after birth, that of Styxl2 is relatively stable in the first month
after birth but starts to drop sharply by two months after birth (Fig. 5A). This is also
consistent with our findings that Styxl2 is not required for the maintenance of the sarcomeres
in adult muscles (Fig. 3A-D).

Materials and Methods

Animals

ABSR wild-type zebrafish were maintained following standard procedures and the zygotes were collected for microinjection experiments. $Pax7^{CreERT2/+}$ mice (Stock No: 017763) were purchased from the Jackson Laboratory (Bar Harbor, ME, USA). $Myf5^{Cre/+}$, $Pax7^{Cre/+}$, and Tg $(HSA-MerCreMer)$ mice were generous gifts from Drs. Michael Rudnicki (Ottawa Hospital Research Institute, ON, Canada), Mario Capecchi (University of Utah, UT, USA), and Karyn Esser (University of Florida, FL, USA), respectively. $Styxl2^{flox/flox}$ mice were generated by the Model Animal Research Centre (Nanjing University, Nanjing, China). To delete $Styxl2$ in adult mice, tamoxifen dissolved in corn oil (15 mg/ml) was injected into adult mice intraperitoneally at 5 µl per gram of body weight for 5 consecutive days followed by another 5 doses of tamoxifen injection every 3-4 days. All animals were maintained and handled following the protocols approved by the Animal Ethics Committee of the Hong Kong University of Science and Technology.

Cell culture and DNA constructs

C2C12 cells and HEK 293T cells were purchased from ATCC (Manassas, VA, USA). C2C12 cells were maintained in growth medium (GM) (Dulbecco's modified Eagle's medium supplemented with 20% fetal bovine serum, 100 Units/ml penicillin, and 100 µg/ml streptomycin) and differentiation was induced with differentiation medium (DM) (Dulbecco's modified Eagle's medium supplemented with 2% horse serum, 100 Units/ml penicillin, and 100 µg/ml streptomycin). HEK 293T cells were maintained in Dulbecco's modified Eagle's medium supplemented with 10% fetal bovine serum, 100 Units/ml penicillin, and 100 µg/ml streptomycin. Mouse Styxl2, Styxl2ΔN509 and Styxl2N513 were amplified from a mouse muscle cDNA library and inserted into a modified pcDNA3.0 vector containing a Flag-expressing cassette to generate constructs encoding Flag-Styxl2, Flag-Styxl2ΔN509 and Flag-Styxl2N513. BirA* was amplified from pcDNA3.1 mycBioID (addgene, #35700) and an HA tag was added to its C-terminus. The cDNA encoding BirA*-HA was fused in-frame with that encoding mouse Styxl2 and cloned into the pRetroX-Tight-Pur vector (kindly provided by Dr. Yusong Guo, the Hong Kong University of Science and Technology) to get pTight-Styxl2-BirA*-HA. Mouse Myh9, Myh9-head and Myh9-tail were amplified from a plasmid encoding Myosin-IIA-GFP (addgene, #38297) and inserted into a modified pcDNA3.0 vector containing an HA-expressing cassette to generate constructs

encoding HA-Myh9, HA-Myh9-head and HA-Myh9-tail. Mouse Myh10-head and Myh10-

tail were amplified from a mouse C2C12 cell cDNA library and inserted into a modified

pcDNA3.0 vector containing an HA-expressing cassette to generate constructs encoding HA-

Myh10-head and HA-Myh10-tail. Fish Styxl2 and Styxl2ΔN493 were amplified from a

zebrafish cDNA library and inserted into either pCS2+ vector or pCS2+-GFP vector to

generate pCS2+-zStyxl2 and pCS2+-zStyxl2ΔN493 or pCS2+-GFP-zStyxl2.

RNA extraction and RT-qPCR

Total RNA was extracted from C2C12 cells using the TRIzol Reagent (Invitrogen) following

the manufacturer's instructions. The reverse transcription was performed using an oligo-dT

primer and the ImRrom-II Reverse Transcription System (Promega). The ABI 7700 Sequence

Detection System (PE Applied-Biosystems) was used for qPCR and the relative fold change

of genes of interest was analysed based on the $2^{-\Delta\Delta Ct}$ method (Livak and Schmittgen, 2001).

The qPCR primers used were listed below:

Styxl2-F: 5'- GCCCATCCACCTCTCCTC-3'

Styxl2-R: 5'- GCTCCTGTCATCCATCTTCTC-3'

Myh9-F: 5'-GGCCCTGCTAGATGAGGAGT-3'

Myh9-R: 5'-CTTGGGCTTCTGGAACTTGG-3'

Myh10-F: 5'-GGAATCCTTTGGAAATGCGAAGA-3'

Myh10-R: 5'-GCCCCAACAATATAGCCAGTTAC-3'

Gapdh-F: 5'-CCCACTCTTCCACCTTCG-3'

Gapdh-R: 5'-TCCTTGGAGGCCATGTAG-3'

siRNA transfection

The following siRNAs were used to knock down specific genes. They were transfected into

C2C12 cells using the Lipofectamine RNAiMax reagent (Thermo Fisher Scientific) following

manufacturer's instructions.

GFP (negative control): 5'-GCTGACCCTGAAGTTCATC-3'

Jak1: 5'-GCCTGAGAGTGGAGGTAAC-3'

Stat1: 5'-GGATCAAGTCATGTGCATA-3'

ATG5: 5'-GGCTCCTGGATTATGTCAT-3'

Microarray

Total mRNA were extracted from duplicated samples and subjected to microarray analysis

using the Affymetrix GeneChip® Mouse Genome 430 2.0 Array. The data were analysed by

the GeneSpring software.

Plasmid transfection

The Lipofectamine Transfection Reagent (Thermo Fisher Scientific) and the PLUS Reagent

(Thermo Fisher Scientific) were used to transfect plasmids into C2C12 myoblasts following

manufacturer's instructions. Polyethylenimine (PEI, Polysciences) was used to transfect

plasmids into HEK 293T cells.

Western blot

For cultured cells, Western blot was performed according to the standard procedures. For

mouse tissues, the samples were homogenized using a T-25 digital IKA®ULTRA-TURRAX

disperser in a cell lysis buffer (25 mM Tris-HCl pH7.4, 150 mM NaCl, 1 mM EDTA, 1%

NP-40, 5% glycerol) with a cocktail of protease inhibitors (20 mM p-nitrophenylphosphate,

20 mM β-glycerolphosphate, 0.5 mM phenylmethylsulfonyl fluoride, 2 μg/ml aprotinin, 0.5

μg/ml leupeptin, 0.7 μg/ml pepstatin, 50 μM sodium vanadate). Then the soluble lysates were

used for Western blot. The primary antibodies used include anti-Jak1 (Upstate, 06-272), anti-

Stat1 (Upstate, 06-501), anti-β-Tubulin (Sigma, T4026), anti-sarcomere myosin heavy chain

(MHC) (DSHB, MF20), anti-GAPDH (Ambion, AM4300), anti-Myh9 (Cell Signaling, 3403),

anti-Myh10 (Santa Cruz, sc-376942), anti-Flag (Sigma, F3165), and anti-HA (Roche,

12CA5). The Styxl2 antiserum was generated in-house by injecting rabbits with recombinant

proteins spanning aa 2-142 of Styxl2 fused in-frame with glutathione S-transferase. The

antiserum was purified with protein-G beads. The streptavidin-HRP was purchased from GE

Healthcare (#RPN1231).

Immunostaining

C2C12 myoblasts were fixed with 4% paraformaldehyde (PFA) for 10 min at room

temperature, and then washed with 0.1% PBST. Then samples were permeabilized with 0.5%

PBST for 20 min. Zebrafishes were fixed with PFA for 2-4 h at room temperature and

washed with 0.1% PBST. Then the fishes were permeabilized with cold acetone for 8 min at -

20°C. The following procedures are the same for two types of samples. After

permeabilization, the samples were washed with 0.1% PBST and blocked with 4% IgG-free

BSA for 1 h and then incubated with the primary antibody at 4°C overnight. Then samples
were washed with 0.1% PBST and incubated with the secondary antibody at room
temperature for 1 h. The DNA was stained with 100 ng/ml 4′,6 diamidino 2 phenylindole
(DAPI). For fish slow muscles, the nucleus was stained with an anti-Prox1 antibody. The
primary antibodies used include anti-Flag (Sigma, F7425), anti-HA (Santa Cruz, sc-805),
anti-α-Actinin (Sigma, A7732), anti-Prox1 (Angiobio, 11-002P), anti-slow myosin heavy
chain (DSHB, F59), anti-myosin heavy chain (DSHB, A4.1025), and anti-sarcomere myosin
heavy chain (MHC) (DSHB, MF20).

Microinjection of zebrafish zygotes
The zygotes of zebrafish were collected in the morning. 2 nl of either plasmid DNA- or
morpholino-containing solution was injected into fish zygotes as report before (Jin et al.,
2009). The concentration of pCS2+-GFP-zStyxl2 was 25 ng/μl. The concentration of
morpholinos used for microinjection was 0.5 mM. A control morpholino targeting a non-
related gene was used as negative control (Ctrl-MO). Morpholinos used were synthesized by
Gene Tools (USA) and their sequences were listed below:
Ctrl-MO: 5’-AGCACACAAAGGCGAAGGTCAACAT-3’
Styxl2-MO: 5’-GCTGATCCTCCACAGACGACGCCAT-3’
Myh10-MO: 5’-CTTCACAAATGTGGTCTTACCTTGA-3’
The DNA constructs of pCS2+-zStyxl2 (with several synonymous mutations near the start
codon of zStyxl2 to make it MO-resistant) and pCS2+-zStyxl2ΔN493 were linearized and
used as templates for *in vitro* transcription. The mRNA was synthesized using the
mMESSAGE mMACHINE® SP6 Kit (Ambion, AM1340) followed by purification with the
MEGAclear™ Transcription Clean-Up Kit (Invitrogen, AM1908). The purified mRNA was
dissolved in RNase-free water and diluted to 150 ng/μl immediately before microinjection.

Transmission Electron Microscopy
The heart and skeletal muscles were processed following the published protocols (Graham
and Orenstein, 2007). Then the samples were examined by a Hitachi H7650 transmission
electron microscope (Hitachi High-Technologies). The sarcomere length was measured from
electron microscopic images with a magnification of 50,000 X. The length of 3-5 sarcomeres
was measured for each mouse and a total of six mice were examined. The results were

presented as mean ± SD. The unpaired Student's t-test was performed and $p < 0.05$ was

considered statistically significant.

Measurement of muscle force generation *In vitro*

The gastrocnemius muscle of left hindlimb was carefully dissected from mice together with

femur condyle and Achilles tendon. The twitch and tetanic force generated by the dissected

gastrocnemius muscle and the muscle cross-sectional area (MCSA) were measured following

published protocols (Guo et al., 2015). The force generated was recorded by the Dynamic

Muscle Control system (DMC v5.4; Aurora Scientific) and analysed by the Dynamic Muscle

Analysis system (DMA v3.2; Aurora Scientific). Both twitch and tetanic force were then

normalized to the MCSA.

BioID and mass spectrometry

C2C12 myoblasts were infected by a retrovirus carrying pRetroX-Tet-On Advanced (kindly

provided by Dr. Yusong Guo, the Hong Kong University of Science and Technology) and

selected by 1 mg/ml G418 to get the Tet-on cells. Then the Tet-on cells were infected by a

retrovirus carrying pTight-Styxl2-BirA*-HA and selected by 1.5 μg/ml puromycin to obtain

stable Tet-Styxl2-BirA*-HA cells. Thus, the expression of Styxl2-BirA*-HA is controlled by

the Tet-on system in these stable C2C12 myoblasts. These stable cells were seeded in GM

and cultured for 24 h before being induced to differentiate in DM for 48 h. The medium was

then replaced with DM containing 50 μM biotin (control) or 50 μM biotin plus 1 μg/ml

doxycycline. After another 24 h, the control and doxycycline-treated cells were processed for

mass spectrometry analysis following the published protocols (Roux et al., 2018). Briefly, the

biotinylated proteins were enriched by Dynabeads MyOne Streptavidin C1 (Invitrogen,

#65001). The proteins bound to the beads were reduced with 10 mM Tris(2-

carboxyethyl)phosphine hydrochloride (Sigma, C4706) at 40°C for 30 min and alkylated with

20 mM Iodoacetamide (Sigma, I1149) in the dark for 30 min to prevent re-formation of the

disulfide bond. The samples were digested with Trypsin Gold (Promega, V5280) overnight at

37°C on a Thermomixer at 700 rpm followed by centrifugation. The supernatant was

transferred to a new tube, acidified to pH2-3 by trifluoroacetic acid (Thermofisher, 85183),

and desalted with a Pierce C18 spin column (Thermo Fisher Scientific, 89873). The tryptic

peptides were labelled with iTRAQ reagents (AB Sciex, 4390812) following the

manufacturer's instructions. Briefly, peptides from control cells were labelled with iTRAQ

tag 114 and those from doxycycline-treated cells were labelled with tag 115. The labelled samples were mixed and fractionated by an UltiMate 3000 RSLCnano system (Thermo Fisher Scientific). A C18 Acclaim PepMap RSLC analytical column (75 μm × 250 mm, 2 μm, 100Å) with a C18 nano Viper trap-column (0.3 mm × 5 mm, Thermo Fisher Scientific) was used for peptide elution and separation at a flow rate of 300 nl/min . The mobile phase contained buffer A (0.1% formic acid, 2% acetonitrile) and B (0.1% formic acid, 98% acetonitrile). The gradient was set as follows: 0 min, 3% B; 10 min, 3% B; 12 min, 7% B; 62 min, 20% B; 64 min, 30% B; 65 min, 80% B; 73 min, 80% B; 73.1 min, 3% B; and 90 min, 3% B. MS data were then acquired using an Orbitrap Fusion Lumos mass spectrometer (Thermo Fisher Scientific) in a positive mode, with the following settings: 1) MS1 scan: 400 to 1,600 m/z; resolution: 120,000; automatic gain control (AGC): 400,000; and maximum injection time: 50 ms; 2) The collision energy was set at 35% and orbitrap was used for MS2 scan as well; 3) MS2 scan starts from 100 m/z; resolution: 30,000; AGC: 50,000; and maximum injection time: 250 ms; 4) Exclusion window was set for 40 s; 5) The intensity threshold was set at 25,000. Data analysis was carried out using Mascot Deamon version 2.5.0 (Matrix Science) with standard settings for each instrument type and the data were searched against the *Mus musculus* database downloaded from SwissProt. A peptide tolerance of 10 ppm and fragment ion tolerance of 20 ppm were used. Carbamidomethylation of cysteine was specified as fixed modification, while oxidation of methionine and acetylation of protein N termini were set as variable modifications. The false discovery rate was set at 0.01 for proteins and peptide spectrum match. The 115/114 iTRAQ ratio of one protein was calculated based on the peak intensity of reporter ions by the software, and defined as Enrichment score for proteins in doxycycline-treated cells over that in control cells. The *p* values for each protein generated by the software were used for gene ontology analysis.

Gene ontology enrichment analysis

All the biotinylated proteins enriched (with Enrichment score > 1) in cells expressing Styxl2-BirA*-HA were imported and analyzed at Geneontology (http://geneontology.org/). The terms in the cellular component category were presented.

Co-immunoprecipitation

C2C12 cells or HEK293T cells were lysed in the lysis buffer (25 mM Tris-HCl pH7.4, 150 mM NaCl, 1 mM EDTA, 1% NP-40, 5% glycerol) with a cocktail of protease inhibitors. The soluble whole cell lysates were first immunoprecipitated with an antibody targeting Styxl2,

702 MEF2A (negative control), or Flag (Roche, F3165). The proteins co-precipitated were
703 analysed by Western blot.
704

Injury-induced muscle regeneration
Avertin was used to anaesthetize mice at 0.5 mg per gram of body weight by intraperitoneal
injection. Then 120 µl (for females) or 150 µl (for males) of 10 µM Cardiotoxin was injected
into muscles of each hind limb to induce acute injury. After >30 days, the muscles were
dissected and tested. To examine protein levels at different time points, each tibialis anterior
muscle was injected with 30 µl of 1.2% $BaCl_2$ to induce acute damage. The regenerating
muscles were collected for Western blot at different time points.
712

Acknowledgements
The project was supported by research grants from the Hong Kong Research Grant Council
(C6018-19G, AoE/M-604/16, T13-602/21-N, and T13-605/18-W), the State Key Laboratory
of Molecular Neuroscience at HKUST, the Hong Kong Center for Neurodegenerative
Diseases (HKCeND), and the Shenzhen Bay Laboratory (S201101002, Guangdong, China).
We thank Drs. K Esser, M Rudnicki, and M Capecchi for providing mouse Cre lines used in
this study. We also thank Dr. Y Guo for providing the BioID vector and Drs. Y Dai and S
Zhao for their help with zebrafish work.

Competing interests
The authors declare no competing interests.

Figure Legends

Figure 1. Styxl2 regulates sarcomere integrity during zebrafish muscle development. (A) A schematic of mouse Styxl2 protein. DSPc (red): dual-specificity phosphatase, catalytic domain. The numbers denote the positions of various amino acids in mouse Styxl2 including Ser (S)-225. (B) The expression of Styxl2 protein in different mouse tissues. 100 μg of soluble mouse tissue lysates were subjected to Western blot analysis. GAPDH: glyceraldehyde-3-phosphate dehydrogenase. SM: skeletal muscles. (C) The expression of Styxl2 protein in C2C12 myoblasts before and after differentiation in culture. MHC: myosin heavy chain. (D) Zebrafish zygotes were either mock-injected or injected with a Styxl2-specific morpholino (MO). Sarcomeres in muscles at 48 hpf were revealed by transmission electron microscopy (TEM). Representative images were shown. Scale bar: 500 nm. (E) Sarcomeres in slow and fast muscles of zebrafish embryos at 24 hpf were revealed by immunostaining using specific antibodies as indicated. Ctrl: control (non-treated); MO: *Styxl2*-morpholino-treated. The nuclei of slow and fast muscle fibers were counter-stained with an anti-Prox1 antibody and DAPI, respectively. Scale bar: 10 μm.

Figure 2. Conditional deletion of mouse *Styxl2* leads to defective sarcomeres in striated muscles. (A) The expression of Styxl2 protein in skeletal muscles of wild-type (+/+), heterozygous (+/-) and *Styxl2* knockout (KO) (-/-) (driven by *EIIA*-Cre) mice. Soluble tissue lysates from limbs of E14.5 mouse embryos were subjected to Western blot analysis. N.S.: non-specific. (B) E14.5 embryos of control (Ctrl) and *EIIA*-Cre;*Styxl2^f/f^* KO mice. *: edema was found in some *Styxl2* KO fetuses. (C) The sarcomere structures of cardiomyocytes in the heart of E14.5 embryos from control and *EIIA*-Cre;*Styxl2^f/f^* KO mice were revealed by TEM. Scale bar: 1 μm. (D) The expression of Styxl2 protein in skeletal muscles of control (Ctrl) and *Styxl2* CKO P1 mice. (E) The newborn pups (P1) from control (Ctrl) and *Pax7*^Cre/+^;*Styxl2^f/f^* KO mice (CKO). (F) Lung sections of control and Styxl2 CKO P1 mice. Scale bar: 50 μm. (G) The sarcomere structure of diaphragm muscles from control (Ctrl) and *Styxl2* CKO P1 mice revealed by TEM. Scale bar: 1 μm.

Figure 3. Styxl2 is not required for sarcomere maintenance but involved in *de novo* sarcomere assembly in adult muscles. (A) The schematic to assess the role of Styxl2 in sarcomere maintenance in adult muscles. TMX: tamoxifen. (B) Styxl2 protein levels in gastrocnemius muscles of Ctrl and MF-iKO mice. (C, D) The twitch and tetanic force

generated by gastrocnemius muscles of three Ctrl and four MF-iKO mice were measured. In

(A-D), Ctrl: *Styxl2^{f/f}* mice; MF-iKO: *Tg: HSA-MerCreMer;Styxl2^{f/f}* mice. (E) The schematic

to assess the role of Styxl2 in *de novo* sarcomere assembly in adult muscles. CTX:

cardiotoxin. (F) Styxl2 protein levels in regenerated gastrocnemius muscles. N.S.: non-

specific. (G, H) Twitch and tetanic force by regenerated gastrocnemius muscles were

measured from four pairs of control and SC-iKO mice. (I) TEM images of the muscle used

for force generation test. Scale bar: 500 nm. (J) Quantification of sarcomere length in (I).

TEM images from three pairs of mice were examined and the length of 3-5 sarcomeres from

each   mouse   was   measured.   In   (E-J),   Ctrl: *Pax7^{CreERT2/CreERT2}*   mice;   SC-iKO:

Pax7^{CreERT2/CreERT2};Styxl2^{f/f} mice. n.s.: not significant. *, p-value <0.05. **, p-value <0.01.

Figure 4. Styxl2 binds to non-muscle myosin IIs. (A) The schematic of the fusion protein

Styxl2-BirA*-HA. (B) The work flow of sample preparation for BioID. (C) Enriched (with

Enrichment score >1) biotinylated proteins in cells expressing Styxl2-BirA*-HA relative to

that in control cells were identified by mass spectrometry followed by Gene Ontology (GO)

analysis. The top ten groups were shown. (D) C2C12 cells were harvested after 2 days in DM

and the soluble whole cell lysates were subjected to immunoprecipitation (IP) using either an

anti-MEF2A (negative control) or an anti-Styxl2 antibody. The endogenous Myh9 or Myh10

was detected by Western blot. (E) The schematic showing the domains of non-muscle myosin

IIs (NM II). (F) HEK 293T cells were transfected with Flag-Styxl2 (black arrow) together

with HA-NM II-head (red arrowheads) or HA-NM II-tail (black arrowheads). The soluble

whole cell lysates were subjected to IP with an anti-Flag antibody and the co-

immunoprecipitated proteins were detected by Western blot.

Figure 5. Styxl2 protein levels inversely correlates with that of non-muscle myosin IIs.

(A) Protein levels of Styxl2 and non-muscle myosin IIs in skeletal muscles of wild-type mice

of different ages were determined by Western blot. (B) Protein levels of Styxl2 and non-

muscle myosin IIs in skeletal muscles of control and *Styxl2* CKO P1 mice were determined

by Western blot. (C) Relative ratios of non-muscle myosin II protein levels in control and

Styxl2 CKO P1 mice were quantified by ImageJ based on the data from four (for Myh9) or

five (for Myh10) sets of Western blots. (D) Protein levels of Styxl2 and non-muscle myosin

IIs in regenerating tibialis anterior muscles were determined by Western blot. The

experiments were done for four times with similar results and a representative gel was shown.

Ctrl: *Pax7^{CreERT2/CreERT2}* mice; SC-iKO: *Pax7^{CreERT2/CreERT2}; Styxl2^{f/f}* mice; dpi: days post-

injury. (E) Zebrafish zygotes were separately injected with the following morpholinos (MO): control (Ctrl-MO), *Styxl2*-MO, and mixed morpholinos targeting both *Styxl2* and *Myh10* (dMO). At 48 hpf, zebrafishes were poked by a fine needle and their responses over 400 milliseconds (mS) were recorded. (E' and E"): representative images showing positive and no response of zebrafish to needle touch, respectively. (F) Quantification of responses of zebrafish to needle touch. In each group, at least 8 zebrafishes were tested and the percentage of zebrafishes responding to needle touch was calculated. (G) Zebrafishes from each group in (E) were subjected to immunostaining for α-actinin to visualize sarcomere structures. Representative images were shown. Scale bar: 10 μm. *, p-value < 0.05. ****, p-value < 0.0001.

Figure 6. The DSPc domain of Styxl2 is dispensable for degradation of non-muscle myosin IIs. (A) DNA constructs encoding Myh9, Styxl2 or Mst1 (negative control) were co-expressed in C2C12 cells. Soluble whole cell extracts were subjected to Western blot analysis. OE: overexpression. (B) Zebrafish zygotes were injected with various morpholinos as indicated with or without the co-injected *Styxl2* mRNAs. z*Styxl2*FL: the full-length fish *Styxl2* mRNA; z*Styxl2*ΔN493: the truncated fish *Styxl2* mRNA missing the 5' region encoding the N-terminal 493 aa. Representative α-actinin immunofluorescent images were shown. The number in numerator at the bottom left corner of each image represents fish embryos showing normal α-actinin staining pattern, while that in denominator represents the total number of embryos examined. Scale bar: 10 μm. (C) HEK 293T cells were transfected with different plasmids as indicated. 24 h after transfection, cells were harvested and soluble whole cell extracts were subjected to Western blot analysis. The arrow indicates the head domain of Myh9, and the arrowhead indicates Styxl2 missing the N-terminal 509 aa (Styxl2ΔN509). Mst1: negative control. N.S.: non-specific.

Figure 7. The Styxl2-mediated degradation of non-muscle myosin IIs is autophagy dependent. (A) Plasmids encoding Myh9, Styxl2 or Mst1 (negative control) were co-expressed in C2C12 cells together with various siRNAs as indicated. Soluble whole cell extracts were subjected to Western blot analysis. OE, overexpression. N.S.: non-specific. (B) HEK 293T cells were transfected with various constructs as indicated. Various inhibitors of the autophagy-lysosome pathway (50 μM LY294002, 100 μM Chloroquine, 100 nM BafA1, 20 mM NH$_4$Cl) were added to the cell culture 6 h before harvest. OE, overexpression. (C)

826 The schematic showing that Styxl2 promotes sarcomere assembly by binding to and targeting

827 non-muscle myosin IIs for degradation, which facilitates their eventual replacement by

828 muscle myosin II during sarcomere maturation.

Reference:

Alonso, A., Sasin, J., Bottini, N., Friedberg, Ilan, Friedberg, Iddo, Osterman, A., Godzik, A., Hunter, T., Dixon, J., Mustelin, T., 2004. Protein tyrosine phosphatases in the human genome. Cell 117, 699–711. https://doi.org/10.1016/j.cell.2004.05.018

Arrington Cammon B., Bleyl Steven B., Matsunami Norisada, Bonnell Gabriel D., Otterud Brith E.M., Nielsen Douglas C., Stevens Jeffrey, Levy Shawn, Leppert Mark F., Bowles Neil E., 2012. Exome Analysis of a Family With Pleiotropic Congenital Heart Disease. Circ. Cardiovasc. Genet. 5, 175–182. https://doi.org/10.1161/CIRCGENETICS.111.961797

Avellaneda, J., Rodier, C., Daian, F., Brouilly, N., Rival, T., Luis, N.M., Schnorrer, F., 2021. Myofibril and mitochondria morphogenesis are coordinated by a mechanical feedback mechanism in muscle. Nat. Commun. 12, 2091. https://doi.org/10.1038/s41467-021-22058-7

Azzedine, H., Bolino, A., Taïeb, T., Birouk, N., Di Duca, M., Bouhouche, A., Benamou, S., Mrabet, A., Hammadouche, T., Chkili, T., Gouider, R., Ravazzolo, R., Brice, A., Laporte, J., LeGuern, E., 2003. Mutations in MTMR13, a new pseudophosphatase homologue of MTMR2 and Sbf1, in two families with an autosomal recessive demyelinating form of Charcot-Marie-Tooth disease associated with early-onset glaucoma. Am. J. Hum. Genet. 72, 1141–1153. https://doi.org/10.1086/375034

Bellec, Y., Harrar, Y., Butaeye, C., Darnet, S., Bellini, C., Faure, J.-D., 2002. Pasticcino2 is a protein tyrosine phosphatase-like involved in cell proliferation and differentiation in Arabidopsis. Plant J. 32, 713–722. https://doi.org/10.1046/j.1365-313X.2002.01456.x

Berger, P., Berger, I., Schaffitzel, C., Tersar, K., Volkmer, B., Suter, U., 2006. Multi-level regulation of myotubularin-related protein-2 phosphatase activity by myotubularin-related protein-13/set-binding factor-2. Hum. Mol. Genet. 15, 569–579. https://doi.org/10.1093/hmg/ddi473

Bloor, J.W., Kiehart, D.P., 2001. zipper Nonmuscle Myosin-II Functions Downstream of PS2 Integrin in Drosophila Myogenesis and Is Necessary for Myofibril Formation. Dev. Biol. 239, 215–228. https://doi.org/10.1006/dbio.2001.0452

Bogomolovas, J., Fleming, J.R., Franke, B., Manso, B., Simon, B., Gasch, A., Markovic, M., Brunner, T., Knöll, R., Chen, J., Labeit, S., Scheffner, M., Peter, C., Mayans, O., 2021. Titin kinase ubiquitination aligns autophagy receptors with mechanical signals in the sarcomere. EMBO Rep. 22, e48018. https://doi.org/10.15252/embr.201948018

Bönnemann, C.G., Laing, N.G., 2004. Myopathies resulting from mutations in sarcomeric proteins. Curr. Opin. Neurol. 17, 529–537.

Chen, M.J., Dixon, J.E., Manning, G., 2017. Genomics and evolution of protein phosphatases. Sci. Signal. 10, eaag1796. https://doi.org/10.1126/scisignal.aag1796

Cheng, K.C.-C., Klancer, R., Singson, A., Seydoux, G., 2009. Regulation of MBK-2/DYRK by CDK-1 and the Pseudophosphatases EGG-4 and EGG-5 during the Oocyte-to-Embryo Transition. Cell 139, 560–572. https://doi.org/10.1016/j.cell.2009.08.047

Chopra, A., Kutys, M.L., Zhang, K., Polacheck, W.J., Sheng, C.C., Luu, R.J., Eyckmans, J., Hinson, J.T., Seidman, J.G., Seidman, C.E., Chen, C.S., 2018. Force Generation via β-Cardiac Myosin, Titin, and α-Actinin Drives Cardiac Sarcomere Assembly from Cell-Matrix Adhesions. Dev. Cell 44, 87-96.e5. https://doi.org/10.1016/j.devcel.2017.12.012

Clark, K.A., McElhinny, A.S., Beckerle, M.C., Gregorio, C.C., 2002. Striated Muscle Cytoarchitecture: An Intricate Web of Form and Function. Annu. Rev. Cell Dev. Biol. 18, 637–706. https://doi.org/10.1146/annurev.cellbio.18.012502.105840

Conway, S.J., Kruzynska☐Frejtag, A., Kneer, P.L., Machnicki, M., Koushik, S.V., 2003.
What cardiovascular defect does my prenatal mouse mutant have, and why? genesis
35, 1–21. https://doi.org/10.1002/gene.10152

Cooper, L.M., Waddell, D.S., 2020. A tale of two DUSP27s: proposed resolution for the
naming of distinct dual-specificity phosphatases. Am. J. Physiol.-Cell Physiol. 319,
C148–C150. https://doi.org/10.1152/ajpcell.00201.2020

Craig, R., Padrón, R., 2004. Chapter 7 Molecular Structure of the Sarcomere, in: Myology.
pp. 129–166.

Da Costa, M., Bach, L., Landrieu, I., Bellec, Y., Catrice, O., Brown, S., De Veylder, L.,
Lippens, G., Inzé, D., Faure, J.-D., 2006. Arabidopsis PASTICCINO2 Is an
Antiphosphatase Involved in Regulation of Cyclin-Dependent Kinase A. Plant Cell 18,
1426–1437. https://doi.org/10.1105/tpc.105.040485

Das, A.T., Tenenbaum, L., Berkhout, B., 2016. Tet-On Systems For Doxycycline-inducible
Gene Expression. Curr. Gene Ther. 16, 156–167.
https://doi.org/10.2174/1566523216666160524144041

Dlugosz, A.A., Antin, P.B., Nachmias, V.T., Holtzer, H., 1984. The relationship between
stress fiber-like structures and nascent myofibrils in cultured cardiac myocytes. J. Cell
Biol. 99, 2268–2278. https://doi.org/10.1083/jcb.99.6.2268

Doyotte, A., Mironov, A., McKenzie, E., Woodman, P., 2008. The Bro1-related protein HD-
PTP/PTPN23 is required for endosomal cargo sorting and multivesicular body
morphogenesis. Proc. Natl. Acad. Sci. U. S. A. 105, 6308–6313.
https://doi.org/10.1073/pnas.0707601105

Dumont, N., Bentzinger, C., Sincennes, M.-C., Rudnicki, M., 2015. Satellite Cells and
Skeletal Muscle Regeneration. Compr. Physiol. 5, 1027–1059.
https://doi.org/10.1002/cphy.c140068

Fenix, A.M., Neininger, A.C., Taneja, N., Hyde, K., Visetsouk, M.R., Garde, R.J., Liu, B.,
Nixon, B.R., Manalo, A.E., Becker, J.R., Crawley, S.W., Bader, D.M., Tyska, M.J.,
Liu, Q., Gutzman, J.H., Burnette, D.T., 2018. Muscle-specific stress fibers give rise to
sarcomeres in cardiomyocytes. eLife 7, e42144. https://doi.org/10.7554/eLife.42144

Fero, K., Bergeron, S.A., Horstick, E.J., Codore, H., Li, G.H., Ono, F., Dowling, J.J., Burgess,
H.A., 2014. Impaired embryonic motility in dusp27 mutants reveals a developmental
defect in myofibril structure. Dis. Model. Mech. 7, 289–298.
https://doi.org/10.1242/dmm.013235

Firestein, R., Nagy, P.L., Daly, M., Huie, P., Conti, M., Cleary, M.L., 2002. Male infertility,
impaired spermatogenesis, and azoospermia in mice deficient for the
pseudophosphatase Sbf1. J. Clin. Invest. 109, 1165–1172.
https://doi.org/10.1172/JCI12589

Friedberg, Ilan, Nika, K., Tautz, L., Saito, K., Cerignoli, F., Friedberg, Iddo, Godzik, A.,
Mustelin, T., 2007. Identification and characterization of DUSP27, a novel dual-
specific protein phosphatase. FEBS Lett. 581, 2527–2533.
https://doi.org/10.1016/j.febslet.2007.04.059

Graham, L., Orenstein, J.M., 2007. Processing tissue and cells for transmission electron
microscopy in diagnostic pathology and research. Nat. Protoc. 2, 2439–2450.
https://doi.org/10.1038/nprot.2007.304

Guo, A., Leung, K., Qin, J., Chow, S.K., Cheung, W., 2015. Effect of Low-Magnitude, High-
Frequency Vibration Treatment on Retardation of Sarcopenia: Senescence-
Accelerated Mouse-P8 Model. Rejuvenation Res. 19, 293–302.
https://doi.org/10.1089/rej.2015.1759

Gutzman, J.H., Sahu, S.U., Kwas, C., 2015. Non-muscle myosin IIA and IIB differentially
928   regulate cell shape changes during zebrafish brain morphogenesis. Dev. Biol. 397,
929   103–115. https://doi.org/10.1016/j.ydbio.2014.10.017
930   Hasty, P., Bradley, A., Morris, J.H., Edmondson, D.G., Venuti, J.M., Olson, E.N., Klein,
931   W.H., 1993. Muscle deficiency and neonatal death in mice with a targeted mutation in
932   the myogenin gene. Nature 364, 501–506. https://doi.org/10.1038/364501a0
933   Hinton, S.D., 2020. Pseudophosphatase MK-STYX: the atypical member of the MAP kinase
934   phosphatases. FEBS J. 287, 4221–4231. https://doi.org/10.1111/febs.15426
935   Hinton, S.D., 2019. The role of pseudophosphatases as signaling regulators. Biochim.
936   Biophys. Acta BBA - Mol. Cell Res., Protein Phosphatases as Critical Regulators for
937   Cellular Homeostasis 1866, 167–174. https://doi.org/10.1016/j.bbamcr.2018.07.021
938   Hinton, S.D., Myers, M.P., Roggero, V.R., Allison, L.A., Tonks, N.K., 2010. The
939   pseudophosphatase MK-STYX interacts with G3BP and decreases stress granule
940   formation. Biochem. J. 427, 349–357. https://doi.org/10.1042/BJ20091383
941   Holtzer, H., Hijikata, T., Lin, Z.X., Zhang, Z.Q., Holtzer, S., Protasi, F., Franzini-Armstrong,
942   C., Sweeney, H.L., 1997. Independent assembly of 1.6 microns long bipolar MHC
943   filaments and I-Z-I bodies. Cell Struct. Funct. 22, 83–93.
944   https://doi.org/10.1247/csf.22.83
945   Jin, H., Sood, R., Xu, J., Zhen, F., English, M.A., Liu, P.P., Wen, Z., 2009. Definitive
946   hematopoietic stem/progenitor cells manifest distinct differentiation output in the
947   zebrafish VDA and PBI. Development 136, 647–654.
948   https://doi.org/10.1242/dev.029637
949   Keller, C., Hansen, M.S., Coffin, C.M., Capecchi, M.R., 2004. Pax3:Fkhr interferes with
950   embryonic Pax3 and Pax7 function: implications for alveolar rhabdomyosarcoma cell
951   of origin. Genes Dev. 18, 2608–2613. https://doi.org/10.1101/gad.1243904
952   Kharitidi, D., Apaja, P.M., Manteghi, S., Suzuki, K., Malitskaya, E., Roldan, A., Gingras, M.-
953   C., Takagi, J., Lukacs, G.L., Pause, A., 2015. Interplay of Endosomal pH and Ligand
954   Occupancy in Integrin α5β1 Ubiquitination, Endocytic Sorting, and Cell Migration.
955   Cell Rep. 13, 599–609. https://doi.org/10.1016/j.celrep.2015.09.024
956   Kharitidi, D., Manteghi, S., Pause, A., 2014. Pseudophosphatases: Methods of analysis and
957   physiological functions. Methods, Protein Tyrosine Phosphatase Methods 65, 207–
958   218. https://doi.org/10.1016/j.ymeth.2013.09.009
959   Kirkin, V., McEwan, D.G., Novak, I., Dikic, I., 2009. A Role for Ubiquitin in Selective
960   Autophagy. Mol. Cell 34, 259–269. https://doi.org/10.1016/j.molcel.2009.04.026
961   Lakso, M., Pichel, J.G., Gorman, J.R., Sauer, B., Okamoto, Y., Lee, E., Alt, F.W., Westphal,
962   H., 1996. Efficient in vivo manipulation of mouse genomic sequences at the zygote
963   stage. Proc. Natl. Acad. Sci. U. S. A. 93, 5860–5865.
964   https://doi.org/10.1073/pnas.93.12.5860
965   Lange, S., Xiang, F., Yakovenko, A., Vihola, A., Hackman, P., Rostkova, E., Kristensen, J.,
966   Brandmeier, B., Franzen, G., Hedberg, B., Gunnarsson, L.G., Hughes, S.M.,
967   Marchand, S., Sejersen, T., Richard, I., Edström, L., Ehler, E., Udd, B., Gautel, M.,
968   2005. The Kinase Domain of Titin Controls Muscle Gene Expression and Protein
969   Turnover. Science 308, 1599–1603. https://doi.org/10.1126/science.1110463
970   Livak, K.J., Schmittgen, T.D., 2001. Analysis of relative gene expression data using real-time
971   quantitative PCR and the 2(-Delta Delta C(T)) Method. Methods San Diego Calif 25,
972   402–408. https://doi.org/10.1006/meth.2001.1262
973   Loison, O., Weitkunat, M., Kaya-Çopur, A., Alves, C.N., Matzat, T., Spletter, M.L.,
974   Luschnig, S., Brasselet, S., Lenne, P.-F., Schnorrer, F., 2018. Polarization-resolved
975   microscopy reveals a muscle myosin motor-independent mechanism of molecular

actin ordering during sarcomere maturation. PLOS Biol. 16, e2004718.
https://doi.org/10.1371/journal.pbio.2004718

Lu, M.H., DiLullo, C., Schultheiss, T., Holtzer, S., Murray, J.M., Choi, J., Fischman, D.A.,
Holtzer, H., 1992. The vinculin/sarcomeric-alpha-actinin/alpha-actin nexus in
cultured cardiac myocytes. J. Cell Biol. 117, 1007–1022.
https://doi.org/10.1083/jcb.117.5.1007

Luis, N.M., Schnorrer, F., 2021. Mechanobiology of muscle and myofibril morphogenesis.
Cells Dev. 168, 203760. https://doi.org/10.1016/j.cdev.2021.203760

Ma, H., Wardega, P., Mazaud, D., Klosowska-Wardega, A., Jurek, A., Engström, U.,
Lennartsson, J., Heldin, C.-H., 2015. Histidine-domain-containing protein tyrosine
phosphatase regulates platelet-derived growth factor receptor intracellular sorting and
degradation. Cell. Signal. 27, 2209–2219.
https://doi.org/10.1016/j.cellsig.2015.07.020

Ma, X., Takeda, K., Singh, A., Yu, Z.-X., Zerfas, P., Blount, A., Liu, C., Towbin, J.A.,
Schneider, M.D., Adelstein, R.S., Wei, Q., 2009. Conditional Ablation of Nonmuscle
Myosin II-B Delineates Heart Defects in Adult Mice. Circ. Res. 105, 1102–1109.
https://doi.org/10.1161/CIRCRESAHA.109.200303

Martin, T.G., Kirk, J.A., 2020. Under construction: The dynamic assembly, maintenance, and
degradation of the cardiac sarcomere. J. Mol. Cell. Cardiol. 148, 89–102.
https://doi.org/10.1016/j.yjmcc.2020.08.018

Maruyama, R., Velarde, N.V., Klancer, R., Gordon, S., Kadandale, P., Parry, J.M., Hang, J.S.,
Rubin, J., Stewart-Michaelis, A., Schweinsberg, P., Grant, B.D., Piano, F., Sugimoto,
A., Singson, A., 2007. EGG-3 Regulates Cell-Surface and Cortex Rearrangements
during Egg Activation in Caenorhabditis elegans. Curr. Biol. 17, 1555–1560.
https://doi.org/10.1016/j.cub.2007.08.011

Masiero, E., Agatea, L., Mammucari, C., Blaauw, B., Loro, E., Komatsu, M., Metzger, D.,
Reggiani, C., Schiaffino, S., Sandri, M., 2009. Autophagy Is Required to Maintain
Muscle Mass. Cell Metab. 10, 507–515. https://doi.org/10.1016/j.cmet.2009.10.008

McCarthy, J.J., Srikuea, R., Kirby, T.J., Peterson, C.A., Esser, K.A., 2012. Inducible Cre
transgenic mouse strain for skeletal muscle-specific gene targeting. Skelet. Muscle 2,
8. https://doi.org/10.1186/2044-5040-2-8

Mizushima, N., Yoshimori, T., Levine, B., 2010. Methods in Mammalian Autophagy
Research. Cell 140, 313–326. https://doi.org/10.1016/j.cell.2010.01.028

Mukund, K., Subramaniam, S., 2020. Skeletal muscle: A review of molecular structure and
function, in health and disease. Wiley Interdiscip. Rev. Syst. Biol. Med. 12, e1462.
https://doi.org/10.1002/wsbm.1462

Murphy, M.M., Lawson, J.A., Mathew, S.J., Hutcheson, D.A., Kardon, G., 2011. Satellite
cells, connective tissue fibroblasts and their interactions are crucial for muscle
regeneration. Development 138, 3625–3637. https://doi.org/10.1242/dev.064162

Nakai, A., Yamaguchi, O., Takeda, T., Higuchi, Y., Hikoso, S., Taniike, M., Omiya, S.,
Mizote, I., Matsumura, Y., Asahi, M., Nishida, K., Hori, M., Mizushima, N., Otsu, K.,
2007. The role of autophagy in cardiomyocytes in the basal state and in response to
hemodynamic stress. Nat. Med. 13, 619–624. https://doi.org/10.1038/nm1574

Nakhro, K., Park, J.-M., Hong, Y.B., Park, J.H., Nam, S.H., Yoon, B.R., Yoo, J.H., Koo, H.,
Jung, S.-C., Kim, H.-L., Kim, J.Y., Choi, K.-G., Choi, B.-O., Chung, K.W., 2013.
SET binding factor 1 (SBF1) mutation causes Charcot-Marie-Tooth disease type 4B3.
Neurology 81, 165–173. https://doi.org/10.1212/WNL.0b013e31829a3421

Nandi, D., Tahiliani, P., Kumar, A., Chandu, D., 2006. The ubiquitin-proteasome system. J.
Biosci. 31, 137–155. https://doi.org/10.1007/BF02705243

Newell-Litwa, K.A., Horwitz, R., Lamers, M.L., 2015. Non-muscle myosin II in disease: mechanisms and therapeutic opportunities. Dis. Model. Mech. 8, 1495–1515. https://doi.org/10.1242/dmm.022103

Pankiv, S., Clausen, T.H., Lamark, T., Brech, A., Bruun, J.-A., Outzen, H., Øvervatn, A., Bjørkøy, G., Johansen, T., 2007. p62/SQSTM1 Binds Directly to Atg8/LC3 to Facilitate Degradation of Ubiquitinated Protein Aggregates by Autophagy*. J. Biol. Chem. 282, 24131–24145. https://doi.org/10.1074/jbc.M702824200

Parker, F., Baboolal, T.G., Peckham, M., 2020. Actin Mutations and Their Role in Disease. Int. J. Mol. Sci. 21, E3371. https://doi.org/10.3390/ijms21093371

Parker, F., Peckham, M., 2020. Disease mutations in striated muscle myosins. Biophys. Rev. 12, 887–894. https://doi.org/10.1007/s12551-020-00721-5

Parkinson, M.D.J., Piper, S.C., Bright, N.A., Evans, J.L., Boname, J.M., Bowers, K., Lehner, P.J., Luzio, J.P., 2015. A non-canonical ESCRT pathway, including histidine domain phosphotyrosine phosphatase (HD-PTP), is used for down-regulation of virally ubiquitinated MHC class I. Biochem. J. 471, 79–88. https://doi.org/10.1042/BJ20150336

Radhakrishna, U., Albayrak, S., Alpay-Savasan, Z., Zeb, A., Turkoglu, O., Sobolewski, P., Bahado-Singh, R.O., 2016. Genome-Wide DNA Methylation Analysis and Epigenetic Variations Associated with Congenital Aortic Valve Stenosis (AVS). PLOS ONE 11, e0154010. https://doi.org/10.1371/journal.pone.0154010

Reiterer, V., Fey, D., Kolch, W., Kholodenko, B.N., Farhan, H., 2013. Pseudophosphatase STYX modulates cell-fate decisions and cell migration by spatiotemporal regulation of ERK1/2. Proc. Natl. Acad. Sci. 110, E2934–E2943. https://doi.org/10.1073/pnas.1301985110

Reiterer, V., Pawłowski, K., Desrochers, G., Pause, A., Sharpe, H.J., Farhan, H., 2020. The dead phosphatases society: a review of the emerging roles of pseudophosphatases. FEBS J. 287, 4198–4220. https://doi.org/10.1111/febs.15431

Reiterer, V., Pawłowski, K., Farhan, H., 2017. STYX: a versatile pseudophosphatase. Biochem. Soc. Trans. 45, 449–456. https://doi.org/10.1042/BST20160279

Rhee, D., Sanger, J.M., Sanger, J.W., 1994. The premyofibril: Evidence for its role in myofibrillogenesis. Cell Motil. 28, 1–24. https://doi.org/10.1002/cm.970280102

Robinson, F.L., Dixon, J.E., 2005. The phosphoinositide-3-phosphatase MTMR2 associates with MTMR13, a membrane-associated pseudophosphatase also mutated in type 4B Charcot-Marie-Tooth disease. J. Biol. Chem. 280, 31699–31707. https://doi.org/10.1074/jbc.M505159200

Roux, K.J., Kim, D.I., Burke, B., May, D.G., 2018. BioID: A Screen for Protein-Protein Interactions. Curr. Protoc. Protein Sci. 91, 19.23.1-19.23.15. https://doi.org/10.1002/cpps.51

Roux, K.J., Kim, D.I., Raida, M., Burke, B., 2012. A promiscuous biotin ligase fusion protein identifies proximal and interacting proteins in mammalian cells. J. Cell Biol. 196, 801–810. https://doi.org/10.1083/jcb.201112098

Rudnicki, M.A., Schnegelsberg, P.N.J., Stead, R.H., Braun, T., Arnold, H.-H., Jaenisch, R., 1993. MyoD or Myf-5 is required for the formation of skeletal muscle. Cell 75, 1351–1359. https://doi.org/10.1016/0092-8674(93)90621-V

Rui, Y., Bai, J., Perrimon, N., 2010. Sarcomere Formation Occurs by the Assembly of Multiple Latent Protein Complexes. PLOS Genet. 6, e1001208. https://doi.org/10.1371/journal.pgen.1001208

Sanger, J.W., Kang, S., Siebrands, C.C., Freeman, N., Du, A., Wang, J., Stout, A.L., Sanger, J.M., 2006. How to build a myofibril. J. Muscle Res. Cell Motil. 26, 343–354. https://doi.org/10.1007/s10974-005-9016-7

Sanger, J.W., Wang, J., Fan, Y., White, J., Mi-Mi, L., Dube, D.K., Sanger, J.M., Pruyne, D.,
2017. Assembly and Maintenance of Myofibrils in Striated Muscle. Handb. Exp.
Pharmacol. 235, 39–75. https://doi.org/10.1007/164_2016_53
Sanger, J.W., Wang, J., Holloway, B., Du, A., Sanger, J.M., 2009. Myofibrillogenesis in
skeletal muscle cells in zebrafish. Cell Motil. 66, 556–566.
https://doi.org/10.1002/cm.20365
Schwander, M., Leu, M., Stumm, M., Dorchies, O.M., Ruegg, U.T., Schittny, J., Müller, U.,
2003. β1 Integrins Regulate Myoblast Fusion and Sarcomere Assembly. Dev. Cell 4,
673–685. https://doi.org/10.1016/S1534-5807(03)00118-7
Shaid, S., Brandts, C.H., Serve, H., Dikic, I., 2013. Ubiquitination and selective autophagy.
Cell Death Differ. 20, 21–30. https://doi.org/10.1038/cdd.2012.72
Sparrow, J.C., Schöck, F., 2009. The initial steps of myofibril assembly: integrins pave the
way. Nat. Rev. Mol. Cell Biol. 10, 293–298. https://doi.org/10.1038/nrm2634
Squire, J.M., Al☐khayat, H.A., Knupp, C., Luther, P.K., 2005. Molecular Architecture in
Muscle Contractile Assemblies, in: Advances in Protein Chemistry, Fibrous Proteins:
Muscle and Molecular Motors. Academic Press, pp. 17–87.
https://doi.org/10.1016/S0065-3233(04)71002-5
Sun, L., Ma, K., Wang, H., Xiao, F., Gao, Y., Zhang, W., Wang, K., Gao, X., Ip, N., Wu, Z.,
2007. JAK1–STAT1–STAT3, a key pathway promoting proliferation and preventing
premature differentiation of myoblasts. J. Cell Biol. 179, 129–138.
https://doi.org/10.1083/jcb.200703184
Tallquist, M.D., Weismann, K.E., Hellstrom, M., Soriano, P., 2000. Early myotome
specification regulates PDGFA expression and axial skeleton development.
Development 127, 5059–5070.
Tomar, V.S., Baral, T.K., Nagavelu, K., Somasundaram, K., 2019. Serine/threonine/tyrosine-
interacting-like protein 1 (STYXL1), a pseudo phosphatase, promotes oncogenesis in
glioma. Biochem. Biophys. Res. Commun. 515, 241–247.
https://doi.org/10.1016/j.bbrc.2019.05.093
Tonks, N.K., 2006. Protein tyrosine phosphatases: from genes, to function, to disease. Nat.
Rev. Mol. Cell Biol. 7, 833–846. https://doi.org/10.1038/nrm2039
Tullio, A.N., Accili, D., Ferrans, V.J., Yu, Z.-X., Takeda, K., Grinberg, A., Westphal, H.,
Preston, Y.A., Adelstein, R.S., 1997. Nonmuscle myosin II-B is required for normal
development of the mouse heart. Proc. Natl. Acad. Sci. 94, 12407–12412.
https://doi.org/10.1073/pnas.94.23.12407
Wang, J., Fan, Y., Sanger, J.M., Sanger, J.W., 2018. Nonmuscle myosin II in cardiac and
skeletal muscle cells. Cytoskeleton 75, 339–351. https://doi.org/10.1002/cm.21454
Weinfeld, M., Mani, R.S., Abdou, I., Aceytuno, R.D., Glover, J.N.M., 2011. Tidying up loose
ends: the role of polynucleotide kinase/phosphatase in DNA strand break repair.
Trends Biochem. Sci. 36, 262–271. https://doi.org/10.1016/j.tibs.2011.01.006
White, J., Barro, M.V., Makarenkova, H.P., Sanger, J.W., Sanger, J.M., 2014. Localization of
Sarcomeric Proteins During Myofibril Assembly in Cultured Mouse Primary Skeletal
Myotubes. Anat. Rec. 297, 1571–1584. https://doi.org/10.1002/ar.22981
White, J., Wang, J., Fan, Y., Dube, D.K., Sanger, J.W., Sanger, J.M., 2018. Myofibril
Assembly in Cultured Mouse Neonatal Cardiomyocytes. Anat. Rec. 301, 2067–2079.
https://doi.org/10.1002/ar.23961
Wiese, S., Reidegeld, K.A., Meyer, H.E., Warscheid, B., 2007. Protein labeling by iTRAQ: A
new tool for quantitative mass spectrometry in proteome research. PROTEOMICS 7,
340–350. https://doi.org/10.1002/pmic.200600422

Wishart, M.J., Denu, J.M., Williams, J.A., Dixon, J.E., 1995. A Single Mutation Converts a
Novel Phosphotyrosine Binding Domain into a Dual-specificity Phosphatase. J. Biol.
Chem. 270, 26782–26785. https://doi.org/10.1074/jbc.270.45.26782
Wishart, M.J., Dixon, J.E., 2002. The archetype STYX/dead-phosphatase complexes with a
spermatid mRNA-binding protein and is essential for normal sperm production. Proc.
Natl. Acad. Sci. 99, 2112–2117. https://doi.org/10.1073/pnas.251686198
Worby, C.A., Gentry, M.S., Dixon, J.E., 2006. Laforin, a Dual Specificity Phosphatase That
Dephosphorylates Complex Carbohydrates *. J. Biol. Chem. 281, 30412–30418.
https://doi.org/10.1074/jbc.M606117200
Xiao, F., Wang, H., Fu, X., Li, Y., Ma, K., Sun, L., Gao, X., Wu, Z., 2011. Oncostatin M
inhibits myoblast differentiation and regulates muscle regeneration. Cell Res. 21,
350–364. https://doi.org/10.1038/cr.2010.144
Yamamoto, Y., Chino, H., Tsukamoto, S., Ode, K.L., Ueda, H.R., Mizushima, N., 2021.
NEK9 regulates primary cilia formation by acting as a selective autophagy adaptor for
MYH9/myosin IIA. Nat. Commun. 12, 3292. https://doi.org/10.1038/s41467-021-
23599-7
Zaffagnini, G., Martens, S., 2016. Mechanisms of Selective Autophagy. J. Mol. Biol.,
Molecular Mechanisms of Autophagy, Part A 428, 1714–1724.
https://doi.org/10.1016/j.jmb.2016.02.004
1142

Fig. 1

Fig. 2

Fig. 3

Fig. 4

Fig. 5

Fig. 6

Fig. 7