Distinction between small RNA-bound and free ARGONAUTE via an N-terminal

protein-protein interaction site

Simon Bressendorff[1,4], Ida Marie Zobbe Sjøgaard[1,4], Andreas Prestel[2], Birthe B.

Kragelund[2], Christian Poulsen[1,3] and Peter Brodersen[1,5*]

[1]Copenhagen Plant Science Center, University of Copenhagen, Ole Maaløes Vej 5, DK-

2200 Copenhagen N, Denmark.

[2]Structural Biology and NMR Laboratory, Department of Biology, University of

Copenhagen, Ole Maaløes Vej 5, DK-2200 Copenhagen N, Denmark.

[3]Present address: Novo Nordisk A/S, Novo Nordisk Park 1, DK-2760 Måløv, Denmark.

[4]These authors contributed equally to this work

[5] Lead contact, email pbrodersen@bio.ku.dk

* Corresponding author

ABSTRACT

ARGONAUTE (AGO) proteins bind to small non-coding RNAs to form RNA Induced Silencing Complexes (RISCs). In the RNA-bound state, AGO proteins are stable while RNA-free AGOs turn over rapidly. Molecular determinants unique to RNA-free AGO that allow its specific recognition and degradation remain unknown. Here, we show that a confined, linear region in Arabidopsis AGO1, the N-coil, is accessible to antibodies preferentially in the RNA-free state of AGO1. Reanalysis of hydrogen-deuterium exchange data on human Ago2 indicates similar structural flexibility of the N-coil depending on small RNA binding. Unloaded Arabidopsis AGO1 interacts with the autophagy cargo receptor ATI1 via direct contact to specific amino acid residues in the N-coil, and mutation of residues required for ATI1 interaction reduces the degradation rate of unloaded AGO1 *in vivo*. These results provide insight into the molecular basis for specific recognition and degradation of the RNA-free state of eukaryotic AGO proteins.

INTRODUCTION

RNA Induced Silencing Complexes (RISCs) are central components in the processing of genetic information in eukaryotes. The core of RISC is an ARGONAUTE (AGO) protein bound to a small non-coding RNA that guides RISC to complementary target RNAs by base pairing. In turn, RISCs repress gene expression sequence-specifically using a variety of mechanisms operating at either transcriptional or post-transcriptional levels. In animals, two distinct phylogenetic clades of AGO proteins exist. The Ago clade carries out endogenous gene regulation, in the main via microRNAs (miRNAs)[1], and, in particular in invertebrates, basal antiviral defense via virus-derived small interfering RNAs (siRNAs)[2]. The Piwi clade is primarily used for silencing of transposable elements in the germline via Piwi-interacting RNAs (piRNAs)[3]. Plants do not encode Piwi-clade proteins, and use Ago-clade proteins to effect transposon silencing, basal antiviral defense, and endogenous gene regulation via miRNAs and siRNAs[4]. AGO1, the founding member of the AGO protein family, has widespread functions in small RNA-guided regulation at the post-transcriptional level, and is implicated in endogenous gene regulation via most miRNAs and in antiviral defense via virus-derived siRNAs[5].

All AGO proteins consist of four separate domains called N, PAZ, MID and Piwi[6]. They fold into a two-lobed structure in the small RNA-bound state with N and PAZ together connected by the linker region L1 in one lobe and MID/Piwi in the other[7-10] (Figure 1A,B). The ends of the small RNA are tethered in dedicated binding pockets in the MID and PAZ domains such that the 5'-end is bound to MID[11,12] and the 3'-end to PAZ[13]. PAZ and MID are connected by a long linker sequence called L2 that also interacts extensively with the small RNA backbone and with other AGO domains in the small RNA-bound state [8-10] (Figure 1A,B).

The chaperone-assisted assembly of RISC by incorporation of a small RNA into

AGO, referred to as "RISC loading", is a critical event in all RNA silencing pathways. It is

rate-limiting for maturation of many miRNAs *in vivo*[14], and in both metazoans and plants, it

is decisive for the stability of AGO proteins. Small RNA-bound AGO is stable, but unloaded

AGO is rapidly degraded *in vivo*[15-17], unless it is bound to Hsp90[17,18]. Both the ubiquitin-

proteasome pathway and autophagy are involved in turnover of unloaded AGO in plants

and animals[15-20], and progress has been made on identifying molecular components

mediating such degradation. For example, the RING finger protein Iruka ubiquitinates the

fly miRNA effector Ago1 in the L2 region specifically in its unloaded form [21]. This

ubiquitination triggers binding of an adaptor complex, ultimately leading to association with

the valosin-containing protein (VCP) that mediates degradation of the ubiquitinated,

unloaded Ago1 via autophagy[20]. A clear identification of sites differentially accessible

between loaded and unloaded Ago1 was not achieved, however, and it remains possible

that Iruka recognizes unloaded Ago1 outside of the ubiquitinated L2 region. Moreover, the

Ago1-Iruka case is unlikely to be easily generalized to other systems, because the site of

ubiquitination is non-conserved, and is situated in a *Dm*Ago1-specific insertion in L2

compared to other miRNA-interacting AGO proteins (Figure S1). In arabidopsis, mutation

of the F-box protein FBW2 restores steady-state levels of a largely unloaded mutant AGO1

protein with a destabilized fold of the PAZ domain[22]. It is, however, unclear whether FBW2

targets AGO1 directly, and whether it specifically mediates degradation of unloaded, but

functional AGO1, or plays a more general role in chaperone-assisted protein degradation

elicited as a consequence of partial unfolding of the PAZ domain in the point mutant

analyzed. Thus, despite considerable progress on identifying the downstream pathways

mediating degradation of unloaded Ago proteins, the molecular determinants of unloaded

AGO that distinguish it from loaded AGO and allow its selective recognition remain

unknown.

In this study, we use arabidopsis AGO1 to define a region of crucial importance in

the distinction between eukaryotic AGO proteins in the RNA-bound and in the free state.

We show that the N-coil, a small, highly conserved linear region in the N domain, yet part

of the MID-Piwi lobe in the loaded conformation, is preferentially accessible in unloaded

arabidopsis AGO1 and human Ago2. The N-coil in arabidopsis AGO1 functions as a direct

binding site for the autophagy cargo receptor ATG8 INTERACTING PROTEIN1 (ATI1),

and mutations disrupting ATI1 binding reduce the degradation rate of unloaded AGO1 *in*

vivo. These results highlight the N-coil as a key determinant of the distinction between

loaded and unloaded AGO1, and, given its conserved structural properties, probably in

other eukaryotic AGO proteins in both Ago and Piwi clades.

RESULTS

Definition of the N-coil and other N-terminal parts of arabidopsis AGO1

Arabidopsis thaliana (At) AGO1 has a 176-aa extension at its N-terminus, in addition to the

canonical N-PAZ-MID-PIWI domains found in all AGO proteins. Because this extension is

predicted to be an intrinsically disordered region (IDR, Figure S2), we term this part of

AGO1 the N-IDR. The N-IDR is followed by a 14-aa region (F177-K190) that we previously

termed the N-coil[5]. Structurally, the N-coil is of interest, because it is part of the MID-PIWI

lobe in AGO-small RNA complexes, despite its location in the N-terminal part of AGO in

sequence. (Figure 1A-B). Furthermore, the N-coil assumes an extended conformation with

no secondary structure, yet has low structural flexibility, and reaches across the width of

the entire AGO-small RNA complex (Figure 1B). The N-coil is followed by two β-strands

that connect it to the remainder of the N-domain in the N-PAZ lobe (Figure 1B). We term

this latter part of the N-domain the globular N-domain to emphasize its globular fold, in

contrast to the N-coil.

Properties of the N-coil and a hypothesis for its function

Two additional properties of the N-coil make this part of AGO proteins attractive for further

study. First, residues in the conserved N-coil are structurally fixed by numerous inter-

domain interactions in the AGO-small RNA complex, but the N-coil/Piwi interface is only

moderately hydrophobic with several residues calculated to prefer a hydrophilic

environment (Figure 1C). For example, Arg28 in human Ago2, equivalent to Arg182 in

arabidopsis AGO1, participates in a salt bridge deep inside of the PIWI domain (Figure

1D). This may indicate that the conformation of the N-coil revealed in the crystal structures

of AGO-small RNA complexes is but one of several possible conformations of this region.

Second, specific lysine residues in the N-coil of Arabidopsis AGO1 (Lys178, Lys185 and

Lys190) are required for yeast two-hybrid interaction of the N-domain of AGO1 with factors

implicated in regulated proteolysis, including the autophagy cargo receptor ATI1 ().

Combining the indication of alternative N-coil conformations, the known instability of

unloaded AGO proteins, and the N-coil-dependent two-hybrid interactions with regulated

proteolysis factors, we hypothesize that the N-coil is accessible in the unloaded

conformation of AGO1, but becomes attached to the Piwi domain as part of the loading

process. We further hypothesize that the accessibility of the N-coil in the unloaded form is

used for its specific recognition by factors implicated in regulated proteolysis, thereby

providing a molecular explanation for the rapid turnover of unloaded AGO1 *in vivo* (Figure

1E). The remainder of this paper describes the results of tests of predictions that follow

immediately from this hypothesis for N-coil function.

The AGO1 Y691E protein mutated in the 5'-phosphate binding pocket is unloaded

Since all tests of our model require manipulation of unloaded AGO1, we first engineered

the Y691E mutant of AGO1, mutated in a key residue in the MID domain that coordinates

the 5'-phosphate of the 5'-nucleotide of the small RNA and stacks with its nucleobase[11,12],

typically a uracil. In other AGO proteins, the equivalent mutation leads to complete loss of

small RNA binding [20,23]. We used stable transgenic lines in arabidopsis to establish that

the Y691E mutant is indeed unloaded. Immunoprecipitation of wild type and Y691E mutant

protein followed by polynucleotide kinase-mediated labeling of co-purified small RNA

showed that in contrast to AGO1^{WT}, no small RNA was bound by the AGO1^{Y691E} protein

(Figure 2A). Thus, the AGO1^{Y691E} mutant protein is unloaded. Consistent with this

molecular result, expression of the AGO1^{Y691E} mutant protein in the *ago1-3* knockout

background failed to rescue the knockout mutant phenotype, demonstrating total loss of

function (Figure 2B). Identical results were obtained with the AGO1^{Y691E} protein containing

additional mutations in the N-coil and N-domain residues required for ATI1 interaction in

yeast (Figure 2A,B).

The N-coil is uniquely accessible in unloaded AGO1

We decided to test accessibility of the N-coil in small RNA-bound and free AGO1 using

antibody probing. Hence, we raised antibodies against a 16-aa N-coil peptide (F177-

C192). This polyclonal antibody specifically recognized the N-coil as demonstrated by its

reactivity towards FLAG immuno-purified, denatured AGO1 with a wild type N-coil, but

substantial loss of reactivity with FLAG-AGO1 carrying two mutations (K185E/K190E) in

the N-coil (Figure 3A-B). We next used this antibody to probe the accessibility of the N-coil

in AGO1 in two different ways. First, we observed that binding of equal amounts of N-coil

antibody required 10-fold higher amounts of immuno-purified, native, immobilized FLAG-

AGO1^{WT} (largely small RNA-bound, see below) than of similarly purified unloaded AGO1

(FLAG-AGO1^{Y691E}) (Figure 3B-C). We note that this difference cannot be explained by

potentially different patterns of post-translational modifications in the N-coil, because the

antibody recognized denatured forms of the two AGO1 variants equally efficiently (Figure

3B). Second, direct immunoprecipitation from total lysates using the N-coil antibody

resulted in markedly more efficient recovery (~10 fold) of FLAG-AGO1^{Y691E} than of FLAG-

AGO1^{WT} (Figure 3D), in contrast to the result obtained with FLAG antibodies (Figure 3B).

These observations provide clear evidence that the N-coil of AGO1 is preferentially

accessible in the unloaded state.

The N-coil-Piwi interface of human Ago2 exhibits reduced solvent exposure upon small

RNA binding

We next wished to assess whether differential N-coil accessibility applies more generally

to eukaryotic AGO proteins. To this end, we turned to recently published hydrogen-

deuterium exchange data on RNA-free and small RNA-associated human Ago2[24]. Amide

protons exchange with the solvent, and their rate of hydrogen-deuterium exchange upon

dissolution in D_2O is, therefore, a sensitive measure of solvent exposure and engagement

in hydrogen bonds[25]. Indeed, comparison of RNA-free, guide RNA-associated and

guide/target RNA-associated human Ago2 identified regions with differential hydrogen-

deuterium exchange rates between the three states[24]. Interestingly, reanalysis of this data

with an eye towards the N-coil and interacting residues in the Piwi domain in the loaded

conformation showed that regions of slower amide proton exchange in loaded versus

unloaded human Ago2 include the region of interest: the N-coil and, in particular, the

nearby hydrophilic environment in the Piwi domain caging the guanidium group of the

conserved Arg28 side chain in the RNA-bound state (Figure 3E, Figure 1A). Taken

together, the antibody probing of arabidopsis AGO1 and the differential hydrogen-

deuterium exchange rates in human Ago2 upon loading show that structural flexibility and

preferential accessibility of the N-coil in the unloaded state is a recurrent feature of

eukaryotic AGO proteins.

The interaction between ATI1 and the N-domain is direct and involves the N-coil

To test the second prediction that regulated proteolysis factors directly bind the N-coil, we

chose the interaction between a fragment of AGO1 comprising the N-coil and the globular

N-domain (NcGN) and the N-terminal IDR of the autophagy cargo receptor ATI1[26] as

model system. Using yeast two-hybrid assays, this interaction was shown to depend on

the N-coil residues Lys178, Lys185 and Lys190 (). We first expressed and purified the IDR

of ATI1[26] and wild type and K178E mutant versions of the NcGN of AGO1 fused to His_6-

SUMO (His_6-SUMO-NcGN^{AGO1}, Figure 4A), and used these purified proteins for

quantitative binding assays by microscale thermophoresis. This assay uses the distinct

thermophoretic mobility of different molecular entities to measure the percentage of

complex formed between a fluorescently labeled component, and a partner protein added

in increasing concentrations from none (0% complex) to an estimated 100-fold molar

excess (~100% complex)[27]. In our experimental set-up, His_6-SUMO-NcGN^{AGO1} (WT or

K178E) was fluorescently labeled on the unique Cys192 residue close to the N-coil, and

the unlabeled IDR of ATI1 was added in increasing quantities. Using this set-up, we

showed that the interaction between NcGN^{AGO1} and the IDR of ATI1 is direct, and

measured the dissociation constant to be 11±2 µM (Figure 4B). Remarkably, these affinity

measurements did not necessitate application of a thermal field to measure differential

molecular movement, because the fluorescence intensity itself was sensitive to addition of

the IDR of ATI1 (Figure 4C). This observation suggests that in the NcGN^{AGO1}-ATI1(IDR)

complex, the C192-linked fluorophore is located in a distinct chemical environment

compared to the free NcGN^{AGO1}, and thus, close to an interaction site. Indeed, when the

N-coil mutant K178E was analyzed in the same assay, both the affinity of the ATI1-IDR

interaction (47±12 µM), and, especially, the change of fluorescence upon complex

formation were substantially reduced (Figure 4B-C). These observations indicate that the

N-coil is a direct site of interaction between NcGN^{AGO1} and ATI1-IDR, and that it

contributes substantially to the affinity of the interaction.

Residues in the N-coil interact directly with ATI1

To show conclusively that the N-coil is a direct interaction site of ATI1-IDR, we expressed

and purified an ^{15}N-labeled N-coil peptide (172-SSSKAFKFPMRPGKGQSGKRC-192) to

perform 2D ^{1}H-^{15}N heteronuclear single quantum coherence nuclear magnetic resonance

(NMR) spectroscopy in the presence and absence of the ATI1-IDR. These experiments

showed that the signal intensities from backbone amide N-H nuclei in the N-terminal part

of the N-coil (Phe177-Lys185), and, to a lesser extent, the C-terminal Lys190, were

strongly reduced upon addition of ATI1-IDR (Figure 4D,E), indicating that these residues

participate directly in the interaction. Consistent with the importance of Lys178 for the

ATI1-IDR interaction, such loss of signal was markedly less pronounced upon addition of

ATI1-IDR to an ^{15}N-labeled N-coil peptide containing the K178E mutation, in particular for

the amide group of residue 178 (K/E) itself (Figure 4F-H). We conclude that the N-coil is a

direct interaction site of the ATI1-IDR, and that the interaction involves contacts along the

entire N-coil, including the Lys178, Lys185, and Lys190 residues.

ATI1 interacts with unloaded AGO1 in vivo and the binding implicates N-coil residues

Having established that the N-coil interacts directly with a protein implicated in

regulated proteolysis, we next wished to address if this interaction could occur in the

context of full-length AGO1, in particular in the unloaded state. Because heterologous

expression of AGO1 failed, even with the use of a variety of expression hosts (*E. coli*,

baculovirus/*Sf-9, Schizosaccharomyces pombe*) and affinity/solubilization tags, we used

bimolecular fluorescence complementation (BiFC) assays upon transient expression in

Nicotiana benthamiana for assessment of ATI1 interaction with AGO1^{WT} and AGO1^{Y691E}.

The close AGO1 paralogue, AGO10, whose NcGN part interacts much less strongly with

ATI1 in the two-hybrid assay than AGO1 (), was used as a control expected to yield

substantially less specific signal than AGO1. With AGO1^{WT} fused to the C-terminal part of

YFP and ATI1 fused to its N-terminal part (C-YFP-AGO1^{WT} and N-YFP-ATI1), weak, but

specific signal was detected in association with the endoplasmic reticulum (ER; Figure

5A), consistent with previous reports of ATI1-localization to ER-associated ATI-bodies, and

with peripheral ER-association of AGO1[28,29]. We detected a consistent and statistically

significant reduction in intensity of the BiFC signal when the C-YFP-AGO1$^{K185E/K190E}$

mutant containing substitutions in N-coil residues implicated in ATI1 binding was used

(Figure 5A-B; Figure S3). We next compared patterns of BiFC signal obtained with C-YFP-

AGO1^{Y691E}/N-YFP-ATI1 and C-YFP-AGO1$^{Y691E/K185E/K190E}$/N-YFP-ATI1. With both forms,

specific signal was detected, and, similar to what we observed with the wild type AGO1,

the K185E/K190E N-coil mutations resulted in significantly reduced BiFC signal intensity

(Figure 5A-B). These results demonstrate that ATI1 interacts with the unloaded form of

AGO1, and that N-coil residues contribute to this interaction *in vivo*. We note that complete

disruption of ATI1-AGO1 interaction is not expected upon mutation of N-coil residues,

because a previous study showed yeast two-hybrid interactions between ATI1 and AGO1

involving both N-PAZ and MID-Piwi lobes of the protein[30]. We also note that the results

show that ATI1 is capable of interacting with unloaded AGO1, and may possibly only

interact with the unloaded form, because wild type AGO1 exists as a mixed population *in*

vivo, mostly of the RNA-bound form, but also with a minor fraction in the unloaded form

(see below). Clearly, however, such possible exclusive interaction with unloaded AGO1 is

not established by the BiFC assays.

Defective ATI1 interaction sites in NcGN^{AGO1} reduce the degradation rate of unloaded

AGO1

We finally tested the relevance of residues in the NcGN^{AGO1} necessary for ATI1 interaction

for degradation of the full-length, unloaded AGO1 protein using pulse-labeling analysis. To

maximize chances of measuring an effect *in vivo*, we used the

K185E/K190E/M304E/E307A (henceforth, KKME) mutant with residues important for ATI1

interaction as identified by two-hybrid assays mutated both in the N-coil and in the globular

N-domain (). Thus, stable transgenic lines expressing equal steady-state levels of FLAG-

AGO1^{WT}, FLAG-AGO1^{Y691E}, and FLAG-AGO1$^{Y691E/KKME}$ were selected as material for

kinetic analyses. We first considered the possible kinetic pathways of newly synthesized

AGO1^{WT} and AGO1^{Y691E}. Because AGO1^{WT} can be degraded either prior to or after small

RNA loading (Figure 6A), the labeling data must be fitted to a sum of two exponentials,

with half-lives corresponding to unloaded (minor fraction, 8.5±1.8%, see Methods) and

loaded states (major fraction, 91.5±1.8%, see Methods), respectively. In contrast, labeling

of the unloaded AGO1^{Y691E} is expected to follow mono-exponential kinetics. Initial

analyses indicated that ^{35}S-Met/Cys labeling of entire seedlings resulted in a fairly uniform

distribution of label within 15 minutes (Figure S4), but that a classical pulse-chase set-up

was not feasible, because incorporation continued after addition of the chase. Since

synthesis and degradation rates are equal at steady state, we decided to do pulse-labeling

experiments and measure protein half-lives from incorporation rates of ^{35}S-label/total

AGO1 protein in FLAG immunoprecipitations following addition of ^{35}S-Met/Cys label (see

Methods). We first compared FLAG-AGO1^{WT} and FLAG-AGO1^{Y691E} lines expressing

comparable levels of FLAG-AGO1 at steady state (Figure 6B-C, Figure S5). This analysis

showed that wild type AGO1 indeed exhibits two-phase kinetic behavior. Gratifyingly, the

half-life of the rapid phase roughly matches that of unloaded AGO1 measured with the

Y691E mutant [$t_{1/2}$(WT, rapid) = 1.6±0.6 h, $t_{1/2}$(Y691E) = 2.5±0.5 h]. We therefore repeated

the pulse-labeling experiments with wild type and NcGN interaction site mutants in the

unloaded FLAG-AGO1^{Y691E} version, i.e. FLAG-AGO1^{Y691E} vs FLAG-AGO1$^{Y691E/KKME}$.

These experiments showed a modest, but reproducible reduction in the degradation rate

of the unloaded AGO1 protein with mutations in NcGN residues important for ATI1

interaction (Figure 6B,D; Figure S5; results of two independent experiments shown to

convey their quantitative variability). We note that the reduced degradation rate of FLAG-

AGO1$^{Y691E/KKME}$ compared to FLAG-AGO1^{Y691E} was not due partial loading of the FLAG-

AGO1$^{Y691E/KKME}$ protein, because small RNAs were undetectable in immunopurified

fractions of FLAG-AGO1$^{Y691E/KKME}$ and because the mutant showed complete loss of

function when expressed in *ago1-3* knockout backgrounds, indistinguishable from FLAG-

AGO1^{Y691E} (Figure 2A,B). Hence, we conclude that intact NcGN^{AGO1} interaction sites,

including the N-coil, are necessary for maximal turnover rates of unloaded AGO1 *in vivo*.

DISCUSSION

Our results demonstrate that the N-coil is exposed for interaction preferentially in the

unloaded state of AGO1. This is important, because it defines a conformational change

that accompanies the AGO loading process, and, thereby, confers distinct protein-protein

interaction properties on the small RNA-bound and free states of AGO. In turn, this

property offers a molecular explanation of the longstanding observation that unloaded

AGO proteins in plants and animals turn over much more rapidly than the small RNA-

bound form, as demonstrated here for arabidopsis AGO1. Given the structural

conservation of the N-coil in different eukaryotic AGO proteins, in particular its structural

flexibility dependent on loading status, it is likely that this same molecular principle

operates more generally to confer rapid turnover specifically on unloaded AGO proteins.

Of the many questions related to N-coil function that now call for answers, we select two

for further discussion. (1) How widespread a phenomenon across AGO proteins is the

existence of a structurally flexible, MID-Piwi associated N-coil? In particular, is there

evidence for its existence in eukaryotic Piwi-clade proteins and in prokaryotic AGO

proteins? (2) Could the N-coil have functional significance in addition to serving as a

molecular handle for specific recognition of AGO proteins in the free state?

An N-coil in Piwi-clade proteins, but not in prokaryotic Ago proteins

Piwi-clade proteins are also likely to exhibit instability in the unloaded state, because their

steady-state accumulation is reduced in the absence of cognate piRNAs[31]. Thus, it is a

question of interest whether Piwi-clade proteins have an N-coil and what properties it may

have. Structures of three Piwi-clade proteins in the piRNA-bound state (silkworm (*Bombyx*

mori) Siwi, *Drosophila melanogaster* (Dm) Piwi, and the sea sponge *Ephydatia fluviatilis*

(Ef) Piwi) have been reported[32-34]. The Siwi structure shows an N-coil (Ile130 to Gly148,

ISILRTRPEAVTSKKGTSG) associated with the MID-Piwi lobe much like in Ago-clade

structures (Figure 7A). Thus, N-coil interaction may also be used for molecular recognition

of unloaded Piwi proteins, an exciting possibility given that recognition of the unloaded

state has significance beyond regulated proteolysis for Piwi proteins. For instance, in

insect germline cells, Piwi proteins engage in a so-called ping-pong amplification cycle of

piRNAs derived from transposable elements. Ping-pong amplification entails cleavage of a

sense piRNA precursor transcript by a mature piRISC containing the Piwi protein

Aubergine (Aub) bound to a primary, anti-sense piRNA, and subsequent loading of the

sense cleavage fragment onto another Piwi-clade protein, Ago3[35-37]. Ago3 is precluded

from primary piRNA association by the protein Krimper that associates with Ago3

specifically in the free state[38]. The molecular determinants of the recognition of the

unloaded state of Ago3 by Krimper may include unmethylated Arg residues in the N-

terminal IDR of Ago3[38], but it is now a plausible possibility that the N-coil of Ago3 is an

additional such determinant. In contrast, structures of prokaryotic AGO proteins in the

nucleic acid-bound state do not show indication of an N-coil[39-42]. Indeed, in prokaryotic

AGO proteins, the area in the Piwi domain corresponding to the one surrounding the

conserved N-coil residue Arg28 in human Ago2 is markedly different from eukaryotic AGO

proteins. In eukaryotic AGO proteins, this region appears optimized for existence in

alternative states depending on N-coil/Piwi interaction, because a loop containing the

conserved His742 and Asp747 in *Hs*Ago2 is held in place by the buried Arg28 N-coil

residue. We note that this conformation also involves a characteristic cluster of His

residues (Figure 7B) whose degree of protonation may favor one state over the other. In

contrast, in prokaryotic AGO proteins, this region is structurally fixed (Figure 7C),

consistent with the absence of a flexible, Piwi-interacting N-coil.

Relevance of N-coil-Piwi interactions for the mature RISC

While accessibility of the N-coil for protein-protein interaction in the unloaded state can be

exploited for specific recognition of the unloaded form, it is also relevant to consider

possible functions of the N-coil in the mature RISC. Since the N-coil-Piwi interactions seen

in structures of small RNA-bound AGO proteins occur exclusively in the loaded

conformation as judged by the antibody probing results of arabidopsis AGO1 and

differential hydrogen-deuterium exchange rates in human Ago2, it is possible that N-coil-

Piwi interactions are important to reach the compact conformation of mature RISC. Two

observations are compatible with this proposition. First, the N-coil is clearly important for

RISC function because two point mutations in this region have been recovered in forward

screens for defective miRNA function in arabidopsis: *ago1-38* (G186R) that exhibits

relatively weak loss of function[43], and *ago1-55* (G189E) that exhibits stronger loss of

AGO1 function[44]. The exact molecular defects of these point mutants have not been

defined, however, and it remains possible that RISC activity rather than loading could be

affected in these mutants. Second, closer inspection of the available Piwi-clade structures

reveals intriguing differences. While the Siwi protein analyzed has a full N-coil as

discussed above, *Ef* Piwi and *Dm* Piwi harbor N-terminal truncations, presumably due to

proteolytic cleavage prior to structural characterization[33,34], such that the analyzed *Ef* Piwi

protein has a partial N-coil, and *Dm* Piwi no N-coil at all (Figure 7A). Strikingly, the

compactness of the Piwi:piRNA complexes increases progressively from the *Dm* Piwi

structure (no N-coil present) to the Siwi structure (full N-coil present) (Figure 7A,D),

lending support to the idea that the N-coil/MID-Piwi lobe association may be an important

step in reaching the final, compact conformation of mature RISC across both Piwi and Ago

clades. Finally, we note that a key functional importance of the N-coil and the β-strands

connecting it to the rest of the N domain has been observed in human Ago proteins where

substitution of this part of the catalytically inactive human Ago3 by the corresponding part

of the catalytically active Ago2 is sufficient to confer activity on Ago3[45]. This observation

suggests that the relevance of the N-coil interactions with the Piwi domain may not be

limited to the loading process, but may also be important for the extensive conformational

changes required to reach the catalytically active conformation of AGO proteins[46].

MATERIALS AND METHODS

Plant material and growth conditions

The Columbia-0 (Col-0) *Arabidopsis thaliana* accessions were used. *ago1-3* was

described previously[47]. Seeds were sterilized as described in [48] prior to being sown on

Murashige-Skoog (MS) agar medium (4.3 g/L MS salts, 0.8 % agar, 1% sucrose) where

they were grown for 10 days at constant temperature (21 °C) and a 16h light (80 µE m^{-2} s^{-1})/8h darkness cycle.

Antibodies

Anti-FLAG® M2 Affinity Gel and monoclonal anti-FLAG® M2-Peroxidase (HRP) were

purchased from Sigma, anti-HSP70 from Agrisera, and anti-Rabbit IgG (HRP) from Sigma.

The anti-AGO1-N-coil antibody was affinity purified by Eurogentec from sera collected

from a rabbit immunized with a 16-amino acid peptide from the N-coil of AGO1 (F177-

C192, H-FKFPMRPGKGQSGKRC-NH$_2$), synthesized by Schafer-N, Copenhagen,

Denmark. The cysteine corresponding to C192 in AGO1 was used for KLH coupling prior

to immunization.

Site-directed mutagenesis

Site-directed mutagenesis was performed according to the Quickchange protocol

(Stratagene) using Phusion High-Fidelity DNA Polymerase (NEB) and primers containing

the appropriate mutations (Supplemental Table 2). The reaction mix was subsequently

digested with DpnI (Thermo Scientific) to remove methylated template DNA before

transformation to DH5α competent *E. coli* (NEB).

Plant transformation and selection

Transgenic lines were generated by floral dip transformation with *Agrobacterium*

tumefaciens GV3101[49]. The FLAG-AGO1^{Y691E} and FLAG-AGO1$^{KKME/Y691E}$

(K185E/K190E/M304E/E307A/Y691E) mutations were introduced to pCAMBIA3300U

AGO1P:FLAG-AGO1-AGO1T previously described in [50] using site-directed mutagenesis

(primers listed in Supplemental Table 2). FLAG-AGO1^{Y691E} and FLAG-AGO1$^{KKME/Y691E}$

were transformed into *ago1-3/+* lines. Primary transformants (T1) were selected on MS

agar medium supplemented with 100 µg/ml ampicillin and 7.5 µg/ml glufosinate

ammonium (Fluka). Transformed plants were transferred to soil after approximately 10

days of growth on selective medium. For every construct, lines with a single T-DNA locus

and comparable protein expression levels were selected in T2, and descendants

homozygous for the insertion were identified in T3.

Genotyping

Genomic DNA from aerial tissue of rosette leaves was genotyped using Phire Plant Direct

PCR Kit (Thermofischer). The primers used for genotyping are listed in Supplemental

Table 1.

PNK labeling of AGO1 bound sRNA

RNA was extracted directly from FLAG IP beads by addition of TRI-Reagent (Sigma) as

described above except that RNA precipitation was performed in the presence of 20 µg of

glycogen. The RNA was 5'-end labeled with [γ-^{32}P]-ATP by T4-polynucleotide kinase

according to manufacturer (Fermentas). Unbound [γ-^{32}P]-ATP was removed by MicroSpin

G-25 columns (GE Healthcare). The RNA was separated by electrophoresis in a

denaturing 18 % polyacrylamide gel. The gel was exposed to a phosphor screen and

imaged on a Typhoon FLA 7000 scanner (GE Healthcare).

Protein expression and purification

Purification of ATI1N M1-E180 was carried out as previously described [26]. AGO1 NcGN

(S172-Y335) CDS was cloned into a modified pET24a vector encoding His$_6$-SUMO N-

terminally. The K178E substitution was generated by site-directed mutagenesis. An

additional three substitutions (L318A/L321A/L325S) were made by site-directed

mutagenesis to enhance solubility (primers listed in Supplemental Table 1). Plasmids of

His$_6$-SUMO-tagged AGO1 NcGN and His$_6$-SUMO-AGO1 NcGN^{K178E} were transformed into

Escherichia coli BL21 (DE3 7tRNA) Rosetta cells and grown over night at 37 °C in 50 ml

LB media supplemented with 50 µg/ml kanamycin and 33 µg/ml chloramphenicol. 2 ml of

culture were added to 1 liter LB medium supplemented with 50 µg/ml kanamycin and 33

µg/ml chloramphenicol and grown at 37°C, 150 rpm to an OD$_{600}$ of 0.6-0.8 before induced

with 0.5 mM isopropyl β-D-1-thiogalactopyranoside (IPTG) for expression o/n at 18°C, 150

rpm. Cells were harvested by centrifugation at 8,000 x g for 15 min, 4 °C and resuspended

in 25 ml cold lysis buffer (25 mM tris-HCl, 10 mM imidazole, 300 mM NaCl pH 8, 1mM

TCEP) supplemented with 1 tablet complete protease inhibitor EDTA-free (Roche) before

lysed using a French Press (>10.000 psi). The cell lysate was cleared by centrifugation at

30,000 x g for 30 min, 4 °C and the supernatant was passed through a 0.45 µm filter

before incubated on 2.5 ml Ni-NTA agarose beads for 1 h at 4°C. The bead suspension

was transferred to an empty gravity column and washed with 50 ml 25mM Tris-HCl, 20mM

imidazole, 300mM NaCl pH 8, then eluted in 5 ml 25mM Tris-HCl, 300mM imidazole,

150mM NaCl pH 8 and subsequently dialyzed o/n against buffer depleted for imidazole (20

mM Tris-HCl pH 8, 150 mM NaCl, 1 mM β-ME). His$_6$-SUMO-AGO1 NcGN was further

purified by application on a Superdex 200 16 600 (GE Healthcare) size exclusion

chromatographic column connected to an Äkta Purifier system (GE Healthcare) in 20 mM

Tris-HCl pH 7.2, 1 mM KCl, 0.05 mM NaCl, 0.01 mM MgCl$_2$, 1 mM TCEP. Eluting protein

was detected by measurement of absorbance at 280 nm, and fractions of 1 ml were

collected and analyzed by SDS-PAGE. Fractions containing His-SUMO-AGO1 were kept

at 4°C and used fresh.

For expression of AGO1 N-coil, a cDNA encoding residues S172-C192 was cloned into a

pGEX4t1 vector encoding a GST-tag (primers listed in Supplemental Table 1) and

transformed into *Escherichia coli* BL21 (DE3 7tRNA) Rosetta cells. For production of the

stable isotope-labeled AGO1 N-coil, a culture of the expression strain was added to 1 L of

freshly prepared M9 minimal media (22mM KH_2PO_4, 42mM $Na_2HPO_4 \cdot 2H2O$, 17mM NaCl,

1mM $MgSO_4$) supplemented with 1:1000 M2 trace element solution, 0.4 % (v/v) glucose,

18 mM NH_4Cl-^{15}N and 100 μg/ml ampicillin and 33 μg/ml chloramphenicol and grown to

OD_{600} of 0.6-0.8 before induction with 1mM IPTG for overnight expression at 18°C. Cells

were harvested by centrifugation at 8,000 g 15 min at 4 °C and lysed by 2 passages

through a French press (>10,000 psi) in ice cold 1x PBS (137 mM NaCl, 2.7 mM KCl, 10

mM Na_2HPO_4, and 1.8 mM KH_2PO_4, pH 7.4) with 1 mM TCEP and 1 tablet complete

EDTA-free protease inhibitor (Roche). The lysate was cleared by centrifugation at 30,000

g for 30 min at 4 °C and subsequently filtered through a 0.45 μm filter before incubation

with 2.5 ml glutathione-conjugated sepharose beads (GE Healthcare) for 1h at 18 °C. The

beads were packed in an empty gravity column and washed with 50 ml cold 1xPBS. AGO1

N-coil was liberated from the GST-tag by on-bead cleavage o/n at 4 °C with His_6-TEV in

estimated 1:50 molar ratio in 50mM Tris-HCl, 50mM NaCl, 0.5mM EDTA, 1.75mM TCEP,

pH 7.2. The flow-through was collected and His_6-TEV was captured on 0.1 ml Ni-NTA

agarose beads for 1h at 4°C. AGO1 N-coil was further purified by reversed-phase

chromatography using a 17.35ml Zorbax 300 Å Stable-bond C18 column connected to an

Äkta Purifier HPLC system. Trifluoroacetic acid (TFA) was added to the AGO1 sample to a

final concentration of 0.1% and centrifuged at 20,000 g for 15 min to remove insoluble

aggregates prior to application. The column was equilibrated with 100% buffer A (0.1%

TFA [v/v] in MQ water) and AGO1 N-coil was eluted by a gradient into 100% buffer B (70

% [v/v] acetonitrile, 0.1% TFA [v/v] in MQ water) over 10 CV using a flow rate of 3 ml/min.

2 ml fractions were collected by A_{220} detection. AGO1-containing fractions were collected,

flash-frozen in nitrogen and subsequently lyophilized overnight. The samples were

resuspended in milli-Q water, adjusted to pH 6.6, before again being flash-frozen and

lyophilized o/n and stored at -20 °C. Aliquots of AGO1 N-coil-containing fractions were pH-

adjusted by the addition of TA30 (30:70 [v/v] ACN:0.1%TFA) and verified by MALDI-TOF

mass spectrometry on an α-cyano-4-hydroxycinnamic acid (HCCA)-matrix.

Microscale thermophoresis (MST)

All MST data was recorded on a Monolith NT.115 (NanoTemper Technologies) at 25 °C

with medium MST-power exciting the fluorophore in the red excitation channel. The buffer

composition was 20mM tris-HCl pH 7.2, 1mM KCl, 0.05mM NaCl, 0.01mM $MgCl_2$, 1mM

TCEP, 0.05 % [v/v] tween20. 20 µM His_6-SUMO-AGO1-NcGN was orthogonally labeled on

Cystein-192 using the Monolith NT^{TM} Protein Labeling Kit Red – Maleimide (NanoTemper

Technologies). The labeling was performed according to the manufacturer's instructions.

From a stock solution of 271 µM ATI1 IDR, a 1:1 dilution series was prepared and

subsequently mixed 1:1 with SUMO-AGO1 NcGN keeping the concentration and hence

the fluorescence constant. The protein series was loaded into Monolith NT.115 Premium

Capillaries (NanoTemper Technologies) and thermophoresis was measured using

MO.Control v1.6.1 software. The raw data recorded at 0.5-1 sec of thermophoresis was

retrieved and analyzed with GraphPadPrism 8.0 fitting by the law-of-mass-action

$Y = F_0 + \frac{F_{max} \times X}{(K_d + X)}$

where Y is the fluorescence at binding, F_0 is the fluorescence when no complex has

formed, F_{max} is the fluorescence at maximum binding, X is the concentration of ATI1 IDR,

and K_d is the binding constant. All fluorescence intensities were normalized to the initial

fluorescence prior to fitting.

NMR spectroscopy

NMR spectra were recorded on Bruker Avance III 600 MHz or 750 MHz spectrometer

equipped with TCI cryoprobes. Spectra were recorded at 5 °C in buffer composition 20mM

Tris-HCl pH 6.6, 100mM KCl, 5mM NaCl, 1mM $MgCl_2$, 3mM TCEP, 10% (v/v) D_2O, 250

µM sodium 2,2-dimethyl-2-silapentane-5-sulfonate (DSS). For titration experiments, [1]H-

[15]N-HSQC spectra were recorded of 50 µM [15]N-labeled AGO1 N-coil in the absence or

presence of increasing concentrations of up to 265 µM ATI1 IDR. The intensity ratios of

178K and 178E for different concentrations of ATI1 IDR were fitted as an exponential

function in GraphPadPrism 8.0

$$Y = e^{-K*X}$$

Assignment of NMR spectra

Assignment spectra were recorded on a Varian Inova 750 MHz magnet equipped with a

Bruker Avance III console and a Bruker TCI cryoprobe. Backbone amide resonances of

free 1.4 mM AGO1 N-coil and 4.5 mM AGO1 N-coil[K178E] were assigned manually in

CCPNmr Analysis 2.2.4 [51] using [15]N-TOCSY-HSQC and [15]N-NOESY-HSQC spectra [52-55].

In vitro binding assay of AGO1 variants to the N-coil antibody

1.6 g of inflorescence tissue from the indicated genotypes were ground in liquid nitrogen

and extracts were prepared in 6.4 ml IP buffer (50 mM Tris-HCl, pH 7.5, 150 mM NaCl,

10% glycerol, 5 mM $MgCl_2$, and 0.1% Nonidet P-40, 4 mM DTT) supplemented with 2x

Complete EDTA free protease inhibitor (Roche). Samples were spun at 30,000 *g* for 10

min at 4°C and the cleared lysate was incubated with 40 µl anti-FLAG agarose bead slurry

(Sigma) for 30 min at 4°C. The beads were washed three times in IP buffer and divided

into four Eppendorf tubes for each genotype. The anti-FLAG beads were incubated with

50 µl of IP buffer containing the indicated concentrations of anti-AGO1N antibody for 20

min at 4°C. The beads were washed three times in IP buffer and eluted twice in 20 µl IP

buffer with 100 ng/µl FLAG-peptide (Sigma) for 15 min at 4°C. The samples were heated

for 5 min at 75°C in Laemmli buffer, subjected to SDS-PAGE and blotted onto a

nitrocellulose membrane. The FLAG-AGO1 proteins were detected with anti-FLAG-HRP

(1:4000 [Sigma] in 5 % milk in PBS-T [0.05% tween20]) or anti-AGO1-N-coil (1:1000 in 5%

milk in PBS-T) followed by anti-Rabbit antibody (1:10,000 [Sigma] in 5% milk in PBS-T).

The anti-AGO1-N-coil antibody was detected with anti-Rabbit antibody conjugated to HRP

(1:2,000 [Sigma] in 5% milk in PBS-T). The intensity of the signal from retained anti-

AGO1-N-coil antibody was quantified using Image J and fitted using Nonlinear Least

Square (NLS) in R (version 4.2.0) with an exponential equation:

$Y = a * exp(b * conc)$

Immunoprecipitation using the AGO1 N-coil antibody

30 µl pf Protein A-agarose beads (Sigma) were washed three times in PBS and incubated

with 500 µl PBS with 7.5 µg of anti-AGO1-Ncoil antibody (1:20 dilution of the fraction

affinity purified against the N-coil peptide) for one hour at 4°C and then washed in PBS. In

parallel, 200 mg of inflorescence tissue of each genotype were ground in liquid nitrogen

and extracts were prepared in 800 µl IP buffer supplemented with 2x Complete EDTA free

protease inhibitor (Roche). Samples were spun at 17,000 *g* for 10 min at 4°C and

transferred to a new tube. 15 µl of Agarose A beads with anti-AGO1-Ncoil antibody were

added to the samples and incubated for 30 min at 4°C. The beads were washed three

times in 500 µl IP buffer and heated for 5 min in Laemmli. After SDS-PAGE and blotting

onto a nitrocellulose membrane, the FLAG-AGO1 proteins were detected with anti-FLAG-

HRP antibody (1:4000 [Sigma] in 5 % milk in PBS-T). The membrane was developed as

described above with homemade ECL or SuperSignal™ West Femto Maximum Sensitivity

Substrate (Thermo Fisher) and images taken with a Sony A7S camera.

Bimolecular Fluorescence Complementation (BiFC) assays

BiFC was done using the 2in1 vector system which expressed both the YFPn and YFPc

tagged protein as well as free RFP from the same vector[56]. ATI1 was amplified from oligo

(dT) primed arabidopsis cDNA and subcloned into pDONR221 P3P2 entry vector with the

BP clonase II kit (Invitrogen) and similarly, AGO10 and AGO1 were subcloned into

pDONR221 P1P4 (primers listed in Supplemental Table 1). The $AGO1^{K185E/K190E}$ and

$AGO1^{Y691E}$ mutations were introduced using site directed mutagenesis (primers listed in

Supplemental Table 1). Transfer of the Gateway cassette from these two entry vectors to

the pBIFCt-2in1-NN destination vector [56] was performed through LR clonase II kit

(Agilent).

Agrobacterium tumefaciens transformed with the pBIFCt-2in1-NN vector[57] containing the

YFPn- and YFPc-fused cDNAs were infiltrated at an OD_{600} of 0.5 into leaves of *Nicotiana*

benthamiana defective in co-suppression achieved by RNAi of *RDR6*[58]. Three days post

infiltration, epidermal cells were observed for fluorescence emission using a Zeiss LSM700

confocal laser scanning microscope with a C-Apochromat 63x and a 20x objective. YFP

and RFP were excited using laser light of 488 nm 555, respectively. The emitted light was

captured with preset filters of the microscope software. All images are shown as z-stack

projections of 7-9 slices covering 18-23 μm. Quantification was done on eight z-stack

images from individual leaf discs of each construct. The YFP and RFP signal from distinct

interaction sites were quantified using Photoshop and the ratio was plotted on a Log2

scale. Log2 transformation was done to obtain normal distribution for the differences in

YFP/RFP ratios to be tested statistically using one way ANOVA followed by post hoc

Tukey HSD multiple comparison.

Pulse labeling

Seedlings were grown for 7 days on MS agar under a 16h light/8h darkness light regime.

For each time point, 8 seedlings were transferred to 1.5 ml liquid MS in a 12-well culture

plate and grown for two more days. EasyTag™ Express35S Protein Labeling-Mix (Perkin

Elmer) was then added to a final count of 50 µCi/ml and at given time points, seedlings

were briefly washed in water and flash frozen in liquid nitrogen. The tissue was ground to a

fine powder before it was solubilized in 500 µl IP buffer (50 mM Tris-HCl pH 7.5, 150mM

NaCl, 10 % glycerol [v/v], 5mM $MgCl_2$, 0.1% Nonidet P40 [v/v], 4mM DTT, 2 tablets

complete EDTA-free protease inhibitor (Roche)) followed by 5 min incubation at 4°C. The

lysate was cleared by centrifugation at 16,000 *g* for 5 min at 4 °C before it was

immobilized on anti-FLAG (Sigma) agarose bead slurry for 30 min at 4 °C. The beads

were transferred to an empty gravity column and washed with 1.5 ml IP buffer.

Immunoprecipitated protein was released from the beads in Laemmli buffer by heating at

75 °C for 5 min.

Pulse quantification

After Western blot development, the nitrocellulose membranes were exposed to a storage

phosphor screen and the screen was imaged using a Typhoon FLA 7000 scanner (GE

Healthcare). Total lysate and IP Western blot and [35]S signal were quantified using Image J

and ^{35}S signal/IP Western blot signal were plotted against pulse time. Due to the unstable

nature of the unloaded FLAG-AGO1^{Y691E} *in vitro*, the radioactive signals of the degradation

fragments are added to the signal of the intact FLAG-AGO1^{Y691E} protein signal, except for

a band identified as HSP70. For all genotypes except FLAG-AGO1^{WT} the pulse data was

fitted to an equation of the type,

$$y = a - \left(\frac{a}{2^{\frac{time}{t_{1/2}}}} \right)$$

where $t_{1/2}$ is the half-life of the immuno-precipitated protein and a is the maximum ^{35}S

value of radioactivity per protein that the sample will converge to when the pulse has

reached equilibrium, i.e. when all unlabeled protein has been degraded and the newly

synthesized proteins are radio-labelled. Subsequently, the pulse values were adjusted to

converge to the value 100 to facilitate comparison between samples and experiments and

the data refitted to the equation,

$$y = 100 - \left(\frac{100}{2^{\frac{time}{t_{1/2}}}} \right)$$

The degradation of FLAG-AGO1^{WT} can be described as a sum of two exponential decays;

one for the unloaded fraction and one for the small RNA loaded fraction. Since the steady

state protein levels of FLAG-AGO1^{WT} and FLAG-AGO1^{Y691E} and FLAG-AGO1$^{Y691E/KKME}$

are similar, the maximum ^{35}S/IP value of FLAG-AGO1^{WT} will converge to the same value

as the FLAG-AGO1^{Y691E} and FLAG-AGO1$^{Y691E/KKME}$. Therefore, the FLAG-AGO1^{WT} pulse

data can be fitted to an equation of the type,

$$y = a - \left(\left(\frac{a}{2^{\frac{time}{t_{1/2}U}}} \right) + \left(\frac{100 - a}{2^{\frac{time}{t_{1/2}L}}} \right) \right)$$

where *a* is the percentage of unloaded AGO1 at steady state, $t_{1/2}U$ is the half-life of the

unloaded fraction and $t_{1/2}L$ is the half-life of the loaded fraction. The data analysis was

done using Nonlinear Least Square (NLS) in R (version 4.2.0).

Structural analyses

The alignment of N-coil residues between human Ago2 and plant AGO1 did not require

alignment tools. The intra-molecular interactions between the N-coil and the remaining part

of *Hs*Ago2 was calculated using the web-service PDBe PISA at the European

Bioinformatics Institute (http://www.ebi.ac.uk/pdbe/prot_int/pistart.html)[59] by treating the N-

coil as a separate polypeptide chain. The calculations were done using the human Ago2

structure (PDB 4F3T) and buried surface area (Å) and solvation energy estimates

(kcal/mol) are presented for each N-coil residue.

The Fractional Uptake Difference in hydrogen/deuterium exchange between empty and

guide RNA-bound human Ago2 was presented on the human Ago2 structure (PDB 4F3T).

For this a Pymol script was generated using the available data set [24] (MSV000088648 at

https://massive.ucsd.edu) at the 5 minute timepoint and the DECA software ver1.12 [60]. All

the presented structural models were made using CCP4mg [61].

DECLARATION OF INTERESTS

The authors declare no competing interests.

ACKNOWLEDGEMENTS

Theo Bølsterli and his team are thanked for plant care. Signe A. Sjørup is thanked for

assistance with MALDI-TOF experiments and purification of proteins. We thank Laura

Arribas-Hernández and Diego López-Marquez for critical reading of the manuscript. This

work was supported by a project grant (NNF17OC0029194) from the Novo Nordisk

Foundation, and a Consolidator Grant (ERC-2016-CoG 726417 PATHORISC) from the

European Research Council, all to PB. The work was also supported by the Novo Nordisk

Foundation Challenge program (REPIN; grant number NNF18OC0033926 to BBK). The

NMR spectra were recorded at cOpenNMR, an infrastructure initiative supported by the

Novo Nordisk Foundation grant number NNF18OC0032996.

AUTHOR CONTRIBUTIONS

SB constructed plant lines expressing unloaded FLAG-AGO1 and its combinations with N-

coil mutants in *ago1-3* knockout backgrounds, conducted bimolecular fluorescence

complementation analyses, pulse-labeling experiments and resulting data analysis,

developed the AGO1 N-coil antibody and conducted binding experiments and subsequent

data analysis. IMZS expressed and purified recombinant proteins, conducted MST assays

with SB and NMR experiments with AP, and participated with SB in preliminary tests and

AGO1 binding experiments with a lower-affinity N-coil antibody not used in the final

experiments reported here. CP conducted analysis of structures and suggested that the N-

coil may adopt conformations different from the one seen in crystal structures of *Hs*Ago2.

BBK helped design NMR experiments, and, together with AP, supervised all work and data

analysis related to those experiments. PB designed and supervised the study and wrote

the manuscript with contributions from all authors.

FIGURE LEGENDS

Figure 1. Structural properties and conserved interactions of the N-coil in Hs-Ago2

(A)-(B) Overview of the *Hs*Ago2 domain structure with surface outline of the globular N

domain and the N-coil. (A) "side view"; (B) "bottom view".

(C) Intramolecular interactions between the N-coil and the remaining part of *Hs*Ago2 as

calculated by the Protein Interfaces, Surfaces and Assemblies service PISA[59]. The

solvation energy gain of the N-coil is calculated as the difference between free energy of

the Piwi-associated and -dissociated N-coil. A negative solvation energy, Δ_iG, of a residue

makes a positive contribution to the solvation energy gain of the interface, thus favoring

the dissociated state. The color gradient (blue to red) indicates the effect sizes of buried

surface area and solvation energy and highlights the unique properties of Arg-28 in

*Hs*Ago2.

(D) Inlet showing conserved residues and bonds between a buried arginine side chain in

the N-coil (Arg-28 in *Hs*Ago2) and nearby residues.

(E) Model illustrating a hypothetical function of the N-coil in distinguishing loaded from

unloaded AGO protein and in conferring rapid proteolysis on unloaded AGO. Left, in the

loaded form, the N-coil is inaccessible for protein-protein interaction due to its association

with the Piwi domain. In the unloaded form (middle and right), the N-coil is accessible for

interactors, leading to rapid proteolysis (right) that can be slowed down if N-coil-dependent

interactions are precluded by mutation (middle).

Presentations and PISA calculations were made using the crystal structure coordinates of

the miR20a-bound *Hs*Ago2 (PDB 4F3T)[10].

Figure 2. AGO1[Y691E] does not associate with small RNA.

(A) Immunoblot analysis with α-FLAG antibody of total protein and α-FLAG

immunoprecipitated protein from inflorescence tissue from the indicated genotypes. The

lower panel shows polynucleotide kinase labeling of RNA extracted from α-FLAG IPs.

Ponceau stain shows total protein. Y, Y691E; KKME, K185E/K190E/M304E/E307A.

Numbers (Y-2, Y-5, KKME/Y-2, KKME/Y-6) denote independent transgenic lines.

(B) Images 8 days after germination of homozygous *ago1-3* (null) plants and transgenic

plant lines expressing FLAG-AGO1^{Y691E} or FLAG-AGO1$^{KKME/Y691E}$ in the *ago1-3*

homozygous background.

703 Figure 3. Preferential accessibility of the N-coil in the unloaded state of AGO

704 proteins

(A) Schematic representation of the *in vitro* binding experiment shown in (B) and (C).

(B) Protein blots of identical amounts of immobilized, FLAG immuno-affinity purified FLAG-

AGO1 variants (WT, Y691E, KKME/Y691E) incubated with the indicated concentrations of

AGO1 N-coil antibody (α-AGO1-N) and competitively eluted after extensive washing. Top

panel, detection with FLAG antibodies (α-FLAG) to document equal amounts of FLAG-

AGO1 variants in each of the binding assays. Middle panel, detection with α-AGO1-N to

document specificity of the antibody (compare AGO1^{Y691E} (intact N-coil) to

AGO1$^{Y691E/}$KKME (two point mutations (K185E/K190E) in N-coil) and equal reactivity

towards loaded and unloaded AGO1 in the denatured state. Bottom panel, detection with

anti-rabbit IgG to quantify the amount of α-AGO1-N retained by the immobilized FLAG-

AGO1 variants. KKME, K185E/K190E/M304E/E307A.

(C) Quantification of the amount of α-rabbit IgG signal in (B) at different α-AGO1-N

concentrations. The lines show an exponential fit to the data as a trendline.

(D) Immunoprecipitation of FLAG-AGO1^{WT} and FLAG-AGO^{Y691E} with α-AGO1-N antibody

from total lysates of plants expressing comparable levels of FLAG-AGO1^{WT} and FLAG-

AGO^{Y691E}. The Ponceau stain of the lanes containing aliquots of total lysates shows a crop

around the large subunit of RuBisCO.

(E) Projection onto the *Hs*Ago2 structure of the published fractional differential hydrogen-

deuterium exchange (HDX) between unloaded and guide loaded hAgo2[24]. Increasing

differences in exchange rate are indicated with progressively darker purple shading. All

regions with differential rates of exchange between unloaded and guide-loaded *Hs*Ago2

show higher exchange rate in the unloaded state, most notably areas buried by the N-coil

and engaging in interactions with N-coil residue Arg28 in the guide-loaded *Hs*Ago2.

Figure 4. The N-coil is a direct binding site for ATI1 *in vitro*

(A) SDS-PAGE of heterologously expressed His$_6$-SUMO-NcGN^{AGO1} protein used for

fluorescence labeling and microscale thermophoresis (MST) assays.

(B) MST measurements of fluorescence intensity of His$_6$-SUMO-NcGN^{AGO1} WT (left) and

K178E (right) after application of a temperature gradient. Fluorescence data obtained from

the thermophoretic acquisition period were normalized to the initial fluorescence before

being fitted to a one-site binding curve with increasing concentrations of ATI1-IDR. The

reported binding constants (K$_D$) are calculated with standard errors.

(C) Simple fluorescence intensity of Cys192-labeled His$_6$-SUMO-NcGN^{AGO1} WT (left) and

K178E (right) as a function of the concentration of ATI1-IDR.

(D, F) ^{1}H,^{15}N-HSQC nuclear magnetic resonance spectra of 50 µM of ^{15}N-labeled AGO1

N-coil WT (left, black) or K178E peptide (right, black), with 19 µM ATI1-IDR added (green

with N-coil WT, blue with N-coil K178E). Zooms of contour levels of residues K178 and

C192 are highlighted in the boxes. Minor peaks were assigned to a C-terminally truncated

version of the AGO1 N-coil peptide whose presence in the sample was verified by MALDI-

TOF mass spectrometry.

(E, G) Peak intensity plots of ^{15}N-labeled AGO1 N-coil peptide (50 µM) upon titration of

ATI1-IDR. Peak intensities are normalized to peak intensities observed with free AGO1 N-

coil peptide.

(H) Peak intensity of residues K178 (for the AGO1 wild type N-coil, green) and E178 (for

the K178E N-coil mutant, blue) as a function of the ATI1-IDR concentration. The data

points are fitted to an exponential function as a trendline.

Figure 5. Unloaded AGO1 interacts with ATI1 *in vivo*

(A) Confocal images showing results of bimolecular fluorescence complementation (BiFC)

assays to probe the AGO1-ATI1 interaction *in vivo*. The C-terminal half of YFP was fused

to AGO10, or to wild type or unloaded (Y691E) AGO1 with either an intact or a mutated

(K185E/K190E) N-coil. The N-terminal half of YFP was fused to ATI1. Both fused halves of

YFP were expressed from the same plasmid also expressing free RFP. For each BiFC

pair, possible interaction is detected by fluorescence from reconstituted YFP. RFP

fluorescence serves as a control for transformation. Merged channel images demonstrate

the presence of yellow fluorescence only in transformed cells. Scale bar, 50 µm.

(B) Quantification of YFP signal normalized to RFP signal in the given number of YFP-

positive cells for the interactions shown in (A). Letters indicate groups with statistically

significantly different $\log_2$(YFP/RFP) ratios ($p < 0.01$; one-way ANOVA with post-hoc

Tukey Honestly Significant Difference test). See Figure S3 for image data underlying the

quantification in (B). KK, K185E/K190E.

Figure 6. An intact AGO1 N-coil is required for maximal turnover rate of unloaded AGO1 *in vivo*

(A) Kinetic model of the two different fates of newly synthesized AGO1: direct degradation (κ_1), or small RNA loading (κ_2) followed by degradation (κ_3).

(B) Quantification of [35]S-Met/Cys incorporation into FLAG-AGO1[WT], FLAG-AGO1[Y691E] (FLAG-AGO1[Y]) or FLAG-AGO1[KKME/Y691E] (FLAG-AGO1 [KKME/Y]) as a function of pulse time. The lines represent a fit to an exponential equation for unloadable AGO1 containing the Y691E mutation and a fit to a double exponential equation for the WT AGO1. The half-lives calculated by the fitted models are given in hours +/- standard deviation, as detailed in Methods. KKME, K185E/K190E/M304E/E307A.

(C) Zoom in of the dashed area of (B).

(D) Result of an independent experiment done as in (C) to verify the slower incorporation kinetics in FLAG-AGO1[KKME/Y691E] compared to FLAG-AGO1[Y691E]. See Figure S5 for raw data underlying the graphs in Figure 6B-D.

Figure 7. Piwi-clade, but not prokaryotic, AGO proteins have a structurally flexible N-coil

(A) Structural overview of *Hs*Ago2 and Piwi proteins, outlining their N-coils in pink. The *Ephydatia fluviatilis* and *Drosophila melanogaster* Piwi proteins harbour N-coil truncations, probably generated by proteolysis during sample preparation for structural studies[33,34]. The structural representations are based on the published coordinates (*Hs*Ago2, PDB: 4F3T[10]; *Bm*Siwi PDB: 5GUH[32]; *Ef*Piwi, PDB: 7KX7[34]; *Dm*Piwi PDB: 6KR6[33]).

(B) Close-up of region in the Piwi domain in eukaryotic *Hs*Ago2 (PDB: 4F3T[10]) that cages the buried N-coil Arg28 side chain in the small RNA-bound conformation. Contacts

between Arg28 and the loop containing the conserved His742, Glu745 and Asp747

residues are shown by dashed lines. A cluster of histidines formed by residues in β-

strands of the Piwi-domain (His764, His766) and in the Arg28-interacting loop (His742) is

highlighted in blue.

(C) Close-up of the region in the Piwi domain of the prokaryotic *Mj*Ago (PDB: 5G5T[42]) that

corresponds to the region of *Hs*Ago2 shown in (B).

(D)   Secondary structure matching of the *B. mori* and *D. melanogaster* Piwi domains

reveals a more relaxed state of the *D. melanogaster* Piwi molecule (green) in which the

structurally fixed N-coil seen in *B. mori* Siwi is absent.

DATA AND CODE AVAILABILITY

NMR data have been deposited at the Biological Magnetic Resonance Bank (BMRB)

under BMRB entry 51660. The study did not develop new code. Standard procedures for

mathematical and statistical analysis of data are described in Methods.

SUPPLEMENTAL MATERIAL TITLES AND LEGENDS

Figure S1. Alignment of *Dm*Ago1 with *At*AGO1 and *Hs*Ago2.

The alignment highlights the position of the Lys514 residue ubiquitinated in unloaded

DmAgo1 by the E3 ligase Iruka[21]. Secondary structure designation is adopted from

*Hs*Ago2 as represented in [5].

Figure S2. The N-terminus of AtAGO1 is predicted to be an intrinsically disordered

region.

(A) Schematic representation of AtAGO1 with color-coding of the folded domains as

indicated. Residues 1-268 were included in the IDR predictions in (B), (C), and (D).

(B) Intrinsic disorder prediction by IUPred[62].

(C) Structure of the N-terminus predicted by Globplot[63].

(D) Structure of the N-terminus predicted by PONDR[64]. The position of the N-coil is

highlighted by the box.

Figure S3. N-coil mutations reduce the strength of bimolecular fluorescence

complementation mediated by AGO1-ATI1 and AGO1^{Y691E}-ATI1.

Half of the confocal microscopy images used for the quantification shown in Figure 5B

have been assembled in this panel. KK, K185E/K190E.

Figure S4. ^{35}S-Met/Cys uptake in seedlings.

Autoradiogram made from herbarium of *ago1-3$^{-/-}$*/FLAG-*AGO1* seedlings 9 days after

germination pulsed with ^{35}S-Met/Cys for the indicated periods of time.

Figure S5. Pulse labeling of AGO1.

(A) Immunoblot analysis with α-FLAG antibody and autoradiogram of FLAG-AGO1, FLAG-

AGO1^{Y691E} and FLAG-AGO1$^{KKME/Y691E}$ pulsed with ^{35}S-Met/Cys for the indicated periods of

time. Amounts of FLAG-tagged AGO1 in total lysates and in immunoprecipitated fractions

from seedlings using α-FLAG antibody are shown. * indicates a protein co-

immunoprecipitating specifically with unloaded AGO1. The protein was identified as

HSP70 as indicated by probing with α-HSP70 antibody. For this reason, ^{35}S label from the

~70 kDa band was not included in the AGO1 quantifications, see also Methods. KKME,

K185E/K190E/M304E/E307A. Raw data from the gels in this panel underlie the graphical

representations shown in Figure 6B,C.

(B) An independent repeat of the same type of experiment as in (C), with FLAG-*ago1*^{Y691E}

and FLAG-*ago1*$^{KKME/Y691E}$. Raw data from the gels in this panel underlie the graphical

representation shown in Figure 6D.

Table S1 Oligonucleotides used in the study

REFERENCES

1.   Bartel, D.P. Metazoan MicroRNAs. *Cell* **173**, 20-51 (2018).
2.   Ding, S.W. & Voinnet, O. Antiviral immunity directed by small RNAs. *Cell* **130**, 413-
26 (2007).
3.   Czech, B. et al. piRNA-Guided Genome Defense: From Biogenesis to Silencing.
Annual Review of Genetics **52**, 131-157 (2018).
4.   Fang, X. & Qi, Y. RNAi in Plants: An Argonaute-Centered View. *The Plant Cell* **28**,
272-285 (2016).
5.   Poulsen, C., Vaucheret, H. & Brodersen, P. Lessons on RNA silencing mechanisms
in plants from eukaryotic argonaute structures. *Plant Cell* **25**, 22-37 (2013).
6.   Meister, G. Argonaute proteins: functional insights and emerging roles. *Nat Rev
Genet* **14**, 447-59 (2013).
7.   Wang, Y., Sheng, G., Juranek, S., Tuschl, T. & Patel, D.J. Structure of the guide-
strand-containing argonaute silencing complex. *Nature* (2008).
8.   Schirle, N.T. & MacRae, I.J. The crystal structure of human Argonaute2. *Science*
336, 1037-40 (2012).
9.   Nakanishi, K., Weinberg, D.E., Bartel, D.P. & Patel, D.J. Structure of yeast
Argonaute with guide RNA. *Nature* **486**, 368-74 (2012).
10.   Elkayam, E. et al. The structure of human argonaute-2 in complex with miR-20a.
Cell **150**, 100-10 (2012).
11.   Ma, J.B. et al. Structural basis for 5'-end-specific recognition of guide RNA by the A.
fulgidus Piwi protein. *Nature* **434**, 666-70 (2005).
12.   Parker, J.S., Roe, S.M. & Barford, D. Structural insights into mRNA recognition from
a PIWI domain–siRNA guide complex. *Nature* **434**, 663-666 (2005).
13.   Ma, J.B., Ye, K. & Patel, D.J. Structural basis for overhang-specific small interfering
RNA recognition by the PAZ domain. *Nature* **429**, 318-22 (2004).
14.   Reichholf, B. et al. Time-Resolved Small RNA Sequencing Unravels the Molecular
Principles of MicroRNA Homeostasis. *Mol Cell* **75**, 756-768 e7 (2019).
15.   Derrien, B. et al. Degradation of the antiviral component ARGONAUTE1 by the
autophagy pathway. *Proc Natl Acad Sci U S A* **109**, 15942-6 (2012).

16. Smibert, P., Yang, J.-S., Azzam, G., Liu, J.-L. & Lai, E.C. Homeostatic control of
Argonaute stability by microRNA availability. *Nature Structural & Molecular Biology*
20, 789-795 (2013).
17. Martinez, N.J. & Gregory, R.I. Argonaute2 expression is post-transcriptionally
coupled to microRNA abundance. *RNA* **19**, 605-612 (2013).
18. Johnston, M., Geoffroy, M.-C., Sobala, A., Hay, R. & Hutvagner, G. HSP90 Protein
Stabilizes Unloaded Argonaute Complexes and Microscopic P-bodies in Human
Cells. *Molecular Biology of the Cell* **21**, 1462-1469 (2010).
19. Gibbings, D. et al. Selective autophagy degrades DICER and AGO2 and regulates
miRNA activity. *Nat Cell Biol* **14**, 1314-21 (2012).
20. Kobayashi, H., Shoji, K., Kiyokawa, K., Negishi, L. & Tomari, Y. VCP Machinery
Mediates Autophagic Degradation of Empty Argonaute. *Cell Reports* **28**, 1144-
1153.e4 (2019).
21. Kobayashi, H., Shoji, K., Kiyokawa, K., Negishi, L. & Tomari, Y. Iruka Eliminates
Dysfunctional Argonaute by Selective Ubiquitination of Its Empty State. *Molecular
Cell* **73**, 119-129.e5 (2019).
22. Hacquard, T. et al. The Arabidopsis F-box protein FBW2 targets AGO1 for
degradation to prevent spurious loading of illegitimate small RNA. *Cell Reports*
39(2022).
23. Rudel, S. et al. Phosphorylation of human Argonaute proteins affects small RNA
binding. *Nucleic Acids Res* **39**, 2330-43 (2011).
24. Bibel, B., Elkayam, E., Silletti, S., Komives, E.A. & Joshua-Tor, L. Target binding
triggers hierarchical phosphorylation of human Argonaute-2 to promote target
release. *eLife* **11**, e76908 (2022).
25. Maier, C.S. & Deinzer, M.L. Protein Conformations, Interactions, and H/D
Exchange. in *Methods in Enzymology*, Vol. 402 312-360 (Academic Press, 2005).
26. Sjøgaard, I.M.Z. et al. The transmembrane autophagy cargo receptors ATI1 and
ATI2 interact with ATG8 through intrinsically disordered regions with distinct
biophysical properties. *Biochemical Journal* **476**, 449-465 (2019).
27. Wienken, C.J., Baaske, P., Rothbauer, U., Braun, D. & Duhr, S. Protein-binding
assays in biological liquids using microscale thermophoresis. *Nat Commun* **1**, 100
(2010).
28. Li, S. et al. MicroRNAs inhibit the translation of target mRNAs on the endoplasmic
reticulum in Arabidopsis. *Cell* **153**, 562-74 (2013).
29. Sjögren, L. et al. Farnesylated heat shock protein 40 is a component of membrane-
bound RISC in Arabidopsis. *Journal of Biological Chemistry* **293**, 16608-16622
(2018).
30. Michaeli, S. et al. The viral F-box protein P0 induces an ER-derived autophagy
degradation pathway for the clearance of membrane-bound AGO1. *Proceedings of
the National Academy of Sciences* **116**, 22872-22883 (2019).
31. Li, C. et al. Collapse of Germline piRNAs in the Absence of Argonaute3 Reveals
Somatic piRNAs in Flies. *Cell* **137**, 509-521 (2009).
32. Matsumoto, N. et al. Crystal Structure of Silkworm PIWI-Clade Argonaute Siwi
Bound to piRNA. *Cell* **167**, 484-497.e9 (2016).
33. Yamaguchi, S. et al. Crystal structure of Drosophila Piwi. *Nature Communications*
11, 858 (2020).
34. Anzelon, T.A. et al. Structural basis for piRNA targeting. *Nature* **597**, 285-289
(2021).

35.   Brennecke, J. et al. Discrete small RNA-generating loci as master regulators of
transposon activity in Drosophila. *Cell* **128**, 1089-103 (2007).

36.   Gunawardane, L.S. et al. A slicer-mediated mechanism for repeat-associated
siRNA 5' end formation in Drosophila. *Science* **315**, 1587-90 (2007).

37.   Brennecke, J. et al. An Epigenetic Role for Maternally Inherited piRNAs in
Transposon Silencing. *Science* **322**, 1387-1392 (2008).

38.   Sato, K. et al. Krimper Enforces an Antisense Bias on piRNA Pools by Binding
AGO3 in the Drosophila Germline. *Molecular Cell* **59**, 553-563 (2015).

39.   Yuan, Y.-R. et al. Crystal Structure of A. aeolicus Argonaute, a Site-Specific DNA-
Guided Endoribonuclease, Provides Insights into RISC-Mediated mRNA Cleavage.
Molecular Cell **19**, 405-419 (2005).

40.   Wang, Y., Sheng, G., Juranek, S., Tuschl, T. & Patel, D.J. Structure of the guide-
strand-containing argonaute silencing complex. *Nature* **456**, 209-213 (2008).

41.   Miyoshi, T., Ito, K., Murakami, R. & Uchiumi, T. Structural basis for the recognition
of guide RNA and target DNA heteroduplex by Argonaute. *Nature Communications*
7, 11846 (2016).

42.   Willkomm, S. et al. Structural and mechanistic insights into an archaeal DNA-guided
Argonaute protein. *Nature Microbiology* **2**, 17035 (2017).

43.   Gregory, B.D. et al. A link between RNA metabolism and silencing affecting
Arabidopsis development. *Dev Cell* **14**, 854-66 (2008).

44.   Fang, X., Cui, Y., Li, Y. & Qi, Y. Transcription and processing of primary microRNAs
are coupled by Elongator complex in Arabidopsis. *Nature Plants* **1**, 15075 (2015).

45.   Hauptmann, J. et al. Turning catalytically inactive human Argonaute proteins into
active slicer enzymes. *Nature Structural & Molecular Biology* **20**, 814-817 (2013).

46.   Xiao, Y., Maeda, S., Otomo, T. & MacRae, I.J. Structural basis for RNA slicing by a
plant Argonaute. *bioRxiv*, 2022.07.15.500266 (2022).

47.   Bohmert, K. et al. AGO1 defines a novel locus of Arabidopsis controlling leaf
development. *Embo J* **17**, 170-80 (1998).

48.   Barghetti, A. et al. Heat-shock protein 40 is the key farnesylation target in meristem
size control, abscisic acid signaling, and drought resistance. *Genes Dev* **31**, 2282-
2295 (2017).

49.   Clough, S.J. & Bent, A.F. Floral dip: a simplified method for Agrobacterium-
mediated transformation of Arabidopsis thaliana. *Plant J* **16**, 735-43 (1998).

50.   Arribas-Hernandez, L. et al. The Slicer Activity of ARGONAUTE1 Is Required
Specifically for the Phasing, Not Production, of Trans-Acting Short Interfering RNAs
in Arabidopsis. *Plant Cell* **28**, 1563-80 (2016).

51.   Vranken, W.F. et al. The CCPN data model for NMR spectroscopy: Development of
a software pipeline. *Proteins: Structure, Function, and Bioinformatics* **59**, 687-696
(2005).

52.   Palmer, A.G., Cavanagh, J., Wright, P.E. & Rance, M. Sensitivity improvement in
proton-detected two-dimensional heteronuclear correlation NMR spectroscopy.
Journal of Magnetic Resonance (1969) **93**, 151-170 (1991).

53.   Bax, A. & Davis, D.G. MLEV-17-based two-dimensional homonuclear magnetization
transfer spectroscopy. *Journal of Magnetic Resonance (1969)* **65**, 355-360 (1985).

54.   Zhang, O., Kay, L.E., Olivier, J.P. & Forman-Kay, J.D. Backbone 1H and 15N
resonance assignments of the N-terminal SH3 domain of drk in folded and unfolded
states using enhanced-sensitivity pulsed field gradient NMR techniques. *Journal of*
Biomolecular NMR **4**, 845-858 (1994).

55. Schleucher, J. et al. A general enhancement scheme in heteronuclear
multidimensional NMR employing pulsed field gradients. *Journal of Biomolecular*
NMR **4**, 301-306 (1994).
56. Grefen, C. & Blatt, M.R. A 2in1 cloning system enables ratiometric bimolecular
fluorescence complementation (rBiFC). *BioTechniques* **53**, 311-314 (2012).
57. Mehlhorn, D.G., Wallmeroth, N., Berendzen, K.W. & Grefen, C. 2in1 Vectors
Improve In Planta BiFC and FRET Analyses. in *The Plant Endoplasmic Reticulum :*
Methods and Protocols (eds. Hawes, C. & Kriechbaumer, V.) 139-158 (Springer
New York, New York, NY, 2018).
58. Qu, F. et al. RDR6 has a broad-spectrum but temperature-dependent antiviral
defense role in Nicotiana benthamiana. *J Virol* **79**, 15209-17 (2005).
59. Krissinel, E. & Henrick, K. Inference of Macromolecular Assemblies from Crystalline
State. *Journal of Molecular Biology* **372**, 774-797 (2007).
60. Lumpkin, R.J. & Komives, E.A. DECA, A Comprehensive, Automatic Post-
processing Program for HDX-MS Data*. *Molecular & Cellular Proteomics* **18**, 2516-
2523 (2019).
61. McNicholas, S., Potterton, E., Wilson, K.S., and Noble, M.E.M. The CCP4mg
molecular-graphics software. *Acta Cryst.* **D67**, 386-394 (2011).
62. Mészáros, B., Erdős, G. & Dosztányi, Z. IUPred2A: context-dependent prediction of
protein disorder as a function of redox state and protein binding. *Nucleic Acids*
Research **46**, W329-W337 (2018).
63. Linding, R., Russell, R.B., Neduva, V. & Gibson, T.J. GlobPlot: Exploring protein
sequences for globularity and disorder. *Nucleic Acids Res* **31**, 3701-8 (2003).
64. Dosztanyi, Z., Csizmok, V., Tompa, P. & Simon, I. The pairwise energy content
estimated from amino acid composition discriminates between folded and
intrinsically unstructured proteins. *J Mol Biol* **347**, 827-39 (2005).
998

C. N-coil conservation and intra-molecular interaction analysis

N-coil	Phe177 *	Lys178 *	Phe179 *	Pro180 *	Met181	Arg182 *	Pro183 *	Gly184 ·	Lys185 *	Gly186 *	Gln187 *	Ser188 *	Gly189 *	Lys190 ·
Arabidopsis Ago1 / Human Ago2	Phe23	Lys24	Pro25	Pro26	Pro27	Arg28	Pro29	Asp30	Phe31	Gly32	Thr33	Ser34	Gly35	Arg36
Hydrogen bond (H), Salt bridge (S)			H			HS		H			H	H	H	HS
Buried Surface Area, Å²	74	15	78	87	16	140	55	42	61	70	19	81	33	76
Δ_iG Solvation energy effect, kcal/mol	0,43	-0,11	1,16	0,95	0,23	-1,36	0,88	-0,47	0,96	0,35	-0,08	0,02	0,44	-1,48

Figure 1. Structural properties and conserved interactions of the N-coil in Hs-Ago2

(A)-(B) Overview of the *Hs*Ago2 domain structure with surface outline of the globular N domain and the N-coil. (A) "side view"; (B) "bottom view".

(C) Intramolecular interactions between the N-coil and the remaining part of *Hs*Ago2 as calculated by the Protein Interfaces, Surfaces and Assemblies service PISA[59]. The solvation energy gain of the N-coil is calculated as the difference between free energy of the Piwi-associated and -dissociated N-coil. A negative solvation energy, Δ_iG, of a residue makes a positive contribution to the solvation energy gain of the interface, thus favoring the dissociated state. The color gradient (blue to red) indicates the effect sizes of buried surface area and solvation energy and highlights the unique properties of Arg-28 in *Hs*Ago2.

(D) Inlet showing conserved residues and bonds between a buried arginine side chain in the N-coil (Arg-28 in Hs-Ago2) and nearby residues.

(E) Model illustrating a hypothetical function of the N-coil in distinguishing loaded from unloaded AGO protein and in conferring rapid proteolysis on unloaded AGO. Left, in the loaded form, the N-coil is inaccessible for protein-protein interaction due to its association with the Piwi domain. In the unloaded form (middle and right), the N-coil is accessible for interactors, leading to rapid proteolysis (right) that can be slowed down if N-coil-dependent interactions are precluded by mutation (middle).

Presentations and PISA calculations were made using the crystal structure coordinates of the miR20a-bound *Hs*Ago2 (PDB 4F3T)[10].

Figure 2. AGO1^{Y691E} does not associate with small RNA.

(A) Immunoblot analysis with α-FLAG antibody of total protein and α-FLAG immunoprecipitated protein from inflorescence tissue from the indicated genotypes. The lower panel shows polynucleotide kinase labeling of RNA extracted from α-FLAG IPs. Ponceau stain shows total protein. Y, Y691E; KKME, K185E/K190E/M304E/E307A. Numbers (Y-2, Y-5, KKME/Y-2, KKME/Y-6) denote independent transgenic lines.

(B) Images 8 days after germination of homozygous *ago1-3* (null) plants and transgenic plant lines expressing FLAG-AGO1^{Y691E} or FLAG-AGO1$^{KKME/Y691E}$ in the *ago1-3* homozygous background.

Figure 3. Preferential accessibility of the N-coil in the unloaded state of AGO proteins

(A) Schematic representation of the *in vitro* binding experiment shown in (B) and (C).

(B) Protein blots of identical amounts of immobilized, FLAG immuno-affinity purified FLAG-AGO1 variants (WT, Y691E, KKME/Y691E) incubated with the indicated concentrations of AGO1 N-coil antibody (α-AGO1-N) and competitively eluted after extensive washing. Top panel, detection with FLAG antibodies (α-FLAG) to document equal amounts of FLAG-AGO1 variants in each of the binding assays. Middle panel, detection with α-AGO1-N to document specificity of the antibody (compare AGO1[Y691E] (intact N-coil) to AGO1[Y691E]/KKME (two point mutations (K185E/K190E) in N-coil) and equal reactivity towards loaded and unloaded AGO1 in the denatured state. Bottom panel, detection with anti-rabbit IgG to quantify the amount of α-AGO1-N retained by the immobilized FLAG-AGO1 variants. KKME, K185E/K190E/M304E/E307A.

(C) Quantification of the amount of α-rabbit IgG signal in (B) at different α-AGO1-N concentrations. The lines show an exponential fit to the data as a trendline.

(D) Immunoprecipitation of FLAG-AGO1[WT] and FLAG-AGO[Y691E] with α-AGO1-N antibody from total lysates of plants expressing comparable levels of FLAG-AGO1[WT] and FLAG-AGO[Y691E]. The Ponceau stain of the lanes containing aliquots of total lysates shows a crop around the large subunit of RuBisCO.

(E) Projection onto the *Hs*Ago2 structure of the published fractional differential hydrogen-deuterium exchange (HDX) between unloaded and guide loaded hAgo2[24]. Increasing differences in exchange rate are indicated with progressively darker purple shading. All regions with differential rates of exchange between unloaded and guide-loaded *Hs*Ago2 show higher exchange rate in the unloaded state, most notably areas buried by the N-coil and engaging in interactions with N-coil residue Arg28 in the guide-loaded *Hs*Ago2.

Figure 4. The N-coil is a direct binding site for ATI1 *in vitro*

(A) SDS-PAGE of heterologously expressed His_6-SUMO-NcGN^AGO1 protein used for fluorescence labeling and microscale thermophoresis (MST) assays.

(B) MST measurements of fluorescence intensity of His_6-SUMO-NcGN^AGO1 WT (left) and K178E (right) after application of a temperature gradient. Fluorescence data obtained from the thermophoretic acquisition period were normalized to the initial fluorescence before being fitted to a one-site binding curve with increasing concentrations of ATI1-IDR. The reported binding constants (K_D) are calculated with standard errors.

(C) Simple fluorescence intensity of Cys192-labeled His_6-SUMO-NcGN^AGO1 WT (left) and K178E (right) as a function of the concentration of ATI1-IDR.

(D, F) ^{1}H,^{15}N-HSQC nuclear magnetic resonance spectra of 50 μM of ^{15}N-labeled AGO1 N-coil WT (left, black) or K178E peptide (right, black), with 19 μM ATI1-IDR added (green with N-coil WT, blue with N-coil K178E). Zooms of contour levels of residues K178 and C192 are highlighted in the boxes. Minor peaks were assigned to a C-terminally truncated version of the AGO1 N-coil peptide whose presence in the sample was verified by MALDI-TOF mass spectrometry.

(E, G) Peak intensity plots of ^{15}N-labeled AGO1 N-coil peptide (50 μM) upon titration of ATI1-IDR. Peak intensities are normalized to peak intensities observed with free AGO1 N-coil peptide.

(H) Peak intensity of residues K178 (for the AGO1 wild type N-coil, green) and E178 (for the K178E N-coil mutant, blue) as a function of the ATI1-IDR concentration. The data points are fitted to an exponential function as a trendline.

Figure 5. Unloaded AGO1 interacts with ATI1 *in vivo*

(A) Confocal images showing results of bimolecular fluorescence complementation (BiFC) assays to probe the AGO1-ATI1 interaction *in vivo*. The C-terminal half of YFP was fused to AGO10, or to wild type or unloaded (Y691E) AGO1 with either an intact or a mutated (K185E/K190E) N-coil. The N-terminal half of YFP was fused to ATI1. Both fused halves of YFP were expressed from the same plasmid also expressing free RFP. For each BiFC pair, possible interaction is detected by fluorescence from reconstituted YFP. RFP fluorescence serves as a control for transformation. Merged channel images demonstrate the presence of yellow fluorescence only in transformed cells. Scale bar, 50 µm.

(B) Quantification of YFP signal normalized to RFP signal in the given number of YFP-positive cells for the interactions shown in (A). Letters indicate groups with statistically significantly different $\log_2$(YFP/RFP) ratios ($p < 0.01$; one-way ANOVA with post-hoc Tukey Honestly Significant Difference test). See Figure S3 for image data underlying the quantification in (B). KK, K185E/K190E.

Figure 6. An intact AGO1 N-coil is required for maximal turnover rate of unloaded AGO1 *in vivo*

(A) Kinetic model of the two different fates of newly synthesized AGO1: direct degradation (κ_1), or small RNA loading (κ_2) followed by degradation (κ_3).

(B) Quantification of ^{35}S-Met/Cys incorporation into FLAG-AGO1^{WT}, FLAG-AGO1^{Y691E} (FLAG-AGO1^Y) or FLAG-AGO1$^{KKME/Y691E}$ (FLAG-AGO1 $^{KKME/Y}$) as a function of pulse time. The lines represent a fit to an exponential equation for unloadable AGO1 containing the Y691E mutation and a fit to a double exponential equation for the WT AGO1. The half-lives calculated by the fitted models are given in hours +/- standard deviation, as detailed in Methods. KKME, K185E/K190E/M304E/E307A.

(C) Zoom in of the dashed area of (B).

(D) Result of an independent experiment done as in (C) to verify the slower incorporation kinetics in FLAG-AGO1$^{KKME/Y691E}$ compared to FLAG-AGO1^{Y691E}. See Figure S5 for raw data underlying the graphs in Figure 6B-D.

A

Homo sapiens - Ago2   *Bombyx mori* - Siwi   *Ephydatia fluviatilis* - Piwi   *Drosophila melanogaster* - Piwi

N coil ■ N domain ■ L1 □ PAZ ■ L2 ■ MID ■ PIWI ■ Surface outline ▬

B *Homo sapiens* - Ago2   *Methanocaldcoccus janaschii* - Ago   **C** Superposition of PIWI domain

Figure 7. Piwi-clade, but not prokaryotic, AGO proteins have a structurally flexible N-coil

(A) Structural overview of *Hs*Ago2 and Piwi proteins, outlining their N-coils in pink. The *Ephydatia fluviatilis* and *Drosophila melanogaster* Piwi proteins harbour N-coil truncations, probably generated by proteolysis during sample preparation for structural studies[33,34]. The structural representations are based on the published coordinates (*Hs*Ago2, PDB: 4F3T[10]; *Bm*Siwi PDB: 5GUH[32]; *Ef*Piwi, PDB: 7KX7[34]; *Dm*Piwi PDB: 6KR6[33]).

(B) Close-up of region in the Piwi domain in eukaryotic *Hs*Ago2 (PDB: 4F3T[10]) that cages the buried N-coil Arg28 side chain in the small RNA-bound conformation. Contacts between Arg28 and the loop containing the conserved His742, Glu745 and Asp747 residues are shown by dashed lines. A cluster of histidines formed by residues in β-strands of the Piwi-domain (His764, His766) and in the Arg28-interacting loop (His742) is highlighted in blue.

(C) Close-up of the region in the Piwi domain of the prokaryotic *Mj*Ago (PDB: 5G5T[42]) that corresponds to the region of *Hs*Ago2 shown in (B).

(D) Secondary structure matching of the *B. mori* and *D. melanogaster* Piwi domains reveals a more relaxed state of the *D. melanogaster* Piwi molecule (green) in which the structurally fixed N-coil seen in *B. mori* Siwi is absent.