1 **Title**

2 Single-fibre proteomics reveals a greater fibre-type specific mitochondrial remodelling with moderate
3 than high-intensity training.
4
Author list

E. Reisman[1,2,12], J. Botella[1,3,12], C. Huang[4], R. B. Schittenhelm[4], D. A. Stroud[5,6], C. Granata[1,7,8,9], S.

Chandrasiri[1], G. Ramm[10], V. Oorschot[10], N. J. Caruana[1,5*] & D. J. Bishop[1*]

Affiliations

1. Institute for Health and Sport (iHeS), Victoria University, Melbourne, Victoria, 3011, Australia
2. Mary MacKillop Institute for Health Research, Australian Catholic University, Melbourne, Victoria, 3000,
Australia
3. Metabolic Research Unit, School of Medicine and Institute for Mental and Physical Health and Clinical
Translation (IMPACT), Deakin University, Waurn Ponds, Victoria, 3216, Australia.
4. Monash Proteomics & Metabolomics Facility, Biomedicine Discovery Institute and Department of
Biochemistry and Molecular Biology, Monash University, Clayton, Victoria, 3168, Australia
5. Department of Biochemistry and Pharmacology, Bio21 Molecular Science and Biotechnology Institute, The
University of Melbourne, Parkville, Victoria, 3052, Australia
6. Murdoch Children's Research Institute, Royal Children's Hospital, Parkville, Victoria, 3052, Australia
7. Department of Diabetes, Central Clinical School, Monash University, Melbourne, Victoria, 3004, Australia
8. Institute for Clinical Diabetology, German, Diabetes Center, Leibniz Center for Diabetes Research at
Heinrich-Heine-University, Düsseldorf, Düsseldorf, 40225, Germany
9. German Center for Diabetes Research, Partner Düsseldorf, München-Neuherberg, 40225, Germany
10. Ramaciotti Centre for Cryo EM, Biomedicine Discovery Institute and Department of Biochemistry and
Molecular Biology, Monash University, Clayton, Victoria, 3168, Australia
11. Electron Microscopy Core Facility, European Molecular Biology Laboratory, Heidelberg, 69117,
Germany
28
29 12. These authors contributed equally to this work

30 * Correspondence: David.Bishop@vu.edu.au, nikeisha.caruana@vu.edu.au
31
32

Summary

Analyses of mitochondrial adaptations in human skeletal muscle have mostly used whole-muscle

samples, where results may be confounded by the presence of a mixture of type I and II muscle

fibres. Using our MS-based proteomics workflow, we provide new insights into fibre-specific

mitochondrial adaptations following two types of exercise training that were very different in terms

of metabolic demands and fibre recruitment patterns – moderate-intensity-continuous training

(MICT) and sprint interval training (SIT). There was a larger number of differentially expressed

proteins in each fibre type following MICT than SIT. Our novel mitochondrial normalisation

strategy highlighted that most training-induced changes in mitochondrial protein abundances were

stoichiometrically linked to the overall increase in mitochondrial content, except for the decreased

abundance of complex IV subunits in both type I and II fibres following SIT, and the increase in

proteins associated with fatty acid oxidation in type I fibres.

Keywords: Proteomics, single fibre, exercise, complex IV, fatty acid oxidation

48
49

Introduction

The remarkable ability of skeletal muscle to adapt to repeated contractile stimuli is one of the most fundamental and intriguing aspects of physiology (Egan and Zierath, 2013). Repeated exercise sessions (i.e., exercise training) alter the expression of proteins, which can lead to improved maintenance of cellular homeostasis (Hawley et al., 2014), altered metabolism (Hargreaves and Spriet, 2020), and the prevention and treatment of many chronic diseases (Neufer et al., 2015). Exercise is also a potent stimulus to activate mitochondrial biogenesis (Bishop et al., 2019a; Miller and Hamilton, 2012), with resulting increases in mitochondrial content and respiratory function (Granata et al., 2021; Holloszy and Coyle, 1984). These adaptations appear to depend on the characteristics of the exercise stimulus (Jamnick et al., 2020; MacInnis and Gibala, 2017).

One limitation of most human training studies to date is the analysis of whole-muscle samples, where the results may be confounded by diverse changes in different fibre types (Murphy, 2011). Human skeletal muscle is composed of three major fibre types (type I, IIa, and IIx), which differ in their contractile, metabolic, and bioenergetic characteristics; this is reflected by a greater reliance on mitochondrial oxidative enzyme activity in type I fibres (Howald et al., 1985; Ingjer, 1979; Kiessling et al., 1974) and greater glycolytic enzyme activity in type II fibres (Jackman and Willis, 1996). Furthermore, Henneman's size principle (Henneman and Olson, 1965) proposes that type I fibres have a lower threshold of activation, and will be recruited at lower exercise intensities, whereas fast-twitch fibres (type IIa and IIx) have a higher activation threshold and will be increasingly recruited at greater exercise intensities (Sale, 1987; Vollestad and Blom, 1985). Skeletal muscle fibre recruitment is therefore related to exercise intensity (Sale, 1987; Vollestad and Blom, 1985), and it has been hypothesised that higher exercise intensities might stimulate greater adaptations in type II fibres relative to lower intensities (Henneman, 1957; Henneman and Olson, 1965; Mendell, 2005). It has also been proposed that recruitment of type II fibres may help to explain the greater mitochondrial adaptations reported following high-intensity training (Dudley et al., 1982; MacInnis and Gibala, 2017; Sale, 1987; Vollestad and Blom, 1985). Until recently, however, technical limitations have prevented the direct assessment of fibre-type-specific adaptations to different types of exercise training.

Modifications to immunoblotting protocols have enabled researchers to identify training-induced adaptations of mitochondrial proteins in single skeletal muscle fibres (MacInnis et al., 2017; Skelly et al., 2021; Tan et al., 2018; Wyckelsma et al., 2015). However, due to limited starting material,

only five of the more than 1100 known mitochondrial proteins have been investigated to date

(Kristensen et al., 2015; MacInnis *et al.*, 2017; Murphy, 2011; Skelly *et al.*, 2021; Tan *et al.*, 2018;

Wyckelsma *et al.*, 2015). Improvements in mass-spectrometry-based proteomic techniques have

allowed greater resolution of the contractile and metabolic features of single skeletal muscle fibre

types – first in mice (Murgia et al., 2015) and then also in humans (Murgia et al., 2017). The

detection of functionally important low-abundance proteins, in addition to thousands of other

proteins, was a significant development that has advanced our understanding of the breath and

complexity of proteins at the fibre level in skeletal muscle. Recent work has built on this research

and adapted this technique to highlight fibre-type-specific protein changes to moderate-intensity

continuous training (MICT) in the skeletal muscle of a small number of males (n=5) (Deshmukh et

al., 2021). To date, however, no proteomic study has investigated the effects of divergent exercise

intensities, which differ in their recruitment of the major fibre types, on fibre-type-specific

mitochondrial adaptations. A greater understanding of fibre-type-specific adaptations to different

exercise intensities may help to optimise the prescription of exercise to prevent and treat diseases, as

well as counteract some of the detrimental effects of ageing.

In the present study, we developed a sensitive mass spectrometry-based (MS) proteomics workflow

that allowed for the pooling of single muscle fibres, and the utilisation of tandem mass tag (TMT)

labelling, to facilitate the identification of low abundant mitochondrial proteins. To investigate

potential fibre-type-specific changes in protein abundances following training, we employed two

very different training interventions - MICT and sprint-interval training (SIT). These two types of

training were chosen as they differ in intensity (i.e., moderate versus very high intensity) and have

been reported to require different skeletal muscle fibre recruitment patterns (Dudley *et al.*, 1982;

Kristensen *et al.*, 2015; Suriano et al., 2010). Our findings reveal distinct fibre-type-specific protein

responses to these two different types of exercise. Following MICT, changes in proteins (including

mitochondrial proteins) were primarily observed in type I fibres. Despite the hypothesis that SIT

would target type II fibres, we observed few differentially expressed proteins following SIT. There

was, however, a decreased abundance of complex IV subunits in both type I and II fibres following

SIT. We also observed that most training-induced changes in mitochondrial protein abundances were

stoichiometrically linked to the overall increase in mitochondrial content. Our study provides a new

analytical approach that expands our fundamental understanding of fibre-type-specific adaptations to

different types of exercise training. These findings challenge and expand our current knowledge of

fibre-type-specific changes to mitochondrial proteins in response to exercise training and provide an

important resource that may help to reveal the contribution of fibre-type-specific differences to
health and disease.

RESULTS

Single-fibre proteomics

Only two studies in young men, both with small sample sizes (n = 4 and 5, respectively), have
employed proteomics to describe fibre-type-specific differences for the contractile and metabolic
features of human skeletal muscle fibres (Deshmukh *et al.*, 2021; Murgia *et al.*, 2017). In the present
study, twenty-three, recreationally active, healthy men completed one of two 8-week training
interventions, consisting of either moderate-intensity continuous training (MICT) or very high-
intensity sprint-interval training (SIT) (see Table S1 – Tab 1 for participant characteristics). Pre and
post training, skeletal muscle biopsies were collected from the vastus lateralis muscle and single
fibres were isolated, fibre-typed, pooled, and analysed utilising the proteomic workflow described in
Figure 1a.

One of the challenges of analysing skeletal muscle is the large dynamic range of protein
concentrations caused by the highly abundant sarcomeric proteins, which account for over 50% of
the total proteome (Deshmukh et al., 2015; Murgia *et al.*, 2015; Ohlendieck, 2010; 2013), while
proteins from all other muscle compartments, including the mitochondria, are confined to the lower
half of the abundance range (Deshmukh *et al.*, 2015). Compared to typical whole-tissue proteomic
studies, single-muscle fibres provide very low protein yields (Galpin et al., 2012; Murgia *et al.*,
2015), which further complicates the identification of lower abundant proteins (Deshmukh *et al.*,
2015; Ohlendieck, 2011). While previous studies have used label-free quantitative shotgun
approaches after filter-aided sample preparation (Deshmukh *et al.*, 2021; Murgia *et al.*, 2017), we
chose a workflow incorporating tandem mass tag (TMT) labelling, which not only allowed us to
increase sample size and throughput to accommodate our complex experimental design but also
exhibits higher quantitative precision than label-free quantification (O'Connell et al., 2018;
Pappireddi et al., 2019).

Our approach identified a total of 3141 proteins across all pooled samples, with 81% (2534 proteins)
of these identified with high confidence (false discovery rate [FDR] < 0.05) and > 1 unique peptide
(Table S1-Tab 2). On average, we were able to reliably identify 1857 ± 59 (1477 to 2000) high-
confidence proteins per pooled single-fibre sample (Figures S2a, b, & c, Table S1-Tab 2; see also

Table S9 – Tab 1 & Tab 2). Proteins were removed from the dataset if they contained more than 70%

missing values, leaving a total dataset across all samples of 1623 proteins (Table S1-Tab 3).

To further understand the characteristics of the fibre-specific proteome, we examined the cellular

localisation of the proteins identified in each individual sample of pooled type I or type II fibres

based on Gene Ontology Cellular Components (GOCCs). GO enrichment analysis of our dataset

(Figure 1b, Table S1-Tab 4) demonstrates good coverage of the structural and metabolic features of

the muscle fibre proteome, including contractile, sarcomeric, ribosomal, and mitochondrial proteins.

To assess the comprehensiveness of our protein detection in pooled single fibres, we compared our

dataset to the two previous studies (Deshmukh *et al.*, 2021; Murgia *et al.*, 2017) investigating the

single muscle fibre proteome in humans (Figure 1c, Table S1-Tab 5). There were 846 proteins in

common between the three studies, which may represent proteins that can more confidently be used

to compare fibre types in human skeletal muscle. One of the most significantly enriched cellular

components for these commonly identified proteins was the mitochondrion (Figure 1d, Table S1-Tab

6).

Fibre Type Differences in the Pre-Training samples

Multi-dimensional scaling revealed a good separation of the protein profiles between type I and type

II fibres for most, but not all, pooled single fibre samples expressed between fibre types pre-training

(Figure 1e). Although the separation between fibre types was not as distinct as previously reported

(Deshmukh *et al.*, 2021), this may reflect the approximately 20% greater aerobic fitness of our

participants (V$\Box$O$_{2max}$ = 52 vs ~ 44 mL$^.$min$^{-1.}$kg^{-1}) and previous reports that type II fibres become

more like type I fibres with endurance training (Bergh et al., 1978; Jansson et al., 1978; Wilson et al.,

2012).

Proteomic features of the different fibre types are presented in a volcano plot (Figure 1f). A total of

120 proteins were identified as differentially expressed between fibre types pre-training (PRE), using

an empirical Bayes method and with statistical significance set at an adjusted P value cut-off of 0.05

(Table S1-Tab 7). Our results are in agreement with previous work (Deshmukh *et al.*, 2021),

although their study did identify a greater number of differentially expressed proteins between fibre

types when they adopted a posteriori information fusion scheme – a statistical approach that

combines biological significance (fold change) and statistical significance (P value from a single

paired t-test) (Xiao et al., 2014). Unsupervised hierarchical clustering revealed a clear separation of
fibre types, although, as expected, there was some biological variability amongst individuals (Figure
1g).
186
Our analysis revealed new contractile proteins with fibre-type-specific expression profiles (Table S1-
Tab 8). These included MYH3 and MYL4 – two poorly understood members of the myosin family
(Schiaffino et al., 2015). We also observed many of the expected differences between fibre types.
Type I fibres were characterised by a greater abundance of contractile proteins belonging to the
myosin heavy (MYH7) and light (MYL6B) chains, as well as troponin (TNNT1, TNNC1, TNNI1)
and an isoform of the sarcoplasmic reticulum Ca^{2+}-ATPase (ATP2A2, also named SERCA2;
(Lamboley et al., 2015; Morissette et al., 2014)), while type II fibres possessed a greater abundance
of MYH2, MYL1, TNNT3, TNNC2, TNNI2, and also ACTN3.
195
Most studies on muscle fibre heterogeneity have focused on contractile proteins (Ohlendieck, 2013;
Schiaffino, 2011). However, as confirmed in our study, the diversity between muscle fibres is not
only restricted to contractile proteins but extends to a wide range of subcellular systems - including
ionic transport, cellular calcium signalling, and metabolism (Schiaffino, 2011). Approximately 25%
of the proteins identified as differentially expressed between fibre types were mitochondrial – with
most (~ 95%) having a greater abundance in type I fibres. These differentially expressed
mitochondrial proteins were enriched in pathways associated with fatty acid oxidation, as well as
other proteins responsible for the provision and transfer of electrons to support energy provision
(Table S1-Tab 7).
205
206 **Influence of mitochondrial content on fibre-type-specific expression profiles**
207
Of the 395 mitochondrial proteins quantified in the present study, 273 (~ 70%) were in common with
the previous two single-fibre proteomic studies (Figure 2a, Table S2-Tab 1). The top 5 GO
Biological Process (BP) terms for these commonly detected mitochondrial proteins were related to
mitochondrial electron transport (NADH to ubiquinone and respiratory chain complex I assembly),
the tricarboxylic acid cycle, fatty acid beta-oxidation, and branched-chain amino acid catabolic
processes (Figure 2b, Table S2-Tab 2). In line with their superior capacity for oxidative
phosphorylation and fatty acid oxidation (Sahlin et al., 2007), and in agreement with previous work
(Deshmukh *et al.*, 2021), we also observed that type I fibres had a significantly greater abundance of
many mitochondrial proteins (Figure 2c, Table S1-Tab 7).

218 An important question that has not been adequately addressed is whether these fibre-type-specific

219 expression profiles for mitochondrial proteins are independent of the greater mitochondrial content

220 in type I versus type IIa human skeletal muscle fibres (Howald *et al.*, 1985; Kiessling et al., 1971;

221 Ørtenblad et al., 2018). To examine the potential influence of mitochondrial content on fibre-type-

222 specific expression profiles for mitochondrial proteins, we first calculated the mitochondrial protein

223 enrichment (MPE) – a value that identifies the contribution of mitochondrial protein intensities to the

224 overall detectable proteome (Granata *et al.*, 2021). Consistent with previous research (Deshmukh *et*

225 *al.*, 2021), there was a greater abundance of proteins associated with the mitochondria cellular

226 component in type I compared to type II fibres (P < 0.05, Figure 2d, Table S2-Tab 5). As this

227 supported an overall greater content of mitochondria in type I versus type II fibres in our

228 participants, we next employed our previously described normalisation strategy (Granata *et al.*,

229 2021), which corrects for the bias introduced by differences in total mitochondrial content, to

230 identify the relative abundance of individual mitochondrial proteins within each fibre type. This

231 strategy allows us to identify fibre-type differences in mitochondrial protein abundance that are

232 disproportionate to fibre-type differences in the total mitochondrial proteome. Post normalisation,

233 only 8 of the 27 mitochondrial proteins we had previously observed to have fibre-type-specific

234 expression profiles remained significantly different (see Figure 2c and compare Figure 2e, Table S1-

235 Tab 7, with Figure 2f, Table S2-Tab 7). This novel analysis suggests that many of the differences in

236 mitochondrial protein abundances between fibre types reported by us, and others (Deshmukh *et al.*,

237 2021; Murgia *et al.*, 2017), might be due to differences in total mitochondrial content rather than

238 fibre-type-specific remodelling of the mitochondrial proteome.

240 **Proteomic responses to different types of training in type I and type II skeletal muscle fibres**

242 Skeletal muscle has a remarkable ability to remodel in response to repeated contractile stimuli, but

243 little is known about fibre-type-specific changes in protein abundances following exercise training.

244 To date, only one proteomic study has reported fibre-type-specific adaptations to one type of

245 exercise training (1 h at 75 to 90% of maximum heart rate, 4x/week for 12 weeks) in the skeletal

246 muscle of five males (Deshmukh *et al.*, 2021). However, it is not known if fibre-type-specific

247 changes to the proteome depend on the nature of the exercise stimulus. The size principle states that

248 type I fibres are mostly utilised at lower exercise intensities, whereas type II fibres are increasingly

249 recruited at higher intensities (Henneman and Olson, 1965; Mendell, 2005). Therefore, a long-

250 standing hypothesis is that training-induced changes to the proteome in response to low-intensity

exercise will mostly be confined to the type I fibres that are predominantly recruited with this type of

exercise (Godin et al., 2010; Sale, 1987). In contrast, higher-intensity exercise (e.g., all-out sprinting)

has been proposed to stimulate greater adaptations in type II fibres compared to exercise performed

at lower intensities (Greenhaff et al., 1994).

To investigate fibre-type-specific changes to the proteome in response to divergent exercise stimuli,

we had 23 healthy young men train 3 to 4 times per week for 8 weeks after random assignment to

one of two very different types of training – MICT or SIT (Figure 3a). By design, both exercise

groups had very different exercise prescription characteristics; there was a 4.7-fold greater exercise

intensity for SIT and a 5.2-fold greater training volume for MICT (Figure 3b). The divergent

exercise prescriptions, and their associated effects on the metabolic response to exercise, can be

observed by the large increase in blood lactate concentration following SIT (9.9-fold; 1.6 ± 0.6 to

15.7 ± 5.9 mmol$^.$L^{-1}; P < 0.001) when compared to MICT (1.6-fold; 1.38 ± 0.43 to 2.18 ± 0.38

mmol$^.$L^{-1}; P = 0.001) (Figure 3b; Table S3-Tab 1). Similarly, blood pH levels were only decreased

following SIT (from 7.35 ± 0.03 to 7.05 ± 0.09; P < 0.001) and not MICT (from 7.35 ± 0.02 to 7.37

± 0.02; P > 0.05) (reflected by fold differences in [H$^+$] in Figure 3b). These results agree with the

well-described effect of exercise intensity on increasing glycolytic flux through anaerobic pathways

and the associated increase in lactate and decrease in pH (Hargreaves and Spriet, 2020; van Loon et

al., 2001). Immunohistochemical analysis of muscle sections confirmed there was a greater depletion

of muscle glycogen in type I than type II fibres following the first MICT session, and greater muscle

glycogen depletion in the type II fibres following SIT versus MICT (Figure 3c). This is consistent

with previous research (Gollnick et al., 1974; Scribbans et al., 2014) and demonstrates that the two

different types of training employed were associated with very different skeletal muscle fibre

recruitment patterns (i.e., MICT recruited predominantly type I fibres while SIT recruited both type I

& type II fibres; Figure 3d).

As with previous research (Deshmukh *et al.*, 2021), to identify fibre-type-specific changes in protein

abundances following exercise training we chose a posteriori information fusion scheme that

combines biological relevance (fold change) and statistical significance (P value) (Xiao *et al.*, 2014).

The fusion scheme yields a protein significance score (π value) and is robust in selecting

differentially expressed genes/proteins while protecting against false discoveries. Using this fusion

scheme, we identified 31 and 16 proteins that were differentially altered by 8 weeks of MICT in type

I and type II fibres, respectively (Figure 4a, b, left panels; Table S4-Tabs 1 & 3), with no overlap

between the proteins altered in each fibre type (Figure 4c; Table S4-Tab 5). Many of the proteins

with a lower abundance following MICT in type I fibres were related to skeletal muscle contraction
(Table S4-Tab 5). The decrease in the abundance of these proteins following MICT raises the
possibility that the fibre-type differences for these contraction-related proteins observed pre-training
may be a consequence of habitual physical activity rather than an intrinsic characteristic of the
different fibre types. In contrast to our hypothesis, there were few proteins that changed their
abundance in either type I or type II fibres following SIT (i.e., 4 to 8 x 30-s all-out sprints performed
3 to 4 times a week for 8 weeks) (Figure 4a,b, right panels; Table S4-Tabs 2 & 4). As with MICT,
there was no overlap between the proteins altered in each fibre type following SIT (Figure 4d; Table
S4-Tab 6).

The fibre-type-specific adaptations to MICT are strengthened by the greater number of differentially
expressed proteins between type I and type II fibres post-training compared to pre-training (compare
Figure 5a; Table S5-Tab 1 with Figure 1f). Similarly, the limited effects of SIT on the proteome are
reinforced by the small difference in the number of differentially expressed proteins between type I
and type II fibres in the SIT group after training versus pre-training (compare Figure 5b; Table S5-
Tabs 2 with Figure 1f). There were also far fewer proteins with altered expression in either type I or
type II fibres following SIT compared with MICT (Figure 5c; Table S5-Tab 3). Approximately 75%
of the proteins identified as differentially expressed between type I and type II fibres following SIT
also had fibre-specific differential expression pre-training (Figure 5d; Table S5-Tab 4), further
supporting the limited effects of SIT on the proteome. These results are consistent with those from a
study that used immunoblotting on pooled single fibres and observed greater fibre-type-specific
changes following MICT compared with SIT (Skelly *et al.*, 2021).

Of note, we identified an additional 80 differentially expressed proteins between the two fibre types
post-MICT that were not differentially expressed between the two fibre types pre-training (Figure
5d; Table S5-Tab 4). This again suggests greater fibre-type-specific changes to the proteome in
response to MICT than SIT. However, only six of these 80 proteins were also identified as being
significantly altered by 8 weeks of MICT (CES2, CLIC5, ETFB, LASP1, QDPR, TGM2; Table S4-
Tabs 1 & 3). Enrichment analysis was completed on all detected proteins in the POST samples for
each fibre type and for each training type (using the WikiPathways database; Figure 5e; Table S5-
Tab 5). Several pathways presented differences between fibre types in both directions (i.e., some
proteins within a pathway were higher in abundance in type I fibres and some proteins within the
same pathway were higher in type II fibres) – including pathways associated with striated muscle
contraction and proteasome degradation. Fibre-type-specific pathways included glycolysis and

gluconeogenesis, which had a greater enrichment in type II fibres for both groups. In contrast, the

pathways related to fatty acid oxidation and the TCA cycle were more prevalent in type I than type II

fibres.

We narrowed the analysis to look specifically at pathways related to muscle contraction, metabolism,

and glycolysis, and observed that specific proteins (and pathways) were differentially expressed in

the two fibre types after both training interventions (i.e., in MICT and SIT) (Figure5f; Table S5-Tab

6). The network analysis demonstrates that protein differences were more common between the same

fibre type than the same exercise training type. For example, for both types of training striated

muscle contraction proteins associated with slow-twitch skeletal muscle isoforms were

predominately higher post-training in type I fibres (TNNT1, TNNI1, TNNC1, MYL2 and MYL3),

while proteins associated with fast-twitch skeletal muscle isoforms were predominately more

abundant post-training in type II fibres (TNNC2, MYH6, ACTN3, TNNT3, MYBPC2, TPM1,

TNNI2, MYL1 and MYL4). There was also an influence of training type, with the greater abundance

of fast-twitch skeletal muscle proteins in type II fibres attributable to a lower abundance of these

proteins in type I fibres post MICT and no change in either fibre type post SIT (compare Figure 5f to

Figures 4a & 4b). The network analysis also indicated that proteins associated with glycolysis were

mostly differentially expressed only in type II fibres after both types of training; proteins belonging

to the glycogen synthesis pathway were only observed to be greater in type II fibres post MICT and

not post SIT. Finally, although TCA and fatty acid oxidation pathway proteins were predominately

differentially expressed in type I fibres following training, most of these differences were also

present in the PRE samples.

Fibre-type-specific effects of different types of exercise training on mitochondrial proteins

Mitochondrial content is greater in type I compared with type II fibres (Glancy and Balaban, 2011;

Leary et al., 2003) (see also Fig 2d). However, while research has reported that markers of

mitochondrial content increase with exercise training (Bishop *et al.*, 2019a; Bishop et al., 2014;

Larsen et al., 2012; Morgan et al., 1971), little is known about the fibre-specific effects of different

types of exercise training. Consistent with our morphological (mitochondrial volume density

obtained from TEM; Figures 6a & b) and biochemical (citrate synthase [CS] activity; Figure 6c)

markers of mitochondrial content (Table S6-Tab 1), we observed a significant increase in the mean

abundance of known mitochondrial proteins only following MICT and not SIT (Figure 6d) (Table

S6-Tab 2). This can likely be attributed to the five-fold greater volume of training completed by the

MICT group (Bishop *et al.*, 2019a; Granata et al., 2018), although a consensus has yet to be reached

on the relative importance of exercise intensity versus volume on training-induced changes in

markers of mitochondrial content (Bishop et al., 2019b; MacInnis et al., 2019). Changes in the mean

abundance of known mitochondrial proteins following MICT occurred predominantly in the type I

fibres, although there was also a significant but smaller increase in the type II fibres (Figure 6d).

There were no significant increases in known mitochondrial protein abundance in either fibre type

following SIT (Figure 6d).

We observed fibre-type-specific adaptations of mitochondrial proteins to training, consistent with

the only other single-fibre proteomic study investigating the effects of exercise training in single

fibres (Deshmukh *et al.*, 2021) (Figure S1, Table S8 – Tab 1). Our results suggest that the overall

mean abundance of many mitochondrial functional groups is enhanced by moderate-intensity

endurance training, but only in type I fibres. As some of these proteins have been reported to

decrease in aged fibres regardless of fibre type (Murgia *et al.*, 2017), it will be important to explore if

other types of exercise training are able to increase the expression of these proteins in type II fibres.

While changes in most mitochondrial proteins were largely confined to type I fibres following

MICT, the abundance of mitochondrial ribosome proteins increased in both fibre types from PRE to

POST MICT (Figure S1b, Table S8-Tab 2). These results are similar to those of another proteomic

study, which revealed changes in mitochondrial ribosomal proteins in whole muscle following

interval training (Robinson et al., 2017). These authors hypothesised that translational level

regulation is a predominant factor controlling mitochondrial biogenesis in humans in response to

exercise training. No changes in mitochondria ribosome proteins were observed in either fibre type

for SIT. This suggests the volume of MICT was sufficient to induce significant mitochondrial protein

synthesis in both fibre types, whereas SIT may not provide a sufficient stimulus to significantly alter

mitochondrial protein synthesis (Scalzo et al., 2014).

Fibre-type-specific expression profiles for mitochondrial proteins independent of training-

induced changes in mitochondrial content.

Given the greater changes in mitochondrial content that occur following MICT compared with SIT

(Bishop *et al.*, 2014; Granata *et al.*, 2018; Granata et al., 2016a; Granata et al., 2016b); see also

Figure 6d), we again turned our attention to investigating whether the fibre-type-specific changes in

the abundance of individual mitochondrial proteins we observed with the different training types

were independent of changes in total mitochondrial content. Once again, we employed our

previously described normalisation strategy to remove the bias introduced by the changes in
mitochondrial content in each fibre type following both types of training (Granata *et al.*, 2021). This
enabled us to investigate differences in mitochondrial protein abundances between the two fibre
types without the bias introduced by training-induced changes in total mitochondrial content (Figure
7, Table S7 – Tabs 1 & 2). Post normalisation, very few training-induced changes in mitochondrial
protein abundances remained in either fibre type, following either training intervention (Figure 7a).
One exception was the relative decrease in the abundance of complex IV subunits in both type I and
II fibres following SIT (Figure 7b & c). Thus, when changes in mitochondrial content are controlled
for, very high-intensity SIT appears to deprioritise the OXPHOS components involved in the
catalysis of the final step of the mitochondrial electron transport chain (i.e., complex IV). The other
exception was the increase of proteins associated with fatty acid oxidation in type I fibres following
MICT, which our results suggest was greater than, and not stoichiometrically linked, to the overall
training-induced increase in mitochondrial content (Figure 7d). This adds to the evidence that MICT
provides a powerful stimulus to increase fatty acid oxidation (Martin et al., 1993; Shaw et al., 2020);
we add that this adaptation occurs predominantly in type I skeletal muscle fibres (with the exercise
stimulus employed in our study).
419
420
421   **Discussion**
422
Proteomics applied to human skeletal muscle remains in its infancy but has introduced new
opportunities to explore the complexity of the biological networks underlying fibre-specific
differences and responses to exercise training. In the present study, we developed a sensitive MS-
based proteomics workflow that allowed for the pooling of single muscle fibres, and the utilisation of
tandem mass tag (TMT) labelling, to accommodate our complex experimental design and increase
the quantitative accuracy of our results. In addition, we incorporated a much larger sample size than
previous studies (i.e., 23 versus 4 to 5 participants) to better account for the individual variability in
human skeletal muscle and to increase the robustness of our findings. Consistent with previous
research (Deshmukh *et al.*, 2021), we identified many differentially expressed proteins between type
I and type II fibres (Figure 1f). However, further analysis of these results allowed us to make two
new and important observations. The first is that only 8 of the 27 mitochondrial proteins with fibre-
type-specific expression profiles remained significantly different when we used our previously
published normalisation strategy to account for the overall greater content of mitochondria in type I
versus type II fibres (Figure 2c). Thus, many of the differences in mitochondrial protein abundances

between fibre types reported by us (Figure 2e), and others (Deshmukh *et al.*, 2021; Murgia *et al.*,
2017), are likely due to differences in total mitochondrial content rather than representing a fibre-
type-specific mitochondrial proteome. The second new finding is that some of the fibre-type
differences observed pre-training (especially for contraction-related proteins) were increased by
moderate-intensity training. This highlights the challenge of distinguishing between protein
expression profiles that are intrinsic to different fibre types and those that are a consequence of
adaptations to habitual physical activity.

Repeated muscle contractions, via habitual physical activity or exercise training, provide a potent
stimulus to alter the expression of proteins and confer many health benefits (Neufer *et al.*, 2015). The
development of mass-spectrometry-based proteomics has revolutionised our understanding of
training-induced changes in protein abundance (Egan et al., 2011; Granata *et al.*, 2021; Holloway et
al., 2009; Hostrup et al., 2018), including at a fibre-specific level (Deshmukh *et al.*, 2021). However,
while it is well known that adaptations depend on the characteristics of the exercise stimulus
(Jamnick *et al.*, 2020), no proteomic study has investigated the effects of different exercise stimuli
on fibre-type-specific adaptations. Therefore, an important feature of our study design was the use of
two types of training that were very different in terms of volume, intensity, metabolic stress, and
fibre recruitment patterns (Figure 3). There was a larger number of differentially expressed proteins
following MICT than SIT (Figures 4 & 5), which can likely be attributed to the five-fold greater
volume of training completed by the MICT group (Bishop *et al.*, 2019a; Granata *et al.*, 2018). There
were also fibre-specific adaptations to both types of training, with no overlap between the changes
observed in each fibre type. Most notably, increases in proteins associated with complex I of the
ETC, fatty acid oxidation, the TCA cycle, and mitochondrial dynamics were mostly confined to type
I fibres following MICT. This improved understanding of how mitochondrial proteins adapt to
different exercise interventions may lead to more personalised exercise prescriptions to reverse some
of the fibre-specific changes that occur with ageing (Murgia *et al.*, 2017) and, potentially, some
chronic diseases.

Given the well-known effects of different types of training on mitochondrial content (Granata *et al.*,
2018); see also Figure 6), a further important feature of our study was to investigate whether fibre-
type-specific changes in mitochondrial protein abundance were independent of training-induced
changes in total mitochondrial content. To do this, we used our previously described normalisation
strategy to control for the bias introduced by changes in mitochondrial content in each fibre type
following both types of training (Granata *et al.*, 2021). This analysis revealed that most training-
induced changes in mitochondrial protein abundances were stoichiometrically linked to the overall

increase in mitochondrial content (compare Figure 7 with Figure S1). However, a surprising
observation was the decrease in the abundance of complex IV subunits, relative to the overall
increase in mitochondrial protein content, in both type I and II fibres following SIT. This
deprioritisation of the OXPHOS components involved in the final step of the mitochondrial electron
transport chain is intriguing given that, when also controlling for changes in mitochondrial content,
we have previously reported greater changes in mitochondrial respiration following SIT than MICT
(Granata *et al.*, 2016a). The results of our novel normalisation strategy also revealed a discordant
regulation of proteins associated with fatty acid oxidation, with a non-stoichiometric increase of
these proteins in type I but not type II fibres following both types of training. Interestingly, a
previous study has also reported greater increases in perilipin (PLIN) 2 and 5 in type I compared
with type II fibres following both MICT and SIT (Shepherd et al., 2017). These adaptations may
facilitate greater intramuscular triglyceride (IMTG) lipolysis and fatty-acid oxidation in type I fibres
when metabolic demand increases (Shepherd et al., 2012; 2013).

In summary, our study incorporated several unique elements (a sensitive MS-based proteomics
workflow, the largest sample size to date, two very different types of training stress, and a novel
mitochondrial normalisation strategy) to identify many fibre-specific adaptations to different types of
training. This provides an important resource to better understand how different fibre types adapt to
different training interventions, with potential insights for how best to prescribe exercise to improve
both health and human performance. In contrast to the decreased abundance of complex IV subunits
in both type I and II fibres following SIT, and the increase in proteins associated with fatty acid
oxidation only in type I fibres, most training-induced changes in mitochondrial protein abundances
were stoichiometrically linked to the overall increase in mitochondrial content.

Limitations of the study

This study recruited healthy, young, active men and, therefore, some of the conclusions may not
apply equally well to individuals of different genders, ages (e.g., the elderly), ethnicity, or health
status. Another potential limitation is the use of pooled muscle fibre samples; while this assists in
reducing biological variability amongst samples, it may not be a true representation of single-fibre
differences both pre and post-training. Finally, it was not possible for us to determine if the fibre-
specific adaptations to the two different types of exercise training had functional consequences (e.g.,
differences in fibre-specific mitochondrial respiratory function). It is also important to note that there

are several quantitative mass-spectrometry-based methods used to detect changes in the proteome,

each with their advantages and disadvantages, and the results are not always directly comparable.

STAR METHODS

KEY RESOURCES TABLE

REAGENT or RESOURCE	SOURCE	IDENTIFIER
Antibodies		
Myosin heavy chain Type IIA	Developmental Studies Hybridoma Bank (DSHB)	A4-74 RRID: AB_528383
Anti-Myosin heavy chain Type IIB	Developmental Studies Hybridoma Bank (DSHB)	A4-840; RRID: AB_528384
Goat anti-Mouse IgG (H+L) Highly Cross-Adsorbed Secondary Antibody, Alexa Fluor Plus 488	ThermoFisher Scientific	A32723; RRID: AB_2633275
Goat anti-Mouse IgM (Heavy chain) Cross-Adsorbed Secondary Antibody, Alexa Fluor 488	ThermoFisher Scientific	A-21042; RRID: AB_2535711
Myosin heavy chain (slow, alpha- and beta-)	Developmental Studies Hybridoma Bank (DSHB)	BA-F8, RRID: AB_10572253
Myosin heavy chain (all but 2X) antibody	Developmental Studies Hybridoma Bank (DSHB)	BF-35, RRID: AB_2274680
Goat anti-Mouse IgG2b Cross-Adsorbed Secondary Antibody, Alexa Fluor 350	ThermoFisher Scientific	A-21140, RRID: AB_2535777
Goat anti-Mouse IgG1 Cross-Adsorbed Secondary Antibody, Alexa Fluor 488	ThermoFisher Scientific	A-21121, RRID: AB_2535764
Critical commercial assays		
TMT16plex Isobaric Label Reagent Set	ThermoFisher Scientific	A44521
TMTpro™ Zero Label Reagent	ThermoFisher Scientific	A44519
96-well S-Trap™ plate	Protifi	C02-96well
Deposited data		
Proteomics Reveals a Fibre-Type Specific Mitochondrial Remodelling with Moderate but not Very High Intensity Training	This paper	PRIDE: PXD036010
Software and algorithms		
ImageJ	(Schneider et al., 2012)	https://imagej.nih.gov/ij/
R version 4.1.1	R Development Core Team, 2016	https://www.R-project.org/
Proteome Discoverer 2.4	Thermofisher Scientific	
Limma 3.32.2	(Ritchie et al., 2015)	https://bioconductor.org/packages/release/bioc/html/limma.html

Resource availability

Lead contact

515 Further information and requests for reagent and resources may be directed to and will be fulfilled by
516 the lead contact, Professor David J Bishop (David.Bishop@vu.edu.au).

517
518

Experimental model and Subject details

Participants and ethics approval

Twenty-eight healthy men initially volunteered to take part in this study. The data from twenty-three participants (27.2 ± 5.4 y; 178 ± 6 cm; 75.2 ± 9.7 kg; 23.6 ± 2.7 BMI; Table S1 – Tab 1) was included in the final analysis; two participants from the MICT group and two from the SIT group withdrew from the study due to time constraints, and the data from one participant was excluded from the final analysis as their muscle samples were of poor quality (see study recruitment flow chart and final group sizes; Figure S2f). Participants were provided with the information to participants' document, and were informed of the study requirements, benefits, and risks involved before giving their written informed consent. Ethics approval for the study was obtained from the Victoria University Human Research Ethics Committee (HRE17-075) and was registered as a clinical trial under Australian New Zealand Clinical Trials Registry (ANZCTR; ACTRN12617001105336).

Study design and training. During the first two weeks, participants performed two exercise familiarisation sessions and multiple testing familiarisation sessions. Participants were then allocated into one of the training groups, based on their age, body mass index (BMI), maximum rate of oxygen consumption (VO_{2max}), peak aerobic power output (W_{max}; $W.kg^{-1}$), and 20-km time trial (TT) performance ($W.kg^{-1}$) (all $P > 0.05$), to match the two groups for baseline endurance characteristics. The participants then underwent another week of testing and completed 8 weeks of training before the last muscle biopsy sample and final testing was performed (Figure 3a).

Sprint-Interval Training (SIT).

Twelve participants were allocated to this group (27.5 ± 5.1 y; 179 ± 6 cm; 75.5 ± 9.3 kg; 23.5 ± 2.7 BMI). The SIT group completed 4 to 8 30-second "all-out" cycling sprints against a resistance set at 0.075 kg/kg body mass (BM), interspersed with a 4-min recovery period between sprints (Gillen et al., 2016; Granata *et al.*, 2016a). The training load was increased to 0.080 $kg.kg^{-1}$ BM in week 3, to 0.085 $kg.kg^{-1}$ BM in week 5, and to 0.090 $kg.kg^{-1}$ BM in week 7. During the recovery, participants remained on the bikes and were allowed to either rest or cycle against no resistance. Participants started the training with 4 sprints per session and this increased up to 8 sprints per session in week 7.

Moderate-Intensity Continuous Training (MICT).

Twelve participants were allocated to this group (26.9 ± 5.6 y; 178 ± 6 cm; 74.9 ± 10.2 kg; 23.7 ± 2.6 BMI). The training intensity for the MICT training group was established as 10% less than the

first lactate threshold (~ 90% of the lactate threshold 1 (LT1; defined as the first increase of ≥ 0.3

$mmol^.L^{-1}$ of lactate from the previous stage during the submaximal test) for 60-120 min, 3-4/wk over

8 wk) consisting of continuous cycling at a fixed power. The training intensity prescription was

reassessed during the first training session of week 3, 5, and 7 and modified accordingly. At the

commencement of training participants began training for 60 min per session and this increased up to

120 min per session in week 7.

Testing procedures. Participants were required to avoid any vigorous exercise for the 48 h preceding

each performance test (72 h for the skeletal muscle biopsy), and to avoid caffeine consumption for at

least 8 h prior to each test. Tests were performed at a similar time of the day throughout the study to

avoid variations caused by changes in circadian rhythm.

GXT. Graded exercise tests (GXTs) were performed on an electronically-braked cycle ergometer

(Lode Excalibur v2.0, Groningen, The Netherlands) to determine maximal oxygen uptake ($V\Box O_{2max}$)

and maximal aerobic power ($W\Box_{max}$). A GXT design of 1-min incremental stages was utilised,

aiming to attain a total testing duration of 9 to 11 minutes. Breath-by-breath recordings of $V\Box O_2$,

$V\Box CO_2$, and $V\Box_E$ were acquired throughout the GXT using a metabolic analyser (Quark

Cardiopulmonary Exercise Testing, Cosmed, Italy). Upon exhaustion, a 5-min recovery was given to

participants, and a verification exhaustive bout was performed at an intensity equivalent to 90% of

$W\Box_{max}$ to confirm the highest measured $V\Box O_{2max}$ (Jamnick et al., 2018). The $W\Box_{max}$ was determined

as the average power of the last minute completed.

Submaximal test. Once the initial GXT was completed, the ventilatory parameters obtained ($V\Box O_2$,

$V\Box CO_2$, $V\Box_E$) were used to estimate VT1. Once estimated, the submaximal test started at 40 W

lower than the estimated VT1 and the intensity was increased 10 W every 3 min (stage length) until

the LT1 was surpassed and identified. The test was stopped at a blood lactate of 2 $mmol^.L^{-1}$ as the

LT1 had been reached in all cases. Antecubital venous blood was taken in the last 15 s of each stage

and was instantly analysed in duplicate using a blood lactate analyser (YSI 2300 STAT Plus, YSI,

USA).

Physical activity and nutritional controls. Participants were requested to maintain a normal dietary

pattern and physical activity throughout the study. To minimise the variability in muscle metabolism,

participants were provided with a standardised dinner (55 kJ/kg of body mass (BM), providing 2.1 g

carbohydrate/kg BM, 0.3 fat/kg BM, and 0.6 g protein/kg BM) and breakfast (41 kJ/kg BM,

providing 1.8 g carbohydrate/kg BM, 0.2 g fat/kg BM, and 0.3 g protein/kg BM) to be consumed 15

h and 3 h before the biopsies, respectively.

Muscle biopsies and single muscle fibre isolation. A biopsy needle with suction was used to obtain

vastus lateralis muscle biopsies under local anaesthesia (1% xylocaine) at pre training and three days

post training. After being cleaned of excess blood, connective tissue, and fat, muscle biopsies were

divided as follows: one piece of muscle was placed in RNALater® (R0901, Sigma Aldrich,

Australia) on the biopsy day and stored at 4 °C, and then separated on ice into single fibres within a

few days. Forty fibres from each biopsy were individually isolated, and cut into half, with one half of

the fibre stored in RNA*Later™* (R0901, Sigma Aldrich, Australia)) and the remaining fibre halves

were placed in a Laemmli solution (a solubilising buffer) (Murphy, 2011). The Laemmli solution

was composed of 4% (v/v) sodium dodecyl sulfate (SDS), a thiol agent of 10% (v/v) ß-

mercaptoethanol, 20% (v/v) glycerol, 0.125 M tris-hydroxymethyl-aminomethane (tris)-HCl; and

0.015 % (v/v) bromophenol blue, pH 6.8. Each fibre was diluted with 5 µL of 3x the solubilising

buffer 2:1 (v/v) with 10 µL of 1x Tris-Cl (pH 6.8) (Murphy, 2011; Wyckelsma *et al.*, 2015). These

fibre segments were then immediately frozen in liquid nitrogen and stored at ⁻80 °C.

Single Muscle Fibre Analysis – Immunoblotting

Muscle Fibre Typing (Dot Blotting)

To assess muscle fibre types, PVDF membranes were activated in 95% ethanol for 15 to 60 s and

then equilibrated for 2 min in transfer buffer (25 mM Tris, 192 mM glycine, pH 8.3 and 20%

methanol). The wet membrane was then placed on a stack of filter paper (one to two pieces soaked in

transfer buffer on top of one dry piece). The single fibre samples in the Laemmli solution were then

thawed and vortexed, but not centrifuged to avoid pelleting and hence loss of any of the skeletal

muscle protein (Murphy, 2011). Samples were spotted to a specific part of the membrane in aliquots

equating to ~ 1/8 of a fibre segment (i.e., 1 µL) using a multi-channel pipette. This was repeated

twice, once for type I fibre detection and another time for type II fibre detection. After complete

absorption of samples, the membrane was placed on top of a dry piece of filter paper to dry for 2 to 5

min before being reactivated in 95% ethanol for 15 to 60 s and equilibrated in transfer buffer for 2

min. After three quick washes in Tris-buffered saline-Tween (TBST), the membrane was blocked in

5% non-fat milk in TBST (blocking buffer) for 5 min at room temperature. Following blocking, the

membrane was rinsed with TBST and then incubated in MHCIIa (#A4.74, Developmental Studies

Hybridoma Bank [DSHB]) or MHCI (#A4.840, DSHB) antibody overnight at 4 ℃ with gentle

rocking. On the second day, membranes were washed in TBST and then incubated in secondary

antibody IgG (MHCIIa, #A32723, ThermoFisher Scientific) or IgM (MHCI, # Cat # A-21042 4,

ThermoFisher Scientific) at room temperature for 1 h with rocking. Lastly, membranes were washed

in TBST and then exposed to Clarity enhanced chemiluminescence reagent (BioRad, Hercules, CA,

USA), imaged (ChemiDoc MP, BioRad), and analysed for signal density (ImageLab 5.2.1, BioRad)

(See example image in Figure 1a). Using images of all the membranes, it was possible to determine

the fibre type of each sample (I or IIa) or if no MHC protein was present (Christiansen et al., 2019),

which would indicate unsuccessful collection of a fibre segment. Samples with signal density in both

type I and IIa membranes were discarded (i.e., I/IIa hybrid, two different fibres in the same sample).

Muscle glycogen fibre-type histochemistry analysis

Muscles were embedded in O.C.T. compound (Tissue-Tek), frozen in liquid nitrogen-cooled

isopentane, stored at ⁻80 °C, and cut into 10 µm thick cryosections with a cryostat (Leica)

maintained at –20 °C.

Fibre type-specific glycogen depletion was imaged under brightfield using periodic acid-Schiff stain

(PAS), similar to the methodology described elsewhere (Schaart et al., 2004). As such, 10 µm thick

muscle cryosections were serially sectioned onto cleaned glass slides and subsequently fixed in 3.7%

formaldehyde in 90% ethanol for 60 min at 4˚C. After this fixation, sections were then pre-treated

for 5 min with 1% periodic acid (Sigma-Aldrich, Australia) in milliQ-water followed by a washing

step for 1 min in tap water and a wash dip for 5 s in milliQ-water. The slides were then applied with

15% diluted Schiff's reagent in PBS (Sigma-Aldrich, Australia) and incubated for 15 min at room

temperature. The Schiff's reagent was diluted to 15% based on prior optimisation, as when applied

without any dilution the dye intensity saturated the signal. After Schiff's reagent, the sections were

washed for 5 s in milliQ-water followed by a 10-min rinse with tap water. Thereafter, sections were

washed (3 x 5 min) in PBS (137 mmol.L^{-1} sodium chloride, 3 mmol.L^{-1} potassium chloride, 8

mmol.L^{-1} sodium phosphate dibasic and 3 mmol.L^{-1} potassium phosphate monobasic, pH of 7.4).

Slides were then dried before adding ~ 15 µL of PBS per section and mounting with cover slips. The

slides were then viewed and imaged in high resolution using an Olympus BX51 fluorescence

microscope (Olympus Corporation, Tokyo, Japan) and Cell F software (Olympus). All images were

obtained with the x10 objective and captured under bright-field mode.

The second serially sectioned slide underwent immunofluorescence staining of MHC expression
performed with 1:25 primary antibodies against MHCI (BA-F8) and MHCIIa (BF-35)
(Developmental Studies Hybridoma Bank, University of Iowa), whereas secondary antibodies were
Alexa fluor 350 IgG_{2b} 1:500 (blue) and Alexa fluor 488 IgG_{2b} 1:500 (green) (Invitrogen). Antibody
cocktail configurations and immunofluorescence staining procedures were followed according to
Bloemberg et al. (Bloemberg and Quadrilatero, 2012). Once completely dry, the sections were
incubated (blocked) with 10% goat serum (product no: 50197Z, ThermoFisher Scientific) for one
hour. The excess goat serum was then removed, and sections were incubated with primary fibre type
antibodies over night at 4 ºC. The following morning, the primary antibodies were washed off in
milliQ water (3 x 5 minutes). Once the slides were fully dry, they were incubated with fluoro-
conjugated secondary antibodies for 90 minutes. Slides were washed further with milliQ water (3 x 5
minutes) post incubation and dried before adding ~ 15 µL of PBS per section and mounting with
cover slips. The slides were then viewed and imaged in high resolution using an Olympus BX51
fluorescence microscope (Olympus Corporation, Tokyo, Japan) and Cell F software (Olympus). All
images were obtained with the x10 objective and analysis of images was performed using Image J
software (National Institutes of Health, Maryland, USA) (See Figure 3a for example images).

Citrate Synthase (CS) Activity Assay.

CS activity was performed on a 96-well plate by combining 5 µL of a 2 µg$^.$ µL^{-1} muscle lysate, 40 µL
of 3 mM acetyl CoA, and 25 µL of 1 mM DTNB to 165 µL of 100 mM Tris buffer (pH 8.3).
Following 15 µL of 10 mM oxaloacetic acid was added to the plate mixture, protected from the light,
and immediately placed in the spectrophotometer at 30 °C (xMark Microplate Spectrophotometer,
Bio-Rad Laboratories Pty Ltd, Gladesville, NSW, AUS). Absorbance was recorded at 412 nm every
15 s for 6 min after 30 s of mixture agitation. CS activity was calculated from the steepest part of the
curve and reported as mol$^.$kg protein$^{-1.}$h^{-1}.

Blood Lactate and pH measurements
Antecubital venous blood samples (~ 1 mL) were collected pre and post the first exercise training
session from a cannula inserted in the antecubital vein for the determination of venous blood [H$^+$]
and lactate concentrations using a blood-gas analyser (ABL 800 FLEX, Radiometer Copenhagen).
679
680

 Transmission Electron Microscopy.

Skeletal muscle samples were fixed overnight at 4 °C with 0.2 M sodium cacodylate– buffered, 2.5%

glutaraldehyde, and 2% paraformaldehyde. Fixed samples were rinsed with 0.1 M sodium

cacodylate, and postfixed with ferricyanide-reduced osmium tetroxide (1% OsO4, 1.5% K3

[Fe(CN)6], and 0.065 M cacodylate buffer) for 2 h at 4°C. The postfixed samples, were processed by

the Monash Ramaciotti Centre for Cryo-Electron Microscopy, were rinsed with distilled water and

then stored overnight in 70% ethanol. Dehydration was performed by graduated ethanol series (80%,

90%, 95%, 100%, and 100%; 10 min each) and propylene oxide (100% and 100%; 5 min each).

Samples were infiltrated with Araldite 502/Embed 812 by graduated concentration series in

propylene oxide (25% for 1 h, 33% for 1 h, 50% overnight; 66% for 4 h, 75% for 4 h, 100%

overnight; and 100% for 5 h) and then polymerised at 60 °C for 48 h. Embedded samples were

sectioned using an Ultracut UCT ultramicrotome (Leica Biosystems) equipped with a 45° diamond

knife (Diatome) to cut 75-nm ultrathin sections. The grids were stained at room temperature using

2% aqueous uranyl acetate (5 min) and Reynolds lead citrate (3 min) before routine imaging. All

TEM imaging was performed at 80 kV on a Jeol JEM-1400Plus and recorded with a Matataki flash

camera and DigitalMicrograph (Version 1.71.38) acquisition software.

Proteome Sample preparation

Fibres were placed into 5 µL of 2x SDS solubilising buffer (0.125 M Tris·Cl, pH 6.8, 4% SDS, 10%

glycerol, 4 M urea, 5% mercaptoethanol, 0.001% bromophenol blue) diluted 2 times (vol/vol) with 1

x Tris·Cl, pH 6.8. Fibres were pooled according to fibre type and further lysed with heating at 95 °C

for 10 min. The pooled fibres were then sonicated for 20 min (Bioruptor, Diagenode, 20 cycles of 30

s). Reduction and alkylation of disulfides was performed by adding chloroacetamide (40 mM, CAA,

Sigma) and further incubated for 30 min at 50 °C. The SDS lysate was acidified with 12% aqueous

phosphoric acid at 1:10 for a final concentration of ~ 1.2% phosphoric acid and mixed. This step was

essential as the proteins are filtered at this pH. The high percentage of methanol can then precipitate

the protein out, which will be retained on top of the S-Trap™ (ProtiFi). Following this, 350 µL of S-

Trap™ buffer (90% Methanol (MeOH) and 100 mM triethylammounium bicarbonate (TEAB),

$C_7H_{17}NO_3$) was added to the acidified lysis buffer (final pH 7.1). A colloidal protein particulate was

instantly formed in this step. With the S-Trap™ micro column in a 1.7 mL tube for flow through, the

acidified SDS lysate/MeOH S-Trap™ buffer mixture was added into the micro column. The micro

column was then centrifuged at 6500 rpm for 30 s until all SDS lysate/ S-Trap™ buffer had passed

through the S-Trap™ column. Protein was then trapped within the protein-trapping matrix of the spin

column. The captured protein was then washed with 350 µL S-Trap™ buffer with centrifugation and

washing repeated three times. The spin column was then transferred to a fresh 1.7 mL tube (this

aided in preventing contamination of the digestion). For best results, the S-Trap™ micro column was

rotated 180 degrees between the centrifugation washes. The S-Trap™ micro column was then moved

to a clean 1.7 mL sample tube for the digestion with the protease (trypsin and LysC added at a

concentration of 1:50 in 125 µL of 50 mM digestion buffer) into the top of the micro column. The

sample was then centrifuged at a low speed of 1000 rpm for 30 s and any solution that passes

through is returned to the top of the column (the protein-trapping matrix is highly hydrophilic and

will absorb the solution; however, it was important to ensure there was no bubble at the top of the

protein trap). The column was then transferred to another fresh tube and incubated at 37 °C overnight

(~ 16 hours). For 96-well plates, the protease combination was added at a concentration of 1:25 in

125 µL of 50 mM digestion buffer and incubated for 1 hour at 47 °C.

Peptides were eluted with 80 µL each of digestion buffer (50 mM TEAB) and then 0.2% aqueous

formic acid was added to the S-Trap™ protein trapping matrix centrifuged at 3600 rpm for 60 s for

each elution. Hydrophobic peptides were recovered with an elution of 80 µL 60% (v/v) acetonitrile

containing 0.2% formic acid and then centrifuged at 6500 rpm for 60 s. Elutions were pooled.

Further peptide purification was performed as required with 2x SDB-RPS disc stage tips that were

prepared and had sample loaded onto the stage-tip. Peptides were then centrifuged through the

column at 1500 g for 3 min. Stage tips were washed with 100 uL of 90% isopropanol (C_3H_8O)

containing 1% TFA and then washed again with 0.2% TFA in 5% acetonitrile and centrifuged for 4

min following each wash. Peptides were then eluted by adding 100 uL of 60% acetonitrile containing

5% ammonium hydroxide and centrifuged for 4 min. Samples were lyophilised down to dryness with

the SpeedVac (CentriVap Benchtop Centrifugal Vacuum Concentrator, # 7810038, VWR) and

reconstituted for labelling and subsequent MS analysis.

TMTpro Labelling

TMTpro Label Reagents (ThermoFisher) were equilibrated to room temperature and 20 µL of

anhydrous acetonitrile was added to each tube. Dried samples were reconstituted in 0.5 M HEPES

buffer, pH 8.5. The desired amount of TMTpro label was added and incubated at 20 °C for 1 h with

shaking (1000 rpm). The reaction was then quenched by adding a final concentration of 0.25% of

hydroxylamine to the peptide and TMT mixture, and further incubated at 20 °C for 30 min. Samples

were lyophilised down to dryness with the SpeedVac and reconstituted in loading buffer. Each

sample was fractionated into 16 fractions using basic pH reverse phase C18 liquid chromatography.

Peptides were subjected to basic-pH reverse-phase high pressure liquid chromatography (HPLC)

fractionation. Labelled peptides were solubilised in buffer A (10 mM ammonium hydroxide) and

separated on an Agilent HpH Poroshell120 C18 column (2.7 μm particles, 2.1 mm i.d., and 5 cm in

length).

TMT Labelling Strategy

Multiplexing techniques have recently extended from 11 to 16 plex, which not only increases sample

throughput but accommodates complex experimental designs such as this training study. This

provided the ability to label 16 samples in one go and then merge as one sample through the LC-MS.

A 16-plex also allows for an expanded number of treatments, such as replicates, and dose-response

or time-course measurements can be analysed in the same experiment with basically no missing

values across all samples and extending the statistical power across the entire system (Li et al.,

2020). The main study used TMTpro 16-plex version with a different reporter and mass normaliser

than earlier TMT versions (i.e., TMT-6 plex). A limitation of tag-based proteomic strategies is ion

interference-related ratio distortion resulting from fragmentation and analysis of background ions co-

isolated with those of interest. Each sample is differentially labelled, such that when pooled the

signal-to-noise values of sample-specific reporter ions represent the relative abundance of each

protein. As such, the degree of ion interference by the level of TMT signal detected in channels

where a specific protein should be absent can be assessed (Brenes et al., 2019). To best

accommodate for possible reporter ion interferences, we followed a sampling strategy that is

illustrated in Figure S2g. Reporter ion interference (RII) targets were classified according to a typical

product data sheet for 16-plex TMTpro Label Reagents from ThermoFisher Scientific.

Liquid Chromatography-MS/MS Acquisition

Peptides were loaded onto a 2-cm PepMap trap column (ThermoFisher) and separated using a

PepMap 75 μm × 50 cm column (ThermoFisher) with a gradient of 2–30% MeCN containing 0.1%

FA over 120 min at 250 nL/min and at 40 °C. The Orbitrap Fusion mass spectrometer was operated

with the following parameters: an MS1 scan was acquired from 375–1575 m/z (120,000 resolution,

2e5 AGC, 50 ms injection time) followed by MS2 data-dependent acquisition with collision-induced

dissociation (CID) and detection in the ion trap (4e3 AGC, 150 ms injection time, 30 NCE, 1.6 m/z

quadrupole isolation width, 0.25 activation Q). To quantify TMTpro reporter ions, a synchronous

precursor selection MS3 scan was performed with higher-energy collisional dissociation (HCD) and

detection in the orbitrap (120 to 750 m/z, 1e5 AGC, 250 ms injection time, 60 NCE, 2.5 m/z

isolation width). The total cycle time was set to 2.5 s. The acquired raw data was analysed with

Proteome Discoverer 2.4 (ThermoFisher) and the non-normalised protein reporter intensity was

exported to Excel and further analysed in R.

Bioinformatic analysis of proteomics data.

Clean up and normalisation

Only proteins that had a high FDR confidence, were not deemed a contaminant, and had greater than

one unique peptide, were used for analysis. Proteins having < 70% valid values were removed and

the remaining missing data was imputed using k nearest neighbour (knn) method from

the *impute* package in R. Normalisation was performed using a combination of trimmed mean of M

values (TMM), sample loading (SL), and ComBat (v0.0.4) methods (Johnson et al., 2007; Liu et al.,

2004; Plubell et al., 2017; Robinson and Oshlack, 2010). Sample loading adjusts each TMT

experiment to an equal signal per channel and TMM uses the average or median signal that is

typically applied to use a single multiplicative factor to further adjust the samples to each other.

Further to this, ComBat allows users to adjust for batch effects in datasets where the batch covariate

is known, using previously described methodology (Johnson *et al.*, 2007).

Differential expression analysis
800

Differential expression analysis was performed using the R package *limma* (v3.48.3) (Ritchie *et al.*,

2015) on the relative protein abundances after first performing the normalisation technique described

above. Differential expression values were determined by comparing the expression. To focus on

proteins specifically associated with fibre type, the differential expression comparisons were first

filtered to include proteins differentially expressed between type I and type II fibres before exercise

(adjusted P value < 0.05). This established the baseline of the proteins that were significantly

different between the two fibre types prior to exercise and allowed for comparisons between our data

and the data of Murgia et. al. (2017) and Deshmukh et. al. (2020) (Figures 1 and 2). Following this,

comparisons were made to identify any significant effects (Fusion Factor P value < 0.05) exercise

produced on each of the fibre types for each type of exercise (MICT type I Pre vs Post, MICT type II

Pre vs Post, SIT type I Pre vs Post, SIT type II Pre vs Post; Figure 4). Finally, comparisons were

completed between fibre types on samples post exercise (MICT Post type I vs type II and SIT Post

type I vs type II, Figure 4), to distinguish and compare significant (adjusted P value < 0.05)

differences between the fibre types after exercise. Mitochondrial proteins were identified by

annotation using both the Integrated Mitochondrial Protein Index (IMPI) and the Mitocarta 3.0
database (Rath et al., 2020).
The heatmap presented in Figure 1g was completed using the *ComplexHeatmap* package (v.2.8.0),
with hierarchical clustering using the "average" method. The heatmap presented in Figure 7c was
created using the native R heatmap function and clustered according to the OXPHOS complex
subunits as identified in MitoCarta 3.0. For visualisations of differential expression comparisons,
volcano plots were presented in Figure 1f, 2,e,f, 5a,b and used the resulting adjusted P values from
the *limma* analysis, which can be found in Table S1, S2, S3 and S4. Volcanos from Figure 4 a,b used
the P values from the Fusion Factor calculation found in Table S4. Profile Plots presented in Figures
6d, 7a,b,d and S1b,c, were created by calculating a mean of the group in question (complex or
functional group as identified by MitoCarta 3.0) with all proteins identified in that group coloured
and a background displayed of all mitochondrial proteins (with an exception for Figure 7a, which
used all proteins identified). Statistical significance between Pre and Post training values within the
profile plots was completed using a paired t-test and the default adjustment for multiple hypothesis
testing. Venn diagrams featured in Figure 1c, 2a were created using the *VennDiagram* (v1.7.1)
package.

To better characterise the proteomic changes, gene ontology and enrichment analysis were
performed by investigating non-random associations between proteins and over-represented Gene
Ontology (GO). For Figures 1b, we calculated the average number of proteins in each pooled
sample, the proteins identified within the entire dataset of high confidence proteins and compared
them to the entire human genome to determine how many were identified in specific GO CC terms
compared to the human genome. Figure 1d and 2b compared the 846 and 272 proteins respectively,
which were common between the three datasets investigated and the top five GO CC terms for the
846 proteins were visualised, with the top five GO BP terms used for the 273. These analyses were
conducted using Database for Annotation, Visualization and Integrated Discovery (DAVID) (Huang
et al., 2009). Network plots and WikiPathways are presented in Figure 5e and f respectively. The
network plot (Figure 5f) was completed using only statistically significant (adjusted P value < 0.05)
results for the comparisons of type I vs II values post both MICT and SIT, whereas the
Wikipathways BubblePlot uses all results for the same comparisons. These analyses were completed
using the R package *ClusterProfiler* (v4.0.5).

Mitochondrial Normalisation

As the mitochondrial protein expression identified a difference in the mitochondrial content between the two fibre types before exercise, we employed a previously reported normalisation strategy (Granata *et al.*, 2021). This normalisation strategy removes bias introduced by the differences in mitochondrial content and allows the investigation of these adaptations when this bias was removed. Mitochondrial proteins identified through IMPI and MitoCarta 3.0 were subset from the rest of the data, and the same normalisation protocol as reported above was applied.

Data availability and code availability

- Source data to interpret, verify, and extend this research is provided with this paper. The mass spectrometry proteomics data has been deposited in the ProteomeXchange Consortium via the PRIDE partner repository under accession code PXD036010.

- The R scripts used for all omics analyses described above are deposited on GitHub and available through https://doi.org/10.5281/zenodo.7227801. There are no restrictions placed on accessibility of this code.

Author Contributions Statement

D.J.B., J.B, E.R., and C.G. conceptualised and designed the study. J.B. carried out the training study and sample collection. E.R., J.B., and C.H. performed the experiments. E.R and S.C. performed muscle glycogen fibre-type histochemistry analysis. V.O and G.R. performed the electron microscopy experiments. N. J. C. performed the statistical and bioinformatic analysis. D.J.B., E.R., and N.J.C. wrote the manuscript. E.R., J.B., C.H, R.B.S. D.A.S, C.G., N.J.C and D.J.B. edited and revised the manuscript. All persons designated as authors qualify for authorship, and all those qualifying for authorship are listed. All authors have read and approved the final manuscript.

Acknowledgments

We thank Iresha Hanchapola from Monash Proteomics & Metabolomics Facility for proteomic

supplies and instrument support and acknowledge Associate Professor Javier Diaz Lara from the

University de Navarra for support with dot blotting.

Funding

This study was supported by grants from the Australian Research Council (Discovery Projects

DP140104165 and DP160102176 to D.J.B) the Australian Physiological Society (Ph.D. grant to J.B),

and the Australian National Health and Medical Research Council (NHMRC Fellowship

GNT2009732 to D.A.S.). J.B. was supported by a Victoria University International Postgraduate

Research Scholarship.

888
889   **Competing Interests Statement**

890   The authors declare no conflict of interest.

References

Bergh, U., Thorstensson, A., Sjödin, B., Hulten, B., Piehl, K., and Karlsson, J. (1978). Maximal oxygen uptake and muscle fiber types in trained and untrained humans. Med Sci Sports *10*, 151-154.

Bishop, D.J., Botella, J., Genders, A.J., Lee, M.J., Saner, N.J., Kuang, J., Yan, X., and Granata, C. (2019a). High-Intensity Exercise and Mitochondrial Biogenesis: Current Controversies and Future Research Directions. Physiology (Bethesda, Md.) *34*, 56-70. 10.1152/physiol.00038.2018.

Bishop, D.J., Botella, J., and Granata, C. (2019b). CrossTalk opposing view: Exercise training volume is more important than training intensity to promote increases in mitochondrial content. The Journal of Physiology *0*. 10.1113/JP277634.

Bishop, D.J., Granata, C., and Eynon , N. (2014). Can we optimise the exercise training prescription to maximise improvements in mitochondria function and content? . Biochimica et Biophysica Acta *1840*, 1266–1275.

Bloemberg, D., and Quadrilatero, J. (2012). Rapid Determination of Myosin Heavy Chain Expression in Rat, Mouse, and Human Skeletal Muscle Using Multicolor Immunofluorescence Analysis. PLOS ONE *7*, e35273. 10.1371/journal.pone.0035273.

Brenes, A., Hukelmann, J., Bensaddek, D., and Lamond, A.I. (2019). Multibatch TMT Reveals False Positives, Batch Effects and Missing Values. Mol Cell Proteomics *18*, 1967-1980. 10.1074/mcp.RA119.001472.

Christiansen, D., MacInnis, M.J., Zacharewicz, E., Xu, H., Frankish, B.P., and Murphy, R.M. (2019). A fast, reliable and sample-sparing method to identify fibre types of single muscle fibres. Scientific Reports *9*, 6473. 10.1038/s41598-019-42168-z.

Deshmukh, A.S., Murgia, M., Nagaraj, N., Treebak, J.T., Cox, J., and Mann, M. (2015). Deep Proteomics of Mouse Skeletal Muscle Enables Quantitation of Protein Isoforms, Metabolic Pathways, and Transcription Factors Molecular and Cell Proteomics *14*, 841-853.

Deshmukh, A.S., Steenberg, D.E., Hostrup, M., Birk, J.B., Larsen, J.K., Santos, A., Kjøbsted, R., Hingst, J.R., Schéele, C.C., Murgia, M., et al. (2021). Deep muscle-proteomic analysis of freeze-dried human muscle biopsies reveals fiber type-specific adaptations to exercise training. Nature Communications *12*, 304. 10.1038/s41467-020-20556-8.

Dudley, G.A., Abraham, W.M., and Terjung, R.L. (1982). Influence of exercise intensity and duration on biochemical adaptations in skeletal muscle. Journal of applied physiology: respiratory, environmental and exercise physiology *53*, 844-850. 10.1152/jappl.1982.53.4.844.

Egan, B., Dowling, P., O'Connor, P.L., Henry, M., Meleady, P., Zierath, J.R., and O'Gorman, D.J. (2011). 2-D DIGE analysis of the mitochondrial proteome from human skeletal muscle reveals time coursedependent remodelling in response to 14 consecutive days of endurance exercise training Proteomics *11*, 1413–1428.

Egan, B., and Zierath, J.R. (2013). Exercise metabolism and the molecular regulation of skeletal muscle adaptation. Cell Metab. *17*, 162-184. 10.1016/j.cmet.2012.12.012.

Galpin, A.J., Raue, U., Jemiolo, B., Trappe, T.A., Harber, M.P., Minchev, K., and Trappe, S. (2012). Human skeletal muscle fiber type specific protein content. Analytical biochemistry *425*, 175-182. 10.1016/j.ab.2012.03.018.

Gillen, J.B., Martin, B.J., MacInnis, M.J., Skelly, L.E., Tarnopolsky, M.A., and Gibala, M.J. (2016). Twelve Weeks of Sprint Interval Training Improves Indices of Cardiometabolic Health Similar to Traditional Endurance Training despite a Five-Fold Lower Exercise Volume and Time Commitment. PLoS One *11*, e0154075. 10.1371/journal.pone.0154075.

Glancy, B., and Balaban, R.S. (2011). Protein composition and function of red and white skeletal muscle mitochondria. American journal of physiology. Cell physiology *300*, C1280-C1290. 10.1152/ajpcell.00496.2010.

Godin, R., Ascah, A., and Daussin, F.N. (2010). Intensity-dependent activation of intracellular
signalling pathways in skeletal muscle: role of fibre type recruitment during exercise. J Physiol *588*,
4073-4074. 10.1113/jphysiol.2010.195925.
Gollnick, P.D., Piehl, K., and Saltin, B. (1974). Selective glycogen depletion pattern in human
muscle fibres after exercise of varying intensity and at varying pedalling rates. J Physiol *241*, 45-57.
10.1113/jphysiol.1974.sp010639.
Granata, C., Caruana, N.J., Botella, J., Jamnick, N.A., Huynh, K., Kuang, J., Janssen, H.A., Reljic,
B., Mellett, N.A., Laskowski, A., et al. (2021). High-intensity training induces non-stoichiometric
changes in the mitochondrial proteome of human skeletal muscle without reorganisation of
respiratory chain content. Nature Communications *12*, 7056. 10.1038/s41467-021-27153-3.
Granata, C., Jamnick, N.J., and Bishop, D.J. (2018). Training-induced mitochondrial adaptations.
Sports Med., 1-20.
Granata, C., Oliveira, R.S., Little, J.P., Renner, K., and Bishop, D.J. (2016a). Training intensity
modulates changes in PGC-1α and p53 protein content and mitochondrial respiration, but not
markers of mitochondrial content in human skeletal muscle. Faseb j *30*, 959-970. 10.1096/fj.15-
276907.
Granata, C., Oliveira, R.S.F., Little, J.P., Renner, K., and Bishop, D.J. (2016b). Mitochondrial
adaptations to high-volume exercise training are rapidly reversed after a reduction in training
volume in human skeletal muscle   The Federation of American Societies for Experimental Biology
Journal *30*, 1-11.
Greenhaff, P.L., Nevill, M.E., Soderlund, K., Bodin, K., Boobis, L.H., Williams, C., and Hultman, E.
(1994). The metabolic responses of human type I and II muscle fibres during maximal treadmill
sprinting. The Journal of Physiology *478*, 149-155. https://doi.org/10.1113/jphysiol.1994.sp020238.
Hargreaves, M., and Spriet, L.L. (2020). Skeletal muscle energy metabolism during exercise. Nature
Metabolism *2*, 817-828. 10.1038/s42255-020-0251-4.
Hawley, J.A., Hargreaves, M., Joyner, M.J., and Zierath, J.R. (2014). Integrative biology of exercise.
Cell *159*, 738-749. 10.1016/j.cell.2014.10.029.
Henneman, E. (1957). Relation between Size of Neurons and Their Susceptibility to Discharge.
Science (New York, N.Y.) *126*, 1345. 10.1126/science.126.3287.1345.
Henneman, E., and Olson, C.B. (1965). Relations Between Structure and Function in the Design of
Skeletal Muscle. Journal of Neurophysiology *28*, 581-598. 10.1152/jn.1965.28.3.581.
Holloszy, J.O., and Coyle, E.F. (1984). Adaptations of skeletal muscle to endurance exercise and
their metabolic consequences. Journal of applied physiology: respiratory, environmental and exercise
physiology *56*, 831-838. 10.1152/jappl.1984.56.4.831.
Holloway, K.V., O'Gorman, M., Woods, P., Morton, J.P., Evans, L., Cable, N.T., Goldspink, D.F.,
and Burniston, J.G. (2009). Proteomic investigation of changes in human vastus lateralis muscle in
response to interval-exercise training. Proteomics *9*, 5155-5174. 10.1002/pmic.200900068.
Hostrup, M., Onslev, J., Jacobson, G.A., Wilson, R., and Bangsbo, J. (2018). Chronic β2-
adrenoceptor agonist treatment alters muscle proteome and functional adaptations induced by high
intensity training in young men. The Journal of Physiology *596*, 231-252. 10.1113/JP274970.
Howald, H., Hoppeler, H., Claassen, H., Mathieu, O., and Straub, R. (1985). Influences of endurance
training on the ultrastructural composition of the different muscle fiber types in humans. Pflugers
Archiv : European journal of physiology *403*, 369-376. 10.1007/bf00589248.
Huang, D.W., Sherman, B.T., and Lempicki, R.A. (2009). Systematic and integrative analysis of
large gene lists using DAVID bioinformatics resources. Nature protocols *4*, 44-57.
10.1038/nprot.2008.211.
Ingjer, F. (1979). Effects of endurance training on muscle fibre ATP-ase activity, capillary supply
and mitochondrial content in man. The Journal of Physiology *294*, 419-432.
10.1113/jphysiol.1979.sp012938.

Jackman, M.R., and Willis, W.T. (1996). Characteristics of mitochondria isolated from type I and
type IIb skeletal muscle. American Journal of Physiology-Cell Physiology *270*, C673-C678.
10.1152/ajpcell.1996.270.2.C673.
Jamnick, N.A., Botella, J., Pyne, D.B., and Bishop, D.J. (2018). Manipulating graded exercise test
variables affects the validity of the lactate threshold and V̇ O2peak. PLoS One *13*, e0199794.
Jamnick, N.A., Pettitt, R.W., Granata, C., Pyne, D.B., and Bishop, D.J. (2020). An Examination and
Critique of Current Methods to Determine Exercise Intensity. Sports Medicine. 10.1007/s40279-020-
01322-8.
Jansson, E., Sjodin, B., and Tesch, P. (1978). Changes in muscle fibre type distribution in man after
physical training. A sign of fibre type transformation? Acta Physiol Scand *104*, 235-237.
10.1111/j.1748-1716.1978.tb06272.x.
Johnson, W.E., Li, C., and Rabinovic, A. (2007). Adjusting batch effects in microarray expression
data using empirical Bayes methods. Biostatistics *8*, 118-127. 10.1093/biostatistics/kxj037.
Kiessling, K.H., Piehl, K., and Lundquist, C.G. (1971). Effect of Physical Training on Ultrastructural
Features in Human Skeletal Muscle. In Muscle Metabolism During Exercise: Proceedings of a
Karolinska Institutet Symposium held in Stockholm, Sweden, September 6–9, 1970 Honorary guest:
E Hohwü Christensen, B. Pernow, and B. Saltin, eds. (Springer US), pp. 97-101. 10.1007/978-1-
4613-4609-8_9.
Kiessling, K.H., Pilström, L., Bylund, A.C., Saltin, B., and Piehl, K. (1974). Enzyme Activities and
Morphometry in Skeletal Muscle of Middle-Aged Men after Training. Scandinavian journal of
clinical and laboratory investigation *33*, 63-69. 10.3109/00365517409114199.
Kristensen, D.E., Albers, P.H., Prats, C., Baba, O., Birk, J.B., and Wojtaszewski, J.F. (2015). Human
muscle fibre type-specific regulation of AMPK and downstream targets by exercise. J Physiol *593*,
2053-2069. 10.1113/jphysiol.2014.283267.
Lamboley, C.R., Wyckelsma, V.L., Dutka, T.L., McKenna, M.J., Murphy, R.M., and Lamb, G.D.
(2015). Contractile properties and sarcoplasmic reticulum calcium content in type I and type II
skeletal muscle fibres in active aged humans. J Physiol *593*, 2499-2514. 10.1113/jp270179.
Larsen, S., Nielsen, J., Hansen, C.N., Nielsen, L.B., Wibrand, F., Stride, N., Schroder, H.D.,
Boushel, R., Helge, J.W., Dela, F., and Hey-Mogensen, M. (2012). Biomarkers of mitochondrial
content in skeletal muscle of healthy young human subjects. J Physiol *590*, 3349-3360.
10.1113/jphysiol.2012.230185.
Leary, S.C., Lyons, C.N., Rosenberger, A.G., Ballantyne, J.S., Stillman, J., and Moyes, C.D. (2003).
Fiber-type differences in muscle mitochondrial profiles. American Journal of Physiology-
Regulatory, Integrative and Comparative Physiology *285*, R817-R826. 10.1152/ajpregu.00058.2003.
Li, J., Van Vranken, J.G., Pontano Vaites, L., Schweppe, D.K., Huttlin, E.L., Etienne, C.,
Nandhikonda, P., Viner, R., Robitaille, A.M., Thompson, A.H., et al. (2020). TMTpro reagents: a set
of isobaric labeling mass tags enables simultaneous proteome-wide measurements across 16
samples. Nature Methods *17*, 399-404. 10.1038/s41592-020-0781-4.
Liu, H., Sadygov, R.G., and Yates, J.R., 3rd (2004). A model for random sampling and estimation of
relative protein abundance in shotgun proteomics. Analytical chemistry *76*, 4193-4201.
10.1021/ac0498563.
MacInnis, M.J., and Gibala, M.J. (2017). Physiological adaptations to interval training and the role
of exercise intensity. J Physiol *595*, 2915-2930. 10.1113/jp273196.
MacInnis, M.J., Skelly, L.E., and Gibala, M.J. (2019). CrossTalk proposal: Exercise training
intensity is more important than volume to promote increases in human skeletal muscle
mitochondrial content. J Physiol *597*, 4111-4113. 10.1113/jp277633.
MacInnis, M.J., Zacharewicz, E., Martin, B.J., Haikalis, M.E., Skelly, L.E., Tarnopolsky, M.A.,
Murphy, R.M., and Gibala, M.J. (2017). Superior mitochondrial adaptations in human skeletal
muscle after interval compared to continuous single-leg cycling matched for total work. The Journal
of Physiology *595*, 2955-2968. 10.1113/JP272570.

Martin, W.H., 3rd, Dalsky, G.P., Hurley, B.F., Matthews, D.E., Bier, D.M., Hagberg, J.M., Rogers, M.A., King, D.S., and Holloszy, J.O. (1993). Effect of endurance training on plasma free fatty acid turnover and oxidation during exercise. The American journal of physiology *265*, E708-714. 10.1152/ajpendo.1993.265.5.E708.

Mendell, L.M. (2005). The size principle: a rule describing the recruitment of motoneurons. Journal of Neurophysiology *93*, 3024-3026. 10.1152/classicessays.00025.2005.

Miller, B.F., and Hamilton, K.L. (2012). A perspective on the determination of mitochondrial biogenesis. American journal of physiology. Endocrinology and metabolism *302*, E496-E499. 10.1152/ajpendo.00578.2011.

Morgan, T.E., Cobb, L.A., Short, F.A., Ross, R., and Gunn, D.R. (1971). Effects of Long-Term Exercise on Human Muscle Mitochondria. In Muscle Metabolism During Exercise: Proceedings of a Karolinska Institutet Symposium held in Stockholm, Sweden, September 6–9, 1970 Honorary guest: E Hohwü Christensen, B. Pernow, and B. Saltin, eds. (Springer US), pp. 87-95. 10.1007/978-1-4613-4609-8_8.

Morissette, M.P., Susser, S.E., Stammers, A.N., O'Hara, K.A., Gardiner, P.F., Sheppard, P., Moffatt, T.L., and Duhamel, T.A. (2014). Differential regulation of the fiber type-specific gene expression of the sarcoplasmic reticulum calcium-ATPase isoforms induced by exercise training. Journal of applied physiology (Bethesda, Md. : 1985) *117*, 544-555. 10.1152/japplphysiol.00092.2014.

Murgia, M., ,, Nagaraj, N., Deshmukh, A.S., Zeiler, M., Cancellara, P., Moretti, I., Reggiani, C., Schiaffino, S., and Mann, M. (2015). Single muscle fiber proteomics reveals unexpected mitochondrial specialization. EMBO Reports *16*, 387-395.

Murgia, M., Toniolo, L., Nagaraj, N., Ciciliot, S., Vindigni, V., Schiaffino, S., Reggiani, C., and Mann, M. (2017). Single Muscle Fiber Proteomics Reveals Fiber-Type- Specific Features of Human Muscle Aging Cell Reports *19*, 2396–2409.

Murphy, R.M. (2011). Enhanced technique to measure proteins in single segments of human skeletal muscle fibers: fiber-type dependence of AMPK-alpha1 and -beta1. Journal of applied physiology (Bethesda, Md. : 1985) *110*, 820-825. 10.1152/japplphysiol.01082.2010.

Neufer, P.D., Bamman, Marcas M., Muoio, Deborah M., Bouchard, C., Cooper, Dan M., Goodpaster, Bret H., Booth, Frank W., Kohrt, Wendy M., Gerszten, Robert E., Mattson, Mark P., et al. (2015). Understanding the Cellular and Molecular Mechanisms of Physical Activity-Induced Health Benefits. Cell Metabolism *22*, 4-11. https://doi.org/10.1016/j.cmet.2015.05.011.

O'Connell, J.D., Paulo, J.A., O'Brien, J.J., and Gygi, S.P. (2018). Proteome-Wide Evaluation of Two Common Protein Quantification Methods. Journal of proteome research *17*, 1934-1942. 10.1021/acs.jproteome.8b00016.

Ohlendieck, K. (2010). Proteomics of skeletal muscle differentiation, neuromuscular disorders and fiber aging. Expert review of proteomics *7*, 283-296. 10.1586/epr.10.2.

Ohlendieck, K. (2011). Skeletal muscle proteomics: current approaches, technical challenges and emerging techniques. Skeletal Muscle *1*, 1-15.

Ohlendieck, K. (2013). Proteomics of exercise-induced skeletal muscle adaptations. OA Sports Medicine *1*, 1-12.

Ørtenblad, N., Nielsen, J., Boushel, R., Söderlund, K., Saltin, B., and Holmberg, H.-C. (2018). The Muscle Fiber Profiles, Mitochondrial Content, and Enzyme Activities of the Exceptionally Well-Trained Arm and Leg Muscles of Elite Cross-Country Skiers. Frontiers in physiology *9*, 1031-1031. 10.3389/fphys.2018.01031.

Pappireddi, N., Martin, L., and Wühr, M. (2019). A Review on Quantitative Multiplexed Proteomics. Chembiochem *20*, 1210-1224. 10.1002/cbic.201800650.

Plubell, D.L., Wilmarth, P.A., Zhao, Y., Fenton, A.M., Minnier, J., Reddy, A.P., Klimek, J., Yang, X., David, L.L., and Pamir, N. (2017). Extended Multiplexing of Tandem Mass Tags (TMT) Labeling Reveals Age and High Fat Diet Specific Proteome Changes in Mouse Epididymal Adipose Tissue. Mol Cell Proteomics *16*, 873-890. 10.1074/mcp.M116.065524.

Rath, S., Sharma, R., Gupta, R., Ast, T., Chan, C., Durham, T.J., Goodman, R.P., Grabarek, Z., Haas,
M.E., Hung, W.H.W., et al. (2020). MitoCarta3.0: an updated mitochondrial proteome now with sub-
organelle localization and pathway annotations. Nucleic acids research. 10.1093/nar/gkaa1011.
Ritchie, M.E., Phipson, B., Wu, D., Hu, Y., Law, C.W., Shi, W., and Smyth, G.K. (2015). limma
powers differential expression analyses for RNA-sequencing and microarray studies. Nucleic acids
research *43*, e47. 10.1093/nar/gkv007.
Robinson, M.D., and Oshlack, A. (2010). A scaling normalization method for differential expression
analysis of RNA-seq data. Genome Biology *11*, R25. 10.1186/gb-2010-11-3-r25.
Robinson, M.M., Dasari, S., Konopka, A.R., Johnson, M.L., Manjunatha, S., Esponda, R.R., Carter,
R.E., Lanza, I.R., and Nair, K.S. (2017). Enhanced Protein Translation Underlies Improved
Metabolic and Physical Adaptations to Different Exercise Training Modes in Young and Old
Humans. Cell Metab *25*, 581-592. 10.1016/j.cmet.2017.02.009.
Sahlin, K., Mogensen, M., Bagger, M., Fernström, M., and Pedersen, P.K. (2007). The potential for
mitochondrial fat oxidation in human skeletal muscle influences whole body fat oxidation during
low-intensity exercise. Am J Physiol Endocrinol Metab *292*, E223-230.
10.1152/ajpendo.00266.2006.
Sale, D.G. (1987). Influence of exercise and training on motor unit activation. Exercise and sport
sciences reviews *15*, 95-151.
Scalzo, R.L., Peltonen, G.L., Binns, S.E., Shankaran, M., Giordano, G.R., Hartley, D.A., Klochak,
A.L., Lonac, M.C., Paris, H.L., Szallar, S.E., et al. (2014). Greater muscle protein synthesis and
mitochondrial biogenesis in males compared with females during sprint interval training. Faseb j *28*,
2705-2714. 10.1096/fj.13-246595.
Schaart, G., Hesselink, R.P., Keizer, H.A., van Kranenburg, G., Drost, M.R., and Hesselink, M.K.
(2004). A modified PAS stain combined with immunofluorescence for quantitative analyses of
glycogen in muscle sections. Histochem Cell Biol *122*, 161-169. 10.1007/s00418-004-0690-0.
Schiaffino, S., Rossi, A.C., Smerdu, V., Leinwand, L.A., and Reggiani, C. (2015). Developmental
myosins: expression patterns and functional significance. Skeletal Muscle *5*, 22. 10.1186/s13395-
015-0046-6.
Schiaffino, S.a.R., C. (2011). Fiber Types in Mammalian Skeletal Muscles. Physiol Rev *91*, 1447–
1531. 10.1152.
Schneider, C.A., Rasband, W.S., and Eliceiri, K.W. (2012). NIH Image to ImageJ: 25 years of image
analysis. Nature Methods *9*, 671-675. 10.1038/nmeth.2089.
Scribbans, T.D., Edgett, B.A., Vorobej, K., Mitchell, A.S., Joanisse, S.D., Matusiak, J.B., Parise, G.,
Quadrilatero, J., and Gurd, B.J. (2014). Fibre-specific responses to endurance and low volume high
intensity interval training: striking similarities in acute and chronic adaptation. PLoS One *9*, e98119.
10.1371/journal.pone.0098119.
Shaw, C.S., Swinton, C., Morales-Scholz, M.G., McRae, N.L., Erftemeyer, T., Aldous, A., Murphy,
R.M., and Howlett, K.F. (2020). The impact of exercise training status on the fibre type specific
abundance of proteins regulating intramuscular lipid metabolism. Journal of Applied Physiology.
10.1152/japplphysiol.00797.2019.
Shepherd, S.O., Cocks, M., Meikle, P.J., Mellett, N.A., Ranasinghe, A.M., Barker, T.A.,
Wagenmakers, A.J.M., and Shaw, C.S. (2017). Lipid droplet remodelling and reduced muscle
ceramides following sprint interval and moderate-intensity continuous exercise training in obese
males. International Journal of Obesity *41*, 1745-1754. 10.1038/ijo.2017.170.
Shepherd, S.O., Cocks, M., Tipton, K.D., Ranasinghe, A.M., Barker, T.A., Burniston, J.G.,
Wagenmakers, A.J., and Shaw, C.S. (2012). Preferential utilization of perilipin 2-associated
intramuscular triglycerides during 1 h of moderate-intensity endurance-type exercise. Experimental
physiology *97*, 970-980. 10.1113/expphysiol.2012.064592.
Shepherd, S.O., Cocks, M., Tipton, K.D., Ranasinghe, A.M., Barker, T.A., Burniston, J.G.,
Wagenmakers, A.J., and Shaw, C.S. (2013). Sprint interval and traditional endurance training

increase net intramuscular triglyceride breakdown and expression of perilipin 2 and 5. J Physiol *591*,
657-675. 10.1113/jphysiol.2012.240952.
Skelly, L.E., Gillen, J.B., Frankish, B.P., MacInnis, M.J., Godkin, F.E., Tarnopolsky, M.A., Murphy,
R.M., and Gibala, M.J. (2021). Human skeletal muscle fiber type-specific responses to sprint interval
and moderate-intensity continuous exercise: acute and training-induced changes. Journal of applied
physiology (Bethesda, Md. : 1985) *130*, 1001-1014. 10.1152/japplphysiol.00862.2020.
Suriano, R., Edge, J., and Bishop, D. (2010). Effects of cycle strategy and fibre composition on
muscle glycogen depletion pattern and subsequent running economy. Br J Sports Med *44*, 443-448.
10.1136/bjsm.2007.046029.
Tan, R., Nederveen, J.P., Gillen, J.B., Joanisse, S., Parise, G., Tarnopolsky, M.A., and Gibala, M.J.
(2018). Skeletal muscle fiber-type-specific changes in markers of capillary and mitochondrial
content after low-volume interval training in overweight women. Physiological reports *6*.
10.14814/phy2.13597.
van Loon, L.J., Greenhaff, P.L., Constantin-Teodosiu, D., Saris, W.H., and Wagenmakers, A.J.
(2001). The effects of increasing exercise intensity on muscle fuel utilisation in humans. The Journal
of physiology *536*, 295-304. 10.1111/j.1469-7793.2001.00295.x.
Vollestad, N.K., and Blom, P.C. (1985). Effect of varying exercise intensity on glycogen depletion in
human muscle fibres. Acta Physiol Scand *125*, 395-405. 10.1111/j.1748-1716.1985.tb07735.x.
Wilson, J.M., Loenneke, J.P., Jo, E., Wilson, G.J., Zourdos, M.C., and Kim, J.S. (2012). The effects
of endurance, strength, and power training on muscle fiber type shifting. J Strength Cond Res *26*,
1724-1729. 10.1519/JSC.0b013e318234eb6f.
Wyckelsma, V.L., McKenna, M.J., Serpiello, F.R., Lamboley, C.R., Aughey, R.J., Stepto, N.K.,
Bishop, D.J., and Murphy, R.M. (2015). Single-fiber expression and fiber-specific adaptability to
short-term intense exercise training of Na+-K+-ATPase α- and β-isoforms in human skeletal muscle.
Journal of Applied Physiology *118*, 699-706. 10.1152/japplphysiol.00419.2014.
Xiao, Y., Hsiao, T.H., Suresh, U., Chen, H.I., Wu, X., Wolf, S.E., and Chen, Y. (2014). A novel
significance score for gene selection and ranking. Bioinformatics *30*, 801-807.
10.1093/bioinformatics/btr671.

1166

1167 **Figure 1 Legend**
1168

a) Workflow for characterisation of pooled human skeletal muscle fibres. Single fibres
from human biopsy muscle samples were taken and fibre typing was confirmed through
dot blotting. After optimisation, 6 single fibres were pooled and proteins were extracted
using a S-Trap™ sample processing spin column before being labelled with a TMT
labelling protocol. LC-MS/MS was subsequently performed to quantify proteins for data
analysis.

b) Percentage of proteins identified within the samples in comparison with the human
genome. The bars display the percentage of proteins identified in the particular gene
ontology cellular compartment (GOCC) group based on i) the average number of
proteins identified within individual pooled samples (pink), ii) proteins identified within
the entire dataset of high confidence proteins (orange) and iii) known annotated proteins
in the human genome, not detected in entire single fibre dataset (grey). Identification of
GOCC terms was completed using the DAVID functional annotation tool.

c) A Venn diagram displaying the proteins identified in the current study alongside two
published single fibre papers. Due to differences in experimental design, and to make an
appropriate comparison, Murgia et al. (2017) was filtered for young participants only
with 70% valid values and Deshmukh et al. (2021) was filtered for 70% valid values.

d) Proteins identified by all three single fibre studies in Figure 1c underwent GOCC
analysis. The top 5 terms of the 846 proteins identified in the intersect of Deshmukh et
al., Reisman et al. and Murgia et al. are presented. The top GOCC terms and
corresponding log P value were identified using the DAVID software (Huang *et al.*,
2009) and computed by a Fisher's Exact Test.

e) Multidimensional Scaling (MDS) Plot used to show the distances among fibre types for
PRE exercise type I (blue) and type II (red) samples.

f) Volcano plot illustrating the fold change distribution and significance of proteins in the
PRE samples between type I and type II fibres. Points were labelled based on whether
they were identified as differentially expressed uniquely within our study (compared
with the differentially expressed proteins identified in Deshmukh et al. 2021 and Murgia
et al. 2017) or if the corresponding $-\log_{10}P$ value was greater than or equal to 5.0.

g) Heatmap displaying z-score values of differentially expressed proteins when comparing
type II to type I within the PRE samples. Samples predominately cluster together by fibre
type I (blue) or type II (red).

1201

1202 **Figure 2 Legend**

1203

a) A Venn diagram displaying the number of mitochondrial proteins annotated according to
the mitochondrial protein database Mitocarta 3.0 (Rath *et al.*, 2020) in the current study,
alongside two published human single-fibre datasets (Deshmukh et al. 2021 and Murgia
et al. 2017). Only 70% valid values pre intervention were annotated to Mitocarta 3.0.

b) Top biological processes (GOBPs) in which the identified mitochondrial proteins
common to all three studies illustrated in Fig 2a are mainly involved. The bars represent
Log_{10}P values, where *P* represents the significance of each GOBP enriched by the
mitochondrial proteins. The *P*-value was computed using DAVID software (Huang *et*
al., 2009).

c) Illustration of mitochondrial proteins that were differentially expressed between type II
and type I fibres in the PRE samples. Proteins that were identified as differentially
expressed following standard normalisation and mitochondrial normalisation of the data
are also highlighted in green.

d) Comparison of Mean Mitochondrial Protein Expression in PRE exercise samples for type
I (blue) and type II fibres (red).

e) Volcano plot illustrating the fold-change distribution of proteins in the PRE samples and
the significant differences in protein expression between type I and type II fibres. Points
were labelled if they were mitochondrial proteins (according to Mitocarta 3.0; (Rath *et*
al., 2020)) and identified as differentially expressed.

f) Volcano plot illustrating the fold-change distribution of proteins in the PRE samples and
the significant differences in protein expression between type I and type II fibres. Points
were labelled if they were mitochondrial proteins (according to Mitocarta 3.0; (Rath *et*
al., 2020)) and identified as differentially expressed following mitochondrial
normalisation (Granata *et al.*, 2021).

1228

1229

Figure 3 Legend

a) The 8-week exercise training regime and methodology. Skeletal muscle biopsies were taken before week 1 and after week 8. Participants were randomly allocated into either the moderate-intensity continuous training (MICT) or sprint interval training (SIT) group. Grey bars indicate the number of training sessions per week. Following biopsies, muscle fibres were isolated into a single fibres. Single fibre lysates were fibre typed according to dot blotting and subsequently pooled by fibre type (six fibres). Samples then underwent LC-MS/M analysis.

b) Bar graphs displaying fold differences between the moderate-intensity continuous training (MICT) and sprint interval training (SIT) for exercise intensity ($\%W_{max}$) and total training volume (J). Also shown are bar graphs displaying the change in blood lactate concentration ($mmol \bullet L^{-1}$) and hydrogen ion (H^+) concentration ($mol \bullet L^{-1}$) from pre to post exercise following the first moderate-intensity continuous exercise (MICT) and the first sprint interval exercise (SIT) session. Mean ± standard deviation ; * adjusted P value < 0.05.

c) *Histochemistry* – the effects of a single session of exercise on fibre-type specific glycogen depletion patterns in cross-sections of human vastus lateralis muscle samples collected before and after either moderate-intensity continuous exercise (MICT) or sprint interval exercise (SIT). The top panels display immunofluorescent fibre-type staining for myosin heavy chain content, with type I muscle fibres (MHC7) identified as blue and type II fibres (MHC2) identified as green. The bottom panels display serial sections that have been stained for glycogen (periodic acid-Schiff staining). Bars = 200 µm.

d) Illustration of the hierarchal recruitment pattern of type I to IIa to IIx/b skeletal muscle fibres in relation to exercise intensity. Moderate-intensity continuous exercise (MICT) is indicated in purple and sprint interval exercise (SIT) in green, with type I fibres and type II fibres in blue and red, respectively.

Figure 4 Legend

a) Volcano plots illustrating the fold-change distribution of proteins that were
differentially expressed in type I fibres following MICT or SIT. Proteins are coloured
based on a significant fusion factor (Xiao et al., 2014) and a Log2Fold Change of >
0.2.

b) Volcano plots illustrating the fold change distribution of proteins that were
differentially expressed in type II fibres following MICT or SIT. Proteins are coloured
based on a significant fusion factor (Xiao et al., 2014) and a Log2Fold Change of >
0.2.

c) Venn Diagram displaying the number of proteins that were differentially expressed in
type I or type II fibres following MICT, according to the fusion factor (Xiao et al.,
2014).

d) Venn Diagram displaying number of proteins that were differentially expressed in
type I or type II fibres following SIT, according to the fusion factor (Xiao et al.,
2014).

1282

1283 **Figure 5 Legend**

1284
a) Volcano plot illustrating the fold-change distribution of proteins between type I and II
fibres in the POST MICT samples. Points were labelled based on whether they were
identified as differentially expressed, according to adjusted P values from the *limma*
analysis.

b) Volcano plot illustrating the fold-change distribution of proteins between type I and II
fibres in the POST SIT samples. Points were labelled based on whether they were
identified as differentially expressed, according to adjusted P values from the *limma*
analysis.

c) Venn Diagrams displaying number of proteins identified as higher in type I or type II
fibres for both exercise types, with the proteins that appear in both datasets seen in the
overlap. Individual proteins are listed in supplemental Table 5, Tab 3.

d) Venn Diagram displaying the number of proteins that were differentially expressed
between type I and type II fibres in the PRE samples (n=23 participants), compared
with the number of differentially expressed proteins between type I and type II fibres
in the POST MICT (n = 11) and SIT samples (n=12).

e) *EnrichPlot* dot plot results of protein expression in type I vs type II fibres for the POST MICT and SIT samples. Pathway terms were identified using WikiPathways. The top 15 terms according to adjusted P-values are displayed. Red dots identify pathway terms found higher in type II fibres, while blue dots indicate pathway terms found higher in type I fibres. Saturation of each colour is proportional to the average log2 fold change for each exercise and fibre group. The size of each dot indicates how many proteins correspond to each pathway term. For all proteins, the corresponding $\log_2$ fold change and P-values from the differential expression analysis were used. All terms identified can be found in supplemental Table 5, Tab 5.

f) *ClusterProfiler* network plot using differentially expressed (adjusted P value < 0.05) terms. Pathway terms were identified using WikiPathways. Colours indicate in which exercise and fibre group the proteins in that WikiPathway term were found. The size of each circle indicates how many proteins corresponded to each term. The displayed terms were manually selected for readability of the figure. All terms identified can be found in supplemental Table 5, Tab 6.

Figure 6 Legend

a) A representative images of a skeletal muscle samples analysed using transmission electron microscopy (TEM) pre- and post-training for both MICT and SIT. Note the difference in the abundance of mitochondria. Bars = 0.5 µm

b) Violin plots of change (Δ) in mitochondrial volume density (% Mito_{VD}) from pre to post training following moderate-intensity continuous training (MICT - purple) and sprint interval training (SIT - green).

c) Violin plots of change (Δ) in citrate synthase activity (CS; $\text{mol.h}^{-1}.\text{kg protein}^{-1}$) from pre to post training following moderate-intensity continuous training (MICT - purple) and sprint interval training (SIT - green).

d) Scaled profile plots showing the relative abundance of mitochondrial proteins (grey) pre and post exercise in the MICT (purple) and SIT (green) groups for each fibre type, and the mean relative abundance (black) of the proteins in each group.

Data in d & e are shown as Δ mean z-score. * indicates P values for significance (*: P <= 0.01, **: P <= 0.001, ***: P <= 0.0001, ****: P <= 0.00001) by paired t-test from pre to post training.

1335

1336 **Figure 7 Legend**

1337

a) Scaled profile plots with the application of mitochondrial normalisation showing changes in the relative abundance of mitochondrial proteins (grey) pre and post training in the MICT and SIT groups for each fibre type. The mean of each group is indicated by the black line. Data is shown as Δ mean z-score.

b) Scaled profile plots with the application of mitochondrial normalisation showing changes in the relative abundance of the subunits in each of the oxidative phosphorylation (OXPHOS) complexes (CI to CV), as well as the mitochondrial ribosome, in type I and type II fibre types in response to both types of training (MICT and SIT). The mean of each group (all proteins identified in each pathway using MitoCarta 3.0) is indicated by the black line. Data is shown as Δ mean z-score. * indicates P value for significance (*: $P \leq 0.01$, **: $P \leq 0.001$, ***: $P \leq 0.0001$, ****: $P \leq 0.00001$) based on a paired t-test of the pre to post training values.

c) Scaled profile plots with the application of mitochondrial normalisation showing changes in the relative abundance of proteins involved in Fatty Acid Oxidation, the tricarboxylic acid cycle, and mitochondrial dynamics (all proteins identified in pathway using MitoCarta 3.0) in type I and type II fibre types in response to both types of training (MICT and SIT). The mean of each group is indicated by the black line. Data is shown as Δ mean z-score. * indicates P value for significance (*: $P \leq 0.01$, **: $P \leq 0.001$, ***: $P \leq 0.0001$, ****: $P \leq 0.00001$) based on a paired t-test of the pre to post training values.

d) Heatmap displaying z-score values of protein subunits of the OXPHOS complexes, as well as the mitochondrial ribosome, in each of the two training groups (MICT and SIT), in type I and type II fibre types pre and post training, with the application of mitochondrial normalisation.

Supplemental Figure 1 Legend

a) Heatmap displaying z-score values of protein subunits of the OXPHOS complexes, as
well as the mitochondrial ribosome, in each of the two training groups (MICT and
SIT), in type I and type II fibre types pre and post training, without the application of
mitochondrial normalisation.

b) Scaled profile plots without the application of mitochondrial normalisation showing
changes in the relative abundance of the subunits in each of the oxidative
phosphorylation (OXPHOS) complexes (CI to CV), as well as the mitochondrial
ribosome, in type I and type II fibre types in response to both types of training (MICT
and SIT). The mean of each group (all proteins identified in each pathway using
MitoCarta 3.0) is indicated by the black line. Data is shown as Δ mean z-score. *
indicates P value for significance (*: $P \leq 0.01$, **: $P \leq 0.001$, ***: $P \leq 0.0001$, ****:
$P \leq 0.00001$) based on a paired t-test of the pre to post training values.

c) Scaled profile plots without the application of mitochondrial normalisation showing
changes in the relative abundance of proteins involved in Fatty Acid Oxidation, the
tricarboxylic acid cycle, and mitochondrial dynamics (all proteins identified in
pathway using MitoCarta 3.0) in type I and type II fibre types in response to both
types of training (MICT and SIT). The mean of each group is indicated by the black
line. Data is shown as Δ mean z-score. * indicates P value for significance (*: $P \leq$
0.01, **: $P \leq 0.001$, ***: $P \leq 0.0001$, ****: $P \leq 0.00001$) based on a paired t-test of
the pre to post training values.

1385
1386
1387

Supplementary Figure 2 Legend

a) Number of proteins quantified in the training study dataset by our proteomic workflow in all pooled single-fibre samples. Samples are arranged according to TMT batches (A-H). Samples are labelled according to participant ID (Z to A), training type (S or M), Biopsy time (pre or post), and fibre type (I or II))

b) Reproducibility of protein intensities (intensity-based absolute quantification; IBAQ) for replicates of type I skeletal muscle fibre samples. Core mitochondrial proteins (OXPHOS – NDUFA8, SDHA, URCRC2, COX5B, ATP5B, IDH3B and CKM) are highlighted in yellow, in addition to sarcomeric proteins (MYH 7 (blue), ACTA1 (orange)).

c) Reproducibility of protein intensities (intensity-based absolute quantification; IBAQ) for replicates of type II skeletal muscle fibre samples. Core mitochondrial proteins (OXPHOS – NDUFA8, SDHA, URCRC2, COX5B, ATP5B, IDH3B and CKM) are highlighted in yellow, in addition to MYH2 (green).

d) Relative abundance of myosin heavy chains (MYH) in type I samples pooled by dot blotting, according to the percentage of the MYH7 (blue = type I), MYH2 (red = type-IIa) MYH1 (brown = type-IIx), and MYH4 (purple = type IIb) protein isoforms detected.

e) Relative abundance of myosin heavy chains (MYH) in type II samples pooled by dot blotting, according to the percentage of the MYH7 (blue = type I), MYH2 (red = type-IIa) MYH1 (brown = type-IIx), and MYH4 (purple = type IIb) protein isoforms detected.

f) Study recruitment flow chart and final group sizes for the groups that completed 8 weeks of moderate-intensity interval training (MICT) or sprint interval training (SIT). Following the initial recruitment of 28 participants, 2 men from each group dropped out of the study due to time constraints and one fibrotic muscle sample was removed from the MICT group muscle analysis.

g) Sampling strategy when using a 96-well plate to best accommodate for possible reporter ion interferences. Reporter ion interference (RII) targets were classified according to a typical product data sheet for 16-plex TMTpro Label Reagents from ThermoFisher Scientific.

h) Hierarchical visualisation of the statistical comparisons performed in this study.

Description of Additional Supplementary Files

File name: Supplementary Table 1

Description: Participant characteristics, Raw intensities for all proteomic data, and log

intensities for all identified proteins in all samples following the normalisation described in

the methods. Data also includes gene ontology cellular compartment (GOCC) analysis for

pooled single fibre analysis for percentage of cellular compartment identified (shown in

Figure 1b) and the top 5 GOCC and corresponding $\log_{10}$ P value identified using the DAVID

software for proteins identified by key single fibre proteomic studies (shown in Figure 1c).

Further analysis includes the protein lists presented in the Venn diagram of Figure 1c for

identified proteins within the current study, along with the studies of Deshmukh et al. and

Murgia et al. (Tab 5), along with the top GOCC terms for the common proteins identified in

all 3 studies (Fig 1d, Tab 6). Statistical analysis is lastly presented via logged intensities of

proteins within the PRE type I and type II fibre samples. The table also displays calculations

completed to calculate the 'Fusion Factor' used in Deshmukh et al. (2021) and the

corresponding significance. Further detail is provided in the legend tab of the linked Excel

file.

1437

File name: Supplementary Table 2

Description: The list of mitochondrial proteins (according to Mitocarta 3.0) in common to all

previous single fibre human studies pre intervention. Further analysis is presented in the table

output by the DAVID software to identify top gene ontology biological processes (GOBP)

groups of the mitochondrial proteins identified in over 70% of the total sample in all three

studies listed (Deshmukh et al., Murgia et al., and the current study). Further analysis is

presented with the average of all mitochondrial proteins in each individual. The mean of

averages were calculated and compared between PRE training type I fibres and type II fibres.

Mitochondrial protein expression (MPE) calculation is subsequently presented for identified

mitochondrial proteins within the study in each comparison. Data normalised for

mitochondrial content for identified mitochondrial proteins within the study for the PRE type

I vs II comparison and presented in the volcano plots (Figure 2f). Further detail provided in

the legend tab of the linked Excel file.

1451

1452 **File name: Supplementary Table 3**

1453

Description: Fold Difference from Pre to Post exercise for blood concentration of H^+ and
lactate results.

File name: Supplementary Table 4

Description: The normalised Data (Log2) with *limma* and Fusion Factor calculations for
MICT Type I and Type II - PRE vs POST. In addition, Normalised Data (Log$_2$ with *limma*
and Fusion Factor Calculations for SIT type I & type II - PRE vs POST. Proteins identified in
type I vs type II fibres POST MICT and corresponding GO Biological Processes results.
Parallel analysis is presented for POST SIT. Further detail is provided in the legend tab of
the linked Excel file.

1463

File name: Supplementary Table 5

Description: Log intensities for type I versus type II fibres in POST MICT and SIT samples
following normalisation. GO Enrichment (Biological Processes) completed with DAVID on
the common and unique MICT and SIT proteins identified when comparing Post I vs II
samples. Further corresponding GOBP analysis is also made proteins identified as
differentially expressed between type I and II fibres in the PRE samples. Further enrichment
analysis is undertaken via WikiPathways enrichment analysis and Network analysis of
differences in protein abundance between type I & type II fibres in the POST samples for
MICT & SIT (analysed separately), using enrichplot & clusterprofiler on the *limma* results
displayed in Tab 1 and 2 of this file. Further detail provided in the legend tab of the linked
Excel file.

1475

File name: Supplementary Table 6

Description: Differences from Pre to Post training for Mitochondria Volume Density (%)
and Citrate Synthase activity (mol•h^{-1}•kg protein^{-1}) for MICT & SIT. Analysis presented for
scaled intensities (z-scores) and means for known mitochondrial proteins for all samples
following standard normalisation. Scaled means for each complex in each sample group are
also presented in addition to the mean change for each comparison of training and fibre type.
Further detail is provided in the legend tab of the linked Excel file.

1483

File name: Supplementary Table 7

Description: Scaled intensities and means for known mitochondrial proteins for all samples
following mitochondrial normalisation. Protein functional classes and metabolic pathways
presented with the scaled means (z-scores) for each functional group (defined by MitoCarta

3.0) in each sample group in addition to the mean change for each comparison of training and

fibre type with mitochondrial normalisation applied. Further detail provided in the legend tab

of the linked Excel file.

File name: Supplementary Table 8

Description: Protein functional classes and metabolic pathways presented with the scaled

means (z-scores) for each functional group (defined by MitoCarta 3.0) in each sample group

in addition to the mean change for each comparison of training and fibre type. Further detail

is provided in the legend tab of the linked Excel file.

1496

File name: Supplementary Table 9

Description: The number of proteins quantified in the training study dataset by our proteomic

workflow in all pooled single-fibre samples. Raw TMT Batch ID is presented and arranged

according to TMT batches (A-H). Samples are labelled according to participant ID (Z-C),

training type (S for SIT or M for MICT), training time (pre or post) and fibre type (I or II). In

addition, reliability between replicates is presented - List of Intensity reported by intensity-

based absolute quantification (IBAQ) for replicates and corresponding gene names of pooled

type I fibres (n = 6 fibres) and type II fibres. Furthermore, the relative TMT intensity and

percentage abundance of myosin heavy chains (MYH) for type I and type II samples pooled

by dot blotting is presented.

1507

a)

PRE TRAINING BIOPSY — MICT OR SIT — 8 WEEKS OF EXERCISE TRAINING — Week 1, Week 2, Week 3, Week 4, Week 5, Week 6, Week 7, Week 8 — POST TRAINING BIOPSY → Single Fibre Isolation → Single Fibre per Tube (Lysis) → Pooled (Six) Single Muscle Fibres → LC-MS/MS; Fibre Typing by Dot Blotting

b)

Exercise Intensity (% Wmax): MICT, SIT *
Training Volume (J): MICT, SIT *
Δ Blood Lactate Conc. (mmol L⁻¹): MICT, SIT *
Δ Blood [H+] (mol L⁻¹): MICT, SIT *

c)

MICT — PRE, POST; SIT — PRE, POST
200 μm

d)

Moderate Intensity Continuous Training (MICT)
Sprint Interval Training (SIT)

% Muscle Fibre Recruitment vs Exercise Intensity (Low — High)
Type IIx, Type IIa, Type I