miR-148a-3p inhibits osteogenesis by targeting Itga11 via

PI3K/Akt/GSK3/β-catenin pathway

Authors: Xiaojun **Chen**[1#]; Kewei **Rong**[2#]; Wenqing **Han**[1]; Yichuan **Pang**[3*]; Gang **Chai**[1*]

Affiliations:

[1]Department of Plastic and Reconstructive Surgery, Shanghai Ninth People's Hospital, Shanghai Jiao Tong University School of Medicine, Shanghai, 200011, China.

[2]Department of Orthopaedics, Shanghai Ninth People's Hospital, Shanghai Jiao Tong University School of Medicine, Shanghai, 200011, China.

[3]Department of Nuclear Medicine, Shanghai Tenth People's Hospital, Tongji University School of Medicine, Shanghai, 200072, China

[#] These authors contributed equally to this work.

[*] Co-corresponding author

Corresponding Authors:

Gang **Chai**, M.D., Ph.D., Department of Plastic and Reconstructive Surgery, Shanghai Ninth People's Hospital, Shanghai Jiao Tong University School of Medicine, 639 ZhiZaoJu Road, 200011, Shanghai, China, chaig1081@sh9hospital.org.cn.

Yichuan **Pang**, Ph.D., Department of Nuclear Medicine, Shanghai Tenth People's Hospital, Tongji University School of Medicine, 36 Yunxin Road, 200072, Shanghai, China, pangyichuan@126.com

Abstract

In previous research, miR-148a-3p deficiency was observed in bone malformation in hemifacial microsomia. Herein, in this article, we probed into the role of miR-148a-3p in bone physiology by utilizing miR-148a knock-out (KO) mice. Compared with wild-type (WT) or heterozygotic (HE) littermates, miR-148a knock-out mice manifested lower body weight, bone dysplasia with increased bone mass. Through *in-vitro* experiments, in terms of miR-148a-3p overexpression (miRNA mimic transfection) and knockout (primary cells from WT and KO littermates), we found that miR-148a-3p can suppress osteogenesis, either in the ALP activity or bone nodules formation. Afterward, by means of proteomics, combined with RNA-sequencing and prediction databases of microRNA targets (miRDB and TargetScan), nine candidate genes targeted by miR-148a-3p were identified. Among them, only *Itga11* was regulated by mRNA degradation, while the others were modulated via post-transcriptional inhibition. Based on several online databases (GenePaint, BioGPS, STRING), Integrin Subunit Alpha 11 (Itga11) was suggested to play an essential role in osteogenesis and it was confirmed as one direct target of miR-148a-3p by dual-luciferase reporter assay. Meanwhile, gene set enrichment analysis (GSEA) indicated activation of PI3K-Akt signaling pathway and WNT signaling pathway in miR-148a KO mice. The thereafter western blot confirmed that PI3K/Akt/GSK3/β-catenin signaling pathway was involved. Taken together, we demonstrated that miR-148a-3p can inhibit osteogenesis by targeting Itga11 via PI3K/Akt/GSK3/β-catenin pathway.

Keywords: miR-148a-3p; osteogenesis; multi-omics; Itga11

Introduction

miRNAs are small endogenous non-coding RNAs composed of ~22 nucleotides that can direct posttranscriptional modification of mRNA targets. They have been confirmed to be essential in govern stem cell fates during embryonic development[1]. They get involved in various physiological and pathological processes as well[2,3]. In previous research, by taking advantage of the bio-samples collected from bilateral mandibles of hemifacial microsomia (OMIM: 164210) patients, miR-148a-3p was detected to be significantly decreased in the affected mandible compared with the unaffected side.

MiR-148a-3p is highly conserved in various species[4], and has already been suggested as an important regulator in bone microenvironment. As an epigenetic regulator of bone homeostasis, miR-148a-3p was consistently upregulated in osteoporosis, and considered as a potential new plasma-based biomarker osteoporosis/ osteoporotic fracture[5]. It correlated with bone strength and metabolism as well, which had a negative correlation with total body bone mineral density[6].

Multiple targets of miR-148a-3p have already been found based on *in vitro* experiments. Nevertheless, there was little agreement in results across studies. Moreover, complete opposite trends existed. For example, it was reported that miR-148a-3p can inhibit osteogenic differentiation via p300[7],Wnt5a[8], WNT1, TGFB2, IGF1[6], NRP1[9] or Kdm6b[10], while it can also promote osteogenesis via SMURF1[11]. The reasons why various candidates and tendency were obtained might due to the fact that the target genes regulated by the same miRNA may vary depending on cells or tissues[2].

In order to better understand the role of miR-148a-3p in osteogenesis, miR-148a knock-out (KO) mice were bred in this project. Meanwhile, for the first time, we probed into its regulatory mechanism in a natural *in vivo* environment.

Materials and Methods

Generation and genotype identification of miR-148a Knock-Out (KO) mice:

miR-148a KO mice were obtained from GemPharmatech Co., Ltd. (Jiangsu, China) and housed under SPF condition in Shanghai Ninth People's Hospital, Shanghai Jiao Tong University School of Medicine. The Ethics Committee of Shanghai Ninth People's Hospital has approved all the following animal procedures. Heterozygous male and female mice were mating (the male-to-female ratio was 1:2) to generate miR-148a knock-out (KO), heterozygous (HE), and wild-type (WT) littermates. Ear or tail tissues were obtained for genotyping.

Phenotype identification of miR-148a KO mice: Female mice were used here for the phenotype identification. Every mouse was weighed once a week and euthanatized at 12 weeks old when their mandibles and tibias were dissected for further examination. The length of mandibles was measured via vernier caliper, and X-ray scans were performed to assess mandibular bone mineral density. We also applied μCT to the tibias for more information about bone mass.

Double fluorescent labeling: Calcein (8mg/kg) and Alizarin Red (20mg/kg) were administered separately by intraperitoneal injection at 10 or 11 weeks of age. These male mice were euthanatized at 12 weeks old. Mandibles were dissected and fixed in 75% ethanol. The species were then dehydrated with different concentrations of ethanol and embedded. After cutting and grinding, the sections were observed under a fluorescent microscope (OLYMPUS). The distance between the observed fluorescent bands represented the bone mineral apposition rate.

Paraffin Section: 12-week-old male mice without any treatment were sacrificed. After fixing in 4% paraformaldehyde for 48 hours, the tissues were decalcified in 10% Ethylene Diamine Tetra-acetic Acid (EDTA) for about four weeks. Then the tissues were processed by dehydrating in different concentrations of ethanol, infiltrating with xylene, and embedding with paraffin (Leica, Germany). Sections were cut at 5-μm thickness for further histochemical and immunofluorescent staining. In brief, sections were processed in a sequence of deparaffinization, hydration, antigen retrieval,

permeabilization, blocking, primary antibody incubation (OCN, Servicebio, Catalog#GB11233; Runx2, Servicebio, Catalog#GB11264; Osterix, Abcam, Catalog#ab22552; ITGA11, Abcam, Catalog#ab198826), corresponding secondary antibody incubation, and nuclear staining.

Bone Defect Animal Experiment: Eight-week-old male mice were first anesthetized by 1% pentobarbital. Then the right tibia was exposed, and muscles were dissociated in part. A pilot hole was first made with a drill point in the proximal tibia but away from the growth plate. Then the hole was enlarged with a 1 mm reamer. After irrigation, the skin was sutured. μCT was conducted on the 14[th] day after operation to evaluate the bone healing process.

Cell Culture: Ectomesenchymal stem cells (EMSCs) were obtained from embryos of E13.5 pregnant heterozygous mice as described in Zhao's article[12]. In brief, the mandibular process was dissected, minced into pieces, digested with type I collagenase for two hours, followed centrifugation, resuspension and seeding. EMSCs were cultured in complete media (CM), that is, alpha-MEM supplemented with 10% Fetal Bovine Serum (FBS) and 1% penicillin/streptomycin (p./s.) (Gibco). All cultures were maintained in a 37□ incubator with 5% CO_2.

Cell Transfection: EMSCs were transfected with miR-148a-3p mimic reagents (RiboBio, Guangzhou, China) according to the manufacturer's protocol. In brief, the transfection reagents were first mixed with mimic reagents. After being kept stationary for 20 minutes, the mixture was used for cell transfection for six hours. Then the mixture was replaced with complete media or osteogenic media. The transfection procedure was repeated every three days if long-term culture was needed.

Cell Proliferation Assay: Cell Counting Kit-8 (CCK8, Dojindo, China) was used to examine the role of miR-148a-3p in cell proliferation according to the manufacturer's protocol. In brief, EMSCs were seeded on 96-well plates at a density of $3x10^3$ cells

per well. At the specific time point (24h, 48h, 72h, or 96h, respectively), the standard culture medium was replaced with 100μl CM containing 10μL CCK8 reagent, and then the plates were incubated at 37□ for another two hours. Absorbances were measured at 450nm using the Absorbance Microplate Spectrophotometer (Tecan, China). Absorbance readings were normalized to the control group, and the cell proliferation curve was calculated using GraphPad Prism 8 (GraphPad Software, USA).

Osteoblast differentiation and staining: EMSCs were maintained in CM for normal growth and osteogenic media (CM supplemented with 2 mM β-glycerophosphate, 0.05 mg/ml ascorbic acid, and 10 nM dexamethasone) for osteoblast differentiation. For miR-148a-3p mimic experiments, extra transfection procedures were done before every time the osteogenic medium was changed. Alkaline phosphatase (ALP) staining was performed on the 7[th] day to assay the ALP activity, and Alizarin Red (AR) staining was done on the 21[st] day to evaluate bone nodules formation. The staining protocol was as follows. Cells were first washed three times with PBS, fixed in 4% paraformaldehyde for 20 minutes at room temperature, and stained with ALP solution or 1% AR solution (Beyotime, Shanghai, China) for about 30 minutes. Cells were then washed three times with PBS again, and digital images were captured.

Quantitative real-time polymerase chain reaction (qPCR): For mRNA quantification, total cellular RNA was extracted from cultured cells using Multisource Total RNA Extract Reagent (Axygen, Catalog#11818KD1). The RNA purity and concentration were assessed using Nanodrop ND-1000 (Thermo Fisher). Single-stranded cDNA was reverse transcribed from 500ng total RNA using reverse transcriptase and oligo(dT) primer (Takara, Catalog#RR036). Real-time qPCR amplification was performed using the PrimeScript™ RT-PCR Kit (Takara, Catalog#RR420). *Alp, Col1a1, Runx2, Bglap, Spp1, Tnfsf11* were detected for osteogenesis. All the PCR procedures were performed in a QuantStudio 6 Flex

(Applied Biosystems), and the $2^{-\Delta\Delta CT}$ method was used for analysis.

Primers were as follows, *Alp* (Forward: AGAAGTTCGCTATCTGCCTTGCCT; Reverse: TGGCCAAAGGGCAATAACTAGGGA), *Col1a1* (Forward: GCTCCTCTTAGGGGCCACT; Reverse: CCACGTCTCACCATTGGGG), *Runx2* (Forward: GACTGTGGTTACCGTCATGGC; Reverse: ACTTGGTTTTTCATAACAGCGGA); *Bglap* (Forward: TAGCAGACACCATGAGGACCATCT; Reverse: CCTGCTTGGACATGAAGGCTTTGT), *Spp1* (Forward: AGCAAGAAACTCTTCCAAGCAA; Reverse: GTGAGATTCGTCAGATTCATCCG), *Tnfsf11* (Forward: GTACTTTCGAGCGCAGATGG; Reverse: TCCAACCATGAGCCTTCCAT).

Western blot analysis: Total cellular protein extracts were isolated from cultured EMSCs stimulated with osteogenic media using RIPA lysis buffer (Beyotime, Catalog#P0013C) supplemented with 1% PMSF (Beyotime, Catalog#ST506). Lysates were cleared by centrifugation at 15,000*g for 10 mins at 4°C, and the supernatant was collected. The mixture of extracted proteins and loading buffer (Beyotime, Catalog#P0015B) was denatured at 95℃ for 10 minutes and added to the prefabricated PAGE gels (GenScript, Nanjing, China). Separated proteins were then electroblotted onto polyvinylidene difluoride (PVDF) membranes (Millipore, USA). Following the transfer, membranes were blocked with 5% w/v skimmed milk in TBST for one hour and then incubated at 4□ with primary antibodies (p-PI3K, Abcam, Catalog#ab278545; p-AKT(S473), CST, Catalog#4060S; p-AKT(T308), CST, Catalog#13038S; p-GSK3b(S9), CST, Catalog#5558S; pan-PI3K, CST, Catalog#20584-1-AP; pan-AKT, CST, Catalog#4691S; pan-GSK3, Abcam, Catalog#ab185141; β-catenin, CST, Catalog#8480S; β-actin, Abcam, Catalog#ab8227) overnight. Membranes were washed thrice with TBST and then incubated with fluorescent secondary antibodies for one hour at room temperature. After washing thrice with TBST, the membrane was developed using the Odyssey infrared imaging system (Li-COR, USA).

Omics analysis: Bones from four-week-old WT and KO littermates were collected for omics analysis. RNA sequence was performed in order of RNA extraction (miRNeasy Mini Kit, Qiagen#217004), library construction, and sequencing with Illumina NovaSeq 6000. Proteomics was completed following protein extraction, trypsin digestion, HPLC classification, and liquid chromatography-mass spectrometry (LC-MS). Several online databases or tools were applied for further analysis, miRBD (https://mirdb.org/)[13] and TargetScan (https://www.targetscan.org/mmu_80/)[14] for potential targets prediction, GenePaint (https://gp3.mpg.de/)[15,16] for *in situ* hybridization image of E14.5 mouse embryo, BioGPS (http://biogps.org/)[17] for candidate gene expression profiles in various tissues and cells, STRING (https://cn.string-db.org/)[18] for the protein-protein interaction network, and jvenn (http://jvenn.toulouse.inra.fr/)[19] for Venn diagrams.

Dual-Luciferase Reporter Assay: Reporter plasmids containing ITGA11 3'-UTR region with wild type or mutant miR-148a-3p binding sites (Itga11-MUT) were first constructed in pmiR-RB-REPORT™ (RiboBio). Then plasmids were co-transfected into HEK-293T cell lines with miR-148a-3p Mimic or Negative Control (NC). After culturing for 48 hours, Dual-Glo® Luciferase Assay System (Promega, USA) was used to detect the luciferase activities, and firefly luciferase activities were normalized to renilla ones for the statistic.

Statistical analysis: As specified, all data were presented as mean ± standard deviations of three or more pooled independent experiments. Each dot in the histogram represented one independent data. Unpaired t-test was used for two groups, and analysis of variance (ANOVA) was performed for three groups or more. When ANOVA demonstrated significant differences between groups, multiple comparisons were carried out between the control group and every experimental group via Dunnett's t-test in GraphPad Prism 8. Statistical significance was defined as [*] $P<0.05$, [**] $P<0.01$, [***] $P<0.001$ and [****] $P<0.0001$.

Results

miR-148a Knock-Out (KO) mice manifested bone dysplasia with enhanced osteogenic potential.

Since miR-148a-3p is the main product of pre-miR-148a, miR-148a KO mice (**Fig.S1a**) were utilized to investigate the function of miR-148a-3p in bone formation. Compared with heterozygotic (HE) littermates, KO mice were grossly smaller (**Fig.1a**). The bodyweight of 12-week-old mice was lost as the knock-out of the miR-148a allele, with an average weight of 21.92g in WT, 20.24g in HE, and 18.98g in KO (**Fig.1b**). From X-ray images, mandibular hypoplasia can be observed, both in the body length (13.83±0.27mm versus 13.36±0.34mm, *P*=0.0236) and in the ramus height (5.22±0.29mm versus 4.66±0.14mm, *P*=0.0015) (**Fig.1c**). Meanwhile, enhanced mineralization apposition rate can be observed in the KO mice (**Fig.1d**).

As the mandible largely comprises compact bone, next, we utilized long bones for further investigation. We assessed the bone mass changes through tibia µCT scans. KO mice had remarkably higher bone mass than HE ones, with increased values in bone volume/ total volume (BV/TV), trabecular number (Tb. N), trabecular thickness (Tb. Th), while decreased values in trabecular separation (Tb. Sp) (**Fig.1e**). The expression levels of Osteocalcin (OCN) and Runx2, both markers for osteogenesis, were significantly increased when miR-148a alleles were knocked out, especially near the growth plate (**Fig.1f**). Similar immunostaining results were observed in cranium (**Fig.1g**).

Bone defect animal models were also adopted, where the bone healing process was accelerated in KO mice, in comparison to WT littermates. The mean bone volume/total volume (BV/TV) increased from 0.1575 in the WT group to 0.2891 in the KO group (*P*=0.0090) (**Fig.S1b**).

miR-148a-3p inhibited osteogenesis *in-vitro*.

Primary ectomesenchymal stem cells (EMSCs) were isolated from E13.5 embryos to verify the intrinsic ability to osteogenesis *in vitro*. Compared with EMSCs from WT

littermates, primary EMSCs with miR-148a KO had enhanced osteogenic potential, with more abundant and deeper ALP staining on the fourth day (**Fig.2a**).

miR-148a-3p mimic was then adopted to overexpress the miR-148a-3p expression level in EMSCs extracted from WT pups. After transfection, the miR-148a-3p expression level was significantly upregulated, over 150 folds in EMSCs (**Fig.2b**). The proliferation assay demonstrated no significant difference in cell proliferation ability among the control, NC, and Mimic groups (**Fig.2c**). Nevertheless, with miR-148a-3p overexpression, the osteogenesis was impaired, either in the ALP activity or the bone nodules formation stained by Alizarin Red solutions (**Fig.2d**). qPCR results were identical with that of staining, with lower *Alp* (P=0.0424), *Col1a1* (P=0.0016) and *Spp1* (P=0.0112) mRNA expression level just after transfection (without osteogenic induction), and remarkably decreased *Runx2* (P<0.0001), *Bglap* (P<0.0001) and *Spp1* (P<0.0001) mRNA expression level after seven days' transfection and induction (**Fig.2e**).

Potential targets of miR-148a-3p in regulating osteogenesis.

Proteomics of bones from 4-week-old WT and KO littermates were applied to study the potential mechanism of miR-148a-3p in regulating osteogenesis. By setting protein expression level ratio=1.2 (KO/WT≥1.2 or KO/WT≤0.833), 475 candidate proteins were targeted with a P value<0.05 (**Fig.3a**, **Supplementary Table 1**). Among them, 337 candidates were upregulated in KO mice, including ALPL (KO/WT=1.844, P=0.014) and SP7 (KO/WT = 2.326, P=0.022) (**Fig.3b**). Functional enrichment analysis of these differentially expressed proteins revealed that, compared with WT littermates, KO mice had a more powerful ossification process (**Fig.3c**), conforming to our previous *in vivo* and *in vitro* findings. With the help of miRDB and TargetScan database, nine candidate genes were predicted to be directly regulated by miR-148a-3p (**Fig.3d**, **Supplementary Table 2**). The protein expression levels of six candidates (CDK19, RAB34, ITGA11, PDIA3, SSR1, STT3A) were upregulated in condition of miR-148a-3p deficiency (**Fig.3e**), while the rest three (LIPA, MECP2, STXBP5) were downregulated (**Fig.S1c**).

Traditionally, miRNAs suppress the post-transcriptional expression of their target genes via two primary mechanisms, either through mRNA degradation or not[2]. Thus, in order to clarify the exact mechanism of how miR-148a regulate these candidates, mRNA-sequencing was accomplished to find out the gene mRNA expression differences between KO and WT littermates. When Q<0.05 and FC2 (KO/WT≥2 or ≤0.5) were set as filter parameters, 1259 genes were found in all (**Fig.3f**, **Supplementary Table 3**), among which 879 genes were upregulated. Once again, GO enrichment found that "Ossification" was greatly enhanced in KO mice (**Fig.3g**). The mRNA-seq results were then combined into the Venn diagram, only one candidate, *Itga11*, remained (**Fig.3h**, **Supplementary Table 4**), indicating miR-148a-3p can completely complement with Itga11 3'-UTR and modulate its protein expression level by mRNA cleavage. In comparison, the rest candidates were regulated via post-translational and translational repression.

miR-148a inhibit osteogenesis by targeting Itga11

Based on several online databases, Itga11 was suggested to play an essential role in osteogenesis. The *in situ* hybridization image of E14.5 mouse embryo (obtained from GenePaint, https://gp3.mpg.de/) demonstrated high expression level of Itga11 in Meckel's cartilage, ribs and spine during embryonic period (**Fig.4a**). The expression pattern of Itga11 (obtained from BioGPS, http://biogps.org/) revealed that Itga11 was highly expressed in osteoblasts, over 30 times more than in other tissues or cells (**Fig.4b**). In addition, the protein-protein interaction network (obtained from STRING, https://cn.string-db.org/) showed interaction between Itga11 and Type I Collagen (Col1a1 and Col1a2), a specific marker in osteogenesis (**Fig.4c**). What's more, Itga11 has already been reported as a regulator in bone homeostasis, as a receptor for either Collagen XIII[20] or osteolectin[21].

Next, immunostaining results verified the higher expression in KO littermates, in comparison to HE ones (**Fig.4d**). Dual-luciferase reporter assay verified the direct combination of miR-148a-3p and *Itga11* 3'-Untranslated Regions (3'-UTR). The luciferase activity was reduced by 23% when supplemented with miR-148a-3p and

Itga11-WT (*P*=0.0004), while almost no difference with miRNA NC or *Itga11*-MUT (**Fig.4e**).

miR-148a inhibit osteogenesis via PI3K/Akt/GSK3/β-catenin signaling pathway

Gene set enrichment analysis (GSEA) of mRNA-sequencing results indicated activation of some key metabolic pathways, including PI3K-Akt signaling pathway and Wnt signaling pathway (**Fig.4f**). Similar results were observed in proteomics, key proteins, such as AKT and β-catenin, were remarkably upregulated in miR-148a KO samples (**Fig.S1d**).

The molecular mechanism was further studied through western blot. With the stimulation of osteogenic media, the phosphorylation proportion of AKT (p-AKT/ pan-AKT) was obviously increased and reached climax at one hour at a level of about 0.096, then decayed with time. In the absence of miR-148a, the climax time point was not changed but peak value raised to 0.105. In the WT group, the phosphorylation proportion of GSK3β (p-GSK3β/pan-GSK3β), the inactive form of GSK3β, reached peak ratio of 0.801 at 2-hour stimulation. Then the values came down over time. While in the KO group, it remained at a relative high level till the eighth hour (0.907 at the second hour and 0.659 at the eighth hour). Meanwhile, obvious increasement can be observed in β-catenin protein expression level, a terminal protein of the classic WNT signaling pathway[22] (**Fig.4g**).

Discussion

In this article, we utilized miR-148a KO mice to study the role of miR-148a-3p in bone physiology. By taking advantage of miR-148a KO mice, we assessed the role of miR-148a in bone physiology *in vivo* for the first time. miR-148a KO mice manifested bone dysplasia with increased bone mass. Meanwhile, proteomics and RNA-sequencing of bone samples from littermates were performed to look for candidate targets in a natural *in vivo* environment. Itga11 was verified to be a direct target of miR-148a-3p in inhibiting osteogenesis via PI3K/Akt/GSK3/β-catenin

signaling pathway.

Multiple targets of miR-148a-3p that regulate bone physiology have already been reported. However, most of them were identified based on cell lines through *in vitro* experiments. Taking osteogenesis regulation for example, various target genes existed, like p300[7],Wnt5a[8], WNT1, TGFB2, IGF1[6], NRP1[9], Kdm6b[10], and SMURF1[11]. Due to the nature that the target genes regulated by the same miRNA may vary depending on cells or tissues[2], looking for targets based on *in vivo* environment might be a better choice.

Utilizing bone samples from four-week-old littermates, *Itga11* was found as the main effector of miR-148a in regulating osteogenesis. As reported, *Itga11*, a marker of mesenchymal stem cells[23], has an essential role during embryogenesis, especially in mesenchymal cells around the cartilage anlage in the developing skeleton[24,25], and takes part in the regulation of bone homeostasis[26,27]. Itga11 has also been reported to be involved in PI3K/Akt and Wnt signaling pathway, supporting our findings from another view. ITGA11 is one of the core genes that could influence PI3K/Akt signaling pathway once they were activated[28], its expression can affect tumor progression by modulating the PI3K/Akt pathway[29]. Besides, Wnt pathway can be activated by osteolectin/α11β1 signaling[27].

Nevertheless, we failed to unveil the mechanism of bone dysplasia, especially in aspect of shortened length. Osteoclastogenesis may get involved. Besides, according to recent publication, it might be related to abnormal endothelial proteolytic activities, which has been proved to have a vital role in bone elongation instead of mature osteoclasts[30]. ALDH18A1, which was defined as an important candidate in pathological angiogenesis through single-cell RNA sequencing[31], was significantly increased in KO mice (KO/WT=1.471, *P*=0.0119). Literatures have revealed the role of miR-148a in regulating angiogenesis[32-40] and osteoclastogenesis[41,42]. Further related researches should be carried out. What's more, crosstalk between genes and unconventional functions of miRNA[43] should be taken into consideration.

In summary, our findings showed that miR-148a-3p inhibited osteogenesis by targeting Itga11 via PI3K/Akt/GSK3/β-catenin pathway, which might partly be the

mechanism of how miR-148a deficiency caused bone dysplasia.

Acknowledgements:

This work was supported by National Natural Science Foundation of China (81741012); Shanghai Municipal Key Clinical Specialty-shslczdzk00901 (ZWJCB18). Thank PTM BIOLABS for their great help in proteomics analysis.

Conflict of Interest Statement:

The authors have no conflict of interest to declare.

Author contributions

XJ Chen, YC Pang and G Chai designed research; XJ Chen, RK Wei and WQ Han performed research; XJ Chen and YC Pang analyzed the data; XJ Chen drafted the paper; YC Pang and G Chai revised the paper.

References

1.   Cui Y, Lyu X, Ding L, et al. Global miRNA dosage control of embryonic germ layer specification. *Nature.* 2021.

2.   Bartel DP. Metazoan MicroRNAs. *Cell.* 2018;173(1):20-51.

3.   Slack FJ, Chinnaiyan AM. The Role of Non-coding RNAs in Oncology. *Cell.* 2019;179(5):1033-1055.

4.   van Herwijnen MJC, Driedonks TAP, Snoek BL, et al. Abundantly Present miRNAs in Milk-Derived Extracellular Vesicles Are Conserved Between Mammals. *Front Nutr.* 2018;5:81.

5.   Jones TL, Esa MS, Li KHC, et al. Osteoporosis, fracture, osteoarthritis & sarcopenia: A systematic review of circulating microRNA association. *Bone.* 2021;152:116068.

6.   Grieco GE, Cataldo D, Ceccarelli E, et al. Serum Levels of miR-148a and miR-21-5p Are Increased in Type 1 Diabetic Patients and Correlated with Markers of Bone Strength and Metabolism. *Noncoding RNA.* 2018;4(4).

7.   Liu N, Sun Y. microRNA-148a-3p-targeting p300 protects against osteoblast differentiation and osteoporotic bone reconstruction. *Regen Med.* 2021;16(5):435-449.

8.   Duan DY, Tang J, Tian HT, Shi YY, Jia J. Adipocyte-secreted microvesicle-derived miR-148a regulates adipogenic and osteogenic differentiation by targeting Wnt5a/Ror2 pathway. *Life Sci.* 2021;278:119548.

9.   Bao L, Zhang X, Xu Y, et al. Dysfunction of MiR-148a-NRP1 Functional Axis Suppresses Osteogenic Differentiation of Periodontal Ligament Stem Cells Under Inflammatory Microenvironment. *Cell Reprogram.* 2019;21(6):314-322.

10. Tian L, Zheng F, Li Z, et al. miR-148a-3p regulates adipocyte and osteoblast differentiation by targeting lysine-specific demethylase 6b. *Gene.* 2017;627:32-39.

11. Huang S, Li Y, Wu P, et al. microRNA-148a-3p in extracellular vesicles derived from bone marrow mesenchymal stem cells suppresses SMURF1 to

prevent osteonecrosis of femoral head. *J Cell Mol Med.* 2020;24(19):11512-11523.

12. Zhao M, Wang Y, Li G, et al. The role and potential mechanism of p75NTR in mineralization via in vivo p75NTR knockout mice and in vitro ectomesenchymal stem cells. *Cell Prolif.* 2020;53(2):e12758.

13. Chen Y, Wang X. miRDB: an online database for prediction of functional microRNA targets. *Nucleic Acids Res.* 2020;48(D1):D127-D131.

14. Agarwal V, Bell GW, Nam JW, Bartel DP. Predicting effective microRNA target sites in mammalian mRNAs. *Elife.* 2015;4.

15. Visel A, Thaller C, Eichele G. GenePaint.org: an atlas of gene expression patterns in the mouse embryo. *Nucleic Acids Res.* 2004;32(Database issue):D552-556.

16. Eichele G, Diez-Roux G. High-throughput analysis of gene expression on tissue sections by in situ hybridization. *Methods.* 2011;53(4):417-423.

17. Wu C, Jin X, Tsueng G, Afrasiabi C, Su AI. BioGPS: building your own mash-up of gene annotations and expression profiles. *Nucleic Acids Res.* 2016;44(D1):D313-316.

18. Szklarczyk D, Gable AL, Nastou KC, et al. The STRING database in 2021: customizable protein-protein networks, and functional characterization of user-uploaded gene/measurement sets. *Nucleic Acids Res.* 2021;49(D1):D605-D612.

19. Bardou P, Mariette J, Escudie F, Djemiel C, Klopp C. jvenn: an interactive Venn diagram viewer. *BMC Bioinformatics.* 2014;15:293.

20. Koivunen J, Tu H, Kemppainen A, et al. Integrin alpha11beta1 is a receptor for collagen XIII. *Cell Tissue Res.* 2021;383(3):1135-1153.

21. Zhang J, Cohen A, Shen B, et al. The effect of parathyroid hormone on osteogenesis is mediated partly by osteolectin. *Proc Natl Acad Sci U S A.* 2021;118(25).

22. Lerner UH, Ohlsson C. The WNT system: background and its role in bone. *J Intern Med.* 2015;277(6):630-649.

23.	Kaltz N, Ringe J, Holzwarth C, et al. Novel markers of mesenchymal stem cells defined by genome-wide gene expression analysis of stromal cells from different sources. *Exp Cell Res.* 2010;316(16):2609-2617.

24.	Tiger CF, Fougerousse F, Grundstrom G, Velling T, Gullberg D. alpha11beta1 integrin is a receptor for interstitial collagens involved in cell migration and collagen reorganization on mesenchymal nonmuscle cells. *Dev Biol.* 2001;237(1):116-129.

25.	Zhang WM, Popova SN, Bergman C, Velling T, Gullberg MK, Gullberg D. Analysis of the human integrin alpha11 gene (ITGA11) and its promoter. *Matrix Biol.* 2002;21(6):513-523.

26.	Koivunen J, Tu H, Kemppainen A, et al. Integrin alpha11beta1 is a receptor for collagen XIII. *Cell Tissue Res.* 2020.

27.	Shen B, Vardy K, Hughes P, et al. Integrin alpha11 is an Osteolectin receptor and is required for the maintenance of adult skeletal bone mass. *Elife.* 2019;8.

28.	Guo SM, Wang JX, Li J, et al. Identification of gene expression profiles and key genes in subchondral bone of osteoarthritis using weighted gene coexpression network analysis. *J Cell Biochem.* 2018;119(9):7687-7695.

29.	Zhang H, Zhang L, Lu M. Inhibition of integrin subunit alpha 11 restrains gastric cancer progression through phosphatidylinositol 3-kinase/Akt pathway. *Bioengineered.* 2021;12(2):11909-11921.

30.	Romeo SG, Alawi KM, Rodrigues J, Singh A, Kusumbe AP, Ramasamy SK. Endothelial proteolytic activity and interaction with non-resorbing osteoclasts mediate bone elongation. *Nat Cell Biol.* 2019;21(4):430-441.

31.	Rohlenova K, Goveia J, Garcia-Caballero M, et al. Single-Cell RNA Sequencing Maps Endothelial Metabolic Plasticity in Pathological Angiogenesis. *Cell Metab.* 2020;31(4):862-877 e814.

32.	Wang XX, Zhang H, Li Y. Preliminary study on the role of miR148a and DNMT1 in the pathogenesis of acute myeloid leukemia. *Mol Med Rep.* 2019;19(4):2943-2952.

33.	Xu Q, Jiang Y, Yin Y, et al. A regulatory circuit of miR-148a/152 and DNMT1

in modulating cell transformation and tumor angiogenesis through IGF-IR and IRS1. *J Mol Cell Biol.* 2013;5(1):3-13.

34. Yu J, Li Q, Xu Q, Liu L, Jiang B. MiR-148a inhibits angiogenesis by targeting ERBB3. *J Biomed Res.* 2011;25(3):170-177.

35. Kim H, Ko Y, Park H, et al. MicroRNA-148a/b-3p regulates angiogenesis by targeting neuropilin-1 in endothelial cells. *Exp Mol Med.* 2019;51(11):1-11.

36. Mu J, Zhu D, Shen Z, et al. The repressive effect of miR-148a on Wnt/beta-catenin signaling involved in Glabridin-induced anti-angiogenesis in human breast cancer cells. *BMC Cancer.* 2017;17(1):307.

37. Xu Q, Liu LZ, Yin Y, et al. Regulatory circuit of PKM2/NF-kappaB/miR-148a/152-modulated tumor angiogenesis and cancer progression. *Oncogene.* 2015;34(43):5482-5493.

38. Wang J, Yao Y, Wang K, Li J, Chu T, Shen H. MicroRNA-148a-3p alleviates high glucose-induced diabetic retinopathy by targeting TGFB2 and FGF2. *Acta Diabetol.* 2020;57(12):1435-1443.

39. Wong HA, Fatimy RE, Onodera C, et al. The Cancer Genome Atlas Analysis Predicts MicroRNA for Targeting Cancer Growth and Vascularization in Glioblastoma. *Mol Ther.* 2015;23(7):1234-1247.

40. Ge H, Shrestha A, Liu C, Wu P, Cheng B. MicroRNA 148a-3p promotes Thrombospondin-4 expression and enhances angiogenesis during tendinopathy development by inhibiting Kruppel-like factor 6. *Biochem Biophys Res Commun.* 2018;502(2):276-282.

41. Liu M, Sun Y, Zhang Q. Emerging Role of Extracellular Vesicles in Bone Remodeling. *J Dent Res.* 2018;97(8):859-868.

42. Tian G, Hu K, Qiu S, et al. Exosomes derived from PC-3 cells suppress osteoclast differentiation by downregulating miR-148a and blocking the PI3K/AKT/mTOR pathway. *Exp Ther Med.* 2021;22(5):1304.

43. Dragomir MP, Knutsen E, Calin GA. SnapShot: Unconventional miRNA Functions. *Cell.* 2018;174(4):1038-1038 e1031.

Figure Legends

Fig.1 miR-148a knock-out (KO) mice manifested bone dysplasia with enhanced osteogenic potential. (a) Representative gross images showing knock-out (KO) mice were grossly smaller than the heterozygotic (HE) ones. **(b)** Body weights of 12-week-old mice. The weight was lost along with the knock-out of the miR-148a allele (WT, n=6; HE, n=14; KO, n=8). One-way ANOVA tests, followed by Multiple Comparisons between each column and the WT column via Dunnett's t-test. Afterward, student t-tests were utilized between the HE and the KO group. **(c)** X-ray images (left) and length quantification (right) of mandibles from KO and HE mice (n=6 per group). Statistical analysis by student t-test. **(d)** Double fluorescent labeling of mandibles from KO and HE mice showed higher mineralization speed in KO ones. **(e)** µCT scans of tibias from knock-out (KO) and heterozygotic (HE) mice showed a remarkably higher bone mass in KO ones. Left: Representative three-dimensional reconstruction images. Right: Quantification analysis of bone parameters (n=6 per group). BV/TV, Bone Volume/Total Volume; Tb.N., Trabecular Number; Tb.Th., Trabecular Thickness; Tb.Sp., Trabecular Separation. **(f-g)** Representative immunostaining images of tibia sections (n=3 per group). **(f)** Runx2. **(g)** OCN. **(h)** Representative Osterix[+] immunostaining images of cranium sections.

Fig.2 miR-148a-3p inhibited osteogenesis *in-vivo*. (a) Alkaline phosphatase (ALP) staining of primary ectomesenchymal stem cells (EMSCs) extracted from wild-type (WT) or knock-out (KO) pups after 3-day osteogenic induction *in vitro*. **(b)** Validation of miR-148a-3p overexpression after miR-148a-3p mimic transfection in WT EMSCs. **(c)** Proliferation assay revealed that miR-148a-3p had no obvious influence on cell proliferation. **(d)** ALP (after 7-day osteogenic induction) and AR staining (after 21-day osteogenic induction) of WT EMSCs with or without miR-148a-3p mimic transfection. The osteogenic potential was impaired with miR-148a-3p overexpression. **(e)** mRNA expression level after seven days' osteogenic induction with or without miR-148a-3p overexpression. NC, Negative Control. All comparisons were

performed between the NC and the Mimic group via two-sided student t-tests.

Fig.3 Potential targets of miR-148a-3p in regulating osteogenesis. **(a)** Proteomics results of bones from 4-week-old WT and KO littermates (n=3 per group). 475 candidates were targeted by setting protein expression level ratio=1.2 and P value<0.05. **(b)** Osteogenesis-related protein expression levels (ALPL and SP7) were upregulated in KO mice. **(c)** Functional enrichment analysis of differentially expressed proteins revealed that, compared with WT littermates, KO mice had a more powerful ossification process (Red Arrow). **(d)** Venn diagram of proteomics, miRDB, and TargetScan results. Nine candidates that miR-148a-3p may directly regulate were found. **(e)** The protein expression levels of six candidates (CDK19, RAB34, ITGA11, PDIA3, SSR1, STT3A) were upregulated in condition of miR-148a-3p deficiency. **(f)** mRNA-sequencing results of bones from WT and KO littermates (n=3 per group) when Q<0.05 and FC2 were set as filter parameters. **(g)** GO enrichment found that "Ossification" was greatly enhanced in KO mice. **(h)** Venn diagram of proteomics, mRNA-sequencing, miRDB, and TargetScan results, indicating miR-148a-3p can modulate ITGA11 protein expression level by degrading its mRNA.

Fig.4 miR-148a inhibit osteogenesis by targeting *Itga11* via PI3K/Akt/GSK3/β-catenin signaling pathway. **(a)** The *in situ* hybridization image of Itga11 in E14.5 mouse embryo obtained from GenePaint. **(b)** Itga11 gene expression profile of Itga11 obtained from BioGPS, with a rather high expression in osteoblasts. **(c)** The protein-protein interaction network obtained from STRING showed interaction between Itga11 and Type I Collagen, a specific marker in osteogenesis. **(d)** Immunostaining results verified the higher ITGA11 expression in KO littermates. **(e)** Dual-luciferase reporter assay confirmed that miR-148a-3p could bind to the 3'UTR of Itga11 and regulate its expression. **(f)** Gene set enrichment analysis (GSEA) of mRNA-sequencing results indicated activation of PI3K/Akt and Wnt signaling pathway. **(g)** PI3K/Akt/GSK3/β-catenin signaling pathway was promoted in condition of miR-148a knock out.

Supplementary Figure Legends

Fig.S1 (a) miRNA-sequencing of bone samples from littermates confirmed miR-148a knock out in KO mice. **(b)** Bone defect animal models indicated an accelerated bone healing process in KO mice in comparison to WT ones. Left: Representative middle-cross section of μCT scans. Right: Quantification of the bone defect site. (n=6 per group). **(c)** The protein expression levels of three candidates (LIPA, MECP2, STXBP5) were downregulated in condition of miR-148a-3p deficiency. **(d)** Protein expression pattern in PI3K/Akt and Wnt signaling pathway obtained from proteomics. Key proteins, such as AKT and β-catenin, were remarkably upregulated in miR-148a KO samples. Red box: Upregulated proteins; Green box: Downregulated proteins.

Supplementary Table

Supplementary Table 1. Proteomics results of bones from 4-week-old knock-out (KO) and wild-type (WT) littermates.

Supplementary Table 2. Raw data of Venn diagram composed of proteomics, miRDB and TargetScan results.

Supplementary Table 3. mRNA-sequencing results of bones from 4-week-old knock-out (KO) and wild-type (WT) littermates.

Supplementary Table 4. Raw data of Venn diagram composed of proteomics, mRNA-sequencing, miRDB and TargetScan results.

Fig.1

Fig.2

Fig.3

Fig.4

Fig.S1