Synaptopodin-associated ribosomal hubs regulate lesion-induced homeostatic

plasticity at hippocampal mossy fiber synapses

Pia Kruse[1], Andreas Vlachos[1,2,3,§], Maximilian Lenz[1,§]

[1]Department of Neuroanatomy, Institute of Anatomy and Cell Biology, Faculty of Medicine, University of Freiburg,

Freiburg, Germany.

[2]Center for Basics in Neuromodulation (NeuroModulBasics), Faculty of Medicine, University of Freiburg, Freiburg,

Germany.

[3]Center BrainLinks-BrainTools, University of Freiburg, Freiburg, Germany.

[§]Correspondence to andreas.vlachos@anat.uni-freiburg.de and maximilian.lenz@anat.uni-freiburg.de,

Department of Neuroanatomy, Institute of Anatomy and Cell Biology, Faculty of Medicine,

University of Freiburg, 79104 Freiburg, Germany

Keywords: homeostatic plasticity, denervation, synaptopodin, local protein synthesis

ABSTRACT

Neurological diseases can lead to the denervation of brain regions caused by demyelination, traumatic injury or cell death. Nevertheless, the molecular and structural mechanisms underlying lesion-induced reorganization of denervated brain regions are a matter of ongoing investigation. In order to address this issue, we performed an entorhinal cortex lesion (ECL) in mouse organotypic entorhino-hippocampal tissue cultures of both sexes and studied denervation-induced homeostatic plasticity of mossy fiber synapses, which connect dentate granule cells with CA3 pyramidal neurons and play important roles in spatial learning. Partial denervation caused a homeostatic strengthening of excitatory neurotransmission in dentate granule cells (GC), in CA3 pyramidal neurons, and their direct synaptic connections as revealed by paired recordings (GC-to-CA3). These functional changes were accompanied by ultrastructural reorganization of mossy fiber synapses, which regularly contain the plasticity-related protein synaptopodin and the spine apparatus organelle. We demonstrate that the spine apparatus organelle and its associated protein synaptopodin assemble ribosomes in close proximity to synaptic sites and unravel a synaptopodin-related transcriptome, which can be linked to the expression of homeostatic synaptic plasticity. Notably, synaptopodin-deficient tissue preparations that lack the spine apparatus organelle, failed to express homeostatic adjustments. Hence, synaptopodin and the spine apparatus organelle form local protein synthesis hubs that are essential for mediating lesion-induced homeostatic synaptic plasticity.

INTRODUCTION

Denervation is a common feature of various neurological diseases, which are accompanied by neuronal cell death, demyelination or traumatic injury (Mesulam et al., 2019). As a result of the loss-of-input in otherwise healthy brain regions, complex reorganizational processes of neural circuits have been described. Among them, homeostatic synaptic plasticity has been established as an adaptive process of synaptic features that aims at compensating for perturbations of network activity (Turrigiano, 2012; Vitureira et al., 2012; Turrigiano, 2017). Specifically, compensatory excitatory synaptic strengthening upon denervation or activity deprivation has been reported (Turrigiano et al., 1998). Although homeostatic synaptic plasticity has been recognized as a local mechanism that can occur at distinct dendritic regions, the mechanisms that mediate dendritic autonomy in maintaining homeostasis remain widely unknown.

The hippocampal network is a crucial structure in memory acquisition. Here, the excitatory connection between dentate granule cells and CA3 pyramidal cells, i.e., the hippocampal mossy fiber synapse, has been demonstrated to play a crucial role in memory formation, maintenance and retrieval (Weng et al., 2018). Although numerous studies have addressed the unique plasticity features of this synapse (e.g. the onset of synaptic homeostasis and detonating transmission; (Lee et al., 2013)) and its impact on memory and behavior, no studies have addressed whether lesion-induced homeostatic synaptic plasticity occurs at this synapse when the main cortical excitatory input is affected. In this context, organotypic tissue cultures of the entorhino-hippocampal complex provide the opportunity to study synaptic transmission at hippocampal mossy fiber synapses in steady-state condition (Maus et al., 2020). Due to the laminated cortico-hippocampal input, the *in vitro* entorhinal cortex lesion (ECL) has been established as a model for the partial denervation of hippocampal neural circuits (Prang et al., 2003; Vlachos et al., 2012a; Vlachos et al., 2012c; Lenz et al., 2019).

In this study we used an entorhinal cortex lesion (ECL) of organotypic entorhino-hippocampal tissue cultures to study homeostatic adaptations in the dentate gyrus and CA3 network upon partial denervation. We provide evidence that partial denervation leads to excitatory synaptic strengthening in both dentate granule cells and CA3 pyramidal neurons. Using paired whole-cell patch-clamp recordings, we demonstrate changes in the synaptic transmission at hippocampal mossy fiber synapses. These changes are accompanied by an ultrastructural reorganization of the spine apparatus organelle, which has previously been linked to synaptic plasticity (Deller et al., 2003; Jedlicka et al., 2009; Vlachos et al., 2009; Vlachos et al., 2013; Korkotian et al., 2014; Jedlicka and Deller, 2017; Maggio and Vlachos, 2018; Lenz et al., 2021b; Lenz et al., 2021a). We demonstrate that synaptopodin and the spine apparatus organelle form ribosomal hubs that are linked to homeostatic plasticity-related local protein synthesis (Sutton et al., 2006). Consequently, synaptopodin-deficient tissue cultures, which lack the spine apparatus organelle, showed deficits in the expression of homeostatic changes in excitatory neurotransmission and regional trancriptomes upon ECL. We therefore conclude, that the spine

70 apparatus organelle forms plastic local protein synthesis hubs that contribute to homeostatic synaptic

71 adjustments upon partial denervation.

MATERIALS AND METHODS

Ethics statement

Mice were maintained in a 12 h light/dark cycle with food and water available ad libitum. Every effort was made to minimize distress and pain of animals. All experimental procedures were performed according to German animal welfare legislation and approved by the appropriate animal welfare committee and the animal welfare officer of Albert-Ludwigs-University Freiburg, Faculty of Medicine (X-17/07K, X-17/09K, X-21/01B).

Preparation of tissue cultures

Entorhino-hippocampal tissue cultures were prepared at postnatal day 3-5 from C57BL/6J (wildtype, $Synpo^{+/+}$) and B6.129-Synpo^{tm1Mndl}/Dllr [$Synpo^{-/-}$, (Deller et al., 2003; Lenz et al., 2021b; Lenz et al., 2021a)] mice of either sex as previously described (Del Turco and Deller, 2007). Cultivation medium contained 50% (v/v) MEM, 25% (v/v) basal medium eagle, 25% (v/v) heat-inactivated normal horse serum, 25 mM HEPES buffer solution, 0.15% (w/v) bicarbonate, 0.65% (w/v) glucose, 0.1 mg/ml streptomycin, 100 U/ml penicillin, and 2 mM glutamax. The pH was adjusted to 7.3. All tissue cultures were allowed to mature for at least 18 days in a humidified atmosphere with 5% CO_2 at 35°C. Cultivation medium was replaced 3 times per week.

Entorhinal cortex lesion (ECL)

The entorhinal cortex of mature slice cultures (≥18 days *in vitro*) was transected with a sterile scalpel from the rhinal to the hippocampal fissure to achieve a complete lesion of the perforant path. Except for the lesion-induced partial denervation of the hippocampus, cytoarchitecture of both the hippocampus and the entorhinal cortex remained unchanged.

Whole-cell patch-clamp recordings

Whole-cell patch-clamp recordings were carried out at 35°C (3-6 neurons per culture). The bath solution contained (in mM) 126 NaCl, 2.5 KCl, 26 $NaHCO_3$, 1.25 NaH_2PO_4, 2 $CaCl_2$, 2 $MgCl_2$, and 10 glucose (aCSF) and was oxygenated continuously (5% CO_2 / 95% O_2). Patch pipettes contained (in mM) 126 K-gluconate, 10 HEPES, 4 KCl, 4 ATP-Mg, 0.3 GTP-Na$_2$, 10 PO-Creatine, 0.3% (w/v) biocytin (pH 7.25 with KOH, 290 mOsm with sucrose), having a tip resistance of 4-6 MΩ. Pyramidal neurons were visually identified using a LN-Scope (Luigs & Neumann) equipped with an infrared dot-contrast and a 40x water-immersion objective (NA 0.8; Olympus). Spontaneous excitatory postsynaptic currents (sEPSCs) of both dentate granule cells and CA3 pyramidal neurons were recorded in voltage-clamp mode at a holding potential of -70 mV. Series resistance was monitored before and after each recording and recordings were discarded if the series resistance reached ≥ 30 MΩ. For recording of intrinsic cellular properties in current-clamp mode, pipette capacitance of 2.0 pF was corrected and series resistance was compensated using the automated bridge balance tool of the MultiClamp commander. I-

106 V-curves were generated by injecting 1 s square pulse currents starting at −100 pA and increasing in 10
107 pA steps until +500 pA current injection was reached (sweep duration: 2 s).

Paired whole-cell patch-clamp recordings of hippocampal mossy fiber synapses

Action potentials were generated in the presynaptic dentate granule cell by 5 ms square current pulses
(1 nA). For connectivity analysis, 5 action potentials were applied at 20 Hz respectively (inter-sweep-
interval: 10 sec; at least 30 repetitions), while recording unitary excitatory postsynaptic currents
(uEPSCs) from CA3 pyramidal cells. Neurons were considered to be connected if > 5 % of action
potentials evoked time-locked inward uEPSCs (Vlachos et al., 2012b). Recordings were performed in
the presence of (-)-bicuculline-methiodide (10 μM, #ab120108 Abcam) to prevent inhibitory synapse
recruitment. Moreover, glutamine (200 μM, #11539876 Gibco) was added to the recording medium to
avoid synapse depletion (Tani et al., 2014).

Post-hoc staining and imaging

Tissue cultures were fixed overnight in a solution of 4% (w/v) paraformaldehyde (PFA; in PBS (0.1M,
pH 7.4) with 4% (w/v) sucrose). After fixation, cultures were washed in PBS (0.1 M, pH 7.4) and
consecutively incubated for 1 h with 10% (v/v) normal goat serum (NGS) in 0.5% (v/v) Triton X-100
containing PBS to reduce nonspecific staining. For post-hoc visualization of patched granule cells and
CA3 pyramidal cells, Streptavidin Alexa Fluor 488 (Streptavidin A488, 1:1000; #S32354 Invitrogen)
was added to the incubation solution. DAPI nuclear stain (1:5000 in PBS for 10 minutes; #62248
Thermo Scientific) was used to visualize cytoarchitecture. Cultures were washed, transferred onto glass
slides and mounted for visualization with DAKO anti-fading mounting medium (#S302380-2, Agilent).

Transmission electron microscopy

Wildtype (Synpo$^{+/+}$) and Synpo$^{-/-}$ tissue cultures were fixed in 4 % paraformaldehyde (w/v) and 2 %
glutaraldehyde (w/v) in 0.1 M phosphate buffer (PB) overnight and washed for 1 hour in 0.1 M PB.
After fixation, tissue cultures were sliced with a vibratome and the slices were incubated with 1 %
osmium tetroxide for 20 min in 5 % (w/v) sucrose containing 0.1 M PB. The slices were washed 5 times
for 10 min in 0.1 M PB and washed in graded ethanol [10 min in 10 % (v/v) and 10 min in 20 % (v/v)].
The slices were then incubated with uranyl acetate [1 % (w/v) in 70 % (v/v) ethanol)] overnight and
were subsequently dehydrated in graded ethanol 80 % (v/v), 90 % (v/v) and 98 % (v/v) for 10 min.
Finally, slices were incubated with 100 % (v/v) ethanol twice for 15 min followed by two 15 min washes
with propylene oxide. The slices were then transferred for 30 min in a 1:1 mixture of propylene oxide
with durcupan and then for 1 hour in durcupan. The durcupan was exchanged with fresh durcupan and
the slices were transferred to 4 °C overnight. The slices were then embedded between liquid release-
coated slides and coverslips. Cultures were re-embedded and ultrathin sections were collected on copper
grids. Electron microscopy was performed at a LEO 906E microscope (Zeiss) at 4646x magnification.

Regional mRNA library preparations and transcriptome analysis

RNA library preparations for transcriptome analysis were performed using the NEBNext® Single Cell/Low Input RNA Library Prep Kit for Illumina® (New England Biolabs, #E6420) according to the manufacturer's instructions. Briefly, isolation of the dentate gyrus and CA3 area from individual tissue cultures was performed using a scalpel. One isolated dentate gyrus or CA3 area respectively was transferred to 7.5 µl lysis buffer (supplemented with murine RNase inhibitor) and homogenized using a pestill. Samples were centrifuged for 30 seconds at 10,000g and 5 µl of supernatant were collected from individual samples and further processed. After cDNA synthesis, cDNA amplification was performed according to the manufacturer's protocol. cDNA yield was consecutively analyzed by a High Sensitivity DNA assay on a Bioanalyzer instrument (Agilent). cDNA amount was adjusted to 10 ng for further downstream applications. After fragmentation and adaptor ligation, dual index primers (New England Biolabs, #E7600S) were ligated in a library amplification step using 10 PCR cycles. Libraries were finally cleaned up with 0.8X SPRI Beads following a standard bead purification protocol. Library purity and size distribution was assessed with a High Sensitivity DNA assay on a Bioanalyzer instrument (Agilent). We quantified the libraries using the NEBNext Library Quant Kit for Illumina (New England Biolabs, #E7630) based on the mean insert size provided by the Bioanalyzer. A 10 nM sequencing pool (120 µl in Tris-HCl, pH 8.5) was generated for sequencing on the NovaSeq6000 sequencing platform (Illumina; service provided by CeGaT GmbH, Tübingen, Germany). We performed a paired-end sequencing with 101 bp read length. Data analysis was performed at the Galaxy platform (usegalaxy.eu). All files contained more than 10 M high-quality reads (after mapping to the reference genome; mm10) having at least a phred quality of 30 (>90% of total reads).

Synaptopodin co-Immunoprecipitation (SP-coIP) and RT-qPCR analysis

Six tissue cultures per biological replicate were washed in coIP-solution and dounce homogenized on ice in a solution containing (150 µl per sample, coIP-solution, in mM): 10 HEPES, 120 NaCl, 0.05% (v/v) NP-40, cOmplete protease inhibitor (Roche, according to manufacturers' instructions), murine RNase inhibitor (1:1000, New England Biolabs, #M0314) and cycloheximide (0.5 mg/ml, Sigma Aldrich, #C-4859). Input was precleared twice with protein A beads (10 µl per 100 µl sample/supernatant) to reduce unspecific binding. Protein A beads (10 µl, New England Biolabs, #S1425) were washed twice in coIP-solution and consecutively incubated for 1 hour at 4°C with an anti-Synaptopodin antibody (rabbit anti-Synaptopodin, Synaptic Systems, #163002). Beads were carefully washed twice to remove unbound antibodies and incubated for 1 hour with the precleared supernatant ('input') at 4°C with head-over-tail rotation. 10 µl of the precleared supernatant were kept as input control. After incubation, beads were washed twice with coIP-solution and finally dissolved in RNA protection buffer (New England Biolabs). RNA was released from the beads through Protein K lysis (Monarch® Total RNA Miniprep Kit; #T2010S New England Biolabs). After lysis, beads were removed

from the solution in a magnetic rack and the RNA containing supernatant was transferred to fresh tubes.

RNA isolation on coIP- and input-samples was then performed according to the manufacturer's

instructions. RNA content was determined using the Agilent RNA 6000 Pico Kit (#5067-1513 Agilent)

with a 2100 Bioanalyzer (#G2939BA Agilent). For RT-qPCR analysis, purified RNA was consecutively

reverse transcribed (RevertAid RT Kit; #K1691 Thermo Scientific). cDNA was diluted in water to a

final concentration of 0.2 ng/µl. RT-qPCR was performed using a C1000 Touch Thermal Cycler (BIO-

RAD) and the CFX 384 Real-Time PCR system (BIO-RAD). 0.9 ng target cDNA diluted in TaqMan

Gene Expression Master Mix (Applied Biosystems #4369016) were amplified using standard TaqMan

Gene Expression Assays (Applied Biosystems; Assay-IDs: Gapdh: Mm99999915_g1; Gria1:

Mm00433753_m1). RNA library preparations for Synpo-related transcriptome analysis were performed

using the NEBNext® Single Cell/Low Input RNA Library Prep Kit for Illumina® (New England Biolabs,

#E6420) according to the manufacturer's instructions. We quantified the libraries using the NEBNext

Library Quant Kit for Illumina (New England Biolabs, #E7630) based on the mean insert size provided

by the Bioanalyzer. A 10 nM sequencing pool (120 µl in Tris-HCl, pH 8.5) was generated for sequencing

on the NovaSeq6000 Sequencing platform (Illumina; service provided by CeGaT GmbH, Tübingen,

Germany). We performed a paired-end sequencing with 151 bp read length. Data analysis was

performed at the Galaxy platform (usegalaxy.eu). All files contained more than 10 M high-quality reads

(after mapping to the reference genome; mm10) having at least a phred quality of 30 (>90% of total

reads).

Quantification and statistics

Electrophysiological data were analyzed using pClamp 10.7 (Axon Instruments) and MiniAnalysis

(Synaptosoft) software. The fraction of action potentials followed by time-locked excitatory

postsynaptic current responses was considered as synaptic response rate. uEPSC amplitude was assessed

in uEPSCs from successfully transmitted action potentials as well as the mean amplitude of all

successfully evoked postsynaptic responses. In order to ensure analysis of time-locked unitary

responses, events were included if > 5% of presynaptic pulses were successfully transmitted for each

pulse respectively. For individual pulses in pulse trains that caused ≤ 5% successfully transmitted

events, response rate was set to '0'. EPSC properties were analyzed using the automated template search

tool for event detection. In order to perform a hierarchical analysis of sEPSCs, all events of a recorded

cell were sorted based on their peak amplitude. Events above the 50th percentile therefore correspond to

all events above the median amplitude of events. Input resistance was calculated for the injection of –

100 pA current at a time frame of 200 ms with a maximum distance to the initial hyperpolarization.

Resting membrane potential was calculated as the mean baseline value. AP detection was performed

using the input-output curve threshold search event detection tool, and the AP frequency was assessed

by the number of APs detected during the respective current injection time (1s).

RT-qPCR data were analyzed as previously described (Lenz et al., 2020) with the Bio-Rad CFX Maestro 1.0 software package using the ΔΔCq method with Gapdh as reference gene. Values were normalized to the mean value of the respective control group.

Electron microscopy images were saved as TIF-files and analyzed using the ImageSP Viewer software (http://e.informer.com/sys-prog.com). Ultrastructural features were analyzed by an investigator blind to experimental conditions in randomly selected mossy fiber terminals from electron micrographs of the stratum lucidum in non-lesioned and lesioned wildtype tissue cultures.

RNA sequencing data were uploaded to the Galaxy web platform (public server: usegalaxy.eu; (Goecks et al., 2010; Afgan et al., 2018; Galaxy, 2022)) and transcriptome analysis was performed using the Galaxy platform in accordance with the reference-based RNA-seq data analysis tutorial (Batut et al., 2021). Adapter sequencing, low quality, and short reads were removed via the CUTADAPT tool (Galaxy version 3.5+galaxy0). Reads were mapped using RNA STAR (Galaxy version 2.7.8a+galaxy0) with the mm10 full reference genome (Mus musculus). The evidence-based annotation of the mouse genome (GRCm38), version M25 (Ensembl 100), served as gene model (GENCODE). For an initial assessment of gene expression, unstranded FEATURECOUNT (Galaxy version 2.0.1+galaxy2) analysis was performed from RNA STAR output. Only samples that contained >60% uniquely mapping reads (feature: "exon") were considered for further analysis. Statistical evaluation was performed using DESeq2 (Galaxy version 2.11.40.7+ galaxy1) with "pathway integrity" (ECL experiments) or "coIP" (synaptopodin-coIP experiments) as the primary factor that might affect gene expression. Genes with a low number of mean reads (< 300 BaseMean) were excluded from further analysis. Genes were considered as differentially expressed or enriched if the adjusted p-value was < 0.05. Data visualization was performed according to a modified version of a previously published workflow (Reimand et al., 2019). A ranked list for the gene set enrichment analysis was generated based on Wald-Stats from the DeSeq2 output. For the consecutive gene set enrichment analysis, we have used the GSEA software package ((Mootha et al., 2003; Subramanian et al., 2005); BROAD institute; version GSEA_4.2.1) with integrated MSigDB gene sets. Enrichment maps were built using cytoscape ((Shannon et al., 2003); version 3.9.0) with the EnrichmentMap application. Further functional enrichment analyses were performed using g:Profiler (version e107_eg54_p17_bf42210) with g:SCS multiple testing correction method applying significance threshold of 0.05 (Raudvere et al., 2019). Gene sets with 100-500 terms were considered for illustration. Heatmaps were generated based on z-scores of the normalized count table.

Data were statistically analyzed using GraphPad Prism 7 (GraphPad software, USA). All values represent mean ± standard error of the mean. For statistical comparison of two experimental groups, a Mann-Whitney test was employed. EPSC amplitude/frequency plots and AP-frequency plots were statistically assessed by the repeated measure (RM) two-way ANOVA test with Sidak's (two groups) multiple comparisons test. uEPSC amplitude values from individual cells were stacked in subcolumns and the pulse number was defining tabular rows (COLUMN factor: pathway integrity, genetic

background; ROW factor: EPSC amplitude bin or current injection). P-values < 0.05 were considered

statistically significant (*p < 0.05, **p < 0.01, ***p < 0.001); results that did not yield significant

differences are designated 'ns'. Statistical differences in XY-plots were indicated in the legend of the

figure panels (*) when detected through multiple comparisons, irrespective of their localization and the

level of significance. In the text and figures, values represent the mean ± standard error of the mean

(s.e.m.).

Data availability

Sequencing data have been deposited in the Gene Expression Omnibus (GEO) repository (accession

number: GSE216509). Original data are available from the authors upon reasonable request.

Digital illustrations

Confocal image stacks were stored as TIF files. Figures were prepared using the ImageJ software

package (https://imagej.nih.gov/ij/) and Photoshop graphics software (Adobe, San Jose, CA, USA).

Image brightness and contrast were adjusted.

RESULTS

Entorhinal cortex lesion leads to homeostatic synaptic plasticity in both dentate granule cells and CA3 pyramidal neurons

Entorhinal cortex lesion was performed in mature tissue cultures to achieve a complete dissection of the perforant path leading to a partial denervation of both dentate granule cells and CA3 pyramidal neurons (Figure 1A). Spontaneous excitatory postsynaptic currents (sEPSCs) were recorded from dentate granule cells and CA3 pyramidal neurons in non-lesioned control and lesioned tissue cultures (3 days post entorhinal cortex lesion, 3dp ECL) to assess denervation-induced effects on excitatory neurotransmission (Figure 1B). As previously described [e.g., (Vlachos et al., 2012c; Lenz et al., 2019)], partial denervation caused a strengthening of excitatory synapses in dentate granule cells, i.e., a significant increase in both the mean sEPSC amplitude and half width as well as the mean sEPSC area (Figure 1C). Percentile analysis revealed that the lesion-induced increase in sEPSC amplitudes is evident when comparing the 50% biggest events in both groups while no effect was observed when comparing smaller events (Figure 1D). Notably, a significant upregulation in the mean sEPSC frequency was observed after lesion (Figure 1E) that resulted from an increased abundance of high amplitude sEPSCs as revealed by the amplitude/frequency-plot (Figure 1F).

Next, we examined excitatory neurotransmission in CA3 pyramidal neurons in non-lesioned and lesioned cultures (3dp ECL; Figure 1G-J). Again, a significant increase in both the mean sEPSC amplitude and area was observed in CA3 cells from lesioned cultures indicating a denervation-induced excitatory synaptic strengthening (Figure 1G). Notably, the mean sEPSC half width remained unchanged. In the sEPSC amplitude percentile analysis, excitatory synaptic strengthening seems to be restricted to the high amplitude events (Figure 1H). Moreover, the mean sEPSC frequency remained unchanged upon ECL (Figure 1I), which resulted from an increase in the abundance of high amplitude events (in the sEPSC amplitude range from 80-110 pA, RM two-way ANOVA followed by Sidak's multiple comparisons test) at the expense of small amplitude events (Figure 1J). We therefore conclude that ECL leads to a homeostatic increase in excitatory neurotransmission in both dentate granule cells and CA3 pyramidal neurons.

Partial denervation does not cause changes in intrinsic cellular properties

Passive and active membrane properties were assessed in both dentate granule cells (Figure 2A-C) and CA3 pyramidal neurons (Figure 2D-F) to test for homeostatic adaptations of intrinsic cellular properties upon lesion. In dentate granule cells no changes in passive membrane properties (Figure 2A), i.e., the resting membrane potential (RMP) and the input resistance as well as the input-output curve dynamics were observed (Figure 2B). Additionally, the action potential frequency of dentate granule cells was similar in lesioned and non-lesioned cultures (Figure 2C). Noteworthy, RMP assessment confirmed the mature state of the recorded dentate granule cells [c.f. (Strehl et al., 2018)].

295 Concerning CA3 pyramidal neurons, neither the passive membrane properties (Figure 2D) – including

296 input-output curve dynamics (Figure 2E) – nor the active properties (Figure 2F) were changed. Based

297 on these results, we conclude that intrinsic cellular properties were stable following denervation in both

298 tested cell types of the hippocampal network.

Entorhinal cortex lesion induces changes in synaptic transmission at hippocampal mossy fiber

synapses

Hippocampal mossy fiber synapses have previously been linked to learning processes and memory

formation (Lipp et al., 1988; Vandael et al., 2020). Concerning their synaptic features, they have been

described as synapses that generate large postsynaptic currents upon activation (Henze et al., 2002b).

Given that our results demonstrate changes in both investigated regions, which additionally hint towards

a preferential lesion-related plasticity induction in readily strong synapses at CA3 pyramidal neurons,

we next assessed the features of synaptic transmission at hippocampal mossy fiber synapses.

Therefore, we performed paired whole-cell patch-clamp recordings of individual dentate granule cells

and CA3 pyramidal neurons in organotypic tissue cultures (Figure 3A). To test for synaptic connections,

5 consecutive action potentials at 20 Hz frequency were elicited in dentate granule cells (Figure 3B).

Unitary excitatory postsynaptic currents (uEPSCs) were monitored in CA3 pyramidal neurons in non-

lesioned and lesioned tissue cultures. We found that a train of presynaptic action potentials in connected

neurons induced a robust facilitation of postsynaptic responses in both postsynaptic response rate and

uEPSC amplitude under baseline conditions (Figure 3B-D). Notably, the response rate to the first

presynaptic action potential, which is the number of presynaptic action potentials that generate a

postsynaptic response, was low in non-lesioned cultures (~3%; Figure 3C). In lesioned cultures,

however, a significant increase in the synaptic reliability for the first two pulses was observed. For the

last three pulses in one train of stimulation, averaged synaptic response rate levels became similar

(Figure 3C). Moreover, the uEPSC amplitude was significantly increased in response to the individual

pulses 3 days after ECL (Figure 3D). This novel approach of GC-to-CA3 paired recordings in the steady-

state environment of organotypic tissue cultures provided evidence that synaptic transmission at

hippocampal mossy fiber synapses is a target for lesion-induced remodeling of hippocampal networks.

Synaptopodin and the spine apparatus organelle form dendritic ribosomal hubs that are

associated with excitatory synapse related mRNAs

Hippocampal mossy fiber synapses show a complex organization of their pre- and postsynaptic sites.

Among these features, a regular occurrence of the spine apparatus organelle, which is a transmission-

and plasticity-regulating organelle, has been reported at the postsynaptic site (Chicurel and Harris,

1992). Using transmission electron microscopy, a close interaction between ribosomes and the spine

apparatus organelle was readily revealed (Figure 4A). We hypothesized that the spine apparatus

organelle at its marker protein synaptopodin (Deller et al., 2003) might regulate lesion-induced plasticity

at hippocampal mossy fiber synapses by regulating local protein synthesis. To test whether
synaptopodin interacts with local translation machineries, i.e., dendritic ribosomes, we performed
synaptopodin co-immunoprecipitation (Synpo-coIP) experiments with subsequent RT-qPCR analysis.
Using RNA assessment on a Bioanalyzer, we found rRNA peaks in Synpo-coIP experiments whereas
no rRNA was detected in bead-only controls (Figure 4B). RT-qPCR analysis (reference gene: Gapdh)
from input and Synpo-coIP samples revealed an enrichment of Gria1 mRNA (Figure 4C), which has
been linked to synaptopodin-dependent synaptic plasticity (Vlachos et al., 2009) and homeostatic
synaptic adjustments (Sutton et al., 2006; Aoto et al., 2008). We therefore conclude, that synaptopodin
and the spine apparatus organelle might serve as a ribosomal hub that significantly contributes to
plasticity-related local protein synthesis.

To further unravel the plasticity-mediating features of synaptopodin and the spine apparatus organelle,
we performed a transcriptome analysis of Synpo-coIP enriched mRNA (Figure 4D, extended data table
4-1). A considerable amount of mRNAs was enriched after Synpo-coIP compared to input samples (UP;
2355 mRNAs; extended data table 4-2). Subsequently, a gene set enrichment analysis (GSEA) was
performed to elucidate the functional relevance of synaptopodin-associated mRNAs (Figure 4E).
Among others, an enrichment in synapse and actin cytoskeleton-related gene sets was identified. Of
note, these data suggest the synaptopodin-dependent assembly of ribosomal hubs that might be
dedicated to local protein synthesis at synapses.

Ultrastructural reorganization of the spine apparatus organelle
Previous studies have demonstrated that the ultrastructural organization of the spine apparatus organelle
might represent a target for plasticity (Vlachos et al., 2013; Lenz et al., 2021a). To study lesion-induced
effects on the ultrastructural organization of the spine apparatus organelle that might interfere with local
protein synthesis, we performed transmission electron microscopy of mossy fiber synapses in non-
lesioned and lesioned (3dp ECL) entorhino-hippocampal tissue cultures (Figure 4F; c.f., (Vlachos et al.,
2013)). No differences in the structural organization of pre- and postsynaptic sites could be detected
between the groups (presynapse: mossy fiber perimeter: 11.61 ± 0.4 μm (control; n = 120) vs. 10.88 ±
0.35 μm (3dp ECL; n = 90); postsynapse: number of PSDs: 4.91 ± 0.33 (control; n = 120) vs. 4.6 ± 0.27
(3dp ECL; n = 90), length of PSDs: 201 nm ± 4.35 (control; n = 589) vs. 201.7 ± 5.42 nm (3dp ECL; n
= 414); values indicate mean ± sem, Mann-Whitney test). Nevertheless, a substantial reorganization of
the spine apparatus organelle was evident in lesioned cultures 3 days after injury compared to non-
lesioned controls (Figure 4G, H). We found an increased number of dense plates in spine apparatus
organelles from lesioned cultures (Figure 4G). At the same time the length of individual dense plates
was significantly reduced in lesioned cultures (Figure 4H). Therefore, we conclude that the spine
apparatus organelle represents a target for lesion-induced changes at hippocampal mossy fiber synapses.

The synaptopodin-associated transcriptome remains stable upon lesion

Given the ultrastructural reorganization of the spine apparatus organelle, we next tested whether the

composition of the synaptopodin-associated transcriptome changed in lesioned networks 3dp ECL.

Therefore, a differential mRNA enrichment analysis was performed in Synpo-coIP and their respective

input RNA samples from lesioned and non-lesioned wildtype cultures (Figure 5). In mRNA libraries

prepared from input RNA samples, we found a considerable amount of differentially expressed genes

upon ECL (extended data table 5-1). In contrast, a limited amount of mRNAs were differentially

enriched upon ECL in Synpo-coIP samples (extended data table 5-2). We therefore conclude that the

composition of the synaptopodin-associated transcriptome remains widely stable upon partial

denervation.

Synaptopodin-deficient tissue cultures show deficits in homeostatic synaptic plasticity upon

partial denervation

Based on our ultrastructural analysis, we hypothesized that the reorganization of the spine apparatus

organelle might contribute to homeostatic synaptic plasticity at hippocampal mossy fiber synapses. To

gain further insights concerning the role of the spine apparatus organelle, we performed experiments in

synaptopodin-deficient tissue preparations, which lack the spine apparatus organelle (Figure 6; (Deller

et al., 2003)).

Synaptopodin-deficient dentate granule cells showed a significant increase in both the mean sEPSC

amplitude and area as well as the mean sEPSC half width 3 days following ECL (Figure 6A). In contrast,

no changes in the mean sEPSC frequency, which have been observed in wildtype (Synpo$^{+/+}$) cultures,

were detected (Figure 6B). Moreover, passive membrane properties remained stable upon lesion in

synaptopodin-deficient tissue preparations (Figure 6C).

When assessing excitatory neurotransmission onto CA3 pyramidal neurons in lesioned and non-lesioned

cultures, no significant changes in the sEPSC event analysis (Figure 6D) or the mean sEPSC frequency

(Figure 6E) were observed. Again, passive membrane properties remained unchanged (Figure 6F).

Thus, we here provide evidence for deficits in homeostatic synaptic adaptations for both dentate granule

cells and CA3 pyramidal neurons in synaptopodin-deficient tissue cultures.

We extended these findings with GC-to-CA3 paired whole-cell patch-clamp recordings in

synaptopodin-deficient preparations (Figure 6G). In contrast to wildtype cultures, no significant

differences in the synaptic reliability could be detected 3 days after ECL (Figure 6H). When analyzing

the uEPSC amplitude of successfully transmitted events, surprisingly, we found a significant reduction

in the pulse-related uEPSC amplitude in lesioned synaptopodin-deficient preparations (Figure 6I).

Notably, excitatory neurotransmission seemed facilitated under baseline conditions as compared to

wildtype cultures (Figure 7): In addition to significant changes in sEPSC half width, area and

amplitude/frequency-plots (Figure 7 A, B), changes in uEPSCs at mossy fiber synapses were evident in

paired recordings (Figure 7 C, D). Hence, synaptopodin-deficient mossy fiber synapses responded in a

synaptic depression rather than homeostatic synaptic strengthening after the lesion. These findings

demonstrate the role of the postsynaptic spine apparatus organelle on plasticity of mossy fiber synapses.

They are consistent with recent work that suggested a role of synaptopodin and the spine apparatus in

mediating metaplasticity (Hulme et al., 2014; Lenz et al., 2021b).

404   Entorhinal cortex lesion causes synaptopodin-dependent changes in the region-specific

405   hippocampal transcriptome

To further unravel synaptopodin-dependent changes in hippocampal subfields upon lesion, we

performed transcriptome analyses of both the dentate gyrus and the CA3 region in non-lesion and

lesioned wildtype (Synpo$^{+/+}$) and synaptopodin-deficient (Synpo$^{-/-}$) cultures (Figure 8). The expression

analysis revealed a considerable number of genes that are differentially expressed in both hippocampal

regions from wildtype animals (Figure 8A, B). In contrast, synaptopodin-deficient tissue culture

preparations showed deficits in the lesion-induced adjustment of gene expression in the CA3 region

(Figure 8A). Moreover, wildtype and synaptopodin-deficient tissue preparations showed a genotype

dependent gene expression adjustment upon lesion in both regions although a common set of regulated

genes has been identified (Figure 8B; for detailed tabular results see extended data tables 8-1, 8-2, 8-3,

8-4). We therefore conclude, that synaptopodin might act as a sensor that triggers differential gene

expression towards homeostatic adjustments in a region-specific fashion.

Unbiased gene set enrichment analysis for all differentially expressed genes (gene ontology – molecular

function; GO:MF) from wildtype CA3 samples revealed that ECL caused changes in gene expression

related to ubiquitin- and GTP-related signaling (Figure 8C). Additionally, significant enrichments in

vesicle-, glia- and synapse-related gene sets were evident (gene ontology – cellular compartment;

GO:CC). In order to learn more about lesion-induced differential expression of synaptic gene sets in the

CA3 region ("synapse"; GOCC: 0045202), differentially expressed gene lists were filtered according to

the content of the respective gene ontology (Figure 8D). We found 153 differentially expressed synaptic

genes, which correspond to the strengthening of excitatory neurotransmission at mossy fiber synapses.

The same analyses were performed for wildtype DG samples (Figure 8E, F). In addition to changes in

ubiquitin- and GTP-related signaling genes, enrichments were seen for actin and transcription factor

binding (gene ontology – molecular function; Figure 8E). Moreover, endosome and mitochondria

related gene sets were enriched (gene ontology – cellular compartment; GO:CC). Notably, 160

differentially expressed synaptic genes were identified (Figure 8F).

Therefore, we conclude that the lesion-induced change in the region-specific transcriptome might

represent an essential component of homeostatic synaptic plasticity.

DISCUSSION

Denervation is a common phenomenon in numerous medical conditions affecting the central nervous system, which can trigger homeostatic adaptations in neural networks (Vlachos et al., 2012c; Hengen et al., 2013; Lambo and Turrigiano, 2013; Platschek et al., 2016; Barnes et al., 2017; Bissen et al., 2021). In this study, we have unraveled a novel feature of the spine apparatus organelle that crucially contributes to the localized expression of compensatory synaptic changes. In line with previous findings that employed proximity proteomics of synaptopodin (Falahati et al., 2022), we identified a close interaction between synaptopodin and ribosomes. Using synaptopodin co-immunoprecipitation experiments, we elucidated the synaptopodin-related transcriptome, which might contribute to plasticity-related local protein synthesis. Together with findings that indicate interactions between synaptopodin and ER-related proteins (Falahati et al., 2022), we suggest a role for synaptopodin and the spine apparatus organelle as local plasticity hubs near synaptic sites.

To study lesion-induced changes in synaptic transmission, we used the *in vitro* model of ECL in organotypic entorhino-hippocampal tissue cultures. After *in vitro* maturation, tissue organization reaches a steady-state that is characterized an *in vivo* like ultrastructural synaptic architecture (Haber et al., 2006; Heller and Rusakov, 2017). In this context, recent work has demonstrated that organotypic hippocampal tissue cultures have intact mossy fiber pathways and synapses (Maus et al., 2020). Moreover, the excitatory cortical input – the perforant path – terminates at distal apical dendrites of dentate granule cells and pyramidal neurons of the ammonic subregions (Frotscher and Heimrich, 1995; Frotscher et al., 1995; Frotscher et al., 1997). Our results provide evidence for homeostatic excitatory synaptic plasticity in both dentate granule cells and CA3 pyramidal neurons upon lesion, which is accompanied by changes in the region-specific transcriptome. Moreover, we here established a novel method to assess synaptic transmission at individual hippocampal mossy fiber synapses – the monosynaptic excitatory contacts between dentate granule cells and CA3 pyramidal neurons – in a steady-state environment by paired whole-cell patch-clamp recordings of connected neurons. In this experimental approach, we identified synaptic transmission at hippocampal mossy fiber synapses as a target for denervation induced synaptic plasticity. In addition, ultrastructural changes in the spine apparatus organelle, which is well established in the regulation of synaptic plasticity (Vlachos et al., 2013; Lenz et al., 2021a), can be observed after ECL. Since homeostatic adaptations on both synaptic and transcriptomic levels were altered in synaptopodin-deficient tissue cultures which lack the spine apparatus organelle, we conclude a crucial role for synaptopodin in mediating, sensing and triggering compensatory synaptic changes.

The dentate gyrus/CA3 neural network is an integral component of the memory-associated trisynaptic hippocampal pathway. The hippocampal mossy fiber pathway forms large *en passant* synapses and connects one individual granule cell to a limited number of CA3 pyramidal neurons (Acsady et al., 1998). Both regions receive cortical input – the perforant path – from entorhinal layer II stellate cells that can induce, promote and modulate information processing in hippocampal subregions. Moreover,

this pathway is a well-established target for various diseases; among them traumatic, inflammatory and

neurodegenerative conditions like Alzheimer's disease (AD; (Thal et al., 2000; Robinson et al., 2014)).

Here, we demonstrate that both dentate granule cells and CA3 pyramidal neurons express synaptic

plasticity upon partial denervation, which might interfere with information processing.

The expression of synaptic plasticity in the hippocampal neural circuit is involved in crucial aspects of

normal brain function, such as learning and memory. The hippocampal mossy fiber synapse has been

linked to distinct forms of memory acquisition and retrieval (Marr, 1971; Hunsaker et al., 2008).

Specifically, activity-dependent structural and functional changes in synaptic features have been found

to encode distinct input patterns for a prolonged period of time (Lee et al., 2013; Vandael et al., 2020).

Notably, alterations in hippocampal mossy fiber transmission have been reported in animal models of

Alzheimer's disease (AD) and aging related cognitive decline (Villanueva-Castillo et al., 2017; Viana

da Silva et al., 2019) where a particular loss in cortical input pathways onto hippocampal neurons can

be found. Nevertheless, it remains unclear if the lesion-induced synaptic response deteriorates cognitive

performance. Given that (1) homeostatic synaptic plasticity is altered in synaptopodin-deficient tissue

preparations and (2) alterations in synaptopodin expression have been linked to cognitive decline

(Wingo et al., 2019), we suggest that synaptopodin-mediated homeostasis aims at maintaining proper

network function.

Recently, characteristic features of transmission and stabilizing factors at hippocampal mossy fiber

synapses have been revealed (Vyleta and Jonas, 2014; Vyleta et al., 2016; De Bruyckere et al., 2018;

Vandael et al., 2021a). Upon presynaptic activity, a detonating behavior can be observed in which a

limited number of – but not single – action potentials can elicit a suprathreshold postsynaptic

depolarisation (Henze et al., 2002a). Although this phenomenon has been extensively described in

mossy fiber bouton-CA3 pyramidal cell paired recordings (Vandael et al., 2021b), we were not able to

reveal detonating characteristics in granule cell-CA3 pyramidal cell pairs. We assume that both

differences in the experimental model and methodology can account for these differences, which

warrant further clarification in future studies. Nevertheless, a filtering of single action potentials with a

subsequent robust facilitation of synaptic transmission at this synapse was evident, which is in line with

previous reports (Zalutsky and Nicoll, 1990; Gundlfinger et al., 2007). Our results demonstrate that the

partial loss of input following lesion can influence both synaptic reliability and potency, which has been

observed for other plasticity-inducing stimuli at this synapse (Zalutsky and Nicoll, 1990). We suggest

that lesion-induced modulation of filtering/facilitation properties might interact with network

phenomena, such as pattern separation and completion along the dentate gyrus/CA3 axis. The increase

in synaptic reliability and potency might shift the balance towards pattern completion rather than

separation (O'Reilly and McClelland, 1994; Leutgeb et al., 2007; Prince et al., 2016). Moreover, the

presence of synaptopodin will presumably interfere with this network process under baseline and

plasticity-inducing conditions accounting for the previously reported cognitive deficits in synaptopodin-

deficient mice (Deller et al., 2003) and the synaptopodin-related cognitive decline during human aging
(Wingo et al., 2019).

Of note, it is an established view that mossy fiber synaptic plasticity is predominantly mediated by
presynaptic processes (Vandael et al., 2020; Orlando et al., 2021; Monday et al., 2022). Nevertheless,
recent work has suggested that the postsynaptic site shapes the transmission and plasticity through
transsynaptic mechanisms (Vandael et al., 2021a). In this study, we provide evidence that the
postsynaptically located spine apparatus organelle mediates denervation-induced synaptic plasticity at
this synapse. Since its discovery, the function of the spine apparatus organelle and the mechanisms
through which it orchestrates synaptic plasticity remained enigmatic. We here demonstrate that
synaptopodin and the spine apparatus organelle represent dendritic ribosomal hubs that contribute to
plasticity-driven local protein synthesis. The lesion-induced ultrastructural remodeling of the spine
apparatus organelle might therefore represent a novel regulatory mechanism for localized changes in
synaptic transmission (Vlachos et al., 2009; Dubes et al., 2022).

Notably, we found opposing plasticity phenomena at hippocampal mossy fiber synapses on the level of
individual neuron pairs in wildtype and synaptopodin-deficient tissue preparations. Synaptopodin
deficiency did not only cause defects in plasticity expression but hampered the ability of neurons to
maintain synaptic stability upon lesion. While the loss of input in wildtype cultures led to a significant
homeostatic strengthening of DG-CA3 excitatory synaptic transmission, synaptopodin deficiency was
associated with a lesion-related loss of synaptic potency. We therefore conclude that the function of the
spine apparatus organelle and its related protein synaptopodin might not be restricted to the mediation
of receptor accumulation at postsynaptic sites but rather comprises the implementation of plasticity
directionality (Vlachos, 2012). However, the link between synaptopodin and recently described
mechanisms that switch plasticity directionality from homeostatic to associative rules remain to be
determined (Wu et al., 2022). On the other hand, the lack of synaptopodin might decrease synaptic
resilience due to the lack of stabilizing features (Yap et al., 2020), which is required to maintain synaptic
transmission upon lesion. Regardless of these considerations, once more we report that synaptopodin
and the spine apparatus organelle act as crucial regulators of synaptic plasticity.

Finally, synaptic transcriptional changes as seen in wildtype cultures were absent in synaptopodin-
deficient cultures. Due to these observations, we suggest a dual role for synaptopodin as activity sensor
and plasticity effector, which assembles synapse-related mRNAs at dendritic plasticity hubs.

ACKNOWLEDGEMENTS

The Galaxy server that was used for some calculations in this study is partially funded by Collaborative Research Centre 992 Medical Epigenetics (DFG grant SFB 992/1 2012) and German Federal Ministry of Education and Research (BMBF grants 031 A538A/A538C RBC, 031L0101B/031L0101C de.NBI-epi, and 031L0106 de.STAIR (de.NBI)). We thank Bernd Heimrich for helpful and stimulating discussions on ultrastructural mossy fiber architecture. Moreover, we thank Sigrun Nestel for excellent technical assistance in electron microscopy. This work was supported by Else Kröner-Fresenius-Stiftung (EKFS_#2019_A94 to ML) and Deutsche Forschungsgemeinschaft (DFG; Project-ID 259373024 B14–CRC/TRR 167 to AV).

REFERENCES

Acsady L, Kamondi A, Sik A, Freund T, Buzsaki G (1998) GABAergic cells are the major postsynaptic targets of mossy fibers in the rat hippocampus. J Neurosci 18:3386-3403.

Afgan E, Baker D, Batut B, van den Beek M, Bouvier D, Cech M, Chilton J, Clements D, Coraor N, Gruning BA, Guerler A, Hillman-Jackson J, Hiltemann S, Jalili V, Rasche H, Soranzo N, Goecks J, Taylor J, Nekrutenko A, Blankenberg D (2018) The Galaxy platform for accessible, reproducible and collaborative biomedical analyses: 2018 update. Nucleic Acids Res 46:W537-W544.

Aoto J, Nam CI, Poon MM, Ting P, Chen L (2008) Synaptic signaling by all-trans retinoic acid in homeostatic synaptic plasticity. Neuron 60:308-320.

Barnes SJ, Franzoni E, Jacobsen RI, Erdelyi F, Szabo G, Clopath C, Keller GB, Keck T (2017) Deprivation-Induced Homeostatic Spine Scaling In Vivo Is Localized to Dendritic Branches that Have Undergone Recent Spine Loss. Neuron 96:871-882 e875.

Batut B, van den Beek M, Doyle MA, Soranzo N (2021) RNA-Seq Data Analysis in Galaxy. Methods Mol Biol 2284:367-392.

Bissen D, Kracht MK, Foss F, Hofmann J, Acker-Palmer A (2021) EphrinB2 and GRIP1 stabilize mushroom spines during denervation-induced homeostatic plasticity. Cell Rep 34:108923.

Chicurel ME, Harris KM (1992) Three-dimensional analysis of the structure and composition of CA3 branched dendritic spines and their synaptic relationships with mossy fiber boutons in the rat hippocampus. J Comp Neurol 325:169-182.

De Bruyckere E, Simon R, Nestel S, Heimrich B, Katzel D, Egorov AV, Liu P, Jenkins NA, Copeland NG, Schwegler H, Draguhn A, Britsch S (2018) Stability and Function of Hippocampal Mossy Fiber Synapses Depend on Bcl11b/Ctip2. Front Mol Neurosci 11:103.

Del Turco D, Deller T (2007) Organotypic entorhino-hippocampal slice cultures--a tool to study the molecular and cellular regulation of axonal regeneration and collateral sprouting in vitro. Methods Mol Biol 399:55-66.

Deller T, Korte M, Chabanis S, Drakew A, Schwegler H, Stefani GG, Zuniga A, Schwarz K, Bonhoeffer T, Zeller R, Frotscher M, Mundel P (2003) Synaptopodin-deficient mice lack a spine apparatus and show deficits in synaptic plasticity. Proc Natl Acad Sci U S A 100:10494-10499.

Dubes S, Soula A, Benquet S, Tessier B, Poujol C, Favereaux A, Thoumine O, Letellier M (2022) miR-124-dependent tagging of synapses by synaptopodin enables input-specific homeostatic plasticity. EMBO J:e109012.

Falahati H, Wu Y, Feuerer V, Simon HG, De Camilli P (2022) Proximity proteomics of synaptopodin provides insight into the molecular composition of the spine apparatus of dendritic spines. Proc Natl Acad Sci U S A 119:e2203750119.

Frotscher M, Heimrich B (1995) Lamina-specific synaptic connections of hippocampal neurons in vitro. J Neurobiol 26:350-359.

Frotscher M, Heimrich B, Deller T (1997) Sprouting in the hippocampus is layer-specific. Trends Neurosci 20:218-223.

Frotscher M, Heimrich B, Deller T, Nitsch R (1995) Understanding the cortex through the hippocampus: lamina-specific connections of the rat hippocampal neurons. J Anat 187 (Pt 3):539-545.

Galaxy C (2022) The Galaxy platform for accessible, reproducible and collaborative biomedical analyses: 2022 update. Nucleic Acids Res.

Goecks J, Nekrutenko A, Taylor J, Galaxy T (2010) Galaxy: a comprehensive approach for supporting accessible, reproducible, and transparent computational research in the life sciences. Genome Biol 11:R86.

Gundlfinger A, Leibold C, Gebert K, Moisel M, Schmitz D, Kempter R (2007) Differential modulation of short-term synaptic dynamics by long-term potentiation at mouse hippocampal mossy fibre synapses. J Physiol 585:853-865.

Haber M, Zhou L, Murai KK (2006) Cooperative astrocyte and dendritic spine dynamics at hippocampal excitatory synapses. J Neurosci 26:8881-8891.

Heller JP, Rusakov DA (2017) The Nanoworld of the Tripartite Synapse: Insights from Super-Resolution Microscopy. Front Cell Neurosci 11:374.

Hengen KB, Lambo ME, Van Hooser SD, Katz DB, Turrigiano GG (2013) Firing rate homeostasis in visual cortex of freely behaving rodents. Neuron 80:335-342.

Henze DA, Wittner L, Buzsaki G (2002a) Single granule cells reliably discharge targets in the hippocampal CA3
network in vivo. Nat Neurosci 5:790-795.

Henze DA, McMahon DB, Harris KM, Barrionuevo G (2002b) Giant miniature EPSCs at the hippocampal mossy
fiber to CA3 pyramidal cell synapse are monoquantal. J Neurophysiol 87:15-29.

Hulme SR, Jones OD, Raymond CR, Sah P, Abraham WC (2014) Mechanisms of heterosynaptic metaplasticity.
Philos Trans R Soc Lond B Biol Sci 369:20130148.

Hunsaker MR, Rosenberg JS, Kesner RP (2008) The role of the dentate gyrus, CA3a,b, and CA3c for detecting
spatial and environmental novelty. Hippocampus 18:1064-1073.

Jedlicka P, Deller T (2017) Understanding the role of synaptopodin and the spine apparatus in Hebbian synaptic
plasticity - New perspectives and the need for computational modeling. Neurobiol Learn Mem 138:21-
30.

Jedlicka P, Schwarzacher SW, Winkels R, Kienzler F, Frotscher M, Bramham CR, Schultz C, Bas Orth C, Deller
T (2009) Impairment of in vivo theta-burst long-term potentiation and network excitability in the dentate
gyrus of synaptopodin-deficient mice lacking the spine apparatus and the cisternal organelle.
Hippocampus 19:130-140.

Korkotian E, Frotscher M, Segal M (2014) Synaptopodin regulates spine plasticity: mediation by calcium stores.
J Neurosci 34:11641-11651.

Lambo ME, Turrigiano GG (2013) Synaptic and intrinsic homeostatic mechanisms cooperate to increase L2/3
pyramidal neuron excitability during a late phase of critical period plasticity. J Neurosci 33:8810-8819.

Lee KJ, Queenan BN, Rozeboom AM, Bellmore R, Lim ST, Vicini S, Pak DT (2013) Mossy fiber-CA3 synapses
mediate homeostatic plasticity in mature hippocampal neurons. Neuron 77:99-114.

Lenz M, Galanis C, Kleidonas D, Fellenz M, Deller T, Vlachos A (2019) Denervated mouse dentate granule cells
adjust their excitatory but not inhibitory synapses following in vitro entorhinal cortex lesion. Exp Neurol
312:1-9.

Lenz M, Kruse P, Eichler A, Straehle J, Beck J, Deller T, Vlachos A (2021a) All-trans retinoic acid induces
synaptic plasticity in human cortical neurons. Elife 10.

Lenz M, Eichler A, Kruse P, Muellerleile J, Deller T, Jedlicka P, Vlachos A (2021b) All-trans retinoic acid induces
synaptopodin-dependent metaplasticity in mouse dentate granule cells. Elife 10.

Lenz M, Eichler A, Kruse P, Strehl A, Rodriguez-Rozada S, Goren I, Yogev N, Frank S, Waisman A, Deller T,
Jung S, Maggio N, Vlachos A (2020) Interleukin 10 Restores Lipopolysaccharide-Induced Alterations in
Synaptic Plasticity Probed by Repetitive Magnetic Stimulation. Front Immunol 11:614509.

Leutgeb JK, Leutgeb S, Moser MB, Moser EI (2007) Pattern separation in the dentate gyrus and CA3 of the
hippocampus. Science 315:961-966.

Lipp HP, Schwegler H, Heimrich B, Driscoll P (1988) Infrapyramidal mossy fibers and two-way avoidance
learning: developmental modification of hippocampal circuitry and adult behavior of rats and mice. J
Neurosci 8:1905-1921.

Maggio N, Vlachos A (2018) Tumor necrosis factor (TNF) modulates synaptic plasticity in a concentration-
dependent manner through intracellular calcium stores. J Mol Med (Berl) 96:1039-1047.

Marr D (1971) Simple memory: a theory for archicortex. Philos Trans R Soc Lond B Biol Sci 262:23-81.

Maus L, Lee C, Altas B, Sertel SM, Weyand K, Rizzoli SO, Rhee J, Brose N, Imig C, Cooper BH (2020)
Ultrastructural Correlates of Presynaptic Functional Heterogeneity in Hippocampal Synapses. Cell Rep
30:3632-3643 e3638.

Mesulam MM, Lalehzari N, Rahmani F, Ohm D, Shahidehpour R, Kim G, Gefen T, Weintraub S, Bigio E, Geula
C (2019) Cortical cholinergic denervation in primary progressive aphasia with Alzheimer pathology.
Neurology 92:e1580-e1588.

Monday HR, Kharod SC, Yoon YJ, Singer RH, Castillo PE (2022) Presynaptic FMRP and local protein synthesis
support structural and functional plasticity of glutamatergic axon terminals. Neuron 110:2588-2606
e2586.

Mootha VK et al. (2003) PGC-1alpha-responsive genes involved in oxidative phosphorylation are coordinately
downregulated in human diabetes. Nat Genet 34:267-273.

O'Reilly RC, McClelland JL (1994) Hippocampal conjunctive encoding, storage, and recall: avoiding a trade-off.
Hippocampus 4:661-682.

Orlando M, Dvorzhak A, Bruentgens F, Maglione M, Rost BR, Sigrist SJ, Breustedt J, Schmitz D (2021)
Recruitment of release sites underlies chemical presynaptic potentiation at hippocampal mossy fiber
boutons. PLoS Biol 19:e3001149.

Platschek S, Cuntz H, Vuksic M, Deller T, Jedlicka P (2016) A general homeostatic principle following lesion
induced dendritic remodeling. Acta Neuropathol Commun 4:19.

Prang P, Del Turco D, Deller T (2003) Associational sprouting in the mouse fascia dentata after entorhinal lesion
in vitro. Brain Res 978:205-212.

Prince LY, Bacon TJ, Tigaret CM, Mellor JR (2016) Neuromodulation of the Feedforward Dentate Gyrus-CA3
Microcircuit. Front Synaptic Neurosci 8:32.

Raudvere U, Kolberg L, Kuzmin I, Arak T, Adler P, Peterson H, Vilo J (2019) g:Profiler: a web server for
functional enrichment analysis and conversions of gene lists (2019 update). Nucleic Acids Res 47:W191-
W198.

Reimand J, Isserlin R, Voisin V, Kucera M, Tannus-Lopes C, Rostamianfar A, Wadi L, Meyer M, Wong J, Xu C,
Merico D, Bader GD (2019) Pathway enrichment analysis and visualization of omics data using
g:Profiler, GSEA, Cytoscape and EnrichmentMap. Nat Protoc 14:482-517.

Robinson JL, Molina-Porcel L, Corrada MM, Raible K, Lee EB, Lee VM, Kawas CH, Trojanowski JQ (2014)
Perforant path synaptic loss correlates with cognitive impairment and Alzheimer's disease in the oldest-
old. Brain 137:2578-2587.

Shannon P, Markiel A, Ozier O, Baliga NS, Wang JT, Ramage D, Amin N, Schwikowski B, Ideker T (2003)
Cytoscape: a software environment for integrated models of biomolecular interaction networks. Genome
Res 13:2498-2504.

Strehl A, Galanis C, Radic T, Schwarzacher SW, Deller T, Vlachos A (2018) Dopamine Modulates Homeostatic
Excitatory Synaptic Plasticity of Immature Dentate Granule Cells in Entorhino-Hippocampal Slice
Cultures. Front Mol Neurosci 11:303.

Subramanian A, Tamayo P, Mootha VK, Mukherjee S, Ebert BL, Gillette MA, Paulovich A, Pomeroy SL, Golub
TR, Lander ES, Mesirov JP (2005) Gene set enrichment analysis: a knowledge-based approach for
interpreting genome-wide expression profiles. Proc Natl Acad Sci U S A 102:15545-15550.

Sutton MA, Ito HT, Cressy P, Kempf C, Woo JC, Schuman EM (2006) Miniature neurotransmission stabilizes
synaptic function via tonic suppression of local dendritic protein synthesis. Cell 125:785-799.

Tani H, Dulla CG, Farzampour Z, Taylor-Weiner A, Huguenard JR, Reimer RJ (2014) A local glutamate-
glutamine cycle sustains synaptic excitatory transmitter release. Neuron 81:888-900.

Thal DR, Holzer M, Rub U, Waldmann G, Gunzel S, Zedlick D, Schober R (2000) Alzheimer-related tau-
pathology in the perforant path target zone and in the hippocampal stratum oriens and radiatum correlates
with onset and degree of dementia. Exp Neurol 163:98-110.

Turrigiano G (2012) Homeostatic synaptic plasticity: local and global mechanisms for stabilizing neuronal
function. Cold Spring Harb Perspect Biol 4:a005736.

Turrigiano GG (2017) The dialectic of Hebb and homeostasis. Philos Trans R Soc Lond B Biol Sci 372.

Turrigiano GG, Leslie KR, Desai NS, Rutherford LC, Nelson SB (1998) Activity-dependent scaling of quantal
amplitude in neocortical neurons. Nature 391:892-896.

Vandael D, Okamoto Y, Jonas P (2021a) Transsynaptic modulation of presynaptic short-term plasticity in
hippocampal mossy fiber synapses. Nat Commun 12:2912.

Vandael D, Borges-Merjane C, Zhang X, Jonas P (2020) Short-Term Plasticity at Hippocampal Mossy Fiber
Synapses Is Induced by Natural Activity Patterns and Associated with Vesicle Pool Engram Formation.
Neuron 107:509-521 e507.

Vandael D, Okamoto Y, Borges-Merjane C, Vargas-Barroso V, Suter BA, Jonas P (2021b) Subcellular patch-
clamp techniques for single-bouton stimulation and simultaneous pre- and postsynaptic recording at
cortical synapses. Nat Protoc 16:2947-2967.

Viana da Silva S, Zhang P, Haberl MG, Labrousse V, Grosjean N, Blanchet C, Frick A, Mulle C (2019)
Hippocampal Mossy Fibers Synapses in CA3 Pyramidal Cells Are Altered at an Early Stage in a Mouse
Model of Alzheimer's Disease. J Neurosci 39:4193-4205.

Villanueva-Castillo C, Tecuatl C, Herrera-Lopez G, Galvan EJ (2017) Aging-related impairments of hippocampal
mossy fibers synapses on CA3 pyramidal cells. Neurobiol Aging 49:119-137.

Vitureira N, Letellier M, Goda Y (2012) Homeostatic synaptic plasticity: from single synapses to neural circuits.
Curr Opin Neurobiol 22:516-521.

Vlachos A (2012) Synaptopodin and the spine apparatus organelle-regulators of different forms of synaptic
plasticity? Ann Anat 194:317-320.

Vlachos A, Bas Orth C, Schneider G, Deller T (2012a) Time-lapse imaging of granule cells in mouse entorhino-
hippocampal slice cultures reveals changes in spine stability after entorhinal denervation. J Comp Neurol
520:1891-1902.

Vlachos A, Korkotian E, Schonfeld E, Copanaki E, Deller T, Segal M (2009) Synaptopodin regulates plasticity of
dendritic spines in hippocampal neurons. J Neurosci 29:1017-1033.

Vlachos A, Muller-Dahlhaus F, Rosskopp J, Lenz M, Ziemann U, Deller T (2012b) Repetitive magnetic
stimulation induces functional and structural plasticity of excitatory postsynapses in mouse organotypic
hippocampal slice cultures. J Neurosci 32:17514-17523.

Vlachos A, Becker D, Jedlicka P, Winkels R, Roeper J, Deller T (2012c) Entorhinal denervation induces
homeostatic synaptic scaling of excitatory postsynapses of dentate granule cells in mouse organotypic
slice cultures. PLoS One 7:e32883.

Vlachos A, Ikenberg B, Lenz M, Becker D, Reifenberg K, Bas-Orth C, Deller T (2013) Synaptopodin regulates
denervation-induced homeostatic synaptic plasticity. Proc Natl Acad Sci U S A 110:8242-8247.

Vyleta NP, Jonas P (2014) Loose coupling between Ca2+ channels and release sensors at a plastic hippocampal
synapse. Science 343:665-670.

Vyleta NP, Borges-Merjane C, Jonas P (2016) Plasticity-dependent, full detonation at hippocampal mossy fiber-
CA3 pyramidal neuron synapses. Elife 5.

Weng FJ, Garcia RI, Lutzu S, Alvina K, Zhang Y, Dushko M, Ku T, Zemoura K, Rich D, Garcia-Dominguez D,
Hung M, Yelhekar TD, Sorensen AT, Xu W, Chung K, Castillo PE, Lin Y (2018) Npas4 Is a Critical
Regulator of Learning-Induced Plasticity at Mossy Fiber-CA3 Synapses during Contextual Memory
Formation. Neuron 97:1137-1152 e1135.

Wingo AP, Dammer EB, Breen MS, Logsdon BA, Duong DM, Troncosco JC, Thambisetty M, Beach TG, Serrano
GE, Reiman EM, Caselli RJ, Lah JJ, Seyfried NT, Levey AI, Wingo TS (2019) Large-scale proteomic
analysis of human brain identifies proteins associated with cognitive trajectory in advanced age. Nat
Commun 10:1619.

Wu CH, Tatavarty V, Jean Beltran PM, Guerrero AA, Keshishian H, Krug K, MacMullan MA, Li L, Carr SA,
Cottrell JR, Turrigiano GG (2022) A bidirectional switch in the Shank3 phosphorylation state biases
synapses toward up- or downscaling. Elife 11.

Yap K, Drakew A, Smilovic D, Rietsche M, Paul MH, Vuksic M, Del Turco D, Deller T (2020) The actin-
modulating protein synaptopodin mediates long-term survival of dendritic spines. Elife 9.

Zalutsky RA, Nicoll RA (1990) Comparison of two forms of long-term potentiation in single hippocampal
neurons. Science 248:1619-1624.

FIGURES

Figure 1: Entorhinal cortex lesion caused homeostatic synaptic plasticity in both dentate granule

cells and CA3 pyramidal cells.

(A) Entorhinal cortex lesion (ECL) in organotypic entorhino-hippocampal tissue cultures was

performed by dissecting the perforant path (white line, scale bar 300 μm). Cytoarchitecture was

visualized by DAPI nuclear staining.

(B) Spontaneous excitatory postsynaptic currents (sEPSCs) were recorded in both dentate granule cells

and CA3 pyramidal cells in non-lesioned and lesioned cultures (3 days post lesion, 3 dp ECL).

(C-D) Spontaneous EPSC analysis in dentate granule cells revealed a lesion-induced excitatory synaptic

strengthening in the mean sEPSC amplitude, half width and area (C; $n_{control} = 22$ cells; $n_{3dp\ ECL} = 22$ cells;

Mann-Whitney test; Linear regression fit for XY-plot analysis). Percentile analysis of sEPSC

amplitudes (D) demonstrated that the excitatory synaptic strengthening was evident in the 50% largest

events (Mann-Whitney test).

(E-F) The frequency of sEPSCs was significantly increased 3 days after ECL (E; Mann-Whitney test).

Moreover, amplitude/frequency-analysis shows that the abundance of larger sEPSC amplitudes was

higher in lesioned cultures when compared to non-lesioned controls (F; RM-two-way ANOVA followed

by Sidak's multiple comparisons test).

(G-H) Spontaneous EPSC analysis in CA3 pyramidal cells revealed a lesion-induced increase in mean

sEPSC amplitude, whereas half width and area remained unchanged (G; $n_{control} = 48$ cells; $n_{3dp\ ECL} = 49$

cells; Mann-Whitney test; Linear regression fit for XY-plot analysis). Percentile analysis of sEPSC

amplitudes (H) demonstrated that the excitatory synaptic strengthening was only visible in the 50%

largest events (Mann-Whitney test).

(I-J) No lesion-induced changes in mean sEPSC frequency were detectable in CA3 pyramidal neurons.

Amplitude/frequency-analysis showed a reduced number of small events in CA3 pyramidal cells 3 days

after ECL (Mann-Whitney test for column statistics and RM-two-way ANOVA followed by Sidak's

multiple comparisons test for XY-plot analysis).

Individual data points are indicated by colored dots. Values represent mean ± s.e.m (*$p < 0.05$, **$p <$

0.01, ***$p < 0.001$, ns - non significant difference).

Figure 2: Entorhinal cortex lesion did not influence active or passive membrane properties.

(A-B) Both resting membrane potential and input resistance (A) as well as the input-output curve for the first 15 sweeps of current injection (B) in dentate granule cells were unaffected by ECL ($n_{control}$ = 22 cells; $n_{3dp\ ECL}$ = 22 cells; Mann-Whitney test and RM two-way ANOVA followed by Sidak's multiple comparisons test for XY-plot analysis).

(C) Action potential frequency upon increasing current injection in dentate granule cells was similar in non-lesioned and lesioned cultures (RM two-way ANOVA followed by Sidak's multiple comparisons test).

(D-E) Both resting membrane potential and input resistance (D) as well as the input-output curve for the first 15 sweeps of current injection (E) in CA3 pyramidal neurons were unaffected by the ECL ($n_{control}$ = 48 cells; $n_{3dp\ ECL}$ = 49 cells; Mann-Whitney test and RM two-way ANOVA followed by Sidak's multiple comparisons test for XY-plot analysis).

(F) Action potential frequency upon increasing current injection in CA3 pyramidal neurons is similar in non-lesioned and lesioned cultures (RM two-way ANOVA followed by Sidak's multiple comparisons test).

Individual data points are indicated by colored dots. Values represent mean ± s.e.m (ns - non significant difference).

Figure 3: Lesion-induced loss of the cortico-hippocampal projection changed transmission at hippocampal mossy fiber synapses.

(A) Post-hoc visualization of patched neurons in organotypic tissue cultures after paired-electrophysiological recordings between individual presynaptic (pre) dentate granule cells and postsynaptic (post) CA3 pyramidal neurons to analyze synaptic transmission features at hippocampal mossy fiber synapses (scale bar 300 μm).

(B) Five consecutive action potentials were induced in dentate granule cells at 20 Hz (bottom trace) to assess postsynaptic responses in CA3 pyramidal neurons in lesioned and non-lesioned cultures (upper traces).

(C) The response rate, i.e., the fraction of presynaptic pulses that elicited a postsynaptic response in a connected pair of individual neurons, was significantly increased for the first two pulses in lesioned cultures when compared to non-lesioned controls ($n_{control} = 10$; $n_{3dp\ ECL} = 10$; Mann-Whitney test for the respective pulse numbers).

(D) In successfully elicited unitary EPSCs (uEPSCs), a significant increase in uEPSC amplitude was detectable ($n°1_{control} = 20$ events, $n°1_{ECL} = 187$ events; $n°2_{control} = 115$ events, $n°2_{ECL} = 304$ events; $n°3_{control} = 292$ events, $n°3_{ECL} = 394$ events (one data point outside the axis limits); $n°4_{control} = 326$ events, $n°4_{ECL} = 385$ events; $n°5_{control} = 307$ events, $n°5_{ECL} = 384$ events; Mann-Whitney test for the respective pulse numbers).

Individual data points are indicated by colored dots. Values represent mean ± s.e.m (*$p < 0.05$, **$p < 0.01$, ***$p < 0.001$, ns - non significant difference).

Figure 4: The spine apparatus organelle assembles ribosomes and excitatory synapse related mRNAs.

(A) Transmission electron microscopy (TEM) images suggested a close association of ribosomes (arrowheads) and the spine apparatus organelle. Scale bar = 300 nm.

(B) Bioanalyzer traces (picoRNA Chip) from RNA samples prepared from precleared input (left) and Synpo-coIPs (right) demonstrating the co-immunoprecipitation of ribosomal RNA with anti-synaptopodin antibody. Notably, beads-only controls did not show an immunoprecipitation of rRNA.

(C) RT-qPCR analysis demonstrated a significant enrichment of Gria1 mRNA in synaptopodin-coIP compared to input samples indicating an association between the spine apparatus organelle and synaptic plasticity related mRNAs (normalized to Gapdh; $n_{input/SP\text{-}coIP}$ = 10 samples, Mann-Whitney test).

(D) Analysis of differential mRNA abundance in Synpo-coIP RNA samples compared to their respective input samples to reveal the synaptopodin related transcriptome (extended data table 4-1). A log2(FC) value > 0 indicates an mRNA enrichment in Synpo-coIP samples (UP: 2355 genes; extended data table 4-2). Differentially enriched genes ($p_{adj.}$ < 0.05) are indicated by orange dots (n = 5 samples input and Synpo-coIP respectively; 5-6 tissue cultures pooled from one filter per sample; DESeq2-analysis, usegalaxy.eu platform).

(E) A gene set enrichment analysis for a cellular compartment gene ontology was performed for significantly enriched genes in Synpo-coIP samples and visualized by a cytoscape map (node cutoff p = 0.05, edge cutoff (similarity): 0.8; node size reflects gene set size; node positions have been manually adjusted; original data set provided in extended data figure 4-3).

(F) Transmission electron microscopy was used to visualize ultrastructural properties of mossy fiber synapses in lesioned and non-lesioned control cultures (scale bar 1 μm).

(G) Upon lesion, the abundance of spine apparatus organelles (SA) and their number of dense plates were significantly increased in the postsynaptic compartment of mossy fiber synapses ($n_{control}$ = 120 (no SA = 97 postsynapses, 1 dense plate = 22 postsynapses, 2 dense plates = 1 postsynapse, 3 dense plates = 0 postsynapses); n_{ECL} = 90 (no SA = 56 postsynapses, 1 dense plate = 21 postsynapses, 2 dense plates = 12 postsynapses, 3 dense plates = 1 postsynapses)).

(H) While the number of dense plates increased, their length significantly decreased 3 days after ECL ($n_{control}$ = 24, one data point outside the axis limits; n_{ECL} = 48; Mann-Whitney test).

Individual data points are indicated by colored dots. Values represent mean ± s.e.m (**p < 0.01, ns - non significant difference).

Figure 5: Entorhinal cortex lesion did not promote major changes in synaptopodin-associated

mRNA.

Volcano plots to visualize the differential abundance of mRNAs in lesioned and non-lesioned cultures.

Both input and Synpo-coIP samples have been analyzed. Differentially expressed/enriched genes ($p_{adj.}$

< 0.05) are indicated by orange dots (n = 5 samples for both lesioned and non-lesioned cultures; 5-6

tissue cultures pooled from one filter per sample; DESeq2-analysis, usegalaxy.eu platform; extended

data tables 5-1 and 5-2).

Figure 6: Lesion-induced homeostatic synaptic plasticity at hippocampal mossy fiber synapses was impaired in synaptopodin-deficient tissue cultures.

(A-B) sEPSCs were recorded from dentate granule cells in synaptopodin-deficient tissue cultures (Synpo$^{-/-}$). Here, the mean sEPSC amplitude, half width and area significantly increased 3 days after the ECL (A; $n_{control}$ = 29; $n_{3dp\ ECL}$ = 26; Mann-Whitney test; Linear regression fit for XY-plot analysis). In contrast to wildtype cultures, no changes in the mean sEPSC frequency or the sEPSC amplitude/frequency-distribution were observed (B; Mann-Whitney test for column statistics and RM-two-way ANOVA followed by Sidak's multiple comparisons test for XY-plot analysis).

(C) No changes in passive membrane properties of dentate granule cells were detected in synaptopodin-deficient cultures upon lesion (Mann-Whitney test for column statistics and RM-two-way ANOVA followed by Sidak's multiple comparisons test for XY-plot analysis).

(D-E) The mean sEPSC amplitude in CA3 pyramidal neurons was not significantly different in lesioned and non-lesioned control cultures. Noteworthy, the mean sEPSC half width significantly decreased in lesioned cultures (D; $n_{control}$ = 25; $n_{3dp\ ECL}$ = 25; Mann-Whitney test; linear regression fit for XY-plot analysis). No lesion-induced changes in the mean sEPSC frequency or the amplitude/frequency-interdependence were detectable (E; Mann-Whitney test for column statistics and RM-two-way ANOVA followed by Sidak's multiple comparisons test for XY-plot analysis).

(F) No changes in passive membrane properties of CA3 pyramidal neurons were detected in synaptopodin-deficient cultures upon lesion (Mann-Whitney test for column statistics and RM-two-way ANOVA followed by Sidak's multiple comparisons test for XY-plot analysis).

(G-H) Paired whole-cell patch-clamp recordings were carried out between individual dentate granule cells (GC) and CA3 pyramidal neurons in lesioned and non-lesioned control cultures prepared from synaptopodin-deficient animals (G). Upon injection of 5 current pulses in the presynaptic granule cells, no changes in the pulse-number-dependent response rate of connected mossy fiber synapses were detectable after the ECL (H; $n_{control}$ = 12; $n_{3dp\ ECL}$ = 10; Mann-Whitney test for the respective pulse numbers).

(I) Analyzing properties of successfully transmitted uEPSCs, a significant decrease in uEPSC amplitude was evident in lesioned cultures when compared to non-lesioned controls (n°1$_{control}$ = 144 events, n°1$_{ECL}$ = 82 events; n°2$_{control}$ = 300 events, n°2$_{ECL}$ = 263 events; n°3$_{control}$ = 426 events (one data point outside the axis limits), n°3$_{ECL}$ = 402 events; n°4$_{control}$ = 486 events, n°4$_{ECL}$ = 364 events; n°5$_{control}$ = 472 events, n°5$_{ECL}$ = 317 events; Mann-Whitney test for the respective pulse numbers).

Individual data points are indicated by colored dots. Values represent mean ± s.e.m (*p < 0.05, ***p < 0.001, ns - non significant difference).

Figure 7: Synaptopodin deficiency shifted the hippocampal mossy fiber synapse to an excitable
state under baseline conditions.

(A-B) sEPSCs from CA3 pyramidal cells of both synaptopodin-deficient and wildtype tissue
preparations respectively. While the mean sEPSC amplitude was unchanged, sEPSC area and half width
were significantly increased in synaptopodin-deficient tissue cultures (A; $n_{Synpo+/+} = 48$; $n_{Synpo-/-} = 25$;
Mann-Whitney test; Linear regression fit for XY-plot analysis). Additionally, the sEPSC
amplitude/frequency-distribution was significantly changed between the two groups (B; RM-two-way
ANOVA followed by Sidak's multiple comparisons test).

(C-D) When comparing the response rate of postsynaptic CA3 neurons upon presynaptic action
potential generation from paired recordings in $Synpo^{+/+}$ and $Synpo^{-/-}$ tissue preparations, no significant
increase was observed (C; $n_{Synpo+/+} = 10$; $n_{Synpo-/-} = 12$; Mann-Whitney test for the respective pulse
numbers; c.f., Figures 3 and 6). However, when analyzing the amplitudes of successfully transmitted
uEPSCs at the postsynaptic CA3 pyramidal cells, a significant increase in uEPSC amplitudes became
evident in synaptopodin-deficient tissue preparations after each of the five pulses respectively (D;
$n°1_{Synpo+/+} = 20$ events, $n°1_{Synpo-/-} = 144$ events; $n°2_{Synpo+/+} = 115$ events, $n°2_{Synpo-/-} = 300$ events; $n°3_{Synpo+/+}$
$= 292$ events, $n°3_{Synpo-/-} = 426$ events (one data point outside the axis limits); $n°4_{Synpo+/+} = 326$ events,
$n°4_{Synpo-/-} = 486$ events; $n°5_{Synpo+/+} = 307$ events, $n°5_{Synpo-/-} = 472$ events; Mann-Whitney test for the
respective pulse numbers).

Individual data points are indicated by colored dots. Values represent mean ± s.e.m (*$p < 0.05$, **$p <$
0.01, ***$p < 0.001$, ns - non significant difference).

Figure 8: Entorhinal cortex lesion induced synaptopodin-dependent transcriptomic changes.

(A) Volcano Plots to visualize differential gene expression in wildtype (Synpo$^{+/+}$) and synaptopodin-

deficient (Synpo$^{-/-}$) tissue preparations based on region-specific transcriptome analysis of the dentate

gyrus and the CA3 area. Colored dots indicate differentially expressed genes ($p_{adj.} < 0.05$; CA3-Synpo$^{+/+}$:

$n_{control} = 2$, $n_{3dpECL} = 3$; DG-Synpo$^{+/+}$: $n_{control} = 4$, $n_{3dpECL} = 7$; CA3-Synpo$^{-/-}$: $n_{control} = 6$, $n_{3dpECL} = 6$; DG-

Synpo$^{-/-}$: $n_{control} = 7$, $n_{3dpECL} = 4$; detailed DESeq2-outputs provided in extended data table 8-1, 8-2, 8-3

and 8-4).

(B) VENN-diagrams illustrate the magnitude of lesion-induced differential gene expression and the

overlap in commonly regulated genes between wildtype and synaptopodin-deficient tissue preparations.

Synaptopodin-deficient preparations showed deficits in lesion-induced gene expression mainly in the

CA3 region. Despite the identification of commonly regulated genes, a considerable number of

genotype-specific changes can be identified in DG samples.

(C) For wildtype CA3 samples (Synpo$^{+/+}$-CA3), a gene set enrichment analysis (gene ontologies:

molecular function (MF) and cellular compartment (CC)) was performed and visualized (TOP-5) by the

g:Profiler web platform.

(D) Heatmap-visualization demonstrates the differential expression of synapse related genes (GOCC:

0045202) upon lesion-induced cellular remodeling in the wildtype CA3 region. TOP-10 differentially

expressed genes were additionally illustrated.

(E) For wildtype DG samples (Synpo$^{+/+}$-DG), a gene set enrichment analysis (gene ontologies:

molecular function (MF) and cellular compartment (CC)) was performed and visualized (TOP-5) by the

g:Profiler web platform.

(F) Heatmap-visualization demonstrates the differential expression of synapse related genes (GOCC:

0045202) upon lesion-induced cellular remodeling in the wildtype DG region. TOP-10 differentially

expressed genes for both UP and DOWN regulation were additionally illustrated.

Analysis was performed on the Galaxy platform, Differential gene expression was determined using the

DESeq2 application.

EXTENDED DATA TABLES

Extended data table 4-1: DESeq2 results table Synpo-coIP vs. input (BaseMean > 300; c.f. Fig. 4)

Extended data table 4-2: Synpo-coIP enriched genes (log2FC > 0; c.f. Fig. 4)

Extended data figure 4-3: Cytoscape map from Synpo-coIP enrichment experiments (c.f. Fig. 4)

Extended data table 5-1: DESeq2 results table ECL vs. control (input; Base Mean > 300; c.f. Figure 5)

Extended data table 5-2: DESeq2 results table ECL vs. control (Synpo-coIP; Base Mean > 300; c.f.
Figure 5)

Extended data table 8-1: DESeq2 results table ECL vs. control (Synpo$^{+/+}$, CA3; Base Mean > 300; c.f.
Figure 8)

Extended data table 8-2: DESeq2 results table ECL vs. control (Synpo$^{+/+}$, DG; Base Mean > 300; c.f.
Figure 8)

Extended data table 8-3: DESeq2 results table ECL vs. control (Synpo$^{-/-}$, CA3; Base Mean > 300; c.f.
Figure 8)

Extended data table 8-4: DESeq2 results table ECL vs. control (Synpo$^{-/-}$, DG; Base Mean > 300; c.f.
Figure 8)

A streptavidin DAPI

pre
DG
CA3 post

B control
CA3
10pA

3dp ECL
10pA

GC
30mV
50ms

C

response rate [fraction of total pulses] vs number of pulses

control
3dp ECL

D

uEPSC amplitude [pA] vs number of pulses

control
3dp ECL

n°1 ***
n°2 **
n°3 *
n°4 **
n°5 ***

A

B

Input

Synpo-coIP

tRNA 18S 28S

tRNA 18S 28S

Input

Beads-only-coIP

tRNA 18S 28S

tRNA 18S 28S

C

Gria1 mRNA normalized to Gapdh

**

input   coIP

D

Synpo-coIP vs. input

-log10(adj.p-value)

UP: 2355

DOWN: 3406

Gria1
Grin1
Synpo
Grin2a
Grin2b

log2(FC)

E

GOCC
Synpo-coIP
enriched

Synapse:
- Synapse
- Neuron-to-Neuron Synapse
- Postsynapse
- Cell-Cell-Contact-Zone
- Somatodendritic Compartment
- Synaptic Vesicle
- Neuron Projection
- Neuron-to-Neuron-Cytoplasm
- Dendritic Tree

Axon:
- Axon
- Distal Axon
- Axon Initial Segment
- Axon Cytoplasm
- Polarized Growth

Plasma Membrane:
- Anchoring Junction
- Plasma-Membrane-Region
- Cytoplasma-Region
- Lateral Plasma Membrane
- Tight Junction
- Cell Body

Actin Cytoskeleton:
- Cortical Actin Cytoskeleton
- Cortical Cytoskeleton
- Cell Cortex
- Transporter Complex
- T-Tubule

Microtubules:
- Microtubule Cytoskeleton
- Supramolecular Polymer
- Supramolecular Complex
- Midbody
- Microtubule Associated Complex
- Polymeric Cytoskeletal Fiber

Others:
- Histon Deacetylase Complex
- Inclusion Body

F control | 3dp ECL

G control | 3dp ECL

no SA
SA: 1 dense plate
SA: 2 dense plates
SA: 3 dense plates

H

● control ● 3dp ECL

**

dense plate length [nm]

Dentate Gyrus

A

sEPSC amplitude [pA]

sEPSC area [pA*ms]:
control: 101 ± 4.8
3dp ECL: 114 ± 3.6*

sEPSC half width [ms]

control
3dp ECL

B

sEPSC frequency [Hz] ns

control
3dp ECL

sEPSC frequency [Hz]

sEPSC amplitude [pA]

control
3dp ECL

C   Synpo$^{-/-}$

resting membrane potential [RMP, mV] ns

membrane response [mV]

current injection [pA]

control
3dp ECL

CA3

D

sEPSC amplitude [pA] ns

sEPSC area [pA*ms]:
control: 461 ± 40.1
3dp ECL: 357 ± 31.1

sEPSC half width [ms]

control
3dp ECL

E

sEPSC frequency [Hz] ns

control
3dp ECL

sEPSC frequency [Hz]

sEPSC amplitude [pA]

control
3dp ECL

F   Synpo$^{-/-}$

resting membrane potential [RMP, mV] ns

membrane response [mV]

current injection [pA]

control
3dp ECL

Paired Recordings

G

control
CA3   10pA

3dp ECL   10pA

GC   30mV
50ms

H

response rate [fraction of total pulses]

number of pulses

I   Synpo$^{-/-}$

control
3dp ECL

uEPSC amplitude [pA]

n°1   n°2   n°3   n°4   n°5
***   ***   ***   ***   ***

number of pulses