1 **Synaptopodin-associated ribosomal hubs regulate lesion-induced homeostatic**

2 **plasticity at hippocampal mossy fiber synapses**

Pia Kruse[1], Andreas Vlachos[1,2,3,§], Maximilian Lenz[1,§]

[1]Department of Neuroanatomy, Institute of Anatomy and Cell Biology, Faculty of Medicine, University of Freiburg,

Freiburg, Germany.

[2]Center for Basics in Neuromodulation (NeuroModulBasics), Faculty of Medicine, University of Freiburg, Freiburg,

Germany.

[3]Center Brain Links Brain Tools, University of Freiburg, Freiburg, Germany.

[§]Correspondence to andreas.vlachos@anat.uni-freiburg.de and maximilian.lenz@anat.uni-freiburg.de,

Department of Neuroanatomy, Institute of Anatomy and Cell Biology, Faculty of Medicine,

University of Freiburg, 79104 Freiburg, Germany

Keywords: homeostatic plasticity, denervation, synaptopodin, local protein synthesis

ABSTRACT

Neurological diseases can lead to the denervation of brain regions caused by demyelination, traumatic injury or cell death. Nevertheless, the molecular and structural mechanisms underlying the lesion-induced reorganization of denervated brain regions are a matter of ongoing investigation. In order to address this issue, we performed an entorhinal cortex lesion (ECL) in organotypic entorhinal-hippocampal tissue cultures and studied denervation-induced homeostatic plasticity of mossy fiber synapses, which connect dentate granule cells with CA3 pyramidal neurons and play important roles in spatial learning. Partial denervation caused a homeostatic strengthening of excitatory neurotransmission in dentate granule cells (GC), in CA3 pyramidal neurons, and their direct synaptic connections as revealed by paired recordings (GC-to-CA3). These functional changes were accompanied by ultrastructural reorganization of mossy fiber synapses, which regularly contain the plasticity-related protein synaptopodin and the spine apparatus organelle. We demonstrate that the spine apparatus organelle and its associated protein synaptopodin assemble ribosomes in close proximity to synaptic sites and moreover we unravel synaptopodin-related transcriptome, which can be linked to the expression of homeostatic synaptic plasticity. Notably, synaptopodin-deficient tissue preparations that lack the spine apparatus organelle, failed to express homeostatic adjustments of both excitatory neurotransmission and the region-specific transcriptome. Hence, synaptopodin and the spine apparatus organelle form local protein synthesis hubs that are essential for mediating lesion-induced homeostatic synaptic plasticity.

INTRODUCTION

Denervation is a common feature of various neurological diseases, which are accompanied by neuronal cell death, demyelination or traumatic injury of the central nervous system (Mesulam et al., 2019). As a result of the loss-of-input in otherwise healthy brain regions, complex reorganizational processes of neural circuits have been described. Among them, homeostatic synaptic plasticity has been established as an adaptive process of synaptic features that aims at compensating for perturbations of network activity (Turrigiano, 2012; Turrigiano, 2017; Vitureira et al., 2012). Specifically, excitatory synaptic strengthening upon denervation or activity deprivation has been reported as a compensatory synaptic adjustment (Turrigiano et al., 1998). Although homeostatic synaptic plasticity has been recognized as a local mechanism that can occur at distinct dendritic regions, the mechanisms that mediate dendritic autonomy in maintaining homeostasis remain unknown.

The hippocampal network is a crucial structure in memory acquisition. Here, the excitatory connection between dentate granule cells and CA3 pyramidal cells, i.e., the hippocampal mossy fiber synapse, has been demonstrated to play a crucial role in memory formation, maintenance and retrieval (Weng et al., 2018). Although numerous studies have addressed the unique plasticity features of this synapse (e.g. the onset of synaptic homeostasis and detonating transmission; (Lee et al., 2013)) and its impact on memory and behavior, no studies have addressed whether lesion-induced homeostatic synaptic plasticity occurs at this synapse when the main cortical excitatory input is affected, as seen for example in Alzheimer's disease (AD). In this context, organotypic tissue cultures of the entorhino-hippocampal complex provide the opportunity to study synaptic transmission at hippocampal mossy fiber synapses in a steady-state condition (Maus et al., 2020). Due to the laminated cortico-hippocampal input, the *in vitro* entorhinal-cortex lesion has been established as a model for the partial denervation of hippocampal neural circuits (Lenz et al., 2019; Prang et al., 2003; Vlachos et al., 2012a; Vlachos et al., 2012b).

In this study we used an entorhinal cortex lesion (ECL) of organotypic entorhino-hippocampal tissue cultures to study homeostatic adaptations in the dentate gyrus and CA3 network upon partial denervation. We provide evidence that partial denervation leads to excitatory synaptic strengthening in both dentate granule cells and CA3 pyramidal neurons. Using paired whole-cell patch-clamp recordings, we demonstrate changes in the synaptic transmission at hippocampal mossy fiber synapses that are accompanied by an ultrastructural reorganization of the spine apparatus organelle, which has previously been firmly linked to synaptic plasticity (Deller et al., 2003; Jedlicka and Deller, 2017; Jedlicka et al., 2009; Korkotian et al., 2014; Lenz et al., 2021a; Lenz et al., 2021b; Maggio and Vlachos, 2018; Vlachos et al., 2013; Vlachos et al., 2009). We demonstrate that synaptopodin and the spine apparatus organelle form ribosomal hubs that are linked to homeostatic plasticity-related local protein synthesis (Sutton et al., 2006). Consequently, synaptopodin-deficient tissue cultures, which lack the spine apparatus organelle, showed alterations in their ability to express homeostatic plasticity. Moreover, we demonstrate that synaptopodin-deficient tissue cultures have deficits in the denervation-induced adaptation of region-specific, i.e., dentate gyrus and CA3 transcriptomes. We therefore conclude, that

72   the spine apparatus organelle forms plastic local protein synthesis hubs that promote and mediate

73   homeostatic synaptic adjustments upon partial denervation.

MATERIALS AND METHODS

Ethics statement

Mice were maintained in a 12 h light/dark cycle with food and water available ad libitum. Every effort was made to minimize distress and pain of animals. All experimental procedures were performed according to German animal welfare legislation and approved by the appropriate animal welfare committee and the animal welfare officer of Albert-Ludwigs-University Freiburg, Faculty of Medicine (X-17/07K, X-17/09K, X-21/01B).

Preparation of tissue cultures

Entorhino-hippocampal tissue cultures were prepared at postnatal day 3-5 from C57BL/6J (wildtype, Synpo$^{+/+}$) and B6.129-Synpo^{tm1Mndl}/Dllr [Synpo$^{-/-}$, (Deller et al., 2003; Lenz et al., 2021a; Lenz et al., 2021b)] mice of either sex as previously described (Del Turco and Deller, 2007). Cultivation medium contained 50% (v/v) MEM, 25% (v/v) basal medium eagle, 25% (v/v) heat-inactivated normal horse serum, 25 mM HEPES buffer solution, 0.15% (w/v) bicarbonate, 0.65% (w/v) glucose, 0.1 mg/ml streptomycin, 100 U/ml penicillin, and 2 mM glutamax. The pH was adjusted to 7.3. All tissue cultures were allowed to mature for at least 18 days in a humidified atmosphere with 5% CO_2 at 35°C. Cultivation medium was replaced 3 times per week.

Entorhinal cortex lesion (ECL)

The entorhinal cortex of mature slice cultures (≥18 days in vitro) was transected with a sterile scalpel from the rhinal to the hippocampal fissure to achieve a complete lesion of the perforant path. Except for the lesion-induced partial denervation of the hippocampus, cytoarchitecture of both the hippocampus and the entorhinal cortex remained unchanged.

Whole-cell patch-clamp recordings

Whole-cell patch-clamp recordings were carried out at 35°C (3-6 neurons per culture). The bath solution contained (in mM) 126 NaCl, 2.5 KCl, 26 $NaHCO_3$, 1.25 NaH_2PO_4, 2 $CaCl_2$, 2 $MgCl_2$, and 10 glucose (aCSF) and was oxygenated continuously (5% CO_2 / 95% O_2). Patch pipettes contained (in mM) 126 K-gluconate, 10 HEPES, 4 KCl, 4 ATP-Mg, 0.3 GTP-Na$_2$, 10 PO-Creatine, 0.3% (w/v) biocytin (pH 7.25 with KOH, 290 mOsm with sucrose), having a tip resistance of 4-6 MΩ. Pyramidal neurons were visually identified using a LN-Scope (Luigs & Neumann) equipped with an infrared dot-contrast and a 40x water-immersion objective (NA 0.8; Olympus). Spontaneous excitatory postsynaptic currents (sEPSCs) of both dentate granule cells and CA3 pyramidal neurons were recorded in voltage-clamp mode at a holding potential of −70 mV. Series resistance was monitored before and after each recording and recordings were discarded if the series resistance reached ≥ 30 MΩ. For recording of intrinsic cellular properties in current-clamp mode, pipette capacitance of 2.0 pF was corrected and series resistance was compensated using the automated bridge balance tool of the MultiClamp commander. I-

111 V-curves were generated by injecting 1 s square pulse currents starting at −100 pA and increasing in 10
112 pA steps until +500 pA current injection was reached (sweep duration: 2 s).

Paired whole-cell patch-clamp recordings of hippocampal mossy fiber synapses

Action potentials were generated in the presynaptic dentate granule cell by 5 ms square current pulses
(1 nA). For connectivity analysis, 5 action potentials were applied at 20 Hz respectively (inter-sweep-
interval: 10 sec; at least 30 repetitions), while recording unitary excitatory postsynaptic currents
(uEPSCs) from CA3 pyramidal cells. Neurons were considered to be connected if > 5 % of action
potentials evoked time-locked inward uEPSCs. Recordings were performed in the presence of (-)-
bicuculline-methiodide (10 µM, #ab120108 Abcam) to prevent inhibitory synapse recruitment.
Moreover, glutamine was added to the recording medium to avoid synapse depletion.

Post hoc staining and imaging

Tissue cultures were fixed overnight in a solution of 4% (w/v) paraformaldehyde (PFA; in PBS (0.1M,
pH 7.4) with 4% (w/v) sucrose). After fixation, cultures were washed in PBS (0.1 M, pH 7.4) and
consecutively incubated for 1 h with 10% (v/v) normal goat serum (NGS) in 0.5% (v/v) Triton X-100
containing PBS to reduce nonspecific staining. For post-hoc visualization of patched granule cells and
CA3 pyramidal cells, Streptavidin Alexa Fluor 488 (Streptavidin A488, 1:1000; #S32354 Invitrogen)
was added to the incubation solution. DAPI nuclear stain (1:5000 in PBS for 10 minutes; #62248
Thermo Scientific) was used to visualize cytoarchitecture. Cultures were washed, transferred onto glass
slides and mounted for visualization with DAKO anti-fading mounting medium (#S302380-2, Agilent).

Transmission electron microscopy

Wildtype (Synpo$^{+/+}$) and Synpo$^{-/-}$ tissue cultures were fixed in 4 % paraformaldehyde (w/v) and 2 %
glutaraldehyde (w/v) in 0.1 M phosphate buffer (PB) overnight and washed for 1 hour in 0.1 M PB.
After fixation, tissue cultures were sliced with a vibratome and the slices were incubated with 1 %
osmium tetroxide for 20 min in 5 % (w/v) sucrose containing 0.1 M PB. The slices were washed 5 times
for 10 min in 0.1 M PB and washed in graded ethanol [10 min in 10 % (v/v) and 10 min in 20 % (v/v)].
The slices were then incubated with uranyl acetate [1 % (w/v) in 70 % (v/v) ethanol)] overnight
and were subsequently dehydrated in graded ethanol 80 % (v/v), 90 % (v/v) and 98 % (v/v) for 10
min. Finally, slices were incubated with 100 % (v/v) ethanol twice for 15 min followed by two
15 min washes with propylene oxide. The slices were then transferred for 30 min in a 1:1
mixture of propylene oxide with durcupan and then for 1 hour in durcupan. The durcupan was
exchanged with fresh durcupan and the slices were transferred to 4 °C overnight. The slices were then
embedded between liquid release-coated slides and coverslips. Cultures were re-embedded in blocks
and ultrathin sections were collected on copper grids. Electron microscopy was performed at a LEO
906E microscope (Zeiss) at 4646x magnification.

Regional mRNA library preparations and transcriptome analysis

RNA Library preparations for transcriptome analysis were performed using the NEBNext® Single Cell/Low Input RNA Library Prep Kit for Illumina® (New England Biolabs, #E6420) according to the manufacturer's instructions. Briefly, isolation of the dentate gyrus and CA3 area from individual tissue cultures was performed using a scalpel. One isolated dentate gyrus or CA3 area respectively was transferred to 7.5 µl lysis buffer (supplemented with murine RNase inhibitor) and homogenized using a pestill. Samples were centrifuged for 30 seconds at 10,000g and 5 µl of supernatant were collected from individual samples and further processed. After cDNA synthesis, cDNA amplification was performed according to the manufacturer's protocol with 12 PCR cycles. cDNA yield was consecutively analyzed by a High Sensitivity DNA assay on a Bioanalyzer instrument (Agilent). cDNA amount was adjusted to 10 ng for further downstream applications. After fragmentation and adaptor ligation, dual index primers (New England Biolabs, #E7600S) were ligated in a library amplification step using 10 PCR cycles. Libraries were finally cleaned up with 0.8X SPRI Beads following a standard bead purification protocol. Library purity and size distribution was assessed with a High Sensitivity DNA assay on a Bioanalyzer instrument (Agilent). We quantified the libraries using the NEBNext Library Quant Kit for Illumina (New England Biolabs, #E7630) based on the mean insert size provided by the Bioanalyzer. A 10 nM sequencing pool (120 µl in Tris-HCl, pH 8.5) was generated for sequencing on the NovaSeq6000 Sequencing platform (Illumina; service provided by CeGaT GmbH, Tübingen, Germany). We performed a paired-end sequencing with 101 bp read length. Data analysis was performed at the Galaxy platform (usegalaxy.eu). All files contained more than 10 M high-quality reads (after mapping to the reference genome; mm10) having at least a phred quality of 30 (>90% of total reads).

Synaptopodin co-Immunoprecipitation (SP-coIP) and RT-qPCR analysis

Six tissue cultures per biological replicate were washed in coIP-solution and dounce homogenized on ice in a solution containing (150 µl per sample, coIP-solution, in mM): 10 HEPES, 120 NaCl, 0.05% (v/v) NP-40, cOmplete protease inhibitor (Roche, according to manufacturers' instructions), murine RNase inhibitor (1:1000, New England Biolabs, #M0314) and cycloheximide (0.5 mg/ml, Sigma Aldrich, #C-4859). Input was precleared twice with protein A beads (10 µl per 100 µl sample/supernatant) to reduce unspecific binding. Protein A beads (10 µl, New England Biolabs, #S1425) were washed twice in coIP-solution and consecutively incubated for 1 hour at 4°C with an anti-Synaptopodin antibody (rabbit anti-Synaptopodin, Synaptic Systems, #163002). Beads were carefully washed twice to remove unbound antibodies and incubated for 1 hour with the precleared supernatant ('input') at 4°C with head-over-tail rotation. 10 µl of the precleared supernatant were kept as input control. After incubation, beads were washed twice with coIP-solution and finally dissolved in RNA protection buffer (New England Biolabs). RNA was released from the beads through Protein K lysis (Monarch® Total RNA Miniprep Kit; #T2010S New England Biolabs). After lysis, beads were removed from the solution in a magnetic rack and the RNA containing supernatant was transferred to fresh tubes.

RNA isolation on coIP- and input-samples was then performed according to the manufacturer's

instructions. RNA content was determined using the Agilent RNA 6000 Pico Kit (#5067-1513 Agilent)

with a 2100 Bioanalyzer (#G2939BA Agilent). Purified RNA was consecutively reverse transcribed

(RevertAid RT Kit; #K1691 Thermo Scientific). cDNA was diluted in water to a final concentration of

0.2 ng/µl. RT-qPCR was performed using a C1000 Touch Thermal Cycler (BIO-RAD) and the CFX

384 Real-Time PCR system (BIO-RAD). 0.9 ng target cDNA diluted in TaqMan Gene Expression

Master Mix (Applied Biosystems #4369016) were amplified using standard TaqMan Gene Expression

Assays (Applied Biosystems; Assay-IDs: Gapdh: Mm99999915_g1; Gria1: Mm00433753_m1). RNA

Library preparations for Synpo-related transcriptome analysis were performed using the NEBNext®

Single Cell/Low Input RNA Library Prep Kit for Illumina® (New England Biolabs, #E6420) according

to the manufacturer's instructions. We quantified the libraries using the NEBNext Library Quant Kit for

Illumina (New England Biolabs, #E7630) based on the mean insert size provided by the Bioanalyzer. A

10 nM sequencing pool (120 µl in Tris-HCl, pH 8.5) was generated for sequencing on the NovaSeq6000

Sequencing platform (Illumina; service provided by CeGaT GmbH, Tübingen, Germany). We

performed a paired-end sequencing with 151 bp read length. Data analysis was performed at the Galaxy

platform (usegalaxy.eu). All files contained more than 10 M high-quality reads (after mapping to the

reference genome; mm10) having at least a phred quality of 30 (>90% of total reads).

Quantification and statistics

Electrophysiological data were analyzed using pClamp 10.7 (Axon Instruments) and MiniAnalysis

(Synaptosoft) software. The fraction of action potentials followed by time-locked excitatory

postsynaptic current responses was considered as synaptic response rate. uEPSC amplitude was assessed

in uEPSCs from successfully transmitted action potentials as well as the mean amplitude of all

successfully evoked postsynaptic responses. In order to ensure selective analysis of time-locked

responses, events were included if > 5% of presynaptic pulses were successfully transmitted for each

pulse respectively. For individual pulses that caused ≤ 5% successfully transmitted events, response rate

was set to '0'. EPSC properties were analyzed using the automated template search tool for event

detection (Lenz et al., 2021b). In order to perform a hierarchical analysis of sEPSCs, all events of a

recorded cell were sorted based on their peak amplitude. Events above the 50th percentile therefore

correspond to all events above the cells median amplitude of events. Input resistance was calculated for

the injection of –100 pA current at a time frame of 200 ms with a maximum distance to the initial

hyperpolarization. Resting membrane potential was calculated as the mean baseline value. AP detection

was performed using the input-output curve threshold search event detection tool, and the AP frequency

was assessed by the number of APs detected during the respective current injection time. Ultrastructural

features were analyzed by an investigator blind to experimental conditions in randomly selected mossy

fiber terminals from electron micrographs of the stratum lucidum in non-lesioned and lesioned wildtype

tissue cultures.

RT-qPCR data were analyzed as previously described (Lenz et al., 2020) with the Bio-Rad CFX Maestro
1.0 software package using the ΔΔCq method with Gapdh as reference gene. Values were normalized
to the mean value of the respective control group.

Electron microscopy images were saved as TIF-files and analyzed using the ImageSP Viewer software
(http://e.informer.com/sys-prog.com).

RNA sequencing data were uploaded to the Galaxy web platform (public server: usegalaxy.eu; (Afgan
et al., 2018; Galaxy, 2022; Goecks et al., 2010)) and transcriptome analysis was performed using the
Galaxy platform in accordance with the reference-based RNA-seq data analysis tutorial (Batut et al.,
2021). Adapter sequencing, low quality, and short reads were removed via the CUTADAPT tool
(Galaxy version 3.5+galaxy0). Reads were mapped using RNA STAR (Galaxy version 2.7.8a+galaxy0)
with the mm10 full reference genome (Mus musculus). The evidence-based annotation of the mouse
genome (GRCm38), version M25 (Ensembl 100), served as gene model (GENCODE). For an initial
assessment of gene expression, unstranded FEATURECOUNT (Galaxy version 2.0.1+galaxy2)
analysis was performed from RNA STAR output. Only samples that contained >60% uniquely mapping
reads (feature: "exon") were considered for further analysis. Statistical evaluation was performed using
DESeq2 (Galaxy version 2.11.40.7+ galaxy1) with "pathway integrity" as the primary factor that might
affect gene expression. Genes with a low number of mean reads (< 300 BaseMean) were excluded from
further analysis. Genes were considered as differentially expressed if the adjusted p-value was < 0.05.
Data visualization was performed according to a modified version of a previously published workflow
(Reimand et al., 2019). A ranked list for the gene set enrichment analysis was generated based on Wald-
Stats from the DeSeq2 output. For the consecutive gene set enrichment analysis, we have used the GSEA
software package ((Mootha et al., 2003; Subramanian et al., 2005); BROAD institute; version
GSEA_4.2.1) with integrated MSigDB gene sets. Enrichment maps were built using cytoscape
((Shannon et al., 2003); version 3.9.0) with the EnrichmentMap application. Heatmaps were generated
based on z-scores of the normalized count table.

Data were statistically analyzed using GraphPad Prism 7 (GraphPad software, USA). All values
represent mean ± standard error of the mean. For statistical comparison of two experimental groups, a
Mann-Whitney test was employed. Short-term plasticity experiments were statistically assessed by the
repeated measure (RM) two-way ANOVA test with Sidak's (two groups) or Tukey's (three groups)
multiple comparisons test. uEPSC amplitude values from individual cells were stacked in subcolumns
and the pulse number was defining tabular rows (COLUMN factor: pathway, genetic background or
recombination; ROW factor: pulse number). P-values < 0.05 were considered statistically significant
(*p < 0.05, **p < 0.01, ***p < 0.001); results that did not yield significant differences are designated
'ns'. Statistical differences in XY-plots were indicated in the legend of the figure panels (*) when
detected through multiple comparisons, irrespective of their localization and the level of significance.
In the text and figures, values represent the mean ± standard error of the mean (s.e.m.).

Digital illustrations

Confocal image stacks were stored as TIF files. Figures were prepared using the ImageJ software

package (https://imagej.nih.gov/ij/) and Photoshop graphics software (Adobe, San Jose, CA, USA).

Image brightness and contrast were adjusted.

RESULTS

Entorhinal cortex lesion leads to homeostatic synaptic plasticity in both dentate granule cells and CA3 pyramidal neurons

Entorhinal cortex lesion was performed in mature tissue cultures to achieve a complete dissection of the perforant path leading to a partial denervation of both dentate granule cells and CA3 pyramidal neurons (Figure 1A). Spontaneous excitatory postsynaptic currents (sEPSCs) were recorded from dentate granule cells and CA3 pyramidal neurons in non-lesioned control and lesioned tissue cultures (3 days post entorhinal cortex lesion, 3dp ECL) to assess denervation-induced effects on excitatory neurotransmission (Figure 1B). As previously described [e.g., (Lenz et al., 2019; Vlachos et al., 2012b)], partial denervation caused a strengthening of excitatory synapses in dentate granule cells, i.e., a significant increase in both the mean sEPSC amplitude and half width as well as the mean sEPSC area (Figure 1C). A consecutive percentile analysis revealed that the lesion-induced increase in sEPSC amplitudes is evident when comparing the 50% biggest events in both groups while no effect was observed when comparing smaller events (Figure 1D). Notably, a significant upregulation in the mean sEPSC frequency was observed after lesion (Figure 1E) that resulted from an increased abundance of high amplitude sEPSC events as revealed by the amplitude/frequency-plot (Figure 1F).

Next, we examined excitatory neurotransmission in CA3 pyramidal neurons in non-lesioned and lesioned cultures (3dp ECL; Figure 1G-J). Again, a significant increase in both the mean sEPSC amplitude and area was observed in CA3 cells from lesioned cultures indicating a denervation-induced excitatory synaptic strengthening (Figure 1G). Notably, the mean sEPSC half width remained unchanged. In the sEPSC amplitude percentile analysis, excitatory synaptic strengthening seems to be restricted to the high amplitude events (Figure 1H). Moreover, the mean sEPSC frequency remained unchanged upon ECL (Figure 1I), which resulted from an increase in the abundance of high amplitude events (in the sEPSC amplitude range from 80-110 pA, RM two-way ANOVA followed by Sidak's multiple comparisons test) at the expense of small amplitude events (Figure 1J). We therefore conclude that ECL leads to a homeostatic increase in excitatory neurotransmission in both dentate granule cells and CA3 pyramidal neurons.

Partial denervation does not cause changes in intrinsic cellular properties

Passive and active membrane properties were assessed in both dentate granule cells (Figure S1A-C) and CA3 pyramidal neurons (Figure S1D-F) to test for homeostatic adaptations of intrinsic cellular properties upon lesion. In dentate granule cells no changes in passive membrane properties (Figure S1A), i.e., the resting membrane potential (RMP) and the input resistance as well as the input-output curve dynamics were observed (Figure S1B). Additionally, the action potential frequency of dentate granule cells was similar in lesioned and non-lesioned cultures (Figure S1C). Noteworthy, RMP assessment confirmed the mature state of the recorded dentate granule cells [c.f. (Strehl et al., 2018)].

299 Concerning CA3 pyramidal neurons, neither the passive membrane properties (Figure S1D) – including
300 input-output curve dynamics (Figure S1E) – nor the active properties (Figure S1F) were changed. Based
301 on these results, we conclude that intrinsic cellular properties were stable following denervation in both
302 tested cell types of the hippocampal network.
303
Entorhinal cortex lesion induces changes in synaptic transmission at hippocampal mossy fiber
synapses

Hippocampal mossy fiber synapses have previously been linked to learning processes and memory
formation (Lipp et al., 1988; Vandael et al., 2020). Concerning their synaptic features, they have been
described as synapses that generate large postsynaptic currents upon activation (Henze et al., 2002a).
Given that our results demonstrate changes in both investigated regions, which additionally hint towards
a preferential lesion-related plasticity induction in readily strong synapses at CA3 pyramidal neurons,
we next assessed the features of synaptic transmission at hippocampal mossy fiber synapses.
Therefore, we performed paired whole-cell patch-clamp recordings of individual dentate granule cells
and CA3 pyramidal neurons in organotypic tissue cultures (Figure 2A). To test for synaptic connections,
5 consecutive action potentials at 20 Hz frequency were elicited in dentate granule cells (Figure 2B).
Unitary excitatory postsynaptic currents (uEPSCs) were monitored in CA3 pyramidal neurons in non-
lesioned and lesioned tissue cultures. We found that a train of presynaptic action potentials in connected
neurons induced a robust facilitation of postsynaptic responses in both postsynaptic response rate and
uEPSC amplitude under baseline conditions (Figure 2B-D). Notably, the response rate to the first
presynaptic action potential, which is the number of presynaptic action potentials that generate a
postsynaptic response, was very low (~3%; Figure 2C). In lesioned cultures, however, a significant
increase in the synaptic reliability for the first two pulses was observed. For the last three pulses in one
train of stimulation, averaged synaptic response rate levels became similar (Figure 3C). Moreover, the
uEPSC amplitude was significantly increased in response to the individual pulses 3 days after ECL
(Figure 2D). This novel approach of GC-to-CA3 paired recordings in the steady-state environment of
organotypic tissue cultures provided evidence that synaptic transmission at hippocampal mossy fiber
synapses is a target for lesion-induced remodeling of hippocampal networks.
327
Synaptopodin and the spine apparatus organelle represent dendritic ribosomal hubs that are
associated with excitatory synapse related mRNAs

Hippocampal mossy fiber synapses show a complex organization of their pre- and postsynaptic sites.
Among these features, a regular occurrence of the spine apparatus organelle, which is a transmission-
and plasticity-regulating organelle, has been reported at the postsynaptic site (Chicurel and Harris,
1992). We hypothesized that the spine apparatus organelle and its associated protein synaptopodin
mediate their effects on mossy fiber plasticity by interfering with local protein synthesis. Using
transmission electron microscopy, a close interaction between ribosomes and the spine apparatus

organelle was readily revealed (Figure 3A). To test whether synaptopodin, which is a marker protein
and essential component of the spine apparatus organelle (Deller et al., 2003), interacts with local
translation machineries, i.e., dendritic ribosomes, we performed synaptopodin co-immunoprecipitation
(Synpo-coIP) experiments with subsequent RT-qPCR analysis. Using RNA assessment on a
Bioanalyzer, we found rRNA peaks in Synpo-coIP experiments whereas no rRNA was detected in bead-
only controls (Figure 3B). RT-qPCR analysis (reference gene: Gapdh) from input and Synpo-coIP
samples revealed that the Gria1 mRNA, which has been linked to synaptopodin-dependent synaptic
plasticity (Vlachos et al., 2009) and homeostatic synaptic plasticity (Aoto et al., 2008; Sutton et al.,
2006), is significantly enriched in Synpo-coIP samples (Figure 3C). We therefore conclude, that
synaptopodin and the spine apparatus organelle might serve as a ribosomal hub that significantly
contributes to plasticity-related local protein synthesis.

To further unravel the plasticity-mediating features of synaptopodin and the spine apparatus organelle,
we performed a transcriptome analysis of Synpo-coIP enriched mRNA (Figure 3D, Table S1). A
considerable amount of mRNAs were enriched after Synpo-coIP compared to input samples: 2355
mRNAs showed a higher abundance in Synpo-coIP samples compared to the respective input samples
(UP). Subsequently, a gene set enrichment analysis (GSEA) was performed to elucidate the functional
relevance of synaptopodin-associated mRNAs (Figure 3E, Table S2). We identified distinct clusters of
cellular compartment gene sets that have been enriched by Synpo-coIP. Among them, synapse and actin
cytoskeleton-related gene sets were identified that have been linked to synaptic plasticity. Of note, these
data suggest the synaptopodin-dependent assembly of localized organelle associated transcriptome hubs
that can be dedicated to local protein synthesis.

The synaptopodin-associated transcriptome remains stable upon lesion
We next tested whether the composition of the synaptopodin-associated transcriptome changes in
lesioned networks 3dp ECL. Therefore, a differential mRNA abundance analysis was performed in
Synpo-coIP and their respective input RNA samples from lesioned and non-lesioned wildtype cultures
(Figure S2). In mRNA libraries prepared from input RNA samples, we found a considerable amount of
differentially expressed genes upon ECL (Table S3). In contrast, a limited amount of mRNAs were
differentially enriched upon ECL in Synpo-coIP samples (Table S4). We therefore conclude that the
composition of the synaptopodin-associated transcriptome remain stable upon partial denervation.

Ultrastructural reorganization of the ribosome associated spine apparatus organelle
Previous studies have demonstrated that the ultrastructural organization of the spine apparatus organelle
might represent a target for plasticity (Lenz et al., 2021b; Vlachos et al., 2013). To study lesion-induced
effects on the ultrastructural organization of the ribosome associated spine apparatus organelle that
might interfere with local protein synthesis, we performed transmission electron microscopy of mossy
fiber synapses in non-lesioned and lesioned (3dp ECL) entorhino-hippocampal tissue cultures (Figure

3F; c.f., (Vlachos et al., 2013)). No differences in the structural organization of pre- and postsynaptic

sites could be detected between the groups (presynapse: mossy fiber perimeter: 11.61 ± 0.4 μm (control;

n = 120) vs. 10.88 ± 0.35 μm (3dp ECL; n = 90); postsynapse: number of PSDs: 4.91 ± 0.33 (control; n

= 120) vs. 4.6 ± 0.27 (3dp ECL; n = 90), length of PSDs: 201 nm ± 4.35 (control; n = 589) vs. $201.7 \pm$

5.42 nm (3dp ECL; n = 414); values indicate mean $\pm$ sem, Mann-Whitney test). Nevertheless, a

substantial reorganization of the spine apparatus organelle was evident in lesioned cultures 3 days after

injury compared to non-lesioned controls (Figure 3G, H). We found a significantly increased number

of dense plates in spine apparatus organelles from lesioned cultures (Figure 3G). At the same time the

length of individual dense plates was significantly reduced in lesioned cultures (Figure 3H). Therefore,

we conclude that the spine apparatus organelle, which is an mRNA hub for local protein synthesis,

represents a target for lesion-induced changes at hippocampal mossy fiber synapses.

Synaptopodin-deficient tissue cultures show deficits in homeostatic synaptic plasticity upon

partial denervation

Based on our ultrastructural analysis, we hypothesized that the reorganization of the spine apparatus

organelle might contribute to homeostatic synaptic plasticity at hippocampal mossy fiber synapses. To

gain further insights concerning the role of the spine apparatus organelle in this adaptive process, we

performed experiments in synaptopodin-deficient tissue preparations, which lack the spine apparatus

organelle (Figure 4; (Deller et al., 2003)).

Synaptopodin-deficient dentate granule cells showed a significant increase in both the mean sEPSC

amplitude and area as well as the mean sEPSC half width 3 days following ECL (Figure 4A). In contrast,

no changes in the mean sEPSC frequency, which have been observed in wildtype (Synpo$^{+/+}$) cultures,

were detected (Figure 4B). Moreover, passive membrane properties remained stable upon lesion in

synaptopodin-deficient tissue preparations (Figure 4C).

When assessing excitatory neurotransmission onto CA3 pyramidal neurons in lesioned and non-lesioned

cultures, no significant changes in the sEPSC event analysis (Figure 4D) or the mean sEPSC frequency

(Figure 4E) were observed. Again, passive membrane properties remained unchanged (Figure 4F).

Thus, we here provide evidence for deficits in the implementation of homeostatic synaptic adaptations

for both dentate granule cells and CA3 pyramidal neurons in synaptopodin-deficient tissue cultures.

We extended these findings with GC-to-CA3 paired whole-cell patch-clamp recordings in

synaptopodin-deficient preparations (Figure 4G). In contrast to wildtype cultures, no significant

differences in the synaptic reliability could be detected 3 days after ECL (Figure 4H). When analyzing

the uEPSC amplitude of successfully transmitted events, surprisingly, we found a significant reduction

in the pulse-related uEPSC amplitude in lesioned synaptopodin-deficient preparations (Figure 4I), while

excitatory neurotransmission seemed facilitated under baseline conditions as compared to wildtype

cultures (Figure S3). Hence, synaptopodin-deficient mossy fiber synapses responded in a synaptic

depression rather than homeostatic synaptic strengthening after the lesion. These findings demonstrate

the role of the postsynaptic spine apparatus organelle on plasticity of mossy fiber synapses. They are
consistent with recent work that suggested a role of synaptopodin and the spine apparatus in mediating
metaplasticity (Hulme et al., 2014; Lenz et al., 2021a).
413

Entorhinal cortex lesion causes synaptopodin-dependent changes in the region-specific

hippocampal transcriptome

To further unravel lesion-induced changes in hippocampal subfields that might depend on synaptopodin,
we performed transcriptome analysis of both the dentate gyrus and the CA3 region in non-lesion and
lesioned cultures from wildtype (Synpo$^{+/+}$) and synaptopodin-deficient (Synpo$^{-/-}$) animals (Figure 5).
The expression analysis revealed a considerable number of genes that are differentially expressed in
both hippocampal regions from wildtype animals (Figure 5A, B). In contrast, synaptopodin-deficient
tissue culture preparations showed deficits in the lesion-induced adjustment of gene expression in CA3
whereas in the dentate gyrus it remained evident (Figure 5A). Moreover, we demonstrate that wildtype
and synaptopodin-deficient tissue preparations show a genotype dependent gene expression adjustment
upon lesion in both regions although a common set of regulated genes has been identified (Figure 6B;
for detailed tabular results see supplementary tables S5-S8). We therefore conclude, that synaptopodin
might act as a sensor that triggers differential gene expression towards homeostatic adjustments in a
region-specific fashion.
The subsequently performed geneset enrichment analysis with an additional enrichment map
visualization (GSEA, Figure 5C) provided evidence that the ECL caused an inflammatory response in
the hippocampus of wildtype cultures that might result from the accumulation of microglia. Moreover,
partial denervation is accompanied by changes in metabolic gene sets (metabolic clusters), differences
in membrane organization and a potential impact on neurogenesis. Further assessment of the
differentially expressed genes suggested that these changes might be the result of a region-specific
microglia accumulation in the denervated areas.
In order to learn more about the lesion-induced differential expression of genes that have been linked to
synaptic plasticity ("synapse" GOCC: 0045202), the region-specific lists of differentially expressed
genes were filtered according to the content of the respective gene ontology (Figure 5D). We found
numerous synaptic genes in both regions that show a differential expression upon lesion, which
corresponds to the functional assessment of synaptic transmission where a strengthening of excitatory
neurotransmission was observed. Therefore, we conclude that the lesion-induced change in the region-
specific transcriptome might represent an essential component of the homeostatic synaptic response.
442

DISCUSSION

Denervation is a common phenomenon in numerous medical conditions affecting the central nervous system, which can trigger homeostatic adaptations in neural networks (Barnes et al., 2017; Bissen et al., 2021; Hengen et al., 2013; Lambo and Turrigiano, 2013; Platschek et al., 2016; Vlachos et al., 2012b). In this study, we have unraveled a novel feature of the spine apparatus organelle that crucially contributes to the localized expression of compensatory synaptic changes. In line with previous findings that employed proximity proteomics of synaptopodin (Falahati et al., 2022), we identified a close interaction between synaptopodin and ribosomes. Using subsequent synaptopodin co-immunoprecipitation experiments, we here elucidated the synaptopodin-related transcriptome, which might contribute to plasticity-related local protein synthesis. Together with findings that indicate interactions between synaptopodin and ER-related proteins (Falahati et al., 2022), we suggest a role for synaptopodin and the spine apparatus organelle as local plasticity hubs in neuronal dendrites.

To study lesion-induced changes in synaptic transmission, we used the *in vitro* model of ECL in organotypic entorhino-hippocampal tissue cultures to study the effects of a partial loss of input on hippocampal synaptic transmission. After *in vitro* maturation, tissue organization has reached a steady-state that is characterized by structural and functional stability of excitatory synapses and that resembles the ultrastructural synaptic architecture *in vivo* (Haber et al., 2006; Heller and Rusakov, 2017). In this context, recent work has demonstrated that organotypic hippocampal tissue cultures have intact mossy fiber pathways and synapses (Maus et al., 2020). Moreover, tissue cultures of the hippocampal formation show an *in vivo* like cellular architecture and a preserved lamination in which the excitatory cortical input – the perforant path – terminates at distal apical dendrites of dentate granule cells and pyramidal neurons of the ammonic subregions (Frotscher and Heimrich, 1995; Frotscher et al., 1997; Frotscher et al., 1995). Our results provide evidence for homeostatic excitatory synaptic plasticity in both dentate granule cells and CA3 pyramidal neurons upon lesion, which is accompanied by changes in the region-specific transcriptome and the abundance of synapse-related mRNAs. Moreover, we here established a novel method to assess synaptic transmission at individual hippocampal mossy fiber synapses – the monosynaptic excitatory contacts between dentate granule cells and CA3 pyramidal neurons – in a steady-state environment by paired whole-cell patch-clamp recordings of connected neurons. In this experimental approach, we identified synaptic transmission at hippocampal mossy fiber synapses as a target for denervation induced synaptic plasticity. In addition, ultrastructural changes in the spine apparatus organelle, which is well established in the regulation of synaptic plasticity (Lenz et al., 2021b; Vlachos et al., 2013), can be observed after ECL. Since homeostatic adaptations on both synaptic and transcriptomic levels were altered in synaptopodin-deficient animals which lack the spine apparatus organelle, we conclude its crucial role in mediating as well as sensing and triggering compensatory synaptic changes.

The dentate gyrus/CA3 neural network is an integral component of the memory-associated trisynaptic hippocampal pathway. The hippocampal mossy fiber pathway forms large *en passant* synapses and

connects one individual granule cell to a limited number of CA3 pyramidal neurons (Acsady et al.,
1998). Both regions receive cortical input – the perforant path – from entorhinal layer II stellate cells
that can induce, promote and modulate information processing in hippocampal subregions. Moreover,
this pathway is a well-established target for various diseases; among them traumatic, inflammatory and
neurodegenerative conditions like Alzheimer's disease (AD; (Robinson et al., 2014; Thal et al., 2000)).
Here, we demonstrate that both dentate granule cells and CA3 pyramidal neurons express synaptic
plasticity upon partial denervation, which suggests that information processing at the connection
between these cells might be affected.

The expression of synaptic plasticity in the hippocampal neural circuit is involved in crucial aspects of
normal brain function, such as learning and memory acquisition. The hippocampal mossy fiber synapse,
which is the monosynaptic excitatory connection between dentate granule cells and CA3 pyramidal
neurons, has previously been described as an integral component of this network that relates to distinct
forms of memory acquisition and retrieval (Hunsaker et al., 2008; Marr, 1971). Specifically, activity-
dependent structural and functional changes in synaptic architecture and transmission have been found
to encode distinct input patterns for a prolonged period of time (Lee et al., 2013; Vandael et al., 2020).
Notably, alterations in hippocampal mossy fiber transmission have been reported in animal models of
Alzheimer's disease (AD) and aging related cognitive decline (Viana da Silva et al., 2019; Villanueva-
Castillo et al., 2017) where a particular loss in cortical input pathways onto hippocampal neurons can
be found. Nevertheless, it remains unclear if the lesion-induced synaptic response deteriorates cognitive
performance. Given that (1) homeostatic synaptic plasticity is altered in synaptopodin-deficient tissue
preparations and (2) alterations in synaptopodin expression have been linked to cognitive decline
(Wingo et al., 2019), we suggest that synaptopodin-mediated synaptic stability as well as compensatory
plasticity aim at maintaining proper network function.

Recently, characteristic features of transmission and stabilizing factors at hippocampal mossy fiber
synapses have been reported (De Bruyckere et al., 2018; Vandael et al., 2021b; Vyleta et al., 2016;
Vyleta and Jonas, 2014). Upon presynaptic activity, a detonating behavior can be observed in which a
limited number of – but not single – action potentials can elicit a suprathreshold postsynaptic
depolarisation (Henze et al., 2002b). Although this phenomenon has been extensively described in
mossy fiber bouton-CA3 pyramidal cell paired recordings (Vandael et al., 2021a), we were not able to
reveal detonating characteristics in granule cell-CA3 pyramidal cell pairs. Here, we assume that both
differences in the experimental model and methodology can account for these differences, which
warrant further clarification in future studies. Nevertheless, a filtering of single action potentials with a
subsequent robust facilitation of synaptic transmission at this synapse was evident, which is in line with
previous reports demonstrating an initially low presynaptic release probability with consecutive
facilitation (Gundlfinger et al., 2007; Zalutsky and Nicoll, 1990). Our results demonstrate that the partial
loss of input following lesion can influence both synaptic reliability and potency, which has been
observed for other plasticity-inducing stimuli at this synapse (Zalutsky and Nicoll, 1990). We suggest

that lesion-induced modulation of dual filtering/facilitation properties at connected pairs of dentate
granule cells and CA3 pyramidal neurons might interact with both pattern separation and completion
along the dentate gyrus/CA3 axis in a way that the increase in synaptic reliability and potency may shift
the balance towards pattern completion rather than separation (Leutgeb et al., 2007; O'Reilly and
McClelland, 1994; Prince et al., 2016). Moreover, the presence of synaptopodin will presumably
interfere with this network process under baseline and plasticity-inducing conditions which might
account for the previously reported cognitive deficits in synaptopodin-deficient mice (Deller et al.,
2003) and the synaptopodin-related cognitive decline during human aging (Wingo et al., 2019).

Furthermore, it is an established view that mossy fiber synaptic plasticity is predominantly mediated by
presynaptic processes (Monday et al., 2022; Orlando et al., 2021; Vandael et al., 2020). Nevertheless,
recent work has suggested that the postsynaptic site shapes the transmission and plasticity through
transsynaptic mechanisms (Vandael et al., 2021b). In this study, we provide evidence that the
postsynaptically located spine apparatus organelle mediates denervation-induced synaptic plasticity at
this synapse. Since its discovery, the function of the spine apparatus organelle and the mechanisms
through which it orchestrates synaptic plasticity remained enigmatic. We here demonstrate that
synaptopodin and the spine apparatus organelle represent dendritic ribosomal hubs that contribute to
plasticity-driven local protein synthesis. The lesion-induced ultrastructural remodeling of the spine
apparatus organelle might therefore represent a novel regulatory mechanism for localized changes in
synaptic transmission (Dubes et al., 2022; Vlachos et al., 2009).

Notably, we found opposing plasticity phenomena at hippocampal mossy fiber synapses on the level of
individual neuron pairs in wildtype and synaptopodin-deficient tissue preparations where synaptopodin
deficiency did not only cause defects in plasticity expression but hampered the ability of neurons to
maintain synaptic stability upon lesion. While the loss of input in wildtype cultures led to a significant
homeostatic strengthening of DG-CA3 excitatory synaptic transmission, synaptopodin deficiency was
associated with a lesion-related loss of synaptic potency. We therefore conclude that the function of the
spine apparatus organelle and its related protein synaptopodin might not be restricted to the mediation
of receptor accumulation at postsynaptic sites but rather comprises the implementation of plasticity
directionality (Vlachos, 2012). However, the link between synaptopodin and recently described
mechanisms that switch plasticity directionality from homeostatic to associative rules remain to be
determined (Wu et al., 2022). On the other hand, the lack of synaptopodin might decrease synaptic
resilience due to the lack of stabilizing features (Yap et al., 2020), which is required to maintain synaptic
transmission upon lesion. Regardless of these considerations, once more we report that synaptopodin
and the spine apparatus organelle act as crucial regulators of synaptic plasticity.

Moreover, synaptic transcriptional changes as seen in wildtype cultures were absent in synaptopodin-
deficient cultures. Since the differential expression of lesion-related inflammatory genes that might
promote plasticity induction was not observed in synaptopodin-deficient cultures, a link between
synaptopodin and proinflammatory plasticity promotion might exist (Eichler et al., 2022; Strehl et al.,

2014). Due to these observations, we suggest a dual role for synaptopodin as activity sensor and

plasticity effector, which assembles synapse-related mRNAs at dendritic plasticity hubs.

ACKNOWLEDGEMENTS

The Galaxy server that was used for some calculations in this study is partially funded by Collaborative Research Centre 992 Medical Epigenetics (DFG grant SFB 992/1 2012) and German Federal Ministry of Education and Research (BMBF grants 031 A538A/A538C RBC, 031L0101B/031L0101C de.NBI-epi, and 031L0106 de.STAIR (de.NBI)). We thank Bernd Heimrich for helpful and stimulating discussions on ultrastructural mossy fiber architecture. Moreover, we thank Sigrun Nestel for excellent technical assistance in electron microscopy. This work was supported by Else Kröner-Fresenius-Stiftung (EKFS_#2019_A94 to ML) and Deutsche Forschungsgemeinschaft (DFG; Project-ID 259373024 B14–CRC/TRR 167 to AV).

REFERENCES

Acsady, L., Kamondi, A., Sik, A., Freund, T., and Buzsaki, G. (1998). GABAergic cells are the major postsynaptic targets of mossy fibers in the rat hippocampus. J Neurosci *18*, 3386-3403.

Afgan, E., Baker, D., Batut, B., van den Beek, M., Bouvier, D., Cech, M., Chilton, J., Clements, D., Coraor, N., Gruning, B.A*., et al.* (2018). The Galaxy platform for accessible, reproducible and collaborative biomedical analyses: 2018 update. Nucleic Acids Res *46*, W537-W544.

Aoto, J., Nam, C.I., Poon, M.M., Ting, P., and Chen, L. (2008). Synaptic signaling by all-trans retinoic acid in homeostatic synaptic plasticity. Neuron *60*, 308-320.

Barnes, S.J., Franzoni, E., Jacobsen, R.I., Erdelyi, F., Szabo, G., Clopath, C., Keller, G.B., and Keck, T. (2017). Deprivation-Induced Homeostatic Spine Scaling In Vivo Is Localized to Dendritic Branches that Have Undergone Recent Spine Loss. Neuron *96*, 871-882 e875.

Batut, B., van den Beek, M., Doyle, M.A., and Soranzo, N. (2021). RNA-Seq Data Analysis in Galaxy. Methods Mol Biol *2284*, 367-392.

Bissen, D., Kracht, M.K., Foss, F., Hofmann, J., and Acker-Palmer, A. (2021). EphrinB2 and GRIP1 stabilize mushroom spines during denervation-induced homeostatic plasticity. Cell Rep *34*, 108923.

Chicurel, M.E., and Harris, K.M. (1992). Three-dimensional analysis of the structure and composition of CA3 branched dendritic spines and their synaptic relationships with mossy fiber boutons in the rat hippocampus. J Comp Neurol *325*, 169-182.

De Bruyckere, E., Simon, R., Nestel, S., Heimrich, B., Katzel, D., Egorov, A.V., Liu, P., Jenkins, N.A., Copeland, N.G., Schwegler, H*., et al.* (2018). Stability and Function of Hippocampal Mossy Fiber Synapses Depend on Bcl11b/Ctip2. Front Mol Neurosci *11*, 103.

Del Turco, D., and Deller, T. (2007). Organotypic entorhino-hippocampal slice cultures--a tool to study the molecular and cellular regulation of axonal regeneration and collateral sprouting in vitro. Methods Mol Biol *399*, 55-66.

Deller, T., Korte, M., Chabanis, S., Drakew, A., Schwegler, H., Stefani, G.G., Zuniga, A., Schwarz, K., Bonhoeffer, T., Zeller, R*., et al.* (2003). Synaptopodin-deficient mice lack a spine apparatus and show deficits in synaptic plasticity. Proc Natl Acad Sci U S A *100*, 10494-10499.

Dubes, S., Soula, A., Benquet, S., Tessier, B., Poujol, C., Favereaux, A., Thoumine, O., and Letellier, M. (2022). miR-124-dependent tagging of synapses by synaptopodin enables input-specific homeostatic plasticity. EMBO J, e109012.

Eichler, A., Kleidonas, D., Turi, Z., Fliegauf, M., Kirsch, M., Pfeifer, D., Masuda, T., Prinz, M., Lenz, M., and Vlachos, A. (2022). Microglia mediate synaptic plasticity induced by 10 Hz repetitive transcranial magnetic stimulation. bioRxiv.

Falahati, H., Wu, Y., Feuerer, V., Simon, H.G., and De Camilli, P. (2022). Proximity proteomics of synaptopodin provides insight into the molecular composition of the spine apparatus of dendritic spines. Proc Natl Acad Sci U S A *119*, e2203750119.

Frotscher, M., and Heimrich, B. (1995). Lamina-specific synaptic connections of hippocampal neurons in vitro. J Neurobiol *26*, 350-359.

Frotscher, M., Heimrich, B., and Deller, T. (1997). Sprouting in the hippocampus is layer-specific. Trends Neurosci *20*, 218-223.

Frotscher, M., Heimrich, B., Deller, T., and Nitsch, R. (1995). Understanding the cortex through the hippocampus: lamina-specific connections of the rat hippocampal neurons. J Anat *187 (Pt 3)*, 539-545.

Galaxy, C. (2022). The Galaxy platform for accessible, reproducible and collaborative biomedical analyses: 2022 update. Nucleic Acids Res.

Goecks, J., Nekrutenko, A., Taylor, J., and Galaxy, T. (2010). Galaxy: a comprehensive approach for supporting accessible, reproducible, and transparent computational research in the life sciences. Genome Biol *11*, R86.

Gundlfinger, A., Leibold, C., Gebert, K., Moisel, M., Schmitz, D., and Kempter, R. (2007). Differential modulation of short-term synaptic dynamics by long-term potentiation at mouse hippocampal mossy fibre synapses. J Physiol *585*, 853-865.

Haber, M., Zhou, L., and Murai, K.K. (2006). Cooperative astrocyte and dendritic spine dynamics at hippocampal excitatory synapses. J Neurosci *26*, 8881-8891.

Heller, J.P., and Rusakov, D.A. (2017). The Nanoworld of the Tripartite Synapse: Insights from Super-Resolution Microscopy. Front Cell Neurosci *11*, 374.

Hengen, K.B., Lambo, M.E., Van Hooser, S.D., Katz, D.B., and Turrigiano, G.G. (2013). Firing rate homeostasis in visual cortex of freely behaving rodents. Neuron *80*, 335-342.

Henze, D.A., McMahon, D.B., Harris, K.M., and Barrionuevo, G. (2002a). Giant miniature EPSCs at the hippocampal mossy fiber to CA3 pyramidal cell synapse are monoquantal. J Neurophysiol *87*, 15-29.

Henze, D.A., Wittner, L., and Buzsaki, G. (2002b). Single granule cells reliably discharge targets in the hippocampal CA3 network in vivo. Nat Neurosci *5*, 790-795.

Hulme, S.R., Jones, O.D., Raymond, C.R., Sah, P., and Abraham, W.C. (2014). Mechanisms of heterosynaptic metaplasticity. Philos Trans R Soc Lond B Biol Sci *369*, 20130148.

Hunsaker, M.R., Rosenberg, J.S., and Kesner, R.P. (2008). The role of the dentate gyrus, CA3a,b, and CA3c for detecting spatial and environmental novelty. Hippocampus *18*, 1064-1073.

Jedlicka, P., and Deller, T. (2017). Understanding the role of synaptopodin and the spine apparatus in Hebbian synaptic plasticity - New perspectives and the need for computational modeling. Neurobiol Learn Mem *138*, 21-30.

Jedlicka, P., Schwarzacher, S.W., Winkels, R., Kienzler, F., Frotscher, M., Bramham, C.R., Schultz, C., Bas Orth, C., and Deller, T. (2009). Impairment of in vivo theta-burst long-term potentiation and network excitability in the dentate gyrus of synaptopodin-deficient mice lacking the spine apparatus and the cisternal organelle. Hippocampus *19*, 130-140.

Korkotian, E., Frotscher, M., and Segal, M. (2014). Synaptopodin regulates spine plasticity: mediation by calcium stores. J Neurosci *34*, 11641-11651.

Lambo, M.E., and Turrigiano, G.G. (2013). Synaptic and intrinsic homeostatic mechanisms cooperate to increase L2/3 pyramidal neuron excitability during a late phase of critical period plasticity. J Neurosci *33*, 8810-8819.

Lee, K.J., Queenan, B.N., Rozeboom, A.M., Bellmore, R., Lim, S.T., Vicini, S., and Pak, D.T. (2013). Mossy fiber-CA3 synapses mediate homeostatic plasticity in mature hippocampal neurons. Neuron *77*, 99-114.

Lenz, M., Eichler, A., Kruse, P., Muellerleile, J., Deller, T., Jedlicka, P., and Vlachos, A. (2021a). All-trans retinoic acid induces synaptopodin-dependent metaplasticity in mouse dentate granule cells. Elife *10*.

Lenz, M., Galanis, C., Kleidonas, D., Fellenz, M., Deller, T., and Vlachos, A. (2019). Denervated mouse dentate granule cells adjust their excitatory but not inhibitory synapses following in vitro entorhinal cortex lesion. Exp Neurol *312*, 1-9.

Lenz, M., Kruse, P., Eichler, A., Straehle, J., Beck, J., Deller, T., and Vlachos, A. (2021b). All-trans retinoic acid induces synaptic plasticity in human cortical neurons. Elife *10*.

Leutgeb, J.K., Leutgeb, S., Moser, M.B., and Moser, E.I. (2007). Pattern separation in the dentate gyrus and CA3 of the hippocampus. Science *315*, 961-966.

Lipp, H.P., Schwegler, H., Heimrich, B., and Driscoll, P. (1988). Infrapyramidal mossy fibers and two-way avoidance learning: developmental modification of hippocampal circuitry and adult behavior of rats and mice. J Neurosci *8*, 1905-1921.

Maggio, N., and Vlachos, A. (2018). Tumor necrosis factor (TNF) modulates synaptic plasticity in a concentration-dependent manner through intracellular calcium stores. J Mol Med (Berl) *96*, 1039-1047.

Marr, D. (1971). Simple memory: a theory for archicortex. Philos Trans R Soc Lond B Biol Sci *262*, 23-81.

Maus, L., Lee, C., Altas, B., Sertel, S.M., Weyand, K., Rizzoli, S.O., Rhee, J., Brose, N., Imig, C., and Cooper, B.H. (2020). Ultrastructural Correlates of Presynaptic Functional Heterogeneity in Hippocampal Synapses. Cell Rep *30*, 3632-3643 e3638.

Mesulam, M.M., Lalehzari, N., Rahmani, F., Ohm, D., Shahidehpour, R., Kim, G., Gefen, T., Weintraub, S., Bigio, E., and Geula, C. (2019). Cortical cholinergic denervation in primary progressive aphasia with Alzheimer pathology. Neurology *92*, e1580-e1588.

Monday, H.R., Kharod, S.C., Yoon, Y.J., Singer, R.H., and Castillo, P.E. (2022). Presynaptic FMRP and local protein synthesis support structural and functional plasticity of glutamatergic axon terminals. Neuron *110*, 2588-2606 e2586.

Mootha, V.K., Lindgren, C.M., Eriksson, K.F., Subramanian, A., Sihag, S., Lehar, J., Puigserver, P., Carlsson, E., Ridderstrale, M., Laurila, E*., et al.* (2003). PGC-1alpha-responsive genes involved in oxidative phosphorylation are coordinately downregulated in human diabetes. Nat Genet *34*, 267-273.

O'Reilly, R.C., and McClelland, J.L. (1994). Hippocampal conjunctive encoding, storage, and recall: avoiding a trade-off. Hippocampus *4*, 661-682.

Orlando, M., Dvorzhak, A., Bruentgens, F., Maglione, M., Rost, B.R., Sigrist, S.J., Breustedt, J., and Schmitz, D. (2021). Recruitment of release sites underlies chemical presynaptic potentiation at hippocampal mossy fiber boutons. PLoS Biol *19*, e3001149.

Platschek, S., Cuntz, H., Vuksic, M., Deller, T., and Jedlicka, P. (2016). A general homeostatic principle following lesion induced dendritic remodeling. Acta Neuropathol Commun *4*, 19.

Prang, P., Del Turco, D., and Deller, T. (2003). Associational sprouting in the mouse fascia dentata after entorhinal lesion in vitro. Brain Res *978*, 205-212.

Prince, L.Y., Bacon, T.J., Tigaret, C.M., and Mellor, J.R. (2016). Neuromodulation of the Feedforward Dentate Gyrus-CA3 Microcircuit. Front Synaptic Neurosci *8*, 32.

Reimand, J., Isserlin, R., Voisin, V., Kucera, M., Tannus-Lopes, C., Rostamianfar, A., Wadi, L., Meyer, M., Wong, J., Xu, C*., et al.* (2019). Pathway enrichment analysis and visualization of omics data using g:Profiler, GSEA, Cytoscape and EnrichmentMap. Nat Protoc *14*, 482-517.

Robinson, J.L., Molina-Porcel, L., Corrada, M.M., Raible, K., Lee, E.B., Lee, V.M., Kawas, C.H., and Trojanowski, J.Q. (2014). Perforant path synaptic loss correlates with cognitive impairment and Alzheimer's disease in the oldest-old. Brain *137*, 2578-2587.

Shannon, P., Markiel, A., Ozier, O., Baliga, N.S., Wang, J.T., Ramage, D., Amin, N., Schwikowski, B., and Ideker, T. (2003). Cytoscape: a software environment for integrated models of biomolecular interaction networks. Genome Res *13*, 2498-2504.

Strehl, A., Galanis, C., Radic, T., Schwarzacher, S.W., Deller, T., and Vlachos, A. (2018). Dopamine Modulates Homeostatic Excitatory Synaptic Plasticity of Immature Dentate Granule Cells in Entorhino-Hippocampal Slice Cultures. Front Mol Neurosci *11*, 303.

Strehl, A., Lenz, M., Itsekson-Hayosh, Z., Becker, D., Chapman, J., Deller, T., Maggio, N., and Vlachos, A. (2014). Systemic inflammation is associated with a reduction in Synaptopodin expression in the mouse hippocampus. Exp Neurol *261*, 230-235.

Subramanian, A., Tamayo, P., Mootha, V.K., Mukherjee, S., Ebert, B.L., Gillette, M.A., Paulovich, A., Pomeroy, S.L., Golub, T.R., Lander, E.S*., et al.* (2005). Gene set enrichment analysis: a knowledge-based approach for interpreting genome-wide expression profiles. Proc Natl Acad Sci U S A *102*, 15545-15550.

Sutton, M.A., Ito, H.T., Cressy, P., Kempf, C., Woo, J.C., and Schuman, E.M. (2006). Miniature neurotransmission stabilizes synaptic function via tonic suppression of local dendritic protein synthesis. Cell *125*, 785-799.

Thal, D.R., Holzer, M., Rub, U., Waldmann, G., Gunzel, S., Zedlick, D., and Schober, R. (2000). Alzheimer-related tau-pathology in the perforant path target zone and in the hippocampal stratum oriens and radiatum correlates with onset and degree of dementia. Exp Neurol *163*, 98-110.

Turrigiano, G. (2012). Homeostatic synaptic plasticity: local and global mechanisms for stabilizing neuronal function. Cold Spring Harb Perspect Biol *4*, a005736.

Turrigiano, G.G. (2017). The dialectic of Hebb and homeostasis. Philos Trans R Soc Lond B Biol Sci *372*.

Turrigiano, G.G., Leslie, K.R., Desai, N.S., Rutherford, L.C., and Nelson, S.B. (1998). Activity-
dependent scaling of quantal amplitude in neocortical neurons. Nature *391*, 892-896.

Vandael, D., Borges-Merjane, C., Zhang, X., and Jonas, P. (2020). Short-Term Plasticity at
Hippocampal Mossy Fiber Synapses Is Induced by Natural Activity Patterns and Associated with
Vesicle Pool Engram Formation. Neuron *107*, 509-521 e507.

Vandael, D., Okamoto, Y., Borges-Merjane, C., Vargas-Barroso, V., Suter, B.A., and Jonas, P. (2021a).
Subcellular patch-clamp techniques for single-bouton stimulation and simultaneous pre- and
postsynaptic recording at cortical synapses. Nat Protoc *16*, 2947-2967.

Vandael, D., Okamoto, Y., and Jonas, P. (2021b). Transsynaptic modulation of presynaptic short-term
plasticity in hippocampal mossy fiber synapses. Nat Commun *12*, 2912.

Viana da Silva, S., Zhang, P., Haberl, M.G., Labrousse, V., Grosjean, N., Blanchet, C., Frick, A., and
Mulle, C. (2019). Hippocampal Mossy Fibers Synapses in CA3 Pyramidal Cells Are Altered at an Early
Stage in a Mouse Model of Alzheimer's Disease. J Neurosci *39*, 4193-4205.

Villanueva-Castillo, C., Tecuatl, C., Herrera-Lopez, G., and Galvan, E.J. (2017). Aging-related
impairments of hippocampal mossy fibers synapses on CA3 pyramidal cells. Neurobiol Aging *49*, 119-
137.

Vitureira, N., Letellier, M., and Goda, Y. (2012). Homeostatic synaptic plasticity: from single synapses
to neural circuits. Curr Opin Neurobiol *22*, 516-521.

Vlachos, A. (2012). Synaptopodin and the spine apparatus organelle-regulators of different forms of
synaptic plasticity? Ann Anat *194*, 317-320.

Vlachos, A., Bas Orth, C., Schneider, G., and Deller, T. (2012a). Time-lapse imaging of granule cells
in mouse entorhino-hippocampal slice cultures reveals changes in spine stability after entorhinal
denervation. J Comp Neurol *520*, 1891-1902.

Vlachos, A., Becker, D., Jedlicka, P., Winkels, R., Roeper, J., and Deller, T. (2012b). Entorhinal
denervation induces homeostatic synaptic scaling of excitatory postsynapses of dentate granule cells in
mouse organotypic slice cultures. PLoS One *7*, e32883.

Vlachos, A., Ikenberg, B., Lenz, M., Becker, D., Reifenberg, K., Bas-Orth, C., and Deller, T. (2013).
Synaptopodin regulates denervation-induced homeostatic synaptic plasticity. Proc Natl Acad Sci U S A
110, 8242-8247.

Vlachos, A., Korkotian, E., Schonfeld, E., Copanaki, E., Deller, T., and Segal, M. (2009). Synaptopodin
regulates plasticity of dendritic spines in hippocampal neurons. J Neurosci *29*, 1017-1033.

Vyleta, N.P., Borges-Merjane, C., and Jonas, P. (2016). Plasticity-dependent, full detonation at
hippocampal mossy fiber-CA3 pyramidal neuron synapses. Elife *5*.

Vyleta, N.P., and Jonas, P. (2014). Loose coupling between Ca2+ channels and release sensors at a
plastic hippocampal synapse. Science *343*, 665-670.

Weng, F.J., Garcia, R.I., Lutzu, S., Alvina, K., Zhang, Y., Dushko, M., Ku, T., Zemoura, K., Rich, D.,
Garcia-Dominguez, D.*, et al.* (2018). Npas4 Is a Critical Regulator of Learning-Induced Plasticity at
Mossy Fiber-CA3 Synapses during Contextual Memory Formation. Neuron *97*, 1137-1152 e1135.

Wingo, A.P., Dammer, E.B., Breen, M.S., Logsdon, B.A., Duong, D.M., Troncosco, J.C., Thambisetty,
M., Beach, T.G., Serrano, G.E., Reiman, E.M.*, et al.* (2019). Large-scale proteomic analysis of human
brain identifies proteins associated with cognitive trajectory in advanced age. Nat Commun *10*, 1619.

Wu, C.H., Tatavarty, V., Jean Beltran, P.M., Guerrero, A.A., Keshishian, H., Krug, K., MacMullan,
M.A., Li, L., Carr, S.A., Cottrell, J.R.*, et al.* (2022). A bidirectional switch in the Shank3
phosphorylation state biases synapses toward up- or downscaling. Elife *11*.

Yap, K., Drakew, A., Smilovic, D., Rietsche, M., Paul, M.H., Vuksic, M., Del Turco, D., and Deller, T.
(2020). The actin-modulating protein synaptopodin mediates long-term survival of dendritic spines.
Elife *9*.

Zalutsky, R.A., and Nicoll, R.A. (1990). Comparison of two forms of long-term potentiation in single
hippocampal neurons. Science *248*, 1619-1624.

FIGURES

Figure 1: Entorhinal cortex lesion causes homeostatic synaptic plasticity in both dentate granule

cells and CA3 pyramidal cells.

(A) Entorhinal cortex lesion (ECL) in organotypic entorhino-hippocampal tissue cultures was

performed by dissecting the perforant path (white line, scale bar 300 µm). Hippocampal cells were

visualized by post-hoc staining of neurons after electrophysiological assessment.

(B) Spontaneous excitatory postsynaptic currents (sEPSCs) were recorded in both dentate granule cells

and CA3 pyramidal cells in non-lesioned and lesioned cultures (3 days post lesion, 3 dp ECL).

(C-D) Spontaneous EPSC analysis in dentate granule cells reveals a lesion-induced excitatory synaptic

strengthening in the mean sEPSC amplitude, half width and area (C; $n_{control} = 22$ cells; $n_{3dp\ ECL} = 22$ cells;

Mann-Whitney test; Linear regression fit for XY-plot analysis). Percentile analysis of sEPSC

amplitudes (D) demonstrates that the excitatory synaptic strengthening is detected in the 50% largest

events (Mann-Whitney test).

(E-F) The frequency of sEPSCs is significantly increased 3 days after ECL (E; Mann-Whitney test).

Moreover, amplitude/frequency-analysis shows that the abundance of larger sEPSC amplitudes is higher

in lesioned cultures when compared to non-lesioned controls (F; RM-two-way ANOVA followed by

Sidak's multiple comparisons test).

(G-H) Spontaneous EPSC analysis in CA3 pyramidal cells reveals a lesion-induced increase in mean

sEPSC amplitude, whereas half width and area remained unchanged (G; $n_{control} = 48$ cells; $n_{3dp\ ECL} = 49$

cells; Mann-Whitney test; Linear regression fit for XY-plot analysis). Percentile analysis of sEPSC

amplitudes (H) demonstrates that the excitatory synaptic strengthening is only visible in the 50% largest

events (Mann-Whitney test).

(I-J) No lesion-induced changes in mean sEPSC frequency was detectable in CA3 pyramidal neurons.

Amplitude/frequency-analysis shows a reduced portion of smaller events in CA3 pyramidal cells 3 days

after ECL (Mann-Whitney test for column statistics and RM-two-way ANOVA followed by Sidak's

multiple comparisons test for XY-plot analysis).

Individual data points are indicated by colored dots. Values represent mean ± s.e.m (*$p < 0.05$, **$p <$

0.01, ***$p < 0.001$, ns - non significant difference).

Figure 2: Lesion-induced loss of the cortico-hippocampal projection leads to changed
transmission at hippocampal mossy fiber synapses.

(A) Post-hoc visualization of patched neurons in organotypic tissue cultures after paired-
electrophysiological recordings between individual presynaptic (pre) dentate granule cells and
postsynaptic (post) CA3 pyramidal neurons to analyze synaptic transmission features at hippocampal
mossy fiber synapses (scale bar 300 µm).

(B) Five consecutive action potentials were induced in dentate granule cells at 20 Hz (bottom trace) to
assess postsynaptic responses in CA3 pyramidal neurons in lesioned and non-lesioned control cultures
(upper traces).

(C) The response rate, i.e., the fraction of presynaptic pulses that elicited a postsynaptic response in a
connected pair of individual neurons, was significantly increased for the first two pulses in lesioned
cultures when compared to non-lesioned controls ($n_{control} = 10$; $n_{3dp\ ECL} = 10$; Mann-Whitney test for the
respective pulse numbers).

(D) In successfully elicited unitary EPSCs (uEPSCs), a significant increase in uEPSC amplitude was
detectable ($n°1_{control} = 20$ events, $n°1_{ECL} = 187$ events; $n°2_{control} = 115$ events, $n°2_{ECL} = 304$ events;
$n°3_{control} = 292$ events, $n°3_{ECL} = 394$ events (one data point outside the axis limits); $n°4_{control} = 326$ events,
$n°4_{ECL} = 385$ events; $n°5_{control} = 307$ events, $n°5_{ECL} = 384$ events; Mann-Whitney test for the respective
pulse numbers).

Individual data points are indicated by colored dots. Values represent mean ± s.e.m (*$p < 0.05$, **$p <$
0.01, ***$p < 0.001$, ns - non significant difference).

Figure 3: The spine apparatus organelle is a hub for ribosomes and excitatory synapse related mRNAs.

(A) Transmission electron microscopy (TEM) images suggest a close association of ribosomes (arrowheads) and the spine apparatus organelle. Scale bar = 300 nm.

(B) Bioanalyzer traces (picoRNA Chip) from RNA samples prepared from precleared input (left) and Synpo-coIPs (right). Traces demonstrate the co-immunoprecipitation of ribosomal RNA with anti-synaptopodin antibody. Notably, beads-only controls did not show an immunoprecipitation of rRNA.

(C) RT-qPCR analysis demonstrates a significant enrichment of Gria1 mRNA in synaptopodin-coIP compared to input samples indicating an association between the spine apparatus organelle and synaptic plasticity related mRNAs (normalized to Gapdh; $n_{input/SP-coIP}$ = 10 samples, Mann-Whitney test).

(D) Analysis of differential mRNA abundance in Synpo-coIP RNA samples compared to their respective input samples to reveal the synaptopodin related transcriptome. A log2(FC) value > 0 indicates an mRNA enrichment in Synpo-coIP samples (UP: 2355 genes). Differentially enriched genes ($p_{adj.}$ < 0.05) are indicated by orange dots (n = 5 samples input and Synpo-coIP respectively; 5-6 tissue cultures pooled from one filter per sample; DESeq2-analysis, usegalaxy.eu platform).

(E) A gene set enrichment analysis for a cellular compartment gene ontology was performed for significantly enriched genes in Synpo-coIP samples and visualized by a cytoscape map (node cutoff p = 0.05, edge cutoff (similarity): 0.8; node size represents gene set size; node positions have been manually adjusted).

(F) Transmission electron microscopy was used to visualize ultrastructural properties of mossy fiber synapses in lesioned and non-lesioned control cultures (scale bar 1 μm).

(G) Upon lesion, the abundance of spine apparatus organelles (SA) and their number of dense plates were significantly increased in the postsynaptic compartment of mossy fiber synapses ($n_{control}$ = 120 (no SA = 97 postsynapses, 1 dense plate = 22 postsynapses, 2 dense plates = 1 postsynapse, 3 dense plates = 0 postsynapses); n_{ECL} = 90 (no SA = 56 postsynapses, 1 dense plate = 21 postsynapses, 2 dense plates = 12 postsynapses, 3 dense plates = 1 postsynapses)).

(H) While the number of dense plates increased, their length significantly decreased 3 days after ECL ($n_{control}$ = 24 (one data point outside the axis limits; n_{ECL} = 48, Mann-Whitney test).

Individual data points are indicated by colored dots. Values represent mean ± s.e.m (*p < 0.05, **p < 0.01, ns - non significant difference).

Figure 4: Lesion-induced homeostatic synaptic plasticity at hippocampal mossy fiber synapses is impaired in synaptopodin-deficient mice.

(A-B) sEPSCs were recorded from dentate granule cells in synaptopodin-deficient tissue cultures (Synpo$^{-/-}$). Here, the mean sEPSC amplitude, half width and area significantly increased 3 days after the ECL (A; $n_{control}$ = 29; $n_{3dp\ ECL}$ = 26; Mann-Whitney test; Linear regression fit for XY-plot analysis). In contrast to wildtype cultures, no changes in the mean sEPSC frequency or the sEPSC amplitude/frequency-distribution were observed (B; Mann-Whitney test for column statistics and RM-two-way ANOVA followed by Sidak's multiple comparisons test for XY-plot analysis).

(C) No changes in passive membrane properties of dentate granule cells were detected in synaptopodin-deficient cultures upon lesion (Mann-Whitney test for column statistics and RM-two-way ANOVA followed by Sidak's multiple comparisons test for XY-plot analysis).

(D-E) The mean sEPSC amplitude in CA3 pyramidal neurons was not significantly different in lesioned and non-lesioned control cultures. Noteworthy, the mean sEPSC half width significantly decreased in lesioned cultures (D; $n_{control}$ = 25; $n_{3dp\ ECL}$ = 25; Mann-Whitney test; linear regression fit for XY-plot analysis). No lesion-induced changes in the mean sEPSC frequency or the amplitude/frequency-interdependence were detectable (E; Mann-Whitney test for column statistics and RM-two-way ANOVA followed by Sidak's multiple comparisons test for XY-plot analysis).

(F) No changes in passive membrane properties of CA3 pyramidal neurons were detected in synaptopodin-deficient cultures upon lesion (Mann-Whitney test for column statistics and RM-two-way ANOVA followed by Sidak's multiple comparisons test for XY-plot analysis).

(G-H) Paired whole-cell patch-clamp recordings were carried out between individual dentate granule cells (GC) and CA3 pyramidal neurons in lesioned and non-lesioned control cultures prepared from synaptopodin-deficient animals (G). Upon injection of 5 presynaptic current pulses in the presynaptic granule cells, no changes in the pulse-number-dependent response rate of connected mossy fiber synapses were detectable after the ECL (H; $n_{control}$ = 12; $n_{3dp\ ECL}$ = 10; Mann-Whitney test for the respective pulse numbers).

(I) Analyzing properties of successfully transmitted uEPSCs, a significant decrease in uEPSC amplitude was evident in lesioned cultures when compared to non-lesioned controls (n°1$_{control}$ = 144 events, n°1$_{ECL}$ = 82 events; n°2$_{control}$ = 300 events, n°2$_{ECL}$ = 263 events; n°3$_{control}$ = 426 events (one data point outside the axis limits), n°3$_{ECL}$ = 402 events; n°4$_{control}$ = 486 events, n°4$_{ECL}$ = 364 events; n°5$_{control}$ = 472 events, n°5$_{ECL}$ = 317 events; Mann-Whitney test for the respective pulse numbers).

Individual data points are indicated by colored dots. Values represent mean ± s.e.m (*p < 0.05, ***p < 0.001, ns - non significant difference).

Figure 5: Entorhinal cortex lesion leads to synaptopodin-dependent transcriptomic changes in the CA3 area of organotypic tissue cultures.

(A) Volcano Plots to visualize differential gene expression in wildtype (Synpo$^{+/+}$) and synaptopodin-deficient (Synpo$^{-/-}$) tissue preparations based on region-specific transcriptome analysis of the dentate gyrus and the CA3 area. Red dots indicate differentially expressed genes ($p_{adj.} < 0.05$; CA3-Synpo$^{+/+}$: $n_{control} = 2$, $n_{3dpECL} = 3$; DG-Synpo$^{+/+}$: $n_{control} = 4$, $n_{3dpECL} = 7$; CA3-Synpo$^{-/-}$: $n_{control} = 6$, $n_{3dpECL} = 6$; DG-Synpo$^{-/-}$: $n_{control} = 7$, $n_{3dpECL} = 4$).

(B) VENN-Diagrams illustrate the magnitude of lesion-induced differential gene expression and the overlap in commonly regulated genes between wildtype and synaptopodin-deficient tissue preparations. The illustration indicates that (1) synaptopodin-deficient preparations show deficits in lesion-driven gene expression mainly in CA3 and (2) despite the identification of commonly regulated genes, a considerable number of genotype-specific genes can be identified.

(C) For wildtype cultures (Synpo$^{+/+}$), a gene set enrichment analysis was performed and visualized by cytoscape maps and enrichment plots for both hippocampal regions. In both regions, a transcriptional inflammatory response that is accompanied by metabolic changes, can be detected.

(D) To identify differentially expressed genes that relate to synaptic transmission in wildtype cultures (Synpo$^{+/+}$), DESeq2-output was filtered for genes listed in the cellular compartment genes ontology "synapse" (GOCC: 0045202). Heatmap-visualization demonstrate that the expression of synapse related genes is a target for lesion-induced cellular remodeling.

Analysis was performed on the Galaxy platform, Differential gene expression was determined using the DESeq2 application.

Figure S1: Entorhinal cortex lesion does not influence active or passive membrane properties.

(A-B) Both resting membrane potential and input resistance (A) as well as the input-output curve for

the first 15 sweeps of current injection (B) in dentate granule cells were unaffected by the ECL ($n_{control}$

= 22 cells; $n_{3dp\ ECL}$ = 22 cells; Mann-Whitney test and RM two-way ANOVA followed by Sidak's

multiple comparisons test for XY-plot analysis).

(C) Action potential frequency upon increasing current injection in dentate granule cells is similar in

non-lesioned and lesioned cultures (RM two-way ANOVA followed by Sidak's multiple comparisons

test).

(D-E) Both resting membrane potential and input resistance (D) as well as the input-output curve for

the first 15 sweeps of current injection (E) in CA3 pyramidal neurons were unaffected by the ECL

($n_{control}$ = 48 cells; $n_{3dp\ ECL}$ = 49 cells; Mann-Whitney test and RM two-way ANOVA followed by Sidak's

multiple comparisons test for XY-plot analysis).

(F) Action potential frequency upon increasing current injection in dentate granule cells is similar in

non-lesioned and lesioned cultures (RM two-way ANOVA followed by Sidak's multiple comparisons

test).

Individual data points are indicated by colored dots. Values represent mean ± s.e.m (ns - non significant

difference).

Figure S2: Entorhinal cortex lesion does not promote major changes in synaptopodin-associated

mRNA.

Volcano plots to visualize the differential abundance of mRNAs in lesioned and non-lesioned cultures.

Both input and Synpo-coIP samples have been analyzed. Differentially expressed/enriched genes ($p_{adj.}$

< 0.05) are indicated by orange dots (n = 5 samples for both lesioned and non-lesioned cultures; 5-6

tissue cultures pooled from one filter per sample; DESeq2-analysis, usegalaxy.eu platform).

Figure S3: Synaptopodin deficiency shifts the hippocampal mossy fiber synapse to an excitable
state under baseline conditions.

(A-B) sEPSCs were recorded from CA3 pyramidal cells of both synaptopodin-deficient and wildtype
tissue preparations respectively. While the mean sEPSC amplitude remained unchanged, sEPSC area
and half width were significantly increased in synaptopodin-deficient tissue cultures (A; $n_{Synpo+/+}$ = 48;
$n_{Synpo-/-}$ = 25; Mann-Whitney test; Linear regression fit for XY-plot analysis). Additionally, the sEPSC
amplitude/frequency-distribution was significantly changed between the two groups (B; RM-two-way
ANOVA followed by Sidak's multiple comparisons test).

(C-D) When comparing the response rate of postsynaptic CA3 neurons upon presynaptic action
potential generation between paired recordings of $Synpo^{+/+}$ and $Synpo^{-/-}$ tissue preparations, no
significant increase was observed (C; $n_{Synpo+/+}$ = 10; $n_{Synpo-/-}$ = 12; Mann-Whitney test for the respective
pulse numbers). However, when analyzing the amplitudes of successfully transmitted uEPSCs at the
postsynaptic CA3 pyramidal cells, a significant increase in uEPSC amplitudes became evident in
synaptopodin-deficient tissue preparations after each of the five pulses respectively (D; $n°1_{Synpo+/+}$ = 20
events, $n°1_{Synpo-/-}$ = 144 events; $n°2_{Synpo+/+}$ = 115 events, $n°2_{Synpo-/-}$ = 300 events; $n°3_{Synpo+/+}$ = 292 events,
$n°3_{Synpo-/-}$ = 426 events (one data point outside the axis limits); $n°4_{Synpo+/+}$ = 326 events, $n°4_{Synpo-/-}$ = 486
events; $n°5_{Synpo+/+}$ = 307 events, $n°5_{Synpo-/-}$ = 472 events; Mann-Whitney test for the respective pulse
numbers).

Individual data points are indicated by colored dots. Values represent mean ± s.e.m (*$p < 0.05$, **$p <$
0.01, ***$p < 0.001$, ns - non significant difference).

SUPPLEMENTARY TABLES

Table S1: DESeq2 results table Synpo-coIP vs. input (BaseMean > 300; c.f. Fig. 3)

Table S2: Synpo-coIP enriched genes (log2FC > 0; c.f. Fig. 3)

Table S3: DESeq2 results table ECL vs. control (input; Base Mean > 300; c.f. Figure S2)

Table S4: DESeq2 results table ECL vs. control (Synpo-coIP; Base Mean > 300; c.f. Figure S2)

Table S5: DESeq2 results table ECL vs. control (Synpo$^{+/+}$, CA3; Base Mean > 300; c.f. Figure 5)

Table S6: DESeq2 results table ECL vs. control (Synpo$^{+/+}$, DG; Base Mean > 300; c.f. Figure 5)

Table S7: DESeq2 results table ECL vs. control (Synpo$^{-/-}$, CA3; Base Mean > 300; c.f. Figure 5)

Table S8: DESeq2 results table ECL vs. control (Synpo$^{-/-}$, DG; Base Mean > 300; c.f. Figure 5)

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure S1

Figure S2

CA3

A

ns

B

sEPSC area [pA*ms]:
Synpo+/+: 341 ± 24.8
Synpo-/-: 461 ± 40.1**

Paired Recordings

C

D

Figure S3