Amyloid-β$_{1\text{-}42}$ oligomers enhance mGlu$_5$R-dependent synaptic weakening via NMDAR

activation and complement C3aR/C5aR signaling

Ai Na Ng[1,*], Eric W. Salter[2,3,*], John Georgiou[2,4], Zuner A. Bortolotto[1], Graham L.

Collingridge[1,2,3,4,5]

[1]School of Physiology, Pharmacology and Neuroscience, University of Bristol,

Biomedical Sciences Building, University Walk, Bristol, BS8 1TD, United Kingdom.

[2] Lunenfeld-Tanenbaum Research Institute, Mount Sinai Hospital, Sinai Health

System, 600 University Avenue, Toronto, ON, M5G 1X5, Canada.

[3] Department of Physiology, University of Toronto, Toronto, Ontario, M5S 1A8,

Canada

[4] Tanz Centre for Research in Neurodegenerative Diseases, University of Toronto,

Krembil Discovery Tower, 60 Leonard Avenue, Toronto, ON, M5T 0S8, Canada.

[5] Correspondence: Graham L. Collingridge, collingridge@lunenfeld.ca

* These authors contributed equally

Abstract

Synaptic dysfunction, weakening, and loss of synapses are well-correlated with the

pathology of Alzheimer's disease (AD). Oligomeric amyloid beta (oAβ) is considered

a major synaptotoxic trigger for AD. Recent studies have implicated hyperactivation of

the complement cascade as the driving force for loss of synapses caused by oAβ.

However, the initial synaptic cues that trigger pathological complement activity remain

elusive. Here, we examined a form of synaptic long-term depression (LTD) mediated

by metabotropic glutamate receptors (mGluR) which is disrupted in rodent models of

AD. Exogenous application of oAβ (1-42) to mouse hippocampal slices enhanced the

magnitude of mGlu subtype 5 receptor (mGlu$_5$R)-dependent LTD. We found that the

enhanced synaptic weakening occurred via both NMDARs and complement

C3aR/C5aR signaling. Our findings reveal a mechanistic interaction between mGlu$_5$R,

NMDARs, and the complement cascade in synaptic weakening induced by oAβ, which

could represent an early trigger of synaptic loss and degeneration in AD.

Introduction

Alzheimer's disease (AD) is a progressive neurodegenerative disease characterized

by memory loss, cognitive deficits and changes to personality and behavior (Breijyeh

& Karaman, 2020). Although it is the most common cause of dementia affecting the

aging population worldwide, therapeutic options are limited. Weakening and

subsequent loss of synapses are early events in AD progression that precede the

deterioration of memory and cognitive functions (Selkoe, 2002; Koffie, Hyman, &

Spires-Jones, 2011; Li & Selkoe, 2020). The development of improved treatment

options requires a deeper understanding of how synapses are initially impaired and

how this leads to overt neuronal degeneration.

A key pathological feature of AD are plaques, which are made up primarily of insoluble

aggregates of amyloid beta (Aβ) protein derived from the amyloid precursor protein

(APP). The original amyloid cascade hypothesis posited that these Aβ plaques are the

initial trigger of AD pathology (Hardy & Higgins, 1992). However, subsequent research

found that the amount of soluble oligomers of Aβ (oAβ) better correlates with the

degree of cognitive impairment in AD patients (DeKosky & Scheff, 1990; Terry et al.,

1991). Interestingly, oAβ binds to excitatory synapses (Lacor et al., 2004; Lacor et al.,

2007; Koffie et al., 2009), leading to disruption of synaptic function. Normally, synaptic

strength is modulated through long-term potentiation (LTP) and long-term depression

(LTD), which are thought to provide the cellular and molecular basis of learning and

memory (Bliss & Collingridge, 1993). oAβ mediates a shift in the LTP/LTD balance,

favoring LTD over LTP, leading to eventual net synapse loss (Lambert et al., 1998;

Kim, Anwyl, Suh, Djamgoz, & Rowan, 2001; Hsieh et al., 2006; Shankar et al., 2007;

Shankar et al., 2008; Li et al., 2009; Jo et al., 2011; Cline, Bicca, Viola, & Klein, 2018;

Li & Selkoe, 2020; Taylor, Emptage, & Jeans, 2021). Therefore, the study of oAβ-

dependent changes to LTD provides a means to investigate the initial synaptic

changes relevant to the earliest stages of AD.

Synapse damage and loss caused by oAβ does not occur in a neuron-autonomous

manner. The complement cascade is an innate immune pathway which has emerged

as a central driver of synapse loss in AD (Wu, Dissing-Olesen, MacVicar, & Stevens,

2015; Hammond, Robinton, & Stevens, 2018). Complement cascade activation leads

to phagocytosis by microglia via C3b/CR3, as well as chemotactic and inflammatory

signaling via C3a-C3aR and C5a-C5aR axes (Ricklin, Hajishengallis, Yang, &

Lambris, 2010). Tissue from post-mortem samples of patients with AD demonstrates

activation of complement activity, including increased deposition of C1q and

proteolytic cleavage of C3 (Afagh, Cummings, Cribbs, Cotman, & Tenner, 1996;

Yasojima, Schwab, McGeer, & McGeer, 1999; Wu et al., 2019). Importantly, oAβ acts

as a potent activator of the complement cascade and oAβ-mediated synapse loss is

dependent on the complement pathway (Hong et al., 2016; Bie, Wu, Foss, & Naguib,

2019). However, the upstream synaptic signals that lead to aberrant activation of

complement and subsequent synapse loss induced by oAβ are not well understood.

LTP and LTD of glutamatergic synapses is mediated by activation of N-methyl-D-

aspartate receptors (NMDARs) and/or metabotropic glutamate receptors (mGluRs:

Collingridge, Peineau, Howland, & Wang, 2010; Nicoletti et al., 2011; Volianskis et al.,

2015). LTD, induced either by NMDARs or mGluRs, can trigger subsequent synapse

elimination (Shinoda, Tanaka, Tominaga-Yoshino, & Ogura, 2010; Wiegert & Oertner,

2013; Hasegawa, Sakuragi, Tominaga-Yoshino, & Ogura, 2015; Wiegert, Pulin, Gee,

& Oertner, 2018). Both mGluR- and NMDAR-dependent signaling are affected by

oAβ, leading to synaptic plasticity impairments and aberrant synapse loss (Hsieh et

al., 2006; Shankar et al., 2007; Shankar et al., 2008; Jo et al., 2011; Cavallucci et al.,

2013; Beraldo et al., 2016; Jackson et al., 2019; Abd-Elrahman et al., 2020; Li &

Selkoe, 2020).

Synaptic plasticity involving the mGluR subtype 5 (mGlu$_5$R) is particularly pertinent to

the understanding of AD for multiple reasons. For example, mGlu$_5$R was been found

to be the only co-receptor of cellular prion protein (PrP$_c$) necessary for oAβ to activate

intracellular signaling in neurons (Um et al., 2013). Further, both knockout of mGlu$_5$R

and pharmcological antagonism is protective in oAβ-based and genetic models of AD

(Hamilton, Esseltine, DeVries, Cregan, & Ferguson, 2014; Hu et al., 2014; Hamilton

et al., 2016; Abd-Elrahman et al., 2020). Interestingly, the complement cascade has

recently been implicated in oAβ-dependent synapse loss mediated by mGluR

activation (Bie et al., 2019). A silent allosteric modulator (SAM) of mGlu$_5$R has also

been found to restore synapse density through reduced tagging by C1q and microglia

engulfment in a genetic mouse model of AD (Spurrier et al., 2022).

Previously, we identified a simple method to study $mGlu_5R$-mediated LTD, via the brief

application, S-3,5-dihydroxyphenylglycine (DHPG) which causes lasting depression of

AMPAR-mediated synaptic transmission (Palmer, Irving, Seabrook, Jane, &

Collingridge, 1997). This DHPG-induced LTD (DHPG-LTD) has since been used

extensively to uncover many aspects of mGluR physiological and pathological

function, including the regulation of mGluR function in AD mouse models (Hsieh et al.,

2006; Shankar et al., 2008; Li et al., 2009; Cavallucci et al., 2013; Mango & Nisticò,

2018; Yang, Zhou, Ryazanov, & Ma, 2021). Here, we have used DHPG-LTD to

investigate how the complement cascade may be rapidly recruited by mGluR

activation. We have applied oAβ as a standard way to induce synaptotoxicity that is

relevant to AD pathology. We found that oAβ enhanced $mGlu_5R$-mediated LTD via a

mechanism that involves both the activation of NMDARs and the C3aR and C5aR

arms of the complement cascade. These findings reveal a signaling axis between

glutamate receptors and the complement cascade which triggers early synaptic

dysfunction that may underlie the initial stages of dementia.

Materials and Methods

Animals

Adult C57BL/6JOlaHsd male mice (10-16 weeks old) from Envigo (Bicester, U.K.) and

a $mGlu_5R$ knockout line (Lu et al., 1997), backcrossed more than 10 generations to a

C57BL/6J background, were used in this study. We chose to use male mice due to the

recently reported sex specificity of oAβ-mGlu$_5$R binding which is present only in males (Abd-Elrahman et al., 2020). Animals were housed four per cage in a room with designated 12 h light/dark cycle (lights on at 08:15/ off at 20:15) with temperature at 21 °C ± 2 °C and access to food and water *ad libitum.* All procedures were performed in accordance with the U.K. Animals (Scientific Procedures) Act of 1986. Genomic DNA from mGlu$_5$R knockout litters were isolated from tail tips with DNeasy blood and tissue kit (Qiagen; Hilden, Germany; cat. no. 69504) and amplified as described (Lu et al., 1997). The amplified products were resolved and visualized in ethidium bromide-stained agarose gel.

Oligomeric amyloid-β$_{1-42}$ (oAβ) preparation

oAβ was prepared as described (Whitcomb et al., 2015). Briefly, Aβ peptide$_{(1-42)}$ (MilliporeSigma; Burlington, Massachusetts; TFA recombinant human, ultra-pure; cat. no. AG912) was dissolved in 100 % 1,1,1,3,3,3-hexafluoro-2-propanol (HFIP; MilliporeSigma; cat. no. 105228) to a final concentration of 1 mg/mL. The HFIP/peptide mixture was incubated at room temperature for 1 h and sonicated for 10 min in a water-bath sonicator. The mixture was then air-dried under a gentle stream of nitrogen gas for 1.5 h. The peptide crystal was re-suspended in 100 % DMSO, aliquoted to 10 µL per tube and stored at -80 °C until experimental use (see next section).

Hippocampal slice preparation

Mice were deeply anaesthetized via inhalation with a mixture of 5 % isofluorane / 95 % oxygen and euthanized by cervical dislocation. The brain was rapidly removed (<1 min) and chilled in cold ACSF solution (containing, in mM: 124 NaCl, 3 KCl, 26

NaHCO$_3$, 1.25 NaH$_2$PO$_4$, 10 D-glucose, 2 MgSO$_4$ and 2 CaCl$_2$; bubbled continuously

with 95% O$_2$ / 5% CO$_2$). Hippocampi were isolated from both hemispheres and

transverse slices (400 µm thickness) were prepared from the dorsal ends of both

hippocampi using a McIlwain tissue chopper (Ted Pella; Redding, California). The CA3

region was left intact for LTP but removed via a surgical cut in LTD experiments. Slices

were transferred to a holding chamber with room temperature ACSF and allowed to

recover for at least 1 h. On the day of experiment, the Aβ peptide/DMSO mixture was

diluted to 100 µM in D-PBS, vortexed briefly and allowed to aggregate for 2 h at room

temperature to form oAβ. The oAβ stock was then diluted to a final concentration of

500 nM in oxygenated ACSF and acutely applied to hippocampal slices for 2 h at room

temperature. As an inclusion criterion to account for batch-to-batch variability, we

routinely assessed the bioactivity of our oAβ preparation by confirming the ability of

each oAβ batch to impair hippocampal LTP (Figure 1A,B).

Electrophysiology

After vehicle or oAβ incubation, a dorsal hippocampal slice was submerged in a

Slicemate recording chamber (Scientifica; Uckfield, U.K.) with ACSF perfusing at a

rate of 2.5 mL/min and temperature maintained at 30.0 ± 0.5 °C. Extracellular field

potentials were recorded from the stratum radiatum area of CA1 using a glass

microelectrode (~2-3 MΩ) filled with ACSF. Responses were evoked by stimulating

the Schaffer collaterals using a platinum-iridium bipolar electrode (FHC; Bowdoin,

Maine) positioned at the border between area CA2/CA1. Stimuli of 100 µs in duration

were delivered once every 15 s via a DS2 constant voltage isolator (Digitimer;

Hertfordshire, U.K.). Responses were amplified using a Multiclamp 700B amplifier

(Molecular Devices; San Jose, California), digitized at 40 kHz and monitored in real time using WinLTP software (WinLTP Ltd.: Bristol, U.K: Anderson & Collingridge, 2007). After a period of stable baseline recordings, datapoints for input/output (I/O) plots were obtained with fixed stimulus intensity at 1x, 1.5x, 2x, 3x, and 4.5x the threshold for evoking a visually detectable field excitatory postsynaptic potential (fEPSP) (≤ 0.05 mV). For plasticity experiments a stimulus intensity of 3x the threshold intensity was employed. LTP was induced by a theta-burst stimulation (TBS) protocol, with four stimuli delivered at 100 Hz, repeated 10 times at a frequency of 5 Hz. LTD was chemically induced by bath-application of (S)-3,5-DHPG (50 µM; Abcam; Cambridge, U.K.; cat. no. ab120007) for 10 min. For offline analysis, responses were low-pass filtered at 2 kHz and the mean of four consecutive responses was quantified.

Quantification and statistical analysis

Data in the time course plots were normalized to the mean of the entire baseline period (defined as 100 %) and presented as mean ± standard error of mean (SEM). The magnitude of LTP or LTD was quantified and presented as the percentage change of the fEPSP slope within the last 5 min of the recording *versus* the baseline period. Fiber volleys (FVs) and fEPSPs were quantified using peak amplitude and initial slope measurements, respectively, using WinLTP. The I/O plots were analysed using Microsoft Excel linear regression software. The respective slope values from a fitted line between FV *versus* stimulation intensity and fEPSP slope *versus* FV were used for quantification. Statistical comparisons between -oAβ and +oAβ conditions were assessed using an unpaired two-tailed Student's t-test. The level of significance was set at $p < 0.05$.

Results

Acute oAβ exposure enhances mGlu$_5$R-dependent LTD

To investigate the impact of oAβ on synaptic function, we incubated acute hippocampus slices for 2 h with oAβ (500 nM). We subsequently measured synaptic plasticity with field potential recordings at Schaffer collateral-CA1 synapses. As a positive control for the bioactivity of our oAβ preparation, we confirmed previous findings (Shankar et al., 2008; Li et al., 2011) that oAβ was able to impair LTP in slices pretreated with oAβ (+oAβ) *versus* interleaved non-treated slices (-oAβ; Figure 1A,B). Although LTP was impaired by acute application of oAβ, basal synaptic transmission was unaltered, indicated by the overlapping input-output relationship of +oAβ compared to -oAβ slices (Figure S1). As such, this paradigm allowed us to study early disruptions to plasticity that precedes the loss of synapses.

To investigate how oAβ alters mGluR-dependent synaptic plasticity mechanisms, we induced LTD using DHPG, a selective agonist for group I mGluRs (mGlu$_1$R and mGlu$_5$R), that is widely used to study mGluR function. A short application of DHPG (10 min) induced a stable LTD (DHPG-LTD) in -oAβ slices from wild-type mice (Figure 1C,D). The magnitude of DHPG-LTD was significantly enhanced in the interleaved +oAβ slices (Figure 1C,D), consistent with previous observations (Hsieh et al., 2006; Shankar et al., 2008; Li et al., 2009; Cavallucci et al., 2013; Mango & Nisticò, 2018; Yang et al., 2021). To determine the group I mGluR subtype-specificity of our observed phenotype, we used hippocampal slices from mGlu$_5$R$^{-/-}$ mice (Lu et al., 1997) which were littermates to the mGlu$_5$R$^{+/+}$ mice in Figure 1C,D. DHPG-LTD was absent in both -oAβ and +oAβ slices (Figure 1E,F). This indicates that the form of LTD which is

enhanced by oAβ requires the activation of mGlu$_5$ receptors (referred to hereafter as

mGlu$_5$R-LTD). That is, both the LTD under control conditions and the additional LTD

observed in the presence of oAβ are both fully dependent on the activation of

mGlu$_5$Rs.

NMDARs are required for oAβ-mediated enhancement of mGlu$_5$R-LTD

There are two distinct forms of DHPG-LTD that can be distinguished by their

dependence upon the activation of NMDARs (Palmer et al., 1997; Sanderson et al.,

2022). To determine whether NMDARs are required for mGlu$_5$R-LTD either under

control conditions or the enhanced mGlu$_5$R-LTD observed in the presence of oAβ, we

used a selective NMDAR antagonist, L689,560 (5 µM). L689,560 was used as it acts

at the glycine site of the NMDAR and so is independent of L-glutamate concentration,

which might be altered by oAβ treatment (Li et al., 2009). In these experiments, we

interleaved a new set of controls and again observed the potentiation of mGlu$_5$R-LTD

by oAβ (Figure 2A,B). We found that NMDAR activation was not required for mGlu$_5$R-

LTD in control (-oAβ) conditions. However, the enhancement of mGlu$_5$R-LTD by oAβ

was completely prevented by L689,560 (Figure 2 C,D) and therefore requires NMDAR

signaling.

oAβ enhances mGlu$_5$R-LTD via complement cascade signaling

Early studies of the effects of oAβ on synaptic plasticity found that inhibition of LTP

was prevented by application of minocycline, an inhibitor of microglia activation (Wang,

Rowan, & Anwyl, 2004). Further, it has recently been found that agonism of group I

mGluRs *in vivo* activates the complement cascade leading to synapse loss (Bie et al.,

2019). We therefore hypothesized that complement signaling is necessary for the

enhancement of mGlu$_5$R-LTD by oAβ. We antagonized either C3aR or C5aR, as both

receptors have been implicated in the pathology of genetic mouse models of AD

(Fonseca et al., 2009; Lian et al., 2016; Hernandez et al., 2017; Litvinchuk et al., 2018;

Carvalho et al., 2022; Gomez-Arboledas et al., 2022). For these experiments we also

interleaved a new set of controls (Figure 3A,B) and again observed the potentiation of

mGlu$_5$R-LTD by oAβ. To inhibit C5aR, we bath-applied PMX205 (0.3 µM), a peptide-

based antagonist (March et al., 2004). PMX205 had no effect on the level of mGlu$_5$R-

LTD induced under control conditions but eliminated the enhancement of LTD induced

by oAβ (Figure 3C,D). We next used W54011, a non-peptide based C5aR antagonist

(Sumichika et al., 2002). As with PMX205, bath-application of W54011 (0.3 µM) had

no effect on the level of mGlu$_5$R-LTD induced under control conditions but eliminated

the potentiation induced by oAβ (Figure 3E,F). Lastly, we investigated the role of C3aR

using the antagonist SB290157 (Ames et al., 2001). As with the C5aR antagonists,

SB290157 (1 µM) abolished the enhanced mGlu$_5$R-LTD induced by oAβ without

affecting the level of mGlu$_5$R-LTD under control conditions (Figure 3G,H). Together,

these data indicate that the oAβ-mediated enhancement of mGlu$_5$R-LTD occurs via

activation of complement C3aR/C5aR signaling.

Discussion

260 The complement cascade is emerging as a key innate immune pathway for shaping

261 synapses during brain development, as well as driving synapse loss in various disease

262 states. However, the synaptic signals that lead to complement activation remain poorly

263 understood. In the present study, we uncover a mechanistic link between innate

immune signaling and the amplification of glutamatergic synapse depression induced by oAβ. We found that mGlu$_5$R-dependent LTD induced by DHPG application was enhanced in slices treated with oAβ (Figure 1). Unlike vehicle-treated slices, the oAβ-mediated enhanced portion of mGlu$_5$R-LTD was dependent on NMDAR signaling (Figure 2). Similarly, antagonism of either C3aR or C5aR prevented oAβ from enhancing mGlu$_5$R -LTD (Figure 3), without affecting the underlying mGlu$_5$R-LTD *per se*. Given that basal synaptic transmission was unaltered by oAβ (Figure S1) our findings reveal a signaling axis between mGluRs, NMDARs, and the complement cascade which mediates early synaptic plasticity alterations, upstream of overt synapse loss.

oAβ-induced glutamatergic synapse dysfunction

Although AD is an extremely slow progressing disease, it is likely that at the level of a single synapse there is acute damage, which in many cases may be orchestrated by toxic oAβ species. This would presumably occur when the processes that ordinarily prevent the accumulation of oAβ are sufficiently compromised such that the local concentration increases to a level that can impair synaptic function and structure. Therefore, the transient application of oAβ has been widely employed to model this early, critical stage of the disease. Consistent with the validity of this model, mechanisms that are engaged by the acute application of oAβ to impact synaptic plasticity, such as the involvement of GSK-3, caspase-3 (Jo et al., 2011), tau (Shipton et al., 2011; Taylor et al., 2021) and microglia (Wang et al., 2004) are all directly relevant to human AD pathology.

Multiple studies have found that oAβ causes a shift in the LTP-LTD balance, favoring

LTD (Cullen, Suh, Anwyl, & Rowan, 1997; Walsh et al., 2002; Hsieh et al., 2006;

Shankar et al., 2007; Shankar et al., 2008; Li et al., 2009; Jo et al., 2011; Li et al.,

2011; Cavallucci et al., 2013). Altered activity of both group I mGluRs and NMDARs

by oAβ is central to this glutamatergic synaptic failure. mGlu$_5$R was identified as a co-

receptor of cellular prion protein (PrP$_c$) for binding oAβ and is necessary for oAβ-

induced intracellular signal transduction and synapse loss (Um et al., 2013; Haas,

Kostylev, & Strittmatter, 2014; Beraldo et al., 2016; Brody & Strittmatter, 2018).

Additionally, oAβ binds to NMDARs (De Felice et al., 2007; Decker et al., 2010) and

impairs NMDAR-dependent LTP while enhancing NMDAR-dependent LTD (Cullen et

al., 1997; Walsh et al., 2002; Wang et al., 2004; Hsieh et al., 2006; Shankar et al.,

2007; Shankar et al., 2008; Jo et al., 2011; O'Riordan, Hu, & Rowan, 2018; Abd-

Elrahman et al., 2020; Taylor et al., 2021). Despite these important advances, whether

NMDARs and mGluRs are synergistically involved in oAβ-induced synaptic weakening

and loss is poorly understood. Of relevance to the present study, at least two distinct

forms of DHPG-LTD exist, which can be distinguished by the dependence on

NMDARs (Palmer et al., 1997; Sanderson et al., 2022). In our current study, DHPG-

LTD induction under control conditions was not affected by NMDAR antagonism. In

contrast the oAβ-enhanced portion of the LTD fully depended on the activation of

NMDARs. As such, promotion of mGluR-NMDAR signaling during synaptic depression

may be a key aspect of the shift in LTP/LTD balance caused by oAβ.

Both mGlu$_5$R and NMDARs are also implicated in AD pathogenesis in genetic mouse

models and humans. Antagonists of mGlu$_5$R and genetic deletion have both been

found to reduce synaptic and behavioral deficits in AD mouse models (Hamilton et al.,

2014; Hamilton et al., 2016; Haas et al., 2017; Abd-Elrahman et al., 2020; Spurrier et

al., 2022). Indeed, the mGlu$_5$R SAM BMS-984923 (Haas et al., 2017), is currently in

phase I clinical trials. Evidence that altered NMDAR signaling is causally related to the

human condition is the use of memantine, a non-competitive NMDAR antagonist, for

the treatment of AD. Memantine slows the progression of dementia and its underlying

therapeutic mechanism has been attributed to the normalization of synaptic plasticity

(Parsons, Stöffler, & Danysz, 2007; Wang & Reddy, 2017; Liu, Chang, Song, Li, & Wu,

2019). Our experiments reveal one way in which NMDARs and mGlu$_5$Rs may act

synergistically to control synaptic integrity.

C3aR/C5aR signaling in AD

In both post-mortem human samples as well as numerous AD mouse models, many

components of the complement cascade have been found to be upregulated

(Hammond et al., 2018; Schartz & Tenner, 2020). Importantly, pharmacological and

genetic manipulations of C3aR and C5aR have provided evidence that both receptors

are necessary for synaptic dysfunction and cognitive impairments in AD genetic

mouse models. In particular, the C3aR and C5aR antagonists used in the current

study, SB290157 and PMX205, respectively, have each been found to reduce

pathology in genetic mouse models of AD (Fonseca et al., 2009; Lian et al., 2016;

Gomez-Arboledas et al., 2022). Furthermore, synapse engulfment by microglia and

neuron loss are reduced in C3aR$^{-/-}$ mice in the PS19 tauopathy AD model (Litvinchuk

et al., 2018) and C5aR1 genetic ablation is protective in AD mouse models (Hernandez

et al., 2017; Carvalho et al., 2022). However, the role of oAβ in aberrant C3aR/C5aR

signaling, and the intersection with glutamatergic synaptic signaling to cause this

dysfunction was previously not understood. Our study provides insight into this communication, by demonstrating that oAβ enhances synaptic weakening, via NMDAR and mGlu$_5$R synergistic mechanisms that require both C3aRs and C5aRs.

One important question to be addressed by future studies is which arm(s) of the complement cascade initiates the cleavage of C3 following mGlu$_5$R-LTD in the presence of oAβ? Proteolytic cleavage of C3 generates both C3a, as well as components of the C5 convertase which produces C5a. C3 conversion can be driven by either the classical arm, initiated by C1q, the lectin arm via mannose-binding lectin (MBL) or the alternative pathway through C3 "tickover" (Ricklin et al., 2010). Hong et al. (2016) found that the ability of oAβ to inhibit LTP requires the classical complement pathway, as co-incubation of hippocampal slices with oAβ and a function-blocking antibody of C1q restored LTP. Furthermore, *in vivo* injections of oAβ into rat hippocampus upregulated C1q, and increased synapse phagocytosis by microglia (Bie et al., 2019). C1q upregulation by oAβ and subsequent synapse phagocytosis required group I mGluR signaling, and *in vivo* DHPG administration alone increased C1q. Finally, a recent study found that a SAM of mGlu$_5$R reduced C1q co-localization with PSD-95 and microglia engulfment of synapses in a mouse model of AD (Spurrier et al., 2022). As such, our findings suggest that C3aR/C5aR activation links the induction of the classical arm of the complement cascade, via mGluRs and NMDARs, to downstream synaptic depression.

Cleavage of C3 also generates the C3b fragment, which can covalently attach to synapses where it is converted to iC3b and binds to the phagocytic receptor CR3

(Ricklin et al., 2010). Genetic deletion of CR3 has also been found to spare synapses

following oAβ administration *in vivo* (Hong et al., 2016), indicating that phagocytosis

through iC3b-CR3 binding is also critical for the deleterious effects of complement in

pathology. As weaker synapses are targeted by complement for phagocytic removal

(Schafer et al., 2012), an intriguing question is whether synapse weakening by

C3aR/C5aR activation also tags those synapses with iC3b for subsequent engulfment

by microglia. CR3 can also impact synapses by non-phagocytic mechanisms in

neuroinflammatory conditions. LPS and hypoxia activates CR3 which induces LTD of

AMPAR transmission in the hippocampus (Zhang et al., 2014). Thus, microglia and

the complement cascade have a multi-faceted role in synaptic dysfunction in disease,

being critical for both changes to AMPAR-mediated transmission strength as well as

structural synapse elimination.

Relevance to other neurodegenerative diseases

Upregulation of complement cascade proteins and activity is not exclusive to AD and

has been detected in several other neurodegenerative diseases. For example, A1

reactive astrocytes, which express high levels of C3, have been identified in human

post-mortem tissue from patients with amyotrophic lateral sclerosis, Huntington's

disease, Parkinson's disease, and multiple sclerosis (Liddelow et al., 2017). Further,

studies from *in vitro* and mouse models of these diseases have demonstrated the

necessity of C3 and/or C5 for their pathology (Woodruff et al., 2008; McGeer, Lee, &

McGeer, 2017; Wang, Lee, Lee, Woodruff, & Noakes, 2017; Hammond et al., 2020;

Bourel et al., 2021; Chen, Hu, Ding, Du, & Hu, 2021; Gregersen et al., 2021; Lopez-

Sanchez et al., 2022). Interestingly, mGlu$_5$R has separately been identified as an

emerging therapeutic target in these complement-associated neurodegenerative

diseases (Geurts et al., 2003; Abd-Elrahman et al., 2017; Bonifacino et al., 2017; Doria

et al., 2018; Bonifacino et al., 2019; Farmer et al., 2020; Zhang et al., 2021; Azam et

al., 2022; Li, Colson, Abd-Elrahman, & Ferguson, 2022; Su, Wang, Han, & Shen,

2022). Therefore, the pathological mGlu$_5$R-NMDAR-C3aR/C5aR signaling axis

identified in our study may have relevance for multiple brain disorders.

Concluding Remarks

mGluRs, NMDARs, and the complement cascade have been independently studied

for their role in mediating the deleterious effects of oAβ. Our study has found a direct

link which places glutamate receptor-innate immune interactions at the center of early

synaptic dysfunction. As oAβ is thought to be a key trigger in AD, a deeper

understanding of the cellular and molecular mechanisms of these interactions has the

potential to elucidate therapeutic targets which halt AD progression.

Acknowledgements

This work was supported by a CIHR (Canadian Institutes of Health Research) Foundation Grant to GLC (#154276). GLC is the holder of the Krembil Family Chair in Alzheimer's Research. We thank the trainee support provided by the C.R. Younger Foundation. The work was supported by the MS Society of Canada as well as gifts from Jon and Nancy Love and the Dani Reiss Family Foundation (Neurodegeneration and Aging Research Program). We are grateful for the long-standing support of the Amalgamated Transit Union (ATU) Local 113 (Toronto, Canada). EWS was supported by a CIHR Doctoral Award (Frederick Banting and Charles Best Canada Graduate Scholarship, CGS-D) and an Ontario Graduate Scholarship (OGS).

Competing interests

The authors have no competing interests to declare.

References

Abd-Elrahman, K. S., Albaker, A., de Souza, J. M., Ribeiro, F. M., Schlossmacher, M.

G., Tiberi, M., Hamilton, A., & Ferguson, S. S. G. (2020). Abeta oligomers

induce pathophysiological mGluR5 signaling in Alzheimer's disease model

mice in a sex-selective manner. *Sci Signal, 13*(662).

doi:10.1126/scisignal.abd2494

Abd-Elrahman, K. S., Hamilton, A., Hutchinson, S. R., Liu, F., Russell, R. C., &

Ferguson, S. S. G. (2017). mGluR5 antagonism increases autophagy and

prevents disease progression in the zQ175 mouse model of Huntington's

disease. *Sci Signal, 10*(510). doi:10.1126/scisignal.aan6387

Afagh, A., Cummings, B. J., Cribbs, D. H., Cotman, C. W., & Tenner, A. J. (1996).

Localization and cell association of C1q in Alzheimer's disease brain. *Exp*

Neurol, 138(1), 22-32. doi:10.1006/exnr.1996.0043

Ames, R. S., Lee, D., Foley, J. J., Jurewicz, A. J., Tornetta, M. A., Bautsch, W.,

Settmacher, B., Klos, A., Erhard, K. F., Cousins, R. D., Sulpizio, A. C., Hieble,

J. P., McCafferty, G., Ward, K. W., Adams, J. L., Bondinell, W. E., Underwood,

D. C., Osborn, R. R., . . . Sarau, H. M. (2001). Identification of a selective

nonpeptide antagonist of the anaphylatoxin C3a receptor that demonstrates

antiinflammatory activity in animal models. *J Immunol, 166*(10), 6341-6348.

doi:10.4049/jimmunol.166.10.6341

Anderson, W. W., & Collingridge, G. L. (2007). Capabilities of the WinLTP data

acquisition program extending beyond basic LTP experimental functions. *J*

Neurosci Methods, 162(1-2), 346-356. doi:10.1016/j.jneumeth.2006.12.018

Azam, S., Jakaria, M., Kim, J., Ahn, J., Kim, I. S., & Choi, D. K. (2022). Group I mGluRs
in Therapy and Diagnosis of Parkinson's Disease: Focus on mGluR5 Subtype.
Biomedicines, 10(4). doi:10.3390/biomedicines10040864

Beraldo, F. H., Ostapchenko, V. G., Caetano, F. A., Guimaraes, A. L., Ferretti, G. D.,
Daude, N., Bertram, L., Nogueira, K. O., Silva, J. L., Westaway, D., Cashman,
N. R., Martins, V. R., Prado, V. F., & Prado, M. A. (2016). Regulation of Amyloid
beta Oligomer Binding to Neurons and Neurotoxicity by the Prion Protein-
mGluR5   Complex.   *J   Biol   Chem,   291*(42),   21945-21955.
doi:10.1074/jbc.M116.738286

Bie, B., Wu, J., Foss, J. F., & Naguib, M. (2019). Activation of mGluR1 Mediates C1q-
Dependent Microglial Phagocytosis of Glutamatergic Synapses in Alzheimer's
Rodent Models. *Mol Neurobiol, 56*(8), 5568-5585. doi:10.1007/s12035-019-
1467-8

Bliss, T. V., & Collingridge, G. L. (1993). A synaptic model of memory: long-term
potentiation   in   the   hippocampus.   *Nature,   361*(6407),   31-39.
doi:10.1038/361031a0

Bonifacino, T., Cattaneo, L., Gallia, E., Puliti, A., Melone, M., Provenzano, F., Bossi,
S., Musante, I., Usai, C., Conti, F., Bonanno, G., & Milanese, M. (2017). In-vivo
effects of knocking-down metabotropic glutamate receptor 5 in the
SOD1(G93A) mouse model of amyotrophic lateral sclerosis.
Neuropharmacology, 123, 433-445. doi:10.1016/j.neuropharm.2017.06.020

Bonifacino, T., Provenzano, F., Gallia, E., Ravera, S., Torazza, C., Bossi, S.,
Ferrando, S., Puliti, A., Van Den Bosch, L., Bonanno, G., & Milanese, M. (2019).
In-vivo genetic ablation of metabotropic glutamate receptor type 5 slows down

disease progression in the SOD1(G93A) mouse model of amyotrophic lateral

sclerosis. *Neurobiol Dis, 129*, 79-92. doi:10.1016/j.nbd.2019.05.007

Bourel, J., Planche, V., Dubourdieu, N., Oliveira, A., Sere, A., Ducourneau, E. G.,

Tible, M., Maitre, M., Leste-Lasserre, T., Nadjar, A., Desmedt, A., Ciofi, P.,

Oliet, S. H., Panatier, A., & Tourdias, T. (2021). Complement C3 mediates early

hippocampal neurodegeneration and memory impairment in experimental

multiple sclerosis. *Neurobiol Dis, 160*, 105533. doi:10.1016/j.nbd.2021.105533

Breijyeh, Z., & Karaman, R. (2020). Comprehensive Review on Alzheimer's Disease:

Causes and Treatment. *Molecules, 25*(24). doi:10.3390/molecules25245789

Brody, A. H., & Strittmatter, S. M. (2018). Synaptotoxic Signaling by Amyloid Beta

Oligomers in Alzheimer's Disease Through Prion Protein and mGluR5. *Adv*

Pharmacol, 82, 293-323. doi:10.1016/bs.apha.2017.09.007

Carvalho, K., Schartz, N. D., Balderrama-Gutierrez, G., Liang, H. Y., Chu, S. H.,

Selvan, P., Gomez-Arboledas, A., Petrisko, T. J., Fonseca, M. I., Mortazavi, A.,

& Tenner, A. J. (2022). Modulation of C5a-C5aR1 signaling alters the dynamics

of AD progression. *J Neuroinflammation, 19*(1), 178. doi:10.1186/s12974-022-

02539-2

Cavallucci, V., Berretta, N., Nobili, A., Nisticò, R., Mercuri, N. B., & D'Amelio, M.

(2013). Calcineurin inhibition rescues early synaptic plasticity deficits in a

mouse model of Alzheimer's disease. *Neuromolecular Med, 15*(3), 541-548.

doi:10.1007/s12017-013-8241-2

Chen, M. M., Hu, Z. L., Ding, J. H., Du, R. H., & Hu, G. (2021). Astrocytic Kir6.1 deletion

aggravates neurodegeneration in the lipopolysaccharide-induced mouse model

of Parkinson's disease via astrocyte-neuron cross talk through complement C3-

C3R   signaling. *Brain   Behav   Immun,   95*,   310-320.
doi:10.1016/j.bbi.2021.04.003

Cline, E. N., Bicca, M. A., Viola, K. L., & Klein, W. L. (2018). The Amyloid-β Oligomer
Hypothesis: Beginning of the Third Decade. *J Alzheimers Dis, 64*(s1), S567-
s610. doi:10.3233/jad-179941

Collingridge, G. L., Peineau, S., Howland, J. G., & Wang, Y. T. (2010). Long-term
depression in the CNS. *Nat Rev Neurosci, 11*(7), 459-473.
doi:10.1038/nrn2867

Cullen, W. K., Suh, Y. H., Anwyl, R., & Rowan, M. J. (1997). Block of LTP in rat
hippocampus in vivo by beta-amyloid precursor protein fragments. *Neuroreport,*
8(15), 3213-3217. doi:10.1097/00001756-199710200-00006

De Felice, F. G., Velasco, P. T., Lambert, M. P., Viola, K., Fernandez, S. J., Ferreira,
S. T., & Klein, W. L. (2007). Abeta oligomers induce neuronal oxidative stress
through an N-methyl-D-aspartate receptor-dependent mechanism that is
blocked by the Alzheimer drug memantine. *J Biol Chem, 282*(15), 11590-
11601. doi:10.1074/jbc.M607483200

Decker, H., Jürgensen, S., Adrover, M. F., Brito-Moreira, J., Bomfim, T. R., Klein, W.
L., Epstein, A. L., De Felice, F. G., Jerusalinsky, D., & Ferreira, S. T. (2010). N-
methyl-D-aspartate receptors are required for synaptic targeting of Alzheimer's
toxic amyloid-β peptide oligomers. *J Neurochem, 115*(6), 1520-1529.
doi:10.1111/j.1471-4159.2010.07058.x

DeKosky, S. T., & Scheff, S. W. (1990). Synapse loss in frontal cortex biopsies in
Alzheimer's disease: correlation with cognitive severity. *Ann Neurol, 27*(5), 457-
464. doi:10.1002/ana.410270502

Doria, J. G., de Souza, J. M., Silva, F. R., Olmo, I. G., Carvalho, T. G., Alves-Silva, J.,

Ferreira-Vieira, T. H., Santos, J. T., Xavier, C. Q. S., Silva, N. C., Maciel, E. M.

A., Conn, P. J., & Ribeiro, F. M. (2018). The mGluR5 positive allosteric

modulator VU0409551 improves synaptic plasticity and memory of a mouse

model of Huntington's disease. *J Neurochem, 147*(2), 222-239.

doi:10.1111/jnc.14555

Farmer, K., Abd-Elrahman, K. S., Derksen, A., Rowe, E. M., Thompson, A. M., Rudyk,

C. A., Prowse, N. A., Dwyer, Z., Bureau, S. C., Fortin, T., Ferguson, S. S. G., &

Hayley, S. (2020). mGluR5 Allosteric Modulation Promotes Neurorecovery in a

6-OHDA-Toxicant Model of Parkinson's Disease. *Mol Neurobiol, 57*(3), 1418-

1431. doi:10.1007/s12035-019-01818-z

Fonseca, M. I., Ager, R. R., Chu, S. H., Yazan, O., Sanderson, S. D., LaFerla, F. M.,

Taylor, S. M., Woodruff, T. M., & Tenner, A. J. (2009). Treatment with a C5aR

antagonist decreases pathology and enhances behavioral performance in

murine models of Alzheimer's disease. *J Immunol, 183*(2), 1375-1383.

doi:10.4049/jimmunol.0901005

Geurts, J. J., Wolswijk, G., Bö, L., van der Valk, P., Polman, C. H., Troost, D., &

Aronica, E. (2003). Altered expression patterns of group I and II metabotropic

glutamate receptors in multiple sclerosis. *Brain, 126*(Pt 8), 1755-1766.

doi:10.1093/brain/awg179

Gomez-Arboledas, A., Carvalho, K., Balderrama-Gutierrez, G., Chu, S. H., Liang, H.

Y., Schartz, N. D., Selvan, P., Petrisko, T. J., Pan, M. A., Mortazavi, A., &

Tenner, A. J. (2022). C5aR1 antagonism alters microglial polarization and

mitigates disease progression in a mouse model of Alzheimer's disease. *Acta*

Neuropathol Commun, 10(1), 116. doi:10.1186/s40478-022-01416-6

Gregersen, E., Betzer, C., Kim, W. S., Kovacs, G., Reimer, L., Halliday, G. M., Thiel,

S., & Jensen, P. H. (2021). Alpha-synuclein activates the classical complement

pathway and mediates complement-dependent cell toxicity. *J*

Neuroinflammation, 18(1), 177. doi:10.1186/s12974-021-02225-9

Haas, L. T., Kostylev, M. A., & Strittmatter, S. M. (2014). Therapeutic molecules and

endogenous ligands regulate the interaction between brain cellular prion

protein (PrPC) and metabotropic glutamate receptor 5 (mGluR5). *J Biol Chem,*

289(41), 28460-28477. doi:10.1074/jbc.M114.584342

Haas, L. T., Salazar, S. V., Smith, L. M., Zhao, H. R., Cox, T. O., Herber, C. S.,

Degnan, A. P., Balakrishnan, A., Macor, J. E., Albright, C. F., & Strittmatter, S.

M. (2017). Silent Allosteric Modulation of mGluR5 Maintains Glutamate

Signaling while Rescuing Alzheimer's Mouse Phenotypes. *Cell Rep, 20*(1), 76-

88. doi:10.1016/j.celrep.2017.06.023

Hamilton, A., Esseltine, J. L., DeVries, R. A., Cregan, S. P., & Ferguson, S. S. (2014).

Metabotropic glutamate receptor 5 knockout reduces cognitive impairment and

pathogenesis in a mouse model of Alzheimer's disease. *Mol Brain, 7*, 40.

doi:10.1186/1756-6606-7-40

Hamilton, A., Vasefi, M., Vander Tuin, C., McQuaid, R. J., Anisman, H., & Ferguson,

S. S. (2016). Chronic Pharmacological mGluR5 Inhibition Prevents Cognitive

Impairment and Reduces Pathogenesis in an Alzheimer Disease Mouse Model.

Cell Rep, 15(9), 1859-1865. doi:10.1016/j.celrep.2016.04.077

Hammond, J. W., Bellizzi, M. J., Ware, C., Qiu, W. Q., Saminathan, P., Li, H., Luo, S.,

Ma, S. A., Li, Y., & Gelbard, H. A. (2020). Complement-dependent synapse loss

and microgliosis in a mouse model of multiple sclerosis. *Brain Behav Immun,*

87, 739-750. doi:10.1016/j.bbi.2020.03.004

Hammond, T. R., Robinton, D., & Stevens, B. (2018). Microglia and the Brain:
Complementary Partners in Development and Disease. *Annu Rev Cell Dev*
Biol, 34, 523-544. doi:10.1146/annurev-cellbio-100616-060509

Hardy, J. A., & Higgins, G. A. (1992). Alzheimer's disease: the amyloid cascade
hypothesis. *Science, 256*(5054), 184-185. doi:10.1126/science.1566067

Hasegawa, S., Sakuragi, S., Tominaga-Yoshino, K., & Ogura, A. (2015). Dendritic
spine dynamics leading to spine elimination after repeated inductions of LTD.
Sci Rep, 5, 7707. doi:10.1038/srep07707

Hernandez, M. X., Jiang, S., Cole, T. A., Chu, S. H., Fonseca, M. I., Fang, M. J.,
Hohsfield, L. A., Torres, M. D., Green, K. N., Wetsel, R. A., Mortazavi, A., &
Tenner, A. J. (2017). Prevention of C5aR1 signaling delays microglial
inflammatory polarization, favors clearance pathways and suppresses cognitive
loss. *Mol Neurodegener, 12*(1), 66. doi:10.1186/s13024-017-0210-z

Hong, S., Beja-Glasser, V. F., Nfonoyim, B. M., Frouin, A., Li, S., Ramakrishnan, S.,
Merry, K. M., Shi, Q., Rosenthal, A., Barres, B. A., Lemere, C. A., Selkoe, D. J.,
& Stevens, B. (2016). Complement and microglia mediate early synapse loss
in Alzheimer mouse models. *Science, 352*(6286), 712-716.
doi:10.1126/science.aad8373

Hsieh, H., Boehm, J., Sato, C., Iwatsubo, T., Tomita, T., Sisodia, S., & Malinow, R.
(2006). AMPAR removal underlies Abeta-induced synaptic depression and
dendritic spine loss. *Neuron, 52*(5), 831-843. doi:10.1016/j.neuron.2006.10.035

Hu, N. W., Nicoll, A. J., Zhang, D., Mably, A. J., O'Malley, T., Purro, S. A., Terry, C.,
Collinge, J., Walsh, D. M., & Rowan, M. J. (2014). mGlu5 receptors and cellular
prion protein mediate amyloid-β-facilitated synaptic long-term depression in
vivo. *Nat Commun, 5*, 3374. doi:10.1038/ncomms4374

Jackson, J., Jambrina, E., Li, J., Marston, H., Menzies, F., Phillips, K., & Gilmour, G. (2019). Targeting the Synapse in Alzheimer's Disease. *Front Neurosci, 13*, 735. doi:10.3389/fnins.2019.00735

Jo, J., Whitcomb, D. J., Olsen, K. M., Kerrigan, T. L., Lo, S. C., Bru-Mercier, G., Dickinson, B., Scullion, S., Sheng, M., Collingridge, G., & Cho, K. (2011). Aβ(1-42) inhibition of LTP is mediated by a signaling pathway involving caspase-3, Akt1 and GSK-3β. *Nat Neurosci, 14*(5), 545-547. doi:10.1038/nn.2785

Kim, J. H., Anwyl, R., Suh, Y. H., Djamgoz, M. B., & Rowan, M. J. (2001). Use-dependent effects of amyloidogenic fragments of (beta)-amyloid precursor protein on synaptic plasticity in rat hippocampus in vivo. *J Neurosci, 21*(4), 1327-1333. doi:10.1523/jneurosci.21-04-01327.2001

Koffie, R. M., Hyman, B. T., & Spires-Jones, T. L. (2011). Alzheimer's disease: synapses gone cold. *Mol Neurodegener, 6*(1), 63. doi:10.1186/1750-1326-6-63

Koffie, R. M., Meyer-Luehmann, M., Hashimoto, T., Adams, K. W., Mielke, M. L., Garcia-Alloza, M., Micheva, K. D., Smith, S. J., Kim, M. L., Lee, V. M., Hyman, B. T., & Spires-Jones, T. L. (2009). Oligomeric amyloid beta associates with postsynaptic densities and correlates with excitatory synapse loss near senile plaques. *Proc Natl Acad Sci U S A, 106*(10), 4012-4017. doi:10.1073/pnas.0811698106

Lacor, P. N., Buniel, M. C., Chang, L., Fernandez, S. J., Gong, Y., Viola, K. L., Lambert, M. P., Velasco, P. T., Bigio, E. H., Finch, C. E., Krafft, G. A., & Klein, W. L. (2004). Synaptic targeting by Alzheimer's-related amyloid beta oligomers. *J Neurosci, 24*(45), 10191-10200. doi:10.1523/JNEUROSCI.3432-04.2004

Lacor, P. N., Buniel, M. C., Furlow, P. W., Clemente, A. S., Velasco, P. T., Wood, M., Viola, K. L., & Klein, W. L. (2007). Abeta oligomer-induced aberrations in

synapse composition, shape, and density provide a molecular basis for loss of connectivity in Alzheimer's disease. *J Neurosci, 27*(4), 796-807. doi:10.1523/JNEUROSCI.3501-06.2007

Lambert, M. P., Barlow, A. K., Chromy, B. A., Edwards, C., Freed, R., Liosatos, M., Morgan, T. E., Rozovsky, I., Trommer, B., Viola, K. L., Wals, P., Zhang, C., Finch, C. E., Krafft, G. A., & Klein, W. L. (1998). Diffusible, nonfibrillar ligands derived from Abeta1-42 are potent central nervous system neurotoxins. *Proc Natl Acad Sci U S A, 95*(11), 6448-6453. doi:10.1073/pnas.95.11.6448

Li, S. H., Colson, T. L., Abd-Elrahman, K. S., & Ferguson, S. S. G. (2022). Metabotropic Glutamate Receptor 5 Antagonism Reduces Pathology and Differentially Improves Symptoms in Male and Female Heterozygous zQ175 Huntington's Mice. *Front Mol Neurosci, 15,* 801757. doi:10.3389/fnmol.2022.801757

Li, S., Hong, S., Shepardson, N. E., Walsh, D. M., Shankar, G. M., & Selkoe, D. (2009). Soluble oligomers of amyloid Beta protein facilitate hippocampal long-term depression by disrupting neuronal glutamate uptake. *Neuron, 62*(6), 788-801. doi:10.1016/j.neuron.2009.05.012

Li, S., Jin, M., Koeglsperger, T., Shepardson, N. E., Shankar, G. M., & Selkoe, D. J. (2011). Soluble Aβ oligomers inhibit long-term potentiation through a mechanism involving excessive activation of extrasynaptic NR2B-containing NMDA receptors. *J Neurosci, 31*(18), 6627-6638. doi:10.1523/jneurosci.0203-11.2011

Li, S., & Selkoe, D. J. (2020). A mechanistic hypothesis for the impairment of synaptic plasticity by soluble Aβ oligomers from Alzheimer's brain. *J Neurochem, 154*(6), 583-597. doi:10.1111/jnc.15007

Lian, H., Litvinchuk, A., Chiang, A. C., Aithmitti, N., Jankowsky, J. L., & Zheng, H.
(2016). Astrocyte-Microglia Cross Talk through Complement Activation
Modulates Amyloid Pathology in Mouse Models of Alzheimer's Disease. *J
Neurosci, 36*(2), 577-589. doi:10.1523/jneurosci.2117-15.2016

Liddelow, S. A., Guttenplan, K. A., Clarke, L. E., Bennett, F. C., Bohlen, C. J.,
Schirmer, L., Bennett, M. L., Münch, A. E., Chung, W. S., Peterson, T. C.,
Wilton, D. K., Frouin, A., Napier, B. A., Panicker, N., Kumar, M., Buckwalter, M.
S., Rowitch, D. H., Dawson, V. L., . . . Barres, B. A. (2017). Neurotoxic reactive
astrocytes are induced by activated microglia. *Nature, 541*(7638), 481-487.
doi:10.1038/nature21029

Litvinchuk, A., Wan, Y. W., Swartzlander, D. B., Chen, F., Cole, A., Propson, N. E.,
Wang, Q., Zhang, B., Liu, Z., & Zheng, H. (2018). Complement C3aR
Inactivation Attenuates Tau Pathology and Reverses an Immune Network
Deregulated in Tauopathy Models and Alzheimer's Disease. *Neuron, 100*(6),
1337-1353.e1335. doi:10.1016/j.neuron.2018.10.031

Liu, J., Chang, L., Song, Y., Li, H., & Wu, Y. (2019). The Role of NMDA Receptors in
Alzheimer's Disease. *Front Neurosci, 13*, 43. doi:10.3389/fnins.2019.00043

Lopez-Sanchez, C., Poejo, J., Garcia-Lopez, V., Salazar, J., Garcia-Martinez, V., &
Gutierrez-Merino, C. (2022). Kaempferol prevents the activation of complement
C3 protein and the generation of reactive A1 astrocytes that mediate rat brain
degeneration induced by 3-nitropropionic acid. *Food Chem Toxicol, 164*,
113017. doi:10.1016/j.fct.2022.113017

Lu, Y. M., Jia, Z., Janus, C., Henderson, J. T., Gerlai, R., Wojtowicz, J. M., & Roder,
J. C. (1997). Mice lacking metabotropic glutamate receptor 5 show impaired

learning and reduced CA1 long-term potentiation (LTP) but normal CA3 LTP. *J Neurosci, 17*(13), 5196-5205. doi:10.1523/jneurosci.17-13-05196.1997

Mango, D., & Nisticò, R. (2018). Role of ASIC1a in Aβ-induced synaptic alterations in the hippocampus. *Pharmacol Res, 131*, 61-65. doi:10.1016/j.phrs.2018.03.016

March, D. R., Proctor, L. M., Stoermer, M. J., Sbaglia, R., Abbenante, G., Reid, R. C., Woodruff, T. M., Wadi, K., Paczkowski, N., Tyndall, J. D., Taylor, S. M., & Fairlie, D. P. (2004). Potent cyclic antagonists of the complement C5a receptor on human polymorphonuclear leukocytes. Relationships between structures and activity. *Mol Pharmacol, 65*(4), 868-879. doi:10.1124/mol.65.4.868

McGeer, P. L., Lee, M., & McGeer, E. G. (2017). A review of human diseases caused or exacerbated by aberrant complement activation. *Neurobiol Aging, 52*, 12-22. doi:10.1016/j.neurobiolaging.2016.12.017

Nicoletti, F., Bockaert, J., Collingridge, G. L., Conn, P. J., Ferraguti, F., Schoepp, D. D., Wroblewski, J. T., & Pin, J. P. (2011). Metabotropic glutamate receptors: from the workbench to the bedside. *Neuropharmacology, 60*(7-8), 1017-1041. doi:10.1016/j.neuropharm.2010.10.022

O'Riordan, K. J., Hu, N. W., & Rowan, M. J. (2018). Aß Facilitates LTD at Schaffer Collateral Synapses Preferentially in the Left Hippocampus. *Cell Rep, 22*(8), 2053-2065. doi:10.1016/j.celrep.2018.01.085

Palmer, M. J., Irving, A. J., Seabrook, G. R., Jane, D. E., & Collingridge, G. L. (1997). The group I mGlu receptor agonist DHPG induces a novel form of LTD in the CA1 region of the hippocampus. *Neuropharmacology, 36*(11-12), 1517-1532. doi:10.1016/s0028-3908(97)00181-0

Parsons, C. G., Stöffler, A., & Danysz, W. (2007). Memantine: a NMDA receptor antagonist that improves memory by restoration of homeostasis in the

glutamatergic system--too little activation is bad, too much is even worse. *Neuropharmacology, 53*(6), 699-723. doi:10.1016/j.neuropharm.2007.07.013

Ricklin, D., Hajishengallis, G., Yang, K., & Lambris, J. D. (2010). Complement: a key system for immune surveillance and homeostasis. *Nat Immunol, 11*(9), 785-797. doi:10.1038/ni.1923

Sanderson, T. M., Ralph, L. T., Amici, M., Ng, A. N., Kaang, B. K., Zhuo, M., Kim, S. J., Georgiou, J., & Collingridge, G. L. (2022). Selective Recruitment of Presynaptic and Postsynaptic Forms of mGluR-LTD. *Front Synaptic Neurosci, 14*, 857675. doi:10.3389/fnsyn.2022.857675

Schafer, D. P., Lehrman, E. K., Kautzman, A. G., Koyama, R., Mardinly, A. R., Yamasaki, R., Ransohoff, R. M., Greenberg, M. E., Barres, B. A., & Stevens, B. (2012). Microglia sculpt postnatal neural circuits in an activity and complement-dependent manner. *Neuron, 74*(4), 691-705. doi:10.1016/j.neuron.2012.03.026

Schartz, N. D., & Tenner, A. J. (2020). The good, the bad, and the opportunities of the complement system in neurodegenerative disease. *J Neuroinflammation, 17*(1), 354. doi:10.1186/s12974-020-02024-8

Selkoe, D. J. (2002). Alzheimer's disease is a synaptic failure. *Science, 298*(5594), 789-791. doi:10.1126/science.1074069

Shankar, G. M., Bloodgood, B. L., Townsend, M., Walsh, D. M., Selkoe, D. J., & Sabatini, B. L. (2007). Natural oligomers of the Alzheimer amyloid-beta protein induce reversible synapse loss by modulating an NMDA-type glutamate receptor-dependent signaling pathway. *J Neurosci, 27*(11), 2866-2875. doi:10.1523/jneurosci.4970-06.2007

Shankar, G. M., Li, S., Mehta, T. H., Garcia-Munoz, A., Shepardson, N. E., Smith, I., Brett, F. M., Farrell, M. A., Rowan, M. J., Lemere, C. A., Regan, C. M., Walsh, D. M., Sabatini, B. L., & Selkoe, D. J. (2008). Amyloid-beta protein dimers isolated directly from Alzheimer's brains impair synaptic plasticity and memory. *Nat Med, 14*(8), 837-842. doi:10.1038/nm1782

Shinoda, Y., Tanaka, T., Tominaga-Yoshino, K., & Ogura, A. (2010). Persistent synapse loss induced by repetitive LTD in developing rat hippocampal neurons. *PLoS One, 5*(4), e10390. doi:10.1371/journal.pone.0010390

Shipton, O. A., Leitz, J. R., Dworzak, J., Acton, C. E., Tunbridge, E. M., Denk, F., Dawson, H. N., Vitek, M. P., Wade-Martins, R., Paulsen, O., & Vargas-Caballero, M. (2011). Tau protein is required for amyloid {beta}-induced impairment of hippocampal long-term potentiation. *J Neurosci, 31*(5), 1688-1692. doi:10.1523/jneurosci.2610-10.2011

Spurrier, J., Nicholson, L., Fang, X. T., Stoner, A. J., Toyonaga, T., Holden, D., Siegert, T. R., Laird, W., Allnutt, M. A., Chiasseu, M., Brody, A. H., Takahashi, H., Nies, S. H., Pérez-Cañamás, A., Sadasivam, P., Lee, S., Li, S., Zhang, L., . . . Strittmatter, S. M. (2022). Reversal of synapse loss in Alzheimer mouse models by targeting mGluR5 to prevent synaptic tagging by C1Q. *Sci Transl Med, 14*(647), eabi8593. doi:10.1126/scitranslmed.abi8593

Su, L. D., Wang, N., Han, J., & Shen, Y. (2022). Group 1 Metabotropic Glutamate Receptors in Neurological and Psychiatric Diseases: Mechanisms and Prospective. *Neuroscientist, 28*(5), 453-468. doi:10.1177/10738584211021018

Sumichika, H., Sakata, K., Sato, N., Takeshita, S., Ishibuchi, S., Nakamura, M., Kamahori, T., Ehara, S., Itoh, K., Ohtsuka, T., Ohbora, T., Mishina, T., Komatsu, H., & Naka, Y. (2002). Identification of a potent and orally active non-

peptide C5a receptor antagonist. *J Biol Chem, 277*(51), 49403-49407.

doi:10.1074/jbc.M209672200

Taylor, H. B. C., Emptage, N. J., & Jeans, A. F. (2021). Long-term depression links

amyloid-beta to the pathological hyperphosphorylation of tau. *Cell Rep, 36*(9),

109638. doi:10.1016/j.celrep.2021.109638

Terry, R. D., Masliah, E., Salmon, D. P., Butters, N., DeTeresa, R., Hill, R., Hansen,

L. A., & Katzman, R. (1991). Physical basis of cognitive alterations in

Alzheimer's disease: synapse loss is the major correlate of cognitive

impairment. *Ann Neurol, 30*(4), 572-580. doi:10.1002/ana.410300410

Um, J. W., Kaufman, A. C., Kostylev, M., Heiss, J. K., Stagi, M., Takahashi, H., Kerrisk,

M. E., Vortmeyer, A., Wisniewski, T., Koleske, A. J., Gunther, E. C., Nygaard,

H. B., & Strittmatter, S. M. (2013). Metabotropic glutamate receptor 5 is a

coreceptor for Alzheimer aβ oligomer bound to cellular prion protein. *Neuron,*

79(5), 887-902. doi:10.1016/j.neuron.2013.06.036

Volianskis, A., France, G., Jensen, M. S., Bortolotto, Z. A., Jane, D. E., & Collingridge,

G. L. (2015). Long-term potentiation and the role of N-methyl-D-aspartate

receptors. *Brain Res, 1621*, 5-16. doi:10.1016/j.brainres.2015.01.016

Walsh, D. M., Klyubin, I., Fadeeva, J. V., Cullen, W. K., Anwyl, R., Wolfe, M. S.,

Rowan, M. J., & Selkoe, D. J. (2002). Naturally secreted oligomers of amyloid

beta protein potently inhibit hippocampal long-term potentiation in vivo. *Nature,*

416(6880), 535-539. doi:10.1038/416535a

Wang, H. A., Lee, J. D., Lee, K. M., Woodruff, T. M., & Noakes, P. G. (2017).

Complement C5a-C5aR1 signalling drives skeletal muscle macrophage

recruitment in the hSOD1(G93A) mouse model of amyotrophic lateral sclerosis.

Skelet Muscle, 7(1), 10. doi:10.1186/s13395-017-0128-8

Wang, Q., Rowan, M. J., & Anwyl, R. (2004). Beta-amyloid-mediated inhibition of NMDA receptor-dependent long-term potentiation induction involves activation of microglia and stimulation of inducible nitric oxide synthase and superoxide. *J Neurosci, 24*(27), 6049-6056. doi:10.1523/jneurosci.0233-04.2004

Wang, R., & Reddy, P. H. (2017). Role of Glutamate and NMDA Receptors in Alzheimer's Disease. *J Alzheimers Dis, 57*(4), 1041-1048. doi:10.3233/jad-160763

Whitcomb, D. J., Hogg, E. L., Regan, P., Piers, T., Narayan, P., Whitehead, G., Winters, B. L., Kim, D. H., Kim, E., St George-Hyslop, P., Klenerman, D., Collingridge, G. L., Jo, J., & Cho, K. (2015). Intracellular oligomeric amyloid-beta rapidly regulates GluA1 subunit of AMPA receptor in the hippocampus. *Sci Rep, 5*, 10934. doi:10.1038/srep10934

Wiegert, J. S., & Oertner, T. G. (2013). Long-term depression triggers the selective elimination of weakly integrated synapses. *Proc Natl Acad Sci U S A, 110*(47), E4510-4519. doi:10.1073/pnas.1315926110

Wiegert, J. S., Pulin, M., Gee, C. E., & Oertner, T. G. (2018). The fate of hippocampal synapses depends on the sequence of plasticity-inducing events. *Elife, 7*. doi:10.7554/eLife.39151

Woodruff, T. M., Costantini, K. J., Crane, J. W., Atkin, J. D., Monk, P. N., Taylor, S. M., & Noakes, P. G. (2008). The complement factor C5a contributes to pathology in a rat model of amyotrophic lateral sclerosis. *J Immunol, 181*(12), 8727-8734. doi:10.4049/jimmunol.181.12.8727

Wu, T., Dejanovic, B., Gandham, V. D., Gogineni, A., Edmonds, R., Schauer, S., Srinivasan, K., Huntley, M. A., Wang, Y., Wang, T. M., Hedehus, M., Barck, K. H., Stark, M., Ngu, H., Foreman, O., Meilandt, W. J., Elstrott, J., Chang, M. C.,

. . . Hanson, J. E. (2019). Complement C3 Is Activated in Human AD Brain and

Is Required for Neurodegeneration in Mouse Models of Amyloidosis and

Tauopathy. *Cell Rep, 28*(8), 2111-2123.e2116.

doi:10.1016/j.celrep.2019.07.060

Wu, Y., Dissing-Olesen, L., MacVicar, B. A., & Stevens, B. (2015). Microglia: Dynamic

Mediators of Synapse Development and Plasticity. *Trends Immunol, 36*(10),

605-613. doi:10.1016/j.it.2015.08.008

Yang, W., Zhou, X., Ryazanov, A. G., & Ma, T. (2021). Suppression of the kinase for

elongation factor 2 alleviates mGluR-LTD impairments in a mouse model of

Alzheimer's disease. *Neurobiol Aging, 98*, 225-230.

doi:10.1016/j.neurobiolaging.2020.11.016

Yasojima, K., Schwab, C., McGeer, E. G., & McGeer, P. L. (1999). Up-regulated

production and activation of the complement system in Alzheimer's disease

brain. *Am J Pathol, 154*(3), 927-936. doi:10.1016/s0002-9440(10)65340-0

Zhang, J., Malik, A., Choi, H. B., Ko, R. W., Dissing-Olesen, L., & MacVicar, B. A.

(2014). Microglial CR3 activation triggers long-term synaptic depression in the

hippocampus via NADPH oxidase. *Neuron, 82*(1), 195-207.

doi:10.1016/j.neuron.2014.01.043

Zhang, Y. N., Fan, J. K., Gu, L., Yang, H. M., Zhan, S. Q., & Zhang, H. (2021).

Metabotropic glutamate receptor 5 inhibits α-synuclein-induced microglia

inflammation to protect from neurotoxicity in Parkinson's disease. *J

Neuroinflammation, 18*(1), 23. doi:10.1186/s12974-021-02079-1

Figures

Figure 1: oAβ enhances mGlu₅R-LTD at CA1 synapses.

(**A-B**) Validation of oAβ preparation, confirming its ability to impair LTP. Delivery of a theta-burst protocol (TBS; see Methods), at the time indicated by the arrow, induced LTP in -oAβ slices (51 ± 7 %, n = 28). LTP was impaired in the interleaved +oAβ slices (17 ± 3 %, n = 28). Representative fEPSP traces for both conditions are shown to the right of the time course plot (superimposed for sample experiment, before and after TBS, at times indicated by the numbers 1-4). **B** Bar graph shows the level of LTP in each experiment. p = 0.00002.

(**C-D**) Application of DHPG (50 µM, 10 min), induced LTD in -oAβ slices from

mGlu$_5$R$^{+/+}$ mice (21 ± 2 %, n = 9). Interleaved +oAβ slices showed an enhancement

of DHPG-LTD (31 ± 4 %, n= 9) with summary of LTD experiments in **D**. p = 0.032.

Representative fEPSP traces demonstrating enhanced DHPG-LTD in +oAβ compared

to -oAβ slices.

(**E-F**) In slices from littermate mGlu$_5$R$^{-/-}$ mice, DHPG failed to induce LTD in either

condition (+oAβ = 2 ± 3 %, n=10; -oAβ = -3 ± 2 %, n=10) with summary in **F**.

Representative fEPSP traces showing the absence of DHPG-LTD in both conditions.

Sample fEPSP traces in all panels are the mean of 4 consecutive responses at the

indicated time points (1-4). For each experiment, sample traces before and after

DHPG treatment are superimposed. Two-tailed t-tests were used in **B**, **D** and **F**.

Figure 2: Enhancement of mGlu$_5$R-LTD by oAβ requires NMDARs.

(**A-B**) In a separate cohort, pretreatment with oAβ also resulted in an enhancement of DHPG-LTD (34 ± 2 %, n=22) compared to the interleaved -oAβ slices (21 ± 2 %, n=22). p=0.00031.

(**C-D**) In the presence of L689,560 (5 µM), the magnitude of DHPG-LTD was similar in +oAβ slices (26 ± 4 %, n=9) and -oAβ slices (22 ± 2 %, n=9).

Sample fEPSP traces in all panels are the mean of 4 consecutive responses at the indicated time points (1-4). For each experiment, sample traces before and after DHPG treatment are superimposed. A two-tailed t-test was used in **B** and **D**.

Figure 3: Complement C5aR and C3aR signaling is necessary for oAβ to enhance

mGlu$_5$R-LTD.

(**A-B**) In a third cohort, pretreatment with oAβ enhanced DHPG-LTD (43 ± 3 %, n=16)

compared to the interleaved -oAβ slices (26 ± 3 %, n=16). p=0.00064.

(**C-D**) In the presence of the C5aR antagonist PMX205 (0.3 µM), the level of DHPG-

LTD was similar in +oAβ slices (35 ± 4 %, n=8) and -oAβ slices (29 ± 6 %, n=8).

(**E-F**) In the presence of a separate C5aR antagonist, W54011 (0.3 µM) the magnitude

of DHPG-LTD was similar in +oAβ slices (32 ± 5 %, n=6) compared to -oAβ slices (29

± 4 %, n=6).

(**G-H**) In the presence of a C3aR antagonist, SB290157 (1 µM), DHPG-LTD magnitude

was not significantly different between +oAβ slices (35 ± 3 %, n=8) and -oAβ slices

(28 ± 6 %, n=8).

Sample fEPSP traces in all panels are the mean of 4 consecutive responses at the

indicated time points. For each experiment, sample traces before and after DHPG

treatment are superimposed. Two-tailed t-tests were used in **B**, **D**, **F** and **H**.

<u>Supplemental Figure 1: Acute oAβ application does not affect basal synaptic strength</u>

<u>at CA1 synapses.</u>

(**A-C**) CA3-CA1 basal synaptic transmission is similar in +oAβ slices and -oAβ slices,

as assessed by input-output curves. **A** Fiber volley (FV) amplitude *versus* stimulus

intensity (line slopes: +oAβ = -0.026 ± 0.003, n = 13; -oAβ = -0.030 ± 0.003, n = 13).

B fEPSP slope *versus* FV amplitude (line slopes: +oAβ = 2.517 ± 0.237, n = 13; black,

versus -oAβ = 2.513 ± 0.362, n = 13; white). **C** Representative fEPSP traces at

different stimulus intensity (multiples of threshold) for both conditions.