Ai Na Ng[1,*], Eric W. Salter[2,3,*], John Georgiou[3], Zuner A. Bortolotto[1], Graham L.

Collingridge[1,2,3,4,5]

[1] Glutamate Research Group, School of Physiology, Pharmacology and Neuroscience,

University of Bristol, Biomedical Sciences Building, University Walk, Bristol, BS8 1TD,

United Kingdom.

[2] Lunenfeld-Tanenbaum Research Institute, Mount Sinai Hospital, Sinai Health

System, 600 University Avenue, Toronto, ON, M5G 1X5, Canada.

[3] Department of Physiology, University of Toronto, Toronto, Ontario, M5S 1A8,

Canada

[4] Tanz Centre for Research in Neurodegenerative Diseases, University of Toronto,

Krembil Discovery Tower, 60 Leonard Avenue, Toronto, ON, M5T 0S8, Canada.

[5]Correspondence: Graham L. Collingridge, collingridge@lunenfeld.ca

*These authors contributed equally

Abstract

Synaptic dysfunction, weakening, and loss of synapses are well-correlated with the

pathology of Alzheimer's disease (AD). Recent studies have implicated hyper-

activation of the complement cascade as the driving force for loss of synapses in AD.

However, the synaptic cues to trigger pathological complement activity remain elusive.

To address this, we studied the actions of oligomeric, 42 residue length amyloid β

(oAβ) on metabotropic glutamate receptor (mGluR) long-term depression (mGluR-

LTD) at hippocampal CA1 synapses, a form of synaptic plasticity that is disrupted in

rodent models of AD. We found that exogenous application of oAβ to mouse

hippocampal slices enhanced the magnitude of mGluR subtype 5 (mGlu$_5$R)-

dependent LTD. This enhanced synaptic weakening occurred via both NMDARs and

complement C3aR/C5aR signaling, as revealed by pharmacological inhibition

experiments. Our findings reveal a novel mechanistic interaction between mGlu$_5$R,

NMDARs, and the complement cascade in synaptic weakening induced by oAβ, which

could represent an early trigger of synaptic degeneration in AD.

Introduction

Alzheimer's disease (AD) is a progressive neurodegenerative disease characterized

by memory loss, cognitive deficits and changes to personality and behavior (Breijyeh

& Karaman, 2020). Although it is the most common cause of dementia affecting the

aging population worldwide, therapeutic options are limited. The development of better

treatments necessitates a deeper understanding of pathophysiological mechanisms.

Weakening and subsequent loss of synapses are early events in AD progression that

precede the deterioration of memory and cognitive functions (Koffie, Hyman, & Spires-

Jones, 2011; Li & Selkoe, 2020). As such, targeting these events has the potential to

prevent AD before irreversible neuronal damage can occur. Normally, synaptic

strength is balanced by long-term potentiation (LTP) and long-term depression (LTD),

which are thought to provide the cellular and molecular basis of learning and memory

(Bliss & Collingridge, 1993). Substantial evidence implicates amyloid precursor protein

(APP) in the pathogenesis of AD (O'brien & Wong, 2011). APP is cleaved by

secretases to yield amyloid beta (Aβ) peptides of 30-51 amino acids, including Aβ$_{1-42}$

which is highly prone to aggregation and rapidly forms oligomeric Aβ (oAβ) which accumulates in the brain (Lue, et al., 1999). *In vitro* and *in vivo* studies of AD models have found that oAβ mediates a shift in the LTP/LTD balance, favoring LTD over LTP, leading to net synapse loss (Kim, Anwyl, Suh, & Djamgoz, 2001; Hsieh, et al., 2006; Shankar, et al., 2007; Shankar, et al., 2008; Li, et al., 2009; Jo, et al., 2011; Cline, Bicca, Viola, & Klein, 2018; Li & Selkoe, 2022). Therefore, understanding the mechanisms by which oAβ alters synaptic plasticity to bias towards LTD and synapse elimination is essential to determining the pathological events underlying AD.

The complement cascade is an innate immune pathway which has emerged as a central driver of synapse loss in AD (Wu, Dissing-Olesen, MacVicar, & Stevens, 2015; Hammond, Robinton, & Stevens, 2018). Complement cascade activation leads to phagocytosis by microglia via C3b/CR3, as well as chemotactic and inflammatory signaling via C3a-C3aR and C5a-C5aR axes (Ricklin, Hajishengallis, Yang, & Lambris, 2010). Tissue from post-mortem samples of patients with AD demonstrates activation of complement activity, including increased deposition of C1q and proteolytic cleavage of C3 (Afagh, Cummings, Cribbs, Cotman, & Tenner, 1996; Yasojima, Schwab, McGeer, & McGeer, 1999; Wu, et al., 2019). Further, oAβ acts as a potent activator of the complement cascade and oAβ-mediated synapse loss is dependent on the complement pathway (Hong, et al., 2016; Bie, Wu, Foss, & Naguib, 2019). Finally, in multiple mouse models of AD, there is aberrant activation of complement-mediated synapse phagocytosis. Genetic deletion or pharmacological inhibition of complement cascade proteins prevents synapse and neuronal loss, preserving cognitive function (Fonseca, et al., 2009; Hong, et al., 2016; Dejanovic, et al., 2018; Litvinchuk, et al., 2018; Wu, et al., 2019; Carvalho, et al., 2022). However,

the upstream synaptic signals that lead to aberrant activation of complement and
subsequent synapse loss are not well understood.

LTP and LTD of glutamatergic synapses is mediated by activation of N-methyl-D-
aspartate receptors (NMDARs) and/or metabotropic glutamate receptors (mGluRs;
Collingridge, Peineau, Howland, & Wang, 2010; Nicoletti, et al., 2011; Volianskis, et
al., 2015). LTD, induced either by NMDARs or mGluRs, can trigger subsequent
synapse elimination (Shinoda, Tanaka, Tominaga-Yoshino, & Ogura, 2010; Wiegert &
Oertner, 2013; Hasegawa, Sakuragi, Tominaga-Yoshino, & Ogura, 2015; Wiegert,
Pulin, Gee, & Oertner, 2018). Both mGluR- and NMDAR-dependent signaling are
affected in AD, leading to synaptic plasticity impairments and aberrant synapse loss
(Hsieh, et al., 2006; Shankar, et al., 2007; Shankar, et al., 2008; Jo, et al., 2011; Li, et
al., 2011; Cavallucci, et al., 2013; Beraldo, et al., 2016; Jackson, et al., 2019; Li &
Selkoe, 2020; Abd-Elrahman, et al., 2020). While it is known that mGluRs form protein
complexes with NMDARs (Tu, et al., 1999; Perroy, et al., 2008), the role of this
interaction in mediating synaptic impairments in AD is not well understood.
Interestingly, the complement cascade has recently been implicated in synapse loss
mediated by mGluRs in AD mouse models. Intra-hippocampal activation of mGluRs
with the group I ($mGlu_1R$ and $mGlu_5R$) agonist (S)-3,5-dihydroxyphenyl glycine (S
isomer, abbreviated herein as DHPG) *in vivo* increased C1q synaptic tagging and
synapse elimination. Additionally, knockout of $mGlu_1R$ in a mouse model of AD
prevented microglial synapse phagocytosis (Bie, Wu, Foss, & Naguib, 2019). A silent
allosteric modulator of $mGlu_5R$ has also recently been found to restore synapse
density through reduced tagging by C1q and microglia engulfment in a mouse model
of AD (Spurrier, 2022).

However, the interaction between NMDARs and mGluRs in promoting complement activation in AD is not defined. To address this, we used hippocampus slices treated with oAβ as an *in situ* model of AD. We found that oAβ enhanced mGlu$_5$R-mediated LTD via a mechanism that involves both the activation of NMDARs and the C3aR and C5aR arms of the complement cascade. These findings have implications pertinent to the understanding of the mechanism that leads to synaptic deficits that may underlie the initial stages of dementia.

Methods

Animals

Adult C57BL/6JOlaHsd male mice (10-16 weeks old) from Envigo (U.K.) or mGlu$_5$R knockout line (Lu, et al., 1997) backcrossed more than 10 generations to the C57BL/6J strain were used in this study. We chose to use male mice due to the recently reported sex specificity of oAβ-mGlu$_5$R binding which is present only in males (Abd-Elrahman, et al., 2020). Animals were housed four per cage in a room with designated 12 h light/dark cycle (lights on at 08:15/ off at 20:15) with temperature at 21 °C ± 2 °C and access to food and water *ad libitum.* All procedures were performed in accordance with the U.K. Animals (Scientific Procedures) Act of 1986. Genomic DNA from mGlu5$_R$ knockout litters were isolated from tail tips with DNeasy blood and tissue kit (Qiagen; cat. no. 69504) and amplified as described (Lu, et al., 1997). The amplified products were resolved and visualized in ethidium bromide-stained agarose gel.

Oligomeric amyloid-β$_{1-42}$ (oAβ) preparation

oAβ was prepared as described (Whitcomb, et al., 2015). Briefly, Aβ peptide$_{(1-42)}$ (Millipore; TFA recombinant human, ultra-pure; cat. no. AG912) was dissolved in 100

% 1,1,1,3,3,3-hexafluoro-2-propanol (HFIP; Sigma Aldrich; cat. no. 105228) to a final concentration of 1 mg/mL. The HFIP/peptide mixture was incubated at room temperature for 1 h and sonicated for 10 min in a water-bath sonicator. The mixture was then air-dried under a gentle stream of nitrogen gas for 1.5 h. The peptide crystal was re-suspended in 100 % DMSO, aliquoted to 10 μL per tube and stored at -80 °C until experimental use (see next section).

Hippocampal slice preparation

Mice were deeply anaesthetized via inhalation with a mixture of 5 % isofluorane / 95 % oxygen and euthanized by cervical dislocation. The brain was rapidly removed (<1 min) and chilled in cold ACSF solution (in mM: 124 NaCl, 3 KCl, 26 $NaHCO_3$, 1.25 NaH_2PO_4, 10 D-glucose, 2 $MgSO_4$ and 2 $CaCl_2$; bubbled continuously with 95% O_2 /5% CO_2). Hippocampi were isolated from both hemispheres in cold ACSF. Transverse slices of 400 μm in thickness were prepared from the dorsal ends of hippocampi using a McIlwain tissue chopper. The CA3 region was left intact for LTP but removed via a surgical cut in LTD experiments. Slices were transferred to a holding chamber with room temperature ACSF and allowed to recover for at least 1 h. On the day of experiment, the Aβ peptide/DMSO mixture was diluted to 100 μM in D-PBS, vortexed briefly and allowed to aggregate for 2 h at room temperature. The heterogenous mixture of oAβs was diluted to a final concentration of 500 nM in oxygenated ACSF and acutely applied to hippocampal slices for 2 h at room temperature. As an inclusion criterion to account for batch-to-batch variability, we routinely assessed the bioactivity of our aggregated oAβs by confirming the ability of each oAβ batch to impair hippocampal LTP (Figure S1A,B).

Electrophysiology

After vehicle or oAβ incubation, a dorsal hippocampal slice was submerged in a Slicemate recording chamber (Scientifica, U.K.) with ACSF perfusing at a rate of 2.5 mL/min and temperature maintained at 30.0 ± 0.5 °C. Extracellular field potentials were recorded from the stratum radiatum area of CA1 using a glass microelectrode (~2-3 MΩ) filled with ACSF. Responses were evoked by stimulating the Schaffer collaterals using a platinum-iridium bipolar electrode (FHC) positioned at the border between area CA2-CA1. Stimuli of 100 μs in duration were delivered once every 15 s via a DS2 constant voltage isolator (Digitimer, U.K.). Responses were amplified using a Multiclamp 700B amplifier (Molecular Devices, U.S.A.), digitized at 40 kHz and monitored in real time using WinLTP (WinLTP Ltd, U.K.) software (Anderson & Collingridge, 2007). After a period of stable baseline recordings, datapoints for input/output (I/O) plots were obtained with fixed stimulus intensity at 1x, 1.5x, 2x, 3x, and 4.5x. The lowest intensity at 1x was defined as the threshold for evoking a visually detectable field excitatory postsynaptic potential (fEPSP) (≤ 0.05 mV). fEPSPs for plasticity experiments were recorded using a stimulus intensity of 3x the threshold intensity. LTP was induced by a theta-burst stimulation (TBS) protocol, with four stimuli delivered at 100 Hz, repeated 10 times at a frequency of 5 Hz. LTD was chemically induced by bath-application of (S)-3,5-DHPG (50 μM; Abcam; cat. no. ab120007) for 10 min. For offline analysis, responses were filtered at 2 kHz and expressed as the mean of four consecutive responses.

Quantification and statistical analysis

Data in the time course plots were normalized to the mean of the entire baseline period, defined as 100 % and presented as mean ± standard error of mean (SEM). The magnitude of LTP or LTD was presented as percentage of potentiation or depression of fEPSP from baseline and quantified as mean of the last 5 min of time

course recordings. Fiber volleys (FVs) and fEPSPs were quantified using peak

amplitude and initial slope measurements, respectively, using WinLTP. The I/O plots

were analysed using Excel linear regression software (Microsoft). The respective

slope values from a fitted line between FV *versus* stimulation intensity and fEPSP

slope *versus* FV were used for quantification. Statistical comparisons between -oAβ

and +oAβ conditions were assessed using unpaired two-tailed Student's t-test

(Microsoft Excel). The level of significance was set at $p<0.05$.

Results

Acute oAβ exposure enhances mGlu$_5$R-LTD

To investigate the impact of oAβs on synaptic function, we incubated acute

hippocampus slices for 2 h with oAβ. We subsequently measured synaptic plasticity

with field potential recordings at Schaffer collateral-CA1 synapses. As a positive

control for the bioactivity of our oAβ preparation, we confirmed previous findings

(Shankar, et al., 2008; Li, et al., 2011) that oAβ was able to impair LTP in slices pre-

treated with oAβ (+oAβ) *versus* interleaved non-treated slices (-oAβ; Figure S1A,B).

Although LTP was impaired by acute application of oAβ, basal synaptic transmission

was unaltered, indicated by the overlapping input-output relationship of +oAβ

compared to -oAβ slices (Figure S1C-E). As such, this paradigm allowed us to study

early disruptions to plasticity that precedes the loss of synapses.

To investigate how oAβ alters mGluR-dependent synaptic plasticity mechanisms, we

induced LTD using DHPG (S isomer), a selective agonist for group I mGluRs (mGlu$_1$R

and mGlu$_5$R), that is widely used to study mGluR function. A short application of DHPG

(10 min) induced a stable LTD in -oAβ slices from WT mice (Figure 1A,B). The

magnitude of DHPG-LTD was significantly enhanced in the interleaved +oAβ slices (Figure 1A, B), consistent with published observations (Hsieh, et al., 2006; Shankar, et al., 2008; Li, et al., 2009; Cavallucci, et al., 2013). To determine the group I mGluR subtype-specificity of our observed phenotype, we used hippocampal slices from mGlu$_5$R$^{-/-}$ mice (Lu, et al., 1997) which were littermates to the mGlu$_5$R$^{+/+}$ mice in Figure 1A, B. DHPG-LTD was absent in both -oAβ and +oAβ slices (Figure 1C,D). This indicates that the form of LTD which is enhanced by oAβ requires the activation of mGlu$_5$R receptors (referred to hereafter as mGlu$_5$R LTD).

NMDARs are required for oAβ-mediated enhancement of mGlu$_5$R-LTD

There are two distinct forms of DHPG-LTD that can be distinguished by their dependence upon the activation of NMDARs (Palmer, Irving, Seabrook, Jane, & Collingridge, 1997; Sanderson, et al., 2022). To determine whether NMDARs are required for either mGlu$_5$R-LTD under control conditions or the enhanced mGlu$_5$R-LTD observed in the presence of oAβ, we used a selective NMDAR antagonist, L689,560. L689,560 was used as it acts at the glycine site of the NMDAR and so is independent of L-glutamate concentration, which might be altered by oAβ treatment (Li, et al., 2009). In these experiments, we interleaved a new set of controls and again observed the potentiation of mGlu$_5$R-LTD by oAβ (Figure 1E,F). We found that NMDAR activation was not required for mGlu$_5$R-LTD in control mice. However, the enhancement of DHPG-LTD by oAβs was completely prevented by L689,560 (Figure 1G,H) and therefore requires NMDARs.

oAβ enhanced mGlu$_5$R-LTD via complement cascade signaling

Early studies of the effects of oAβ on LTP found that inhibition of LTP was prevented

by application of minocycline, an inhibitor of microglia activation (Wang, Rowan, &

Anwyl, 2004). Further, it has recently been found that agonism of group I mGluR *in*

vivo activates the complement cascade leading to synapse loss (Bie, Wu, Foss, &

Naguib, 2019). We therefore hypothesized that complement signaling is necessary for

the enhancement of mGlu$_5$R-LTD by oAβs. We antagonized either C3aR or C5aR, as

both receptors have been implicated in the pathology of mouse models of AD

(Fonseca, et al., 2009; Lian, et al., 2016; Hernandez, et al., 2017; Litvinchuk, et al.,

2018; Carvalho, et al., 2022; Gomez-Arboledas, et al., 2022). For these experiments

we also interleaved a new set of controls (Figure 2A,B) and again observed the

potentiation of mGlu$_5$R-LTD by oAβ. To inhibit C5aR, we bath-applied PMX205, a

peptide-based antagonist (March, et al., 2004). PMX205 had no effect on the level of

mGlu$_5$R-LTD induced under control conditions but eliminated the potentiation induced

by oAβ (Figure 2C,D). We next used W54011, a non-peptide based C5aR antagonist

(Sumichika, et al., 2002). As with PMX205, bath-application of W54011 had no effect

on the level of mGlu$_5$R-LTD induced under control conditions but eliminated the

potentiation induced by oAβ (Figure 2E,F). Lastly, we investigated the role of C3aR

using the antagonist SB290157 (Ames, et al., 2001). As with the C5aR antagonists,

SB290157 abolished the enhanced mGlu$_5$R-LTD induced by oAβ without affecting the

level of mGlu$_5$R-LTD under control conditions (Figure 2G,H). Together, these data

indicate that the oAβ-mediated enhancement of mGlu$_5$R-LTD occurs via activation of

complement C3aR/C5aR signaling.

Discussion

The complement cascade is emerging as a key innate immune pathway for shaping

glutamatergic synapses during brain development, as well as driving synapse loss in

various disease states. However, the synaptic signals that lead to complement

activation remain poorly understood. In the present study, we uncovered a novel

signaling mechanism through which both complement C3aR and C5aR activation

amplified LTD of glutamatergic synapses. As oAβ has been identified as a key

mediator of synaptic toxicity in AD (Cline, Bicca, Viola, & Klein, 2018; Li & Selkoe,

2020), we used exogenous application of oAβ to acute hippocampus slices to model

the early stages of oAβ-mediated synaptic impairment. Our findings that basal synaptic

transmission is unaltered confirms that we engaged an early stage of synaptic

plasticity impairment (Figure S1). Thus, our findings reveal mechanisms which are

upstream of overt synapse loss. We uncovered that mGlu$_5$R-dependent LTD, induced

by DHPG application, was enhanced in slices treated with oAβ (Figure 1). Unlike

vehicle treated slices, the oAβ-mediated enhanced portion of mGlu$_5$R-LTD was

dependent on NMDAR signaling. Similarly, antagonism of either C3aR or C5aR

prevented oAβ from enhancing DHPG-LTD (Figure 2), without affecting the underlying

mGlu$_5$R-LTD *per se*. Together, our results have uncovered a novel interaction

between mGluRs, NMDARs, and the complement cascade, which drive pathological

levels of synaptic depression.

Glutamatergic synapse dysfunction in AD

A foundational hypothesis in AD research over the past two decades has centered

around synaptic failure, which posits that oAβs initially reduce synaptic efficacy and

subsequently cause neuronal loss (Selkoe, 2002; Li & Selkoe, 2020). Reductions in

AMPAR- and NMDAR-mediated synaptic transmission, as well as a shift in the LTP-

LTD balance favoring LTD, caused by oAβ (Cullen, Suh, Anwyl, & Rowan, 1997;

Walsh, et al., 2002; Hsieh, et al., 2006; Shankar, et al., 2007; Shankar, et al., 2008; Li, et al., 2009; Jo, et al., 2011; Li, et al., 2011; Cavallucci, et al., 2013) has provided converging evidence supporting the idea that a primary triggering event in AD is excess synapse weakening. Altered activity of both group I mGluRs and NMDARs by oAβ and in AD mouse models is central to glutamatergic synaptic failure. Interestingly, $mGlu_5R$ was identified as a co-receptor of cellular prion protein (PrP_c) for binding oAβ and is necessary for oAβ-induced intracellular signal transduction and synapse loss (Um, et al., 2013; Haas, Kostylev, & Strittmatter, 2014; Beraldo, et al., 2016; Brody & Strittmatter, 2018). Additionally, antagonists of $mGlu_5R$ and genetic deletion have both been found to reduce synaptic and behavioral deficits in AD mouse models (Hamilton, Esseltine, DeVries, Cregan, & Ferguson, 2014; Hamilton et al., 2016; Haas, et al., 2017; Abd-Elrahman, et al., 2020; Spurrier, 2022). Evidence that altered NMDAR signaling is causally related to the human condition is the use of memantine, a non-competitive NMDAR antagonist, for the treatment of AD. Memantine slows the progression of dementia and its underlying mechanism has been attributed to the normalization of synaptic plasticity (Parsons, Stöffler, & Danysz, 2007; Wang & Reddy, 2017; Liu, Chang, Song, Li, & Wu, 2019).

Despite these important advances, how NMDARs and mGluRs, individually or synergistically, are involved in triggering synaptic weakening and loss in AD is poorly understood. Of relevance to the present study, at least two distinct forms of DHPG-LTD exist, which can be distinguished by the dependence on NMDARs (Palmer, Irving, Seabrook, Jane, & Collingridge, 1997; Sanderson, et al., 2022). In our current study, DHPG-LTD under control conditions was not affected by NMDAR antagonism. In contrast the oAβ-enhanced portion of the LTD totally depended on the activation of

NMDARs. As such, promotion of mGluR-NMDAR signaling during synaptic depression may be a key aspect of the shift in LTP/LTD balance and may constitute a trigger towards AD.

C3aR/C5aR signaling in AD

In both post-mortem human samples as well as numerous AD mouse models, many components of the complement cascade have been found to be upregulated (Hammond, Robinton, & Stevens, 2018; Schartz, 2020). Importantly, pharmacological and genetic manipulations of both C3aR and C5aR have provided evidence that both receptors are necessary for synaptic dysfunction and cognitive impairments in AD mouse models. SB290157, the C3aR antagonist used in our current study, reduces amyloid pathology in the APP/TTA mouse model (Lian, et al., 2016), and synapse engulfment by microglia and neuron loss are reduced in C3aR$^{-/-}$ mice in the PS19 tauopathy AD model (Litvinchuk, et al., 2018). Similarly, C5aR1 genetic ablation is protective in AD mouse models (Hernandez, et al., 2017; Carvalho, et al., 2022). Additionally, the C5aR antagonist used in our study, PMX205, improved amyloid pathologies, reactive gliosis, and behavioral outcomes in multiple AD mouse models (Fonseca, et al., 2009; Gomez-Arboledas, et al., 2022). However, the way in which glutamatergic synapses communicate with the complement cascade to cause this dysfunction was not understood. Our study provides insight into this communication, by demonstrating that oAβ enhances synaptic weakening, via NMDAR and mGlu$_5$R synergistic mechanisms that requires both C3aRs and C5aRs.

One important question to be addressed by future studies pertains to which arm of the complement cascade initiates the cleavage of C3 following mGluR-LTD in the presence of oAβ? Proteolytic cleavage of C3 generates both C3a, as well as

components of the C5 convertase which produces C5a. C3 conversion can be driven by either the classical arm, initiated by C1q, the lectin arm via mannose-binding lectin (MBL) or the alternative pathway through C3 "tickover" (Ricklin, Hajishengallis, Yang, & Lambris, 2010). Hong et al. (2016) found that the ability of oAβ to inhibit LTP requires the classical complement pathway, as co-incubation of hippocampal slices with oAβ and a function-blocking antibody of C1q restored LTP. Further, *in vivo* injections of oAβ into rat hippocampus upregulated C1q, and increased synapse phagocytosis by microglia (Bie, Wu, Foss, & Naguib, 2019). C1q upregulation by oAβ and subsequent synapse phagocytosis required group I mGluR signaling, and *in vivo* DHPG administration alone increased C1q. Finally, a recent study found that a silent allosteric modulator of mGlu$_5$R reduced C1q co-localization with PSD-95 and microglia engulfment of synapses in a mouse model of AD (Spurrier, 2022). As such, our findings suggest that NMDAR and C3aR/C5aR activation link C1q induction via mGluRs to downstream synaptic depression and elimination. Cleavage of C3 also generates the C3b fragment, which can covalently attach to synapses where it is converted to iC3b and binds to the phagocytic receptor CR3 (Ricklin, Hajishengallis, Yang, & Lambris, 2010). Genetic deletion of CR3 has also been found to spare synapses and improve memory in AD models (Hong, et al., 2016), indicating that phagocytosis through iC3b-CR3 binding is also critical for the deleterious effects of complement in pathology. As weaker synapses are targeted by complement for phagocytic removal (Schafer, et al., 2012), an intriguing question is whether synapse weakening by C3aR/C5aR activation also tags those synapses with iC3b for subsequent engulfment by microglia. CR3 can also impact synapses by non-phagocytic mechanisms in neuroinflammatory conditions. LPS and hypoxia activated CR3 which induces LTD of AMPAR transmission in the hippocampus (Zhang, et al.,

2014). Thus, microglia and the complement cascade have a multi-faceted role in

synaptic dysfunction in disease conditions, being critical for both changes to AMPAR-

mediated transmission strength as well as structural synapse elimination.

Relevance to other neurodegenerative diseases

Upregulation of complement cascade proteins and activity is not exclusive to AD and

has been detected in several other neurodegenerative diseases. For example, A1

reactive astrocytes, which express high levels of C3, have been identified in human

post-mortem tissue from patients with amyotrophic lateral sclerosis, Huntington's

disease, Parkinson's disease, and multiple sclerosis (Liddelow, et al., 2017). Further,

studies from *in vitro* and mouse models of these diseases have demonstrated the

necessity of C3 and/or C5 for their pathology (Woodruff, et al., 2008; McGeer, Lee, &

McGeer, 2017; Wang, Lee, Lee, Woodruff, & Noakes, 2017; Hammond, et al., 2020;

Bourel, et al., 2021; Chen, Hu, Ding, Du, & Hu, 2021; Gregersen, et al., 2021; Lopez-

Sanchez, et al., 2022). Interestingly, $mGlu_5R$ has separately been identified as an

emerging therapeutic target in these complement-associated neurodegenerative

diseases (Geurts, et al., 2003; Abd-Elrahman, et al., 2017; Bonifacino, et al., 2017;

Doria, et al., 2018; Bonifacino, et al., 2019; Farmer, et al., 2020; Su, Wang, Han, &

Shen, 2021; Zhang, et al., 2021; Azam, et al., 2022; Li, Colson, Abd-Elrahman, &

Ferguson, 2022). We therefore speculate that the $mGlu_5R$-NMDAR-C3aR/C5aR

signaling axis identified in our study may be a common feature of many brain

disorders.

Concluding Remarks

mGluRs, NMDARs, and the complement cascade have been independently studied

for their role in the pathogenesis of AD. Our study has found a novel link which places

glutamate receptor-innate immune interactions at the center of early synaptic

dysfunction mediated by oAβ, which is overproduced in AD. As such, a deeper

understanding of the cellular and molecular mechanisms of these interactions has the

potential to elucidate therapeutic targets which prevent synapse loss to halt AD

progression.

Acknowledgements

This work was supported by a CIHR (Canadian Institutes of Health Research)

Foundation Grant to GLC (#154276). GLC is the holder of the Krembil Family Chair in

Alzheimer's Research. We thank the trainee support provided by the C.R. Younger

Foundation. The work was supported by the MS Society of Canada and a donation

from Jon and Nancy Love. We are grateful for the long-standing support of the

Amalgamated Transit Union (ATU) Local 113 (Toronto, Canada). EWS was supported

by a CIHR Doctoral Award (Frederick Banting and Charles Best Canada Graduate

Scholarship, CGS-D) and an Ontario Graduate Scholarship (OGS).

Competing interests

The authors have no competing interests to declare.

References

Abd-Elrahman, K. S., Albaker, A., de Souza, J. M., Ribeiro, F. M., Schlossmacher, M.

G., Tiberi, M., & ... & Ferguson, S. S. (2020). Aβ oligomers induce

pathophysiological mGluR5 signaling in Alzheimer's disease model mice in a

sex-selective manner. *Science Signaling*, eabd2494.

Abd-Elrahman, K. S., Hamilton, A., Hutchinson, S. R., Liu, F., Russell, R. C., &

Ferguson, S. S. (2017). mGluR5 antagonism increases autophagy and

prevents disease progression in the zQ175 mouse model of Huntington's

disease. *Science Signaling*, eaan6387.

Afagh, A., Cummings, B. J., Cribbs, D. H., Cotman, C. W., & Tenner, A. J. (1996).

Localization and cell association of C1q in Alzheimer's disease brain.

Experimental Neurology, 22-32.

Ames, R. S., Lee, D., Foley, J. J., Jurewicz, A. J., Tornetta, M. A., Bautsch, W., & ... &

Sarau, H. M. (2001). Identification of a selective nonpeptide antagonist of the

anaphylatoxin C3a receptor that demonstrates antiinflammatory activity in

animal models. *The Journal of Immunology*, 6341-6348.

Anderson, W. W., & Collingridge, G. L. (2007). Capabilities of the WinLTP data

acquisition program extending beyond basic LTP experimental functions.

Journal of neuroscience methods, 346-256.

Azam, S., Jakaria, M., Kim, J., Ahn, J., Kim, I. S., & Choi, D. K. (2022). Group I mGluRs

in Therapy and Diagnosis of Parkinson's Disease: Focus on mGluR5 Subtype.

Biomedicines.

Beraldo, F. H., Ostapchenko, V. G., Caetano, F. A., Guimaraes, A. L., Ferretti, G. D.,

Daude, N., & ... & Prado, M. A. (2016). Regulation of Amyloid β Oligomer

Binding to Neurons and Neurotoxicity by the Prion Protein-mGluR5 Complex.

Journal of Biological Chemistry, 21945-21955.

Bie, B., Wu, J., Foss, J. F., & Naguib, M. (2019). Activation of mGluR1 mediates C1q-

dependent microglial phagocytosis of glutamatergic synapses in Alzheimer's

rodent models. *Molecular Neurobiology*, 5568-5585.

Bliss, T. V., & Collingridge, G. L. (1993). A synaptic model of memory: long-term potentiation in the hippocampus. *Nature*, 31-39.

Bonifacino, T., Cattaneo, L., Gallia, E., Puliti, A., Melone, M., Provenzano, F., & ... & Milanese, M. (2017). In-vivo effects of knocking-down metabotropic glutamate receptor 5 in the SOD1 G93A mouse model of amyotrophic lateral sclerosis. *Neuropharmacology*, 433-445.

Bonifacino, T., Provenzano, F., Gallia, E. R., Torazza, C., Bossi, S., & ... & Milanese, M. (2019). In-vivo genetic ablation of metabotropic glutamate receptor type 5 slows down disease progression in the SOD1 G93A mouse model of amyotrophic lateral sclerosis. *Neurobiology of Disease*, 79-92.

Bourel, J., Planche, V., Dubourdieu, N., Oliveira, A., Séré, A., Ducourneau, E. G., & ... & Tourdias, T. (2021). Complement C3 mediates early hippocampal neurodegeneration and memory impairment in experimental multiple sclerosis. *Neurobiology of Disease*, 105553.

Breijyeh, Z., & Karaman, R. (2020). Comprehensive Review on Alzheimer's Disease: Causes and Treatment. *Molecules*.

Brody, A. H., & Strittmatter, S. M. (2018). Synaptotoxic Signaling by Amyloid Beta Oligomers in Alzheimer's Disease Through Prion Protein and mGluR5. *Advances in Pharmacology*, 293-323.

Carvalho, K., Schartz, N. D., Balderrama-Gutierrez, G., Liang, H. Y., Chu, S. H., Selvan, P., & ... & Tenner, A. J. (2022). Modulation of C5a-C5aR1 signaling alters the dynamics of AD progression. *Journal of Neuroinflammation*.

Cavallucci, V., Berretta, N., Nobili, A., Nistico, R., Mercuri, N. B., & D'Amelio, M.

(2013). Calcineurin inhibition rescues early synaptic plasticity deficits in a

mouse model of Alzheimer's disease. *Neuromolecular Medicine*, 541-548.

Chen, M. M., Hu, Z. L., Ding, J. H., Du, R. H., & Hu, G. (2021). Astrocytic Kir6. 1

deletion aggravates neurodegeneration in the lipopolysaccharide-induced

mouse model of Parkinson's disease via astrocyte-neuron cross talk through

complement C3-C3R signaling. *Brain, Behaviour and Immunity*, 310-320.

Cline, E. N., Bicca, M. A., Viola, K. L., & Klein, W. L. (2018). The Amyloid-β Oligomer

Hypothesis: Beginning of the Third Decade. *Journal of Alzheimer's Disease*,

S567-S610.

Collingridge, G. L., Peineau, S., Howland, J. G., & Wang, Y. T. (2010). Long-term

depression in the CNS. *Nature Reviews Neuroscience*, 459-473.

Cullen, W. K., Suh, Y. H., Anwyl, R., & Rowan, M. J. (1997). Block of LTP in rat

hippocampus in vivo by beta-amyloid precursor protein fragments. *Neuroreport*,

3212-3217.

Dejanovic, B., Huntley, M. A., De Mazière, A., Meilandt, W. J., Wu, T., Srinivasan, K.,

& ... & Sheng, M. (2018). Changes in the synaptic proteome in tauopathy and

rescue of tau-induced synapse loss by C1q antibodies. *Neuron*, 1322-1336.

Doria, J. G., de Souza, J. M., Silva, F. R., Olmo, I. G., Carvalho, T. G., Alves-Silva, J.,

& ... & Ribeiro, F. M. (2018). The mGluR5 positive allosteric modulator

VU0409551 improves synaptic plasticity and memory of a mouse model of

Huntington's disease. *Journal of Neurochemistry*, 222-239.

Farmer, K., Abd-Elrahman, K. S., Derksen, A., Rowe, E. M., Thompson, A. M., Rudyk,

C. A., & ... & Hayley, S. (2020). mGluR5 Allosteric Modulation Promotes

Neurorecovery in a 6-OHDA-Toxicant Model of Parkinson's Disease. *Molecular*

Neurobiology, 1418-1431.

Fonseca, M. I., Ager, R. R., Chu, S. H., Yazan, O., Sanderson, S. D., LaFerla, F. M.,

& ... & Tenner, A. J. (2009). Treatment with a C5aR antagonist decreases

pathology and enhances behavioral performance in murine models of

Alzheimer's disease. *The Journal of Immunology*, 1375-1383.

Geurts, J. J., Wolswijk, G., Bö, L., Van Der Valk, P., Polman, C. H., Troost, D., &

Aronica, E. (2003). Altered expression patterns of group I and II metabotropic

glutamate receptors in multiple sclerosis. *Brain*, 1755-1766.

Gomez-Arboledas, A., Carvalho, K., Balderrama-Gutierrez, G., Chu, S. H., Liang, H.

Y., Schartz, N. D., & ... & Tenner, A. J. (2022). C5aR1 antagonism alters

microglial polarization and mitigates disease progression in a mouse model of

Alzheimer's disease. *Acta Neuropathologica Communications*.

Gregersen, E., Betzer, C., Kim, W. S., Kovacs, G., Reimer, L., Halliday, G. M., & ... &

Jensen, P. H. (2021). Alpha-synuclein activates the classical complement

pathway and mediates complement-dependent cell toxicity. *Journal of*

Neuroinflammation, 1-10.

Haas, L. T., Kostylev, M. A., & Strittmatter, S. M. (2014). Therapeutic molecules and

endogenous ligands regulate the interaction between brain cellular prion

protein (PrPC) and metabotropic glutamate receptor 5 (mGluR5). *Journal of*

Biological Chemistry, 28460-28477.

Haas, L. T., Salazar, S. V., Smith, L. M., Zhao, H. R., Cox, T. O., Herber, C. S., & ... &

Strittmatter, S. M. (2017). Silent allosteric modulation of mGluR5 maintains

glutamate signaling while rescuing Alzheimer's mouse phenotypes. *Cell*

Reports, 76-88.

Hamilton, A., Esseltine, J. L., DeVries, R. A., Cregan, S. P., & Ferguson, S. S. (2014).

Metabotropic glutamate receptor 5 knockout reduces cognitive impairment and

pathogenesis in a mouse model of Alzheimer's disease. *Molecular Brain*, 1-12.

Hamilton, A., Vasefi, M., Vander Tuin, C., McQuaid, R. J., & Anisman, H. F. (2016).

Chronic Pharmacological mGluR5 Inhibition Prevents Cognitive Impairment

and Reduces Pathogenesis in an Alzheimer Disease Mouse Model. *Cell*

Reports, 1859-1865.

Hammond, J. W., Bellizzi, M. J., Ware, C., Qiu, W. Q., Saminathan, P., Li, H., & ... &

Gelbard, H. A. (2020). Complement-dependent synapse loss and microgliosis

in a mouse model of multiple sclerosis. *Brain, Behaviour and Immunity*, 739-

750.

Hammond, T. R., Robinton, D., & Stevens, B. (2018). Microglia and the Brain:

Complementary Partners in Development and Disease. *Annu Rev Cell Dev*

Biol, 523-544.

Hasegawa, S., Sakuragi, S., Tominaga-Yoshino, K., & Ogura, A. (2015). Dendritic

spine dynamics leading to spine elimination after repeated inductions of LTD.

Scientific Reports, 1-6.

Hernandez, M. X., Jiang, S., Cole, T. A., Chu, S. H., Fonseca, M. I., Fang, M. J., & ...

& Tenner, A. J. (2017). Prevention of C5aR1 signaling delays microglial

inflammatory polarization, favors clearance pathways and suppresses cognitive loss. *Molecular Neurodegeneration*, 1-19.

Hong, S., Beja-Glasser, V. F., Nfonoyim, B. M., Frouin, A., Li, S., Ramakrishnan, S., & ... & Stevens, B. (2016). Complement and microglia mediate early synapse loss in Alzheimer mouse models. *Science*, 712-716.

Hsieh, H., Boehm, J., Sato, C., Iwatsubo, T., Tomita, T., Sisodia, S., & Malinow, R. (2006). AMPAR removal underlies Aβ-induced synaptic depression and dendritic spine loss. *Neuron*, 831-843.

Jackson, J., Jambrina, E., Li, J., Marston, H., Menzies, F., Phillips, K., & Gilmour, G. (2019). Targeting the synapse in Alzheimer's disease. *Frontiers in Neuroscience*.

Jo, J., Whitcomb, D. J., Olsen, K. M., Kerrigan, T. L., Lo, S. C., B.-M. G., . . . Cho, K. (2011). Aβ1–42 inhibition of LTP is mediated by a signaling pathway involving caspase-3, Akt1 and GSK-3β. *Nature Neuroscience*, 545-547.

Kim, J. H., Anwyl, R., Suh, Y. H., & Djamgoz, M. B. (2001). Use-dependent effects of amyloidogenic fragments of (beta)-amyloid precursor protein on synaptic plasticity in rat hippocampus in vivo. *Journal of Neuroscience*, 1327-1333.

Koffie, R. M., Hyman, B. T., & Spires-Jones, T. L. (2011). Alzheimer's disease: synapses gone cold. *Molecular Neurodegeneration*, 1-9.

Li, S. H., Colson, T. L., Abd-Elrahman, K. S., & Ferguson, S. S. (2022). Metabotropic Glutamate Receptor 5 Antagonism Reduces Pathology and Differentially Improves Symptoms in Male and Female Heterozygous zQ175 Huntington's Mice. *Frontiers in Molecular Neuroscience*.

Li, S., & Selkoe, D. J. (2020). A mechanistic hypothesis for the impairment of synaptic plasticity by soluble Aβ oligomers from Alzheimer's brain. *Journal of neurochemistry*, 583-597.

Li, S., Hong, S., Shepardson, N. E., Walsh, D. M., Shankar, G. M., & Selkoe, D. (2009). Soluble oligomers of amyloid β protein facilitate hippocampal long-term depression by disrupting neuronal glutamate uptake. *Neuron*, 788-801.

Li, S., Jin, M., Koeglsperger, T., Shepardson, N. E., Shankar, G. M., & Selkoe, D. J. (2011). Soluble Aβ oligomers inhibit long-term potentiation through a mechanism involving excessive activation of extrasynaptic NR2B-containing NMDA receptors. *Journal of Neuroscience*, 6627-6638.

Lian, H., Litvinchuk, A., Chiang, A. C., Aithmitti, N., Jankowsky, J. L., & Zheng, H. (2016). Astrocyte-Microglia Cross Talk through Complement Activation Modulates Amyloid Pathology in Mouse Models of Alzheimer's Disease. *Journal of Neuroscience*, 577-589.

Liddelow, S. A., Guttenplan, K. A., Clarke, L. E., Bennett, F. C., Bohlen, C. J., Schirmer, L., & ... & Barres, B. A. (2017). Neurotoxic reactive astrocytes are induced by activated microglia. *Nature*, 481-487.

Litvinchuk, A., Wan, Y. W., Swartzlander, D. B., Chen, F., Cole, A., Propson, N. E., & ... & Zheng, H. (2018). Complement C3aR inactivation attenuates tau pathology and reverses an immune network deregulated in tauopathy models and Alzheimer's disease. *Neuron*, 1337-1353.

Liu, J., Chang, L., Song, Y., Li, H., & Wu, Y. (2019). The Role of NMDA Receptors in Alzheimer's Disease. *Frontiers in neuroscience*.

Lopez-Sanchez, C., Poejo, J., Garcia-Lopez, V., Salazar, J., Garcia-Martinez, V., &

Gutierrez-Merino, C. (2022). Kaempferol prevents the activation of complement

C3 protein and the generation of reactive A1 astrocytes that mediate rat brain

degeneration induced by 3-nitropropionic acid. *Food and Chemical Toxicology*,

113017.

Lu, Y. M., Jia, Z., Janus, C., Henderson, J. T., Gerlai, R., Wojtowicz, J. M., & Roder,

J. C. (1997). Mice lacking metabotropic glutamate receptor 5 show impaired

learning and reduced CA1 long-term potentiation (LTP) but normal CA3 LTP.

Journal of Neuroscience, 5196-5205.

Lu, Y. M., Jia, Z., Janus, C., Henderson, J. T., Gerlai, R., Wojtowicz, J. M., & Roder,

J. C. (1997). Mice lacking metabotropic glutamate receptor 5 show impaired

learning and reduced CA1 long-term potentiation (LTP) but normal CA3 LTP.

Journal of Neuroscience, 5196-5205.

Lue, L. F., Kuo, Y. M., Roher, A. E., Brachova, L., Shen, Y., Sue, L., & ... & Rogers, J.

(1999). Soluble Amyloid β Peptide Concentration as a Predictor of Synaptic

Change in Alzheimer's Disease. *The American journal of pathology*, 853-862.

March, D. R., Proctor, L. M., Stoermer, M. J., Sbaglia, R., Abbenante, G., Reid, R. C.,

& ... & Fairlie, D. P. (2004). Potent cyclic antagonists of the complement C5a

receptor on human polymorphonuclear leukocytes. Relationships between

structures and activity. *Molecular Pharmacology*, 868-879.

McGeer, P. L., Lee, M., & McGeer, E. G. (2017). A review of human diseases caused

or exacerbated by aberrant complement activation. *Neurobiology of Aging*, 12-

22.

Nicoletti, F., Bockaert, J., Collingridge, G. L., Conn, P. J., Ferraguti, F., Schoepp, D.
D., & ... & Pin, J. P. (2011). Metabotropic glutamate receptors: from the
workbench to the bedside. *Neuropharmacology*, 1017-1041.

O'brien, R. J., & Wong, P. C. (2011). Amyloid Precursor Protein Processing and
Alzheimer's Disease. *Annual review of neuroscience*, 185-204.

Palmer, M. J., Irving, A. J., Seabrook, G. R., Jane, D. E., & Collingridge, G. L. (1997).
The group I mGlu receptor agonist DHPG induces a novel form of LTD in the
CA1 region of the hippocampus. *Neuropharmacology*, 1517-1532.

Parsons, C. G., Stöffler, A., & Danysz, W. (2007). Memantine: a NMDA receptor
antagonist that improves memory by restoration of homeostasis in the
glutamatergic system--too little activation is bad, too much is even worse.
Neuropharmacology, 699-723.

Perroy, J., Raynaud, F., Homburger, V., Rousset, M. C., Telley, L., Bockaert, J., &
Fagni, L. (2008). Direct interaction enables cross-talk between ionotropic and
group I metabotropic glutamate receptors. *Journal of Biological Chemistry*,
6799-6805.

Ricklin, D., Hajishengallis, G., Yang, K., & Lambris, J. D. (2010). Complement: a key
system for immune surveillance and homeostasis. *Nature Immunology*, 785-
797.

Sanderson, T. M., Ralph, L. T., Amici, M., Ng, A. N., Kaang, B. K., Zhuo, M., & ... &
Collingridge, G. L. (2022). Selective Recruitment of Presynaptic and
Postsynaptic Forms of mGluR-LTD. *Frontiers in Synaptic Neuroscience*.

Schafer, D., Lehrman, E., Kautzman, A., Koyama, R., Mardinly, A., Yamasaki, R., & ... & Stevens, B. (2012). Microglia sculpt postnatal neural circuits in an activity and complement-dependent manner. *Neuron*, 691-705.

Schartz, N. T. (2020). The good, the bad, and the opportunities of the complement system in neurodegenerative disease. *Journal of Neuroinflammation*, 1-25.

Selkoe, D. (2002). Alzheimer's disease is a synaptic failure. *Science*, 789-791.

Shankar, G. M., Bloodgood, B. L., Townsend, M., Walsh, D. M., Selkoe, D. J., & Sabatini, B. L. (2007). Natural oligomers of the Alzheimer amyloid-beta protein induce reversible synapse loss by modulating an NMDA-type glutamate receptor-dependent signaling pathway. *Journal of Neuroscience*, 2866-2875.

Shankar, G. M., Li, S., Mehta, T. H., Garcia-Munoz, A., Shepardson, N. E., Smith, I., & ... Selkoe, D. J. (2008). Amyloid-β protein dimers isolated directly from Alzheimer's brains impair synaptic plasticity and memory. *Nature Medicine*, 837-842.

Shinoda, Y., Tanaka, T., Tominaga-Yoshino, K., & Ogura, A. (2010). Persistent synapse loss induced by repetitive LTD in developing rat hippocampal neurons. *PLOS One*, e10390.

Spurrier, J. N. (2022). Reversal of synapse loss in Alzheimer mouse models by targeting mGluR5 to prevent synaptic tagging by C1Q. *Science Translational Medicine*.

Su, L. D., Wang, N., Han, J., & Shen, Y. (2021). Group 1 Metabotropic Glutamate Receptors in Neurological and Psychiatric Diseases: Mechanisms and Prospective. *Neuroscientist*.

Sumichika, H., Sakata, K., Sato, N., Takeshita, S., Ishibuchi, S., Nakamura, M., & ...

& Naka, Y. (2002). Identification of a potent and orally active non-peptide C5a

receptor antagonist. *Journal of Biological Chemistry*, 49403-49407.

Tu, J. C., Xiao, B., Naisbitt, S., Yuan, J. P., Petralia, R. S., Brakeman, P., & ... &

Worley, P. F. (1999). Coupling of mGluR/Homer and PSD-95 complexes by the

Shank family of postsynaptic density proteins. *Neuron*, 583-592.

Um, J. W., Kaufman, A. C., Kostylev, M., Heiss, J. K., Stagi, M., Takahashi, H., & ... &

Strittmatter, S. M. (2013). Metabotropic glutamate receptor 5 is a coreceptor for

Alzheimer aβ oligomer bound to cellular prion protein. *Neuron*, 887-902.

Volianskis, A., France, G., Jensen, M. S., Bortolotto, Z. A., Jane, D. E., & Collingridge,

G. L. (2015). Long-term potentiation and the role of N-methyl-D-aspartate

receptors. *Brain Research*, 5-16.

Walsh, D. M., Klyubin, I., Fadeeva, J. V., Cullen, W. K., Anwyl, R., Wolfe, M. S., & ...

& Selkoe, D. J. (2002). Naturally secreted oligomers of amyloid beta protein

potently inhibit hippocampal long-term potentiation in vivo. *Nature*, 535-539.

Wang, H. A., Lee, J. D., Lee, K. M., Woodruff, T. M., & Noakes, P. G. (2017).

Complement C5a-C5aR1 signalling drives skeletal muscle macrophage

recruitment in the hSOD1G93A mouse model of amyotrophic lateral sclerosis.

Skeletal Muscle, 1-11.

Wang, Q., Rowan, M. J., & Anwyl, R. (2004). Beta-amyloid-mediated inhibition of

NMDA receptor-dependent long-term potentiation induction involves activation

of microglia and stimulation of inducible nitric oxide synthase and superoxide.

Journal of Neuroscience, 6049-6056.

Wang, R., & Reddy, P. H. (2017). Role of glutamate and NMDA receptors in

Alzheimer's disease. *Journal of Alzheimer's Disease*, 1041-1048.

Whitcomb, D. J., Hogg, E. L., Regan, P., Piers, T., Narayan, P., Whitehead, G., & ...

& Cho, K. (2015). Intracellular oligomeric amyloid-beta rapidly regulates GluA1

subunit of AMPA receptor in the hippocampus. *Scientific Reports*, 1-12.

Wiegert, J. S., Pulin, M., Gee, C. E., & Oertner, T. G. (2018). The fate of hippocampal

synapses depends on the sequence of plasticity-inducing events. *Elife*, e39151.

Wiegert, J., & Oertner, T. (2013). Long-term depression triggers the selective

elimination of weakly integrated synapses. *Proceedings of the National*

Academy of Sciences, E4510-E4519.

Woodruff, T. M., Costantini, K. J., Crane, J. W., Atkin, J. D., Monk, P. N., Taylor, S.

M., & Noakes, P. G. (2008). The complement factor C5a contributes to

pathology in a rat model of amyotrophic lateral sclerosis. *The Journal of*

Immunology, 8727-8734.

Wu, T., Dejanovic, B., Gandham, V. D., Gogineni, A., Edmonds, R., Schauer, S., & ...

& Hanson, J. E. (2019). Complement C3 is activated in human AD brain and is

required for neurodegeneration in mouse models of amyloidosis and tauopathy.

Cell Reports, 2111-2123.

Wu, Y., Dissing-Olesen, L., MacVicar, B. A., & Stevens, B. (2015). Microglia: Dynamic

Mediators of Synapse Development and Plasticity. *Trends in immunology*, 605-

613.

Yasojima, K., Schwab, C., McGeer, E. G., & McGeer, P. L. (1999). Up-regulated

production and activation of the complement system in Alzheimer's disease

brain. *American Journal of Pathology*, 927-936.

Zhang, J., Malik, A., Choi, H. B., Ko, R. W., Dissing-Olesen, L., & MacVicar, B. A.

(2014). Microglial CR3 activation triggers long-term synaptic depression in the

hippocampus via NADPH oxidase. *Neuron*, 195-207.

Zhang, Y. N., Fan, J. K., Gu, L., Yang, H. M., Zhan, S. Q., & Zhang, H. (2021).

Metabotropic glutamate receptor 5 inhibits α-synuclein-induced microglia

inflammation to protect from neurotoxicity in Parkinson's disease. *Journal of*

Neuroinflammation, 1-24.

Figures

<u>Figure 1: oAβs enhances mGlu₅R-LTD at CA1 synapses.</u>

(**A-B**) Application of s-DHPG (50 µM, 10 min), induced LTD in -oAβ slices from mGlu$_5$R$^{+/+}$ mice

(21 ± 2 %, n = 9). Interleaved +oAβ slices showed an enhancement of DHPG-LTD (31 ± 4 %,

n= 9) with summary of LTD experiments in (**B**). p=0.032 *versus* -oAβ. Representative fEPSP

traces showing enhanced DHPG-LTD in +oAβ compared to -oAβ slices.

(**C-D**) In slices from littermate mGlu$_5$R$^{-/-}$ mice, DHPG failed to induce LTD responses in both

conditions (+oAβs = 2 ± 3 %, n=10; -oAβ= -3 ± 2 %, n=10) with summary in (**D**).

Representative fEPSP traces showing the absence of DHPG-LTD in both conditions.

(**E-F**) In a separate cohort from (**A**), pre-treatment of oAβs also resulted in an enhancement

of DHPG-LTD (34 ± 2 %, n=22) compared to the interleaved -oAβ slices (21 ± 2 %, n=22).

p=0.00031, +oAβ *versus* -oAβ.

(**G-H**) Bath application of L689,560 prevented the enhancement of DHPG-LTD in +oAβ slices

(26 ± 4 %, n=9) *versus* the interleaved -oAβ slices (22 ± 2 %, n=9).

Sample fEPSP traces are the mean of 4 consecutive responses at the indicated time points

(1-4). The number of slices (n) used for each experimental condition is shown. For each

experiment, sample traces before and after DHPG treatment are superimposed. Two-tailed t-

test was used in **B, D, F, H**. Scale bar: 10 ms / 0.2 mV. Source data are provided as a Source

Data file.

Figure 2: Complement C5aR and C3aR signaling is necessary for oAβs to enhance mGlu$_5$R-

LTD.

(**A-B**) In a third, separate cohort from Figure 1A and Figure 1E, pretreatment with oAβ

enhanced DHPG-LTD (43 ± 3 %, n=16) compared to the interleaved -oAβ slices (26 ± 3 %,

n=16). p=0.00064, +oAβ compared to -oAβ.

(**C-D**) Antagonism of C5aRs with PMX205 (0.3 µM) prevented the enhancement of DHPG-

LTD in +oAβ slices (35 ± 4 %, n=8) without affecting the level of DHPG-LTD in interleaved -

oAβ slices (29 ± 6 %, n=8).

(**E-F**) Antagonism of C5aRs with W54011 (0.3 µM) prevented the enhancement of DHPG-LTD

in +oAβ slices (32 ± 5 %, n=6) without affecting the level of DHPG-LTD in interleaved -oAβ

slices (29 ± 4 %, n=6).

(**G-H**) Antagonism of C3aRs with SB290157 (1 µM) prevented the enhancement of DHPG-

LTD in +oAβ slices (35 ± 3 %, n=8) without affecting the level of DHPG-LTD in interleaved -

oAβ slices (28 ± 6 %, n=8).

Sample fEPSP traces are the mean of 4 consecutive responses at the indicated time points.

The number of slices (n) used for each experimental condition is shown. Scale bar: 10 ms /

0.2 mV. Source data are provided as a Source Data file.

 Supplemental Figure 1: oAßs impair LTP at CA1 synapses.

(**A-B**) Delivery of a theta-burst protocol (TBS; see Methods), at the time indicated by the arrow,

induced LTP in -oAβ slices (51 ± 7 %, n = 28). LTP was impaired in the interleaved +oAβ

slices (17 ± 3 %, n = 28). Representative fEPSP traces for both conditions are shown to the

right of the time course plot (superimposed for sample experiment, before and after TBS, at

times indicated by the numbers 1-4). **B** Bar graph shows the level of LTP in each experiment.

Two-tailed t-test was used in **B**. p = 0.00002. Scale bar: 10 ms / 0.2 mV. Source data are

provided as a Source Data file.

(**C-E**) Basal synaptic transmission is similar in +oAβ slices and -oAβ slices, as assessed by

input-output curves. **C** Fiber volley (FV) amplitude *versus* stimulus intensity (line slopes: +oAβ

= -0.026 ± 0.003, n = 13; -oAβ = -0.030 ± 0.003, n = 13). **D** fEPSP slope *versus* FV amplitude

(line slopes: +oAβ = 2.517 ± 0.237, n = 13; black, *versus* -oAβ = 2.513 ± 0.362, n = 13; white).

E Representative fEPSP traces at different stimulus intensity (multiples of threshold) for both

conditions.