Pregranulosa cells upregulate cardiac troponin I *Tnni3* upon cell cycle resumption during primordial follicle activation

Güneş Taylor [a,g,1,*] & Emily R Frost [a,b,c,1], Brendan N Crow [a,b,c], Arthur Radley [d], Stefan Boeing [e,f], Christophe Galichet [a,h], Mark A Baker [b,c],
Jessie M Sutherland [b,c,2,*], Robin Lovell-Badge [a,2,*]

[1] These authors contributed equally and should be regarded as co-first authors
[2] These authors contributed equally and should be regarded as co-last authors

[a] Stem Cell Biology and Developmental Genetics Lab, The Francis Crick Institute, London, United Kingdom, NW1 1AT
[b] School of Biomedical Science & Pharmacy, University of Newcastle, Ring Road, Callaghan, New South Wales, Australia, 2308
[c] Hunter Medical Research Institute, New Lambton Heights, New South Wales, Australia, 2305
[d] Developmental Dynamics Lab, The Francis Crick Institute, London, United Kingdom, NW1 1AT
[e] Bioinformatics and Biostatistics Facility, The Francis Crick Institute, London, United Kingdom, NW1 1AT
[f] Scientific Computing – Digital Development Team, The Francis Crick Institute, London, United Kingdom, NW1 1AT
[g] Centre of Reproductive Health, Institute of Regeneration and Repair, University of Edinburgh, Scotland, EH16 4UU
[h] Neurobiological Research Facility, Sainsbury Wellcome Centre, London, United Kingdom, W1T 4JG

* Corresponding authors:
Dr Güneş Taylor, Centre of Reproductive Health, University of Edinburgh, Scotland, EH16 4UU, gunes.taylor@ed.ac.uk
Prof Robin Lovell-Badge, Stem Cell Biology and Developmental Genetics Lab, The Francis Crick Institute, London, United Kingdom, NW1 1AT, robin.lovell-badge@crick.ac.uk
Dr Jessie Sutherland, School of Biomedical Science & Pharmacy, University of Newcastle, Ring Road, Callaghan, New South Wales, Australia, 2308, jessie.sutherland@newcastle.edu.au

Keywords: ovary, oocyte, infertility, granulosa cells, cell cycle, cdkn1b, p27, TNNI3, primordial follicle activation, entropy sorting

Abstract

Primordial follicles are quiescent ovarian structures comprised of a single oocyte surrounded by a layer of somatic supporting pregranulosa cells. Primordial follicle activation is the first step towards oocyte maturation and, ultimately, ovulation. As there is a finite number of primordial follicles within the ovary, their rate of activation is a critical parameter of the female reproductive lifespan. Follicle activation is characterised by morphological and molecular changes in both pregranulosa cells and oocytes. However, the signal initiating activation, and even the cell-type responding to this signal, remain unidentified. Through single-cell transcriptomic profiling of neonatal mouse ovaries we identify a putative gene expression signature of activating pregranulosa cells. We find precocious expression of this putative pregranulosa cell signature in the *Cdkn1b*$^{-/-}$ or p27^{kip1}-null model, which has ubiquitous follicle activation neonatally. We confirm that the protein product of one of these genes, *Tnni3,* is present within mouse granulosa cells, which has also been described recently for human granulosa cells. We demonstrate that expression of cardiac troponin I (TNNI3), an actin-binding protein, is upregulated upon cell-cycle resumption in activating pregranulosa cells. This indicates pregranulosa cells likely initiate primordial follicle activation and support the broader hypothesis that changes in cell tension are involved in triggering follicle activation.

Introduction

Critical parameters of female reproductive lifespans are the finite number of quiescent primordial follicles within their ovaries and the rate of primordial follicle activation for maturation and eventual ovulation. Primordial follicles are formed during gestation and consist of an oocyte with a single layer of surrounding pregranulosa cells (1, 2). Once primordial follicles activate, the pregranulosa cells become steroidogenic granulosa cells and provide structural and metabolic support to the oocyte throughout folliculogenesis (3). Primordial follicles are initially formed and distributed throughout the ovary. However, typically during late foetal development or neonatally in mammals, the majority of follicles within the central ovarian medulla are lost subsequent to a synchronised activation process termed first wave activation (4, 5). In rodents, the sex steroids produced by supporting cells during first wave activation are hypothesised to be important for establishing aspects of puberty and the ovarian–hypothalamic–pituitary axis (4, 6, 7). Following first wave activation, the remaining primordial follicles reside in the outer ovarian cortex and are cyclically activated for eventual ovulation or atresia. The establishment of this 'ovarian reserve' of cortical primordial follicles, and their gradual release from it, sustains fertility over several decades in humans and many months in mice (8).

Consequently, stringent control of follicle activation is required to ensure that the ovarian reserve persists for the entire reproductive lifespan. To date, key indicators of primordial follicle activation are changes in granulosa cell morphology (from squamous to cuboidal) and re-entry into the cell-cycle and increased oocyte size (9-12). The morphological changes to pregranulosa cells precede that of the oocyte, suggesting that primordial follicle activation may be initiated by the pregranulosa cells (13-15). While many pathways have roles in follicle activation, including PI3K, mTOR, KITL, TGFB and Hippo signalling (9, 10, 16-19), the molecular signals initiating follicle activation remain unresolved. Primordial follicle activation is challenging to interrogate due to the asynchronous nature of activation throughout the adult ovarian cortex. Advances in single-cell RNA sequencing (scRNAseq) have accelerated our understanding of many ovarian processes in the mouse (20-22), however the difficulty of knowing which cortical follicle is initiating activation has impeded the identification of a specific gene or process involved in the earliest steps of this critical process.

Here we leveraged the abundance of activating follicles during first wave activation to investigate the transcriptomic changes of pregranulosa cells in neonatal mice. Using scRNAseq we identify a putative signature of activating pregranulosa cells. Importantly, we find this signature of activating pregranulosa cells is observed precociously at E18.5 in a mouse mutant with dysregulated first wave activation (*Cdkn1b*$^{-/-}$ or *Cdkn1b* / p27^{kip1}-null) (23). We confirm presence of one of these putative activation genes, TNNI3, in granulosa cells during first wave activation in neonatal wildtype ovaries. In mutant *Cdkn1b*$^{-/-}$ ovaries we find TNNI3 expressing granulosa cells appear precociously at E18.5. These data support the notion that pregranulosa cell maturation into granulosa cells is a

key driver of primordial follicle activation and integrates actin-binding protein TNNI3 and cell-cycle regulator *Cdkn1b* / p27[kip1] within the molecular mechanism driving this important cell transition.

Results

Identification of putative regulators of primordial follicle activation within pregranulosa cells.

To profile the transcriptomic dynamics of first wave activation in wildtype (WT) mice, scRNAseq was conducted on ovaries at embryonic day 18.5 (E18.5), post-natal day 4 (PD4), and post-natal day 7 (PD7) (Figure 1a). These timepoints captured prior to, during and after first wave activation respectively. Using our previously described protocol (24), we enriched for the somatic cell compartment in the single-cell suspension. The dataset contained 24,810 cells in total with 8,210 cells at E18.5, 7,587 cells at PD4 and 9,013 cells at PD7. Data from all three timepoints were integrated and, subsequent to standard highly variable gene selection, uniform manifold approximation and projection (UMAP) plots were generated with 18 distinct clusters. The six major cell types (granulosa cells Gc_1 – Gc_5), mesenchymal cells (Mc_1 – Mc_5), oocytes (Oo_1), epithelial cells (Ep_1 – Ep_2), immune cells (Im_1) and blood cells (Bl_1 – Bl_2)) in the ovary were annotated based on their expression of established cell type markers (Figure 1a and Supplemental Figure 1a) (5).

We focused on establishing differences between Gc_1 through Gc_4 as cluster Gc_5 had high expression of *Pax8* (Figure 1b), a recently reported marker of supporting-like cells from the *rete ovarii* (25). Using a detailed panel of pregranulosa and granulosa cell markers, we classified Gc_1 as an embryonic population of epithelial-derived pregranulosa cells due to their expression of *Lgr5*, *Gng13* and *Lgas7* (Figure 1b), and the decreased cell numbers in Gc_1 between E18.5 and PD7 (Figure 1a) (4, 5, 26, 27). We identified Gc_2 as a population of pregranulosa cells derived from the Nr5a1+ somatic cell lineage based on their high expression of *Cdkn1b* and *Foxl2* alongside low expression of *Hsd3b1* and *Hmgcs2* (Figure 1b) (5). Interestingly, Gc_3 had similar gene expression to Gc_2 (shown by the expression of *Foxl2* and *Hmgcs2* amongst others), but with elevated expression of well-established markers of activated granulosa cells like *Kitl* and *Fst* (Figure 1b) (10, 28). This suggested that Gc_3 is a population of pregranulosa cells that have initiated primordial follicle activation. Cluster Gc_4 expressed mature granulosa cell marker genes including *Amhr2* and *Hmgcs2* and showed high expression of proliferation genes (Figure 1a, Figure 1b) (9, 29). When interrogated by timepoint and cell cycle phase, we observed that clusters Gc_1, Gc_2 and Gc_3 were present from E18.5, while Gc_4 only appeared at PD4 during first wave activation and was in the proliferative G2M / S phase of the cell cycle (Supplemental Figure 1b). Overall, two distinct gene expression dynamics emerged: genes that increase in expression during activation and persist in

mature granulosa cells (Gc_3 and Gc_4), and genes that decreased upon entry into the activating pregranulosa cell pool (Gc_1 and Gc_2) (Figure 1b).

To interrogate the granulosa cell transcriptome more closely, we subclustered clusters Gc_1- Gc_5 and Ep_1, the surface epithelium from which many supporting cells are derived (Supplemental Figure 1b) (30). Differential expression analysis showed there was significant overlap in gene expression across all Gc clusters, and highlighted that distinct marker genes exist for Gc_4 but not the remaining Gc clusters (Supplemental Figure 1c, Dataset S1-4). As an alternative to the standard differential expression analysis for feature selection we applied a new approach called continuous Entropy Sort Feature Weighting (cESFW) (31) to the Ep_1 and Gc_1 – Gc_5 scRNAseq clusters. Subsetting the scRNA-seq data to the 4376 top ranked genes identified by cESFW generated a higher resolution UMAP of granulosa cell development (Figure 1c). Unsupervised Kmeans clustering revealed 14 distinct clusters (Supplemental Figure 1d). To compare against the differential expression analysis, we manually assigned the cESFW Kmeans clusters to the cell type annotations identified using the conventional scRNAseq workflow (Figure 1c). We then used entropy sorting (ES) (31) to calculate the entropy sort score (ESS) correlation metric to get ranked gene lists for each of these clusters (Figure 1d, Dataset S5). The top ten markers of each cluster differed from those found by standard differential gene expression analysis, but also revealed Gc_3 markers are maintained in Gc_4 from PD4 onwards. As such, cESFW feature selection separates the transcriptional signature of the onset of pregranulosa cell activation (e.g. *Nr5a2*, *Fam13a* and *Tnni3*) from that of mature granulosa cells only (e.g. *Aurka*, *Kif23*).

Transcription of *Tnni3* is upregulated in *Cdkn1b*[-/-] mice with accelerated primordial follicle activation.

To establish the functional involvement of the genes putatively underlying pregranulosa cell activation we queried our activation signature genes in an established model of precocious and aberrant follicle activation, *Cdkn1b*[-/-] (*Cdkn1b* / p27[kip1]-null) mice. *Cdkn1b*/p27[kip1] is a cyclin dependent kinase inhibitor and therefore blocks the cell cycle. *Cdkn1b*[-/-] mice show an uncontrolled activation of follicles perinatally, inducing infertility prior to sexual maturation (23, 32). We performed bulk RNA-sequencing of control and *Cdkn1b*[-/-] ovaries at three timepoints: E13.5, E18.5 and PD8 (Supplemental Figure 2b, Dataset S6). As ovarian supporting cells express *Cdkn1b*/ p27[kip1] prior to expression of the pregranulosa cell specification marker *Foxl2,* the E13.5 timepoint was chosen to confirm pregranulosa specification is unperturbed by the loss of *Cdkn1b* (12, 33). The E18.5 timepoint was chosen as it is prior to first wave activation in WT mice and is the first timepoint of our scRNAseq timecourse. First wave activation has concluded by PD8 and was investigated in the previous report of *Cdkn1b* / p27[kip1]-null mice and aligns with our final scRNAseq timepoint.

We found just 4 differentially expressed genes at E13.5 (p < 0.05), confirming pregranulosa cell specification is essentially unperturbed. As previously reported, larger numbers of primary and secondary follicles were observed in $Cdkn1b^{-/-}$ ovaries at PD8 (Figure 2a) and we found 746 differentially expressed genes at PD8 in our bulk RNAseq data (p < 0.05 Dataset S6). Contrary to the previous report, we observed cuboidal granulosa cells in $Cdkn1b^{-/-}$ ovaries already at E18.5 (Supplemental Figure 2a) (23). This was further supported by the RNAseq data showing 896 upregulated and 520 downregulated differentially expressed genes in $Cdkn1b^{-/-}$ ovaries compared to WT controls at E18.5 (p < 0.05) (Figure 2b, Supplemental Figure 2c and Dataset S6). This suggested that absence of $Cdkn1b$ was causing precocious first wave activation around the time of germ cell nest breakdown and primordial follicle formation.

Since $Cdkn1b$ / p27^{kip1} is exclusively expressed in pregranulosa cells at this developmental stage, we hypothesised that the specific maturation of pregranulosa cells to granulosa cells was driving this precocious primordial follicle activation. Therefore, we compared the 1416 differentially expressed genes from the bulk RNA-sequencing of E18.5 $Cdkn1b^{-/-}$ ovaries with the genes expressed in clusters Gc_1 to Gc_4 from our scRNAseq timecourse (Figure 2c). This revealed that indeed the majority of differentially expressed genes in $Cdkn1b^{-/-}$ ovaries at E18.5 were expressed in the activating pregranulosa cell cluster Gc_3 and granulosa cell cluster Gc_4 in WT ovaries. Importantly, the most statistically significantly upregulated differentially expressed gene at E18.5 in $Cdkn1b^{-/-}$ ovaries was $Tnni3$ (1.70logFC, padj=1.06E-38), a gene also identified by the ESS as one of the top ten markers of Gc_3 activating pregranulosa cells (Figure 1d). $Tnni3$ expression is restricted to Gc_3 and Gc_4 cell clusters, contrary to the high $Cdkn1b$ expression in clusters Gc_1 and Gc_2 (Figure 2d). Together, this suggested engaging the cell-cycle via reduction of $Cdkn1b$ expression and increased $Tnni3$ expression may drive pregranulosa cell maturation into granulosa cells in the mouse.

TNNI3 protein found in granulosa cells of follicles undergoing activation.

$Tnni3$ encodes for an actin-binding protein involved in the troponin complex in cardiac tissues and is also present in the granulosa cells of maturing follicles in humans (34, 35). However, these studies looked at late-stage growing follicles and not the transition of pregranulosa to granulosa cells. Considering our data linking $Tnni3$ to supporting cell activation, we probed WT mouse ovaries at E18.5, PD4 and PD7 for protein expression. TNNI3 staining was low to undetectable within pregranulosa cells at E18.5 (Figure 3a). However, at PD4 and PD7, TNNI3 staining was apparent within granulosa cells (Figure 3a). Importantly, transitioning follicles contain both cuboidal granulosa cells and pregranulosa cells. We were able to capture follicles containing both cell states and show TNNI3 staining only in the activated granulosa cells of a single follicle (Figure 3a). TNNI3 staining in the granulosa cells was largely localised around the outer edge of granulosa cells as previously reported in the human (36) (Figure 3b). We found the relative presence of TNNI3 was significantly greater (Welch's T-test p=<0.0001) in granulosa cells of

activated follicles when compared to the pregranulosa cells of primordial follicles by mean fluorescent intensity (Figure 3c). We also confirmed TNNI3 staining in granulosa cells within activated follicles in cortically derived adult mouse ovaries (Supplemental Figure 3a). This suggests TNNI3 upregulation in activating pregranulosa cells occurs both in medullary follicles activated during first wave activation and in the cortical follicles of the adult. TNNI3 staining was also observed in oocytes of growing follicles but minimal staining was seen in the oocytes of primordial follicles (Figure 3a). This was reflected in the scRNAseq data with the oocyte cluster showing transcription of TNNI3 at the PD4 and PD7 timepoints (average log10 expression at E18.5 = 0.02, PD4 = 0.34 and PD7 = 0.32).

Taken together, our data show pregranulosa cells upregulate *Tnni3* expression upon cell cycle resumption, in a *Cdkn1b*/p27^{kip1} responsive manner, during primordial follicle activation (Figure 3c).

Discussion

Primordial follicle activation is critical to the establishment and duration of mammalian female fertility. Understanding the molecular mechanisms underpinning activation holds the potential to improve reproductive longevity, assist in the diagnosis of infertility and develop targeted treatments for patients with ovarian dysfunction. To interrogate this mechanism, we generated a scRNAseq dataset of first-wave primordial follicle activation. We focused on four distinct clusters of granulosa cells that undergo morphological changes associated with initiation of primordial follicle activation. One cluster, Gc_3, represented a distinct population of activating pregranulosa cells, which was interrogated through entropy sorting (31). This identified several strong candidate genes for the initiation of follicle activation, including *Tnni3*. Confirmation that *Tnni3* may play a functional role in this process was elucidated using the *Cdkn1b*$^{-/-}$ mouse line, a model for accelerated primordial follicle activation, where *Tnni3* was precociously upregulated. These data implicated the pregranulosa cell gene, *Tnni3,* as a potentially novel marker of primordial follicle activation.

The ability to track individual cell transcriptomes makes scRNAseq a powerful tool in understanding primordial follicle activation. Several studies have now used scRNAseq both in the cycling (21, 35) and ageing ovary (37, 38), where the focus is commonly on the oocyte. The transcriptomic datasets we provide here highlight the relevance of somatic cells in follicle activation. Our data provide further evidence that follicle activation is likely initiated within the pregranulosa cells (15). This is in agreement with studies illustrating activation changes in response to mechanical pressure (39) and metabolic changes (40). Consulting this dataset alongside previously published scRNAseq studies of the developing ovary will create a more comprehensive picture of granulosa cell changes across germ cell nest breakdown, primordial follicle formation and activation (5, 20, 21, 33, 41-43). Potential opportunities also exist to correlate this data with human

ovary scRNAseq studies and those from other species to discover how mechanisms of follicle activation are conserved (22).

The bioinformatic approaches within the study provide new opportunities to discriminate transcriptionally similar cell types. Herein, we utilised two complementary approaches for dissecting activating pregranulosa cells from activated granulosa cells, namely Entropy Sorting (ES) and combining scRNAseq with bulk RNAseq from established infertility mouse models. As in previous uses of ES (31, 44, 45), our work demonstrates how the mathematical framework of ES can elucidate subtle changes in transcriptional profiles along developmental trajectories that conventional bioinformatics approaches may struggle to identify. Other developmental questions with dynamic and terminal trajectories could benefit from the application of this approach as a way of more accurately selecting gene targets underpinning development. The second bioinformatic approach was to functionally interrogate genes identified via ES using an infertile mouse model. We utilised a $Cdkn1b^{-/-}$ mouse line that has an accelerated primordial follicle activation phenotype in the neonatal mouse ovary. Contrary to previous reports, our transcriptomic analysis of these ovaries revealed distinct gene expression changes as early as E18.5, and we also find apparent morphological changes of supporting cells in mutant ovaries. Genes upregulated in $Cdkn1b^{-/-}$ ovaries were part of the novel gene expression signature of activating pre-granulosa cells identified in our scRNAseq data, along with mature granulosa cell genes. The combination of scRNAseq data with bulk RNA-sequencing supports and strengthens our characterisation of the genes involved in primordial follicle activation, and as such we highlight this methodology as an under-utilised approach. This type of comparison has previously been informative in studies on embryonic somatic gonadal cells (46). In the context of this study, the combination of these bioinformatic tools was effective in identifying TNNI3 as a novel putative regulator of pregranulosa-driven primordial follicle activation.

$Tnni3$ is an 8-exon gene located on chromosome 7 that encodes a 211-residue protein. TNNI3 functions in cardiac tissue as the inhibitory part of the three-subunit troponin complex. TNNI3 is thought to act as a molecular switch which regulates the interaction of actin and myosin in response to calcium. The function of TNNI3 in the ovary is currently unknown. A recent study of human growing follicles reported TNNI3 expression by granulosa cells and noted a characteristic ring-shaped staining pattern at the outer edge of these granulosa cells (34). This study verifies the presence of TNNI3 protein in mouse granulosa cells of primary and growing follicles and finds a similar staining pattern. Taken with our findings that $Tnni3$ expression is upregulated in the granulosa cells of activating follicles, these data suggest a possible role of TNNI3 in inducing and maintaining the cuboidal shape of pregranulosa cells. Another explanation may be a requirement for TNNI3 to interact with protein kinase C (PKC), an established primordial follicle regulator, as TNNI3 is known to be phosphorylated by PKC in the mouse myocardium (47-49). Other evidence that TNNI3 may interact with other known follicle activation pathways exists where $Tnni3$ is upregulated in the absence of $Foxl2$ in mice, however this is indirect as FOXL2 doesn't appear to bind to the promoter of $Tnni3$ in mouse ovaries

one week after birth (50, 51). Future work may provide a link between *Tnni3* and other biological pathways, however conditional experiments will be required as *Tnni3*[-/-] mice die perinatally from cardiac complications (52).

Questions remain as to whether the loss of cell-cycle inhibition in pregranulosa cells solely drives the precocious primordial follicle activation in *Cdkn1b*[-/-] mutants. Evidence from a closer examination of *Cdkn1b*[-/-] mutants suggests that germ cell nest breakdown and follicle formation may also be affected, as these mice form multi-oocyte follicles at higher rates than their control littermates (32). In this study, the phenotype of *Cdkn1b*[-/-] mutants might also be exacerbated by the loss of p27[kip1] in the oocytes, where expression comes on postnatally in wildtype ovaries. A conditional approach deleting *Cdkn1b* in pregranulosa cells specifically would provide a clear answer if primordial follicle activation is initiated in the pregranulosa cells and dysregulated by the absence of p27[kip1]. Regardless, it is clear from this body of work that further interrogation is necessary to find the master regulators of follicle activation.

In conclusion, this study presents transcriptomic datasets to interrogate primordial follicle activation in the mouse. We identify a gene expression signature within pregranulosa cells that are undergoing primordial follicle activation and highlight p27[kip1]-dependent genes within this signature. We utilise a recently described mathematical framework, Entropy Sorting, to discriminate activation-specific pregranulosa genes as high confidence imputed regulators of primordial follicle activation. Using this method, combined with an established infertile mouse model, we reveal a likely role for *Tnni3* as a regulator of activation within pregranulosa cells. Our data provide further evidence that it is the pregranulosa cell changes and not the oocytes, that underpin primordial follicle activation. Moreover, we hypothesise that the genes identified in Gc_3 by the Entropy Sort Score (ESS) are a critical part of the molecular control initiating this process. Our datasets, along with the bioinformatic approaches we have employed, will be valuable resources for the interrogation of mechanisms underlying mammalian follicle activation and female fertility.

Materials and Methods

Mice
All experiments carried out on mice were approved under the UK Animal (scientific procedures) Act 1986 (Project license 70/8560 and PP8826065). C57BL/6J females (RRID: IMSR_JAX:000664) were supplied by the Francis Crick Institute Biological Research Facility. *Cdkn1b*[tm1Mlf] (*Cdkn1b*[-/-]) (RRID: IMSR_JAX:002781) (53) were maintained on a C57BL/6J/CBA mixed background and mated heterozygously. The Cdkn1b[tm1Mlf] Foxl2[tm1(EGFP)Rlb] (Cdkn1b[-/-] ; FoxL2[P2A-eGFP/+]) were likewise maintained on a C57BL/6J/CBA mixed background and mated homozygously for the FoxL2[P2A-eGFP] allele and heterozygously for the Cdkn1b[-/-] allele (51).

Tissue dissociation and scRNAseq library preparation
Ovarian tissue was dissociated as previously described (24). After mechanical dissociation, and filtration through a 20 μm pre-separation filter, approximately 10,000 live cells were used to generate libraries using the Chromium Single Cell 3' kit v3.1, according to the manufacturer's instructions (54). The libraries were quantified using the TapeStation (Agilent) to confirm quantity and purity, before sequencing on an Hiseq4000 (Illumina). Sequencing outputs were processed using the 10x CellRanger (version.3.0.2) to generate single-cell count data for each time point, using a mouse reference index provided by 10x Genomics (refdata-cellranger-mm10v3.0).

Tissue dissociation and bulkRNA sequencing library preparation
Ovaries from *Cdkn1b*[+/+] and *Cdkn1b*[-/-] mice were collected at E13.5, E18.5 and PD8. A single ovary was dissociated as described above and RNA was extracted using the RNeasy Plus Micro kit (Qiagen). RNA quality was assessed the TapeStation (Agilent). Libraries were constructed using the NEBNext Low Input RNA Library Prep Kit (E6420L, New England Biolabs) according to the manufacturer's instructions. All libraries were quantified using the TapeStation (Agilent) and sequenced on an Hiseq400 (Illumina) to achieve an average of 25 million reads per sample.

Single-cell RNA sequencing analysis
Raw single-cell sequencing data from all three timepoints were processed using the 10x CellRanger pipeline (10x Genomics) and analysed with the Seurat package in R (55). All cells with fewer than 1500 RNA features were removed. Principal component analysis was performed on variable genes and 20 principal components were used to generate the UMAP plots using the default method from the Seurat package. Feature selection for UMAP generation was done using standard highly variable selection approaches (30) and continuous Entropy Sort Feature Weighting as previously described (31). The cESFW workflow used to identify the set of 4376 genes used to produce a high resolution UMAP embedding of pregranulosa cell differentiation may be found at the following GitHub repository, https://github.com/aradley/Taylor_Pregranulosa_Cells.

Bulk RNA sequencing analysis

Libraries were sequenced on an Illumina HiSeq 4000 machine. The 'Trim Galore!' utility version 0.4.2 was used to remove sequencing adaptors and to trim individual reads with the q-parameter set to 20 (https://www.bioinformatics.babraham.ac.uk/projects/trim_galore/). The sequencing reads were then aligned to the mouse genome and transcriptome (Ensembl GRCm38release-89) using RSEM version 1.3.0 (56) in conjunction with the STAR aligner version 2.5.2 (57). The sequencing quality of individual samples was assessed using FASTQC version 0.11.5 (https://www.bioinformatics.babraham.ac.uk/projects/fastqc/) and RNA-SeQC version 1.1.8 (58). Differential gene expression was determined using the R-bioconductor package DESeq2 version 1.14.1 (59, 60). Log fold changes above 1 and below -1 with a padj of less than 0.05 were considered significant.

Data availability
The data supporting the findings of this study are available in the main text and the supplementary materials. Datasets have been submitted to NCBI GEO (GSE246098).

Histology staining
Ovaries for hematoxylin and eosin (H&E) staining were fixed in chilled 4% (w/v) paraformaldehyde at 4 °C for 1 to 2 hrs. Following fixation, ovaries were washed in PBS 3 times for 1 hr each, before cryoprotection in 30% (w/v) sucrose in PBS overnight. The ovaries were then embedded in OCT mounting medium, frozen on an ethanol/dry ice slurry and stored at -80 °C. Samples were cryosectioned at 10 μm and sections were placed on Superfrost Plus glass slides and air dried for 5 minutes. H&E staining was performed by the Francis Crick Institute Experimental Histopathology facility. H&E images were taken at 10x magnification and 40x magnification, with scale bars of 100 μm and 25 μm respectively.

Cryosectioning ovaries
Ovaries were collected from Foxl2$^{P2A-eGFP}$ mice (51), fixed in 4% paraformaldehyde solution on ice for 30 mins followed by a further 1hr incubation at room temperature. Samples were subsequently cryopreserved 30% sucrose in PBS overnight at 4°C and embedded in OCT media prior to cryosectioning. OCT-embedded ovaries were serially sectioned into 12μm sections.

Immunofluorescent staining of cryosectioned ovaries
Cryosectioned ovaries were washed in PBS, blocked with 10% donkey serum (blocking buffer) for 2hrs at room temperature. The primary antibody against TNNI3 (ab47003), FOXL2 (AB5096), and DDX4 (AB13840) were incubated at 4°C overnight at 1:200 dilution blocking buffer. Subsequent to washes, the secondary antibody (Donkey anti-Rabbit IgG Alexa Fluor 488 and Donkey anti-Goat Alexa Fluor 647) was diluted 1:500 in blocking buffer and incubated at room temperature for 2hrs. The nanobody against GFP (GBA-647n) was at 1:200 in blocking buffer for both primary and secondary incubations. Sections were then counterstained for 30 seconds with 4',6-diamidino-2-phenylindole (DAPI) diluted 1:10000. Sections were mounted with Aqua-Poly/Mount and covered

with glass coverslips. Imaging was done using a Zeiss Invert880 with Airyscan 2019 confocal microscope.

Mean fluorescent intensity

Quantification of relative TNNI3 presence in granulosa vs pregranulosa cells was calculated using mean fluorescent intensity (MFI) imageJ analysis software (61). Granulosa and pregranulosa cells were segmented using free hand drawing tool on the green (FOXL2[p2A-eGFP]) channel to segment the whole cell body. MFI was calculated by taking the mean pixel intensity of the TNNI3 channel for both granulosa (total of 372) and pregranulosa cells (total of 144) for 3 biological replicates of PD7 *Foxl2*[P2A-eGFP] mouse ovaries. Each biological replicate was normalised through feature scaling to allow unbiased comparison between replicates and results were analysed via Welches T-test.

Acknowledgements and Funding Sources

Acknowledgements

This work was supported by the Francis Crick Institute, which receives its core funding from Cancer Research UK (CC2116), the UK Medical Research Council (CC2116), and the Wellcome Trust (CC2116). We acknowledge the generous financial assistance to R.L.B. by the UK Biotechnology and Biological Sciences Research Council (BB/ N018680/1). We also gratefully acknowledge the financial assistance to J.M.S. and M.A.B. by the Australian National Health and Medical Research Council (G1600095) and the Hunter Medical Research Institute (G1501433 and G1801335). E.R.F. and B.N.C. are recipients of Australian Government Research Training Program Scholarships. We thank the Advanced Sequencing Facility at the Francis Crick Institute for their help in performing the sequencing studies in this manuscript. We would like to thank members of the Lovell-Badge and Sutherland laboratories for their comments on the manuscript.

Author contributions

G.T. and E.R.F. conceived the study and wrote the manuscript. J.M.S., C.G., M.A.B. and R.L.B. provided expertise and feedback on the study design. All experiments were performed by E.R.F., G.T., B.N.C. and C.G.. RNAseq and scRNAseq analyses were performed by A.R. and S.B.. All the authors critically revised the manuscript. R.L.B. and J.M.S. provided resources for this work.

Declaration of interests

The authors declare no competing interests.

References

1.   M. E. Pepling, A. C. Spradling, Mouse ovarian germ cell cysts undergo programmed breakdown to form primordial follicles. *Developmental biology* 234, 339-351 (2001).

2.   C. Tingen, A. Kim, T. K. Woodruff, The primordial pool of follicles and nest breakdown in mammalian ovaries. *Molecular human reproduction* 15, 795-803 (2009).

3.   R. Li, D. F. Albertini, The road to maturation: somatic cell interaction and self-organization of the mammalian oocyte. *Nat. Rev. Mol. Cell Biol.* 14, 141-152 (2013).

4.   W. Zheng *et al.*, Two classes of ovarian primordial follicles exhibit distinct developmental dynamics and physiological functions. *Human molecular genetics* 23, 920-928 (2014).

5.   W. Niu, A. C. Spradling, Two distinct pathways of pregranulosa cell differentiation support follicle formation in the mouse ovary. *Proceedings of the National Academy of Sciences* 117, 20015-20026 (2020).

6.   A. N. Hirshfield, Heterogeneity of cell populations that contribute to the formation of primordial follicles in rats. *Biol. Reprod.* 47, 466-472 (1992).

7.   W. Zheng, H. Zhang, K. Liu, The two classes of primordial follicles in the mouse ovary: their development, physiological functions and implications for future research. *Molecular human reproduction* 20, 286-292 (2014).

8.   D. Monniaux *et al.*, The ovarian reserve of primordial follicles and the dynamic reserve of antral growing follicles: what is the link? *Biol. Reprod.* 90, 85 (2014).

9.   K. Hardy *et al.*, Nuclear exclusion of SMAD2/3 in granulosa cells is associated with primordial follicle activation in the mouse ovary. *J. Cell Sci.* 131 (2018).

10.   K. J. Hutt, E. A. McLaughlin, M. K. Holland, KIT/KIT Ligand in Mammalian Oogenesis and Folliculogenesis: Roles in Rabbit and Murine Ovarian Follicle Activation and Oocyte Growth1. *Biology of Reproduction* 75, 421-433 (2006).

11.   G. B. John, T. D. Gallardo, L. J. Shirley, D. H. Castrillon, Foxo3 is a PI3K-dependent molecular switch controlling the initiation of oocyte growth. *Developmental biology* 321, 197-204 (2008).

12.   E. R. Frost, G. Taylor, M. A. Baker, R. Lovell-Badge, J. M. Sutherland, Establishing and maintaining fertility: the importance of cell cycle arrest. *Genes Dev.* 35, 619-634 (2021).

13.   S. Lintern-Moore, G. P. Moore, The initiation of follicle and oocyte growth in the mouse ovary. *Biol. Reprod.* 20, 773-778 (1979).

14.   A. Gougeon, G. B. Chainy, Morphometric studies of small follicles in ovaries of women at different ages. *J Reprod Fertil* 81, 433-442 (1987).

15.   H. Zhang *et al.*, Somatic cells initiate primordial follicle activation and govern the development of dormant oocytes in mice. *Current biology : CB* 24, 2501-2508 (2014).

16.   S. Y. Kim *et al.*, Cell autonomous phosphoinositide 3-kinase activation in oocytes disrupts normal ovarian function through promoting survival and overgrowth of ovarian follicles. *Endocrinology* 156, 1464-1476 (2015).

17. P. Reddy *et al.*, Oocyte-specific deletion of Pten causes premature activation of the primordial follicle pool. *Science* **319**, 611-613 (2008).

18. Y. Tong *et al.*, Rapamycin-sensitive mTORC1 signaling is involved in physiological primordial follicle activation in mouse ovary. *Molecular reproduction and development* **80**, 1018-1034 (2013).

19. J. Grosbois, I. Demeestere, Dynamics of PI3K and Hippo signaling pathways during in vitro human follicle activation. *Human reproduction (Oxford, England)* **33**, 1705-1714 (2018).

20. M.-C. Meinsohn *et al.*, Single-cell sequencing reveals suppressive transcriptional programs regulated by MIS/AMH in neonatal ovaries. *Proceedings of the National Academy of Sciences* **118**, e2100920118 (2021).

21. M. E. Morris *et al.*, A single-cell atlas of the cycling murine ovary. *Elife* **11**, e77239 (2022).

22. L. Garcia-Alonso *et al.*, Single-cell roadmap of human gonadal development. *Nature* **607**, 540-547 (2022).

23. S. Rajareddy *et al.*, p27kip1 (cyclin-dependent kinase inhibitor 1B) controls ovarian development by suppressing follicle endowment and activation and promoting follicle atresia in mice. *Molecular endocrinology (Baltimore, Md.)* **21**, 2189-2202 (2007).

24. E. R. Frost *et al.*, Two alternative methods for the retrieval of somatic cell populations from the mouse ovary. *Mol. Hum. Reprod.* **27** (2021).

25. C. Mayère *et al.*, Origin, specification and differentiation of a rare supporting-like lineage in the developing mouse gonad. *Science advances* **8**, eabm0972 (2022).

26. R. H. Rastetter *et al.*, Marker genes identify three somatic cell types in the fetal mouse ovary. *Developmental biology* **394**, 242-252 (2014).

27. L. Mork *et al.*, Temporal differences in granulosa cell specification in the ovary reflect distinct follicle fates in mice. *Biol. Reprod.* **86**, 37 (2012).

28. Z. Wang *et al.*, Follistatin288 Regulates Germ Cell Cyst Breakdown and Primordial Follicle Assembly in the Mouse Ovary. *PLoS ONE* **10**, e0129643 (2015).

29. J. E. Fortune, R. A. Cushman, C. M. Wahl, S. Kito, The primordial to primary follicle transition. *Mol. Cell. Endocrinol.* **163**, 53-60 (2000).

30. C. Ahlmann-Eltze, W. Huber, glmGamPoi: fitting Gamma-Poisson generalized linear models on single cell count data. *Bioinformatics* **36**, 5701-5702 (2020).

31. A. Radley, S. Boeing, A. Smith, Branching topology of the human embryo transcriptome revealed by Entropy Sort Feature Weighting. *Development* **151** (2024).

32. J. Perez-Sanz *et al.*, Increased number of multi-oocyte follicles (MOFs) in juvenile p27Kip1 mutant mice: potential role of granulosa cells. *Human reproduction (Oxford, England)* **28**, 1023-1030 (2013).

33. I. Stevant *et al.*, Dissecting Cell Lineage Specification and Sex Fate Determination in Gonadal Somatic Cells Using Single-Cell Transcriptomics. *Cell reports* **26**, 3272-3283.e3273 (2019).

34. X. Fan *et al.*, Single-cell reconstruction of follicular remodeling in the human adult ovary. *Nature communications* **10**, 3164 (2019).

35.   A. S. K. Jones *et al.*, Cellular atlas of the human ovary using morphologically guided spatial transcriptomics and single-cell sequencing. *Science advances* **10**, eadm7506 (2024).

36.   X. Fan *et al.*, Single-cell reconstruction of follicular remodeling in the human adult ovary. *Nature communications* **10**, 3164 (2019).

37.   J. V. V. Isola *et al.*, A single-cell atlas of the aging mouse ovary. *Nature Aging* **4**, 145-162 (2024).

38.   S. Wang *et al.*, Single-Cell Transcriptomic Atlas of Primate Ovarian Aging. *Cell* **180**, 585-600.e519 (2020).

39.   G. Nagamatsu, S. Shimamoto, N. Hamazaki, Y. Nishimura, K. Hayashi, Mechanical stress accompanied with nuclear rotation is involved in the dormant state of mouse oocytes. *Science advances* **5**, eaav9960 (2019).

40.   X. Zhang *et al.*, Enhanced glycolysis in granulosa cells promotes the activation of primordial follicles through mTOR signaling. *Cell Death & Disease* **13**, 87 (2022).

41.   J.-J. Wang *et al.*, Single-cell transcriptome landscape of ovarian cells during primordial follicle assembly in mice. *PLoS Biol.* **18**, e3001025 (2020).

42.   Y.-Y. Chen *et al.*, Single-cell transcriptomics of staged oocytes and somatic cells reveal novel regulators of follicle activation. *Reproduction* **164**, 55-70 (2022).

43.   J. Zhou *et al.*, Dissecting the fate of Foxl2-expressing cells in fetal ovary using lineage tracing and single-cell transcriptomics. *Cell Discov* **8**, 139 (2022).

44.   A. Dattani *et al.*, Naive pluripotent stem cell-based models capture FGF-dependent human hypoblast lineage specification. *Cell Stem Cell* **31**, 1058-1071.e1055 (2024).

45.   A. R. Libby, T. Rito, A. Radley, J. Briscoe, An in vivo CRISPR screen in chick embryos reveals a role for MLLT3 in specification of neural cells from the caudal epiblast. *bioRxiv* 10.1101/2024.05.16.594506, 2024.2005.2016.594506 (2024).

46.   N. Gonen *et al.*, In vitro cellular reprogramming to model gonad development and its disorders. *Science advances* **9**, eabn9793 (2023).

47.   D. E. Montgomery *et al.*, alpha-Adrenergic response and myofilament activity in mouse hearts lacking PKC phosphorylation sites on cardiac TnI. *Am J Physiol Heart Circ Physiol* **282**, H2397-2405 (2002).

48.   Y. Pi, D. Zhang, K. R. Kemnitz, H. Wang, J. W. Walker, Protein kinase C and A sites on troponin I regulate myofilament Ca2+ sensitivity and ATPase activity in the mouse myocardium. *J. Physiol.* **552**, 845-857 (2003).

49.   S. Sakthivel *et al.*, In vivo and in vitro analysis of cardiac troponin I phosphorylation. *The Journal of biological chemistry* **280**, 703-714 (2005).

50.   J. E. Garcia-Ortiz *et al.*, Foxl2 functions in sex determination and histogenesis throughout mouse ovary development. *BMC developmental biology* **9**, 36 (2009).

51.   R. Migale *et al.*, FOXL2 interaction with different binding partners regulates the dynamics of ovarian development. *Science advances* **10**, eadl0788 (2024).

52.   X. Huang *et al.*, Cardiac Troponin I Gene Knockout. *Circ. Res.* **84**, 1-8 (1999).

53.   M. L. Fero *et al.*, A syndrome of multiorgan hyperplasia with features of gigantism, tumorigenesis, and female sterility in p27(Kip1)-deficient mice. *Cell* **85**, 733-744 (1996).

54. G. X. Y. Zheng *et al.*, Massively parallel digital transcriptional profiling of single cells. *Nat. Commun.* **8**, 14049 (2017).

55. T. Stuart *et al.*, Comprehensive Integration of Single-Cell Data. *Cell* **177**, 1888-1902.e1821 (2019).

56. B. Li, C. N. Dewey, RSEM: accurate transcript quantification from RNA-Seq data with or without a reference genome. *BMC Bioinformatics* **12**, 323 (2011).

57. A. Dobin *et al.*, STAR: ultrafast universal RNA-seq aligner. *Bioinformatics* **29**, 15-21 (2013).

58. D. S. DeLuca *et al.*, RNA-SeQC: RNA-seq metrics for quality control and process optimization. *Bioinformatics* **28**, 1530-1532 (2012).

59. M. I. Love, W. Huber, S. Anders, Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. *Genome Biol.* **15**, 550 (2014).

60. R. C. Development, Team, *R: A language and environment for statistical computing.* (R Foundation for Statistical Computing, Vienna, Austria, 2008).

61. J. Schindelin *et al.*, Fiji: an open-source platform for biological-image analysis. *Nature Methods* **9**, 676-682 (2012).

Figure 1. Identification of activating pregranulosa cell gene expression signature in mouse ovaries.

a UMAPs from scRNAseq on whole mouse ovaries before (E18.5), during (PD4) and after (PD7) primordial follicle activation. 18 clusters representing major ovarian cell types were identified.

b Dotplot showing known markers of pregranulosa cells are found in Gc_1, Gc_2, Gc_3 and Gc_4.

c UMAP from cESFW workflow. Middle panel: coloured by developmental timepoints - E18.5 (yellow), PD4 (magenta) and PD7 (blue). Bottom panel: Unsupervised clusters (Supplemental Figure 1d) coloured by cell type labels from conventional scRNAseq analysis workflow (Ep_1, Gc_1, Gc_2, Gc_3, Gc_4 and Gc_5).

d Heatmap top 10 entropy sort score genes within three pregranulosa cell clusters (Gc_1, Gc_2, Gc_3) and granulosa cells (Gc_4).

Figure 2. Disruption of p27^{Kip1} highlights functional aspects of activating pregranulosa cell gene expression signature.
a Haematoxylin and Eosin staining of E18.5 and PD8 ovaries in WT and *Cdkn1b*$^{-/-}$ mice. Imaged at 10x magnification.
b Volcano plots showing differential expression of gene transcripts between E18.5 WT and *Cdkn1b*$^{-/-}$ ovarian cells (upregulated genes in red, downregulated in blue, n = 4 replicates per group). Black dots indicate genes below the p ≤ 0.05 cutoff.
c Heatmap showing differentially expressed genes in *Cdkn1b*$^{-/-}$ ovaries (p ≤ 0.05) overlaid with granulosa cell subclustering analysis from the scRNAseq experiment (upregulated genes in red, downregulated genes in blue).
d UMAP from cESFW workflow coloured by timepoint, by cluster annotation, and by expression levels of *Cdkn1b* and *Tnni3*.

Figure 3. TNNI3 expression increases as pregranulosa cells activate.
a Immunofluorescence on *FoxL2*p2A-eGFP ovarian cryosections at E18.5 (n=3), PD4 (n=3) and PD7 (n=3). Each section is immunolabeled for GFP (green) labelling all granulosa and pregranulosa cells, TNNI3 (purple) and DAPI (white). 20x magnification images shown in top row of each timepoint. E18.5 samples were further imaged at 100x magnification. PD4 and PD7 samples were additionally imaged at 63x magnification. Growing follicles are denoted by a white dashed border while primordial follicles are denoted by a dashed red border. Transitional follicles denoted in orange dashed border. Yellow arrows indicate high TNNI3 expression in granulosa cells. White scale bars show 20µm.
b High magnification image of pregranulosa cells of an E18.5 *FoxL2*p2A-eGFP primordial follicle (red dashed line) and granulosa cells (white dashed line) of a PD7 growing follicle displaying high TNNI3 staining predominantly around the edge of the granulosa cells (indicated by white arrows). Scale bars = 20µm
c Violin box plot graphing the mean fluorescent intensity of TNNI3 staining between pre-granulosa and granulosa cells in PD7 ovaries (n=3). ****p=<0.0001
d Schematic of pregranulosa to granulosa cell activation.