The identification of a gene expression signature of primordial follicle activation in

mouse pregranulosa cells

Emily R Frost[1,2,3]* Güneş Taylor[3]*#, Stefan Boeing[4,5], Christophe Galichet[3], Mark A Baker[1,2],

Jessie M Sutherland[1,2+#], Robin Lovell-Badge[3+#]

1 Priority Research Centre for Reproductive Science, School of Biomedical Science & Pharmacy,

University of Newcastle, Ring Road, Callaghan, New South Wales, Australia, 2308

2 Hunter Medical Research Institute, New Lambton Heights, New South Wales, Australia, 2305

3 Stem Cell Biology and Developmental Genetics Lab, The Francis Crick Institute, London, United

Kingdom, NW1 1AT

4 Bioinformatics and Biostatistics Facility, The Francis Crick Institute, London, United Kingdom

5 Scientific Computing – Digital Development Team, The Francis Crick Institute, London, United

Kingdom

* These authors contributed equally and should be regarded as co-first authors

+ These authors contributed equally and should be regarded as co-last authors

Corresponding authors: Dr Güneş Taylor, The Francis Crick Institute, gunes.taylor@crick.ac.uk

Dr Jessie Sutherland, University of Newcastle, jessie.sutherland@newcastle.edu.au

Prof Robin Lovell-Badge, The Francis Crick Institute, robin.lovell-badge@crick.ac.uk

Graphical abstract

Abstract

Female fertility is dictated by the number of oocyte-containing primordial follicles within the

ovary. These follicles are established early in development and their selective activation

represents the definitive first step towards oocyte maturation and ovulation. The intrinsic

molecular mechanisms that regulate ovarian follicle activation are largely uncharacterised. In

this study, we report a single-cell RNA sequencing (scRNAseq) dataset to examine mouse

embryonic and neonatal ovaries across the developmental window of primordial follicle

formation and activation. We identified five distinct clusters of granulosa cells and

bioinformatic analysis revealed a population of pregranulosa cells that appeared to be

undergoing follicle activation. This cluster was uniquely differentiated by the expression of

Tnni3, *Slc18a2*, *Fam13a* and *Klf2*, all of which are novel to follicle activation. Finally, we

examined the transcriptome of a mouse model where all primordial follicles enter

folliculogenesis simultaneously ($Cdkn1b^{-/-}$) and showed that the signature genes of activating

pregranulosa cells was observed precociously in $Cdkn1b^{-/-}$ ovaries, further demonstrating that

$p27^{kip1}$ acts as an important repressor of follicle activation. Combined, this data provides insight

into how primordial follicle activation is regulated in mammals and could be utilised to identify

pathways to target to improve fertility preservation and infertility conditions in the future.

Introduction

Mammalian ovary development is a tightly regulated process that is crucial for establishing the ovarian reserve of oocytes. The ovarian reserve must be able to sustain fertility over several months in mice and several decades in humans (Monniaux et al., 2014). The ovarian reserve is defined by the number of oocytes present in the ovary in structures called primordial follicles. Primordial follicles consist of an oocyte with a layer of surrounding pregranulosa cells and are formed during gestation (Kezele, Ague, Nilsson, & Skinner, 2005; Pepling & Spradling, 2001; Tingen, Kim, & Woodruff, 2009). To create the ovarian reserve, both mitosis and meiosis occur in co-ordinated events in the pregranulosa cells and the oocytes respectively. Following the determination of ovarian fate in the bipotential gonad, pregranulosa cells and oogonia develop in close contact, throughout germ cell nest formation and breakdown, resulting in primordial follicle formation and activation (Lei & Spradling, 2013; Pepling & Spradling, 1998; Suzuki, Kanai-Azuma, & Kanai, 2015; Wear, McPike, & Watanabe, 2016).

In the mouse ovary, it is well-established that two specific populations of primordial follicles exist, classified by their spatial distribution and origin, namely the medullary and the cortical primordial follicles (Mork et al., 2012; Niu & Spradling, 2020; W. Zheng, Zhang, Gorre, et al., 2014; W. Zheng, Zhang, & Liu, 2014). Two precise populations of pregranulosa cells contribute to the medullary and cortical classes of primordial follicles. Both lineages of pregranulosa cells derive from the ovarian surface epithelium but at different times (Niu & Spradling, 2020; W. Zheng, Zhang, Gorre, et al., 2014). The first pregranulosa cell population that develops express FOXL2 and p27[kip1] and contribute to medullary primordial follicles, termed bipotential pregranulosa cells (BPGs) (Niu & Spradling, 2020; J.-J. Wang et al., 2020). The second population, termed epithelial pregranulosa cells (EPGs) ingress from the epithelium later. They express LGR5 and contribute to cortical primordial follicles, expressing

FOXL2 and p27[kip1] postnatally (Niu & Spradling, 2020; J.-J. Wang et al., 2020). The formation

of these two classes of primordial follicles are asynchronous, with medullary primordial

follicles formed first (J.-J. Wang et al., 2020). Aside from the distinct differences in

developmental timing, it is not yet clear whether other characteristics vary between the two

pregranulosa classes. Recent advances have identified molecular pathways in both the oocytes

and pregranulosa cells that regulate the formation of primordial follicles, including the

NOTCH, KITL, TGFß and PI3K pathways (Fukuda, Muraoka, Kato, & Saga, 2021; Niu &

Spradling, 2020; Rosario et al., 2019; J.-J. Wang et al., 2020).

In contrast to follicle formation, primordial follicle activation is more challenging to

interrogate, due to the continuous selection of follicles for activation throughout prepubertal

and adult life. However, an initial wave of primordial follicle activation occurs postnatally in

mouse, with the activation of medullary follicles entering into folliculogenesis (Cordeiro et al.,

2015; Mork et al., 2012). This wave presents an opportunity to understand the factors and

pathways that govern the activation of primordial follicles into folliculogenesis. Changes in

cell cycle status, granulosa cell morphology (from squamous to cuboidal) and oocyte size are

key indicators of follicle activation and the upregulation of essential factors including PI3K,

KITL and BMP4 are indicative of entry into the developing pool of follicles (Adhikari & Liu,

2009; Hardy et al., 2018; Hutt, McLaughlin, & Holland, 2006; John, Gallardo, Shirley, &

Castrillon, 2008; McLaughlin & McIver, 2009). Protocols have recently been developed for

the dissociation of the neonatal mouse ovary, generating single-cell suspensions for

sequencing, that can be used during this initial wave of activation (Y.-Y. Chen et al., 2022;

Frost et al., 2021a). To date, most studies have utilised genetically modified mouse models and

microarray experiments to explore the expression of gene-driven signals in stimulating or

inhibiting primordial follicle activation (reviewed in (Y. Chen et al., 2020; McLaughlin &

McIver, 2009; Sobinoff, Sutherland, & McLaughlin, 2013)).

In this study, we use single-cell RNA sequencing (scRNA-seq) to examine the transcriptional

profile of mouse embryonic and neonatal ovaries at three timepoints: embryonic day (E) 18.5,

postnatal day (PND) 4 and PND7. In total we sequenced 24,810 cells from the three

developmental stages and distinguished five clusters of granulosa cells. We identified a

transcriptomic signature of pregranulosa cells during primordial follicle activation in granulosa

cell cluster 1 (Gc_1), by comparing with the activated granulosa cells present in granulosa cell

cluster 4 (Gc_4). Moreover, analysis of a genetically modified mouse line with an accelerated

follicle activation phenotype ($Cdkn1b^{-/-}$) revealed that this signature was observed precociously

in the mutant mice as early as E18.5. Combined, this data reveals pregranulosa cell-specific

pathways that control primordial follicle activation in mice. Further interrogation of these

pathways in the infertile mouse model shows disruptions in the follicle activation process,

uncovering key insights into how primordial follicle activation establishes fertility.

Results

Characterisation of mouse ovarian transcriptome before and during primordial follicle

activation

To investigate the transcriptional signature of granulosa cells both prior to and during

primordial follicle activation, we performed single-cell RNA sequencing of the mouse ovary.

We focused on three critical timepoints representative of ovary development before, during

and after follicle activation, namely: embryonic day 18.5 (E18.5), when germ cell nest

breakdown is initiated to form primordial follicles; post-natal day 4 (PND4), when the initial

wave of primordial follicle activation is occurring; and PND7, when populations of activated

follicles (termed primary follicles) are present within the ovary (Figure 1A). A previously

described and validated dissociation protocol using the enzyme papain was utilised to achieve

a single cell suspension of the ovarian tissue at each stage (Frost et al., 2021b). In total, our

dataset contained 24,810 cells, with 8,210 cells from one E18.5 ovary, 7,587 cells from one

PND4 ovary and 9,013 cells from one PND7 ovary (Table S1). Initially, the data from all three

timepoints was integrated and uniform manifold approximation and projection (UMAP)

analysis was used to assess cell contribution to the clustering based on the developmental stage

(Figure 1B). Across the integrated timepoints, 18 distinct clusters were found, and the six major

cell types present in the ovary across these developmental stages were annotated based on their

expression of established cell type specific genetic markers (granulosa cells, mesenchymal

cells, oocytes, epithelial cells, immune cells and blood cells) (Figure 1C and Figure S1). Across

all three timepoints, most cells mapped to either the mesenchymal (43%) or granulosa cell

(39%) clusters. The number of clusters per cell type were as follows: five granulosa cell

clusters, six mesenchymal clusters, one oocyte cluster, two epithelial clusters, one immune

cluster, two blood cell clusters and one cluster unable to be identified (labelled 'other').

Granulosa cells were identified by the expression of *Fst*, *Amhr2*, *Kitl*, *Runx1*, *Foxl2, Hmgcs2*

and *Lgr5* and clusters were labelled as Gc_1 to Gc_5 (Bagheri-Fam et al., 2020; Jones &

Pepling, 2013; Meinsohn et al., 2021; Nicol et al., 2019; Ottolenghi et al., 2005; Rastetter et

al., 2014; Z. Wang et al., 2015). Mesenchymal cells expressed *Nr2f2* and *Col1a1* and clusters

were named Mc_1 to Mc_6 (Rastetter et al., 2014). Epithelial cells were characterised by the

expression of *Upk3b* and *Krt19*, with epithelial clusters labelled as Ep_1 and Ep_2 (Gaytan,

Morales, Roa, & Tena-Sempere, 2018; Niu & Spradling, 2020). Oocytes were distinguished

by *Ddx4*, *Dazl* and *Sycp3* expression and contained within cluster Oo_1 (Cooke, Lee, Kerr, &

Ruggiu, 1996; Cordeiro et al., 2015; Reynolds, Collier, Bingham, Gray, & Cooke, 2007;

Rosario et al., 2019; Yang et al., 2020). Blood cells were characterised by expression of *Cldn5*

136   and *Cx3cr1* and clusters were labelled as Bl_1 and Bl_2 (Landsman et al., 2009; Ma et al.,

137   2017), while immune cells expressed *Cd53*, contained in cluster Im_1 (Greenberg et al., 2020;

138   J.-J. Wang et al., 2020). Taken together, these results validated the successful isolation of

139   mouse ovarian cells during late embryonic and early postnatal development and illustrated

140   distinct cell-specific transcriptomic profiles essential for further interrogation.

Figure 1. Transcriptional profiling of whole ovaries before, during and after primordial follicle

activation in the mouse

(A) A diagram of the mouse ovary during late embryonic and early postnatal development, highlighting

the key events of germ cell nest breakdown, primordial follicle formation and primordial follicle

activation, and the timepoints chosen for this study. The diagram also shows the workflow of sample

preparation for single-cell RNA sequencing. UMAP Dimensionality Reduction Plots showing

individual cells coloured by timepoint (B) and by cell type (C). Clusters were named by cell type as

follows: granulosa cells (Gc_1, Gc_2, Gc_3, Gc_4, and Gc_5), mesenchymal cells (Mc_1, Mc_2,

Mc_3, Mc_4, Mc_5 and Mc_6), epithelial cells (Ep_1 and Ep_2), oocytes (Oo_1), blood cells (Bl_1

and Bl_2), immune cells (Im_1) and other cells (Ot_1). (D) Heat map showing the most distinct marker

genes in each cluster, used to identify the clusters in the UMAP. For this analysis, 5,000 cells were

randomly selected from the 24,810 cells in this experiment.

Activated granulosa cells are transcriptionally distinct postnatally

Next, we analysed the transcriptomic signature of the ovarian cells at each developmental

stage, by generating individual UMAP analyses by timepoint, to examine the gene expression

of cells before and during primordial follicle activation (Figure 2A). Most of the clusters that

were identified were present in all three developmental stages (Figure 2A). However, changes

in cell contribution were observed in the oocyte (Oo_1) and granulosa cell cluster 1 and 4

(Gc_1 and Gc_4) clusters across developmental time. The number of cells contributing to the

Oo_1 and Gc_1 clusters decreased over time (Figure 2A and Figure S2A and S2B). In contrast,

of the five pregranulosa cell clusters, Gc_4 was absent at E18.5 but appeared at PND4 and was

maintained at PND7 (Figure 2A and Figure S2A and S2B). Cluster Gc_4 expressed granulosa

cell marker genes including *Foxl2, Amhr2* and *Hmgcs2* (Figure 2B and 2C). However, unlike

the other granulosa cell clusters Gc_1, Gc_2 and Gc_3, the Gc_4 cluster also expressed high

levels of genes associated with proliferation, including *Mki67* and *Pcna* (Figure 2D and 2E)

(Miller et al., 2018; Xu et al., 2011). Using gene ontology analysis, the top biological processes

from the genes upregulated in Gc_4 are related to proliferation and include cell cycle

(GO:0007049), mitotic nuclear division (GO:0007067), cell division (GO:0051301),

chromosome segregation (GO:0007059) and mitotic cytokinesis (GO:0000281) (Figure 2F).

Combined, the Gc_4 cluster gene expression signature is indicative of differentiated granulosa

cells typical of primary follicles, which have transitioned into cuboidal granulosa cells and

have entered the cell cycle and are proliferating to generate multiple layers of granulosa cells

surrounding the oocyte. Given the appearance of a population of activated granulosa cells in

the postnatal ovary, we then examined the transcriptomic profile of the remaining pregranulosa

cell populations to characterise the key gene expression differences between clusters and

identify a population of pregranulosa cells in a primed state for activation.

Figure 2. A distinct population of highly proliferative granulosa cells appear during primordial follicle activation in the mouse ovary

(A) UMAP Dimensionality Reduction Plots showing individual cells from each individual timepoint coloured by cell type. (B) UMAP Dimensionality Reduction Plots showing cells from all three samples, overlaid with their expression of the following genes associated with granulosa cells: *Amh, Amhr2, Cdkn1b, Esr2, Foxl2, Fst, Hmgcs2, Hsd17b1, Hsd3b1, Inha, Kitl, Nr5a2,* and *Runx1*. Dots are coloured according to their level of gene expression, with high levels of expression shown in purple. Expression was determined by area under the curve (AUC). (C) Dotplot showing the expression of the above genes associated with granulosa cells, with high levels of expression shown in purple and the percentage of cells in each cluster expressing the gene indicated by the size of the dot. (D) UMAP Dimensionality Reduction Plots showing cells from all three timepoints, overlaid with their expression of the following proliferation markers: *mKi67* and *Pcna*. Dots are coloured according to their level of gene expression, with high levels of expression shown in purple. Expression was determined by area under the curve (AUC). (E) Dotplot showing the expression of the above genes associated with proliferation, with high levels of expression shown in purple and the percentage of cells in each cluster expressing the gene indicated by the size of the dot. (F) Gene ontology (GO) information of the biological process categories with highest count of genes from the genes upregulated in granulosa cell cluster 4 (Gc_4). The list of genes upregulated in Gc_4 was identified by differential gene expression analysis and annotated via GO analysis in DAVID (v6.8).

Epithelial and bipotential derived pregranulosa cells exist in distinct clusters

We next asked if we could characterise the remaining granulosa cell clusters by their differences in gene expression. As cluster Gc_5 consisted of fewer cells with variable expression of granulosa cell marker genes, we focused on the four largest pregranulosa cell clusters (Gc_1 through to Gc_4). We compared the genes expressed in our granulosa cell clusters with the published data from (Niu & Spradling, 2020) to similarly classify our heterogenous granulosa cell populations. In granulosa cell cluster 1 (Gc_1), differential gene expression analysis showed that *Gng13*, *Lgas7* and *Apoc1*, among other genes, had higher levels of average expression than in all other clusters (Figure 3A and 3B, S2 Table). The gene list generated from the differential gene expression analysis, along with the proximity of the cluster to an epithelial cell cluster (Ep_1), suggested that this cluster is likely to be a population of EPGs. This was also strengthened by the observation that most of the cells in Gc_1 were from the E18.5 timepoint, as EPGs do not ingress from the ovarian epithelium to contribute to

follicles postnatally (Figure S2B) (Niu & Spradling, 2020). To confirm the identification of the

Gc_1 cluster, we used a selection of genes described as marker genes of EPGs and examined

their expression in all 18 clusters (Niu & Spradling, 2020) (Figure 3C). This supported the

differential gene expression analysis, with specific expression of EPG marker genes in

Gc_1.We used this same approach to identify granulosa cell cluster 2 (Gc_2) as a population

of pregranulosa cells derived from the bipotential *Nr5a1+* somatic cell lineage (BPGs). This

cluster showed more similarity in gene expression to the other granulosa cell clusters (Figure

3D and 3E). However, differential gene expression analysis showed that *Serpinb6a* and

Cdkn1b, among other genes, had higher levels of average expression in Gc_2 than in all other

clusters (Figure 3E). This combined with the position of the cluster between the EPGs and the

mitotically active granulosa cells highlighted the Gc_2 cluster as a likely BPG population. To

verify, we compared our expression with the selection of marker genes for BPGs and found

high levels of *Foxl2* and *Aard*, which are first expressed in BPGs (Niu & Spradling, 2020)

(Figure 3F). Additionally, Gc_2 cells express low levels of *Hsd3b1* and *Hmgcs2*, markers of

pregranulosa cells later in development (Figure 3F). This supported our data indicating Gc_2

cells are BPGs.

Figure 3. Clear origins of two populations of pregranulosa cells based on their gene expression signature

(A) UMAP Dimensionality Reduction Plots showing cells from all three samples, overlaid with the

cells selected for differential gene expression (DGE) analysis, Gc_1 cells are shown in yellow and all

other clusters are shown in grey. DGE analysis performed using the method detailed by Ahlmann-Eltze

(Ahlmann-Eltze & Huber, 2020). (B) MA plot showing the differential expression of gene transcripts

between Gc_1 cells and cells from all other clusters; genes upregulated in Gc_1 cells are shown in

yellow and genes upregulated in all other cells shown in black. Grey dots indicate genes below the $\log_2$

fold change $\pm$ 2; p $\leq$ 0.05 cutoff. (C) Dotplot showing the expression of genes associated with epithelial

pregranulosa cells, with high levels of expression shown in purple and the percentage of cells in each

cluster expressing the gene indicated by the size of the dot. Gene list obtained from (Niu & Spradling,

2020). (D) UMAP Dimensionality Reduction Plots showing cells from all three samples, overlaid with

the cells selected for differential gene expression (DGE) analysis, Gc_2 cells are shown in orange and

all other clusters are shown in grey. DGE analysis performed using the method detailed by Ahlmann-

Eltze (Ahlmann-Eltze & Huber, 2020). (E) MA plot showing the differential expression of gene

transcripts between Gc_2 cells and cells from all other clusters; genes upregulated in Gc_2 cells shown

in orange and genes upregulated in all other cells shown in black. Grey dots indicate genes below the

$\log_2$ fold change $\pm$ 2; p $\leq$ 0.05 cutoff. (F) Dotplot showing the expression of genes associated with

bipotential pregranulosa cells, with high levels of expression shown in purple and the percentage of

cells in each cluster expressing the gene indicated by the size of the dot. Gene list obtained from (Niu

& Spradling, 2020).

Identification of the genetic signature of pre-granulosa cells in activating primordial

follicles

The identification of the Gc_1 and Gc_2 clusters as EPGs and BPGs respectively, led us to

hypothesise the genetic profile of Gc_3 as a potential population of activating granulosa cells.

Cells that contribute to Gc_3 originate from all three timepoints, but a large proportion of cells

arise from the PND7 timepoint (Figure S2B). Interestingly, cells in Gc_3 have similar gene

expression signatures to BPGs in Gc_2 (shown by the expression of *Foxl2* and *Hmgcs2*

amongst others), but also high average levels of expression of genes like *Kitl* and *Fst*, which

are well-established regulators of follicle activation (Figure 4A and 4B) (Hutt et al., 2006; Z.

Wang et al., 2015). Additionally, the cells in the Gc_3 cluster also express markers of

differentiated granulosa cells, such as *Hsd3b1*, *Amh*, *Esr2* and *Nr5a2*, albeit at lower average

levels of expression when compared to the differentiated proliferative granulosa cells in Gc_4

(Figure 2C and 4B). Differential gene expression analysis illustrated that *Hsd3b1, Fst* and *Inha*

among other genes, were on average more highly expressed in Gc_3 than in other cell clusters,

supporting the hypothesis that these cells are activating pregranulosa cells (Figure 2C and 4B

and S2 Table).

To determine if Gc_3 is an intermediate stage between quiescent BPG cells (Gc_2) and

differentiated proliferating granulosa cells (Gc_4), we performed two differential gene

expression (DGE) analyses between Gc_3 and Gc_2, and Gc_3 and Gc_4. The 3,302 cells that

contributed to Gc_3 and 1,803 cells that contributed to Gc_2 were selected and average gene

expression was analysed according to the methods detailed by Ahlmann-Eltze (Ahlmann-Eltze

& Huber, 2020) (Figure 4C). There was a total of 581 differentially expressed genes ($\log_2$ fold

change ± 2; $p \leq 0.05$) between cells from Gc_3 and Gc_2 (Figure 4D and S3 Table). 274 genes

were upregulated in Gc_3 cells and 307 genes were upregulated in Gc_2 cells (Figure 4D and

S3 Table). Gene ontology was used to identify the top biological pathways implicated in Gc_2

cells and Gc_3 respectively (Figure 4E and 4F). In the Gc_2 cluster, the top biological

processes based on 307 differentially expressed genes were ion transport (GO:0006811),

defence response to virus (GO:0051607), transmembrane transport (GO:0055085), immune

system process (GO:0002376) and immune response (GO:0006955) (Figure 4E). In contrast,

the top processes based on the 274 differentially expressed genes in the Gc_3 cluster were cell

cycle (GO:0007049), cellular response to DNA damage stimulus (GO:0006974), DNA

replication (GO:0006260), cell adhesion (GO:0007155) and DNA repair (GO:0006281)

(Figure 4F). These processes illustrate that cell cycle and the associated DNA replication

processes are occurring in the cells of Gc_3, which is a critical cell fate transition towards

differentiated granulosa cells.

Figure 4. Comparison of gene expression between putative activating pregranulosa cells (Gc_3)

and bipotential pregranulosa cells (Gc_2)

(A) UMAP Dimensionality Reduction Plots showing cells from all three samples, overlaid with the

cells selected for differential gene expression (DGE) analysis, Gc_3 cells are shown in pink and all

other clusters are shown in grey. DGE analysis performed using the method detailed by Ahlmann-Eltze

(Ahlmann-Eltze & Huber, 2020). (B) MA plot showing the differential expression of gene transcripts between Gc_3 cells and cells from all other clusters; genes upregulated in Gc_3 cells are shown in pink and genes upregulated in all other cells shown in black. Grey dots indicate genes below the $\log_2$ fold change $\pm$ 2; $p \leq 0.05$ cutoff. (C) UMAP Dimensionality Reduction Plots showing cells from all three samples, overlaid with the cells selected for differential gene expression (DGE) analysis, Gc_3 cells are shown in pink and Gc_2 cells are shown in orange. DGE analysis performed using the method detailed by Ahlmann-Eltze (Ahlmann-Eltze & Huber, 2020). (D) MA plot showing the differential expression of gene transcripts between Gc_3 and Gc_2 cells; genes upregulated in Gc_3 cells shown in pink and genes upregulated in Gc_2 cells shown in orange. Grey dots indicate genes below the $\log_2$ fold change $\pm$ 2; $p \leq 0.05$ cutoff. (E) Gene ontology (GO) information of the biological process categories with highest count of genes from the genes upregulated in Gc_2 shown in orange. (F) Gene ontology (GO) information of the biological process categories with highest count of genes from the genes upregulated in Gc_3 shown in pink. The lists of genes upregulated in Gc_2 and Gc_3 were identified by differential gene expression analysis and annotated via GO analysis in DAVID (v6.8).

To characterise a gene expression signature of primordial follicle activation, we compared the differentially expressed genes from the differential gene expression analysis performed in Figure 4A (S2 Table) and the differentially expressed genes from the Gc_3 vs Gc_2 comparison performed in Figure 4C (S3 Table). Established pregranulosa marker genes of primordial follicle activation including *Amh*, *Inha* and *Nr5a2* were observed when these gene lists were compared (Figure S3). The expression of these genes validated our approach for identifying a gene expression signature of activation, and we then looked for novel genes potentially controlling activation. Genes including *Slc18a2*, *Tnni3* and *Fam13a* are suitable candidates that share similar gene expression profiles to *Amh*, *Inha* and *Nr5a2*, and are representative of cells entering activation (Figure 5).

Figure 5. Expression of key genes upregulated during primordial follicle activation

The UMAP cluster maps show the expression of differentially expressed genes in Gc_3. UMAP cluster maps of novel pregranulosa marker genes (*Slc18a2*, *Tnni3* and *Fam13a*) identified in our differential gene expression analysis and DGE comparisons that are upregulated during primordial follicle activation. UMAP cluster maps are shown for each timepoint individually and only include cells from the timepoints in each panel (E18.5 = yellow, PND4 = pink and PND7 = blue). Dots are coloured according to their level of gene expression, with high levels of expression shown in purple.

We also performed DGE analysis between the activating pregranulosa cells (Gc_3) and the differentiated granulosa cells (Gc_4). The same method was utilised as previously, with 3,302 Gc_3 cells selected and 909 Gc_4 cells selected for gene expression analysis (Figure 6A). 667 differentially expressed genes ($\log_2$ fold change $\pm$ 2; $p \leq 0.05$) were identified between cells

from Gc_3 and Gc_4 (Figure 6B and S4 Table). 389 genes were upregulated in Gc_3 cells,

compared with 278 genes upregulated in Gc_4 cells (Figure 6B and S4 Table). The top

biological processes associated with the genes expressed in Gc_3 cells were cell differentiation

(GO:0030154), multicellular organism development (GO:0007275), spermatogenesis

(GO:0007283), meiotic cell cycle (GO:0051321) and metabolic process (GO:0008152) (Figure

6C). Gc_4 cells had distinct biological processes divergent from Gc_3 cells, namely cell cycle

(GO:0007049), mitotic nuclear division (GO:0007067), cell division (GO:0051301),

chromosome segregation (GO:0007059) and nucleosome assembly (GO:0006334) (Figure

6D). Taken together, these results provided a list of genes that are expressed in the pregranulosa

cells at the initiation of primordial follicle activation, prior to re-entry into the cell cycle.

Figure 6. Comparison of gene expression between granulosa cell cluster 3 (Gc_3) and granulosa cell cluster 4 (Gc_4)

(A) UMAP Dimensionality Reduction Plots showing cells from all three samples, overlaid with the cells selected for differential gene expression (DGE) analysis, Gc_3 cells are shown in pink and Gc_4 cells are shown in maroon. DGE analysis performed using the method detailed by Ahlmann-Eltze (Ahlmann-Eltze & Huber, 2020). (B) MA plot showing the differential expression of gene transcripts between Gc_3 and Gc_4 cells; genes upregulated in Gc_3 cells shown in pink and genes upregulated in Gc_4 cells shown in maroon. Grey dots indicate genes below the $\log_2$ fold change ± 2; $p \leq 0.05$ cutoff. (C) Gene ontology (GO) information of the biological process categories with highest count of genes from the genes upregulated in Gc_3 shown in pink. (D) Gene ontology (GO) information of the biological process categories with highest count of genes from the genes upregulated in Gc_4 shown in maroon. The lists of genes upregulated in Gc_2 and Gc_3 were identified by differential gene expression analysis and annotated via GO analysis in DAVID (v6.8).

Examining the changes in gene expression signature in a model of accelerated primordial follicle activation

Having identified putative genes expressed in the pregranulosa cells during primordial follicle activation, we next looked to query these candidate genes in an established model of abnormal follicle activation. Specifically, the genetically modified mouse line *Cdkn1b*[-/-] which lacks the expression of the cell cycle inhibitor p27[kip1] (gene name *Cdkn1b*). This mutant line shows an uncontrolled activation of follicles which enter folliculogenesis perinatally, inducing infertility prior to sexual maturation (Perez-Sanz et al., 2013; Rajareddy et al., 2007). No previous RNA sequencing experiments have been performed on ovaries from *Cdkn1b*[-/-] mice. We examined the ovaries of *Cdkn1b*[-/-] and littermate wildtype controls at E18.5. This timepoint was selected as it was reported to be prior to primordial follicle activation occurring in both wild-type and *Cdkn1b*[-/-] ovaries (Rajareddy et al., 2007). Additionally, this timepoint covers germ cell nest breakdown and primordial follicle formation and overlaps with the first timepoint of the single-cell sequencing experiment. As previously reported, no gross morphological differences in histology were observed at E18.5 (Figure 7A). The absence of p27kip1 in *Cdkn1b*[-/-] ovaries was confirmed by qRT-PCR (Figure 7B) and by immunofluorescence (Figure 7C). We assessed that the follicle activation defect was not due to a failure of germ cells or pregranulosa cells to specify correctly in *Cdkn1b*[-/-] mice, by staining for DDX4 (germ cell marker) and FOXL2 (granulosa cell marker) at E18.5 (Figure 7D). Having established the *Cdkn1b*[-/-] mouse line as an appropriate model to examine primordial follicle activation defects, we performed bulk RNA-sequencing of WT and *Cdkn1b*[-/-] ovaries at E18.5. Surprisingly, at E18.5, 1,633 transcripts were already differentially expressed between WT and *Cdkn1b*[-/-] ovaries ($p < 0.05$). Of these genes, 1,008 genes were upregulated in *Cdkn1b*[-/-] ovaries and 626 genes were downregulated in *Cdkn1b*[-/-] ovaries (Figure 7E and S5 Table). Gene ontology was utilised to identify the top biological pathways affected in *Cdkn1b*[-/-] ovaries (Figure 7F). The top

processes based on the 626 downregulated DEGs in *Cdkn1b*$^{-/-}$ ovaries were positive regulation

of transcription from RNA polymerase II promoter (GO:0045944), regulation of transcription

from RNA polymerase II promoter (GO:0006357), multicellular organism development

(GO:0007275), cell differentiation (GO:0030154) and regulation of transcription, DNA-

templated (GO:0006355) (Figure 7F). The top processes based on the 1,008 upregulated DEGs

in *Cdkn1b*$^{-/-}$ ovaries were cell cycle (GO:0007049), cell division (GO:0051301), cellular

response to DNA damage stimulus (GO:0006974), DNA repair (GO:0006281) and mRNA

processing (GO:0006397) (Figure 7F).

Figure 7. Examination of ovarian development and gene expression in E18.5 *Cdkn1b⁻/⁻* mice.

(A) Haematoxylin and Eosin (H&E) staining of E18.5 ovaries in both wild-type (WT) and *Cdkn1b*^{-/-}

mice. Images were taken at 10 x magnification and 40 x magnification, with scale bars of 100 μm and

25 μm respectively. (B) Relative mRNA expression of *Cdkn1b* in E18.5 wild-type (WT) and *Cdkn1b*^{-/-}

ovaries. The data are represented as mean $\pm$ SEM (* $p < 0.05$). (C) Cellular localisation of p27^{kip1} in

wild-type (WT) ovaries and *Cdkn1b*^{-/-} ovaries at E18.5. Ovaries were stained for p27^{kip1} at E18.5 by

immunofluorescence. Images were taken at 20 x magnification, with scale bars of 50 μm. (D) Cellular

localisation of DDX4 and FOXL2 in E18.5 wild-type (WT) and *Cdkn1b*^{-/-} ovaries. Ovaries were stained

with the oocyte marker DDX4 and the granulosa cell marker FOXL2 at E18.5 by immunofluorescence.

Images were taken at 20 x magnification, with scale bars of 50 μm. (E) MA plot indicates the differential

expression of gene transcripts between WT E18.5 dissociated ovarian cells and *Cdkn1b*^{-/-} dissociated

ovarian cells (n = 4 replicates per group). Genes upregulated in *Cdkn1b*^{-/-} ovaries are shown in blue and

genes downregulated in *Cdkn1b*^{-/-} ovaries shown in green. Grey dots indicate genes below the $p \leq 0.05$

cutoff. (F) Gene ontology (GO) information of the biological process categories with top 5 highest

count of genes from the genes differentially regulated in *Cdkn1b*^{-/-} ovaries. The list of genes was

identified by differential gene expression analysis ($p \leq 0.05$) and annotated via GO analysis in DAVID

(v6.8).

To confirm if the differentially expressed genes detected in *Cdkn1b*^{-/-} ovaries at E18.5 were

altered due to changes in the activation status of granulosa cells, we overlaid the *Cdkn1b*^{-/-} bulk

sequencing data on the single-cell sequencing data to identify genes that were dysregulated in

Cdkn1b^{-/-} ovaries at E18.5 and are expressed in granulosa cell clusters in the wild-type context

(Figure 8A). This identified that the majority of differentially expressed genes in *Cdkn1b*^{-/-}

ovaries at E18.5 were expressed in granulosa cell clusters Gc_3 and Gc_4 and mesenchymal

clusters Mc_3, Mc_4 and Mc_5 in the wild-type ovary (Figure 8A). To better understand the

genes that are upregulated in granulosa cells of *Cdkn1b*^{-/-} ovaries, we compared the 1,008

differentially expressed genes that were upregulated in *Cdkn1b*^{-/-} ovaries with the 274 genes

upregulated in activating pregranulosa cells (Gc_3) (Figure 8B and S6 Table). 60 genes were

present in both groups, with most of these factors (70%) relating to cell cycle processes. This

suggested that the gene expression signature of activation that we see present in Gc_3 cells in

the single-cell experiment is similarly present in *Cdkn1b*^{-/-} ovaries at E18.5 (Figure 8B and S6

Table). Moreover, this is developmentally earlier than primordial follicle activation in wild-

type ovaries, shown by the proportion of Gc_3 cells originating mostly from PND4 and PND7

(Figure S2). In summary, primordial follicle activation begins earlier in

Cdkn1b^{-/-} ovaries, with the altered expression of genes more representative of activating

pregranulosa cells, rather than BPGs or EPGs (Figure 8B). To check if activated granulosa cell

genes could also be identified in E18.5 *Cdkn1b*^{-/-} ovaries, we compared the 1,008 differentially

expressed genes that were upregulated in E18.5 *Cdkn1b*^{-/-} ovaries with the 278 genes

upregulated in differentiated granulosa cells (Gc_4) (Figure 8C and S7 Table). Indeed, 141

genes were present in both groups, with several of these genes being known targets of $p27^{kip1}$,

including *Ccnb1*, *Aurka* and *Aurkb*. Taken together, these findings show that the genes

associated with both activating and activated pregranulosa cells are precociously expressed in

Cdkn1b^{-/-} ovaries at E18.5, despite no gross morphological changes in germ cell nest structures.

Figure 8. Precocious gene expression of differentiated granulosa cells in *Cdkn1b$^{-/-}$* ovaries at E18.5

(A) Heat map showing the expression of differentially expressed genes in *Cdkn1b*^{-/-} ovaries (p ≤ 0.05) in the context of the cell clusters from the single-cell sequencing experiment. Genes highly expressed in each cluster are shown in red and genes lowly expressed in each cluster are shown in yellow. Clusters are labelled across the top of the heatmap and coloured according to the UMAP. The logFC values for genes in the bulk *Cdkn1b*^{-/-} experiment are shown on the right side of the UMAP, and clustered according to their pattern of expression. Upregulated genes are shown in blue and downregulated genes are shown in green. (B) Venn diagram comparing the 274 genes differentially expressed in Gc_3 cells vs. Gc_2 cells (red circle) with the 1,008 genes upregulated in E18.5 *Cdkn1b*^{-/-} ovaries vs. E18.5 WT ovaries (blue circle). 60 genes upregulated in Gc_3 cells were also upregulated in *Cdkn1b*^{-/-} ovaries. For cell clusters, genes were identified as differentially expressed by single-cell RNA sequencing ($\log_2$ fold change ± 2; p ≤ 0.05) and for *Cdkn1b*^{-/-} ovaries genes were identified by bulk RNA-sequencing (p ≤ 0.05). (C) Venn diagram comparing the 278 genes differentially expressed in Gc_4 cells vs. Gc_3 cells (orange circle) with the 1,008 genes upregulated in E18.5 *Cdkn1b*^{-/-} ovaries vs. E18.5 WT ovaries (red circle). 141 genes that are upregulated in Gc_4 cells are also upregulated in *Cdkn1b*^{-/-} ovaries. For cell clusters, genes were identified as differentially expressed by single-cell RNA sequencing ($\log_2$ fold change ± 2; p ≤ 0.05) and for *Cdkn1b*^{-/-} ovaries genes were identified by bulk RNA-sequencing (p ≤ 0.05).

Discussion

In this study, we have identified a putative primordial follicle activation gene expression signature in the pregranulosa cells of the mouse ovary. By utilising single cell RNA-sequencing, we were able to resolve five distinct clusters of granulosa cells in three timepoints: E18.5, PND4 and PND7. One of these distinct granulosa cell clusters was identified as a population of activating pregranulosa cells, enabling our examination of the pathways that govern the first wave of primordial follicle activation. Genes which showed specific expression in this cluster of pregranulosa cells were identified and hypothesised to interact to promote and control follicle activation. To functionally interrogate this hypothesis, we utilised the *Cdkn1b*^{-/-} mouse line which has an accelerated primordial follicle activation phenotype in the neonatal mouse ovary. Transcriptomic analysis of these ovaries revealed distinct gene expression changes at E18.5, prior to the normal developmental window of primordial follicle activation. These gene changes had not been observed in *Cdkn1b*^{-/-} ovaries previously.

Differentially expressed genes upregulated in *Cdkn1b⁻/⁻* ovaries were also observed in the gene

expression signature of activating pregranulosa cells, illustrating a precocious primordial

follicle activation process in *Cdkn1b⁻/⁻* ovaries, regulated at least in part by p27^{kip1} maintaining

cell cycle arrest and preventing granulosa cell activation.

A transcriptomic dataset to explore primordial follicle activation in mice

In this study, we have investigated the transcriptome of the embryonic and perinatal developing

mouse ovary. We successfully captured the dominant cell types in the ovary, namely the

oocytes, pregranulosa and mesenchymal cells at three developmental timepoints by analysing

24,810 cells. One advantage of this dataset is the ability to interrogate any ovarian cell type

and its contribution to primordial follicle activation. Additionally, this dataset complements

single-cell datasets generated to examine pre-granulosa cell specification (Niu & Spradling,

2020) and primordial follicle formation (J.-J. Wang et al., 2020). This study provides a valuable

resource for interrogating primordial follicle activation at the transcriptomic level either with

a cell-centric view or focusing on cell-cell interactions. Notably we observed a decrease in the

number of cells in the Oo_1 cluster, particularly at PND7 (Figure 2A). The dissociation

protocol used to prepare the ovarian samples for single cell RNA-sequencing in this study

enriches the suspension for somatic cells, rather than germ cells, via size selection, as oocytes

increase in size as activation occurs and primordial follicles transition to primary follicles

(Frost et al., 2021b). This may explain the decrease in oocyte numbers in the postnatal

timepoints (Zhang et al., 2014). While in the current study, our focus was on identifying

granulosa cell activators of follicle activation, opportunities exist to further analyse this dataset

by investigating receptor/ligand pathways and secreted factors that may be initiated in the

granulosa cells and then activated or passed to the oocytes to facilitate their growth.

Pregranulosa cells cluster separately depending on cellular origin and activation status

We were able to identify five distinct clusters of granulosa cells in three timepoints: at E18.5,

PND4 and PND7, providing supporting evidence for heterogeneity in granulosa cell

populations (Cordeiro et al., 2015; Mork et al., 2012; Niu & Spradling, 2020). Activated

granulosa cells are the simplest granulosa cell type to identify, due to the extensive

characterisation of late-stage follicles and the function of granulosa cells to respond to follicle-

stimulating hormone and produce estrogen (Francois et al., 2017; O'Shaughnessy, McLelland,

& McBride, 1997). Activated granulosa cells also resume proliferation to support follicle

growth and ensure multiple layers of granulosa cells to form cumulus-oocyte complexes

necessary for ovulation (Cannon, Cherian-Shaw, Lovekamp-Swan, & Chaffin, 2007; Rebecca

L. Robker & JoAnne S. Richards, 1998; R. L. Robker & J. S. Richards, 1998). In our dataset,

GC_4 expressed *Amh*, *Nr5a1* and *Esr2*, along with higher levels of *Hsd3b1* and *mKi67*,

characteristic of activated granulosa cells (Figure 2C). The identification of activated granulosa

cells illustrates that our dataset captures not only primordial follicles, but also primary follicles,

which are the second stage of folliculogenesis and represent a follicle stage after primordial

follicle activation.

In addition to identifying activated granulosa cells, we also validated the presence of

populations of both BPG and EPG populations (Figure 3). BPGs express *Foxl2* from E12.5 and

create medullary primordial follicles during germ cell nest breakdown (Mork et al., 2012; Niu

& Spradling, 2020; J.-J. Wang et al., 2020). These cells appear primed to support activation

just after birth, based on the expression of genes including *Hmgcs2*, *Aard* and *Akr1cl*, which

are upregulated further in activated granulosa cells (Figure 3D). EPGs were more closely

clustered with epithelial cells, characterised by strong expression of *Lgr5*, *Gng13* and *Cst8*

(Figure 3B). In studies performed using an Lgr5-CreERT2 mouse line, Niu et al. established

that EPGs contribute to cortical primordial follicles, which support fertility throughout

adulthood (Niu & Spradling, 2020). In our study, the number of EPGs in GC_3 decreased

dramatically in the PND4 and PND7 timepoints (Figure S2B). It is unclear whether our

clustering strategy masks the subtle gene expression differences in EPGs when they upregulate

FOXL2 postnatally, or if EPGs become more transcriptionally similar to bipotential

pregranulosa cells (Pisarska, Barlow, & Kuo, 2011). It is of interest to consider whether the

differences in these two pregranulosa cell types persist into adulthood and whether

differentiated granulosa cells retain specific gene expression signatures relating to their origin.

Lineage tracing studies using *Lgr5*-GFP and/or *Foxl2*-GFP, combined with single-cell

sequencing experiments on later timepoints such as PND21 would shed light on the trajectory

of these cells and how similar or distinct their transcriptomes become.

A signature of gene activity regulating primordial follicle activation in pregranulosa cells

In our analysis, one distinct granulosa cell cluster was observed to be a population of activating

pregranulosa cells, allowing us to map and examine the gene networks that govern the first

wave of primordial follicle activation. This represents one of the first single cell RNA-

sequencing dataset to examine the transcriptomic profile of granulosa cells during the first

wave of primordial follicle activation in the mouse ovary (Y.-Y. Chen et al., 2022). We have

identified a gene expression signature of primordial follicle activation in Gc_3 cells, using

novel bioinformatic differential gene expression approaches. This signature includes *Tnni3* and

Slc18a2. *Tnni3* has been identified in human granulosa cells and genetic variants of *Slc18a2*

are associated with polycystic ovary syndrome (Fan et al., 2019; Q. Li et al., 2016). Further

work to piece together where these genes function and how they cooperate with other

primordial follicle activation networks is essential. It is important to note that the mechanisms

controlling primordial follicle activation are poorly understood. Combining transcriptomic and

proteomic approaches, such as chromatin immunoprecipitation followed by selective isolation

of chromatin-associated proteins (ChIP-SICAP) will yield extensive detail on how these

networks translate to the protein level (Rafiee, Girardot, Sigismondo, & Krijgsveld, 2016).

Post-translational modifications can also be investigated using mass spectrometry, which has

not yet been performed on an early postnatal ovarian time-course to examine timepoint specific

modifications. Mechanisms of activation may not be only limited to gene expression

signatures, with mechanical stress and autophagy recently implicated in follicle activation

(Nagamatsu, Shimamoto, Hamazaki, Nishimura, & Hayashi, 2019; Zhihan et al., 2019). This

transcriptomic dataset provides the essential groundwork for future studies into primordial

follicle activation by providing cell-specific transcriptomic profiles and identifying a profile of

cells before and during follicle activation. Combining this dataset with previously published

single cell RNA-sequencing studies of the developing ovary could establish a time course of

gene expression across the processes of germ cell nest breakdown, primordial follicle

formation and activation (Niu & Spradling, 2020; Stevant et al., 2019; J.-J. Wang et al., 2020).

Exciting opportunities also exist to correlate this data with human ovary single cell RNA-

sequencing studies and that from other species to uncover how well the process is conserved.

Disruption of the genetic primordial follicle activation network in Cdkn1b^{-/-} mice

A valuable inclusion in this study was the functional assessment of our newly identified gene

expression signature against the sequencing profile of a mouse model of infertility. This

combination of single cell RNA-sequencing data with bulk RNA-sequencing supports and

strengthens our characterisation of the genes involved in primordial follicle activation. An

important confirmation was the observation that cell cycle control is not just a side effect of

activation, but cell cycle control genes are integrated in the network that activates primordial

follicles. Both in Gc_3 and Gc_4, cell cycle processes dominate the gene ontology analysis and

illustrate the essential role that cell cycle resumption plays in follicle activation. 36% of the

genes identified in activating pregranulosa cells (Gc_3) and differentiated granulosa cells

(Gc_4) were upregulated in $Cdkn1b^{-/-}$ ovaries. It remains to be elucidated if these changes are

primarily due to the loss of a cell cycle inhibitor or alternatively, a downstream effect of

increased activation. Two genes, $Cep55$ and $Prc1,$ are key factors in cytokinesis, which is

highly relevant in the ovary, because germ cells remain attached in nests through incomplete

cytokinesis until primordial follicle formation (Lei & Spradling, 2013; Rosario et al., 2019).

Higher transcript levels from these genes were observed in the $Cdkn1b^{-/-}$ ovary, which suggests

a developmentally early breakdown of germ cell nests, exacerbating the follicle activation

phenotype previously described (Rajareddy et al., 2007). Stringent experiments using inducible

$Cdkn1b$ gene deletion or overexpression of alternate cell cycle inhibitors may reveal the

importance of cell cycle regulators in the developing ovary.

Questions remain as to what initial signal drives the precocious primordial follicle activation

in $Cdkn1b^{-/-}$ mutants. In this study, a non-cell autonomous effect of p27[kip1] cannot be ruled out,

because p27[kip1] functions in multiple tissues, including the pituitary (Moncho-Amor et al.,

2020). A conditional approach, such as using a $Cdkn1b^{fl/fl};Nr5a1-Cre+$ mouse line, would

provide a clear answer if primordial follicle activation is initiated in the pregranulosa cells and

dysregulated by the absence of p27[kip1]. Regardless, it is clear from this body of work that further

interrogation is necessary to find the master regulators of follicle activation. The hierarchy of

the numerous factors that cause follicle activation are still being explored: as most previous

studies have used single gene-deletion models, which makes the construction of a gene

regulatory network challenging (Adhikari & Liu, 2009; Jagarlamudi, Reddy, Adhikari, & Liu,

2010; Uda et al., 2004). For p27[kip1] in particular, potential upstream regulators that may initiate

primordial follicle activation by reducing p27[kip1] expression include FOXL2, WNT4, SMAD2

and JAK2 (Granados-Aparici et al., 2019; Huang et al., 2018; Jakel, Weinl, & Hengst, 2011;

Maatouk, Mork, Chassot, Chaboissier, & Capel, 2013; Nicol et al., 2018). Additionally, given

that spatial and temporal control of activation is required to selectively activate follicles in

adulthood, epigenetic mechanisms and chromatin accessibility are two unexplored areas that

may provide the precise control necessary to activate multiple factors simultaneously. Future

experiments using ChIP-sequencing and ATAC-sequencing on pregranulosa cells during the

window of activation in the mouse will yield fascinating insights into how the genes

upregulated during primordial follicle activation are controlled.

This study describes a transcriptomic dataset to interrogate primordial follicle activation and

reveals key pathways that are controlling follicle activation in mice. This dataset provides a

platform for characterising a gene regulatory network that regulates primordial follicle

activation. The combination of single cell RNA-sequencing with a bulk RNA-sequencing

dataset in an infertility model reveals disruptions in the follicle activation process, but also

similar pathways that are consistently activated. This combination of transcriptomic analyses

adds a novel bioinformatic approach to this field to understand cell-specific changes in

infertility models. An increased understanding of the factors that act co-operatively to stimulate

or inhibit follicle activation will reveal key insights into how the ovarian reserve is established

and maintained into adulthood and how disruptions in this process lead to infertility conditions.

Materials and Methods

Key resources table

Reagent type (species) or resource	Reagent or resource	Source	Identifier
Antibody	Rabbit monoclonal anti-p27^{kip1}	Cell Signalling Technology	Cat#3686S; RRID: AB_2077850
Antibody	Goat polyclonal anti-FOXL2	Abcam	Cat#ab5096; RRID: AB_304750
Antibody	Rat monoclonal anti-GCNA1	Abcam	Cat#ab82527; RRID: AB_1659152
Antibody	Rabbit polyclonal anti-DDX4	Abcam	Cat#ab13840; RRID: AB_443012
Antibody	Mouse monoclonal anti-DDX4	Abcam	Cat#ab27591; RRID: AB_11139638
Chemical	Paraformaldehyde	Thermofisher Scientific	Cat# 50-980-488
Chemical	ProLong™ Gold Antifade Mountant	Thermofisher Scientific	Cat# P36930
Chemical	Target Retrieval Solution, Citrate pH 6.1	Agilent	Cat#S169984-2
Chemical	Triton X-100	Sigma Aldrich	Cat# T8787
Commercial assay or kit	Chromium Next GEM Single Cell 3′ Kit v3.1	10x Genomics	Cat#1000268
Commercial assay or kit	NEBnext Low Input RNA Library Prep Kit	New England Biolabs	Cat# E6420L
Commercial assay or kit	RNeasy Plus Micro Kit	Qiagen	Cat#74034
Commercial assay or kit	Superscript III	Thermofisher Scientific	Cat#18080093
Genetic reagent (*Mus musculus*)	Mouse: Wildtype (WT; C57BL/6J)	The Jackson Laboratory	Cat#JAX000664; RRID:IMSR_JAX:000664
Genetic reagent (*Mus musculus*)	Mouse: *Cdkn1b*$^{-/-}$ (B6.129S4-Cdkn1b^{tm1Mlf}/J)	The Jackson Laboratory	Cat#JAX002781 RRID:IMSR_JAX:002781
Oligonucleotide	PCR forward primer for mouse *Cdkn1b* WT allele GATGGACGCCAGACAAGC	The Jackson Laboratory	N/A
Oligonucleotide	PCR reverse primer for mouse *Cdkn1b* WT allele CTCCTGCCATTCGTATCTGC	The Jackson Laboratory	N/A
Oligonucleotide	PCR forward primer for mouse *Cdkn1b* mutant allele CTTGGGTGGAGAGGCTATTC	The Jackson Laboratory	N/A
Oligonucleotide	PCR reverse primer for mouse *Cdkn1b* mutant allele AGGTGAGATGACAGGAGATC	The Jackson Laboratory	N/A
Oligonucleotide	qPCR forward primer for mouse *Cdkn1b* gene CAGAAGAAAATGTTTCAGACGGT	This paper	N/A
Oligonucleotide	qPCR reverse primer for mouse *Cdkn1b* gene TCTTAATTCGGAGCTGTTTACGT	This paper	N/A

Oligonucleotide	qPCR forward primer for mouse *Gapdh* gene TTCACCACCATGGAGAAGGC	(Nicol et al., 2018)	N/A
Oligonucleotide	qPCR reverse primer for mouse *Gapdh* gene GGCATGGACTGTGGTCATGA	(Nicol et al., 2018)	N/A
Software and algorithm	Adobe Illustrator CC	Adobe Systems Inc.	https://www.adobe.com/illustrator
Software and algorithm	DAVID Functional Annotation	(Dennis et al., 2003)	https://david.ncifcrf.gov
Software and algorithm	Fiji	(Schindelin et al., 2012)	https://fiji.sc/
Software and algorithm	Prism	GraphPad Inc.	https://www.graphpad.com/scientific-software/prism/
Software and algorithm	R version 3.6.0	R Project for Statistical Computing	https://scicrunch.org/resolver/SCR_001905 RRID:SCR_001905
Software and algorithm	Seurat package 3	R toolkit for single-cell genomics	https://satijalab.org/seurat/articles/install.html
Software and algorithm	Zen software	Zeiss	https://www.zeiss.com/microscopy/int/products/microscope-software/zen.html

Mouse strains, husbandry and genotyping

All experiments carried out on mice were approved under the UK Animal (scientific

procedures) Act 1986 (Project license 70/8560). Animals were kept in individually ventilated

cages (IVC) with access to food and water *ad libitum*. Mouse husbandry, ear biopsies, breeding

and vaginal plug (VP) checks were performed by the Biological Research Facility team of the

Francis Crick Institute. The VP day was considered as 0.5 day from time of conception (E0.5)

and the day of birth considered P0. Adult female animals were euthanised by cervical

dislocation and ovaries were collected immediately following euthanasia. Embryos were

euthanised by decapitation and gonads were collected with the adjacent mesonephros. For

RNA-sequencing, the mesonephros was removed from the ovary using dissecting forceps.

C57BL/6J mice (RRID: IMSR_JAX:000664) were supplied by the Biological Research

Facility at the Francis Crick Institute and used for the single-cell sequencing experiments.

Cdkn1b^{tm1Mlf} (*Cdkn1b^{-/-}*) (RRID: IMSR_JAX:002781) (Fero et al., 1996) were maintained on

a mixed background and mated heterozygously. Experiments involving *Cdkn1b^{-/-}* mice were

performed using littermate animals as controls. Genotyping of adult *Cdkn1b*[-/-] mice was

performed by Transnetyx®. Embryos from heterozygous matings were PCR genotyped using

primers for the WT allele and mutant allele as previously published (Fero et al., 1996).

RNA sequencing sample preparation

Tissue dissociation and single-cell library preparation

Female ovaries at E18.5, PND1 and PND7 were harvested and the ovarian bursa and fat were

removed, leaving only ovary tissue. Ovarian tissue was dissociated as previously described

(Frost et al., 2021b). Briefly, two ovaries from each timepoint were incubated separately in a

solution of papain (10108014001, Sigma-Aldrich, 1mg/ml in HBSS) for 15min at 37°C in

presence of 10µg/ml of both DNAse I (10104159001, Thermo Scientific) and Rock Inhibitor

(M1817, Abmole Bioscience) (Moncho-Amor et al., 2020). The papain solution containing the

cells was diluted with 1x HBSS containing 0.0025% BSA and 0.01% 1M Tris pH8.0. The cell

solution was then mechanically dissociated using a 10mL glass serological pipette, and passed

through a 20µM pre-separation filter (130-101-812, Miltenyi Biotec). Cells were counted using

the EVE Cell counter (NanoEnTek, Cambridge Bioscience), stained with trypan blue to

measure viability. Approximately 10,000 live cells were used to generate libraries using the

Chromium Single Cell 3' kit v3.1, according to the manufacturer's instructions (10x Genomics,

1000268) (G. X. Y. Zheng et al., 2017). The libraries were quantified using the TapeStation

(Agilent) to confirm quantity and purity, before sequencing on an Hiseq4000 (Illumina).

Sequencing outputs were processed using the 10x CellRanger (version.3.0.2) to generate

single-cell count data for each time point, using a mouse reference index provided by 10x

Genomics (refdata-cellranger-mm10v3.0). All subsequent analyses were performed in R

v.3.6.0 using the Seurat (v3) package (Stuart et al., 2019).

Tissue dissociation and bulk library preparation

Ovaries from *Cdkn1b*$^{+/+}$ and *Cdkn1b*$^{-/-}$ mice were collected at E13.5, E18.5 and PND8 and the

ovarian bursa and fat were removed. A single ovary was dissociated as described above. Cells

were counted using the EVE Cell counter (NanoEnTek, Cambridge Bioscience) and stained

with trypan blue to measure viability. RNA was extracted from the dissociated cells using the

RNeasy Plus Micro kit (Qiagen) according to the manufacturer's protocol. RNA quality was

assessed the TapeStation (Agilent) to confirm quantity and purity. For E13.5 and E18.5 ovaries,

libraries were constructed using the NEBNext Low Input RNA Library Prep Kit (E6420L, New

England Biolabs) according to the manufacturer's instructions. For PND8 ovaries, libraries

were constructed using the polyA KAPA mRNA Hyper Prep Kit (Roche) according to the

manufacturer's instructions. All libraries were quantified using the TapeStation (Agilent) and

sequenced on an Hiseq400 (Illumina) to achieve an average of 25 million reads per sample.

scRNAseq Bioinformatics analysis

Raw single-cell sequencing data from all three timepoints were processed using the 10x

CellRanger pipeline (10x Genomics). Count tables were loaded into R and analysed with the

Seurat package (Stuart et al., 2019). All cells with fewer than 1500 RNA features were

removed. Principal component analysis was performed on variable genes and 20 principal

components were used to generate the UMAP analysis. The UMAP clustering plot was

performed using the default method from the Seurat package. Differential Gene Expression

(DGE) analysis was performed as previously described (Ahlmann-Eltze & Huber, 2020).

DAVID Gene Ontology (GO) functional annotation tools were utilised to explore the biological

processes associated with the genes differentially expressed in specific clusters (Dennis et al.,

2003).

Bulk RNA sequencing analysis

Twelve libraries were sequenced on an Illumina HiSeq 4000 machine. The 'Trim Galore!'

utility version 0.4.2 was used to remove sequencing adaptors and to trim individual reads with

the q-parameter set to 20 (https://www.bioinformatics.babraham.ac.uk/projects/trim_galore/).

The sequencing reads were then aligned to the mouse genome and transcriptome (Ensembl

GRCm38release-89) using RSEM version 1.3.0 (B. Li & Dewey, 2011) in conjunction with

the STAR aligner version 2.5.2 (Dobin et al., 2013). The sequencing quality of individual

samples was assessed using FASTQC version 0.11.5

(https://www.bioinformatics.babraham.ac.uk/projects/fastqc/) and RNA-SeQC version 1.1.8

(DeLuca et al., 2012). Differential gene expression was determined using the R-bioconductor

package DESeq2 version 1.14.1 (Development & Team, 2008; Love, Huber, & Anders, 2014).

Gene set enrichment analysis (GSEA) was conducted as described (Subramanian et al., 2005).

Immunofluorescence and in situ hybridisation

Ovaries for immunofluorescence were fixed in chilled 4 % paraformaldehyde at 4 °C for 1 to

2 hrs. Following fixation, ovaries were washed in PBS 3 times for 1 hr each, before immersing

in 30 % sucrose in PBS overnight to cryoprotect the sample. The ovaries were then embedded

in OCT mounting medium, frozen on an ethanol/dry ice slurry and stored at -80 °C. Samples

were cryosectioned at 10 μm and sections were placed on Superfrost Plus glass slides and air

dried for 5 minutes, before storing at -80 °C. Before starting immunofluorescence, slides were

air dried for 5 minutes and then washed 3 times for 5 mins each in PBS. Antigen retrieval was

performed by immersing slides in 10 % target retrieval solution pH 6.1 (S169984-2, Agilent)

with distilled water. Antigen retrieval was done for 30 mins at 65 °C. Slides were washed 3

times in PBS for 5 mins and then incubated in blocking solution (10 % donkey serum in 0.1 %

triton X-100 PBS) for 2 hr at room temperature. Primary antibodies were added to the slide,

diluted in 10% blocking solution, and left overnight at 4 °C. Primary antibodies were used at

the following dilutions: rabbit anti-p27^{kip1} (D69C12, Cell Signalling) at 1:200, goat anti-

FOXL2 (ab5096, Abcam) at 1:100, goat anti-SOX9 (AF3075, R&D Systems) at 1:200, rat anti-

GCNA1 (ab82527, Abcam) at 1:200, rabbit anti-DDX4 (ab13840, Abcam) at 1:200 and mouse

anti-DDX4 (ab27591, Abcam) at 1:200. The next day, the slides were washed 3 times in 0.1 %

triton X-100 PBS for 5 minutes. Secondary antibodies were diluted in blocking solution (10 %

donkey serum in 0.1 % triton X-100 PBS) and slides were incubated with the appropriate

secondary antibodies for 1 hr at room temperature. The following secondary antibodies were

used at a 1:200 dilution: Donkey anti-rabbit 488 (Invitrogen, A21206), Donkey anti-goat 647

(Invitrogen, A21447), Donkey anti-mouse 647 (Invitrogen, A-31571) and Donkey anti-rat 647

(Abcam, ab150155). The slides were then washed in 0.1 % triton X-100 PBS for 5 minutes 3

times, then the slides were counter-stained with 4'-6-diamino-2-phenylindole (DAPI). Finally,

sections were washed once in distilled water, air dried and coverslip mounted with Prolong

Gold Antifade Mountant (P36930, Thermo Scientific) and Aqua-Poly/Mount (Polysciences,

Inc., Warrington, PA, USA). Images were taken using a Leica SPE confocal microscope and

processed using Fiji.

Histology staining
721
Ovaries for hematoxylin and eosin (H&E) staining were fixed in chilled 4 % paraformaldehyde

at 4 °C for 1 to 2 hrs. Following fixation, ovaries were washed in PBS 3 times for 1 hr each,

before immersing in 30 % sucrose in PBS overnight to cryoprotect the sample. The ovaries

were then embedded in OCT mounting medium, frozen on an ethanol/dry ice slurry and stored

at -80 °C. Samples were cryosectioned at 10 μm and sections were placed on Superfrost Plus

glass slides and air dried for 5 minutes. H&E staining was performed by the Francis Crick

Institute Experimental Histopathology facility.

RNA extraction and qPCR
730
Total mRNA was extracted from individual ovaries using the RNeasy Plus Micro kit (74034,

Qiagen) according to the manufacturer's protocol. Total mRNA was quantified using the

NanoDrop™ 2000 Spectrophotometer (Thermo Fisher Scientific). After quantification, 50 ng

of RNA was reverse transcribed into cDNA using the Superscript III cDNA synthesis kit

(Thermo Fisher Scientific) as per the manufacturer's instructions. Quantitative PCR was

performed on cDNA, with primer sequences listed in the key resources table. Reactions were

performed in triplicate using GoTaq® qPCR Master Mix (Promega) and 100 nM each of

forward and reverse primers and analysed on the LightCycler® 96 Instrument (Roche).

Relative mRNA levels were determined by calculating $2^{-\Delta\Delta Ct}$ values relative to the

housekeeping gene *Gapdh*. Data is shown as mean ± SEM from four biological replicates.

Statistical analyses

Statistical analyses were performed using Prism v9.0 (GraphPad Software, USA). Student's

two-sided unpaired *t-tests* were used when comparing two groups of normally distributed

values. Mann Whitney tests were used when comparing two groups of non-parametric data.

For multiple comparisons, analysis of variance (ANOVA) was performed. All results are

represented as mean ± standard error of the mean (SEM) and standard significance levels were

used: $*p<0.05$, $**p<0.01$, $***p<0.001$, $****p<0.0001$.

Data availability

The data supporting the findings of this study are available in the main text and the

supplementary materials. All sequencing data will be made publicly available after the

publication of this article.

Acknowledgements

We would like to thank members of the Lovell-Badge and Sutherland laboratories for their

comments on the manuscript. We thank the Advanced Sequencing Facility at the Francis Crick

Institute for their help in performing the sequencing studies in this manuscript. This work was

supported by the Francis Crick Institute, which receives its core funding from Cancer Research

UK (FC001107), the UK Medical Research Council (FC001107), and the Wellcome Trust

(FC001107). We acknowledge the generous financial assistance to R.L.B. by the UK

Biotechnology and Biological Sciences Research Council (BB/ N018680/1). We also

gratefully acknowledge the financial assistance to J.M.S. and M.A.B. by the Australian

National Health and Medical Research Council (G1600095) and the Hunter Medical Research

Institute (G1501433 and G1801335). E.R.F. is the recipient of an Australian Government

Research Training Program Scholarship.

Author contributions

766 E.R.F and G.T conceived the study and wrote the manuscript. J.M.S, C.G, M.A.B and R.L.B

767 provided expertise and feedback on the study design. All the authors critically revised the

768 manuscript. All experiments were performed by E.R.F except for the following: G.T and E.R.F

769 jointly prepared samples for single-cell RNA sequencing. S.B performed both RNA-seq and

770 single-cell RNA-seq analysis. J.M.S and R.L.B provided resources for this work.

Declaration of interests

772 The authors declare no competing interests.

References

Adhikari, D., & Liu, K. (2009). Molecular mechanisms underlying the activation of mammalian
primordial follicles. *Endocr Rev, 30*(5), 438-464. doi:10.1210/er.2008-0048
Ahlmann-Eltze, C., & Huber, W. (2020). glmGamPoi: fitting Gamma-Poisson generalized linear
models on single cell count data. *Bioinformatics, 36*(24), 5701-5702.
doi:10.1093/bioinformatics/btaa1009
Bagheri-Fam, S., Chen, H., Wilson, S., Ayers, K., Hughes, J., Sloan-Bena, F., . . . Wilhelm, D.
(2020). The gene encoding the ketogenic enzyme HMGCS2 displays a unique expression
during gonad development in mice. *PLOS ONE, 15*(1), e0227411.
doi:10.1371/journal.pone.0227411
Cannon, J. D., Cherian-Shaw, M., Lovekamp-Swan, T., & Chaffin, C. L. (2007). Granulosa cell
expression of G1/S phase cyclins and cyclin-dependent kinases in PMSG-induced follicle
growth. *Mol Cell Endocrinol, 264*(1-2), 6-15. doi:10.1016/j.mce.2006.09.015
Chen, Y., Yang, W., Shi, X., Zhang, C., Song, G., & Huang, D. (2020). The Factors and Pathways
Regulating the Activation of Mammalian Primordial Follicles in vivo. *Front Cell Dev Biol,
8*(1018). doi:10.3389/fcell.2020.575706
Chen, Y.-Y., Russo, D. D., Drake, R. S., Duncan, F. E., Shalek, A. K., Goods, B. A., & Woodruff, T.
K. (2022). Single-cell transcriptomics of staged oocytes and somatic cells reveal novel
regulators of follicle activation. *Reproduction, 164*(2), 55-70. doi:10.1530/rep-22-0053
Cooke, H. J., Lee, M., Kerr, S., & Ruggiu, M. (1996). A murine homologue of the human DAZ gene
is autosomal and expressed only in male and female gonads. *Hum Mol Genet, 5*(4), 513-516.
doi:10.1093/hmg/5.4.513
Cordeiro, M. H., Kim, S. Y., Ebbert, K., Duncan, F. E., Ramalho-Santos, J., & Woodruff, T. K.
(2015). Geography of follicle formation in the embryonic mouse ovary impacts activation
pattern during the first wave of folliculogenesis. *Biology of Reproduction, 93*(4), 88.
doi:10.1095/biolreprod.115.131227

DeLuca, D. S., Levin, J. Z., Sivachenko, A., Fennell, T., Nazaire, M.-D., Williams, C., . . . Getz, G. (2012). RNA-SeQC: RNA-seq metrics for quality control and process optimization. *Bioinformatics, 28*(11), 1530-1532. doi:10.1093/bioinformatics/bts196

Dennis, G., Sherman, B. T., Hosack, D. A., Yang, J., Gao, W., Lane, H. C., & Lempicki, R. A. (2003). DAVID: Database for Annotation, Visualization, and Integrated Discovery. *Genome Biology, 4*(5), P3. doi:10.1186/gb-2003-4-5-p3

Development, R. C., & Team. (2008). *R: A language and environment for statistical computing.* Vienna, Austria: R Foundation for Statistical Computing.

Dobin, A., Davis, C. A., Schlesinger, F., Drenkow, J., Zaleski, C., Jha, S., . . . Gingeras, T. R. (2013). STAR: ultrafast universal RNA-seq aligner. *Bioinformatics, 29*(1), 15-21. doi:10.1093/bioinformatics/bts635

Fan, X., Bialecka, M., Moustakas, I., Lam, E., Torrens-Juaneda, V., Borggreven, N. V., . . . Chuva de Sousa Lopes, S. M. (2019). Single-cell reconstruction of follicular remodeling in the human adult ovary. *Nat Commun, 10*(1), 3164. doi:10.1038/s41467-019-11036-9

Fero, M. L., Rivkin, M., Tasch, M., Porter, P., Carow, C. E., Firpo, E., . . . Roberts, J. M. (1996). A syndrome of multiorgan hyperplasia with features of gigantism, tumorigenesis, and female sterility in p27(Kip1)-deficient mice. *Cell, 85*(5), 733-744. Retrieved from https://ac.els-cdn.com/S0092867400812398/1-s2.0-S0092867400812398-main.pdf?_tid=81ed8a97-9c51-4fe7-82d6-725b355e7e1c&acdnat=1548794809_c846c788db4da96843436c6939aab1ba

Francois, C. M., Petit, F., Giton, F., Gougeon, A., Ravel, C., Magre, S., . . . Guigon, C. J. (2017). A novel action of follicle-stimulating hormone in the ovary promotes estradiol production without inducing excessive follicular growth before puberty. *Sci Rep, 7*, 46222. doi:10.1038/srep46222

Frost, E. R., Ford, E. A., Taylor, G., Boeing, S., Beckett, E. L., Roman, S. D., . . . Sutherland, J. M. (2021a). Two alternative methods for the retrieval of somatic cell populations from the mouse ovary. *Molecular Human Reproduction, 27*(6). doi:10.1093/molehr/gaab033

Frost, E. R., Ford, E. A., Taylor, G., Boeing, S., Beckett, E. L., Roman, S. D., . . . Sutherland, J. M. (2021b). Two alternative methods for the retrieval of somatic cell populations from the mouse ovary. *Molecular Human Reproduction.* doi:10.1093/molehr/gaab033

Fukuda, K., Muraoka, M., Kato, Y., & Saga, Y. (2021). Decoding the transcriptome of pre-granulosa cells during the formation of primordial follicles in the mouse. *Biology of Reproduction.* doi:10.1093/biolre/ioab065

Gaytan, F., Morales, C., Roa, J., & Tena-Sempere, M. (2018). Changes in keratin 8/18 expression in human granulosa cell lineage are associated to cell death/survival events: potential implications for the maintenance of the ovarian reserve. *Hum Reprod, 33*(4), 680-689. doi:10.1093/humrep/dey010

Granados-Aparici, S., Hardy, K., Franks, S., Sharum, I. B., Waite, S. L., & Fenwick, M. A. (2019). SMAD3 directly regulates cell cycle genes to maintain arrest in granulosa cells of mouse primordial follicles. *Sci Rep, 9*(1), 6513. doi:10.1038/s41598-019-42878-4

Greenberg, Z. J., Monlish, D. A., Bartnett, R. L., Yang, Y., Shen, G., Li, W., . . . Schuettpelz, L. G. (2020). The Tetraspanin CD53 Regulates Early B Cell Development by Promoting IL-7R Signaling. *The Journal of Immunology, 204*(1), 58-67. doi:10.4049/jimmunol.1900539

Hardy, K., Mora, J. M., Dunlop, C., Carzaniga, R., Franks, S., & Fenwick, M. A. (2018). Nuclear exclusion of SMAD2/3 in granulosa cells is associated with primordial follicle activation in the mouse ovary. *Journal of Cell Science, 131*(17). doi:10.1242/jcs.218123

Huang, K., Wang, Y., Zhang, T., He, M., Sun, G., Wen, J., . . . Wang, C. (2018). JAK signaling regulates germline cyst breakdown and primordial follicle formation in mice. *Biology Open, 7*(1), bio029470. doi:10.1242/bio.029470

Hutt, K. J., McLaughlin, E. A., & Holland, M. K. (2006). KIT/KIT Ligand in Mammalian Oogenesis and Folliculogenesis: Roles in Rabbit and Murine Ovarian Follicle Activation and Oocyte Growth1. *Biology of Reproduction, 75*(3), 421-433. doi:10.1095/biolreprod.106.051516

Jagarlamudi, K., Reddy, P., Adhikari, D., & Liu, K. (2010). Genetically modified mouse models for premature ovarian failure (POF). *Mol Cell Endocrinol, 315*(1-2), 1-10. doi:10.1016/j.mce.2009.07.016

Jakel, H., Weinl, C., & Hengst, L. (2011). Phosphorylation of p27Kip1 by JAK2 directly links
cytokine receptor signaling to cell cycle control. *Oncogene, 30*(32), 3502-3512.
doi:10.1038/onc.2011.68
John, G. B., Gallardo, T. D., Shirley, L. J., & Castrillon, D. H. (2008). Foxo3 is a PI3K-dependent
molecular switch controlling the initiation of oocyte growth. *Dev Biol, 321*(1), 197-204.
doi:10.1016/j.ydbio.2008.06.017
Jones, R. L., & Pepling, M. E. (2013). KIT signaling regulates primordial follicle formation in the
neonatal mouse ovary. *Dev Biol, 382*(1), 186-197. doi:10.1016/j.ydbio.2013.06.030
Kezele, P. R., Ague, J. M., Nilsson, E., & Skinner, M. K. (2005). Alterations in the ovarian
transcriptome during primordial follicle assembly and development. *Biology of Reproduction,*
72(1), 241-255. doi:10.1095/biolreprod.104.032060
Landsman, L., Bar-On, L., Zernecke, A., Kim, K.-W., Krauthgamer, R., Shagdarsuren, E., . . . Jung,
S. (2009). CX3CR1 is required for monocyte homeostasis and atherogenesis by promoting
cell survival. *Blood, 113*(4), 963-972. doi:10.1182/blood-2008-07-170787
Lei, L., & Spradling, A. C. (2013). Mouse primordial germ cells produce cysts that partially fragment
prior to meiosis. *Development, 140*(10), 2075-2081. doi:10.1242/dev.093864
Li, B., & Dewey, C. N. (2011). RSEM: accurate transcript quantification from RNA-Seq data with or
without a reference genome. *BMC Bioinformatics, 12*(1), 323. doi:10.1186/1471-2105-12-323
Li, Q., Yan, Z., Kuang, Y., Zhou, X., Jin, L., He, L., . . . Wang, L. (2016). Genetic variations in the 3'-
untranslated region of SLC18A2 are associated with serum FSH concentration in polycystic
ovary syndrome patients and regulate gene expression in vitro. *Hum Reprod, 31*(9), 2150-
2157. doi:10.1093/humrep/dew162
Love, M. I., Huber, W., & Anders, S. (2014). Moderated estimation of fold change and dispersion for
RNA-seq data with DESeq2. *Genome Biology, 15*(12), 550. doi:10.1186/s13059-014-0550-8
Ma, S. C., Li, Q., Peng, J. Y., Zhouwen, J. L., Diao, J. F., Niu, J. X., . . . Jiang, W. G. (2017). Claudin-
5 regulates blood-brain barrier permeability by modifying brain microvascular endothelial
cell proliferation, migration, and adhesion to prevent lung cancer metastasis. *CNS Neurosci*
Ther, 23(12), 947-960. doi:10.1111/cns.12764
Maatouk, D. M., Mork, L., Chassot, A. A., Chaboissier, M. C., & Capel, B. (2013). Disruption of
mitotic arrest precedes precocious differentiation and transdifferentiation of pregranulosa
cells in the perinatal Wnt4 mutant ovary. *Dev Biol, 383*(2), 295-306.
doi:10.1016/j.ydbio.2013.08.026
McLaughlin, E. A., & McIver, S. C. (2009). Awakening the oocyte: controlling primordial follicle
development. *Reproduction, 137*(1), 1-11. doi:10.1530/rep-08-0118
Meinsohn, M. C., Hughes, C. H. K., Estienne, A., Saatcioglu, H. D., Pépin, D., Duggavathi, R., &
Murphy, B. D. (2021). A role for orphan nuclear receptor liver receptor homolog-1 (LRH-1,
NR5A2) in primordial follicle activation. *Sci Rep, 11*(1), 1079. doi:10.1038/s41598-020-
80178-4
Miller, I., Min, M., Yang, C., Tian, C., Gookin, S., Carter, D., & Spencer, S. L. (2018). Ki67 is a
Graded Rather than a Binary Marker of Proliferation versus Quiescence. *Cell Rep, 24*(5),
1105-1112.e1105. doi:10.1016/j.celrep.2018.06.110
Moncho-Amor, V., Chakravarty, P., Galichet, C., Matheu, A., Lovell-Badge, R., & Rizzoti, K.
(2020). SOX2 is required independently in both stem and differentiated cells for pituitary
tumorigenesis in p27 null mice. *BioRxiv*, 2020.2006.2009.142638.
doi:10.1101/2020.06.09.142638
Monniaux, D., Clement, F., Dalbies-Tran, R., Estienne, A., Fabre, S., Mansanet, C., & Monget, P.
(2014). The ovarian reserve of primordial follicles and the dynamic reserve of antral growing
follicles: what is the link? *Biology of Reproduction, 90*(4), 85.
doi:10.1095/biolreprod.113.117077
Mork, L., Maatouk, D. M., McMahon, J. A., Guo, J. J., Zhang, P., McMahon, A. P., & Capel, B.
(2012). Temporal differences in granulosa cell specification in the ovary reflect distinct
follicle fates in mice. *Biology of Reproduction, 86*(2), 37. doi:10.1095/biolreprod.111.095208
Nagamatsu, G., Shimamoto, S., Hamazaki, N., Nishimura, Y., & Hayashi, K. (2019). Mechanical
stress accompanied with nuclear rotation is involved in the dormant state of mouse oocytes.
Sci Adv, 5(6), eaav9960. doi:10.1126/sciadv.aav9960

Nicol, B., Grimm, S. A., Chalmel, F., Lecluze, E., Pannetier, M., Pailhoux, E., . . . Yao, H. H. (2019). RUNX1 maintains the identity of the fetal ovary through an interplay with FOXL2. *Nat Commun, 10*(1), 5116. doi:10.1038/s41467-019-13060-1

Nicol, B., Grimm, S. A., Gruzdev, A., Scott, G. J., Ray, M. K., & Yao, H. H. (2018). Genome-wide identification of FOXL2 binding and characterization of FOXL2 feminizing action in the fetal gonads. *Hum Mol Genet, 27*(24), 4273-4287. doi:10.1093/hmg/ddy312

Niu, W., & Spradling, A. C. (2020). Two distinct pathways of pregranulosa cell differentiation support follicle formation in the mouse ovary. *Proceedings of the National Academy of Sciences, 117*(33), 20015-20026. doi:10.1073/pnas.2005570117

O'Shaughnessy, P. J., McLelland, D., & McBride, M. W. (1997). Regulation of luteinizing hormone-receptor and follicle-stimulating hormone-receptor messenger ribonucleic acid levels during development in the neonatal mouse ovary. *Biology of Reproduction, 57*(3), 602-608. doi:10.1095/biolreprod57.3.602

Ottolenghi, C., Omari, S., Garcia-Ortiz, J. E., Uda, M., Crisponi, L., Forabosco, A., . . . Schlessinger, D. (2005). Foxl2 is required for commitment to ovary differentiation. *Hum Mol Genet, 14*(14), 2053-2062. doi:10.1093/hmg/ddi210

Pepling, M. E., & Spradling, A. C. (1998). Female mouse germ cells form synchronously dividing cysts. *Development, 125*(17), 3323-3328.

Pepling, M. E., & Spradling, A. C. (2001). Mouse ovarian germ cell cysts undergo programmed breakdown to form primordial follicles. *Dev Biol, 234*(2), 339-351. doi:10.1006/dbio.2001.0269

Perez-Sanz, J., Arluzea, J., Matorras, R., Gonzalez-Santiago, N., Bilbao, J., Yeh, N., . . . de la Hoz, C. (2013). Increased number of multi-oocyte follicles (MOFs) in juvenile p27Kip1 mutant mice: potential role of granulosa cells. *Hum Reprod, 28*(4), 1023-1030. doi:10.1093/humrep/des436

Pisarska, M. D., Barlow, G., & Kuo, F. T. (2011). Minireview: roles of the forkhead transcription factor FOXL2 in granulosa cell biology and pathology. *Endocrinology, 152*(4), 1199-1208. doi:10.1210/en.2010-1041

Rafiee, M. R., Girardot, C., Sigismondo, G., & Krijgsveld, J. (2016). Expanding the Circuitry of Pluripotency by Selective Isolation of Chromatin-Associated Proteins. *Mol Cell, 64*(3), 624-635. doi:10.1016/j.molcel.2016.09.019

Rajareddy, S., Reddy, P., Du, C., Liu, L., Jagarlamudi, K., Tang, W., . . . Liu, K. (2007). p27kip1 (cyclin-dependent kinase inhibitor 1B) controls ovarian development by suppressing follicle endowment and activation and promoting follicle atresia in mice. *Mol Endocrinol, 21*(9), 2189-2202. doi:10.1210/me.2007-0172

Rastetter, R. H., Bernard, P., Palmer, J. S., Chassot, A. A., Chen, H., Western, P. S., . . . Wilhelm, D. (2014). Marker genes identify three somatic cell types in the fetal mouse ovary. *Dev Biol, 394*(2), 242-252. doi:10.1016/j.ydbio.2014.08.013

Reynolds, N., Collier, B., Bingham, V., Gray, N. K., & Cooke, H. J. (2007). Translation of the synaptonemal complex component Sycp3 is enhanced in vivo by the germ cell specific regulator Dazl. *Rna, 13*(7), 974-981. doi:10.1261/rna.465507

Robker, R. L., & Richards, J. S. (1998). Hormonal Control of the Cell Cycle in Ovarian Cells: Proliferation Versus Differentiation. *Biology of Reproduction, 59*(3), 476-482. doi:10.1095/biolreprod59.3.476

Robker, R. L., & Richards, J. S. (1998). Hormone-induced proliferation and differentiation of granulosa cells: a coordinated balance of the cell cycle regulators cyclin D2 and p27Kip1. *Mol Endocrinol, 12*(7), 924-940. doi:10.1210/mend.12.7.0138

Rosario, R., Crichton, J. H., Stewart, H. L., Childs, A. J., Adams, I. R., & Anderson, R. A. (2019). Dazl determines primordial follicle formation through the translational regulation of Tex14. *FASEB Journal, 33*(12), 14221-14233. doi:10.1096/fj.201901247R

Schindelin, J., Arganda-Carreras, I., Frise, E., Kaynig, V., Longair, M., Pietzsch, T., . . . Cardona, A. (2012). Fiji: an open-source platform for biological-image analysis. *Nat Methods, 9*(7), 676-682. doi:10.1038/nmeth.2019

Sobinoff, A. P., Sutherland, J. M., & McLaughlin, E. A. (2013). Intracellular signalling during female gametogenesis. *Mol Hum Reprod, 19*(5), 265-278. doi:10.1093/molehr/gas065

Stevant, I., Kuhne, F., Greenfield, A., Chaboissier, M. C., Dermitzakis, E. T., & Nef, S. (2019).
Dissecting Cell Lineage Specification and Sex Fate Determination in Gonadal Somatic Cells
Using Single-Cell Transcriptomics. *Cell Rep, 26*(12), 3272-3283.e3273.
doi:10.1016/j.celrep.2019.02.069
Stuart, T., Butler, A., Hoffman, P., Hafemeister, C., Papalexi, E., Mauck, W. M., 3rd, . . . Satija, R.
(2019). Comprehensive Integration of Single-Cell Data. *Cell, 177*(7), 1888-1902.e1821.
doi:10.1016/j.cell.2019.05.031
Subramanian, A., Tamayo, P., Mootha, V. K., Mukherjee, S., Ebert, B. L., Gillette, M. A., . . .
Mesirov, J. P. (2005). Gene set enrichment analysis: A knowledge-based approach for
interpreting genome-wide expression profiles. *Proceedings of the National Academy of
Sciences, 102*(43), 15545-15550. doi:10.1073/pnas.0506580102
Suzuki, H., Kanai-Azuma, M., & Kanai, Y. (2015). From Sex Determination to Initial
Folliculogenesis in Mammalian Ovaries: Morphogenetic Waves along the Anteroposterior
and Dorsoventral Axes. *Sex Dev, 9*(4), 190-204. doi:10.1159/000440689
Tingen, C., Kim, A., & Woodruff, T. K. (2009). The primordial pool of follicles and nest breakdown
in mammalian ovaries. *Mol Hum Reprod, 15*(12), 795-803. doi:10.1093/molehr/gap073
Uda, M., Ottolenghi, C., Crisponi, L., Garcia, J. E., Deiana, M., Kimber, W., . . . Pilia, G. (2004).
Foxl2 disruption causes mouse ovarian failure by pervasive blockage of follicle development.
Hum Mol Genet, 13(11), 1171-1181. doi:10.1093/hmg/ddh124
Wang, J.-J., Ge, W., Zhai, Q.-Y., Liu, J.-C., Sun, X.-W., Liu, W.-X., . . . Shen, W. (2020). Single-cell
transcriptome landscape of ovarian cells during primordial follicle assembly in mice. *PLOS
Biology, 18*(12), e3001025. doi:10.1371/journal.pbio.3001025
Wang, Z., Niu, W., Wang, Y., Teng, Z., Wen, J., Xia, G., & Wang, C. (2015). Follistatin288
Regulates Germ Cell Cyst Breakdown and Primordial Follicle Assembly in the Mouse Ovary.
PLOS ONE, 10(6), e0129643. doi:10.1371/journal.pone.0129643
Wear, H. M., McPike, M. J., & Watanabe, K. H. (2016). From primordial germ cells to primordial
follicles: a review and visual representation of early ovarian development in mice. *J Ovarian
Res, 9*(1), 36. doi:10.1186/s13048-016-0246-7
Xu, B., Hua, J., Zhang, Y., Jiang, X., Zhang, H., Ma, T., . . . Shi, Q. (2011). Proliferating Cell Nuclear
Antigen (PCNA) Regulates Primordial Follicle Assembly by Promoting Apoptosis of Oocytes
in Fetal and Neonatal Mouse Ovaries. *PLOS ONE, 6*(1), e16046.
doi:10.1371/journal.pone.0016046
Yang, C.-R., Rajkovic, G., Daldello, E. M., Luong, X. G., Chen, J., & Conti, M. (2020). The RNA-
binding protein DAZL functions as repressor and activator of mRNA translation during
oocyte maturation. *Nature Communications, 11*(1), 1399. doi:10.1038/s41467-020-15209-9
Zhang, H., Risal, S., Gorre, N., Busayavalasa, K., Li, X., Shen, Y., . . . Liu, K. (2014). Somatic cells
initiate primordial follicle activation and govern the development of dormant oocytes in mice.
Curr Biol, 24(21), 2501-2508. doi:10.1016/j.cub.2014.09.023
Zheng, G. X. Y., Terry, J. M., Belgrader, P., Ryvkin, P., Bent, Z. W., Wilson, R., . . . Bielas, J. H.
(2017). Massively parallel digital transcriptional profiling of single cells. *Nature
Communications, 8*(1), 14049. doi:10.1038/ncomms14049
Zheng, W., Zhang, H., Gorre, N., Risal, S., Shen, Y., & Liu, K. (2014). Two classes of ovarian
primordial follicles exhibit distinct developmental dynamics and physiological functions.
Hum Mol Genet, 23(4), 920-928. doi:10.1093/hmg/ddt486
Zheng, W., Zhang, H., & Liu, K. (2014). The two classes of primordial follicles in the mouse ovary:
their development, physiological functions and implications for future research. *Mol Hum
Reprod, 20*(4), 286-292. doi:10.1093/molehr/gau007
Zhihan, T., Xinyi, M., Qingying, L., Rufei, G., Yan, Z., Xuemei, C., . . . Junlin, H. (2019). Autophagy
participates in cyst breakdown and primordial folliculogenesis by reducing reactive oxygen
species levels in perinatal mouse ovaries. *J Cell Physiol, 234*(5), 6125-6135.
doi:10.1002/jcp.27367

1013
1014
1015

 Supplemental figures

1017
1018 **Supplemental Figure 1. Identification of cell types within the mouse ovary**

1019 The UMAP cluster maps show the expression of genes used to mark granulosa cells, mesenchymal

1020 cells, germ cells, epithelial cells and blood and immune cells. Dots are coloured according to their level

1021 of gene expression, with high levels of expression shown in purple. Red dashed lines show the clusters

1022 classified as each cell type.

1023

Supplemental Figure 2. Cell contributions to each cluster

(A) Barchart showing the cell count in each cluster and the number of cells allocated to each

cluster from each timepoint (E18.5, PND4 and PND7). Cells from the E18.5 timepoint are

shown in yellow, cells from the PND4 timepoint are shown in pink and cells from the PND7

timepoint are shown in blue. (B) Barchart showing the percentage of cells in each cluster that

arise from each timepoint (E18.5, PND4 and PND7). Cells from the E18.5 timepoint are shown

in yellow, cells from the PND4 timepoint are shown in pink and cells from the PND7 timepoint

are shown in blue.

1032

1033
1034
Supplemental Figure 3. Expression of key genes upregulated during primordial follicle activation

The UMAP cluster maps show the expression of differentially expressed genes in Gc_3. UMAP cluster

maps of established pregranulosa marker genes (*Amh*, *Inha* and *Nr5a2*) identified in our differential

gene expression analysis and DGE comparisons that are upregulated during primordial follicle

activation. UMAP cluster maps are shown for each timepoint individually and only include cells from

the timepoints in each panel (E18.5 = yellow, PND4 = pink and PND7 = blue). Dots are coloured

according to their level of gene expression, with high levels of expression shown in purple.

1042
1043

Supplemental Tables

Supplemental Table 1. Single cell RNA-seq quality control summary

Stage	Number of Cells	UMI/cell	Genes/cell	Number of reads	Fraction reads in cells	Reads mapped to genome
E18.5	8,210	12,011	3,698	516,012,231	90.3%	90.2%
PND4	7,587	10,267	3,413	468,968,268	91.4%	92.1%
PND7	9,013	9,850	2,982	435,022,244	88.8%	92.1%

Supplementary Table 2: Gene list of differentially expressed transcripts in granulosa cell clusters compared to all other cell clusters from differential gene expression analysis, presented by highest log$_2$FC.

Supplementary Table 3: Gene list of differentially expressed transcripts Gc_3 vs. Gc_2 (log$_2$ fold change ± 2; p ≤ 0.05), presented by highest average differential gene expression. Positive values are upregulated in Gc_3 cells (cells coloured in pink) and negative values are upregulated in Gc_2 cells (cells coloured in orange).

Supplementary Table 4: Gene list of differentially expressed transcripts Gc_3 vs. Gc_4 (log$_2$ fold change ± 2; p ≤ 0.05), presented by highest average differential gene expression. Positive values are upregulated in Gc_3 cells (cells coloured in pink) and negative values are upregulated in Gc_4 cells (cells coloured in maroon).

Supplementary Table 5: Gene list of differentially expressed transcripts between _Cdkn1b_$^{-/-}$ and WT ovaries at E18.5 (p ≤ 0.05), presented by highest log$_2$ fold change. Positive values are upregulated in _Cdkn1b_$^{-/-}$ ovaries (cells coloured in blue) and negative values are upregulated in WT ovaries (cells coloured in green).

Supplementary Table 6: Overlap of differentially expressed genes upregulated in Gc_3 vs Gc_2 and genes upregulated in p27KO E18.5, presented by highest differential gene expression in Gc_3 in the single-cell RNA sequencing dataset. Data taken from Supplementary Tables 3 and 5.

Supplementary Table 7: Overlap of differentially expressed genes upregulated in Gc_4 vs Gc_3 and genes upregulated in p27KO E18.5, presented by highest differential gene expression in Gc_4 in the single-cell RNA sequencing dataset. Data taken from Supplementary Tables 4 and 5.