Complete functional analysis of type IV pilus components of a reemergent plant pathogen
reveals neofunctionalization of paralog genes

Marcus V, Merfa[a], Xinyu Zhu[b], Deepak Shantharaj[a], Laura M. Gomez[a], Eber Naranjo[a], Neha Potnis[a], Paul
A. Cobine[b], Leonardo De La Fuente[a]*

[a]Department of Entomology and Plant Pathology, Auburn University, Auburn, AL 36849, USA.

[b]Department of Biological Sciences, Auburn University, Auburn, AL 36849, USA.

*Corresponding author: Leonardo De La Fuente. Mailing address: Department of Entomology and Plant
Pathology, 209 Rouse Life Sciences Bldg., Auburn University, Auburn, AL 36849. Phone: (334) 844-
2582.

Email: lzd0005@auburn.edu

Abstract

Type IV pilus (TFP) is a multifunctional bacterial structure involved in twitching motility, adhesion, biofilm formation, as well as natural competence. Here, by mutagenesis and functional analysis, we dissected the roles of all genes required for TFP biosynthesis and regulation in the reemergent plant pathogenic fastidious prokaryote *Xylella fastidiosa*. This xylem-limited, insect-transmitted pathogen lives constantly under flow conditions and therefore is highly dependent on TFP for host colonization. In addition, TFP-mediated natural transformation is a process that impacts genomic diversity and environmental fitness. Ten out of the thirty-eight genes analyzed were essential for movement and natural competence. Interestingly, seven sets of paralogs exist, and mutations showed opposing phenotypes, indicating evolutionary neofunctionalization of subunits within TFP. The minor pilin FimT3 was the only protein exclusively required for natural competence. We determined that FimT3 (but not the other two FimT paralogs) is a DNA receptor that is conserved among *X. fastidiosa* strains and binds DNA non-specifically via an electropositive surface. Among plant pathogens, this gene was also found in the genome of strains of the plant associated Xanthomonadaceae family. Overall, we highlight here the complex regulation of TFP in *X. fastidiosa*, providing a blueprint to understand TFP in other bacteria living under flow conditions.

Main

Type IV pili (TFP) are retractable hair-like proteinaceous cell appendages located at one or both poles of a bacterial cell [1]. In addition to playing fundamental roles in twitching motility and natural transformation, they also influence virulence, adhesion, and biofilm formation in many bacterial species [2-4]. During twitching motility, TFP extend, attach to surfaces, then retract pulling cells towards the point of attachment [1,2]. TFP also plays a central part in the DNA-uptake machinery that enables natural transformation [3,5]. Natural transformation is a horizontal gene transfer mechanism that allows bacterial cells to take up free DNA from the environment and incorporate it into their own genomes via homologous recombination (HR) [3,6]. The current model of natural transformation in bacteria involves binding of double-stranded DNA to the tip of TFP, which upon retraction brings this molecule to the outer-membrane surface of the cell [4]. DNA is then transported to the periplasm in a proposed Brownian ratchet mechanism [4,7]. Subsequently, DNA is translocated to the cytosol as a single strand, where recombination occurs if the

incoming DNA shares homology with the genome of the recipient cell [3]. Alternatively, the incoming DNA is

either metabolized or used as template to repair damaged DNA [8].

Natural transformation enables rapid evolution by generating genetic diversity and has been

involved in spreading of antibiotic resistance, adaptation to new environments and emergence of

pathogens [6,8-10]. In addition to animal pathogens [6,9], natural competence has been detected in two plant

pathogens, *Ralstonia solanacearum* [11] and *Xylella fastidiosa* [12], both of which possess a very broad plant

host range and are xylem-colonizers[13]. *X. fastidiosa* is a gram-negative fastidious prokaryote that infects

and causes reemergent diseases in many economically important crops worldwide, such as grapevine,

citrus, almond, peach, blueberry, and olive [14-16]. This pathogen lives constantly under flow conditions

inside the xylem of infected plants and the foregut of xylem-feeding insect vectors [17]. The main

mechanisms of virulence for *X. fastidiosa* are linked to its ability to move systemically within the xylem by

TFP-mediated twitching motility, and formation of biofilm that disrupts sap flow [18].

X. fastidiosa has been classified into subspecies [19-21], and genome analyses revealed widespread

HR events within and among subspecies [21-30]. Moreover, experimental evidence showed that *X. fastidiosa*

strains are able to recombine in vitro during coculture of either live or live and dead strains [31]. This

process occurs under multiple experimental settings including batch and flow culture conditions with

synthetic media or grapevine xylem sap from tolerant and susceptible varieties [12,32,33]. Finally, many of the

genes that are undergoing recombination among strains in vitro or in nature, are annotated as having

roles in virulence and bacterial fitness[26]. These observations led to the hypothesis that HR, a driver of

genetic diversity, is responsible for environmental adaptation [26] and/or host switching/expansion. In fact,

intersubspecific HR has been suggested to have a role in plant host expansion by generating strains that

infect mulberry [20], citrus, coffee [22], blueberry and blackberry [29].

The assembly, function, and regulation of TFP require multiple molecular components. Minor

pilins, which are present in much lower quantities than the major pilin, prime the assembly of TFP [34], and

some species-specific minor pilins may promote additional functions of TFP, such as aggregation and

adherence [35,36]. The functional diversity in pilins is mediated by differences in the C-terminal region of

these proteins, since all pilins (including major and minor pilins) share a highly conserved N-terminal

region [36]. TFP is key to *X. fastidiosa* in both natural competence, which promotes HR and thus generates

large genetic diversity [26,27,31], and twitching motility[18], which is its only mean of movement and

fundamental for a xylem-limited bacterium living under flow conditions. Therefore, in this study we

individually assessed the functional role of a comprehensive set of 38 TFP-related genes by site-directed

mutagenesis. We identified ten core genes that were essential for both natural competence and

movement of this plant pathogen. However, some components, mostly minor pilins, were only essential to

one feature or the other, and interestingly most paralogous genes had opposite functions, suggesting

neofunctionalization. Additionally, our study identified the FimT3 minor pilin as the DNA receptor of the *X.*

fastidiosa pilus. This minor pilin is found among plant pathogens mainly in bacterial species of the

Xanthomonadaceae family. Together, our data provide insight into how TFP paralogs evolve to perform

specific functions within bacterial cells, and how different bacterial species use unique minor pilins to

acquire exogenous DNA through natural competence.

Results

101   **Deletion of TFP-related genes differentially affects natural competence and movement of**

X. fastidiosa. Individual deletion mutant strains were generated by site-directed mutagenesis via natural

transformation to determine the functional roles of 38 TFP-related genes (Table S1) [37]. Genes were

chosen according to genome annotation and searches based on homology to counterparts encoded by

Pseudomonas aeruginosa [38,39]. To the best of our knowledge, these genes represent the entire set of

TFP genes encoded by *X. fastidiosa* strain TemeculaL. Assessment of natural competence was

conducted using the pAX1-Cm plasmid, which recombines into the neutral site 1 of the *X. fastidiosa*

genome and inserts a chloramphenicol (Cm) resistance cassette [31,40], whereas twitching motility was

determined by measuring colony fringe widths on solid agar plates [41]. Although the knockout of genes

studied here did not always modulate natural competence and twitching motility in the same manner, the

correlation between these two traits among mutant strains was positive (R^2 = 0.56, *P*<0.001; Table S3),

indicating that they were usually increased or reduced in a similar manner upon deletion of TFP genes.

113   A set of ten genes were considered core TFP components as deletion of such genes abrogated

both natural competence and movement of *X. fastidiosa* (Fig. 1 and Fig. S1-3). They include the ATPases

pilB and *pilT*, the prepilin peptidase *pilD*, the TFP assembly platform *pilC*, the whole alignment

subcomplex *pilMNOP*, the secretin *pilQ* and the regulatory gene *pilZ*. Deletion of the sensor gene *pilS* of

the PilRS two-component regulatory system (Fig. 1 and Fig. S1-3) significantly increased both natural

competence and twitching motility, while deletion of the regulator *pilR* was essential for twitching motility

but not for transformation. The minor pilin *fimT3* was the only gene found to be essential exclusively for

natural competence (Fig. 1 and Fig. S1). Curiously, this Δ*fimt3* strain showed significantly increased

twitching motility than wild-type (WT) cells (Fig. 1 and Fig. S2-3).

Seven sets of paralogs exist in the *X. fastidiosa* TFP cluster (PilA1-3, PilE/V/W/X1-2, PilY1-3 and

FimT1-3). Intriguingly, the knockout of paralogs presented varied phenotypes suggesting duplication and

neofunctionalization of the genes. The minor pilins *pilE1V1W1X1*, which are encoded in the same operon

(Table S2), were essential for twitching motility but not for transformation (Fig. 1 and Fig. S2-3), although

these mutants did show significantly lower recombination frequencies than WT (Fig. 1 and Fig. S1). In

comparison to the WT, Δ*pilE2* presented significant higher twitching motility and no changes in natural

transformation, Δ*pilV2* had significant higher natural transformation with no altered movement, Δ*pilW2*

showed significant higher natural transformation and significant lower movement, and Δ*pilX2* possessed

both significant lower natural transformation and twitching motility (Fig. 1 and Fig. S1-3). Knockout of

pilA1 significantly increased both natural competence and twitching motility, whereas deletion of *pilA2* and

pilA3 significantly reduced twitching motility and did not alter natural transformation (Fig. 1 and Fig. S1-3).

A strain with deletion of both *pilA1* and *pilA2* abolished twitching motility (Fig. 1 and Fig. S2-3), as

described elsewhere [37]. Additionally, deletion of the minor pilin *fimT1* significantly decreased both natural

competence and movement, while knockout of *fimT2* significantly increased natural competence and did

not change twitching motility, and Δ*fimT3* lost natural competence and presented significantly higher

movement than the WT (Fig. 1 and Fig. S1-3). As for *pilY1-1*, *pilY1-2* and *pilY1-3*, which are tip adhesins,

their deletion significantly reduced twitching motility. However, Δ*pilY1-1* and Δ*pilY1-2* presented

significant lower natural competence whilst Δ*pilY1-3* had significant higher natural competence than the

WT (Fig. 1 and Fig. S1-3).

Lastly, deletion of the pilotin *pilF* and of the retraction ATPase *pilU* significantly reduced natural

competence and movement. Knockout of the chemotaxis genes *pilGHIJL* and *chpBC* significantly

reduced twitching. However, Δ*pilG*, Δ*pilH* and Δ*pilL* showed increased natural competence, while Δ*pilJ*

and ∆*chpC* had lower natural competence, and ∆*pilI* and ∆*chpB* did not change this phenotype in

comparison to the WT (Fig. 1 and Fig. S1-3).

147   **TFP-related genes differentially modulate other virulence traits of *X. fastidiosa*.** We

evaluated phenotypes such as growth, biofilm formation, cell aggregation (measured by settling rate), and

virulence in planta of the individual TFP mutants. Growth rates were measured by change in OD_{600nm} over

time (Fig. S4-S5). Since this growth analysis considered only turbidity that may be affected by cell

attachment (biofilm) to surfaces and cell-to-cell aggregation, the number of viable *X. fastidiosa* cells

(measured as CFU/ml) was also evaluated. Only the non-recombinant and non-motile ∆*pilC* and ∆*pilP*

mutants showed significant changes in the number of cells in comparison to WT, with both having

significantly higher populations (Fig. S6). Biofilm formation is a critical aspect of virulence in *X. fastidiosa*.

Individual deletion of *pilB*, *pilD*, *pilH*, *pilI*, *pilM*, *pilN*, *pilO*, *pilQ*, *pilR*, *pilT*, *pilX1* and *fimT1* significantly

increased biofilm formation, whereas ∆*pilA3* and ∆*fimT3* formed significant less biofilm than WT (Fig. S7).

Planktonic growth was significantly increased for strains with individual deletion of *pilA1*, *pilC*, *pilE1*, *pilF*,

pilM, *pilN*, *pilO*, *pilP*, *pilQ*, *pilR*, *pilV1*, *pilW1*, *pilW2*, *pilX1*, *pilA1pilA2*, *pilY1-1*, *fimT3* and *chpC*, while

individual knockout of *pilD*, *pilT* and *pilY1-3* significantly decreased planktonic growth (Fig. S8). Moreover,

only ∆*pilD*, ∆*pilT* and ∆*pilX2* showed significant higher settling rates, with other mutants not differing

significantly from the WT (Fig. S9). Biofilm formation and settling rate, as well as planktonic growth and

growth rate were positively correlated (Table S3). Meanwhile, biofilm formation and twitching

motility/growth rate, settling rate and planktonic growth/growth rate, and growth rate and total viable *X.*

fastidiosa had significant negative correlations (Table S3). No other phenotype presented a significant

correlation to recombination frequency besides twitching motility.

166   Virulence of a subset of the *X. fastidiosa* mutant strains was assessed by inoculating *Nicotiana*

tabacum L. cv. Petite SR1 plants and scoring disease incidence and severity. Mutants chosen for

analysis had different twitching motility phenotypes, such as higher movement (∆*pilA1* and ∆*pilS*), lower

movement (∆*pilA2*) and non-motility (∆*pilA1pilA2*, ∆*pilQ* and ∆*pilR*). Additionally, mutants in the DNA-

binding *fimT3* (see below) and the paralogs *fimT1* and *fimT2* were tested and no significant change in

virulence was found compared to WT (data not shown). WT-inoculated plants presented the highest

disease severity over the time course of assays (Fig. S10a), with all plants inoculated with the mutant

strains having significantly lower AUDPC (Area Under the Disease Progress Curve) values, except plants

inoculated with Δ*pilA2* (Fig. S10b). All inoculated plants reached 100% disease incidence at the end of

assays, except plants inoculated with Δ*pilA1*, which had 89% disease incidence (±11%, standard error).

However, only WT-inoculated plants consistently presented severe symptoms of leaf scorch, while plants

inoculated with the mutant strains mostly presented mild symptoms (Fig. S10d). In addition, when

evaluating the population of *X. fastidiosa* in planta by qPCR, only the Δ*pilA1pilA2* double mutant showed

a defect in colonizing the top leaf of the plant host, and the Δ*pilQ* and Δ*pilR* mutants presented significant

higher population in the basal leaf of plant hosts (Fig. S10c). Together, results demonstrate that the

deletion of analyzed genes significantly reduced symptoms development in infected tobacco plants

(except for deletion of *pilA2*), which was not explained by bacterial distribution, since most strains could

effectively colonize the entirety of plants (except Δ*pilA1pilA2*).

The minor pilin FimT3 is the DNA receptor of the *X. fastidiosa* type IV pilus. The observation

of the minor pilin FimT3 being the only TFP component that is essential exclusively for natural

transformation (Fig.1 and Fig. S1-3) led us to hypothesize that this component is the DNA receptor of the

TFP of *X. fastidiosa*. We first analyzed the piliation phenotype of Δ*fimT3* in comparison to WT by

transmission electron microscopy (TEM). Both presented long TFP as well as short type I pili, with no

apparent changes in the piliation of *X. fastidiosa* cells upon deletion of *fimT3* (Fig. 2a). Next, the ability of

these cells to take up DNA from the extracellular environment was evaluated by exposing them to Cy-3-

labeled pAX1-Cm plasmid and observing under a fluorescent microscope. After exposure to the

fluorescently labeled DNA fragment, WT cells imported DNA allowing for protection from exogenous

DNaseI, but no fluorescent DNA foci were present in Δ*fimT3* cells (Fig. 2b). DNA foci were mostly

observed at one pole of the cells, which coincides to the described phenotype of *X. fastidiosa* that forms

TFP at only one of its poles [42]. The DNA-uptake ability of the mutant strains of other minor pilins that

decreased the natural transformation ability of *X. fastidiosa* upon deletion was also evaluated for

comparison. These included Δ*pilE1*, Δ*pilV1*, Δ*pilW1*, Δ*pilX1*, Δ*pilX2*, Δ*fimT1* and Δ*fimT2* (Fig.1 and Fig.

S1). The Δ*fimT2* mutant strain was included because this gene is a paralog of *fimT3*, even though its

deletion increased the natural competence of *X. fastidiosa* (Fig.1 and Fig. S1). All analyzed mutant strains

maintained the DNA-uptake (competence) phenotype (Fig. S11), but at different levels (Fig. S12). WT

cells presented the highest percentage of cells with DNA foci, with Δ*pilE1* having the lowest proportion of

cells with DNA acquisition (Fig. S12).

To determine the DNA-binding ability of FimT3, and compare to its paralogs FimT1 and FimT2,

the soluble portion of these proteins (named FimT3s, FimT1s and FimT2s, respectively), which lacked the

conserved N-terminal hydrophobic α-helices of pilins [36], were amplified and cloned for expression. These

proteins were then purified and assayed for DNA binding by agarose electrophoretic mobility shift assay

(EMSA). Results showed FimT3s was the only protein capable of binding DNA, since no shift was

observed with FimT1s and FimT2s when using as much as 10 µM of each protein (Fig. 2c). Moreover,

FimT3s was able to bind to different DNA sequences, independent of homology to the *X. fastidiosa*

TemeculaL genome, indicating that binding is not sequence-specific (Fig. S13). Collectively, *fimT3* being

the only gene essential exclusively for natural competence, the ability of FimT3 to bind to DNA and the

loss of DNA acquisition from the extracellular environment by cells upon its deletion, demonstrate that this

minor pilin is the DNA receptor of the *X. fastidiosa* TFP.

To further characterize FimT3, we searched for specific amino acid residues involved in DNA-

binding. The binding of DNA to TFP via a minor pilin was demonstrated in ComP of *Neisseria,* ComZ of

Thermus thermophilus, and VC0858 of *Vibrio cholerae* [4,43,44]. Amino acids alignments revealed that

FimT3 and ComP share only 28% identity (64% similarity) over 20% of query sequence, while FimT3 and

ComZ have no significant identity. Alignment of FimT3 with VC0858 of *Vibrio cholerae* showed 35%

identity and 62% similarity over 18% of query sequence. In spite of the low overall homology the

sequence alignments showed that an arginine residue at position 162 of FimT3 aligned with a lysine of

ComP (K108) that was previously demonstrated to be essential for its DNA-binding ability [43] (Fig. S14a).

Additionally, an arginine residue at position 160 of FimT3 aligned with an arginine of VC0858 (R168) that

was also important for the DNA-binding activity of the *V. cholerae* pilus [4] (Fig. S14b). These two arginine

residues of FimT3 (R160 and R162) were predicted in silico to have the highest probabilities of binding

DNA among the residues of this protein (Table S4). Moreover, alignment of FimT3 with a FimT ortholog

that is essential for natural competence of *Acinetobacter baylyi* [45] (28% identity and 46% similarity over

91% of query sequence) showed alignment of R162 of FimT3 to an arginine of FimT from *A. baylyi*.

Likewise, FimT3 shares 27% identity (48% similarity) over 80% of query sequence with a recently

described DNA-binding FimT ortholog of *Legionella pneumophila*, in which the aligned R160 and R162

have been demonstrated as important to bind DNA [46] (Fig. S14c).

Analysis of a structural model produced by the Phyre2 algorithm, showed that R160 and R162 of

FimT3 are located within an electropositive surface and could form a pocket to bind DNA (Fig. 2d). Thus,

to assess the role of these arginine residues in the DNA-binding ability of FimT3, we generated a FimT3s

variant by site-directed mutagenesis in which both arginine residues were replaced by alanine residues

(FimT3s-R160AR162A). Agarose EMSAs revealed that WT-FimT3s had a higher affinity for DNA than

FimT3s-R160AR162A based on the amount of protein needed to shift 50% of the input DNA (Fig. 2e).

Similar results were obtained when using the more sensitive acrylamide EMSA and the kanamycin (Km)

resistance cassette sequence as target DNA (previously determined to be bound by FimT3s in Fig. S13).

An abrupt shift was detected when using concentrations of FimT3s ranging between 1 and 5 µM, whereas

a much more subtle and gradual shift was observed for samples treated with FimT3s-R160AR162A within

the same concentration range (Fig. 2f). We then performed a densitometry analysis of the shifted bands

treated with concentrations ranging between 1 and 10 µM of each protein (Fig. 2g) and calculated the

area under the curve to quantify the DNA-binding affinity of FimT3s and FimT3s-R160AR162A. While the

DNA-binding affinity of FimT3s-R160AR162A was lower than that of FimT3s (Fig. 2h), the mutations did

not completely abolish DNA-binding ability suggesting other residues are contributing to this activity.

Nonetheless, these results show that R160 and R162 are important for DNA-binding activity of FimT3.

FimT3 is widely conserved within *X. fastidiosa* strains and, among plant pathogens, is

encoded by other members of the Xanthomonadaceae family. Previously, DNA-binding homologues

of FimT3 have been found in many representative species of γ-Proteobacteria, including the

Xanthomonadaceae family to which *X. fastidiosa* belongs [46]. We performed searches using the

Conditional Reciprocal Best BLAST (CRB-BLAST) [47], as well as phylogenetic analyses, to further explore

the distribution of the *fimT* paralogs within this family. All three paralogs of *fimT* are encoded by *X.*

fastidiosa, *X. taiwanensis*, many clade II *Xanthomonas* spp.[48] and few *Pseudoxanthomonas* spp. (Fig. 3).

Curiously, early branching clade I *Xanthomonas* spp. only encode *fimT1* and *fimT3* (Fig. 3). Specifically,

fimT3 was present in 1,416 out of 3,001 genomes of the Xanthomonadaceae (47%), majority of them

belonging to species within the genus *Xanthomonas*. In the *fimT3* clade, *X. fastidiosa* strains grouped

together, including *X. taiwanensis*, with a similar trend being observed for *Xanthomonas* spp. strains (Fig.

3). However, clade I and clade II *Xanthomonas* spp. and other Xanthomonadaceae species clustered

mostly separated (Fig. 3). The alignment of the different FimT3 sequences encoded by *X. fastidiosa*

strains showed a maximum of six nonconservative substitutions within strains belonging to the

subspecies *pauca*, which encode FimT3 that are 13 amino acids shorter than other strains (Fig. S15a).

Alignment of representative FimT3 sequences among strains belonging to the Xanthomonadaceae family

revealed that the arginine residues at positions 160 and 162 of FimT3 from *X. fastidiosa* strain TemeculaL

were conserved in most orthologous sequences of this protein (Fig. S15b). Additionally, R160 was always

preceded by a glycine residue, constituting a highly conserved GRxR motif (Fig. S15b), as seen

elsewhere [46]. Although amino acid identities ranged from 95.60% to 100% within *X. fastidiosa*, and from

15.5% to 80% compared to other species, only 41 out of 1,416 FimT3 sequences encoded by the

Xanthomonadaceae do not carry the GRxR motif (Table S5). Overall, results indicate that FimT3 is

encoded by representative plant pathogens of the Xanthomonadaceae and that R160 and R162 are

highly conserved within FimT3 sequences.

Discussion

In this study we defined the roles of individual subunits involved in regulation, assembly, and

functioning of the *X. fastidiosa* TFP on natural competence, twitching motility, adhesion, and biofilm (Fig.

4). To our knowledge, our study comprises the largest simultaneous functional description of a

comprehensive set of TFP-related genes. By performing site-directed mutagenesis of all TFP genes

followed by functional studies, we defined a set of ten core genes (*pilBCDMNOPQTZ*) that were essential

for natural competence and movement. Most of these proteins are inner- or outer-membrane associated,

confirming the important role of membrane anchorage for functionality of TFP.

Collectively, our data highlight the pervasive neofunctionalization among TFP components since

seven paralogous gene sets exerted mainly opposing regulatory functions on movement and natural

competence. For instance, paralogs of the minor pilins *pilEVWX* and *fimT* and the TFP tip adhesin *pilY1*

are mostly organized as operons in two different regions of the *X. fastidiosa* TemeculaL genome (Table

S2). Here, we observed that individual mutations in one set of these paralogs abrogated twitching motility

and greatly reduced natural competence, while deletions in the other set of paralogous genes produced

variable phenotypes that were mostly opposite to their counterparts. It has been demonstrated that minor

pilins prime the assembly of TFP [34]. This interaction is specific, in which major pilins require a specific

cognate subset of minor pilins for TFP assembly [49]. Therefore, the presence of paralogs for major and

minor pilins displaying opposite functions within *X. fastidiosa* possibly suggest assembly of multiple TFP

fibers with specific functions. However, this needs to be further assessed (see SI discussion).

Most remarkably among paralogs, is the discovery of the minor pilin FimT3 as the TFP DNA receptor.

This is an outstanding neofunctionalization example since FimT1 and FimT2 did not show DNA

adherence abilities. This finding was also supported by our phylogenetic analysis, which showed

divergent evolution of FimT3 from its paralogs. FimT3 plays an essential role in natural competence by

mediating extracellular DNA uptake that may be integrated into the genome, thereby conferring new

phenotypic traits that may allow an increase in bacterial fitness and/or host expansion of *X. fastidiosa*.

The interaction between DNA molecules and an extracellular specific receptor is a key first step in the

DNA uptake and thus natural competence of bacterial species. Initially, DNA binding by TFP has been

observed in *P. aeruginosa* [50], *Streptococcus pneumoniae* [51], and *V. cholerae* [4], in which pilus retraction

drives DNA uptake by transporting it to the cell surface, which is then imported to the periplasm by the

competence protein ComEA [4,7]. Although these early studies provide evidence that the specialized DNA

receptor resides within type IV competence pilus fibers, no direct binding of DNA by individual purified

major and/or minor pilins was reported. In *V. cholerae*, targeted mutations of positively charged amino

acid residues within the minor pilins VC0858 and VC0859 reduced the DNA-binding affinity of its TFP [4],

but the ability of these minor pilins to directly bind DNA was not assayed. Direct DNA binding has been

first observed for the pilin protein ComP from *N. meningitidis*, with binding occurring at a positively

charged surface of this protein [43]. ComP orthologs are found mostly within the Neisseriaceae family,

where they mediate DNA uptake by binding to specific DNA uptake sequences, as *Neisseria* spp.

preferentially take up homotypic DNA [43,52]. ComZ, encoded by *T. thermophilus*, was the second minor

pilin described as a TFP DNA receptor [44]. However, ComZ is an unusual minor pilin that possesses an

additional large β-solenoid domain inserted into the common β-sheet structure of pilins. Furthermore,

ComZ binds to another minor pilin, PilA2 (PilV in *X. fastidiosa*), and its DNA binding ability does not

appear to require positive charges in its surface [44]. Nonetheless, the binding of ComZ to another minor

pilin suggests that specific interactions within the pilus structure may occur to assemble the competence

pilus fiber.

Here, the deletion of individual paralogs of the major pilin PilA abrogated neither natural

competence nor twitching motility, suggesting functional overlapping among them (see SI Discussion). In

addition to the novel functions described above, our study confirmed the role of core components that

were previously analyzed in other bacterial species including *Acinetobacter baumannii* [53], *A. baylyi* [45], *T.*

thermophilus [54], *Haemophilus influenzae* [55], *P. aeruginosa* [56-59], *L. pneumophila*[46] and *X. fastidiosa* [32,42,60].

The exceptions were *pilP* and *pilZ* in which previous deletion of both greatly reduced natural competence

[61,62], but did not abolish it here. It is of note that very few of these genes have been simultaneously

assayed for their role in both natural transformation and movement mediated by TFP.

Deletion of FimT3 in *X. fastidiosa*, but not of the paralogs FimT1 and FimT2, exclusively disrupted

the natural competence of this bacterium. FimT3 binds to DNA via a surface in which positively charged

arginine residues contribute to the DNA binding ability of this protein. However, the binding appears to be

non-specific, which is in contrast to ComP [43]. FimT from the human pathogen *L. pneumophila* [46], which is

a homologue of the FimT3 studied here, was recently identified as a pilin with DNA-binding ability.

Another FimT homolog was also demonstrated to be specialized in natural competence in *A. baylyi* [45],

although no DNA binding was assessed. In our study, FimT3 seems to be particularly enriched in plant

pathogens of the Xanthomonadaceae family, being encoded by all *X. fastidiosa* strains as well as many

other bacterial species belonging to this family. Thus, our results, together with previous studies,

demonstrate that DNA binding by TFP occurs among bacterial species through a similar component such

as minor pilins, which themselves are not widely distributed or conserved, but are rather specific to

different bacterial groups. For instance, the ability of strain-specific subsets of minor pilins to modulate

TFP dynamics has already been described in *P. aeruginosa* (see SI Discussion) [49]. Understanding how

this minor pilin interacts with different minor and major pilins of *X. fastidiosa* in the competence pilus is an

important area to explore. It is not known whether the other Xanthomonadaceae bacterial species

encoding FimT3 that were found in our analysis are also naturally competent. For instance, *X. fastidiosa*

strains belonging to the *pauca* subspecies have a shorter FimT3 amino acid sequence, missing part of

the C-terminal, and have a few aminoacidic changes (although maintain the arginine residues studied

here). The only reported attempt to transform a *X. fastidiosa* strain belonging to the subspecies *pauca* by

natural competence in vitro has failed [63]. Studies with the *X. fastidiosa* subsp. *pauca* FimT3 variant should

investigate both individual DNA-binding activity and potential roles of the C-terminus in interaction with

the rest of the TFP structure.

In summary, our study highlights the fundamental functions of TFP in virulence and evolution

evidenced by complex regulatory features. Due to the worldwide threat that *X. fastidiosa* presents to

many economically important crops and to the ever-increasing list of plant hosts, it is imperative to

investigate how this plant pathogen acquires foreign DNA and expands its genetic diversity to help

researchers understand its eco-evolutionary history and re-emergence of this pathogen in many

geographic areas. In addition, our results provide a robust blueprint to assess TFP functions in bacteria

living under flow conditions. An extensive discussion including additional genes analyzed in this study and

their functional roles in natural transformation and twitching motility of *X. fastidiosa*, in addition to unique

behaviors and other phenotypes such as biofilm formation, virulence in planta, and diversity in paralogous

genes functions, is included in SI Discussion. All of this indicates that a complex and tightly regulated

machinery is involved in these bacterial functions.

Methods

Methods that are only cited here are described in detail in SI Materials and Methods.

Bacterial strains, plasmids, and culture conditions. All strains and plasmids used in this study

are listed in Table S6. All *X. fastidiosa* mutant strains used here are derivatives of the *X. fastidiosa* subsp.

fastidiosa strain TemeculaL [26], and were originated by site-directed mutagenesis of each gene of interest

using natural competence as described elsewhere [37]. Primers used in this process are listed in Table S7.

X. fastidiosa was recovered from -80 ºC glycerol stocks and routinely cultured for seven days at 28 ºC on

periwinkle wilt (PW) agar plates [64], modified by removing phenol red and using bovine serum albumin (1.8

g/l) (BSA; Gibco Life Sciences Technology), and sub-cultured onto fresh PW agar plates for another

seven days at 28 ºC before use. *X. fastidiosa* mutant strains were grown similarly but using PW plates

amended with Km. All assays were performed using the subcultured *X. fastidiosa* strains. Cells were

suspended and cultured in PD3 broth [64] to perform phenotypic assays, while phosphate-buffered saline

(PBS) was used to suspend cells in liquid for in planta assays. Whenever needed, the antibiotics Km and

Cm were used at concentrations of 50 and 10 µg/ml, respectively. When used together with Cm, the Km

concentration was reduced to 30 µg/ml. Luria-Bertani (LB) medium (BD Difco) was used to culture *E. coli*

cells. When needed, Km, Cm and ampicillin (Amp) were added to LB at concentrations of 50, 35 and 100

µg/ml, respectively.

Phenotypic analyses of *X. fastidiosa* WT and mutant strains. Analyses of natural competence

among strains [31], twitching motility [33], growth curve and growth rate [31], biofilm formation and planktonic

growth [31,65], settling rate [31], piliation observation by TEM [37], and virulence in planta using tobacco as

model system [66] were performed as described elsewhere. All experiments had at least three biological

replicates, unless otherwise stated.

DNA uptake assays. DNA uptake assays were performed as similarly described for the analysis

of natural competence among *X. fastidiosa* strains (see SI Materials and Methods). Briefly, recipient cells

were suspended in PD3 broth to $OD_{600nm} = 0.6$, spotted onto PD3 agar plates and grown for three days at

28 ºC. Then, 1 µg of fluorescently labeled pAX1-Cm plasmid (10-µl volume), labeled using the *Label* IT

Nucleic Acid Labeling kit, Cy3 (Mirus Bio LLC), was added on top of cells, air-dried, and incubated at 28

ºC for another 24 h. After, cells were harvested in 150 µl of PD3 broth, and a 50 µl aliquot was treated

with 10 units of DNase I (New England Biolabs) for 10 minutes at 37 ºC to degrade the remaining

extracellular DNA, and cells were observed using a 100× oil immersion objective in a Nikon Eclipse Ti

inverted microscope (Nikon). Image acquisition was performed using a Nikon DS-Q1 digital camera

(Nikon) controlled by the NIS-Elements software version 3.0 (Nikon), which was also used to create

merged fluorescent images. To detect Cy3, an excitation wavelength of 590 nm was used (tetramethyl

rhodamine isothiocyanate; TRITC filter). The proportion of cells that acquired extracellular DNA was

calculated as the percentage of cells with fluorescent DNA foci to cells without fluorescent DNA foci.

Experiments were performed at least three times independently.

Protein cloning, expression, and purification. Cloning, expression, and purification of FimT1s,

FimT2s and FimT3s were performed similarly as described elsewhere using the pHIS-Parallel1 plasmid

(Table S6) [67,68]. Amino acid exchanges in FimT3s were performed by designing a pair of primers (Table

S7) that exchanged the arginine residues at positions 160 and 162 by alanine residues. This pair of

primers was designed using the QuickChange Primer Design tool (Agilent Technologies, Inc.;

https://www.agilent.com/store/primerDesignProgram.jsp). For amino acid exchanges, the whole pHIS-

Parallel1-*fimT3s* construct was amplified via *Pfu* DNA polymerase (G-Biosciences) using a standard

protocol from the manufacturer and this pair of primers in a S1000 thermal cycler (Bio-Rad). Expression

and purification of FimT3s-R160AR162A were performed as described above (see SI Materials and

Methods) [67]. PCR products were purified using the Gel/PCR DNA Fragments Extraction kit (IBI Scientific).

Correct cloning and amino acid exchanges in all constructs were confirmed by Sanger sequencing

(Sequetech Corporation) using the T7 promoter forward primer (Table S7).

DNA binding assays. Agarose EMSAs were mainly performed to assess the DNA-binding ability

of purified FimT1s, FimT2s and FimT3s. Briefly, 200 ng of DNA (usually pAX1-Cm plasmid) were

incubated for 30 minutes at 28 ºC with increasing concentrations of purified proteins in 20 µL EMSA

reaction buffer (50 mM Tris-HCl, pH 7.5; 50 mM NaCl; 200 mM KCl; 5 mM $MgCl_2$; 5 mM EDTA, pH 8.0; 5

mM DTT; 0.25 mg/ml BSA). Lysis buffer containing 250 µM imidazole, in which proteins were suspended,

was included as blank control. After the incubation period, DNA was separated by electrophoresis on a

0.8% agarose gel containing GelRed nucleic acid gel stain (Biotium) in Tris-acetate-EDTA buffer (120 V

for 40 minutes). Experiments were performed at least three times independently. On the other hand,

native acrylamide EMSAs were used to perform titration of the DNA binding activity of FimT3s and

FimT3s-R160AR162A. In summary, 60 ng of Cy-3 labeled Km resistance cassette were incubated for 30

minutes at 28 ºC with increasing concentrations of purified proteins in 20 µL EMSA reaction buffer and

separated by electrophoresis on a 3.5% native acrylamide gel in Tris-borate-EDTA buffer (40 V for 4

hours). Native acrylamide gels were pre-run at 40 V for 30 minutes before being used. DNA samples

were directly visualized using the ImageQuant LAS 4010 Imaging System (GE Healthcare), since the

used DNA was fluorescently labeled with Cy3. Experiments were performed two times independently.

Densitometry analysis of the shifted bands of Cy-3 labeled DNA in native acrylamide gels was performed

by measuring the fluorescent intensity of these bands when treated with 1, 2, 5, and 10 µM of each

protein using ImageJ [69]. Then, these values were used to calculate the area under the fluorescent curves,

as performed to obtain the AUDPC (see SI Materials and Methods) [70], to quantify the DNA-binding affinity

of FimT3s and FimT3s-R160AR162A by determining the progress of shifted bands in relation to each

protein concentration.

Bioinformatic analyses. FimT3 modeling was performed using the Phyre2 web portal [71] and

visualized via PyMOL version 2.4.0 (Schrödinger, LLC). The surface electrostatics of FimT3 was

predicted using the APBS Electrostatics plugin from PyMOL. Sequence alignments of FimT3 with ComP,

VC0858 and FimT from *A. baylyi* and *L. pneumophila* were performed by retrieving the respective

sequences from NCBI, aligning through the T-Coffee Multiple Sequence Alignment Server [72] and

visualizing using the BoxShade webserver (https://embnet.vital-it.ch/software/BOX_form.html). For FimT3

analyses within the Xanthomonadaceae family, DNA sequences of different FimT homologs from *Xylella*,

Xanthomonas, *Lysobacter* and *Stenotrophomonas* were used as query for screening genomes of

Xanthomonadaceae using CRB-BLAST [47] with default settings. Nucleotide sequences for the CRB-

BLAST hits were then retrieved and aligned using MAFFT [73], followed by RAxML version 8.0.24 [74] to build

a phylogenetic tree, which was visualized using FigTree version 1.4.4 (http://tree.bio.ed.ac.uk/). FimT3-

encoding sequences were retrieved from the phylogenetic tree using the TREE2FASTA Perl script [75],

translated into amino acid sequences and aligned using Clustal Omega (Clustal 12.1) to determine

percentage of identical amino acids [76]. The visualization of the alignment of representative FimT3

sequences among *X. fastidiosa* strains and representative bacterial species belonging to the

Xanthomonadaceae family was performed using the BoxShade webserver. The FimT3 GRxR motif

sequence logo was generated using the WebLogo webserver[77].

Data analysis. Data from natural competence assays (recombination frequency) and the area

under the fluorescent curve were compared by two-tailed Student's *t*-test. Data from twitching motility,

growth rate, viable *X. fastidiosa* CFU/ml obtained during natural competence assays, biofilm formation,

planktonic growth, settling rate, AUDPC, *X. fastidiosa* population in planta and percentage of cells

acquiring DNA from the extracellular environment were individually analyzed by one-way analysis of

variance (ANOVA) followed by Tukey's HSD multiple comparisons of means in R 4.0.0 under the package

multcomp [78]. Correlation among analyzed phenotypes was determined by Pearson's correlation using the

SigmaPlot software version 11.0 (Systat Software Inc.).

References
464
1   Mattick, J. S. Type IV pili and twitching motility. *Annu Rev Microbiol* **56**, 289-314,
doi:10.1146/annurev.micro.56.012302.160938 (2002).
2   Jarrell, K. F. & McBride, M. J. The surprisingly diverse ways that prokaryotes move. *Nat Rev*
Microbiol **6**, 466-476 (2008).
3   Dubnau, D. & Blokesch, M. Mechanisms of DNA uptake by naturally competent bacteria. *Annu Rev*
Genet **53**, 217-237, doi:10.1146/annurev-genet-112618-043641 (2019).
4   Ellison, C. K. *et al.* Retraction of DNA-bound type IV competence pili initiates DNA uptake during
natural transformation in *Vibrio cholerae*. *Nat Microbiol* **3**, 773-780, doi:10.1038/s41564-018-0174-
y (2018).
5   Seitz, P. & Blokesch, M. DNA-uptake machinery of naturally competent *Vibrio cholerae*. *Proc Natl*
Acad Sci U S A **110**, 17987-17992, doi:10.1073/pnas.1315647110 (2013).
6   Johnston, C., Martin, B., Fichant, G., Polard, P. & Claverys, J. P. Bacterial transformation:
distribution, shared mechanisms and divergent control. *Nat Rev Microbiol* **12**, 181-196 (2014).
7   Seitz, P. *et al.* ComEA is essential for the transfer of external DNA into the periplasm in naturally
transformable *Vibrio cholerae* cells. *PLoS Genet* **10**, e1004066, doi:10.1371/journal.pgen.1004066
(2014).
8   Seitz, P. & Blokesch, M. Cues and regulatory pathways involved in natural competence and
transformation in pathogenic and environmental Gram-negative bacteria. *FEMS Microbiol Rev* **37**,
336-363, doi:10.1111/j.1574-6976.2012.00353.x (2013).
9   Johnsborg, O., Eldholm, V. & Håvarstein, L. S. Natural genetic transformation: prevalence,
mechanisms and function. *Res Microbiol* **158**, 767-778, doi:10.1016/j.resmic.2007.09.004 (2007).
10   Wiedenbeck, J. & Cohan, F. M. Origins of bacterial diversity through horizontal genetic transfer and
adaptation to new ecological niches. *FEMS Microbiol Rev* **35**, 957-976, doi:10.1111/j.1574-
6976.2011.00292.x (2011).
11   Bertolla, F., Frostegård, Å., Brito, B., Nesme, X. & Simonet, P. During infection of its host, the plant
pathogen *Ralstonia solanacearum* naturally develops a state of competence and exchanges
genetic material. *Mol Plant Microbe Interact* **12**, 467-472 (1999).
12   Kung, S. H. & Almeida, R. P. P. Natural competence and recombination in the plant pathogen
Xylella fastidiosa. *Appl Environ Microbiol* **77**, 5278-5284, doi:10.1128/AEM.00730-11 (2011).

13   De La Fuente, L., Merfa, M. V., Cobine, P. A. & Coleman, J. J. Pathogen adaptation to the xylem
environment. *Annu Rev Phytopathol* **60**, 8.1-8.24, doi:10.1146/annurev-phyto-021021-041716
(2022).
14   Hopkins, D. L. & Purcell, A. H. *Xylella fastidiosa*: Cause of Pierce's Disease of grapevine and other
emergent diseases. *Plant Dis* **86**, 1056-1066 (2002).
15   Saponari, M., Boscia, D., Nigro, F. & Martelli, G. P. Identification of DNA sequences related to
Xylella fastidiosa in oleander, almond and olive trees exhibiting leaf scorch symptoms in Apulia
(Southern Italy). *J. Plant Pathol.* **95**, 668 (2013).
16   Chang, C. J., Donaldson, R., Brannen, P., Krewer, G. & Boland, R. Bacterial leaf scorch, a new
blueberry disease caused by *Xylella fastidiosa*. *HortScience* **44**, 413-417 (2009).
17   Almeida, R. P. P., Coletta-Filho, H. D. & Lopes, J. R. S. in *Manual of Security Sensitive Microbes*
and Toxins (ed D. Liu) 841-850 (CRC Press, 2014).
18   Chatterjee, S., Almeida, R. P. P. & Lindow, S. Living in two worlds: the plant and insect lifestyles of
Xylella   fastidiosa.   *Annu   Rev   Phytopathol*   **46**,   243-271,
doi:10.1146/annurev.phyto.45.062806.094342 (2008).
19   Schaad, N. W., Postnikova, E., Lacy, G., Fatmi, M. & Chang, C. J. *Xylella fastidiosa* subspecies:
X. fastidiosa subsp. [correction] *fastidiosa* [correction] subsp. nov., *X. fastidiosa* subsp. *multiplex*
subsp. nov., and *X. fastidiosa* subsp. *pauca* subsp. nov. *Syst Appl Microbiol* **27**, 290-300,
doi:10.1078/0723-2020-00263 (2004).
20   Nunney, L., Schuenzel, E. L., Scally, M., Bromley, R. E. & Stouthamer, R. Large-scale
intersubspecific recombination in the plant-pathogenic bacterium *Xylella fastidiosa* is associated
with the host shift to mulberry. *Appl Environ Microbiol* **80**, 3025-3033, doi:10.1128/AEM.04112-13
(2014).
21   Scally, M., Schuenzel, E. L., Stouthamer, R. & Nunney, L. Multilocus sequence type system for the
plant pathogen *Xylella fastidiosa* and relative contributions of recombination and point mutation to
clonal diversity. *Appl Environ Microbiol* **71**, 8491-8499, doi:10.1128/AEM.71.12.8491-8499.2005
(2005).
22   Nunney, L., Yuan, X., Bromley, R. E. & Stouthamer, R. Detecting genetic introgression: high levels
of intersubspecific recombination found in *Xylella fastidiosa* in Brazil. *Appl Environ Microbiol* **78**,
4702-4714, doi:10.1128/AEM.01126-12 (2012).
23   Almeida, R. P. P. *et al.* Genetic structure and biology of *Xylella fastidiosa* strains causing disease
in citrus and coffee in Brazil. *Appl Environ Microbiol* **74**, 3690-3701, doi:10.1128/AEM.02388-07
(2008).
24   Coletta-Filho, H. D., Francisco, C. S., Lopes, J. R. S., Muller, C. & Almeida, R. P. P. Homologous
recombination and *Xylella fastidiosa* host-pathogen associations in South America. *Phytopathology*
107, 305-312, doi:10.1094/PHYTO-09-16-0321-R (2017).
25   Nunney, L. *et al.* Population genomic analysis of a bacterial plant pathogen: novel insight into the
origin   of   Pierce's   disease   of   grapevine   in   the   U.S.   *PLoS   One*   **5**,   e15488,
doi:10.1371/journal.pone.0015488 (2010).
26   Potnis, N. *et al.* Patterns of inter- and intrasubspecific homologous recombination inform eco-
evolutionary dynamics of *Xylella fastidiosa*. *ISME J* **13**, 2319-2333, doi:10.1038/s41396-019-0423-
y (2019).
27   Vanhove, M. *et al.* Genomic diversity and recombination among *Xylella fastidiosa* subspecies. *Appl*
Environ Microbiol **85**, e02972-02918, doi:10.1128/AEM (2019).
28   Castillo, A. I., Chacón-Díaz, C., Rodríguez-Murillo, N., Coletta-Filho, H. D. & Almeida, R. P. P.
Impacts of local population history and ecology on the evolution of a globally dispersed pathogen.
BMC Genomics **21**, 369, doi:10.1186/s12864-020-06778-6 (2020).
29   Nunney, L., Hopkins, D. L., Morano, L. D., Russell, S. E. & Stouthamer, R. Intersubspecific
recombination in *Xylella fastidiosa* strains native to the United States: infection of novel hosts
associated   with   an   unsuccessful   invasion.   *Appl   Environ   Microbiol*   **80**,   1159-1169,
doi:10.1128/AEM.02920-13 (2014).
30   Jacques, M. A. *et al.* New coffee plant-infecting *Xylella fastidiosa* variants derived via homologous
recombination. *Appl Environ Microbiol* **82**, 1556-1568, doi:10.1128/AEM.03299-15 (2016).
31   Kandel, P. P., Almeida, R. P. P., Cobine, P. A. & De La Fuente, L. Natural competence rates are
variable among *Xylella fastidiosa* strains and homologous recombination occurs in vitro between

549 550	subspecies *fastidiosa* and *multiplex*. *Mol Plant Microbe Interact* **30**, 589-600, doi:10.1094/MPMI-02-17-0053-R (2017).
551 552	32 Kung, S. H. & Almeida, R. P. P. Biological and genetic factors regulating natural competence in a bacterial plant pathogen. *Microbiology* **160**, 37-46, doi:10.1099/mic.0.070581-0 (2014).
553 554 555	33 Kandel, P. P., Lopez, S. M., Almeida, R. P. P. & De La Fuente, L. Natural competence of *Xylella fastidiosa* occurs at a high frequency inside microfluidic chambers mimicking the bacterium's natural habitats. *Appl Environ Microbiol* **82**, 5269-5277, doi:10.1128/AEM.01412-16 (2016).
556 557 558	34 Nguyen, Y. *et al. Pseudomonas aeruginosa* minor pilins prime type IVa pilus assembly and promote surface display of the PilY1 adhesin. *J Biol Chem* **290**, 601-611, doi:10.1074/jbc.M114.616904 (2015).
559 560 561	35 Jacobsen, T., Bardiaux, B., Francetic, O., Izadi-Pruneyre, N. & Nilges, M. Structure and function of minor pilins of type IV pili. *Med Microbiol Immunol* **209**, 301-308, doi:10.1007/s00430-019-00642-5 (2020).
562 563	36 Giltner, C. L., Nguyen, Y. & Burrows, L. L. Type IV pilin proteins: versatile molecular modules. *Microbiol Mol Biol Rev* **76**, 740-772, doi:10.1128/MMBR.00035-12 (2012).
564 565 566 567	37 Kandel, P. P., Chen, H. & De La Fuente, L. A short protocol for gene knockout and complementation in *Xylella fastidiosa* shows that one of the type IV pilin paralogs (PD1926) is needed for twitching while another (PD1924) affects pilus number and location. *Appl Environ Microbiol* **84**, e01167-01118 (2018).
568 569	38 Burrows, L. L. *Pseudomonas aeruginosa* twitching motility: type IV pili in action. *Annu Rev Microbiol* **66**, 493-520, doi:10.1146/annurev-micro-092611-150055 (2012).
570 571	39 Nolan, L. M. *et al. Pseudomonas aeruginosa* is capable of natural transformation in biofilms. *Microbiology* **166**, 995-1003, doi:10.1099/mic.0.000956 (2020).
572 573	40 Matsumoto, A., Young, G. M. & Igo, M. M. Chromosome-based genetic complementation system for *Xylella fastidiosa*. *Appl Environ Microbiol* **75**, 1679-1687, doi:10.1128/AEM.00024-09 (2009).
574 575 576	41 Galvani, C. D., Li, Y., Burr, T. J. & Hoch, H. C. Twitching motility among pathogenic *Xylella fastidiosa* isolates and the influence of bovine serum albumin on twitching-dependent colony fringe morphology. *FEMS Microbiol Lett* **268**, 202-208, doi:10.1111/j.1574-6968.2007.00601.x (2007).
577 578	42 Meng, Y. *et al.* Upstream migration of *Xylella fastidiosa* via pilus-driven twitching motility. *J Bacteriol* **187**, 5560-5567, doi:10.1128/JB.187.16.5560-5567.2005 (2005).
579 580	43 Cehovin, A. *et al.* Specific DNA recognition mediated by a type IV pilin. *Proc Natl Acad Sci U S A* **110**, 3065-3070, doi:10.1073/pnas.1218832110 (2013).
581 582	44 Salleh, M. Z. *et al.* Structure and properties of a natural competence-associated pilin suggest a unique pilus tip-associated DNA receptor. *mBio* **10**, e00614-00619, doi:10.1128/mBio (2019).
583 584 585	45 Leong, C. G. *et al.* The role of core and accessory type IV pilus genes in natural transformation and twitching motility in the bacterium *Acinetobacter baylyi*. *PLoS One* **12**, e0182139, doi:10.1371/journal.pone.0182139 (2017).
586 587	46 Braus, S. A. G. *et al.* The molecular basis of FimT-mediated DNA uptake during bacterial natural transformation. *Nature Communications* **13**, 1065, doi:10.1038/s41467-022-28690-1 (2022).
588 589 590 591	47 Aubry, S., Kelly, S., Kümpers, B. M., Smith-Unna, R. D. & Hibberd, J. M. Deep evolutionary comparison of gene expression identifies parallel recruitment of *trans*-factors in two independent origins of C4 photosynthesis. *PLoS Genet* **10**, e1004365, doi:10.1371/journal.pgen.1004365 (2014).
592 593 594	48 Jacques, M. A. *et al.* Using ecology, physiology, and genomics to understand host specificity in *Xanthomonas*. *Annu Rev Phytopathol* **54**, 163-187, doi:10.1146/annurev-phyto-080615-100147 (2016).
595 596 597	49 Giltner, C. L., Rana, N., Lunardo, M. N., Hussain, A. Q. & Burrows, L. L. Evolutionary and functional diversity of the *Pseudomonas* type IVa pilin island. *Environ Microbiol* **13**, 250-264, doi:10.1111/j.1462-2920.2010.02327.x (2011).
598 599	50 van Schaik, E. J. *et al.* DNA binding: a novel function of *Pseudomonas aeruginosa* type IV pili. *J Bacteriol* **187**, 1455-1464, doi:10.1128/JB.187.4.1455-1464.2005 (2005).
600 601	51 Laurenceau, R. *et al.* A type IV pilus mediates DNA binding during natural transformation in *Streptococcus pneumoniae*. *PLoS Pathog* **9**, e1003473, doi:10.1371/journal.ppat.1003473 (2013).
602 603	52 Berry, J. L., Cehovin, A., McDowell, M. A., Lea, S. M. & Pelicic, V. Functional analysis of the interdependence between DNA uptake sequence and its cognate ComP receptor during natural

604		transformation in *Neisseria* species. *PLoS Genet* **9**, e1004014, doi:10.1371/journal.pgen.1004014
605		(2013).
606	53	Vesel, N. & Blokesch, M. Pilus production in *Acinetobacter baumannii* is growth phase dependent
607		and essential for natural transformation. *J Bacteriol* **203**, e00034- 00021, doi:10.1128/JB.00034-21
608		(2021).
609	54	Rumszauer, J., Schwarzenlander, C. & Averhoff, B. Identification, subcellular localization and
610		functional interactions of PilMNOWQ and PilA4 involved in transformation competency and pilus
611		biogenesis in the thermophilic bacterium *Thermus thermophilus* HB27. *FEBS J* **273**, 3261-3272,
612		doi:10.1111/j.1742-4658.2006.05335.x (2006).
613	55	Carruthers, M. D. *et al.* Biological roles of nontypeable *Haemophilus influenzae* type IV pilus
614		proteins encoded by the *pil* and *com* operons. *J Bacteriol* **194**, 1927-1933, doi:10.1128/JB.06540-
615		11 (2012).
616	56	Tammam, S. *et al.* PilMNOPQ from the *Pseudomonas aeruginosa* type IV pilus system form a
617		transenvelope protein interaction network that interacts with PilA. *J Bacteriol* **195**, 2126-2135,
618		doi:10.1128/JB.00032-13 (2013).
619	57	Alm, R. A., Bodero, A. J., Free, P. D. & Mattick, J. S. Identification of a novel gene, *pilZ*, essential
620		for type 4 fimbrial biogenesis in *Pseudomonas aeruginosa*. *J Bacteriol* **178**, 46-53 (1996).
621	58	Marko, V. A., Kilmury, S. L. N., MacNeil, L. T. & Burrows, L. L. *Pseudomonas aeruginosa* type IV
622		minor pilins and PilY1 regulate virulence by modulating FimS-AlgR activity. *PLoS Pathog* **14**,
623		e1007074, doi:10.1371/journal.ppat.1007074 (2018).
624	59	Ayers, M. *et al.* PilM/N/O/P proteins form an inner membrane complex that affects the stability of
625		the *Pseudomonas aeruginosa* type IV pilus secretin. *J Mol Biol* **394**, 128-142,
626		doi:10.1016/j.jmb.2009.09.034 (2009).
627	60	Li, Y. *et al.* Type I and type IV pili of *Xylella fastidiosa* affect twitching motility, biofilm formation and
628		cell-cell aggregation. *Microbiology* **153**, 719-726, doi:10.1099/mic.0.2006/002311-0 (2007).
629	61	Drake, S. L., Sandstedt, S. A. & Koomey, M. PilP, a pilus biogenesis lipoprotein in *Neisseria*
630		*gonorrhoeae*, affects expression of PilQ as a high-molecular-mass multimer. *Mol Microbiol* **23**, 657-
631		668 (1997).
632	62	Hardy, L., Juan, P. A., Coupat-Goutaland, B. & Charpentier, X. Transposon insertion sequencing
633		in a clinical isolate of *Legionella pneumophila* identifies essential genes and determinants of natural
634		transformation. *J Bacteriol* **203**, e00548-00520 (2021).
635	63	D'Attoma, G. *et al.* Phenotypic characterization and transformation attempts reveal peculiar traits
636		of *Xylella fastidiosa* subspecies *pauca* strain De Donno. *Microorganisms* **8**, 1832,
637		doi:10.3390/microorganisms8111832 (2020).
638	64	Davis, M. J., French, W. J. & Schaad, N. W. Axenic culture of the bacteria associated with phony
639		disease of peach and plum leaf scald. *Curr Microbiol* **6**, 309-314 (1981).
640	65	Cruz, L. F., Cobine, P. A. & De La Fuente, L. Calcium increases *Xylella fastidiosa* surface
641		attachment, biofilm formation, and twitching motility. *Appl Environ Microbiol* **78**, 1321-1331,
642		doi:10.1128/AEM.06501-11 (2012).
643	66	Gluck-Thaler, E. *et al.* Repeated gain and loss of a single gene modulates the evolution of vascular
644		plant pathogen lifestyles. *Sci Adv* **6**, eabc4516 (2020).
645	67	Zhu, X. *et al.* Mitochondrial copper and phosphate transporter specificity was defined early in the
646		evolution of eukaryotes. *Elife* **10**, e64690, doi:10.7554/eLife.64690 (2021).
647	68	Sheffield, P., Garrard, S. & Derewenda, Z. Overcoming expression and purification problems of
648		RhoGDI using a family of "Parallel" expression vectors. *Protein Expr. Purif.* **15**, 34-39 (1999).
649	69	Schneider, C. A., Rasband, W. S. & Eliceiri, K. W. NIH Image to ImageJ: 25 years of image analysis.
650		*Nat Methods* **9**, 671-675 (2012).
651	70	Simko, I. & Piepho, H.-P. The area under the disease progress stairs: Calculation, advantage, and
652		application. *Phytopathology* **102**, 381-389 (2012).
653	71	Kelley, L. A., Mezulis, S., Yates, C. M., Wass, M. N. & Sternberg, M. J. E. The Phyre2 web portal
654		for protein modeling, prediction and analysis. *Nat Protoc* **10**, 845-858, doi:10.1038/nprot.2015.053
655		(2015).
656	72	Di Tommaso, P. *et al.* T-Coffee: a web server for the multiple sequence alignment of protein and
657		RNA sequences using structural information and homology extension. *Nucleic Acids Res* **39**, W13-
658		17, doi:10.1093/nar/gkr245 (2011).

659   73   Katoh, K., Misawa, K., Kuma, K. & Miyata, T. MAFFT: a novel method for rapid multiple sequence
660   alignment based on fast Fourier transform. *Nucleic Acids Res* **30**, 3059-3066 (2002).

661   74   Stamatakis, A. RAxML version 8: a tool for phylogenetic analysis and post-analysis of large
662   phylogenies. *Bioinformatics* **30**, 1312-1313, doi:10.1093/bioinformatics/btu033 (2014).

663   75   Sauvage, T., Plouviez, S., Schmidt, W. E. & Fredericq, S. TREE2FASTA: a flexible Perl script for
664   batch extraction of FASTA sequences from exploratory phylogenetic trees. *BMC Res Notes* **11**,
665   164, doi:10.1186/s13104-018-3268-y (2018).

666   76   Madeira, F. *et al.* The EMBL-EBI search and sequence analysis tools APIs in 2019. *Nucleic Acids
667   Res* **47**, W636-W641, doi:10.1093/nar/gkz268 (2019).

668   77   Crooks, G. E., Hon, G., Chandonia, J. M. & Brenner, S. E. WebLogo: a sequence logo generator.
669   *Genome Res* **14**, 1188-1190, doi:10.1101/gr.849004 (2004).

670   78   Ihaka, R. & Gentleman, R. R: A language for data analysis and graphics. *J. Comput. Graph. Stat.*
671   **5**, 299-314 (1996).

Acknowledgments

674   We thank Dr. Michelle Mendonça Pena for helping with phylogenetic analyses. We thank the Alabama

675   Supercomputing Authority for granting access to their high-performance computing platform. We

676   acknowledge funding from Alabama Agricultural Experiment Station (AAES) Hatch Program (L.D., P.A.C.,

677   N.P) and Auburn Internal Grant Program.

Author contributions

680   M.V.M., P.A.C. and L.D.F. designed research; M.V.M., X.Z. D.S., L.M.G., E.N. and N.P. performed

681   research; M.V.M., N.P., P.A.C. and L.D. analyzed data; and M.V.M. and L.D.F. wrote the paper.

Competing Interests statement

The authors declare no conflict of interest.

Figure legends

Fig. 1. Natural transformation and twitching motility phenotypes of *X. fastidiosa* mutant strains used in this

study. Quantification of natural transformation was performed by enumerating total viable cells and

recombinants and results are expressed as the ratio of recipient cells transformed (recombination

frequency; values shown in the x-axis of the chart). Twitching motility was determined by spotting cells on

agar plates and measuring the movement fringe width after 4 days of growth at 28 ºC (values shown in

the y-axis of the chart). Results of mutant strains for paralogous genes are shown using the same color

for text and symbol. The non-motile and non-recombinant mutants are shown as a black dot. The WT is

represented as a dark blue triangle and all other mutants are represented as black triangles. Data

represent means and standard errors. Different letters in parenthesis indicate significant difference in

fringe width as analyzed by ANOVA followed by Tukey's HSD multiple comparisons of means ($P<0.05$; n

= three to 14 independent replicates with eight to 48 internal replicates each). * and ** indicate significant

difference ($P<0.05$ and $P<0.005$, respectively) of recombination frequency through natural competence in

comparison to the WT as determined using Student's *t*-test (n = three to 21 independent replicates with

two internal replicates each). The detection limit for recombination frequency was 10^{-7} and for twitching

motility was 10 μm. Mutants below the detection limit are shown in the non-motile and non-recombinant

group.

Fig. 2. FimT3 DNA binding activity. **a** Transmission electron microscopy micrographs of pilus formation by

X. fastidiosa WT and Δ*fimT3* cells. Arrows are pointing to TFP. Similar events were captured in two

independent experiments. Images were captured at 31,500× magnification. Scale bars, 500 nm. **b** Uptake

of Cy-3-labeled pAX1-Cm plasmid into a DNaseI resistant state by *X. fastidiosa* cells observed using a

fluorescent microscope. The images correspond to the bright field (left), Cy-3 channel (center) and

merged images (right). In merged images, arrows are pointing to fluorescent DNA foci at the cell poles.

Similar events were captured in three independent experiments. Images were captured at 100×

magnification. Scale bars, 1.5 μm. **c** DNA-binding ability of purified FimT1s, FimT2s and FimT3s

assessed by agarose EMSA. Similar events were captured in three independent experiments. **d** Surface

electrostatics representation of FimT3 merged with its ribbon representation. The arginine amino acid

residues (R160 and R162) that were mutated for functional studies are highlighted in the figure. The

electrostatic potential (eV) color scale is shown in the figure. **e** DNA-binding ability of purified FimT3s and

FimT3s-R160AR162A assessed by agarose EMSA. Similar events were captured in three independent

experiments. **f** Titration of the DNA binding activity of FimT3s and FimT3s-R160AR162A by native

acrylamide EMSA. Similar events were captured in two independent experiments. **g** Densitometry

analysis of the DNA binding activity of FimT3s and FimT3s-R160AR162A in acrylamide EMSA. The

fluorescent intensity of the shifted DNA bands of Cy-3-labeled Km resistance cassette presented in **f** was

measured using ImageJ. The fluorescent intensity of shifted bands when treated only with 1, 2, 5, and 10

µM of each protein was measured, since they presented higher shifts in electrophoretic mobility within this

protein concentration range. **h** Area under the fluorescent curve. The area under the fluorescent curves in

g was calculated to quantify the DNA-binding affinity of FimT3s and FimT3s-R160AR162A. Statistical

significance was determined using Student's *t*-test (* indicates $P<0.05$ in comparison to the wild-type

protein; n = two independent replicates).

Fig. 3. Phylogenetic tree based on nucleotide sequences of the three *fimT* paralogs from

Xanthomonadaceae bacterial strains with whole-genome sequences available. The phylogenetic tree was

built using the Maximum-likelihood method and visualized using FigTree. Branches with bootstrap values

below 70% were collapsed, except where indicated. This was performed to keep conciseness of the

figure while indicating the different clusters for *fimT1*, *fimT2* and *fimT3*. *X. fastidiosa* is highlighted in blue

within the tree. When known, clade I *Xanthomonas* spp. are highlighted in orange, and clade II

Xanthomonas spp. are highlighted in purple. Bacterial species encoding all three *fimT* paralogs are

indicated in bold, while those encoding two paralogs are shown in italics and those encoding only one

paralog are underlined within the tree. The order of appearance of bacterial species in collapsed

branches follows the original order of appearance within the tree. For representation purposes, branches

are highlighted by color according to each *fimT* paralog. Green: *fimT1*; Dark blue: *fimT2*; Red: *fimT3*. ***

Indicates clusters within the *fimT3* clade which contain few FimT3 sequences with no GRxR motif (see

Table S5 and Fig. S15). Abbreviations are the following. ***Coralloluteibacterium***: Cst – *C. stylophorae*.

Luteimonas: Lab – *L. abyssi*; Lae – *L. aestuarii*; Laq – *L. aquatica*; Lar – *L. arsenica*; Lch – *L.*

chenhongjianii; Lcu – *L. cucumeris*; Lde – *L. deserti*; Lfr – *L. fraxinea*; Lgi – *L. gilva*; Lgr – *L. granuli*; Lhu –

L. huabeiensis; Llu – *L. lumbrici*; Lma – *L. marina*; Lme – *L. mephitis*; Lpd – *L. padinae*; Lpn – *L.*

panaciterrae; Lsp – *L.* sp.; Lte – *L. terrae*; Lti – *L. terricola*; Lwe – *L. wenzhouensis*; Lyi – *L. yindakuii*.

Lysobacter: Lys – *L.* sp. **Pseudoxanthomonas:** Pbe – *P. beigongshangi*; Pbr – *P. broegbernensis*; Pco

– *P. composti*; Pda – *P. daejeonensis*; Pdo – *P. dokdonensis*; Pge – *P. gei*; Phe – *P. helianthi*; Pin – *P.*

indica; Pja – *P. japonensis*; Pji – *P. jiangjuensis*; Pkl – *P. kalamensis*; Pko – *P. kaohsiungensis*; Pkr – *P.*

koreensis; Pme – *P. mexicana*; Psc – *P. sacheonensis*; Psn – *P. sangjuensis*; Psp – *P.* sp.; Psx – *P.*

spadix; Psu – *P. suwonensis*; Ptw – *P. taiwanensis*; Pwi – *P. winnipegensis*; Pwu – *P. wuyuanensis*; Pye

– *P. yeongjuensis*. **Rehaibacterium:** Rte – *R. terrae*. **Silanimonas:** Sle – *S. lenta*. **Stenotrophomonas:**

Sac – *S. acidaminiphila*; Sbe – *S. bentonitica*; Sch – *S. chelatiphaga*; Scy – *S. cyclobalanopsidis*; Sda –

S. daejeonensis; Sge – *S. geniculata*; Sgi – *S. ginsengisoli*; Shu – *S. humi*; Sin – *S. indicatrix*; Skr – *S.*

koreensis; Sla – *S. lactitubi*; Sma – *S. maltophilia*; Sne – *S. nematodicola*; Sni – *S. nitritireducens*; Spn –

S. panacihumi; Spv – *S. pavanii*; Spi – *S. pictorum*; Srh – *S. rhizophila*; Sse – *S. sepilia*; Ssp – *S.* sp; Ste

– *S. terrae*; Stu – *S. tumulicola*. **Vulcaniibacterium:** Vge – *V. gelatinicum*; Vte – *V. tengchongense*; Vth –

V. thermophilum. **Xanthomonas:** Xab – *X. albilineans*; Xaf – *X. alfalfae*; Xar – *X. arboricola*; Xax – *X.*

axonopodis; Xbr – *X. bromi*; Xcm – *X. campestris*; Xcn – *X. cannabis*; Xcs – *X. cassavae*; Xci – *X.*

cissicola; Xct – *X. citri*; Xco – *X. codiaei*; Xcu – *X. curcubitae*; Xdy – *X. dyei*; Xer – *X. euroxanthea*; Xeu –

X. euvesicatoria; Xfl – *X. floridensis*; Xfr – *X. fragariae*; Xho – *X. hortorum*; Xha – *X. hyacinthi*; Xhd – *X.*

hydrangea; Xml – *X. maliensis*; Xms – *X. massiliensis*; Xme – *X. melonis*; Xna – *X. nasturtii*; Xor – *X.*

oryzae; Xpe – *X. perforans*; Xpi – *X. pisi*; Xpo – *X. populi*; Xpr – *X. prunicola*; Xsa – *X. sacchari*; Xso – *X.*

sontii; Xsp – *X.* sp.; Xth – *X. theicola*; Xtr – *X. translucens*; Xva – *X. vasicola*; Xve – *X. vesicatoria*.

Xylella: Xtw – *X. taiwanensis*.

Fig. 4. Schematic representation of the functional role of TFP molecular components in natural

competence and twitching motility of *X. fastidiosa*. The figure shows the structure of a type IV pilus (PilA1,

PilA2 and PilA3), the alignment subcomplex (PilM, PilN, PilO and PilP), the inner membrane motor

subcomplex (PilB, PilC, PilD, PilT and PilU), the pore subcomplex (PilF and PilQ), minor pilins (PilE1,

PilE2, PilV1, PilV2, PilW1, PilW2, PilX1, PilX2, FimT1, FimT2 and FimT3), TFP tip adhesins (PilY1-1,

PilY1-2 and PilY1-3), as well as regulatory proteins (PilG, PilH, PilI, PilJ, PilL, PilR, PilS, PilZ, ChpB and

ChpC). Proteins in the figure are color-coded according to their phenotype upon knockout mutation and

thus functional role. More information about the role of each protein is described in Table S1, throughout

the manuscript and in SI Discussion. OM – outer membrane; P – periplasm; IM – inner membrane.