Representation and quantification Of Module Activity from omics data with rROMA

Matthieu Najm[1,2,3#], Matthieu Cornet[1,2,3#], Luca Albergante[1,2,3], Andrei Zinovyev[1,2,3], Isabelle Sermet-

Gaudelus[4,5,6], Véronique Stoven[1,2,3], Laurence Calzone[1,2,3] and Loredana Martignetti [1,2,3 *]

[1] INSERM U900, 75428 Paris, France, [2] Center for Computational Biology, Mines ParisTech, PSL

Research University, 75006 Paris, France, [3] Institut Curie, PSL Research University, 75248 Paris,

France, [4] Faculté de Médecine, Université de Paris, Paris, France, [5]Institut Necker Enfants Malades,

INSERM U1151, Paris, France, [6]AP-HP. Centre - Université Paris Cité; Hôpital Necker Enfants Malades,

Centre de Référence Maladie Rare - Mucoviscidose, Paris, France.

[#] These authors contributed equally to this work;

[*] To whom correspondence should be addressed: loredana.martignetti@curie.fr

Abstract

The efficiency of analyzing high-throughput data in systems biology has been demonstrated in

numerous studies, where molecular data, such as transcriptomics and proteomics, offers great

opportunities for understanding the complexity of biological processes.

One important aspect of data analysis in systems biology is the shift from a reductionist approach

that focuses on individual components to a more integrative perspective that considers the system

as a whole, where the emphasis shifted from differential expression of individual genes to

determining the activity of gene sets.

Here, we present the rROMA software package for fast and accurate computation of the activity of

gene sets with coordinated expression. The rROMA package incorporates significant improvements

in the calculation algorithm, along with the implementation of several functions for statistical

analysis and visualizing results. These additions greatly expand the package's capabilities and offer

valuable tools for data analysis and interpretation. It is an open-source package available on github

at: www.github.com/sysbio-curie/rROMA.

Based on publicly available transcriptomic datasets, we applied rROMA to cystic fibrosis, highlighting

biological mechanisms potentially involved in the establishment and progression of the disease and

the associated genes. Results indicate that rROMA can detect disease-related active signaling

pathways using transcriptomic and proteomic data. The results notably identified a significant

mechanism relevant to cystic fibrosis, raised awareness of a possible bias related to cell culture, and

uncovered an intriguing gene that warrants further investigation.

Contact: loredana.martignetti@curie.fr

Keywords

Pathway activity, Principal Component Analysis, Transcriptomics, Proteomics

Introduction

The use of high-throughput molecular techniques, such as transcriptomics and proteomics, is

becoming increasingly easier with the improvement of data acquisition tools, leading to a drastic

decrease in the costs associated with such analyses. This allows for precise measurement of the

molecular profiles of biological systems at several levels. However, the amount of data produced

during such experiments is very important and requires the use of dedicated software and

algorithms to analyze them. Moreover, the ability to interpret the data in terms of biological

processes becomes a crucial issue. Dedicated analyses are needed to synthesize and transform the

data into valuable biological information [1].

A commonly employed approach in genomics involves comparing measurements at the individual

gene or protein level to identify distinctive markers indicative of specific disease states (biomarkers)

or genes that play a causal role in the studied disease [2]. Nonetheless, in numerous systemic

diseases, the disruption of a signaling pathway can arise from distinct genes within that pathway,

and these gene alterations may vary from one patient to another. For example, in cancer It has

become apparent in recent years that the same pathways are affected by defects in different genes

and that the molecular profiles of patient samples are more similar at the pathway level than at the

individual gene level [3]. Therefore, quantification of gene set activity from omics measurements is

now widely used to transform gene-level data into associated sets of genes representing biological

processes [4]. By employing gene set-based approaches in the analysis of omics data, it becomes

possible to capture valuable biological insights that would otherwise remain undetectable when

solely focusing on individual genes.

In this study, we developed an algorithm, implemented as an R package called rROMA, which was

designed to quantify the activity of sets of genes characterized by their involvement in a common

functional role. Gene set approaches have become very popular to summarize individual molecular

measurements into more interpretable pathways and biological processes. Gene sets can thus

correspond to genes with the same functional activities, genes regulated by the same motifs, genes

belonging to the same signaling pathway, target genes of a transcription factor or genes forming a

group of frequently co-expressed genes. The underlying hypothesis of rROMA is to assess the

activity of a gene set by determining the maximum amount of one-dimensional variance, which is

represented by the first principal component (PC1) derived from the gene within the set. This

quantity is proportional to the influence of a single latent factor on the gene expression within the

gene set and reflects the variability of this factor's activity across the studied samples. This setting

corresponds to the uni-factor linear model of gene expression regulation [5].

The strength of our method lies in its ability to calculate a p-value, a statistical measure that serves

as a crucial indicator of the gene set activity's significance. This p-value enables the prioritization of

biological processes that are pertinent to the underlying conditions in question. Another unique

feature of the rROMA algorithm and to our knowledge not implemented in any other algorithm

available in the literature is its ability to estimate the statistical significance of the distribution of

samples along the first component for a gene set in two ways: it distinguishes between *shifted* and

over-dispersed gene sets. The fact that rROMA distinguishes these two situations is particularly

useful because, in many cases, the activity of a gene set does not correspond to overdispersion of

the module in the global gene expression space but to a shift of the genes in a particular direction.

Therefore, analysis of shifted gene sets will highlight findings that would not be identified with

overdispersion analysis alone.

We applied rROMA to examine pathway activities in airway epithelial cells of individuals with Cystic

Fibrosis (CF) and healthy individuals, shedding light on the molecular processes behind CF

phenotypes. Moreover, we illustrate the use of rROMA in estimating cell type abundance through

bulk transcriptomic data analysis. Here, rROMA enables the clear estimation in cell-type proportions,

enhancing the potential accuracy of gene signatures that encompass both upregulated and

downregulated genes. Finally, we show application of rROMA to a mass-spectrometry-based breast

cancer proteomic dataset, showing its capability of identifying significantly dysregulated pathways in

clinical breast cancer subtypes.

Results

Review of existing methods

Over the past decades, numerous tools have emerged to perform gene set analysis from omics data

to explore pathway activity. Sample-wise pathway activity computation and global pathway

enrichment scoring are two distinct approaches used in the analysis of biological pathways. The

primary goal of sample-wise pathway activity analysis is to assess the extent to which a particular

biological pathway is active or dysregulated in each individual sample. Methods for sample-wise

analysis provide a quantitative measure of pathway activity for each pathway in each sample that

can then be interpreted in terms of pathway activity profiles. Global pathway enrichment scoring, on

the other hand, condenses the information from all samples into a single enrichment score,

providing an overview of the pathway's significance or activity across the study. This approach is

commonly used in over-representation analysis (ORA) of differentially expressed genes [6], gene set

enrichment analysis (GSEA) [7] and similar methods that provide an aggregated score of pathway

enrichment between two biological conditions.

A simple method for sample-wise pathway activity quantification consists of calculating the mean or

median of the expression of genes annotated in a given pathway in each sample. Alternatively, one

can rely on the expression of a single marker gene that represents the overall activity of the pathway

in different samples. A more sophisticated gene set quantification method, called PLAGE (Pathway

Level Analysis of Gene Expression) was introduced by Tomfohr and colleagues [8] to compute the

activity of a gene set by calculating the first principal component (PC1) of the expression matrix

restricted to the genes in the gene set. In the study by Bild et al [9], this PCA-based strategy was

exploited to define activity of several cancer-related pathways on a large collection of human cancer

transcriptomes. This PCA-based method is efficient in modeling situations in which individual genes

within the pathway display different expression patterns across samples, either due to their unequal

contributions to pathway activation or to their involvement encompassing both upregulation and

downregulation. This is particularly relevant when some genes have stronger or opposite expression

variation than others, such as transcription factors. However, the PLAGE algorithm has some

limitations. Specifically, it does not provide a statistical assessment of whether the observed activity

in a particular pathway is statistically significant or merely occurring by random fluctuations.

Moreover, the standard PCA calculation implemented in PLAGE lacks the capability to identify

pathway activation when dealing with a specific gene configuration, which is referred to here as a

shifted gene set.

Many other methods have been developed to estimate pathway activity from transcriptomic data. A

recent comprehensive and systematic comparison of existing computational tools evaluates

different approaches [10]: (i) ranking-based methods such as single sample GSEA (ssGSEA) [11],

GSVA [12], (ii) PCA-based methods, (iii) pathway topology-based methods such as PARADIGM [13]

and many others. In this study, PCA-based methods have been shown to perform best in preserving

characteristics of the original gene expression data while transforming the transcriptome data at the

pathway level whereas ranking-based methods show high robustness against noise. In Table 1, we

reported a feature-wise comparison of rROMA to existing bulk-based pathway analysis tools. This

table comprises methods that quantify pathway activity on a sample-wise basis, considering only the

expression matrix. Other methods exist that necessitate further input such as prior knowledge or

inferred networks to infer pathway activity, such as PARADIGM [13], VIPER [14], PROGENy [15].

Recently, multiple algorithms for pathway activity analysis of single cell transcriptomic data have

been proposed, including PCA-based methods such as PAGODA [16] and MAYA [17]. Benchmark

studies [18,19] comparing their performances in interpreting sparse and noisy single cell

transcriptomic datasets in terms of functional pathways suggest that bulk-based pathway analysis

tools can be applied to single cell datasets, partially outperforming dedicated single-cell tools.

Method	Type of algorithm	Overdispersed gene-sets	Shifted gene-sets	Statistical significance	A priori fixed gene weights	Outlier detection	Code availability	Ref
z-score	Standardized expression		X				R	[20]
GSVA	Ranking-based		X				R	[12]
ssGSEA	Ranking-based		X				Java, R	[11]
PLAGE	PCA-based	X					R	[8]
GO-PCA	PCA-based	X		X			Python	[21]
PCGSE	PCA-based		X	X			R	[22]
rROMA	PCA-based	X	X	X	X	X	R	
Pathifier	Principal curve	X	X				Java, R	[23]

143
Table 1. Feature-wise comparison of rROMA to existing tools

Implementation of rROMA

The rROMA algorithm expands the repertoire of existing PCA-based methods for pathway activity

quantification by introducing unique functionalities, notably (i) a statistical assessment of pathway

activity measurement, enabling the determination of the biological relevance of the results and

facilitating their prioritization; (ii) a PCA computation with a fixed center which allows to detect two

different configurations of pathway genes in the global gene expression space, referred to as *shifted*

and *over-dispersed* gene sets.

The package rROMA is an evolution of the algorithm ROMA, a program originally developed in Java

[24]. The new rROMA software, developed in R language, incorporates significant improvements

including an automatic procedure to not only distinguish between *shifted* and *over-dispersed* gene

sets but also automatically detect outlier genes that can affect the PCA computation. Moreover, the

new software includes several novel functions for user-friendly downstream statistical analysis and

graphical visualization of results that are documented in a software accompanying vignette. These

additions greatly expand the package's capabilities and offer valuable tools for data analysis and

interpretation. The rROMA software is open-source and available on github at:

www.github.com/sysbio-curie/rROMA. The workflow of the rROMA algorithm is schematized in

Figure 1. A detailed vignette to reproduce an example of analysis is available on the github

containing the source code of the algorithm.

The algorithm requires as input a genome-wide expression data matrix and a gene set annotation

file in GMT (Gene Matrix Transposed) format. Optionally, it is possible to incorporate prior

knowledge about the significance of certain genes within a pathway by initially assigning gene

weights in a given pathway within the GMT file. These "adjusted weights" will then substitute the

weights acquired from the PC1 calculation for those specific genes.

The rROMA algorithm starts by imputing missing values in the data matrix and centering values, if

not done beforehand. Then, each gene set is analyzed separately. A multistep procedure is

implemented for (i) extracting the expression submatrix corresponding to a given gene set, (ii)

quantifying the PC1-based activity values robust to outlier samples and outlier genes and (iii)

assessing the statistical significance of the activity values.

First, the algorithm extracts the expression submatrix corresponding to a given gene set. On this

submatrix, it computes the PCA in the sample space via the SVD algorithm [25]. Two measures of

interest are then considered: the percentage of variance explained by the PC1 (referred to as *L1*) and

the median value of the gene projections onto PC1 (referred to as *Median Exp*). Based on these two

values, pathway activity significance is estimated in two distinct ways through the definition of

shifted and *overdispersed* gene sets (Figure 2). A *shifted* set of genes corresponds to the case where

the median expression (*Median Exp*) of all the genes in the gene set is significantly different from the

one of all the genes studied, i.e. that the gene set shows a particularly high expression in at least one

sample (Figure 2A). *Over-dispersion* of a gene set corresponds to the situation where the amount of

variance explained by PC1 (*L1*) calculated for only the genes in that gene set is significantly greater

than the variance of a randomly selected set of genes of the same size. Thus, overdispersion means

greater variability in a set of genes among the considered samples (Figure 2B). An automatic

procedure for the identification of outlier genes and samples is applied to increase robustness of the

PC1 computation (see Methods).

The same measures *L1* and *Median Exp* are computed for *N* randomly generated gene sets of the

same size, which will constitute the reference null distribution. The *Median Exp* value is compared to

the distribution of *Median Exp* values obtained for the random gene sets. If less than 5% (value

defined by a tunable hyperparameter) of the values obtained in the null distribution are lower than

the one obtained for the gene set under study, the latter is said to be *shifted* (ppv Median Exp

<0.05). If less than 5% (value defined by a modifiable hyperparameter) of the *L1* values obtained in

the null distribution are lower than the one obtained for the gene set under study, then the latter is

said to be *overdispersed* (ppv L1 <0.05). The ppv L1 and ppv Median Exp values are corrected for

multiple testing by the Benjamini and Hochberg method [26].

As output, the rROMA software generates an R object including multiple results: (i) a module matrix

detailing *L1* and *Median Exp* values and their corresponding *p*-values and *q*-values for each pathway,

(ii) a sample matrix that presents the activity level of each pathway across samples derived from PC1

and (iii) a gene weight list for each gene set, which reports the gene projections in the PC1 space.

For a *shifted* gene set, the measurement of sample activity level is particularly important. It

determines which samples are responsible for the shift of the gene set. If the samples were already

separated into several conditions prior to the analysis, it is possible to verify that this separation into

groups is indeed responsible for the shift, by checking that the activity levels are significantly

different between the conditions. Conversely, if the conditions are not known *a priori*, it is possible

to determine new groups by performing a hierarchical clustering analysis on the pathways that are

shifted, and thus determine potential new groups of interest from the analysis.

The analyses mentioned above for the case of shifted gene sets are also valid for *over-dispersed*

gene sets. However, in the case of the latter, the analysis of gene weights, meaning the loadings

from the PC1, also becomes interesting: the genes associated with the highest weights are the

driving force in the activity scores of the gene sets. They summarize the information of the gene

sets, which can be particularly useful for assessing the importance of some genes. They can further

be used as an input for systems biology approaches.

The analysis of the sign of the gene weights is also particularly relevant. For example, gene sets

containing both activators and inhibitors can be highlighted by rROMA by being overdispersed, and

the associated genes highlighted. Such gene sets would not be detected as overdispersed by

methods based on the average expression of genes in the samples.

Uncovering active pathways in Cystic Fibrosis with rROMA

We applied rROMA to investigate the activity of pathways in airway epithelial cells from CF patients

and healthy donors. More precisely, we compared the transcriptomes of primary cultures of airway

epithelial cells from patients (N=6) with those of healthy controls (N=6), based on RNAseq data

publicly available in the NCBI's GEO database, under the accession ID GSE176121 [27]. rROMA was

run by specifying the pathway database to use and the expression matrix to analyze, as shown in the

accompanying vignette. Here, the Molecular Signature Database MsigDB hallmark gene set

collection [28] was used, a gene set collection of 50 gene sets specifically curated to represent core

biological processes and pathways that are commonly dysregulated in cancer. However, to provide a

more complete view of the biological processes involved in a study, rROMA can be used with

different reference databases.

The results of rROMA highlight pathways that are provided in the *ModuleMatrix* output. Pathways

with a *ppv Median Exp* lower than a given threshold were deemed as *shifted*, while those with a *ppv*

L1 lower than this threshold were *overdispersed*. The *Plot.Genesets.Samples* function allows for the

visualization of activity scores of significantly shifted and overdispersed pathways across samples, in

the form of a heatmap representation (Figure 3). rROMA identified two shifted pathways,

APICAL_SURFACE which is found to have higher activity scores in healthy controls than in CF

patients, and FATTY_ACID_METABOLISM which has higher scores in CF patients than in controls, and

an overdispersed pathway, COAGULATION, with higher activity scores in healthy controls than in CF

patients.

When sample groups have been pre-defined, as CF or control groups in our case, these groups can

be compared based on the activity scores of the gene sets observed in the samples specific to the

two groups. Boxplots of the activity scores based on predefined groups can facilitate the

interpretation and complement a typical differential analysis. In our study, shifted and overdispersed

pathways behaved differently in CF patients versus healthy controls, as shown in Figure 3.

Alternatively, when the groups are not predefined, analyzing the shifted and overdispersed

pathways can reveal clusters of samples exhibiting similar pathway activity.

The analysis of top contributing genes in each pathway also provides crucial information. The

weights assigned to each gene in the PC1 vector allow to identify the key genes that most contribute

to variations in pathway activities as those with the higher weights. For each pathway, gene weights

are provided by the PlotGeneWeight function. For example, plotting the gene weights for the

COAGULATION pathway (see Figure 4) highlighted the GSN gene, encoding the protein *Gelsolin*, as

the highest contributor to the activity score. Notably, *Gelsolin* has been previously reported to play a

role for CFTR activation [29,30], and to promote mucus fluidification in CF [31].

Finally, many hyperparameters can be specified and changed to modify rROMA speed, precision, or

behavior regarding outlier detection. Details about all available hyperparameters are described in

the vignette accompanying the software. The computational time required to run the algorithm

typically depends on the number of considered pathways and their relative sizes. It also depends on

whether parallelization is enabled. In the present work, the algorithm ran in about 3 minutes and 15

seconds on a MacBook Pro equipped with a 2,6 GHz Intel Core i7 6 cores processor. A single analysis

60 pathways took roughly 5 second. Parallelization was not used, but this would have increased the

speed of the analysis.

Comparison with state-of-art methods

We compared rROMA with four sample-wise pathway quantification methods implemented in the

GSVA package [12] that cover different types of algorithms, namely GSVA, PLAGE, ssGSEA and z-

score. We conducted a comparison of these methods using the RNAseq data from the same

transcriptomic study (GSE176121) on airway epithelial cells. We selected the samples to simulate

two distinct analysis scenarios. In the first scenario, as previously described, we compared CF and

healthy control samples. Our goal was to explore a pathway database to identify pathways that

could differ between these two conditions without any a priori hypotheses of the specific pathways

involved. In the second scenario, we compared the gene expression data of six healthy control

samples before and after treatment with IL17+ TNFα. In this case, our aim was to assess whether the

different methods would effectively reveal the activation of the IL17 and TNFα pathways.

Although only rROMA provides a statistical significance of the pathway activity scores, for the other

methods significance with respect to a phenotype can be evaluated using conventional statistical

models. Therefore, for each method we compared the distributions of the pathway activity scores

between the two groups with a Student t-test and considered dysregulated pathways between the

two groups as those with adjusted p-values for the statistical test less than 0.05.

In the first scenario, the GSVA, ssGSEA, and z-score methods did not reveal any pathways as

dysregulated between CF and non-CF samples based on the statistical test. In contrast, PLAGE

identified dysregulation in 45 out of 50 pathways, i.e. a large number of pathways which prevents

meaningful interpretation with respect to the IL17+TNFα treatment. However, rROMA detected

eight pathways with significant activity differences between CF and healthy samples, shedding light

on potential biological mechanisms that could play a role in the disease.

In the second scenario, the TNFα signaling pathway was found significantly active in the samples

treated with IL17+TNFα compared to the control ones with all methods. We then compared the

ranking of this pathway among the significant results for the different methods. The results obtained

for the different methods are provided in Supplementary Tables 1-10. The TNFα pathway's ranking

among significantly dysregulated pathways based on the t-statistics was as follows: 5 out of 24 for

GSVA, 5 out of 27 for ssGSEA, 7 out of 17 for the z-score and 18 out of 45 for PLAGE. In contrast, the

rROMA algorithm highlights TNFα signaling pathway as the top significantly shifted pathway, thus

providing the most consistent result with the expected output. Among the rROMA results in this

scenario, we pointed out an example in which the P53 pathway exhibits significant overdispersion,

with target genes being affected in opposite directions by the transcription factor (Supplementary

Figure 1).

All results obtained for the two scenarios with the different methods are provided in Supplementary

Tables 1-10 and can be reproduced using the codes publicly available in the following github

repository: https://github.com/sysbio-curie/rRoma_comp.git

rROMA estimates cell type abundances from bulk transcriptomic data

Furthermore, we showcase how rROMA can be utilized to explore cell type abundance using bulk

transcriptomic data. Saint-Criq et al. [32] investigated the impact of two differentiation media on

primary cultures of CF and non-CF airway epithelial cells, as determined by transcriptomic data. They

built a gene signature for each cell type using the 50 most expressed markers derived from a single-

cell RNA sequencing (scRNAseq) dataset [33]. They observed a significant overexpression of genes

belonging to the signature of the secretory cell subtype in cultures grown in one of the media

(referred to as UNC), compared to the other medium (referred to as SC). Conversely, gene markers

of the ciliated subtype were overexpressed in primary cultures grown using the SC medium

compared to UNC medium.

We applied rROMA to the Saint-Criq RNA seq dataset to estimate cell type abundance in CF and non-

CF samples. More precisely, the Plasschaert signature [33] for each cell type was used and gene

reference gene sets, and the activity scores of these gene sets across the samples were represented

in the form of a heat map. Samples were found to be clustered according to the differentiation

medium in which they were grown, and our results confirmed the higher abundance of the ciliated

cell subtype in UNC medium and higher abundance of the secretory cell subtype in SC medium

(Figure 5A). We repeated the rROMA analysis using an alternative signature from Okuda and

colleagues [34]. In contrast with the Plasschaert signature, the Okuda signature includes the most

differentially expressed genes in each cell type, encompassing both overexpressed and

underexpressed genes. As illustrated in figure 5B, rROMA analysis using the Okuda signature

consistently revealed the same relative abundances of secretory and ciliated subtypes between UNC

and SC growing media, as observed with the Plasschaert signature. Thus, rROMA allows us to clearly

highlight differences in cell-type abundances, facilitating the use of gene signatures that contain

both upregulated and downregulated genes and thus potentially more accurate.

Functional analysis of proteomic data with rROMA

Finally, we applied rROMA to study a mass-spectrometry-based proteomic dataset of 9771 proteins

quantified in 122 treatment-naive primary breast cancers [35]. Individual samples were

characterized by their activity levels computed by rROMA for 50 pathways in the MsigDB hallmark

gene set collection [28]. We identified 34 shifted and 6 overdispersed pathways with p-value lower

than 0.05. The samples were clustered based on their activity profiles across the significant

pathways, using unsupervised hierarchical clustering and Euclidean distance.

Overall rROMA achieved satisfactory results with the proteomic dataset. Unsupervised clustering

divided breast cancer samples into six clusters (Figure 6), showing a higher heterogeneity of the

disease compared to the clinical stratification based on five intrinsic PAM50 subtypes [36] (Figure 6).

Cluster 1 predominantly contains basal tumors, whereas the Luminal B tumors are mainly associated

with cluster 5. These two clusters are characterized by high activation of TNFα signaling via NFK-β,

epithelial to mesenchymal transition and inflammatory pathways. These observations are consistent

with known facts about inflammatory and basal breast cancer [37,38]. Luminal A tumors are

segmented mainly in two clusters, namely cluster 4 and cluster 6, both showing high activation of

estrogen response, which is consistent with the fact that ER is the defining and driving factor in

luminal breast cancer.

Discussion

Quantifying the activity of biologically related gene sets is a commonly employed approach to

extract valuable biological insights from high-throughput data. The use of gene sets as aggregated

variables from molecular data enables the capture of biological information that may not be

detectable when solely focusing on individual genes. To address this challenge, we introduced the

rROMA algorithm, a PCA-based approach for quantifying pathway activity. Based on a gene

expression data matrix, this algorithm implements a linear model of gene regulation and efficiently

and reliably quantifies the activity of gene sets by computing the first principal component (PC1),

while also evaluating the statistical significance of this approximation.

Importantly, the algorithm provides the activity level of the gene set for each individual sample and

does not require a predefined labeled classification of the samples into various conditions or groups.

These activity levels can be subsequently compared, bringing to light the heterogeneity present in

the dataset in relation to the analyzed gene sets. This may be useful to define groups of samples or

patients when such stratification is unknown. Furthermore, the algorithm capability of identifying

shifted and *over-dispersed* gene sets is very peculiar compared to existing methods.

We applied rROMA to CF transcriptomic datasets, highlighting some biological mechanisms

potentially involved in the initiation or progression of the disease, and their associated genes. In our

study, out of the 50 hallmark pathways tested, 3 were significantly active:

FATTY_ACID_METABOLISM, APICAL_SURFACE, and COAGULATION. The FATTY_ACID_METABOLISM

pathway has significantly different activity scores between CF patients and healthy donors. This

pathway has been extensively studied in CF, and essential fatty acid deficiency is a well-known CF

phenotype (for a review, see [39]). The APICAL_SURFACE can be related to another well-known

hallmark of CF, i.e. a perturbation of airway surface secretory mucus content. Finally, the

COAGULATION pathway, the only overdispersed pathway in our study, seems to be highlighted due

to one specific gene with a very high associated weight, that is by far the most contributing gene to

the activity score of this pathway: gelsolin (GSN). Gelsolin has been reported as playing a role for

CFTR activation [29, 30], which suggests that the role of this gene in CF disease may be interesting to

study in more detail. The goal of this use case was not to undertake a detailed systems biology

approach of CF, which is beyond the scope of the present paper. In particular, it would require us to

include additional transcriptional dataset to take more samples into account and increase the

statistical power of our analyses, and to test several reference databases of gene sets. Overall, this

case study illustrates that rROMA can identify disease-associated pathway deregulations from

transcriptomic data, allowing a more comprehensive and functional interpretation of the data. It is

also a versatile tool that can shed light on various biological questions such as highlight the key

genes driving these deregulations, identify clusters of samples, study samples' cell-type composition,

or other cellular changes in a broader biological perspective. Indeed, the output of rROMA, in

particular the top weighted genes of significantly active pathways, can be interpreted as nodes

comprising genes and proteins of importance in the system under investigation, which can be used

as inputs to construct mathematical models.

Finally, we showcased the application of rROMA for functional analysis of proteomic data. In this

study, rROMA was applied to a proteomic dataset consisting of 9771 proteins from 122 treatment-

naive primary breast cancer samples. The results showed that rROMA performed well with the

proteomic dataset, revealing distinct clusters of breast cancer samples and higher disease

heterogeneity compared to clinical stratification based on the PAM50 subtypes. Furthermore, the

pathways identified as active by rROMA within these clusters are consistent with established

knowledge regarding signaling pathways relevant to the subtypes, such as high activation of TNFα

signaling via NFK-β, epithelial to mesenchymal transition and inflammatory pathways in the basal

subtype and high activation of estrogen response in the luminal subtype.

Methods

First Principal Component and the Simplest Uni-Factor Linear Model of Gene Expression

Regulation

The main idea of rROMA is based on the simplest uni-factor linear model of gene regulation in which

it is assumed that the expression of a gene *G* in sample *S* is proportional to the activity of one latent

biological factor *F* (which can be a transcription factor or any other endogenous or exogenous factor

affecting gene expression) in sample *S* with positive or negative (response) coefficient. Within this

model, the expression of a gene *G* in a sample *S* is proportional to the activity of a factor *F*, so that

we can write:

Expression(gene *G*, sample *S*) $\approx \alpha^F_G$ Activity ^F_S (1)

where α^F_G is the response coefficient of the gene *G* to the factor *F* and Activity ^F_S is the activity of

factor *F* in the sample *S*. These two values can be easily determined by considering the first

component of the PCA of the genes of the considered gene set in the space of the samples. In this

case, the vector containing the activities of the different samples corresponds to the first

eigenvector, i.e. the first column of the weight matrix. The vector containing the response

coefficients of the different genes in the considered gene set corresponds to the projection of the

genes onto the first component.

The uni-factor model underlying rROMA presumes that an unobserved factor (i.e. a latent factor)

acts on the gene expression variables observed in the gene set and that this action is characterized

by the calculated weights. The weights indicate the strength and direction of effect, and can have

opposite signs, as in the case in which a transcription factor has an activating action on some genes

of the set and a repressor action on others.

As many other applications of PCA, rROMA uses SVD [25] to speed up the calculation. rROMA uses

the R irlba package [40, 41], which allows to focus on the first principal component, by computing

only the first column of V and the first value of S. However, if the data are not centered, the

simplification is no longer valid, and we must now consider the fact that the mean is not zero. The

rROMA algorithm therefore starts by centering the data of the global matrix.

Pre-processing of data for rROMA analysis

The input format for gene or protein expression for rROMA is a tab-delimited text file with columns

corresponding to biological samples, and rows corresponding to genes or proteins. The first row is

assumed to contain the sample identifiers while the first column is assumed to contain the non-

redundant gene or protein names.

If the data table contains missing values, they can be imputed using an approximation of the data

matrix with missing values by a full matrix of lower rank. To do this, the user must specify the rank of

the approximate full matrix that he wants to use. Then, the principal components are calculated up

to the specified rank, using an algorithm capable of working with missing data [42]. This PCA

decomposition is used to construct the full approximate matrix of lower rank, from which the

missing values of the original data are imputed. In the rest of the algorithm, the full imputed matrix

is used.

Orientation of the PC1

In standard PCA, all components are calculated with a sign ambiguity: there is mirror symmetry,

which makes it difficult to determine whether a given set of genes is over- or under-activated. In

rROMA, several methods exist to solve this ambiguity.

If knowledge exists about the role of a gene in a given gene set, we recommend using it by

associating a sign with the effect of the gene in the gene set: negative for an inhibitor, and positive

for an activator, for example. rROMA then uses the information about these signs to choose the

orientation that maximizes the number of genes associated with a positive sign whose projection in

PC1 is positive, and the number of genes associated with a negative sign whose projection in PC1 is

negative.

Although less efficient, other methods of orienting the first component exist in the case where there

is no a priori knowledge about the genes in the reference gene sets. The most efficient method

consists in considering only the genes associated with the most extreme weights according to the

first component (according to a percentage defined by a modifiable hyperparameter), and then

multiplying these weights by the expression level of the corresponding genes. If the result is

negative, then the orientation of the PC1 is reversed. The principal behind this method is to orient

the first component to maximize the positive correlation between the expression of genes and the

PC weight.

Filtering of outlier samples

Measurements may have been performed incorrectly in some samples and keeping them may lead

to erroneous results from rROMA. By default, in the algorithm, no sample filtering is performed, as

the matrix in input is assumed to contain only correct samples. However, outlier sample detection

can be activated by a hyperparameter, and a filtering procedure in two steps is then applied based

by (i) checking the samples for a similar number of detected genes and (ii) PCA computation and

projection of samples in the gene space for identifying data points (samples) that deviate

significantly from the data distribution. Here, the number of PCs used to perform the filtering and

the distance threshold for defining outliers are defined by two hyperparameters. It is also possible to

apply only one of the two filtering steps.

Filtering of outlier genes

The calculation of PC1 can be affected by the presence of an outlier gene in the dataset. This outlier

could indeed artificially affect the PC1. To increase the robustness of the PC1 calculation, we use in

rROMA the "leave-one-out" cross-validation approach [43], working as follows:

I.   for each gene in the selected gene set, the method calculates the percentage of

460   variance explained by the first principal component (PC1) when that gene is

461   removed from the dataset. This value is denoted as *L1*, and each gene is associated

462   with its *L1* value;

463   II.   the distribution of these *L1* values is normalized by centering and scaling to obtain z-

464   scores for each gene;

465   III.   those genes whose associated z-scores exceed a specified threshold value are

466   identified as outliers and removed from the analysis.

The idea behind this method is to identify genes that have too much impact on the PC1 on their

own: if the percentage of variance explained by PC1 increases significantly in the absence of a single

gene, this means that this gene does not follow the alignment of all the others. It is then considered

as an outlier gene.

When a gene is considered an outlier for a given gene set, it is only removed from that gene set. This

default behavior can be modified by a hyperparameter, so that genes are completely removed from

all analyses, as soon as they are considered outliers in at least one gene set. However, in some cases,

these genes may still be of interest for analysis. Additional analysis steps for these genes are

therefore available in rROMA. In particular, the greater the number of gene sets analyzed containing

a given gene, the more likely it is that the gene will be considered as an outlier in at least one gene

set, and therefore be removed from the analysis. To avoid such abusive withdrawals, a Fisher test is

performed. This consists of comparing the average proportion of collections in which the genes in

the analysis are considered as outliers to this same proportion for a particular gene. If the proportion

of outliers is close to the average proportion for all the genes, then it is no longer considered an

outlier. Conversely, if the proportion of aberrations is significantly higher (threshold determined by a

modifiable hyperparameter), then it is still considered an outlier.

Instead, a gene present only in a small number of gene sets may be important for the understanding

of these gene sets. In such cases, it is not desirable to remove it, even if it is considered an outlier.

Thus, if a gene is present in less than a defined number of gene sets (set by a modifiable

hyperparameter), then it is not considered an outlier, regardless of the results of the "leave-one-

out" approach for it.

Finally, it is possible that a significant proportion of genes are considered outliers for a given gene

set. However, removing too many genes can totally distort the detectable activity for a gene set. A

final filter therefore exists to limit the maximum proportion of genes that can be considered as

aberrant for a given gene set and removed from the analysis (the proportion is set by a modifiable

hyperparameter). In this way, only genes with the most extreme leave-one-out behavior are

effectively considered outliers for this gene set.

The rROMA algorithm uses PCA calculations at different stages of its workflow, each serving a

specific purpose. These stages include imputing missing values, identifying outlier samples and

genes, and computing pathway activity. For missing value imputation, rROMA utilizes the iterative

PCA method implemented in the *mice* package [44] on the global expression matrix, imputing any

missing values within the dataset. To detect outlier samples, PCA is performed on the global

expression matrix with imputed missing values. Samples are treated as observations, and genes

serve as variables in this space, allowing for the identification of observable sample outliers. To

identify outlier genes and calculate pathway activity, PCA is carried out within a submatrix that

exclusively contains genes related to the specific pathway of interest. In this context, genes are

treated as observations, and samples serve as variables for PCA analysis.

Optimization of the calculation of null distributions

The p-value is computed as the probability of obtaining the observed activity measure for a specific

gene set by chance. The rROMA algorithm uses a random gene set procedure to generate a null

distribution for the L_1 amount of variance explained by the PC1 and calculates the p-value by

comparing the observed L_1 to the null distribution. Usually, a p-value threshold of 0.05 is employed

to determine the significance of the gene set's activity.

In practice, it is often necessary to test many different gene sets available in large reference

databases such as KEGG [45] or MSigDB [46]. Estimating the null distribution for each set of genes

can lead to very time-consuming calculations. In the rROMA algorithm, there is an option to avoid

calculating overdispersion and shift significance scores for all gene sets using the *ApproxSample*

parameter. When this parameter is set to zero, the calculation is conducted for the null distribution

without any approximation for all gene sets. Instead, when the *ApproxSample* parameter is different

from zero, these scores can be approximated based on a predefined grid of values, which depends

on the size of the gene set under consideration. Indeed, since these two values are dependent on

the size of the gene set, it is not possible to use the same null distribution for all gene sets. To rapidly

estimate the importance of the over-dispersion and shifting scores, rROMA constructs null

distributions for a representative list of gene set sizes. These are selected to be uniformly distributed

in the logarithmic scale between the minimum and maximum size of the reference database. For a

given gene set, the null distribution that is closest in size in the log scale is then chosen.

Data availability

The data utilized in this study has been downloaded from the NCBI GEO database (GEO

accession ID GSE176121).

Code availability

The computational analyses were performed in R 4.2.2 and all the code is available under a GNU

General Public License V3 in a GitHub repository at the following url: https://github.com/sysbio-

curie/rROMA.

Acknowledgements

The authors would like to thank Pierre Gestraud for insightful discussions about the R

implementation. This work was done with financial support from ITMO Cancer Aviesan within the

framework of the 2021-2030 Cancer Control Strategy, on funds administered by Inserm (ITMO MIC

project ModPhosphoNet) and with support from the French government under management of

Agence Nationale de la Recherche as part of the "Investissements d'avenir" program, reference

ANR-19-P3IA-0001 (PRAIRIE 3IA Institute). MC was supported by Fondation pour la Recherche

Médicale (FRM), and MN was supported by Vaincre La Mucoviscidose (VLM) grant number

20190502488 and by La Fondation Dassault Systèmes.

Competing Interests

None declared.

Author Contribution Statement

M.N. and M.C. contributed equally to this work. M.N., M.C., A.Z., L.A. and L.M.: conceptualization,

methodology implementation M.N. and M.C.: data curation, data analyses, formal analysis. L.M. and

M.N.: Writing-original draft I.S.,V.S. and L.C.: scientific support, editing and content review.

References

[1] Hawkins, R. D., Hon, G. C., & Ren, B., Next-generation genomics: an integrative approach, Nature

Reviews Genetics, **11(7),** 476-486. (2010)

[2] Barillot, E., Calzone, L., Hupe, P., Vert, J. P., & Zinovyev, A., Computational systems biology of

cancer, CRC Press, (2012)

[3] Wang, K., Li, M., & Hakonarson, H., Analysing biological pathways in genome-wide association

studies. Nature Reviews Genetics, **11(12)**, 843-854, (2010)

[4] Levine, D. M. et al., Pathway and gene-set activation measurement from mRNA expression data:

the tissue distribution of human pathways. Genome biology, **7(10)**, 1-17, (2006)

[5] Schreiber, A. W., & Baumann, U., A framework for gene expression analysis, Bioinformatics,

23(2), 191-197, (2007)

[6] Puthier D., and van Helden J., Statistics for Bioinformatics - Practicals - Gene enrichment

statistics, https://dputhier.github.io/ASG/practicals/go_statistics_td/go_statistics_td_2015.html ,

(2015)

[7] Subramanian, T. et al., Gene set enrichment analysis: A knowledge-based approach for

interpreting genome-wide expression profiles, Nat Genet, **34**, 267-273, (2003)

[8] Tomfohr, J., Lu, J., & Kepler, T. B., Pathway level analysis of gene expression using singular value

decomposition. BMC bioinformatics, **6,** 1-11, (2005)

[9] Bild, A. H., Potti, A., & Nevins, J. R., Linking oncogenic pathways with therapeutic opportunities.

Nature reviews cancer, **6(9)**, 735-741, (2006)

[10] Lim, S., Lee, S., Jung, I., Rhee, S., & Kim, S., Comprehensive and critical evaluation of

individualized pathway activity measurement tools on pan-cancer data. Briefings in bioinformatics,

21(1), 36-46, (2020)

[11] Barbie DA, Tamayo P, et al., Systematic RNA interference reveals that oncogenic KRAS-driven

cancers require TBK1, Nature, **462**:108-112, (2009)

[12] Hänzelmann, S., Castelo, R., & Guinney, J., GSVA: gene set variation analysis for microarray and

RNA-seq data, BMC bioinformatics, **14**, 1-15, (2013)

[13] Vaske, C. J. et al., Inference of patient-specific pathway activities from multi-dimensional cancer

genomics data using PARADIGM, Bioinformatics, **26(12)**, i237-i245, (2010)

[14] Alvarez, M. J. et al. Network-based inference of protein activity helps functionalize the genetic

landscape of cancer, Nature genetics, **48(8)**, 838, (2016)

[15] Schubert, M. et al., Perturbation-response genes reveal signaling footprints in cancer gene

expression, Nature communications, **9(1)**, 20, (2018)

[16] Fan, J. et al., Characterizing transcriptional heterogeneity through pathway and gene set

overdispersion analysis, Nature methods, **13(3)**, 241-244, (2016)

[17] Landais, Y., & Vallot, C., Multi-modal quantification of pathway activity with MAYA, Nature

Communications, **14(1)**, 1668, (2023)

[18] Holland, C. H. et al., Robustness and applicability of transcription factor and pathway analysis

tools on single-cell RNA-seq data, Genome biology, **21**, 1-19, (2020)

[19] Zhang, Y. et al., Benchmarking algorithms for pathway activity transformation of single-cell RNA-

seq data, Computational and structural biotechnology journal, **18**, 2953-2961, (2020)

[20] Lee, E., Chuang, H. Y., Kim, J. W., Ideker, T., & Lee, D., Inferring pathway activity toward precise

disease classification, PLoS computational biology, **4(11)**, e1000217, (2008)

[21] Wagner, F., GO-PCA: an unsupervised method to explore gene expression data using prior

knowledge, PloS one, **10(11)**, e0143196, (2015)

[22] Frost, H. R., Li, Z., & Moore, J. H., Principal component gene set enrichment (PCGSE), BioData

mining, **8(1)**, 1-18, (2015)

[23] Drier, Y., Sheffer, M., & Domany, E. Pathway-based personalized analysis of cancer. Proceedings

of the National Academy of Sciences, **110(16)**, 6388-6393, (2013)

[24] Martignetti, L., Calzone, L., Bonnet, E., Barillot, E., & Zinovyev, A., ROMA: representation and

quantification of module activity from target expression data, Frontiers in genetics, **7**, 18, (2016)

[25] Golub, G. H., & Reinsch, C., Singular value decomposition and least squares solutions, Handbook

for Automatic Computation: Volume **II**: Linear Algebra, 134-151, Springer Berlin Heidelberg, (1971)

[26] Benjamini, Y., & Hochberg, Y. Controlling the false discovery rate: a practical and powerful

approach to multiple testing, Journal of the Royal statistical society: series B (Methodological), **57(1),**

289-300, (1995)

[27] Rehman, T. et al. Inflammatory cytokines TNF-α and IL-17 enhance the efficacy of cystic fibrosis

transmembrane conductance regulator modulators, The Journal of clinical investigation, **131**(16),

(2021)

[28] Liberzon, A. et al. The molecular signatures database hallmark gene set collection, Cell systems,

1(6), 417-425, (2015)

[29] Cantiello, H. Role of actin filament organization in CFTR activation, Pflügers Archiv, **443**, S75-S80,

(2001)

[30] Vasconcellos, C. A. et al. Reduction in viscosity of cystic fibrosis sputum in vitro by gelsolin,

Science, **263(5149)**, 969-971, (1994)

[31] Bucki, R. et al., Enhancement of Pulmozyme activity in purulent sputum by combination with

poly-aspartic acid or gelsolin, Journal of Cystic Fibrosis, **14(5)**, 587-593, (2015)

[32] Saint-Criq, V. et al., Choice of differentiation media significantly impacts cell lineage and

response to CFTR modulators in fully differentiated primary cultures of cystic fibrosis human airway

epithelial cells, Cells, **9(9)**, 2137, (2020)

[33] Plasschaert, L. W. et al., A single-cell atlas of the airway epithelium reveals the CFTR-rich

pulmonary ionocyte, Nature, **560(7718)**, 377-381, (2018)

[34] Okuda, K. et al. Secretory cells dominate airway CFTR expression and function in human airway

superficial epithelia, American journal of respiratory and critical care medicine, **203(10)**, 1275-1289,

(2021)

[35] Krug, K., et al., Proteogenomic landscape of breast cancer tumorigenesis and targeted therapy,

Cell, **183(5)**, 1436-1456, (2020)

[36] Parker, J. S. et al., Supervised risk predictor of breast cancer based on intrinsic subtypes, Journal

of clinical oncology, **27(8)**, 1160, (2009)

[37] Fouad, T. M., Kogawa, T., Reuben, J. M., & Ueno, N. T., The role of inflammation in inflammatory

breast cancer, Inflammation and cancer, 53-73, (2014)

[38] Sarrió, D. et al., Epithelial-mesenchymal transition in breast cancer relates to the basal-like

phenotype, Cancer research, **68(4)**, 989-997, (2008)

[39] Strandvik, B. Fatty acid metabolism in cystic fibrosis, Prostaglandins, leukotrienes and essential

fatty acids, **83(3)**, 121-129, (2010)

[40] Baglama, J., IRLBA: Fast Partial Singular Value Decomposition Method, (2016)

[41] Tsuyuzaki, K., Sato, H., Sato, K., & Nikaido, I., Benchmarking principal component analysis for

large-scale single-cell RNA-sequencing, Genome biology, **21(1)**, 1-17, (2020)

[42] Gorban, A. N., & Zinovyev, A. Y., Principal graphs and manifolds, Handbook of research on

machine learning applications and trends: algorithms, methods and techniques, 28-59, (2010)

[43] Hastie, T., Tibshirani, R., Friedman, J. H., & Friedman, J. H., The elements of statistical learning:

data mining, inference, and prediction, **2**, 1-758. New York: springer, (2009)

[44] Van Buuren, S., & Groothuis-Oudshoorn, K., Mice: Multivariate imputation by chained equations

in R. Journal of statistical software, **45**, 1-67, (2011)

[45] Kanehisa, M., The KEGG database, In Silico Simulation of Biological Processes: Novartis

Foundation Symposium Chichester, UK: John Wiley & Sons, **247**, 91-103, (2002)

[46] Liberzon, A. et al Molecular signatures database (MSigDB) 3.0. Bioinformatics, **27(12)**, 1739-

1740, (2011)

Figure 1. rROMA workflow. Schematic diagram illustrating the workflow of the rROMA algorithm.

Figure 2. Representation of shifted and overdispersed gene sets. Representation of gene sets in the

case of two samples. Each dot represents one gene, its horizontal (resp. vertical) value

corresponding to its expression in sample 1 (resp. sample 2). Genes associated with latent Factor A

are plotted in blue and the corresponding PC1 direction is plotted in red (a). This example

corresponds to a shifted pathway, as assessed by a median of gene projections onto PC1 direction

far from the origin of the distribution. Genes associated with latent Factor B are plotted in green (b)

and the corresponding PC1 direction is plotted in red. This example corresponds to an overdispersed

pathway, as the PC1 is well aligned with the dots' distribution. Genes in yellow are neither

overdispersed nor shifted, as PC1 explains a relatively small fraction of variance (not represented on

the figure) and the median of projections onto PC1 is close to the origin for this group of genes.

Figure 3. Active pathways in Cystic Fibrosis. Heatmap of activity scores for gene sets identified as

significantly shifted (a) or significantly overdispersed (b) in GSE176121 dataset. Samples are in

columns, gene sets are in rows. Horizontal sidebar color encodes true class labels.

Figure 4. Gene weights for the COAGULATION pathway. Plots illustrating the contribution of genes

to the COAGULATION gene set activity score. The weights in panel (a) indicate the gene projections

on PC1, limited to the genes that have the greatest contribution to the observed variation in the

COAGULATION gene set. In panel (b) the genes of the COAGULATION gene set are represented in the

PCA space. Red dots are genes from the gene set, blue dots show randomly selected genes used to

generate a null distribution.

Figure 5. Estimated cell type abundances from CF transcriptomic data. Heatmap of rROMA scores

obtained for Plasschaert (A) and Okuda (B) signatures of cell types in the Saint-Criq RNA seq dataset.

Samples are in columns, gene sets corresponding to cell types are in rows. Horizontal sidebar color

encodes true class labels.

685 **Figure 6. Functional analysis of proteomic data.** Clustering of pathway activity level matrix for the

686 rROMA significantly active pathways. The color bars indicate the cluster assignment based on

687 rROMA pathway activity profiles (top) and the PAM50 subtype classification (bottom).