Title: Intra-annual temperature variability and nutrient enrichment drive seasonal β-diversity in

global grasslands

Authors: Magda Garbowski[1,2], Elizabeth Boughton[3], Anne Ebeling[4], Philip Fay[5], Yann Hautier[6],

Hanna Holz[7], Anke Jentsch[8], Stephanie Jurburg[1,2], Emma Ladouceur[1,2,9], Jason Martina[10],

Timothy Ohlert[11,12], Xavier Raynaud[13], Christiane Roscher[1,2], Grégory Sonnier[3], Pedro

Maximiliano Tognetti[14], Laura Yahdjian[14], Peter Wilfahrt[15], Stan Harpole[1,2,7]

* Corresponding author email: Magda.Garbowski@uwyo.edu

1) German Centre for Integrative Biodiversity Research (iDiv) Leipzig, Germany

2) Department of Physiological Diversity, Helmholtz Centre for Environmental Research - UFZ

Leipzig, Germany

3) Archbold Biological Station, Venus, Florida, USA

4) University Jena, Institute of Ecology and Evolution, Jena, Germany

5) USDA-ARS Grassland Soil and Water Lab, Temple, Texas, USA

6) Ecology and Biodiversity Group, Department of Biology, Utrecht University, Utrecht, The

Netherlands

7) Martin Luther University Halle-Wittenberg, Halle (Saale), Germany

8) Department of Disturbance Ecology and Vegetation Dynamics, Bayreuth Center of Ecology

and Environmental Research (BayCEER), University of Bayreuth, Bayreuth, Germany

9) Department of Biology, University of Leipzig, Germany

10) Department of Biology, Texas State University, San Marcos, Texas 78666, USA

11) Department of Biology, University of New Mexico, Albuquerque, New Mexico, USA

12) Department of Biology, Colorado State University, Fort Collins, Colorado, USA

13) Sorbonne Université, Université Paris-Cité, UPEC, IRD, CNRS, INRA, Institute of Ecology

and Environmental Sciences, iEES Paris, Paris, France

14) IFEVA, Universidad de Buenos Aires, CONICET, Facultad de Agronomía, Catedra de

Ecología, Buenos Aires, Argentina

15) Department of Ecology, Evolution and Behavior, University of Minnesota, St Paul,

Minnesota, USA

* Corresponding author's present address: Botany Department, 1000 E. University Ave.

University of Wyoming, Laramie, WY 82071

Open Research Statement: Estimates of observed, expected, and null seasonal temporal

diversity used in these analyses as well as the novel code used to generate these estimates can be

found at: https://github.com/MagdaGarbowski/Nutrient_Network_Seasonality. Upon acceptance,

these data and finalized code will be publicly available and archived on figshare.

Abstract

In many grasslands, species with specific traits occupy unique temporal positions within communities. Such intra-annual segregation is predicted to be greatest in systems with high intra-annual climate variability because fluctuating environmental conditions provide opportunities for temporal niche partitioning among species. However, because most studies on intra-annual community dynamics have been conducted at individual sites, relationships between intra-annual climate variability and seasonal community dynamics at global scales have not yet been identified. Furthermore, the same characteristics that promote species-specific responses to fluctuations in environmental conditions may also drive species-specific responses to global change drivers such as eutrophication. Research provides evidence that eutrophication alters inter-annual plant community dynamics yet understanding of how it alters intra-annual dynamics remains limited.

We used early-season and late-season compositional data collected from 10 grassland sites around the world to ask how intra-annual variability in precipitation and temperature as well as nutrient enrichment shape intra-annual species segregation, or seasonal β-diversity, in plant communities. We also assessed whether changes in the abundances of specific functional groups including annual forbs, perennial forbs, C3 and C4 graminoids, and legumes underpin compositional differences between early- and late-season communities and treatments. We found that intra-annual temperature variability and seasonal β-diversity were positively related but observed no relationship between intra-annual precipitation variability and seasonal β-diversity. This suggests that positive relationships between α-diversity and intra-annual temperature variability identified in earlier studies may be underpinned by the positive influence of intra-

annual temperature variability on temporal segregation of species within growing seasons. We

found that nutrient enrichment increased seasonal β-diversity via increased turnover of species

between early- and late-season communities. This finding mirrors patterns observed at inter-

annual scales and suggests fertilization can alter compositional dynamics via similar mechanisms

at varied temporal scales. Finally, fertilization reduced the abundance of C4 graminoids and

legumes and eliminated intra-annual differences in these groups. In contrast, fertilization resulted

in intra-annual differences in C3 graminoids which were not observed in control conditions, and

increased abundance of C3 graminoids and annual forbs overall. Our study provides new insight

into how intra-annual climate variability and nutrient enrichment influence biodiversity and

seasonal dynamics in global grasslands.

Keywords: plant community diversity and structure, global change ecology, grasslands, intra-

annual climate variability, nutrient enrichment, Nutrient Network, temporal beta-diversity,

temporal niche segregation

Introduction

Ecologists have investigated the mechanisms by which climate shapes plant diversity for decades (e.g., Rosenzweig 1995, Hawkins et al. 2003, Harrison et al. 2020). While most research examines associations with mean climate conditions, numerous theoretical (e.g., Lewontin & Cohen 1969, Chesson 2000, Adler & Drake 2008) and empirical (e.g., Levine & Reese 2004, Letten et al. 2013) studies have highlighted the importance of temporal climate variability in structuring populations and communities. Central to these studies is the notion that environmental fluctuations provide opportunities for temporal niche partitioning among species (e.g., Chesson &Huntly 1997, Chesson 2000). Intra-annual fluctuations in temperature and precipitation may be particularly important drivers of intra-annual temporal niche partitioning in plant communities as they influence plant physiology and phenology and modulate ecosystem processes related to water availability and nutrient cycling (Luo et al. 2020). Modeling studies have demonstrated that intra-annual variation in precipitation supports coexistence among plant species with slight ecological differences in germination requirements, growing season length, or seasonal growth activity (Mathias and Chesson 2013). Empirical evidence documenting the influence of intra-annual climate variability on temporal niche segregation mainly comes from studies of desert systems in which winter and summer annuals segregate temporally in response to seasonal fluctuations in precipitation (e.g., Mulroy & Rundel 1977, Guo & Brown 1997). But temporal segregation of coexisting species is evident in diverse ecosystems including tropical forests (Sapijanskas et al. 2014) and grasslands (Fargione & Tilman 2005).

In many grasslands, species with specific strategies optimally utilize resources under different environmental conditions and thus coexist by occupying unique temporal positions within communities. For example, in grasslands co-dominated by C3 and C4 grasses, C3 grasses

grow and set seed primarily in early to mid-season and C4 grasses grow and set seed primarily in

late season. Similar dynamics occur in communities without C3 and C4 grasses present. For

instance, in temperate European grasslands early-season grasses and forbs contribute most to

productivity at the beginning of the season, whereas late-season grasses contribute most to

productivity late in the growing season (e.g., Guimarães-Steinicke et al. 2019, Doležal et al.

2017). However, the same characteristics that promote species-specific responses to fluctuations

in environmental conditions often drive species-specific responses to global change drivers such

as eutrophication. For example, subordinate forbs and legumes that thrive at different time

periods from dominant grasses are often reduced in fertilized conditions in European meadows

(e.g., Doležal et al. 2017). Research from the last several years has elucidated how

eutrophication alters inter-annual plant community dynamics (e.g., Hautier et al. 2014, Chen et

al. 2021). Yet, interactions between global change drivers and intra-annual dynamics of plant

communities remain largely unexplored (White & Hastings 2020).

Eutrophication in grasslands leads to reduced plant species richness and diversity (e.g.,

Hautier et al. 2009, Harpole et al. 2016). Evidence for these patterns has primarily been gathered

from studies that measure plant community composition once a year, usually at the peak of

biomass production (e.g., Borer et al. 2014). However, research has demonstrated that diverse

resource-use strategies allow distinct species assemblages to dominate in different points within

a growing season (Doležal et al. 2017, Guimarães-Steinicke et al. 2019, Huang et al. 2019). In

grassland systems, nutrient enrichment may promote the dominance of species with specific

traits for longer periods of time and reduce overall dissimilarity among early- and late-season

communities (Doležal et al. 2017). Further, the addition of multiple limiting nutrients may

reduce the importance of species-specific trade-offs associated with competition for particular

nutrients (Harpole et al. 2016) that are most limiting at different points throughout the growing

season (Klaus et al. 2016). Assessing how nutrient enrichment affects compositional

dissimilarity between early and late-season assemblages would clarify whether patterns of

reduced diversity in response to fertilization at inter-annual scales are underpinned by reduced

dissimilarity at intra-annual scales.

Changes in overall diversity in responses to nutrient enrichment often result from changes

in the abundance of species that are affected by nutrient enrichment in different ways. Nitrogen

enrichment reduces the abundance of legumes and nutrient conservative C4 grasses but increases

the abundance of nutrient acquisitive C3 grasses (e.g., Suding et al. 2005, Tognetti et al. 2021).

Fertilization with multiple nutrients can further alter community composition by causing shifts in

the relative abundance of species from specific functional groups (Wilcots et al. 2021). When

nutrient enriched communities become dominated by specific species or functional groups,

ecosystem stability may be reduced via a loss of asynchronous species responses to

environmental fluctuations (Hector et al. 2010, Hautier et al. 2014). In addition, nutrient

additions that result in the loss of forb and legume species can have cascading effects on

organisms at other trophic levels, such as pollinators, that often depend on these groups for floral

resources at specific times of the year (Burkle & Irwin et al. 2010, Dyer et al. 2021).

Understanding how nutrient enrichment affects the abundances of specific functional groups at

specific timepoints in the growing season (i.e., early vs late) and between timepoints is essential

to identifying when compositional changes take place and which species drive them.

In this study, we used above-ground species composition data collected early and late in the

growing season from 10 grassland sites around the world to assess how intra-annual variability in

temperature and precipitation as well as eutrophication influence early vs. late season

dissimilarly (seasonal β-diversity hereafter) of plant assemblages. The 10 sites are part of the

Nutrient Network – a globally replicated experiment in which herbaceous plant communities are

supplemented with fertilizer. Data from each site spanned between 4 and 11 years. We used data

from untreated plots to assess relationships between intra-annual precipitation and temperature

variability (estimates obtained from WorldClim; Fick & Hijmans 2017) and seasonal β-diversity

across our study sites. We then examined the effects of fertilization on seasonal β-diversity and

its components of nestedness and replacement. Finally, to determine which species underpin

changes in seasonal β-diversity in response to fertilization, we assessed overall changes in

abundance of different functional groups (i.e., C3 graminoids, C4 graminoids, annual forbs,

perennial forbs, legumes) between early and late timepoints and fertilization treatments. Our

hypotheses were that:

H1. Sites with high intra-annual variability in temperature and precipitation would have high

seasonal β-diversity.

H2. Fertilization would decrease seasonal β-diversity because a subset of species would

dominate across early and late sampling timepoints (i.e., fertilized plots would have

higher nestedness).

H3. Differences in early-season vs. late-season community composition in response to

fertilization would be underpinned by increased abundances of resource acquisitive

species (e.g., C3 graminoids, annual forbs) and decreased abundances of species with

more conservative strategies (e.g., C4 graminoids, perennial forbs, legumes) within and

between specific timepoints (i.e., early vs. late)

Methods

Study design and site locations

Data for this study were collected from 10 grassland sites from around the world that are

part of the Nutrient Network – a globally distributed experiment in which plant communities are

supplemented with factorial combinations of nitrogen (N), phosphorous (P), potassium (K) and

micronutrients (μ) (Borer et al. 2014). While most Nutrient Network sites collect compositional

data once a year, data from this subset of sites is collected twice each growing season ("early"

and "late" hereafter). These sites are distributed across five continents, in Africa (ukul.za),

Australia (burrawan.au), Europe (bayr.de, cereep.fr, frue.ch, jena.de), North America (arch.us,

temple.us, sevi.us) and South America (chilcas.ar) (Table S1).

Compositional data used in this study were collected from 3 to 5 replicated control and

NPKμ plots at these sites from between 4 (bayr.de) and 11 years (ukul.za) (mean length of

experiment is 6.5 years, details in Table S1). Fertilized plots at all sites besides cereep.fr received

10 g m^{-2} of N, P, and K annually with a one-time addition of micronutrients; fertilized plots at

the cereep.fr site received 2.5 g m^{-2} annually with a one-time addition of micronutrients. Despite

this different application rate, all results were qualitatively similar whether the cereep.fr site was

or was not included in analyses, so we retain it in the results presented here. For additional

details about experimental design, please see Borer et al. (2014).

Quantifying intra-annual temperature and precipitation variability

Estimates of intra-annual temperature and precipitation variability for each site were

obtained from WorldClim (Fick and Hijmans 2017). Intra-annual temperature and precipitation

variability for each site are based on monthly totals of precipitation and temperature where

"Temperature variability" is the standard deviation of monthly temperature * 100 and

"Precipitation variability" is the coefficient of variation of monthly precipitation totals for a

given year. In both cases, higher values indicate higher variance or greater fluctuations in

temperature and precipitation within a year. We chose to use annual variability values rather than

growing season values because winter precipitation at some of our sites (e.g., sevi.us) is an

important driver of plant community dynamics. Between 18 and 30 years of precipitation and

temperature data were used to obtain average intra-annual variability values for each site based

on data availability in WorldClim.

Quantifying seasonal β-diversity

Dissimilarity indices are useful tools for measuring differences between communities in

space or time. However, dissimilarity indices can be influenced by local community size (i.e., α-

diversity) and overall richness of species at regional scales (i.e., γ-diversity) (e.g., Chase &

Myers 2011). Deviations from null expectations of dissimilarity (e.g., z-scores) can help

determine whether compositional differences, independent of local community size and regional

species pools, drive dissimilarity patterns (e.g., Chase & Myers 2011). More specifically, larger

z-scores indicate higher dissimilarity between communities, whereas lower deviations indicate

lower dissimilarity.

To quantify dissimilarity between early and late communities in control and NPKμ

treatments at each site we used deviations from expected values of the Bray-Curtis dissimilarity

index (Figure 1). We did this by first permuting species abundances (i.e., absolute values of

species cover) from permanent m^2 survey plots 100 times while holding overall richness and total

abundances of species constant within a plot at a given sampling timepoint (i.e., early or late

season). We used 100 permutations for "early" communities and 100 permutations for "late"

communities to obtain mean and standard deviation values for expected (i.e., null) dissimilarity

in each plot. We then calculated z-scores for each early-to-late comparison for each plot, year,

and treatment combination using the following formula:

$Seasonal\ \beta-diversity_{(z-score)} = \dfrac{BC_{observed} - \mu_{null}}{\sigma_{null}}$

where $BC_{observed}$ is the observed Bray-Curtis value from each plot in each year, μ_{null} is the

expected mean for each plot in each year obtained from the distribution of permutation

dissimilarly estimates, and σ_{null} is the variance for each plot in each year obtained from the

distribution of permutation dissimilarly estimates (Figure 1b). We used these z-scores as our

estimates of seasonal β-diversity from each plot in each year. Comparisons from a given plot in a

given year with less than four species at either the early or late sampling time point were

removed prior to all analyses because communities with less than four species could not produce

sufficient permutations to calculate reliable estimates of seasonal β-diversity. This resulted in the

exclusion of 53 early-to-late comparisons from 888 comparisons total (i.e., 835 comparisons

were included in analyses).

We also decomposed observed Bray-Curtis dissimilarity from each early-to-late comparison

for each plot, year, and treatment into components of species "replacement" and "nestedness"

(Baselga 2009). Nestedness captures the component of dissimilarity that results from one

community being a subset of another community, whereas replacement captures the component

of dissimilarity that results from certain species being lost and others gained between compared

communities. We used the beta.part package (Baselga & Orme 2012) to obtain all dissimilarity

estimates.

Statistical models

We used two separate linear regression models to assess relationships between intra-

annual temperature and precipitation variability and seasonal β-diversity of plant communities.

In these models, average intra-annual temperature variability or precipitation variability for each

site were included as a continuous predictor and average seasonal β-diversity estimates from

control treatments from each site were included as a continuous response. We examined how

NPKμ treatment influenced seasonal β-diversity using a multilevel regression model with plot-

level seasonal β-diversity values included as response variables, treatment as a predictor, and a

random effect of block, nested within treatment year, nested within site. Because not all

functional groups are present at all sites, functional group analyses were completed using data

from sites that had specific functional groups present at >1% cover overall. We fit five separate

multilevel regression models to quantify how abundance of specific functional groups (i.e.,

summed cover of all species belonging to annual forbs, perennial forbs, C3 grasses, C4 grasses

and legumes from m^2 plots) differed among early and late sampling timepoints and treatments.

These models included treatment by season combinations as categorical predictors (i.e.,

control_early, control_late, NPKμ_early, NPKμ_late) and block nested within treatment year

nested within site as a random effect (i.e., interactions were not assessed prior to investigating

differences in means).

For Bayesian inferences and estimates of uncertainty, all models described were fitted

using the Hamiltonian Monte Carlo (HMC) sampler using Stan (Carpenter et al. 2017) and coded

using the 'brms' package (Bürkner et al. 2018) in R (version V.2.1 R Core Development Team).

All models were fit with 4 chains and 3000 iterations (Table S2). We used default priors for all

models in which dissimilarly estimates (i.e., seasonal β-diversity, nestedness, turnover) were

included as response variables. In models assessing differences in functional group abundances

between timepoints and treatments, default priors were used for C3 and C4 graminoid and

perennial forb models, whereas weakly regularizing priors were used in models assessing

differences in legume and annual forb abundances (Table S2). We inspected the HMC chains to

assess model convergence. For all models, we estimated the significance of effects by computing

the difference between posterior distributions of interest and assessing whether the 90% credible

interval of the difference contained zero.

Results

We found a positive relationship between intra-annual temperature variability and

seasonal β-diversity in unfertilized control plots (Slope: 0.0028, 90% Credible Interval (CI):

0.0004 to 0.0050) (Figure 2a) and no relationship between intra-annual precipitation variability

and seasonal β-diversity (Slope: -0.0052, CI: -0.0278 to 0.0166) (Figure 2b). Contrary to our

hypothesis, we detected a significant positive effect of fertilization on seasonal β-diversity across

sites (Figure 3). This result indicates greater dissimilarity between early and late season

communities in fertilized conditions, even after accounting for differences in richness between

the two treatments (average number of species per m^2 in control: 12.8; average number of species

per m^2 in NPKμ: 11.5). However, seasonal β-diversity values were negative in most cases,

indicating lower than expected dissimilarity between early-season and late-season communities.

After decomposing observed seasonal β-diversity into components of nestedness and turnover,

we found no differences between treatments in nestedness (Figure 4a) but higher turnover in

fertilized conditions than in controls (Figure 4b).

The effects of fertilization varied by functional group and between early and late

sampling timepoints (Figure 5). At both sampling timepoints, fertilization resulted in higher

cover of annual forbs (Figure 5a) and C3 graminoids (Figure 5c) and lower cover of C4

graminoids (Fig. 5d) and legumes (Fig. 5e). Perennial forb cover was reduced with fertilization,

but only at the late sampling timepoint (Fig. 5b). Patterns of functional group abundances

between early and late sampling points also varied between control and fertilized treatments. In

control but not fertilized treatments, C4 graminoid abundance was higher at the later sampling

point compared to early in the season and legume abundance was higher early in the season

compared to late in the season. In contrast, abundance of C3 graminoids was higher early in the

season compared to late in the season but only in fertilized treatments.

Discussion

We found higher seasonal β-diversity in grasslands with high intra-annual temperature

variability but observed no relationship between intra-annual precipitation variability and

seasonal β-diversity. These finding suggests that positive relationships between α-diversity and

intra-annual temperature variability identified in earlier studies (e.g., Scheiner & Rey-Benayas

1994, Boonman et al. 2021) may be underpinned by the positive influence of intra-annual

temperature variability on temporal segregation of species within seasons. However, our findings

also contradict several studies that implicate intra-annual precipitation variability as a key driver

of seasonal β-diversity (e.g., Mulroy & Rundel 1977, Mathias & Chesson 2013). We suspect

these contradictions arises due to differences in annual systems from which most evidence

relating intra-annual precipitation variation to seasonal community dynamics has been gathered,

and the predominantly perennial systems in our study. In annual systems, population,

community, and ecosystem dynamics are often governed by interactions between intra-annual

precipitation patterns and demographic traits related to dormancy and germination (e.g., Levine

et al. 2011, Kimball et al. 2011, Shaw et al. 2022). In addition, in systems co-dominated by

annuals and perennials, intra-annual variability in precipitation is often necessary for the

persistence of annuals that occupy unique temporal niches in communities (e.g., Pérez-Camacho

et al. 2012). On the other hand, intra-annual temperature variability may be a more important

driver of intra-annual dynamics in perennial systems because temperature regulates plant

physiology and shapes temporal segregation among many perennial plant species (e.g., Kemp &

Williams. 1980, Monson et al. 1983). Given the low abundance of annual cover across our study

sites (*c.* 11%), we are not able to test these patterns robustly. But a positive, non-significant,

relationship between annual seasonal β-diversity and intra-annual precipitation variability

(Supplementary Figure 1a) and a negative, non-significant, relationship between perennial

seasonal β-diversity and intra-annual precipitation variability (Supplementary Figure 1b) invite

new questions about how intra-annual precipitation variability influences seasonal community

dynamics across diverse ecosystems.

The negative impacts of nutrient enrichment on grassland diversity have been extensively

documented (e.g., Borer et al. 2017) and recent studies have identified a positive effect of

fertilization on temporal β-diversity at inter-annual scales (Koerner et al. 2016, Hodapp et al.

2018, Chen et al. 2021). Mirroring these results at intra-annual scales, we found that fertilization

resulted in higher seasonal β-diversity. Because we used deviations from null expectations as our

measure of seasonal β-diversity, we can be confident that these shifts arise from true changes in

composition rather than random processes influenced by richness. Similar to results found at

inter-annual scales by Chen et al. 2021, we found that higher temporal β-diversity with

fertilization was driven by higher turnover of species, and not nestedness of species, between

early and late-season communities. This suggests that fertilization increases temporal β-diversity

among and within years by similar mechanisms, namely by unique species occupying space

within the community at different times (i.e., turnover) rather than species being lost from one

timepoint to another (i.e., nestedness). At inter-annual scales, increased temporal β-diversity and

species turnover through time can result in reduced stability of ecosystem productivity (e.g.,

Koerner et al. 2016, Chen et al. 2021). Additional research focused on how nutrient enrichment

alters composition and associated ecosystem functions within growing seasons would clarify

whether similar patterns manifest at intra-annual scales.

As expected, fertilization resulted in higher abundance of resource acquisitive species

(i.e., C3 graminoids, annual forbs) and lower abundance of resource conservative species (i.e.,

C4 graminoids, legumes, perennial forbs) within and across sampling timepoints. Numerous

other studies have documented similar patterns at single timepoints (e.g., Suding et al. 2005,

Isabell et al. 2013, Wilcots et al. 2021, Tognetti et al. 2021) and results are primarily attributed to

physiological differences between these groups and concomitant shifts from below-ground

competition for nutrients to aboveground competition for light (e.g., Borer et al. 2014, Harpole et

al. 2017). Among other anatomical and physiological differences, compared to C4 species, C3

species (i.e., most of the annuals in our study and C3 graminoids) generally have lower C:N

ratios (e.g., Wedin & Tilman 1990), lower water and photosynthetic nitrogen use efficiencies

(Black 1971, Taylor et al. 2011, Ripley et al. 2010), and earlier phenologies due to lower optimal

temperatures for photosynthesis (Kemp & Williams 1980). These characteristics can lead to

higher sensitivities of resource acquisitive species to nutrient limitation which allow them to

increase in fertilized conditions (e.g., Zhong et al. 2019).

Fertilization made apparent seasonal differences in C3 graminoid abundance that were

not present under control, while eliminating seasonal differences in abundance of C4 graminoids

and legumes that were present in control treatments. Higher C3 graminoid abundance early in the

season in fertilized treatments could reflect a higher capacity for growth of these species early in

the season, when resources are abundant, followed by more pronounced declines later in the

season when resources are scarce (e.g., Yuan et al. 2007). In contrast, both C4 graminoids and

legumes are adapted to resource-limited conditions which can vary throughout growing seasons

(Klaus et al. 2014). By altering resource conditions, fertilization likely modulates the low-

resource periods during which these species thrive resulting in lower abundances of these species

overall and homogenization of these groups across the growing season.

Our understanding of how grasslands respond to various components of global change is

primarily based on studies that document community shifts at inter-annual scales. However,

global changes may also disrupt within-season community dynamics. Here we show that

seasonal β-diversity across 10 global grasslands is shaped by intra-annual temperature variability

and eutrophication. Given that both intra-annual temperature variability (e.g., Xu et al. 2013) and

precipitation variability (e.g., Hajek & Knapp 2021) are expected to increase in coming decades

understanding effects of these understudied components of global change is essential to

understanding how plant communities will shift into the future. Investigations focused on

specific components of climatic variability such as seasonality (i.e., the occurrence of certain

events within a definite limited period, *sensu* Lieth 1974) or predictability (i.e., the regularity of

recurrence of the within cycle distribution of events, *sensu* Tonkin 2017) would further

illuminate how fluctuations in climate shape seasonal dynamics of plant communities. We also show that nutrient enrichment increases seasonal β-diversity and species turnover, enhances the abundance of resource-acquisitive species (i.e., annual forbs, C3 graminoids) within and between timepoints, and decreases abundances of resource-conservative species (i.e., C4 graminoids, legumes) within and between timepoints. In addition, our results suggest that fertilization results in homogenization of abundances of resource conservative species early to late in the growing season. If the effects of intra-annual community change mirror those at inter-annual scales, the community shifts we observed in our study could have cascading impacts on ecosystem stability, multifunctionality, and the ability of these systems to recover from future perturbations. Our study provides new insight into the mechanisms by which climate variability and nutrient enrichment shape within-season community dynamics in global grasslands and highlights how discerning these patterns is essential to our understanding of biodiversity in these valuable ecosystems.

Acknowledgements: Funding for this project was provided by the German Science Foundation (DFG; via University Leipzig/iDiv) for the project "Strategisches Projekt: NEu – Nutrient Network Europe" (Projektnummer 346001057-03). The authors do not have any conflicts of interest to declare.

Author contributions: MG and SH conceptualized this project, MG completed analyses with assistance from HH, SJ, and EL, MG wrote the manuscript, all authors contributed to data collection and editing the manuscript.

References:

Adler, P.B. & Drake, J.M. (2008) Environmental variation, stochastic extinction, and competitive coexistence. *American Naturalist.* **172**, (5) 186 –195

Baselga, A. (2009) Partitioning the turnover and nestedness components of beta-diversity. *Global Ecology and Biogeography*. **19** (1) 134 - 143

Baselga, A. & Orme, D.L. (2012) betapart: an R package for the study of beta diversity. *Methods in Ecology and Evolution* **3**, 808-812

Borer, E. T. et al. (2014) Herbivores and nutrients control grassland plant diversity via light limitation. *Nature* **508**, 517–520

Borer, E.T., Grace, J.B., Harpole, W.S., MacDougall, A.S., Seabloom, E.W. (2017) A decade of insights into grassland ecosystem responses to global environmental change. *Nature Ecology & Evolution* **1**, 0118

Burkle L.A. & Irwin, R.E. (2010) Beyond biomass: measuring the effects of community-level nitrogen enrichment on floral traits, pollinator visitation and plant reproduction. *Journal of Ecology*. **98** (3), 705 – 717

Bürkner P.C. (2017) "brms: An R Package for Bayesian Multilevel Models using Stan." Journal of Statistical Software, 80(1), 1–28

Carpenter B., Gelman A., Hoffman M.D., et al. (2017) Stan: a probabilistic programing language. *Journal of Statistical Software*. **76** (1)

Chase, J. M. & Myers, J. A. (2011) Disentangling the importance of ecological niches from stochastic processes across scales. *Philosophical Transactions Royal Soc B Biological Sci* **366**, 2351–2363

Chen, Q. et al. (2022) Nutrients and herbivores impact grassland stability across spatial scales through different pathways. *Global Change Biol* **28**, 2678–2688

Chesson, P., (2000) Mechanisms of maintenance of species diversity. *Annual Reviews of Ecology and Systematics* **31**, 343-366

Chesson, P. & Huntly N. (1997) The roles of harsh and fluctuating conditions in the dynamics of ecological communities. *American Naturalist* **150**, 519–553

Doležal, J., Lanta, V., Mudrák, O. & Lepš, J. (2019) Seasonality promotes grassland diversity: Interactions with mowing, fertilization and removal of dominant species. *J Ecol* **107**, 203–215

Dyer, A.G., Jentsch, A., Burd, M., Garcia, J.E., Giejsztowt, J., Camargo, M.G.G., Tjørve, E., Tjørve, K.M.C., White, P.S., Shrestha, M. (2021) Fragmentary blue: Resolving the rarity paradox in flower colours. *Frontiers in Plant Science* 11: Article 618203

Fargione, J. & Tilman, D. (2005) Niche differences in phenology and rooting depth promote coexistence with a dominant C4 bunchgrass. *Oecologia* **143**, 598–606

Fick S.E. & Hijmans, R.J. (2017) WorldClim 2: new 1-km spatial resolution climate surfaces for global land areas. *International Journal of Climatology.* **37**, 4302 – 4315

Guimarães-Steinicke, C., Weigelt, A., Ebling, A., Eisenhauer, N., Duque-Lazo, L., Reu, B., Roscher, C., Schumacher, J., Wagg, C., Wirth, C. (2019) Terrestrial laser scanning reveals temporal changes in biodiversity mechanisms driving grassland productivity. *Advances in Ecological Research* **61**, 133-161

Hajek, O. L. & Knapp, A. K. (2022) Shifting seasonal patterns of water availability: ecosystem responses to an unappreciated dimension of climate change. *New Phytol* **233**, 119–125

Harpole, W. S. et al. (2016) Addition of multiple limiting resources reduces grassland diversity. *Nature* 1–9

Harpole, W. S. et al. (2017) Out of the shadows: multiple nutrient limitations drive relationships among biomass, light and plant diversity. *Functional Ecology* **31**, 1839–1846

Hautier, Y., Niklaus, P. A. & Hector, A. (2009) Competition for Light Causes Plant Biodiversity Loss After Eutrophication. *Science* **324**, 636–638

Hautier, Y., Seabloom, E., Borer, E., et al. (2014) Eutrophication weakens stabilizing effects of diversity in natural grasslands. *Nature* **508**, 521 – 525

Harrison, S., Spasojevic M.J., Li D. (2020) Climate and plant community diversity in space and time. *Proc National Acad Sci* **117** (9), 4464 – 4470

Hawkins, B.A., Field, R., Cornell, D.J., et al. (2003) Energy, water, and broad-scale geographic patterns in species richness. *Ecology* **84** (12), 3105-3117

Hector, A. Hautier, Y., Wacker S.L., et al. (2010) General stabilizing effects of plant diversity on grassland productivity through population asynchrony and overyielding. *Ecology* **91** (8), 2213 – 2220

Hodapp, D. (2018) *et al.* Spatial heterogeneity in species composition constrains plant community responses to herbivory and fertilisation. *Ecology Letters* **21**, 1364–1371

Huang, L., Xue, W. & Herben, T. (2019) Temporal niche differentiation among species changes with habitat productivity and light conditions. *Journal of Vegetation Science* **30**, 438–447

Guo, Q., & Brown, J.H. (1997) Interactions between winter and summer annuals in the Chihuahuan desert. *Oecologia* 111:123–128.

Isabell, F. *et al.* (2013) Nutrient enrichment, biodiversity loss, and consequent declines in ecosystem productivity. *Proc National Acad Sci* **110**, 11911–11916

Kemp, P. R. & Williams, G. J. A (1980) Physiological Basis for Niche Separation Between *Agropyron Smithii* (C3) and *Bouteloua Gracilis* (C4). *Ecology* **61**, 846–858

Klaus, V.H. et al. (2016) Temporal and small-scale spatial variation in grassland productivity, biomass quality, and nutrient limitation. *Plant Ecology*. **217**, 843 – 856

Koerner, S. E. et al. (2016) Nutrient additions cause divergence of tallgrass prairie plant communities resulting in loss of ecosystem stability. *J Ecol* **104**, 1478–1487

Letten, A.D., Ashcroft, M.B., Keith, D.A., Gollan J.R., Ramp D. (2013) The importance of temporal climate variability for spatial patterns in plant diversity. *Ecography* **36**, (12) 1341-1349

Levine, J. M., & M. Rees. (2004) Effects of temporal variability on rare plant persistence in annual systems. *American Naturalist* **164**, 350–363

Lewontin, R.C., & Cohen, D. (1969) On population growth in a randomly varying environment. *Proc. National Acad Sci.* **62** (4), 1056-1060

Lieth, H. (1974) Phenology and seasonality modeling. Springer-Verlag, New York, New York, USA.

Luo, Y. *et al.* (2020) Nutrients and water availability constrain the seasonality of vegetation activity in a Mediterranean ecosystem. *Global Change Biol*.

Mathias, A. & Chesson, P. (2013)Coexistence and evolutionary dynamics mediated by seasonal environmental variation in annual plant communities. *Theor Popul Biol* **84**, 56–71

Monson, R. K., Littlejohn, R. O. & Williams, G. J. (1983) Photosynthetic adaptation to temperature in four species from the Colorado shortgrass steppe: a physiological model for coexistence. *Oecologia* **58**, 43–51

Mulroy, T. W. & Rundel, P. W. (1977)Annual Plants: Adaptations to Desert Environments. *Bioscience* **27**, 109–114

Pérez-Camacho L., Rebello, S., Hernández-Santana, V., García-Salgado, J., Pavón-García, J., Gómez-Sal., A. (2012) Plant functional trait responses to interannual rainfall variability, summer drought and seasonal grazing in Mediterranean herbaceous communities. *Functional Ecology* **26**, 740 – 749

Ripley, B., Frole, K. & Gilbert, M. (2010) Differences in drought sensitivities and photosynthetic
limitations between co-occurring C3 and C4 (NADP-ME) Panicoid grasses. *Ann Bot-london* **105**,
493–503

Rosenzweig, M.L., (1995) Species Diversity In Space and Time, Cambridge University Press,
Cambridge, UK., 436 pp.

Sapijanskas, J., Paquette, A., Potvin, C., Kunert, N. & Loreau, M. (2014) Tropical tree diversity
enhances light capture through crown plasticity and spatial and temporal niche differences.
Ecology **95**, 2479–2492

Suding, K. N. *et al.* (2005) Functional- and abundance-based mechanisms explain diversity loss
due to N fertilization. *Proceedings of the National Academy of Sciences* **102**, 4387–4392

Taylor, S. H. *et al.* (2014) Physiological advantages of C4 grasses in the field: a comparative
experiment demonstrating the importance of drought. *Global Change Biol* **20**, 1992–2003

Tognetti, P. M. *et al.* (2021) Negative effects of nitrogen override positive effects of phosphorus
on grassland legumes worldwide. *Proc National Acad Sci* **118**
Tonkin, J. D., Bogan, M. T., Bonada, N., Rios-Touma, B. & Lytle, D. A. (2017) Seasonality and
predictability shape temporal species diversity. *Ecology* **98**, 1201–1216

Wedin, D. A. & Tilman, D. (1990) Species effects on nitrogen cycling: a test with perennial
grasses. *Oecologia* **84**, 433–441

White, E. R. & Hastings, A. (2020) Seasonality in ecology: Progress and prospects in theory.
Ecol Complex **44**, 100867

Wilcots, M. E., Harpole, W. S., Seabloom, E. W. & Borer, E. T. (2021) Community change can
buffer chronic nitrogen impacts, but multiple nutrients tip the scale. *Ecology* **102**

Xu, L., Myneni R.B., Chapin F.S., et al. (2013) Temperature and vegetation seasonality
diminishment over northern lands. *Nature.* **3**, 581 – 586

Yuan, Z., Liu, W., Niu, S. & Wan, S. (2007) Plant Nitrogen Dynamics and Nitrogen-use
Strategies under Altered Nitrogen Seasonality and Competition. *Ann Bot-london* **100**, 821–830

Zhong, S. *et al.* (2019) Nitrogen Addition Increases the Sensitivity of Photosynthesis to Drought
and Re-watering Differentially in C3 Versus C4 Grass Species. *Front Plant Sci* **10**, 815

Figure captions:

Figure 1: Panel a) Bray-Curtis dissimilarity is calculated using the formula $\frac{B+C}{2A+B+C}$ where B

denotes species lost from the early to late community, C denotes species gained from the early to

late community, and A denotes shared species between the two communities. Panel b) Null

expectations of dissimilarity are obtained by permuting species abundances n number of times (3

times shown) while holding richness and total abundance within communities constant and

calculating Bray-Curtis dissimilarity from these permuted communities. These values provide a

distribution from which null mean (μ_{null}) and variance (σ_{null}) values of dissimilarity are

estimated. Deviation (i.e., Seasonal β-diversity (z-scores)) values are then obtained using the

following formula: Seasonal $\beta - \text{diversity}_{z-score}$ $\frac{BC_{observed} - \mu_{null}}{\sigma_{null}}$ where $BC_{observed}$ is the

observed Bray-Curtis value from each early to late comparison, μ_{null} is the expected mean from

the distribution of permutation comparisons, and σ_{null} is the variance from the distribution of

permutation comparisons. Symbols in both panels represent species.

(a) Bray-Curtis (BC) Dissimilarity $= \dfrac{B+C}{2A+B+C}$

Early community

Late community

(B) Losses = 4   (A) Common = 11   (C) Gains = 6

(b) Null Model and Seasonal β-diversity $_{\text{(z-score)}}$

Early community permutations

Late community permutations

BC_1

BC_2

BC_3

μ_{null}

σ_{null}

$BC_1 \ldots BC_2 \ldots BC_i$

Seasonal β-diversity$_{\text{z-score}} = \dfrac{BC_{\text{observed}} - \mu_{\text{null}}}{\sigma_{\text{null}}}$

17   **Figure 2:** Relationships between intra-annual temperature (panel a) and precipitation (panel b)

18   variability and mean site-level seasonal β-diversity in control treatments. Gray lines show 500

19   predicated relationships from posterior model estimates and dark blue lines show averages of

20   these expectations when the distribution of the slope estimate differed from zero at the $\alpha = 0.1$

21   probability level.

Figure 3: Panel a) Seasonal β-diversity in control and fertilized (i.e., NPKμ) treatments. Circle points are seasonal β-diversity values (i.e., z-scores) from each plot in each year for a given site (different colors) and diamond points are average site-level seasonal β-diversity estimates for each site in each treatment. Panel b) Posterior distributions of seasonal β-diversity in control and NPKμ treatments across all sites and years included in analyses. Horizontal lines and points denote estimated means and error bars denote 90% credible intervals of the distributions.

Figure 4: Posterior distributions of nestedness (panel a) and turnover (panel b) components of observed Bray-Curtis dissimilarity in control and fertilized (i.e., NPK) treatments across all sites and years included in analyses. Points and vertical lines denote estimated means and horizonal line segments denote 90% credible intervals of the distributions.

Figure 5: Posterior distributions of annual forbs (panel a, 6 sites included in analyses), perennial

forbs (panel b, 10 sites included in analyses), C3 graminoids (panel c, 6 sites included in

analyses), C4 graminoids (panel d; 6 sites included in analyses), and legumes (panel e; 8 sites

included in analyses) early (dark colors) and late (light colors) in the growing season in control

(grey) and fertilized (red) treatments. Different letters denote differences between groups at the α

= 0.1 probability level, points denote mean estimates, and horizonal line segments denote 90%

credible intervals of the distributions. Please note the different scales on the x-axes.