Title: Temperature seasonality and nutrient enrichment drive intra-annual community turnover

in global grasslands

Authors: Magda Garbowski[1,2], Elizabeth Boughton[3], Anne Ebeling[4], Philip Fay[5], Yann Hautier[6],

Hanna Holz[7], Anke Jentsch[8], Stephanie Jurburg[1,2], Emma Ladouceur[1,2,9], Jason Martina[10],

Timothy Ohlert[11,12], Xavier Raynaud[13], Christiane Roscher[1,2], Grégory Sonnier[3], Pedro

Maximiliano Tognetti[14], Laura Yahdjian[14], Peter Wilfahrt[15], Stan Harpole[1,2,7]

* Corresponding author email: Magda.Garbowski@uwyo.edu

1) German Centre for Integrative Biodiversity Research (iDiv) Leipzig, Germany

2) Department of Physiological Diversity, Helmholtz Centre for Environmental Research - UFZ

Leipzig, Germany

3) Archbold Biological Station, Venus, Florida, USA

4) University Jena, Institute of Ecology and Evolution, Jena, Germany

5) USDA-ARS Grassland Soil and Water Lab, Temple, Texas, USA

6) Ecology and Biodiversity Group, Department of Biology, Utrecht University, Utrecht, The

Netherlands

7) Martin Luther University Halle-Wittenberg, Halle (Saale), Germany

8) Department of Disturbance Ecology and Vegetation Dynamics, Bayreuth Center of Ecology

and Environmental Research (BayCEER), University of Bayreuth, Bayreuth, Germany

9) Department of Biology, University of Leipzig, Germany

10) Department of Biology, Texas State University, San Marcos, Texas 78666, USA

11) Department of Biology, University of New Mexico, Albuquerque, New Mexico, USA

12) Department of Biology, Colorado State University, Fort Collins, Colorado, USA

13) Sorbonne Université, Université Paris-Cité, UPEC, IRD, CNRS, INRA, Institute of Ecology

and Environmental Sciences, iEES Paris, Paris, France

14) IFEVA, Universidad de Buenos Aires, CONICET, Facultad de Agronomía, Catedra de

Ecología, Buenos Aires, Argentina

15) Department of Ecology, Evolution and Behavior, University of Minnesota, St Paul,

Minnesota, USA

* Corresponding author's present address: Botany Department, 1000 E. University Ave.

University of Wyoming, Laramie, WY 82071

Abstract

In many grasslands, species with specific traits occupy unique temporal positions within

communities. Such intra-annual segregation is predicted to be greatest in systems with high

variability (i.e., seasonality) in precipitation and temperature because fluctuating environmental

conditions select for species that capitalize on resources at different times. However, because

most studies on intra-annual community dynamics have been conducted at individual sites,

relationships between seasonality and intra-annual turnover at global scales have not yet been

identified. Furthermore, the same characteristics that promote species-specific responses to

seasonal fluctuations in environmental conditions may also drive species-specific responses to

global change drivers such as eutrophication. Research provides evidence that eutrophication

alters inter-annual plant community dynamics yet understanding of how it alters intra-annual

dynamics remains limited.

We used early-season and late-season compositional data collected from 10 grassland

sites around the world to ask how seasonality in precipitation and temperature as well as nutrient

enrichment shape intra-annual turnover in plant communities. In addition, we assessed whether

changes in the abundances of specific functional groups including annual forbs, perennial forbs,

C3 and C4 graminoids, and legumes underpin compositional differences between early- and late-

season communities and treatments. We found that temperature seasonality and intra-annual

turnover were positively related but observed no relationship between precipitation seasonality

and intra-annual turnover. These results conflict with theoretical predictions from earlier studies

which suggest precipitation seasonality drives intra-annual turnover in plant communities. In

addition, nutrient enrichment increased intra-annual turnover; this pattern was underpinned by

increased replacement of species between early- and late-season communities in fertilized

conditions. This finding mirrors patterns observed at inter-annual scales and suggests fertilization

can alter compositional dynamics via similar mechanisms at varied temporal scales. Finally,

fertilization reduced the abundance of C4 graminoids and legumes and eliminated intra-annual

differences in these groups. In contrast, fertilization resulted in intra-annual differences in C3

graminoids which were not observed in control conditions, and increased abundance of C3

graminoids and annual forbs overall. Our study provides novel insight into how seasonality and

nutrient enrichment shape intra-annual grassland dynamics and highlights how discerning these

patterns is essential to our understanding of biodiversity in these systems.

Keywords: community composition, grasslands, intra-annual climate variability, nutrient

enrichment, Nutrient Network, precipitation seasonality, temporal beta-diversity, temporal niche

segregation, temperature seasonality

Introduction

The importance of temporal separation of resource use by species, often termed "temporal niche segregation", has been recognized as an important component of species coexistence for decades (e.g., Wiens et al. 1977). In many grasslands, species with specific strategies optimally utilize resources under different environmental conditions and thus coexist by occupying unique temporal positions within communities. For example, in grasslands co-dominated by C3 and C4 grasses, C3 grasses grow and set seed primarily in early to mid-season and C4 grasses grow and set seed primarily in late season. Similar dynamics occur in communities without C3 and C4 grasses present. For example, in temperate European grasslands early-season grasses and forbs contribute most to productivity at the beginning of the season, whereas late-season grasses contribute most to productivity late in the growing season (e.g., Guimarães-Steinicke et al. 2019, Doležal et al. 2017). Theory suggests that predictable fluctuations in environmental conditions (i.e., seasonality) should promote unique temporal assemblages of species by allowing different species to capitalize on optimal conditions at different times (Tonkin et al. 2017). However, the same characteristics that promote species-specific responses to seasonal fluctuations in environmental conditions often drive species-specific responses to global change drivers such as eutrophication. For example, subordinate forbs and legumes that thrive at different time periods from dominant grasses are often reduced in fertilized conditions in European meadows (e.g., Doležal et al. 2017). Research from the last several years has elucidated how eutrophication alters inter-annual plant community dynamics (e.g., Hautier et al. 2014, Chen et al. 2021). Yet, interactions between global change drivers and intra-annual dynamics of plant communities remain largely unexplored (White and Hastings 2020).

Seasonality in temperature and precipitation may be particularly important drivers of

temporal niche segregation, or intra-annual turnover, in plant communities as they influence

plant physiology and phenology and modulate ecosystem processes related to water availability

and nutrient cycling (Luo et al. 2020). Modeling studies have demonstrated how intra-annual

variation in precipitation supports coexistence among plant species with slight ecological

differences in germination requirements, growing season length, or seasonal growth activity

(Mathias and Chesson 2013). Empirical evidence documenting the influence of seasonality on

intra-annual turnover in plant communities mainly comes from studies of desert systems in

which winter and summer annuals segregate temporally in response to seasonal fluctuations in

precipitation (e.g., Mulroy and Rundel 1977, Guo and Brown 1997). In addition, studies in

tropical forests (Sapijanskas et al. 2014) and grasslands (Fargione and Tilman 2005) have

demonstrated how within-season niche partitioning enhances complementarity in resource use

and coexistence among co-occurring species. However, because these studies have been

conducted in individual plant communities, relationships between seasonality and intra-annual

turnover at global scales have not yet been identified. In addition, plant communities adapted to

seasonal segregation may be changing in composition and structure in response to global change

drivers such as eutrophication.

Eutrophication in grasslands leads to reduced plant species richness and diversity (e.g.,

Hautier et al. 2009, Harpole et al. 2016). Evidence for these patterns has primarily been gathered

from studies that measure plant community composition once a year, usually at the peak of

biomass production (e.g., Borer et al. 2014). However, research has demonstrated that diverse

resource-use strategies allow distinct species assemblages to dominate in different points within

a growing season (Doležal et al. 2017, Guimarães-Steinicke et al. 2019, Huang et al. 2019). In

grassland systems, nutrient enrichment may promote the dominance of species with specific

traits for longer periods of time and reduce overall turnover among early- and late-season

communities (Doležal et al. 2017). Further, the addition of multiple limiting nutrients may

reduce the importance of species-specific trade-offs associated with competition for particular

nutrients (Harpole et al. 2016) which may be most limiting at different points throughout the

growing season (Klaus et al. 2016). Assessing how nutrient enrichment affects compositional

turnover between early and late-season assemblages would clarify whether patterns of reduced

diversity in response to fertilization at inter-annual scales are underpinned by reduced turnover at

intra-annual scales.

Changes in overall diversity in responses to nutrient enrichment often result from changes

in the abundance of species that are affected by nutrient enrichment in different ways. For

example, nitrogen enrichment reduces the abundance of legumes and nutrient conservative C4

grasses but increases the abundance of nutrient acquisitive C3 grasses (e.g., Suding et al. 2005,

Tognetti et al. 2021). Fertilization with multiple nutrients can further alter community

composition by causing shifts in the relative abundance of species from specific functional

groups (Wilcots et al. 2021). When nutrient enriched communities become dominated by specific

species or functional groups, ecosystem stability may be reduced via a loss of asynchronous

species responses to environmental fluctuations (Hector et al. 2010, Hautier et al. 2014). In

addition, nutrient additions that result in the loss of forb and legume species can have cascading

effects on organisms at other trophic levels, such as pollinators, that often depend on these

groups for floral resources (Burkle and Irwin et al. 2010, Dyer et al. 2021). Understanding how

nutrient enrichment affects the abundances of specific functional groups at specific timepoints in

the growing season (i.e., early vs late) and between timepoints would clarify when compositional

changes or shifts in dominance take place and which species drive them.

In this study, we used above-ground species composition data collected early and late in the

growing season from 10 grassland sites around the world to assess how seasonality in

temperature and precipitation as well as eutrophication influence intra-annual turnover of plant

assemblages. The 10 sites are part of the Nutrient Network – a globally replicated experiment in

which herbaceous plant communities are supplemented with fertilizer. Data from each site

spanned between 4 and 10 years. We used data from untreated plots to assess relationships

between precipitation and temperature seasonality (estimates obtained from WorldClim; Fick

and Hijmans 2017) and intra-annual compositional turnover across our study sites. We then

examined the effects of fertilization on intra-annual turnover and its components of nestedness

and replacement. Finally, to determine which species underpin changes in intra-annual turnover

in response to fertilization, we assessed overall changes in abundance of different functional

groups (i.e., C3 graminoids, C4 graminoids, annual forbs, perennial forbs, legumes) between

early and late timepoints and fertilization treatments. Our hypotheses are that:

H1. Sites with high intra-annual variability in temperature and precipitation (i.e., high

seasonality) have high intra-annual turnover.

H2. Fertilization decreases intra-annual turnover because a subset of species will dominate

across early and late sampling timepoints.

H3. Differences in early-season vs. late-season community composition in response to

fertilization are underpinned by increased abundances of resource acquisitive species

(e.g., C3 graminoids, annual forbs) and decreased abundances of species with more

159   conservative strategies (e.g., C4 graminoids, perennial forbs, legumes) within and

160   between specific timepoints (i.e., early vs. late)

161 **Methods**

162 *Study design and site locations*

163   Data for this study were collected from 10 grassland sites from around the world that are

164 part of the Nutrient Network – a globally distributed experiment in which plant communities are

165 supplemented with factorial combinations of nitrogen (N), phosphorous (P), potassium (K) and

166 micronutrients (μ) (Borer et al. 2014). While most Nutrient Network sites collect compositional

167 data once a year, data from this subset of sites is collected twice each growing season (named

168 "early" and "late" hereafter). These sites are distributed across five continents, in Africa

169 (ukul.za), Australia (burrawan.au), Europe (bayr.de, cereep.fr, frue.ch, jena.de), North America

170 (arch.us, temple.us, sevi.us) and South America (chilcas.ar) (Table S1).

171   Compositional data used in this study were collected from 3 to 5 replicated control and

172 NPKμ plots at these sites from between 4 (bayr.de) and 11 years (ukul.za) (mean length of

173 experiment is 6.5 years, details in Table S1). Fertilized plots at all sites besides cereep.fr received

174 10 g m^{-2} of N, P, and K annually with a one-time addition of micronutrients; fertilized plots at

175 the cereep.fr site received 2.5 g m^{-2} annually with a one-time addition of micronutrients. Despite

176 this different application rate, all results were qualitatively similar whether the cereep.fr site was

177 or was not included in analyses, so we retain it in the results presented here. For additional

178 details about experimental design, please see Borer et al. (2014).

179 *Quantifying temperature and precipitation seasonality*

180   Estimates of temperature and precipitation seasonality for each site were obtained from

181 WorldClim (Fick and Hijmans 2017). Temperature and precipitation seasonality for each site are

182 based on monthly totals of precipitation and temperature where "Temperature seasonality" is the

183 standard deviation of monthly temperature * 100 and "Precipitation seasonality" is the

184 coefficient of variation of monthly precipitation totals for a given year. In both cases, higher

185 values indicate higher variance or greater fluctuations in temperature and precipitation within a

186 year. We chose to use annual seasonality values rather than growing season values because

187 winter precipitation at some of our sites (e.g., sevi.us) is an important driver of plant community

188 dynamics. Between 18 and 30 years of precipitation and temperature seasonality data were used

189 to obtain average seasonality values for each site based on data availability in WorldClim.

190 *Quantifying intra-annual turnover in communities*

191  Dissimilarity indices are useful tools for measuring differences between communities in

192 space or time. However, dissimilarity indices can be influenced by local community size (i.e., α-

193 diversity) and overall richness of species at regional scales (i.e., γ-diversity) (e.g., Chase and

194 Myers 2011). Deviations from null expectations of dissimilarity (e.g., z-scores) can help

195 determine whether compositional differences, independent of local community size and regional

196 species pools, drive dissimilarity patterns (e.g., Chase and Myers 2011). More specifically, larger

197 z-scores indicate higher dissimilarity between communities, whereas lower deviations indicate

198 lower dissimilarity.

199  To quantify intra-annual turnover between early and late communities in control and NPKμ

200 treatments at each site we used deviations from expected values of the Bray-Curtis dissimilarity

201 index (Figure 1). We did this by first permuting species abundances (i.e., absolute values of

species cover) from permanent m^2 survey plots 100 times while holding overall richness and total

abundances of species constant within a plot at a given sampling timepoint (i.e., early or late

season). We used these 200 permuted communities (i.e., 100 permutations for "early"

communities and 100 permutations for "late" communities) to obtain mean and standard

deviation values for expected (i.e., null) dissimilarity in each plot. We then calculated z-scores

for each early-to-late comparison for each plot, year, and treatment combination using the

following formula: $Turnover_z = \frac{BC_{observed} - \mu_{null}}{\sigma_{null}}$ where $BC_{observed}$ is the observed Bray-

Curtis value from each plot in each year, μ_{null} is the expected mean for each plot in each year

obtained from the distribution of permutation dissimilarly estimates, and σ_{null} is the variance for

each plot in each year obtained from the distribution of permutation dissimilarly estimates

(Figure 1b). We used these z-scores (indicated throughout as turnover$_z$) as our estimates of early

to late dissimilarity from each plot in each year, hereafter referred to as "intra-annual turnover".

Comparisons from a given plot in a given year with less than four species at either the early or

late sampling time point were removed prior to all analyses because communities with less than

four species could not produce sufficient permutations to calculate reliable estimates of

turnover$_z$. This resulted in the exclusion of 53 early-to-late comparisons from 888 comparisons

total (i.e., 835 comparisons were included in analyses).

We also decomposed observed Bray-Curtis dissimilarity from each early-to-late comparison

for each plot, year, and treatment into components of species "replacement" and "nestedness"

(Baselga 2009). Nestedness captures the component of dissimilarity that results from one

community being a subset of another community, whereas replacement captures the component

of dissimilarity that results from certain species being lost and others gained between compared

communities. We used the beta.part package (Baselga and Orme 2012) to obtain all dissimilarity

estimates.

Statistical models

We used two separate linear regression models to assess relationships between

temperature and precipitation seasonality and intra-annual turnover of plant communities. In

these models, average site temperature seasonality or precipitation seasonality values were

included as a continuous predictor and average turnover$_z$ estimates from control treatments from

each site were included as a continuous response. We examined how NPKµ treatment influenced

intra-annual turnover using a multilevel regression model with plot-level turnover$_z$ values

included as response variables, treatment as a predictor, and a random effect of block, nested

within treatment year, nested within site. Because not all functional groups are present at all

sites, functional group analyses were completed using data from sites that had specific functional

groups present at >1% cover overall. We fit five separate multilevel regression models to

quantify how abundance of specific functional groups (i.e., summed cover of all species

belonging to annual forbs, perennial forbs, C3 grasses, C4 grasses and legumes from m^2 plots)

differed among early and late sampling timepoints and treatments. These models included

treatment by season combinations as categorical predictors (i.e., control_early, control_late,

NPKµ_early, NPKµ_late) and block nested within treatment year nested within site as a random

effect (i.e., interactions were not assessed prior to investigating differences in means).

For Bayesian inferences and estimates of uncertainty, all models described were fitted

using the Hamiltonian Monte Carlo (HMC) sampler using Stan (Carpenter et al. 2017) and coded

using the 'brms' package (Bürkner et al. 2018) in R (version V.2.1 R Core Development Team).

All models were fit with 4 chains and 3000 iterations (Table S2). We used default priors for all

models in which dissimilarly estimates (i.e., $turnover_z$, nestedness, replacement) were included

as response variables. In models assessing differences in functional group abundances between

timepoints and treatments, default priors were used for C3 and C4 graminoid and perennial forb

models, whereas weakly regularizing priors were used in models assessing differences in legume

and annual forb abundances (Table S2). We inspected the HMC chains to assess model

convergence. For all models, we estimated the significance of effects by computing the

difference between posterior distributions of interest and assessing whether the 90% credible

interval of the difference contained zero.

Results

We found a positive relationship between temperature seasonality and seasonal turnover

in unfertilized control plots (Slope: 0.0028, 90% Credible Interval (CI): 0.0004 to 0.0050)

(Figure 2a) and no relationship between precipitation seasonality and seasonal turnover (Slope: -

0.0052, CI: -0.0278 to 0.0166) (Figure 2b). Contrary to our hypothesis, we detected a significant

positive effect of fertilization on seasonal $turnover_z$ across sites (Figure 3). This result indicates

greater dissimilarity in fertilized conditions, even after accounting for differences in richness

between the two treatments (average number of species per m^2 in control: 12.8; average number

of species per m^2 in NPKμ: 11.5). However, $turnover_z$ values were negative in most cases,

indicating lower than expected dissimilarity between early-season and late-season communities.

After decomposing observed (i.e., Bray-Curtis) intra-annual turnover into components of

nestedness and replacement, we found no differences between treatments in nestedness (Figure

4a) but higher replacement in fertilized conditions than in controls (Figure 4b).

The effects of fertilization varied by functional group and between early and late

sampling timepoints (Figure 5). At both sampling timepoints, fertilization resulted in higher

cover of annual forbs (Figure 5a) and C3 graminoids (Figure 5c) and lower cover of C4

graminoids (Fig. 5d) and legumes (Fig. 5e). Perennial forb cover was reduced with fertilization,

but only at the late sampling timepoint (Fig. 5b). Patterns of functional group abundances

between early and late sampling points also varied between control and fertilized treatments. In

control but not fertilized treatments, C4 graminoid abundance was higher at the later sampling

point compared to early in the season and legume abundance was higher early in the season

compared to late in the season. In contrast, abundance of C3 graminoids was higher early in the

season compared to late in the season but only in fertilized treatments.

Discussion

Contrary to most empirical and theoretical studies to date (e.g., Mulroy and Rundel 1977,

Mathias and Chesson 2013), we found temperature seasonality, not precipitation seasonality, to

be positively related to intra-annual compositional turnover across our study sites. We suspect

this relationship arises due to differences in annual systems, from which most evidence to date

has been gathered, and the predominantly perennial systems in our study. In annual systems,

population, community, and ecosystem dynamics are often governed by interactions between

intra-annual precipitation patterns and demographic traits related to dormancy and germination

(e.g., Levine et al. 2011, Kimball et al. 2011, Shaw et al. 2022). In addition, in systems co-

dominated by annuals and perennials, intra-annual variability in precipitation is often necessary

for the persistence of annuals that occupy unique temporal niches in communities (e.g., Pérez-

Camacho et al. 2012). If precipitation is highly seasonal, distinct assemblages of annual species

may be promoted early vs. late in the growing season. On the other hand, temperature and

temperature seasonality may be more important drivers of intra-annual dynamics in perennial

systems because temperature is a key driver of species physiology which drives species temporal

segregation in many perennial systems (e.g., Kemp and Williams. 1980, Monson et al. 1983).

Given the low abundance of annual cover across our study sites (*c.* 11%), we are not able to test

these patterns robustly. But a positive, non-significant, relationship between annual species

dissimilarity and precipitation seasonality (Supplementary Figure 1a) and a negative, non-

significant, relationship between perennial species dissimilarity and precipitation seasonality

(Supplementary Figure 1b) invite new questions about how precipitation seasonality influences

intra-annual community dynamics across diverse ecosystems. Studies that explicitly manipulate

precipitation seasonality across systems varying in temperature seasonality could help clarify

how interactions between these two understudied components of climate change shape plant

communities.

The negative impacts of nutrient enrichment on grassland diversity have been extensively

documented (e.g., Borer et al. 2017) and recent studies have identified higher turnover of species

at inter-annual scales with nutrient addition (Koerner et al. 2016, Hodapp et al. 2018, Chen et al.

2021). Mirroring these results at intra-annual scales, we found that fertilization resulted in higher

intra-annual turnover in community composition, despite lower richness in fertilized

communities. Because we used deviations from null expectations as our response, we can be

confident that these shifts arise from true changes in composition rather than random processes

influenced by richness. Similar to results found at inter-annual scales (Chen et al. 2021), we

found that higher turnover with fertilization was driven by higher replacement of species, and not

nestedness of species, between early and late-season communities. This suggests that fertilization

increases temporal turnover among and within years by similar mechanisms, namely by unique

species occupying space within the community at different times (i.e., replacement) rather than

species being lost from one timepoint to another (i.e., nestedness). At inter-annual scales,

increased turnover and species replacement through time can result in reduced stability of

ecosystem productivity (e.g., Koerner et al. 2016, Chen et al. 2021). Additional research focused

on how nutrient enrichment alters composition and associated ecosystem functions within

growing seasons would clarify whether similar patterns manifest at intra-annual scales.

As expected, fertilization resulted in higher abundance of resource acquisitive species

(i.e., C3 graminoids, annual forbs) and lower abundance of resource conservative species (i.e.,

C4 graminoids, legumes, perennial forbs) within and across sampling timepoints. Numerous

other studies have documented similar patterns at single timepoints (e.g., Suding et al. 2005,

Isabell et al. 2013, Wilcots et al. 2021, Tognetti et al. 2021) and results are primarily attributed to

physiological differences between these groups and concomitant shifts from below-ground

competition for nutrients to aboveground competition for light (e.g., Borer et al. 2014, Harpole et

al. 2017). Among other anatomical and physiological differences, compared to C4 species, C3

species (i.e., most of the annuals in our study and C3 graminoids) generally have lower C:N

ratios (e.g., Wedin and Tilman 1990), lower water and photosynthetic nitrogen use efficiencies

(Black 1971, Taylor et al. 2011, Ripley et al. 2010), and earlier phenologies due to lower optimal

temperatures for photosynthesis (Kemp and Williams 1980). These characteristics can lead to

higher sensitivities of resource acquisitive species to nutrient limitation which allow them to

increase in fertilized conditions (e.g., Zhong et al. 2019).

Fertilization made apparent seasonal differences in C3 graminoid abundance that were

not present under control, while eliminating seasonal differences in abundance of C4 graminoids

and legumes that were present in control treatments. Higher C3 graminoid abundance early in the

season in fertilized treatments could reflect a higher capacity for growth of these species early in

the season, when resources are abundant, followed by more pronounced declines later in the

season when resources are scarce (e.g., Yuan et al. 2007). In contrast, both C4 graminoids and

legumes are adapted to resource-limited conditions which can vary throughout growing seasons

(Klaus et al. 2014). By altering resource conditions, fertilization likely modulates the low-

resource periods during which these species thrive resulting in lower abundances of these species

overall and homogenization of these groups across the growing season.

Our understanding of how grasslands respond to various components of global change is

primarily based on studies that document community shifts at inter-annual scales. However,

global changes may also disrupt within-season community dynamics. Here we show that intra-

annual turnover of species across 10 global grasslands is related to temperature seasonality, not

precipitation seasonality as predicted by previous studies. Given that both temperature and

precipitation seasonality are expected to change in coming decades (Xu et al. 2013, Hajek and

Knapp 2021) additional research that explicitly manipulates seasonality is needed to clarify how

these understudied components of global change alter plant communities. In addition, we show

that nutrient enrichment increases intra-annual turnover in grasslands via increased replacement

of species between seasons. The resulting community shifts are underpinned by changes in

abundance of specific functional groups. Namely, fertilization increases abundances of resource-

acquisitive species (i.e., annual forbs, C3 graminoids) and decreases abundances of resource-

conservative species (i.e., C4 graminoids, legumes) within and between early and late sampling

timepoints. In addition, fertilization results in homogenization of abundances of resource

conservative species early to late in the growing season. If the effects of intra-annual community

change mirror those at inter-annual scales, the community shifts we observed in our study could

have cascading effects on ecosystem stability, multifunctionality, and the ability of these systems

to recover from future perturbations. Our study provides novel insight into how seasonality and

nutrient enrichment shape within-season community dynamics in global grasslands and

highlights how discerning these patterns is essential to our understanding of biodiversity in these

valuable ecosystems.

Acknowledgements: Funding for this project was provided by the German Science Foundation

(DFG; via University Leipzig/iDiv) for the project "Strategisches Projekt: NEu – Nutrient

Network Europe" (Projektnummer 346001057-03).

Author contributions: MG and SH conceptualized this project, MG completed analyses with

assistance from HH, SJ, and EL, MG wrote the manuscript, all authors contributed to data

collection and editing the manuscript.

References:

Baselga, A. (2009) Partitioning the turnover and nestedness components of beta-diversity. *Global Ecology and Biogeography*. **19** (1) 134 - 143

Baselga, A & Orme D.L. (2012) betapart: an R package for the study of beta diversity. *Methods in Ecology and Evolution* **3**, 808-812

Borer, E. T. et al. (2014) Herbivores and nutrients control grassland plant diversity via light limitation. *Nature* **508**, 517–520

Borer, E.T., Grace, J.B., Harpole, W.S., MacDougall, A.S., Seabloom, E.W. (2017) A decade of insights into grassland ecosystem responses to global environmental change. *Nature Ecology & Evolution* **1**, 0118

Burkle L.A. & Irwin R.E. (2010) Beyond biomass: measuring the effects of community-level nitrogen enrichment on floral traits, pollinator visitation and plant reproduction. *Journal of Ecology*. **98** (3), 705 – 717

Bürkner P.C. (2017) "brms: An R Package for Bayesian Multilevel Models using Stan." Journal of Statistical Software, 80(1), 1–28

Carpenter B., Gelman A., Hoffman M.D., et al. (2017) Stan: a probabilistic programing language. *Journal of Statistical Software*. **76** (1)

Chase, J. M. & Myers, J. A. (2011) Disentangling the importance of ecological niches from stochastic processes across scales. *Philosophical Transactions Royal Soc B Biological Sci* **366**, 2351–2363

Chen, Q. et al. (2022) Nutrients and herbivores impact grassland stability across spatial scales through different pathways. *Global Change Biol* **28**, 2678–2688

Doležal, J., Lanta, V., Mudrák, O. & Lepš, J. (2019) Seasonality promotes grassland diversity: Interactions with mowing, fertilization and removal of dominant species. *J Ecol* **107**, 203–215

Dyer, A.G., Jentsch, A., Burd, M., Garcia, J.E., Giejsztowt, J., Camargo, M.G.G., Tjørve, E., Tjørve, K.M.C., White, P.S., Shrestha, M. (2021) Fragmentary blue: Resolving the rarity paradox in flower colours. *Frontiers in Plant Science* 11: Article 618203

Ehleringer, J. & Pearcy, R. W. (1983) Variation in Quantum Yield for CO_2 Uptake among C_3 and C_4 Plants. *Plant Physiol* **73**, 555–9

Fargione, J. & Tilman, D. (2005) Niche differences in phenology and rooting depth promote coexistence with a dominant C4 bunchgrass. *Oecologia* **143**, 598–606

Fick S.E. & Hijmans, R.J. (2017) WorldClim 2: new 1-km spatial resolution climate surfaces for global land areas. *International Journal of Climatology.* **37**, 4302–4315

Guimarães-Steinicke, C., Weigelt, A., Ebling, A., Eisenhauer, N., Duque-Lazo, L., Reu, B., Roscher, C., Schumacher, J., Wagg, C., Wirth, C. (2019) Terrestrial laser scanning reveals temporal changes in biodiversity mechanisms driving grassland productivity. *Advances in Ecological Research* **61**, 133-161

Hajek, O. L. & Knapp, A. K. (2022) Shifting seasonal patterns of water availability: ecosystem responses to an unappreciated dimension of climate change. *New Phytol* **233**, 119–125

Harpole, W. S. et al. (2016) Addition of multiple limiting resources reduces grassland diversity. *Nature* 1–9

Harpole, W. S. et al. (2017) Out of the shadows: multiple nutrient limitations drive relationships among biomass, light and plant diversity. *Functional Ecology* **31**, 1839–1846

Hautier, Y., Niklaus, P. A. & Hector, A. (2009) Competition for Light Causes Plant Biodiversity Loss After Eutrophication. *Science* **324**, 636–638

Hautier, Y., Seabloom E., Borer E., et al. (2014) Eutrophication weakens stabilizing effects of diversity in natural grasslands. *Nature* **508**, 521 – 525

Hector, A. Hautier, Y., Wacker S.L., et al. (2010) General stabilizing effects of plant diversity on grassland productivity through population asynchrony and overyielding. *Ecology* **91** (8), 2213 – 2220

Hodapp, D. (2018) *et al.* Spatial heterogeneity in species composition constrains plant community responses to herbivory and fertilisation. *Ecology Letters* **21**, 1364–1371

Huang, L., Xue, W. & Herben, T. (2019) Temporal niche differentiation among species changes with habitat productivity and light conditions. *Journal of Vegetation Science* **30**, 438–447

Guo, Q., & J. H. Brown. (1997) Interactions between winter and summer annuals in the Chihuahuan desert. *Oecologia* 111:123–128.

Isabell, F. *et al.* (2013) Nutrient enrichment, biodiversity loss, and consequent declines in ecosystem productivity. *Proc National Acad Sci* **110**, 11911–11916

Kemp, P. R. & Williams, G. J. A (1980) Physiological Basis for Niche Separation Between *Agropyron Smithii* (C3) and *Bouteloua Gracilis* (C4). *Ecology* **61**, 846–858

Klaus, V.H. et al. (2016) Temporal and small-scale spatial variation in grassland productivity, biomass quality, and nutrient limitation. *Plant Ecology.* **217**, 843 – 856

Koerner, S. E. et al. (2016) Nutrient additions cause divergence of tallgrass prairie plant
communities resulting in loss of ecosystem stability. *J Ecol* **104**, 1478–1487

Luo, Y. *et al.* (2020) Nutrients and water availability constrain the seasonality of vegetation
activity in a Mediterranean ecosystem. *Global Change Biol.*

Mathias, A. & Chesson, P. (2013)Coexistence and evolutionary dynamics mediated by seasonal
environmental variation in annual plant communities. *Theor Popul Biol* **84**, 56–71

Monson, R. K., Littlejohn, R. O. & Williams, G. J. (1983) Photosynthetic adaptation to
temperature in four species from the Colorado shortgrass steppe: a physiological model for
coexistence. *Oecologia* **58**, 43–51

Mulroy, T. W. & Rundel, P. W. (1977)Annual Plants: Adaptations to Desert Environments.
Bioscience **27**, 109–114

Myers, J. A. *et al.* (2013) Beta-diversity in temperate and tropical forests reflects dissimilar
mechanisms of community assembly. *Ecol Lett* **16**, 151–157

Myers, J. A., Chase, J. M., Crandall, R. M. & Jiménez, I. (2015) Disturbance alters beta-diversity
but not the relative importance of community assembly mechanisms. *J Ecol* **103**, 1291–1299

Pérez-Camacho L., Rebello, S., Hernández-Santana, V., García-Salgado, J., Pavón-García, J.,
Gómez-Sal., A. (2012) Plant functional trait responses to interannual rainfall variability, summer
drought and seasonal grazing in Mediterranean herbaceous communities. *Functional Ecology* **26**,
740 – 749

Ripley, B., Frole, K. & Gilbert, M. (2010) Differences in drought sensitivities and photosynthetic
limitations between co-occurring C3 and C4 (NADP-ME) Panicoid grasses. *Ann Bot-london* **105**,
493–503

Sapijanskas, J., Paquette, A., Potvin, C., Kunert, N. & Loreau, M. (2014) Tropical tree diversity
enhances light capture through crown plasticity and spatial and temporal niche differences.
Ecology **95**, 2479–2492

Schofield, E. J., Rowntree, J. K., Paterson, E. & Brooker, R. W. (2018) Temporal Dynamism of
Resource Capture: A Missing Factor in Ecology? *Trends in Ecology & Evolution* **33**, 277–286

Suding, K. N. *et al.* (2005) Functional- and abundance-based mechanisms explain diversity loss
due to N fertilization. *Proceedings of the National Academy of Sciences* **102**, 4387–4392

Taylor, S. H. *et al.* (2014) Physiological advantages of C4 grasses in the field: a comparative
experiment demonstrating the importance of drought. *Global Change Biol* **20**, 1992–2003

Tognetti, P. M. *et al.* (2021) Negative effects of nitrogen override positive effects of phosphorus
on grassland legumes worldwide. *Proc National Acad Sci* **118**

Tonkin, J. D., Bogan, M. T., Bonada, N., Rios-Touma, B. & Lytle, D. A. (2017) Seasonality and
predictability shape temporal species diversity. *Ecology* **98**, 1201–1216
509
510 Wedin, D. A. & Tilman, D. (1990) Species effects on nitrogen cycling: a test with perennial
511 grasses. *Oecologia* **84**, 433–441
512
513 White, E. R. & Hastings, A. (2020) Seasonality in ecology: Progress and prospects in theory.
514 *Ecol Complex* **44**, 100867
515
516 Wiens, J. A. (1977) On competition and variable environments: Populations may experience
517 "ecological crunches" in variable climates, nullifying the assumptions of competition theory and
518 limiting the usefulness of short-term studies of population patterns. *American Scientist*
519
520 Wilcots, M. E., Harpole, W. S., Seabloom, E. W. & Borer, E. T. (2021) Community change can
521 buffer chronic nitrogen impacts, but multiple nutrients tip the scale. *Ecology* **102**
522
523 Xu, L., Myneni R.B., Chapin F.S., et al. (2013) Temperature and vegetation seasonality
524 diminishment over northern lands. *Nature*. **3**, 581 – 586
525
526 Yuan, Z., Liu, W., Niu, S. & Wan, S. (2007) Plant Nitrogen Dynamics and Nitrogen-use
527 Strategies under Altered Nitrogen Seasonality and Competition. *Ann Bot-london* **100**, 821–830
528
529 Zhong, S. *et al.* (2019) Nitrogen Addition Increases the Sensitivity of Photosynthesis to Drought
530 and Re-watering Differentially in C3 Versus C4 Grass Species. *Front Plant Sci* **10**, 815
531

Figures:

Figure 1: Panel a) Bray-Curtis dissimilarity is calculated using the formula $\frac{B+C}{2A+B+C}$ where B

denotes species lost from the early to late community, C denotes species gained from the early to

late community, and A denotes shared species between the two communities. Panel b) Null

expectations of dissimilarity are obtained by permuting species abundances n number of times (3

times shown) while holding richness and total abundance within communities constant and

calculating Bray-Curtis dissimilarity from these permuted communities. These values provide a

distribution from which null mean (μ_{null}) and variance (σ_{null}) values of dissimilarity are

estimated. Deviation (i.e., z-scores, Turnover$_z$) values are then obtained using the following

formula: $Turnover_z \frac{BC_{observed} - \mu_{null}}{\sigma_{null}}$ where BC$_{observed}$ is the observed Bray-Curtis value from each

early to late comparison, μ_{null} is the expected mean from the distribution of permutation

comparisons, and σ_{null} is the variance from the distribution of permutation comparisons. Symbols

in both panels represent species.

Figure 2

Figure 2: Relationships between temperature (panel a) and precipitation (panel b) seasonality and mean intra-annual turnover in control treatments. Gray lines show 500 predicated relationships from posterior model estimates and dark blue lines show averages of these expectations when the distribution of the slope estimate differed from zero at the $\alpha = 0.1$ probability level.

Figure 3

Figure 3: Panel a) Intra-annual turnover in control and fertilized (i.e., NPKμ) treatments. Circle points are turnover z-scores (i.e., deviations from null expectations) from each plot in each year for a given site (different colors) and diamond points are average site-level turnover estimates for each site in each treatment. Panel b) Posterior distributions of intra-annual turnover in control and NPKμ treatments across all sites and years included in analyses. Horizontal lines and points denote estimated means and error bars denote 90% credible intervals of the distributions.

Figure 4

Figure 4: Posterior distributions of nestedness (panel a) and replacement (panel b) components of Bray-Curtis dissimilarity in control and fertilized (i.e., NPK) treatments across all sites and years included in analyses. Points and vertical lines denote estimated means and horizonal line segments denote 90% credible intervals of the distributions.

Figure 5

Figure 5: Posterior distributions of annual forbs (panel a, 6 sites included in analyses), perennial forbs (panel b, 10 sites included in analyses), C3 graminoids (panel c, 6 sites included in analyses), C4 graminoids (panel d; 6 sites included in analyses), and legumes (panel e; 8 sites included in analyses) early (dark colors) and late (light colors) in the growing season in control (grey) and fertilized (red) treatments. Different letters denote differences between groups at the $\alpha = 0.1$ probability level, points denote mean estimates, and horizonal line segments denote 90% credible intervals of the distributions. Please note the different scales on the x-axes.