Part-night exposure to artificial light at night has more detrimental effects on aphid colonies than

fully lit nights

Robin Heinen[1], Oriana Sanchez-Mahecha[1], T. Martijn Bezemer[2], Davide M. Dominoni[3], Claudia

Knappe[8], Johannes Kollmann[4], Anton Kopatsch[7], Zoë A. Pfeiffer[1], Michael Schloter[5,6], Sarah Sturm[1],

Jörg-Peter Schnitzler[7], A. Corina Vlot[8,9], Wolfgang W. Weisser[1]

1 Terrestrial Ecology Research Group, Department for Life Science Systems, TUM School of Life Sciences, Technical
University of Munich, Freising, Germany

2 Institute of Biology, Section Plant Ecology and Phytochemistry, Leiden University, Leiden, The Netherlands

3 School of Biodiversity, One Health & Veterinary Medicine, University of Glasgow, Glasgow, Scotland

4 Chair of Restoration Ecology, Department for Life Science Systems, TUM School of Life Sciences, Technical University of
Munich, Freising, Germany

5 Research Unit Comparative Microbiome Analysis, Helmholtz Munich, Neuherberg, Germany

6 Chair of Soil Science, Department for Life Science Systems, TUM School of Life Sciences, Technical University of Munich,
Freising, Germany

7 Research Unit Environmental Simulation, Helmholtz Munich, Neuherberg, Germany

8 Institute of Biochemical Plant Pathology, Helmholtz Munich, Neuherberg, Germany

9 Chair of Crop Plant Genetics, Faculty of Life Sciences: Food, Nutrition and Health, University of Bayreuth, Kulmbach,
Germany

Author for correspondence: Robin Heinen, robin.heinen@tum.de

Author contributions: RH conceived the original ideas for the study, which were further developed

in close discussion with DD, TMB, JK, MS, JPS, ACV and WWW. RH designed the experiment. RH,

OSM, ZAP and SS conducted the experiment and collected data. CK analyzed gene expression in

plant samples. AK measured and visualized spectral composition in the climate chambers. RH

analyzed the data. RH led the writing of the manuscript and all authors commented and approved

the final version of the manuscript.

Abstract

Artificial light at night (ALAN) threatens natural ecosystems globally. While ALAN research is

increasing, little is known about how ALAN affects plants and interactions with other organisms. We

explored the effects of ALAN on plant defence and plant–insect interactions using barley (*Hordeum*

vulgare) and the English grain aphid (*Sitobion avenae*). Plants were exposed to 'full' or 'part' nights

of 15–20 lux ALAN, or no ALAN 'control' nights, to test the effects of ALAN on plant growth and

defence. Although plant growth was only minimally affected by ALAN, aphid colony growth and

aphid maturation were reduced significantly by ALAN treatments. Importantly, we found strong

differences between full-night and part-night ALAN treatments. Contrary to our expectations, part

ALAN had stronger negative effects on aphid colony growth than full ALAN. Defence-associated gene

expression was affected in some cases by ALAN, but also positively correlated with aphid colony size,

suggesting that the effects of ALAN on plant defences are indirect, and regulated via direct

disruption of aphid colonies, rather than via ALAN-induced upregulation of defences. Mitigating

ecological side effects of ALAN is a complex problem, as reducing exposure to ALAN increased its

negative impact on insect herbivores.

Keywords: Light pollution, ALAN, plant defence, induced defences, gene expression, urban ecology

Introduction

Artificial light at night (ALAN) is increasingly common in urban and rural areas of the world (Gaston, Visser & Hölker, 2015; Kyba et al., 2017). It is emitted by various man-made sources including street lights, building lights, and advertising lights. Current levels of ALAN are unprecedented in the ecological history of life on Earth, as light levels followed a predictable periodicity of diurnal and annual rhythms, with little light at night except of stars and the moon. These rhythms are increasingly disrupted by ALAN in many areas of the world, and ALAN therefore is considered an important aspect of global change (Gaston, Visser & Hölker, 2015; Senzaki et al., 2020; Falcón et al., 2020). For instance, ALAN can interfere with natural behaviour and physiology in many taxa (Sanders et al., 2021), including freshwater and marine invertebrates (Duarte et al. 2019; Perkin et al. 2011; Ayalon et al. 2019), terrestrial insects and several vertebrates, such as bats and birds (Desouhant et al., 2019; Stone, Harris & Jones, 2015; Dominoni, Quetting & Partecke, 2013). Surprisingly, given their dependence on the light environment for acquisition of energy, and as signal for photomorphogenesis, phenological and defence processes, much less is known about how ALAN affects plants (Briggs 2006; Bennie et al. 2016). It can for instance interfere with phenology, starch turnover and defence regulation, which may affect interactions with plant-associated organisms (Heinen 2021). As nights will most likely become significantly brighter, it is important to better understand the impacts on natural ecosystems (Rich & Longcore, 2013). Therefore, ALAN effects on plants and associated organisms warrant further studies (Heinen 2021).

Levels of ALAN will increase in the future because of the decreasing production costs, and the development of more energy- and cost-efficient light-emitting diode (LED) technology making light widely available and affordable (Hölker et al., 2010; Kyba et al. 2017). This will likely also increase the ecological consequences of ALAN. Although many forms of ALAN exist and are still in use, a clear trend towards LED technology has been observed in recent years. The most commonly used LEDs today are cool-white LEDs (with a colour temperature of >4200 K) and these are characterized by a

narrow and high peak in the blue part of the irradiation spectrum, and a broad and lower peak in the

red part of the spectrum. Cool-white LEDs can have devastating effects on many nocturnal animal

species, and to mitigate these detrimental effects, several solutions have been suggested, including

the use of different light colours (Van Geffen et al., 2014; Spoelstra et al., 2015), and to reduce light

hours to periods that people are generally more active (International Darksky Association,

www.darksky.org; Gaston et al. 2012; Day et al. 2015). A plausible scenario that could be

implemented would be to strictly limit ALAN to the first part of the night, when human activity is still

relatively high, and switching off the lights during the second half of the night. This would provide

plants and animals with at least several hours of true darkness to recover. However, how ALAN and

specific light schemes that reduce the number of light hours will affect plant defences and plant-

herbivore interactions, is not known.

One key plant parameter that is affected by the quality, quantity and duration of light that a

plant is exposed to is plant defence. First, a large body of knowledge describes how additional light,

i.e., supplementing plants with artificial light *beyond* the daylight hours, influences plant defences.

Generally, light supplementation into the dark hours increases plant defences. For example, tomato

plants supplemented with low-density light at night have significantly higher immunity to pathogens

compared to plants that grow in a dark night (Yang et al., 2015a). In another study, chickpea plants

supplemented with light at night showed increased levels of the antioxidant ß-carotene (Wu et al.,

2007), which may serve as a defence against oxidative stress (Mittler, 2002). One might conclude

from this that a longer day leads to a more defended plant. A second body of scientific literature

examines the effects of light reduction on plants, i.e., shading and light competition, on plant

physiology, including defences, as partly mediated by changes in light intensity and spectral

composition (e.g., Bennie et al. 2016; Smith et al. 2017; Grubisic et al. 2018). Light competition and

shading can alter key processes in the jasmonic (JA) and salicylic acid (SA) signalling pathways

(reviewed in Ballaré, 2009; Ballaré, 2014; Ballaré & Pierik, 2017), which is the backbone of plant

defences against pathogens and insect herbivores. Reductions in light exposure, through

competition and shading, result in decreased sensitivity to and suppressed JA and SA expression (de

Wit et al., 2013; Ballaré & Pierik, 2017), although the SA-related processes are less well-understood

(Ballaré & Pierik, 2017). Light competition and shade may also lead to reduced accumulation of plant

secondary metabolites, and structural and indirect defences (Izaguirre et al., 2006, 2013; Engelen-

Eigles et al., 2006; Moreno et al., 2009; Agrawal et al., 2013; Cargnel et al., 2014). Relevant to

understanding the effects of ALAN on plants, one result of defence suppression under low-light or

dark conditions is that plants are more susceptible to pathogens and herbivory at night (Roden &

Ingle, 2009; Kraiselburd et al., 2017). This literature suggests that ALAN has the potential to affect

plant defences but that this depends on the specifics of radiation at night.

ALAN may also affect interactions between plants and herbivores (Bennie et al., 2015; Yang et al.,

2015b; Sanders et al., 2018; Heinen 2021). Different typical ALAN sources with different spectral

characteristics have differential bottom-up effects on plant–aphid interactions (Bennie et al., 2015).

For example, supplementing *Lotus pedunculatus* plants with amber ALAN, caused a decrease in

densities of the aphid *Acyrthosiphon pisum* compared to no ALAN in the later stages of seasonal

colony dynamics (Bennie et al., 2015). Importantly, even low light intensities at night can have

significant effects, as ALAN can already suppress aphids under densities as low as 0.1 lux, which is far

below commonly observed levels (Sanders et al., 2018). Although the mechanisms are not fully

understood yet, it could be that ALAN results in systemic activation of defence mechanisms in the

plant. For instance, in one study where tomatoes were supplemented with different spectral

irradiance at night, particularly the plants exposed to red light (and to a lesser extent to other light

colours) showed strong induced defence responses against the root-knot nematode *Meloidogyne*

incognita (Yang et al., 2015b). These defence responses were accompanied by high levels of SA in

root tissues, and upregulated transcript levels of SA-associated genes in the roots (Yang et al.,

2015b). How irradiance levels associated with ALAN affects plant defences, however, is unknown.

In this study, we used a well-characterized plant–aphid model system, barley (*Hordeum vulgare* L.)

and the English grain aphid (*Sitobion avenae* Fabricius), and exposed them to full nights of ca. 20 lux

ALAN ('full-night'), half nights of ca. 20 lux ALAN ('part-night') or no ALAN ('control') to investigate

the effects of ALAN on plant–herbivore interactions and plant defence. We hypothesized that: (i)

Exposure to ALAN reduces aphid colony growth compared to dark nights, mediated by an

upregulation of defence responses. We predict that aphid colonization on plants will be negatively

impacted by plants, through an upregulation of plant defences. (ii) Exposure to ALAN affects plant

growth and health status, compared to dark nights. We predict that effects of ALAN will be positive,

but likely minimal, as the levels of ALAN are expected to be too low to increase photosynthesis. (iii)

A partial reduction in ALAN via a midnight elimination of light sources will reduce the hypothesized

effects of ALAN, serving as a dark recovery time that benefits both plants and insects.

Materials and Methods

Soil

Soils for this experiment were derived from a former flower strip bordering an agricultural field of

the Experimental Station Roggenstein (48.17989, 11.32030; altitude 524 m a.s.l.) at the Technical

University of Munich. The soil was classified as sandy loam (50% sand, 33% silt, 14% clay, 2.4% SOM,

pH 6.8, 1130 mg total-N/kg, 31.5 mg available-S/kg, 1.2 mg available-P/kg, 126 mg available-K/kg).

The soil was sieved through a 2-cm mesh sieve to remove large stones, loam aggregates and roots.

Plant–insect model system

Barley plants (*Hordeum vulgare*, cv. 'Scarlett') were used as model plants. This cultivar was selected

as it was used in previous studies in our research group and because it was highly susceptible to

aphids (Sanchez-Mahecha et al. 2022). As model aphids, we used the English grain aphid (*Sitobion*

avenae 'Fescue', Hemiptera: Aphididae), a species that is specialized on grasses and is a common

pest in cereal cropping systems. We have observed that this barley cultivar and aphid clone respond

in terms of growth and population growth, respectively, to brief pulses of light disruption during the

night in a pilot study. Aphid colonies have been maintained in a climate chamber at the institute at

15 °C (night) to 20 °C (day), under 16h:8h L:D conditions, on barley cultivar 'Chanson' for several

years.

Climate chamber conditions

This study was conducted in the TUM Model EcoSystem Analyzer (TUM*mesa*; Roy et al. 2021) in

Freising, Germany. In this facility, eight separate high-performance climatic chambers allow for a

very high level of between-chamber standardization of abiotic parameters and irradiation conditions

(Teixeira et al., in prep.). Plants were grown under 16h:8h (L:D) daylength regimes, reflecting their

summer growth period, with 21 °C and 16 °C temperatures during day and night, respectively.

Daylights were turned on at 08:00, and were shut off at 00.00, which enabled us to sample under

night conditions between 06:00 and 08:00. Relative air humidity was kept constant at 60%. During

the day, plants received a full-spectrum light (Suppl. Fig. S1c), and the Photosynthetic Photon Flux

Density (PPFD) was set to 500 µmol m^{-2}s^{-1}.

Experimental design

We performed a full-factorial experiment involving ALAN treatment, two aphid treatments and four

destructive (independent) harvesting days (Fig. 1). We used 16 biological replicates for ALAN control,

20 for part-night ALAN and 20 for full night ALAN treatment combinations, respectively (totalling 448

plants). In this study, we used seven climate chambers that were assigned a specific light treatment,

and conducted the experiment in two separate runs, leading to 14 chambers (four assigned ALAN

control, and five part-night and full-night ALAN, respectively). ALAN assignments to chambers were

changed between runs (Supplementary Fig. S2) to control for climate chamber effects in our models.

Biological replicates were divided over chambers and runs so that each chamber had four biological

replicates of each combination (four replicates x eight combinations = 32 plants per chamber).

ALAN treatments – Each chamber was equipped with four linked LED modules each consisting of six

white diodes per module (D-15-01-001, 2 W, 12 V, www.leds24.com), coupled to a dimmer to

control illuminance. ALAN control chambers were kept dark at night, whereas part-night ALAN

chambers were illuminated between 00:00 and 04:00. Full night ALAN chambers were lit with ALAN

between 00:00 and 08:00. Illuminance was measured using a HR4000 spectrometer (using

OceanView v1.6.7) paired to an Ocean Optics Custom Fiber Configuration OCF-102313 equipped

with a cosine corrector with spectral diffusing material (Ocean Insight, Orlando, Florida, US) at pot

level and illuminance was set to 15–20 lux, which are levels commonly observed for LED street lights

(Bennie et al. 2016). Measurements with the same spectral meter underneath local LED street lights

showed highly fluctuating and noisy signals of 5–100 lux. Note that under the experimental

nocturnal light conditions, PPFD levels were below the measurable range of the fixed TUM*mesa* light

sensors, indicating fewer than 0.5 μmol photons m^{-2} s^{-1}. The spectral irradiance of the climate

chambers during the day and spectral irradiance of the LEDs under maximum output capacity and

under the dimmed ALAN treatment levels were measured and visualized after the experiment using

an array-radiometer (Newport OSM400 200–800 nm, Sony CCD-Sensor 2046px and 50 μm gap, 1 nm

spectral resolution, Newport Spectra Physics GmbH, Darmstadt, Germany). Supplementary Fig. S1

shows spectral irradiance for the LED modules under maximum capacity and under set ALAN

treatments, and a visualization of a typical daylight spectral irradiance of the used climate chambers.

Plant and aphid treatments – Barley plants were grown with and without aphids in single round

plastic pots (400 mL, 10 cm, Meyer, Göttingen, Germany). Pots were filled with soil, and watered

with 50 mL of tap water and then left to acclimatize at room temperature for four days, then

brought into the climate chambers, where they were left for another three days. Pots were checked

daily and watered (20 mL) when the soil surface was dry. After this week of acclimatization, two

surface-sterilized barley seeds (5% hypochlorite solution for 2 min, followed by a 1-min rinse with

tap water) were planted in 1–2 cm deep holes made in the middle of the pot. Four days after sowing,

most seedlings had germinated, and the smallest seedling was removed from each pot, so that only

one plant per pot remained. Plants were checked daily, and watered to need, three times a week.

One week after sowing, the height of each plant was recorded, and two pre-final (N4) instar

individuals of *S. avenae* 'Fescue' were added to all pots assigned to the aphid treatment. The plants

from both the aphid and no-aphid treatments were subsequently bagged around the pot rim with a

permeable cellophane bag (Kopp Verpackungen, Höheischweiler, Germany), which was secured with

elastic band to contain the aphids.

Temporal harvest – An *a priori* assigned subset of experimental plants from each combination was

destructively sampled 7-, 10-, 14- and 17-days post-aphid infestation. This allowed us to follow aphid

colony growth over time on independent plants. The destructive harvests were needed to sample

for fresh plant material for gene expression.

Harvest and response variables – On the harvest days, total numbers of aphids were recorded.

Among adult aphids, winged and unwinged aphids were distinguished. We measured plant height by

measuring the distance between soil surface and the tip of the longest leaf. Chlorophyll levels were

measured on the youngest fully expanded leaf at three points and recorded the average, using a

chlorophyll meter (Konica Minolta SPAD-502Plus, Tokyo, Japan). Plants were placed on a cart in the

evening, and the next morning, destructive sampling occurred between 06:00 and 07:00, before the

daylights turned on. This ensured that we assessed the effects of the ALAN treatments – which by

design occurred at night – on gene expression. Chambers were opened to remove plants for about

5–10 s. The harvest itself occurred in a dark corner of the corridor, with the only light coming from a

dim emergency light above a door. Shoots were harvested within 5 min of removing plants from

their chamber. To this end, the fresh shoots were clipped with sterilized scissors, folded in labelled

aluminium foil, and flash-frozen in liquid nitrogen. This was done for the first three replicates for

each treatment combination. We clipped and dried shoots of the remaining fourth replicate to

determine shoot biomass. Roots for all four replicates per treatment combination were

subsequently washed and dried to determine root biomass. Roots and shoots were oven-dried at 60

°C for at least three days before weighing.

Gene expression assay – In order to assess the effects of ALAN on plant-aphid interactions, we

performed a reverse transcriptase-quantitative PCR on mRNA extracted from the frozen plant leaf

material. For the gene expression assay we selected the plants harvested at day 10 post-infestation,

as these plants did not yet show signs of aphid stress or yellowing, and hence provided high-quality

samples for our gene expression assay. Before RNA extraction, three replicates from the same

climate chamber were pooled for each treatment combination. Shoots of the three samples were

ground and homogenized in liquid nitrogen. Effectively, the pooling resulted in 4–5 replicates for

each ALAN-aphid combination (28 samples total). Total RNA was extracted from the homogenized

sample using a QIAGEN RNeasy Plant Mini Kit (QIAGEN, Hilden, Germany), following the

manufacturer's protocol. RNA was converted to cDNA using SuperScriptII reverse transcriptase

(Invitrogen, Thermo Fisher, Waltham, Massachusetts, US).

We selected four markers that are associated with defence-related proteins. Two markers amplify

PR1 and *PR17b*, two genes that code for pathogenesis-related proteins, and are generally involved in

responses to biotrophic pathogens (Christensen et al. 2002; van Loon et al. 2006; Zhang et al. 2012),

and are known to be upregulated in response to aphids (Grönberg 2006; Delp et al. 2009). Two other

markers, *HvERF-like* and *HvWRKY22,* are associated with systemic defence against bacterial and

fungal pathogens in barley (Dey et al. 2014; Lenk et al. 2019). The cDNA preparation was followed by

qPCR on a 7500 Fast real-time qPCR system (Applied Biosystems, Thermo Fisher) with three

technical replicates to obtain average Ct values per gene and sample. The qPCR was performed with

the SensiMix SYBR low ROX kit (Bioline, Meridian Bioscience) using primers from Dey et al. (2014)

and Shrestha et al. (2019).

For further statistical analysis, we determined delta Ct (δCt) values by Ct_{target} - $Ct_{reference}$. Under the

assumption that the endogenous reference gene has stable and high expression in all samples,

indicated by a low $Ct_{reference}$, this generally results in a positive δCt value for a sample, where a higher

δCt indicates a lower expression. As we were interested in the relative expression of genes across

treatments, the δCt values were used for further statistical analysis. However, since the experiment

was replicated in two temporally separated runs, and this was an expected source of variation, we z-

transformed the δCt values per run by subtracting the *run mean* δCt from each *sample* δCt, and

divided this by the *standard deviation of the run mean* δCt, effectively taking out the run effect. The

resulting z-score represents an effect size as the distance between run mean δCt and the sample δCt

in units of standard deviation. As a positive δCt z-score indicates downregulation, which is visually

not intuitive, we inverted the z-score signs, so that higher and lower relative gene expression match

with the sign of the plotted values.

Statistical analysis

All analyses were performed in R version 3.6.3 (R Development Core Team, 2020). Linear mixed

models were performed using the package 'lme4' (Bates et al. 2015), and p-values were estimated

using type II Wald Chi-Square tests using car::Anova from the 'car' package (Fox & Weissberg 2019).

Data visualization was done using 'ggplot2' (Wickham 2016).

Plant variables – Plant variables were analysed using linear mixed models; as fixed factors we

included 'ALAN treatment', 'aphid treatment' and 'harvesting day'. We included 'run' as a covariate

in the model, and 'climate chamber' as a random intercept, to correct for potential climate chamber

effects.

Aphid variables – Aphid variables were analysed using generalized linear mixed models, with a

Poisson error distribution; as fixed factors we included 'ALAN treatment' and 'harvesting day'. We

included 'run' as a covariate in the model, and 'climate chamber' as a random intercept, to correct

for potential climate chamber effects. Only pots allocated to the aphid treatment were included in

the analyses.

Significant effects of ALAN were further analysed using post-hoc Tukey tests using the 'glht()'

command from the 'multcomp' package (Hothorn et al. 2008)

RT-qPCR - Gene expression data were analysed by means of z-transformed δCt values using linear

mixed models. As fixed factors we included 'ALAN treatment' and 'aphid treatment', and 'climate

chamber' as a random intercept, to correct for potential climate chamber effects.

We assessed relationships between gene expression and aphid numbers by averaging the aphid

numbers on the samples that were pooled for the gene expression assays. The relationship between

aphid numbers and inverted z-transformed δCt values was assessed by linear regression.

Results

From the total of 448 sowed barley plants in the study, 430 survived until their assigned harvest and

used for the analyses, of which 213 were assigned to aphid treatment, and 217 to no-aphid

treatment. Across all treatment combinations 14-20 replicates were included in the plant-level

analyses.

ALAN and temporal effects on aphid colony growth

Aphid numbers increased with time, i.e., harvest day (Table 1, Fig. 2a). The total number of aphids

per plant was significantly affected by ALAN (χ^2_2 = 103.7, p < 0.001, Fig. 2a). At the time of final

harvest, aphid numbers on plants exposed to full night ALAN were 2.8% higher (mean 89.7 aphids)

than those exposed to dark night controls (mean 87.4 aphids), whereas plants exposed to part-night

ALAN had about 19.7% lower aphid numbers (mean 70.1 aphids) than controls. Part-night ALAN

treatments had 21.9% lower aphid numbers than full-night ALAN treatments, and post-hoc Tukey

tests indicated significant differences between all ALAN levels (Fig. S3a). We also observed a

significant interaction between ALAN and harvest day (χ^2_6 = 31.2, p < 0.001, Fig. 2a), indicating that

ALAN treatments affected aphid colony growth differently. Specifically, both full night and part-night

ALAN treatments suppressed aphid numbers during the first ten days of infestation, after which the

aphids exposed to full ALAN reached numbers similar to aphids exposed to dark night controls,

whereas aphids exposed to part-night ALAN remained substantially lower throughout the

experimental duration (Fig. 2a). When only adult aphids (winged and unwinged combined) were

considered, numbers were affected by ALAN (χ^2_2 = 10.4, p = 0.005, Fig. 2b). At the time of harvest,

adult aphid numbers were 9.8 and 32.2% lower on plants exposed to full night (mean 8.4 aphids)

and part-night ALAN (mean 11.2 aphids) than on control plants exposed to dark nights (mean 12.4

aphids), respectively. Part-night ALAN treatments had 25.0 % lower adult aphid numbers than full

night ALAN. Post-hoc Tukey tests indicated that at harvest time only part-night ALAN treatment was

significantly lower from the two other levels (Fig. S3b). Harvest day also affected the number of

adult aphids in the population (Table 1), as the number of adult aphids in the population increased

over time (Fig. 2b). We observed no significant interaction between harvest day and ALAN treatment

in their effects on adult aphid numbers (Table 1).

ALAN, aphid and temporal effects on plant performance

Plant height and shoot dry biomass were only marginally (positively) affected by ALAN (plant height:

χ^2_2 = 4.7, p = 0.095, shoot dry biomass: χ^2_2 = 5.1, p = 0.078) and plants tended to be higher under full

night ALAN treatment (Fig. 3a,b), although post-hoc Tukey tests revealed no significant differences

between the three ALAN levels across all experimental plants. Root dry biomass was significantly

affected by ALAN (χ^2_2 = 6.9, p = 0.032, Fig. 3c), with post-hoc Tukey tests revealing that part-night

ALAN significantly decreased root size compared to full night ALAN, but not to controls, across all

experimental plants. Plant height and shoot biomass were not significantly affected by aphid

treatment (Table 1), but aphids marginally reduced root biomass (χ^2_1 = 3.71, p = 0.055, Fig. 3c).

Harvest day positively affected plant height, shoot and root dry biomass, as plants increased in size

during the experiment (Table 1, Fig. 3). We observed no significant interactive effects between ALAN

and herbivory on plant height, shoot or root biomass (Table 1). ALAN significantly affected

chlorophyll levels in the plants (χ^2_2 = 8.8, p = 0.017), with plants exposed to full night ALAN exhibiting

on average 9%, and plants exposed to part-night ALAN 4% higher chlorophyll levels than control

plants exposed to dark nights, across all plants (Fig. 3b). Aphid presence did not significantly affect

chlorophyll levels (Table 1), but these were significantly affected by harvest day, as levels decreased

when plants grew older (Fig. 3b).

ALAN and aphid effects on plant gene expression

The expression of some defence genes was marginally or significantly affected by the ALAN

treatments while the presence of aphids marginally or significantly increased expression of most

defence genes (Table 2). The expression of *HvWRKY22* was marginally affected by ALAN treatments

(χ^2_2 = 5.9, p = 0.053, Fig. 4a), with a trend for expression to be downregulated when plants were

exposed to full night or part-night ALAN treatments (Fig. 4a). The expression of *HvWRKY22* was

strongly upregulated in aphid treatments (χ^2_1 = 8.0, p = 0.005, Fig. 4a). No interactive effects were

observed on the expression of *HvWRKY22* (Table 2). The expression of *HvPR1* was marginally

upregulated by aphid treatments (χ^2_1 = 3.6, p = 0.057, Fig. 4b), but effects of ALAN treatments or

interactive effects on the expression of *HvPR1* were not observed (Table 2). The expression of

HvPR17b differed significantly between ALAN treatments (χ^2_2 = 10.1, p = 0.006, Fig. 4c) and was

marginally upregulated by aphid treatments (χ^2_1 = 3.1, p = 0.077, Fig. 4c). Post-hoc Tukey tests

indicated that expression of *HvPR17b* was significantly lower when plants were exposed to full night

or part-night ALAN, compared to ALAN controls across all plants. The expression of the *HvERF-like*

gene was not significantly affected by any of the treatments (Fig. 4d).

The total number of aphids on those plants used for qPCR analyses positively correlated with the

expression of *HvWRKY22* (R^2 = 0.31, $F_{1,12}$ = 5.5, p = 0.037; Fig. 5a), *HvPR1* (R^2 = 0.55, $F_{1,12}$ = 14.8, p =

0.002; Fig. 5b), and *HvPR17b* (R^2 =0.71; $F_{1,12}$ = 29.4, p < 0.001,; Fig. 5c), but not with the *HvERF-like*

gene (R^2 = 0.002, $F_{1,12}$ = 0.02, p = 0.891; Fig. 5d).

Discussion

In this study, we exposed plants and associated aphids to either full nights of ALAN, nights with a

part-night ALAN treatment, where ALAN was switched off halfway through the night, or controls

with completely dark nights. We found that plant growth was only minimally affected by ALAN,

whereas aphid colony growth and the number of adult aphids in the colony were impacted

significantly by ALAN treatments. Importantly, we found strong differences between full and part-

night ALAN treatments. Contrary to our hypotheses, ALAN mitigation – by limiting the duration of

light exposure at night – had stronger negative effects on aphid colony growth than nights with full

exposure to ALAN. Gene expression assays revealed that defence-associated gene expression was

affected by ALAN and aphid presence in three of four genes, but in these three genes, expression

positively correlated with aphid colony size, suggesting that the effects of ALAN on plant defences

are likely regulated through indirect effects of ALAN on plants via direct disruptive effects on aphid

colonies, rather than via regulation by the plant in direct response to ALAN.

In agreement with our first hypothesis, exposure to ALAN suppressed aphid colony growth, although

particularly in the full-night ALAN treatment, we observed a recovery of aphid numbers over time.

Previous studies show suppressive ALAN effects on plant–aphid interactions in other model systems,

where aphid suppression was associated with changes in plant phenology, and increases in aphid

parasitism by parasitoids (Bennie et al. 2015; Sanders et al. 2015; Bennie et al. 2018; Sanders et al.

2018). Importantly, our study shows that aphid colony decline also takes place without such external

confounding factors. Contrary to the present study, a previous field experiment with *H. vulgare* and

S. avenae detected no effects of ALAN on final aphid densities at nine weeks (Sanders et al. 2018),

indicating that the strongest effects may be early after colonization, and may disappear over time,

which is in line with our observations that aphid numbers recover in full night ALAN treatments.

However, it should be noted that the referenced study included parasitoids, which as aphid

antagonists may have affected aphid colony sizes. Other studies that have assessed ALAN effects on

aphid colony size over time have generally included repeated measures of the same colony (Bennie

et al. 2015; Sanders et al. 2015), which allows to assess individual colony responses, but also

elevates the risk of including colony-specific effects that persist over time. In our study, aphid colony

growth was assessed over time on independent samples which provides a higher temporal

resolution, and also reinforces the generality of ALAN impacts on aphid colony formation, especially

during early colonization.

An important question is what drives the ALAN-induced impacts on aphid numbers? – Two answers

are plausible: On the one hand, ALAN could influence aphids indirectly, via changes in plant

nutritional quality or quantity (plant defence compound and nutritional quality or plant size), which

commonly affects insect performance through bottom-up effects (Will & Van Bell 2006). On the

other hand, ALAN may directly impact aphids, through disturbance of natural behaviour such as

settling times that may limit feeding, and ultimately, fitness. Our data suggest that plant size was

only marginally affected by ALAN. Although various studies have suggested that resource availability

could be the main driver through which ALAN impacts insect responses (Sanders et al. 2015; Bennie

et al. 2015), plant size under levels of infestation observed in our study are unlikely to be limiting to

aphid population growth, as plants were large enough to sustain the relatively small aphid numbers.

It thus seems more plausible that potential bottom-up effects are mediated via plant quality, not

quantity. We observed a significant increase in chlorophyll levels – a proxy for plant health and

quality – when plants were exposed to ALAN, but as our data suggested only a weak and positive

relationship with aphid numbers, it is highly unlikely that chlorophyll levels alone are responsible for

aphid suppression under ALAN. Future research should focus on how ALAN may affect plant quality

for aphids, and especially the diurnal variation in the phloem composition (Douglas 2006), in order

to better understand the possible links between ALAN, the plant and ALAN-induced aphid

suppression.

We assessed the expression of four defence-associated genes in barley corresponding with different

defence pathways, to test whether exposure of plants to ALAN would result in a regulation of

specific pathways. We measured expression of two *PR* genes, which are commonly associated with

innate immunity in plants (van Loon et al. 2006; Wang et al. 2018). In particular, *HvPR1* and *HvPR17b*

are induced upon infection of barley with pathogens and are presumed to positively correlate with

defence (Christensen et al. 2002; Grönberg 2006; Delp et al. 2009; Zhang et al. 2012). Further,

HvWRKY22 and *HvERF-like* have previously been positively associated with SAR-like systemic defence

in barley (Dey et al. 2014). We observed a marginally significant effect (p = 0.053) of ALAN on the

expression of *HvWRKY22*, and a strongly significant effect of ALAN on the expression of *HvPR17b*.

We did not find any effect of ALAN or aphid treatments on the expression of *HvERF-like* gene

expression, suggesting that this gene, which is associated with systemic pathways (Dey et al. 2014),

is unlikely to respond to variation in light at night and is not involved in plant defences against

aphids.

Contrary to our expectation that ALAN would upregulate plant defences, ALAN decreased the

expression of *HvWRKY22* and *HvPR17b*, and we observed similar but non-significant patterns in

HvPR1. In fact, for *HvWRKY22*, *HvPR1* and *HvPR17b*, but not for *HvERF-like* gene, the gene

expression correlated tightly and positively with the aphid numbers present on the plant. This

suggests that patterns of expression differing between ALAN treatments were likely not the result of

direct ALAN effects on plant defence regulation, but instead more likely a result of direct disruptive

effects of ALAN on aphid feeding or settling behaviour and resultant suppression of colony growth,

and that this, in turn, is reflected by a reduced transcript accumulation of certain defence-associated

genes. Importantly, from our results, we cannot conclusively state that ALAN does not affect plant

defences. Our analyses focused on a limited number of genes that are involved in constitutive and

induced plant defence responses against aphids. A more holistic approach, for instance using meta-

transcriptomics of plants under ALAN or dark night treatments could reveal how ALAN may affect

transcriptional regulation of phytohormonal defence mediation, as well as other pathways involved,

for instance, in secondary metabolism.

There are various alternative plant physiological pathways through which ALAN might affect the

phytobiome (Heinen 2021). For instance, exposure to ALAN can disrupt stomatal activity, above-

belowground allocation, and the nocturnal starch turnover from above- to below-ground

compartments (Stitt & Zeeman, 2012; Kwak et al. 2017; 2018). Disruptions in starch metabolism may

result in changes in concentration and composition of sugars in phloem tissues, and as aphids are

directly affected by phloem sugar concentrations via the 'sugar barrier', which requires strong

osmoregulation in the aphid gut to excrete it (Douglas 2006), it is entirely plausible that such a

pathway might affect probing, feeding and reproductive success, leading to altered aphid colony

growth. As such, future studies should investigate ALAN effects on sugars, amino acids,

micronutrients and secondary metabolites in the phloem that are important for aphid performance.

For instance, changes in carbon metabolism over diurnal cycles can greatly impact interactions

between plants and their microbiome (Baraniya et al. 2018), and knowing how ALAN may impact on

carbon metabolism and affect interactions between plants and other organisms will be an exciting

direction for future research (Heinen 2021).

ALAN can also directly impact on aphid colony growth, via effects of light on aphid behaviour. Aphids

perceive light intensity and can respond to low light intensities by producing winged adults (Alkhedir,

Karlovsky & Vidal 2010). Aphids also express diurnal rhythmicity in locomotor and metabolic activity

(Beer et al. 2017), even independent of their plant hosts, which are strongly rhythmic (Joschinski et

al. 2016). Despite strong cyclic circadian clock gene expression, aphids continue the various stages of

their feeding cycles throughout the night, which is hypothesized to be an important element for

osmoregulation of high phloem sugar content (Douglas 2006; Nalam et al. 2021). Light has been

shown to disrupt aphid behaviour, and more strongly so during the night (Narayandas & Alyokhin

2006). This effect is likely stronger in plants with more open canopies, where light reaches the insect

directly. We speculate that ALAN led to disruptions of settling and feeding behaviour, lowering the

uptake of resources, which in turn explain smaller aphid colony sizes and number of matured adults,

and the tightly linked defence gene expression, as observed under our treatments. Further tests to

examine how ALAN impacts aphid feeding behaviour, should make use of electrical penetration

graph analyses (Tjallingii 1978), which record the probing behaviour of aphids over time, to compare

aphid behaviour under dark and lit nights.

An important finding is that contrary to our hypothesis, a mitigation of ALAN levels at night through

a midnight shutoff of the light source enhanced adverse ecological effects of ALAN on insects. A

popular belief is that the ecological side effects of light pollution could be mitigated by optimizing

lighting schemes to match human activity patterns, and turning off lights after peak activity.

However, our data suggest that after experiencing a short duration of ALAN, turning off the lights

has more detrimental effects than keeping them on. These findings are difficult to explain. One

possibility is that a midnight shutoff strategy may be experienced as a double ecological stressor.

First, the organisms experience a lit night, which may be seen as the first stressor, which is then

followed by a severely reduced dark recovery period, i.e., further stress, although the mechanism is

unclear. However, we argue that the ecological consequences of specific mitigation strategies need

to be studied before they are readily implemented, as despite good intentions, they may sometimes

have unexpected side effects (Cieraad et al. 2022).

Our study also highlights the physical limitations of ALAN research. One example is the typical choice

to use illuminance – expressed in lux, which is a photometric unit – in most ALAN research. This

means, that the radiometric quantity will be weighted with a visual function describing the

wavelength dependent response of the human eye. The photopic function of the human eye have

the maximum at 550 nm, while 380–780 nm is named as 'light' (Aphalo et al. 2012). However, light is

in the physical sense better described as radiant energy per time per effective receiver surface and

wavelength interval. This is where ALAN research becomes more problematic, as its spectral

irradiance generally falls below thresholds measurable by standard equipment. Another problematic

aspect is that in the lower ranges of spectral irradiance, the ratio of noise to signal is very high (cf.

Suppl. Fig. S1b). Measurements conducted under LED street lights in the field confirm low signals of

irradiance, with comparatively high levels of noise. A better physical characterization of ALAN

irradiance spectra, and its reach in nature is essential to understand the breadth of ecological

impacts, especially on sessile organisms such as plants.

Despite significant advances in the field, the effects of ALAN on plants and plant-associated

organisms are understudied (Heinen 2021). We aimed to generate a better understanding of how

ALAN impacts on aphid colony growth over time, as mediated via plant quality and quantity. In

general, ALAN impacts on barley plants appear to be minor, but regardless exposure to ALAN leads

to substantial effects on aphid colony growth, which appear to be direct disruptive effects of ALAN,

rather than plant-mediated effects. Our study especially highlights the difficulties associated with

mitigation of the ecological side effects of ALAN, as a reduction of the light intervals increased the

negative impact of ALAN on aphids. Rather little is known about how ALAN affects insect

communities on crops and wild plant species. These findings highlight the need to study the impacts

of ALAN in natural ecosystems, and we show that the solution to light pollution may be more

complex than the flip of a switch.

Acknowledgements

We thank Roman Meier for advice and engineering support in the TUM*mesa* phytochamber

facilities. We thank Hans Hausladen and Stefan Kimmelmann for advice and help in obtaining soil for

the pot experiment. We thank Sharon Zytynska for supporting the project. The experiment was

supported by DFG grant 397565003 WE 3081/36-1 awarded to Sharon Zytynska and WWW.

Figure legends:

Figure 1: Schematic representation of the experimental design. The experiment was a full factorial

combination of ALAN treatments, aphid treatment, temporal samplings, with replicates. ALAN

treatment had three levels (control, part-night ALAN, full-night ALAN), two aphid levels

(presence/absence), four destructive sampling dates, with 16–20 replicates each. In ALAN treatment

scheme, red bars indicate ALAN lights on, dark blue bars indicate ALAN lights off, yellow bars

indicate daylights on. The experiment was executed in seven climate chambers, over two

experimental runs. Climate chambers were assigned specific ALAN treatments, and biological

replicates were divided across assigned chambers and runs. ALAN controls had 16 biological

replicates, divided over four different chambers, part-night ALAN and full night ALAN treatments had

20 biological replicates, divided over five different chambers. Each climate chamber contained 32

plants that were used for this study (pictured). The experiment had a total of 448 plants.

Figure 2: Box plots depicting effects of ALAN treatments and sampling time on a) aphid colony size (n

= 16–20), and b) number of adult aphids within the population (n = 16–20). Box colours depict ALAN

treatments, with grey being control, yellow part-night ALAN, and green full-night ALAN. Boxes

represent median values with upper and lower quartiles, and whiskers represent 1.5x the

interquartile range; individual dots represent outliers. Simplified mixed model output is indicated in

the graphs, and detailed model output can be found in Table 1. Post-hoc Tukey tests were

performed on a subset of the data on the final harvest point, and visualized separately for the subset

in Fig. S3.

Figure 3: Box plots depicting effects of ALAN treatments and sampling time on a) plant height (n =

16–20), b) chlorophyll content (SPAD units) (n = 16–20), c) root (n = 16–20) and d) shoot dry biomass

(n = 4–5). Box colours depict ALAN treatments, with grey being control, yellow part-night ALAN, and

green full-night ALAN. Boxes represent median values with upper and lower quartiles, and whiskers

1.5x the interquartile range; individual dots are outliers. Simplified mixed model output is indicated

in the graphs, and detailed model output can be found in Table 1.

Figure 4: Box plots depicting effects of ALAN treatments on standardized gene expression of a)

HvWRKY22, b) *HvPR-1*, c) *HvPR-17b* and d) *HvERF-like gene* markers (each n = 4 (control) or 5

(treatment)). Box colours depict ALAN treatments, with grey being control, yellow part-night ALAN,

and green full-night ALAN. Boxes represent median values with upper and lower quartiles, and

whiskers represent 1.5x the interquartile range; individual dots represent outliers. Simplified mixed

model output is indicated in the graphs, and detailed model output can be found in Table 2.

Figure 5: Relationships between the average number of aphids on barley plants assigned to a pooled

sample, and the standardized gene expression of a) *HvWRKY22,* b) *HvPR-1*, c) *HvPR-17b* and d)

HvERF-like gene markers across ALAN treatments. Linear correlations with 95% confidence intervals

were fit if the linear relationship was significant.

Tables:

531 Table 1: Output of (generalized) linear mixed models testing the effects of Replicate Run, ALAN (A), herbivory (H) and time (T) on the number of total
532 aphids, the number of adult aphids, plant height (cm), shoot biomass (g), root biomass (g), chlorophyll levels (SPAD units). Presented are degrees of
533 freedom, Wald's Chi-square statistics and p-values, obtained from the models using the car::Anova command in R. Significant effects (p < 0.05) are marked
534 in bold, and marginally significant effects (0.05 < p < 0.1) in italics.

	Total aphids		Adult aphids		Plant height		Shoot dry mass		Root dry mass		Chlorophyll level	
	d.f.	χ^2 (p)	d.f.	χ^2 (p)	d.f.	χ^2 (p)	d.f.	χ^2 (p)	d.f.	χ^2 (p)	d.f.	χ^2 (p)
Run	1	**51.2 (<0.001)**	1	*2.9 (0.089)*	1	**63.3 (<0.001)**	1	**13.6 (<0.001)**	1	0.8 (0.383)	1	**36.0 (<0.001)**
ALAN (A)	2	**103.7 (<0.001)**	2	**10.4 (0.005)**	2	*4.7 (0.094)*	2	*5.1 (0.078)*	2	**6.9 (0.032)**	2	**8.1 (0.017)**
Herbivory (H)	-	-	-	-	1	0.6 (0.456)	1	2.1 (0.146)	1	*3.7 (0.055)*	1	2.6 (0.109)
Time (T)	3	**3166.4 (<0.001)**	3	**406.8 (<0.001)**	3	**123.8 (<0.001)**	3	**21.0 (<0.001)**	3	**10.7 (0.013)**	3	**73.7 (<0.001)**
A x H	-	-	-	-	2	1.8 (0.412)	2	0.2 (0.911)	2	2.4 (0.301)	2	1.4 (0.504)
A x T	6	**31.2 (<0.001)**	6	7.8 (0.254)	6	1.2 (0.974)	6	8.4 (0.210)	6	5.5 (0.487)	6	4.8 (0.565)
H x T	-	-	-	-	3	0.1 (0.995)	3	3.3 (0.351)	3	2.2 (0.525)	3	0.3 (0.957)
A x H x T	-	-	-	-	6	1.8 (0.934)	6	3.9 (0.697)	6	0.3 (0.999)	6	6.8 (0.338)

536   Table 2: Output of linear mixed models testing the effects of ALAN (A), herbivory (H) and time (T) on the standardized expression of *HvWRKY22, PR-1, PR-*
537   *17b,* and *HvERF-like* marker genes. Expression data was standardized by replicate run, which was therefore excluded from the model. Presented are
538   degrees of freedom, Wald's Chi-square statistics and p-values, obtained from the models using the car::Anova command in R. Significant effects (p < 0.05)
539   are marked in bold, and marginally significant effects (0.05 < p < 0.1) in italics.

	HvWRKY22		*PR-1*		*PR-17b*		*HvERF-like*	
	d.f.	χ^2 (p)	d.f.	χ^2 (p)	d.f.	χ^2 (p)	d.f.	χ^2 (p)
ALAN (A)	2	*5.9 (0.053)*	2	3.3 (0.192)	2	**10.1 (0.006)**	2	0.5 (0.779)
Herbivory (H)	1	**8.0 (0.005)**	1	*3.6 (0.057)*	1	*3.1 (0.077)*	1	0.0 (0.998)
A x H	2	3.5 (0.170)	2	2.4 (0.302)	2	1.9 (0.381)	2	1.3 (0.531)

540   **Figures:**

541   Figure 1:

543   Figure 2:

       Figure 3:

549   Figure 4:

553   Figure 5:

(a)

(c)

(b)

d)

References

Ahmad, M., Grancher, N., Heil, M., Black, R. C., Giovani, B., Galland, P., & Lardemer, D.

(2002). Action spectrum for cryptochrome-dependent hypocotyl growth inhibition in

Arabidopsis. Plant Physiology, *129*(2), 774-785.

Alkhedir, H., Karlovsky, P., & Vidal, S. (2010). Effect of light intensity on colour morph

formation and performance of the grain aphid *Sitobion avenae* F.(Homoptera: Aphididae). *Journal of*

Insect Physiology, 56(12), 1999-2005.

Aphalo, P. J., A., McLeod, A., Heikkilä, A., Gómez, I., López Figueroa, F., Robson, M. T., &

Strid, Å. (2012). Beyond the visible: a handbook of best practice in plant UV photobiology.

Helsingfors Universitet. p. 35-70.

Ayalon, I., de Barros Marangoni, L. F., Benichou, J. I., Avisar, D., & Levy, O. (2019). Red Sea

corals under Artificial Light Pollution at Night (ALAN) undergo oxidative stress and photosynthetic

impairment. *Global Change Biology*, 25(12), 4194-4207.

Ballare, C. L. (2009). Illuminated behaviour: phytochrome as a key regulator of light foraging

and plant anti-herbivore defence. *Plant, Cell & Environment*, 32(6), 713-725.

Ballaré, C. L. (2014). Light regulation of plant defense. *Annual Review of Plant Biology*, 65.

Ballaré, C. L., & Pierik, R. (2017). The shade-avoidance syndrome: Multiple signals and

ecological consequences. *Plant, Cell & Environment*, 40(11), 2530-2543.

Baraniya, D., Nannipieri, P., Kublik, S., Vestergaard, G., Schloter, M., & Schöler, A. (2018). The

impact of the diurnal cycle on the microbial transcriptome in the rhizosphere of barley. *Microbial*

ecology, 75(4), 830-833.

Bates, D., Maechler, M., Bolker, B., & Walker, S. (2015). Fitting Linear Mixed-Effects Models

Using lme4. Journal of Statistical Software, 67(1), 1-48.

Beer, K., Joschinski, J., Sastre, A. A., Krauss, J., & Helfrich-Förster, C. (2017). A damping

circadian clock drives weak oscillations in metabolism and locomotor activity of aphids

(*Acyrthosiphon pisum*). *Scientific Reports*, 7(1), 1-9.

Bennie, J., Davies, T. W., Cruse, D., & Gaston, K. J. (2016). Ecological effects of artificial light

at night on wild plants. *Journal of Ecology*, 104(3), 611-620.

Bennie, J., Davies, T.W., Cruse, D., Inger, R. & Gaston, K.J. (2015b) Cascading effects of

artificial light at night: resource-mediated control of herbivores in a grassland ecosystem.

Philosophical Transactions of the Royal Society of London B, 370, 20140131

Briggs, W. R., Beck, C. F., Cashmore, A. R., Christie, J. M., Hughes, J., Jarillo, J. A., ... & Okada,

K. (2001). The phototropin family of photoreceptors. *The Plant Cell*, 13(5), 993-997.

Briggs, W. R., & Christie, J. M. (2002). Phototropins 1 and 2: versatile plant blue-light

receptors. *Trends in Plant Science*, 7(5), 204-210.

Chory, J., Chatterjee, M., Cook, R. K., Elich, T., Fankhauser, C., Li, J., ... & Reed, J. (1996). From

seed germination to flowering, light controls plant development via the pigment phytochrome.

Proceedings of the National Academy of Sciences, 93(22), 12066-12071.

Christie, J.M. (2007) Phototropin blue-light receptors. *Annual Review of Plant Biology*, 58,

21–45.

Cieraad, E., Strange, E., Flink, M., Schrama, M., & Spoelstra, K. (2022). Artificial light at night

affects plant-herbivore interactions. *Journal of Applied Ecology*.

Day, J., Baker, J., Schofield, H., Mathews, F., & Gaston, K. J. (2015). Part-night lighting:

Implications for bat conservation. *Animal Conservation*, 18(6), 512-516.

Dey, S., Wenig, M., Langen, G., Sharma, S., Kugler, K. G., Knappe, C., ... & Vlot, A. C. (2014).

Bacteria-triggered systemic immunity in barley is associated with WRKY and ETHYLENE RESPONSIVE

FACTORs but not with salicylic acid. *Plant Physiology*, 166(4), 2133-2151.

Desouhant, E., Gomes, E., Mondy, N., & Amat, I. (2019). Mechanistic, ecological, and

evolutionary consequences of artificial light at night for insects: review and prospective.

Entomologia Experimentalis et Applicata, 167(1), 37-58.

Dominoni, D., Quetting, M., & Partecke, J. (2013). Artificial light at night advances avian

reproductive physiology. *Proceedings of the Royal Society B: Biological Sciences*, 280(1756),

20123017.

Donohue, K., Heschel, M. S., Chiang, G. C., Butler, C. M., & Barua, D. (2007). Phytochrome

mediates germination responses to multiple seasonal cues. *Plant, Cell & Environment*, 30(2), 202-

212.

Douglas, A. (2006). Phloem-sap feeding by animals: problems and solutions. *Journal of*

experimental botany, 57(4), 747-754.

Duarte, C., Quintanilla-Ahumada, D., Anguita, C., Manríquez, P. H., Widdicombe, S., Pulgar,

J., Silva-Rodríguez, E. A., Miranda, C., Manríquez, K., & Quijón, P. A. (2019). Artificial light pollution at

night (ALAN) disrupts the distribution and circadian rhythm of a sandy beach isopod. *Environmental*

Pollution, 248, 565-573.

Falcón, J., Torriglia, A., Attia, D., Viénot, F., Gronfier, C., Behar-Cohen, F., Martinsons, C., &

Hicks, D. (2020). Exposure to artificial light at night and the consequences for flora, fauna, and

ecosystems. *Frontiers in Neuroscience*, 1183.

Fox, J., & Weisberg, S. (2019). An {R} Companion to Applied Regression, Third Edition.

Thousand Oaks CA: Sage. URL: https://socialsciences.mcmaster.ca/jfox/Books/Companion/

Gaston, K. J. (2018). Lighting up the nighttime. *Science*, 362(6416), 744-746.

Gaston, K. J., Visser, M. E., & Hölker, F. (2015). The biological impacts of artificial light at
night: the research challenge. *Philosophical Transactions of the Royal Society B,* 370(1667),
20140133.

van Geffen, K. G., van Grunsven, R. H., van Ruijven, J., Berendse, F., & Veenendaal, E. M.
(2014). Artificial light at night causes diapause inhibition and sex-specific life history changes in a
moth. *Ecology and Evolution*, 4(11), 2082-2089.

Grubisic, M., van Grunsven, R. H., Manfrin, A., Monaghan, M. T., & Hölker, F. (2018). A
transition to white LED increases ecological impacts of nocturnal illumination on aquatic primary
producers in a lowland agricultural drainage ditch. *Environmental Pollution*, 240, 630-638.

Grubisic, M., Haim, A., Bhusal, P., Dominoni, D. M., Gabriel, K., Jechow, A., Kupprat, F.,
Lerner, A., Marchant, P., Riley, W., Stebelova, K., van Grunsven, R. H. A., Zeman, M., Zubidat, A. E. &
Hölker, F. (2019). Light pollution, circadian photoreception, and melatonin in vertebrates.
Sustainability, 11(22), 6400.

Hothorn T, Bretz F, Westfall P (2008). "Simultaneous Inference in General Parametric
Models." Biometrical Journal, 50(3), 346–363.

Howe, G. T., Gardner, G., Hackett, W. P., & Furnier, G. R. (1996). Phytochrome control of
short-day-induced bud set in black cottonwood. *Physiologia Plantarum*, 97(1), 95-103.

Joschinski, J., Beer, K., Helfrich-Förster, C., & Krauss, J. (2016). Pea aphids (Hemiptera:
Aphididae) have diurnal rhythms when raised independently of a host plant. *Journal of Insect*
Science, 16(1).

International Darksky Association (2020). LED Practical Guide. Tucson, AZ, USA
https://www.darksky.org/our-work/lighting/lighting-for-citizens/led-guide/

Kami, C., Lorrain, S., Hornitschek, P., & Fankhauser, C. (2010). Light-regulated plant growth
and development. In *Current Topics in Developmental Biology* (Vol. 91, pp. 29-66). Academic Press.

Kimura, M., & Kagawa, T. (2006). Phototropin and light-signaling in phototropism. *Current*

Opinion in Plant Biology, 9(5), 503-508.

Kraiselburd, I., Moyano, L., Carrau, A., Tano, J., & Orellano, E. G. (2017). Bacterial

photosensory proteins and their role in plant–pathogen interactions. *Photochemistry and*

Photobiology, 93(3), 666-674.

Kyba, C. C. M., Ruhtz, T., Fischer, J., & Hölker, F. (2011). Cloud coverage acts as an amplifier

for ecological light pollution in urban ecosystems. *PloS one*, 6(3), e17307.

Kyba, C. C. M., Kuester, T., Sánchez de Miguel, A., Baugh, K., Jechow, A., Hölker, F., Bennie, J.,

Elvidge, C. J., Gaston, K. J. & Guanter, L. (2017). Artificially lit surface of Earth at night increasing in

radiance and extent. *Science Advances*, 3(11), e1701528.

Kwak, M. J., Lee, S. H., Khaine, I., Je, S. M., Lee, T. Y., You, H. N., ... & Woo, S. Y. (2017).

Stomatal movements depend on interactions between external night light cue and internal signals

activated by rhythmic starch turnover and abscisic acid (ABA) levels at dawn and dusk. *Acta*

Physiologiae Plantarum, 39(8), 1-12.

Kwak, M. J., Je, S. M., Cheng, H. C., Seo, S. M., Park, J. H., Baek, S. G., ... & Woo, S. Y. (2018).

Night light-adaptation strategies for photosynthetic apparatus in yellow-poplar (*Liriodendron*

tulipifera L.) exposed to artificial night lighting. *Forests*, 9(2), 74.

Lenk, M., Wenig, M., Bauer, K., Hug, F., Knappe, C., Lange, B., Timsy, Häußler, F., Mengel, F.,

Dey, S., Schäffner, A., & Vlot, A. C. (2019). Pipecolic acid is induced in barley upon infection and

triggers immune responses associated with elevated nitric oxide accumulation. *Molecular Plant-*

Microbe Interactions, 32(10), 1303-1313.

Matzke, E. B. (1936). The effect of street lights in delaying leaf-fall in certain trees. *American*

Journal of Botany, 446-452.

Mittler, R. (2002). Oxidative stress, antioxidants and stress tolerance. *Trends in Plant Science*,

7(9), 405-410.

Moreno, J. E., Tao, Y., Chory, J., & Ballaré, C. L. (2009). Ecological modulation of plant

defense via phytochrome control of jasmonate sensitivity. *Proceedings of the National Academy of*

Sciences, 106(12), 4935-4940.

Nalam, V., Isaacs, T., Moh, S., Kansman, J., Finke, D., Albrecht, T., & Nachappa, P. (2021).

Diurnal feeding as a potential mechanism of osmoregulation in aphids. *Insect Science*, 28(2), 521-

532.

Narayandas, G. K., & Alyokhin, A. V. (2006). Diurnal patterns in host finding by potato aphids,

Macrosiphum euphorbiae (Homoptera: Aphididae). *Journal of Insect Behavior*, 19(3), 347.

Perkin, E. K., Hölker, F., Richardson, J. S., Sadler, J. P., Wolter, C., & Tockner, K. (2011). The

influence of artificial light on stream and riparian ecosystems: questions, challenges, and

perspectives. *Ecosphere*, 2(11), 1-16.

Poppe, C., Sweere, U., Drumm-Herrel, H., & Schäfer, E. (1998). The blue light receptor

cryptochrome 1 can act independently of phytochrome A and B in *Arabidopsis thaliana*. *The Plant*

Journal, 16(4), 465-471.

R Core Team (2020). R: A language and environment for statistical computing. R Foundation

for Statistical Computing, Vienna, Austria. URL https://www.R-project.org/

Rich, C., & Longcore, T. (Eds.). (2013). *Ecological consequences of artificial night lighting*.

Island Press.

Roden, L. C., & Ingle, R. A. (2009). Lights, rhythms, infection: the role of light and the

circadian clock in determining the outcome of plant–pathogen interactions. *The Plant Cell*, 21(9),

2546-2552.

Roy, J., Rineau, F., De Boeck, H. J., Nijs, I., Pütz, T., Abiven, S., et al. (2021). Ecotrons:

powerful and versatile ecosystem analysers for ecology, agronomy and environmental science.

Global Change Biology, 27(7), 1387-1407.

Sanders, D., Kehoe, R., Cruse, D., van Veen, F. F., & Gaston, K. J. (2018). Low levels of

artificial light at night strengthen top-down control in insect food web. *Current Biology*, 28(15),

2474-2478.

Schulz, E.-D., Beck, E., Buchmann, N., Clemens, S., Müller-Hohenstein, K., & Scherer-

Lorenzen, M. (2019). *Plant Ecology, Second Edition*. Springer.

Senzaki, M., Barber, J. R., Phillips, J. N., Carter, N. H., Cooper, C. B., Ditmer, M. A., Fristrup, K.

M., McClure, C. J. W., Mennitt, D. J., Tyrrell, L. P., Vukomanovic, J., Wilson, A. A., & Francis, C. D.

(2020). Sensory pollutants alter bird phenology and fitness across a continent. *Nature*, 587(7835),

605-609.

Shinomura, T., Nagatani, A., Hanzawa, H., Kubota, M., Watanabe, M., & Furuya, M. (1996).

Action spectra for phytochrome A-and B-specific photoinduction of seed germination in *Arabidopsis*

thaliana. *Proceedings of the National Academy of Sciences*, *93*(15), 8129-8133.

Shrestha, A., Elhady, A., Adss, S., Wehner, G., Böttcher, C., Heuer, H., Ordon, F., & Schikora,

A. (2019). Genetic differences in barley govern the responsiveness to N-acyl homoserine lactone.

Phytobiomes Journal, 3(3), 191-202.

Smith, H. L., McAusland, L., & Murchie, E. H. (2017). Don't ignore the green light: exploring

diverse roles in plant processes. *Journal of Experimental Botany*, 68(9), 2099-2110.

Somers, D. E., Devlin, P. F., & Kay, S. A. (1998). Phytochromes and cryptochromes in the

entrainment of the *Arabidopsis* circadian clock. Science, 282(5393), 1488-1490.

Stone, E. L., Harris, S., & Jones, G. (2015). Impacts of artificial lighting on bats: a review of

challenges and solutions. *Mammalian Biology*, 80(3), 213-219.

Tjallingii, W. F. (1978). Electronic recording of penetration behaviour by aphids. *Entomologia*

Experimentalis et Applicata, 24(3), 721-730.

Wang, X. D., Bi, W. S., Jing, G., Yu, X. M., Wang, H. Y., & Liu, D. Q. (2018). Systemic acquired

resistance, NPR1, and pathogenesis-related genes in wheat and barley. *Journal of Integrative*

Agriculture, 17(11), 2468-2477.

H. Wickham. ggplot2: Elegant Graphics for Data Analysis. Springer-Verlag New York, 2016.

Will, T., & van Bel, A. J. (2006). Physical and chemical interactions between aphids and

plants. *Journal of experimental botany*, 57(4), 729-737.

Wit, M. de, Spoel, S. H., Sanchez-Perez, G. F., Gommers, C. M., Pieterse, C. M., Voesenek, L.

A., & Pierik, R. (2013). Perception of low red: far-red ratio compromises both salicylic acid-and

jasmonic acid-dependent pathogen defences in *Arabidopsis*. *The Plant Journal*, 75(1), 90-103.

Wu, M. C., Hou, C. Y., Jiang, C. M., Wang, Y. T., Wang, C. Y., Chen, H. H., & Chang, H. M.

(2007). A novel approach of LED light radiation improves the antioxidant activity of pea seedlings.

Food Chemistry, 101(4), 1753-1758.

Yang, Y. X., Wang, M. M., Yin, Y. L., Onac, E., Zhou, G. F., Peng, S., ... & Zhou, Y. H. (2015a).

RNA-seq analysis reveals the role of red light in resistance against *Pseudomonas syringae* pv. *tomato*

DC3000 in tomato plants. *BMC Genomics*, 16(1), 120.

Yang, Y. X., Wang, M. M., Ren, Y., Onac, E., Zhou, G., Peng, S., ... & Yu, J. Q. (2015b). Light-

induced systemic resistance in tomato plants against root-knot nematode *Meloidogyne incognita*.

Plant Growth Regulation, 76(2), 167-175.