Modelling the distribution of rare and data-poor diadromous fish at sea for protected area management

Sophie A. M. Elliott[1, 4, 6*], Anthony Acou[1,2], Laurent Beaulaton[1,5], Jérôme Guitton[4], Elodie Réveillac[3], Etienne Rivot[1,4]

[1] Management of Diadromous Fish in their Environment OFB-INRAE-Institut Agro-UPPA, 35042 Rennes, France
[2] UMS OFB-CNRS-MNHN PatriNat, Station marine du Museum National d'Histoire Naturelle, 35800 Dinard, France
[3] Littoral, Environnement et Sociétés (LIENSs), UMR 7266, La Rochelle Université-CNRS, 17000 La Rochelle, France
[4] DECOD (Ecosystem Dynamics and Sustainability), L'Institut Agro, Ifremer, INRAE, Rennes, France
[5] Service Conservation et Gestion Durable des Espèces Exploitées, OFB, DRAS, 35042 Rennes, France
[6] Game & Wildlife Conservation Trust, Salmon & Trout Research Centre, East Stoke, Wareham BH20 6BB, UK

*Corresponding author: E-mail address: SElliott@gwct.org.uk (S.A.M. Elliott)

ABSTRACT
Anthropogenic pressures have resulted in declines in diadromous fish. Many diadromous fish which were commercially important are now threatened and protected. Little is known about their marine life history phases, and no observation-based Species Distribution Model exists for this group of species at sea. Yet, fisheries dependent and independent data could provide new insights into the distribution of diadromous fish at sea.

We collated a database of 168 904 hauls from fisheries observer bycatch data and scientific fisheries surveys, from eastern Atlantic and Mediterranean waters. The distribution of eleven rare and data-poor diadromous fish (shads, lampreys, salmonids, the European eel, the thinlip mullet, smelt and the European flounder) were modelled. A Bayesian site occupancy model, that incorporates imperfect detection to account for repeat detections and non-detections, the non-random nature of fishing gear type and spatial autocorrelation was used. From the model outputs, we explored bycatch risk and the role of MPAs, required under the Marine Strategy Framework Directive and Habitat Directive and assessed.

Diadromous fish were observed within relatively shallow coastal areas. Species specific gear bycatch trends were observed. Core distribution areas corresponded to their known water basin presence, indicating connectivity with their freshwater habitats. Numerous Habitat Directive Marine Protected Areas were found to be of relevance.

Given the coastal distribution of these species, they are exposed to higher anthropogenic pressures from both terrestrial and marine environments. Risk of bycatch at sea for most species appears to be low. Nonetheless, for threatened individuals, even a small amount of bycatch may impact their populations, especially since misreporting is likely to be high. Differences in catchability between gears highlight potential benefits of limiting access of certain gears within protected areas to reduce bycatch.

1. Introduction

Diadromous fish undertake long distance migrations between freshwater and marine ecosystems during their varying life history stages (Lassalle et al., 2008; Legrand et al., 2020; Limburg and Waldman, 2009; McDowall, 2009). They are particularly vulnerable since they are subject to both terrestrial and freshwater pressures (e.g., terrestrial run-off, pollution, habitat destruction, barriers to migration, climate change, fishing, etc.) (Costa et al., 2021; Limburg and Waldman, 2009; Merg et al., 2020; Verhelst et al., 2021). Furthermore, unlike many marine fish that are geographically widespread, numerous anadromous fish (e.g., shad and salmonids) form river-specific populations that are more susceptible to extinction (Limburg and Waldman, 2009; McDowall, 2009). As a result, most diadromous species native to the northern Atlantic Ocean have declined in abundance by at least 90% since the end of the late 19th century (Drouineau et al., 2018; Limburg and Waldman, 2009; Waldman and Quinn, 2022).

To try and halt the loss of diadromous fish populations, various International Agreements (e.g., Bern Convention, Convention of Migratory Species) and legislations (e.g., EU Habitat Directive, EU Marine Strategy Framework Directive (MSFD)) have been enacted (Table 1). Measures implemented focus principally on freshwater or estuarine habitats, such as restoration of river continuity as required by the EU Water Framework Directive (WFD, 2000/60/EC) and freshwater Spatial Areas of Conservation under the Habitat Directive (HD, 92/43/EEC). However, most diadromous species remain threatened at a national level (www.nationalredlist.org; Table 1), and the improvements have yet to be attained throughout their range (Verhelst et al., 2021; Waldman and Quinn, 2022; Wilson and Veneranta, 2019). The marine life history stages and at-sea distribution of diadromous fish, largely remain a black box (Elliott et al., 2021), causing difficulty in evaluating their conservation status required under the various legislations that protect these species (Table 1; Wilson and Veneranta, 2019). Furthermore, few Marine Protected Areas (MPAs) have been designated to protect them (e.g., https://mpa.ospar.org; https://natura2000.eea.europa.eu), and at-sea bycatch is not well understood, even if it is suspected that it could contribute to significant mortality (Kappel, 2005; Stratoudakis et al., 2016; Verhelst et al., 2021; Wilson and Veneranta, 2019).

Under the EU data collection framework, Member States are required to collect fisheries bycatch data through an onboard observer program (Cornou et al., 2015). Fisheries data is a rich set of information which provides year-round catch information and can be a valuable source of information for data-deficient species (e.g., Baum et al., 2003; Bisch et al., 2022; Elliott et al., 2020b). Bycatch data for protected species can be difficult to access, therefore fisheries observer data could help understand the distribution and bycatch mortality of data-poor species and meet EU Habitat Directive and MSFD (92/43/EEC; 2008/56/EC) requirements. Use of fisheries data does, however, require biases from the different gear types, the targeted nature of fishing, and un-balanced sampling to be taken into consideration (e.g. Alglave et al., 2022; Bourdaud et al., 2017).

Species Distribution Models (SDMs) provide a means to 'fill in the gaps' and provide complete coverage maps on which to base conservation and management decisions (Guisan and Thuiller, 2005; Guisan and Zimmermann, 2000; Leathwick et al.,

2005). Modelling the distribution of diadromous fish could help improve protection
measures for threatened and data-poor (insufficient biological information to
determine the current exploitation status (Berkson and Thorson, 2015; Prince and
Hordyk, 2019)) diadromous fish within existing MPAs. SDMs could also be used to
meet Descriptor 1 (Biodiversity – habitat extent) requirements under the MSFD
(2008/56/EC). However, building diadromous fish at sea distribution models
remains challenging because most of these fish are either IUCN red listed at a
national level (www.nationalredlist.org) or data-poor (Limburg and Waldman, 2009;
Merg et al., 2020). Modelling the distribution of data-poor and rare species, can
lead to underestimation of their true distribution due to imperfect detection
(uncertainty in the presence of a species within a site where the data was
recorded) (Guillera-Arroita, 2017; MacKenzie et al., 2002).

To model rare and data-poor species' distribution, specific statistical methods are
required to combine several sources of data that may not be initially designed to
study the species distribution (Engler et al., 2004; Lomba et al., 2010; Simmonds et
al., 2020). Using such data requires correction for detection bias arising from
observer error, species rarity and variability in environmental conditions (Guillera-
Arroita, 2017; Kellner and Swihart, 2014; MacKenzie et al., 2002) to help determine
sites where the probability of presence of species is higher (Belmont et al., 2022).
The low detectability of rare species can also result in a high proportion of false
absences (Guillera-Arroita, 2017). Spatial and temporal variation in observation
effort is another important source of heterogeneity in the observations (Guillera-
Arroita, 2017; Kellner and Swihart, 2014; Meyer et al., 2011). Integrated
hierarchical statistical model for species distributions have a great potential to
combine multiple sources of data to infer a single latent field of abundance or
presence/absence, and to enhance rare species distribution models for
conservation and management purposes (Engler et al., 2004; Lomba et al., 2010;
Simmonds et al., 2020). Another common problem which can arise when modelling
species distribution is that data can be autocorrelated (more similar when closer
together). Residual spatial autocorrelation can arise from population demographic
processes or the influence of unobserved variables. Ignoring spatial autocorrelation
can lead to inaccurate parameter estimations (Latimer et al., 2006). Developing
models that explicitly account for spatial autocorrelation in the latent field of
presence/absence is, therefore, necessary for accurate inferences on the
distribution.

Given the lack of knowledge on the distribution of most diadromous fish during
their marine life history stages, and the need to ensure they are sufficiently
protected, our objective was to model the current distribution (2003-2019) of
diadromous fish at sea. In addition, we wanted to quantify the risk of bycatch from
different gear types and assess the value of Habitat Directive MPAs for their
conservation. Since at sea targeted surveys do not exist for these species, we
collated a database of 168 904 hauls from fisheries dependent (French fisheries
bycatch observer data) and independent data (scientific survey data undertaken to
assess fish populations) in eastern Atlantic and French Mediterranean waters. A
Bayesian site-occupancy model, taking into account spatial autocorrelation in the
probability of presence, and imperfect detection through gear type, was used for
these data-poor species (Dormann et al., 2007; Latimer et al., 2006; MacKenzie et
al., 2002; Moriarty et al., 2020). From the distribution models we wanted to

answer the following questions: 1) Is the distribution of diadromous fish at sea
connected to existing knowledge on their freshwater habitats? 2) Do different gears
have different catchability, and does this provide information on the risk of
bycatch of diadromous fishes? 3) Could existing Habitat Directive MPAs protect
diadromous fish if specific measures are implemented?

Table 1. Diadromous fish observed within north-eastern Atlantic waters and their IUCN conservation status. A = Anadromous, C =
Catadromous, - = not listed, CR = Critical, LC = Least Concern, VU = Vulnerable. Demersal species are found near the seabed,
pelagic species occur in the open sea, demerso-pelagic species are species that migrate between these two water column zones,
host dependent refers to lampreys that are parasitic species during their marine phase. ✓ = protected under the specific
convention or Directive, letters specify the specific appendix the species is protected under.

Latin name	Common name	Type	Water column zone[1]	EU IUCN[3]	International conventions and legislation							
					CITES	WFD	HD	MSFD	Bern	Bonn	OSPAR	Barcelona
Acipenser sturio	Atlantic sturgeon	A	Demersal	CR	AI	✓	II, IV	✓	III	I, II	✓	✓
Alosa alosa	Allis shad	A	Pelagic	LC	-	✓	II, V	✓	III	-	✓	-
Alosa fallax	Twait shad	A	Pelagic	LC	-	✓	II, V	✓	III	-	-	-
Alosa agone	Mediterranean twaite shad	A	Pelagic	LC	-		II, V	✓	III	-	-	✓
Anguilla anguilla	European eel	C	Demerso-pelagic	CR	AII	✓	-	✓	-	-	✓	✓
Lampetra fluviatilis	River lamprey	A	Host dependent	LC	NA	✓	II, V	✓	III	-	-	✓
Petromyzon marinus	Sea lamprey	A	Host dependent	LC	-	✓	II	✓	III	-	✓	✓
*Chelon ramada**	Thinlip mullet	C	Demersal[2]	LC	-	✓	-	-	-	-	-	-
Osmerus eperlanus	Smelt	A	Pelagic	LC	-		-	-	-	-	-	-
Platichthys flesus	European flounder	C	Demersal	LC	-		-	-	-	-	-	-
Salmo salar	Atlantic salmon	A	Pelagic	VU[4]	-	✓	II, V	✓	III	-	✓	-
Salmo trutta	Sea trout	A	Pelagic	LC	-	✓	-		-	-	-	-

171   [1] Elliott and Dewailly, 1995; [2] Almeida, 1996; [3] Freyhof and Brooks, 2011; [4] Nieto et al., 2015 *previously known as *Lisa ramada*.

2. Method

2.1. Combining fisheries independent and dependent surveys

Fisheries dependent and independent data from 1965 to 2019 were collated within eastern Atlantic waters (Greater North Sea, Celtic Sea, Bay of Biscay and the Iberian coast) and French Mediterranean waters). Scientific bottom trawl surveys were extracted from the International Council for the Exploration of the Sea (ICES) Database of Trawl Surveys (DATRAS) portal (https://www.ices.dk/data/data-portals/Pages/DATRAS.aspx; Refer to Elliott et al., 2021 Table S1; Data in brief Table S1 - S2) and French Metropolitan scientific surveys not submitted to ICES DATRAS were obtained from the Institut Français de Recherche pour l'Exploitation de la Mer (IFREMER) (https://campagnes.flotteoceanographique.fr/campaign). Fisheries-dependent data came from French fisheries observer data (ObsMer data; Cornou et al., 2015) that began in 2003. According to the sampling plan, fisheries observers sample fishing vessels and fishing operations when on board (Fauconnet et al., 2015). ObsMer data is held by IFREMER and available on request from the French Ministry of Fisheries and Aquaculture (Direction des Pêches Maritimes et de l'Aquaculture; Refer to Elliott et al, 2021 Table S1; Data in brief Table S1).

From the database, diadromous fish presence and absence, length, gear type, spatial location and year and month of capture were extracted (Fig. 1; Fig. S1). Due to missing data and insufficient information on the length of hauls, size of vessels, mesh size, etc., effort was not possible to calculate. Details of gear types and surveys that were used for the analysis can be found within the supplementary materials of Elliott et al., (2021).

Prior to model analysis significant data cleaning was undertaken to standardise the surveys and extract relevant data to perform statistical analysis (e.g., converting gear types to gear categories, converting alike variables to the same units, etc.; see details within the Data in brief; Appendix A). Data from 2003 were used for spatial analysis, since a large proportion (74%) of the data came from the ObsMer dataset which started in 2003. In addition, prior to this data few presences were observed.

Since the proportion of zeros to presences was very uneven, which is not ideal for predictive modelling (Fielding and Bell, 1997), all surveys, gear types, target species (the intended catch for a particular fishery - from the ObsMer dataset) and ICES statistical divisions (Fig. 1) without diadromous fish presence, were deleted before modelling (Fig. S2 contains a spatial map of all hauls per gear type). Depths outside the ranges the species were observed were also removed. To avoid targeted fishing bias, all species that were targeted were removed. The latter included 2 European eels, *Anguilla anguilla*, 33 thinlip mullets, *Chelon ramada* and 41 flat fish which may have included European flounders, *Platichthys flesus* presences. All species were modelled other than the European sturgeon, *Acipenser sturio* because of too few presences (11 presences from 2003 – 2019). Note, for *L. fluviatilis*, one individual was caught by Set Longline within the Bay of Biscay. Due to its isolated nature of this capture, this caused problems when modelling (over detection of the line caught individual). Line gear type was therefore removed from the hierarchical modelling process.

2.2. Environmental predictor variables

Six environmental variables were considered as potential predictors of the presence of diadromous fish (seabed depth; distance from coast; sediment type; salinity; net primary production; and sea surface temperature; Table 2; Fig. S3). Depth and distance from the coast are thought to indicate key diadromous fish migratory periods (Taverny et al., 2012; Taverny and Elie, 2001; Trancart et al., 2014). Salinity and temperature are known to have direct physiological effects on diadromous fish, and their changes can indicate migration timing (Arevalo et al., 2020; Trancart et al., 2014). Net primary production is the gross primary production by autotrophs over the rate at which they respire, which is a measure of marine ecosystem functioning (ERSEM, 2020). Sediment type can be considered a proxy for food availability and shelter (Elliott et al., 2016; Trancart et al., 2014). As of result of too few presences and less information on length than the presence of individuals, distribution changes in length at stage, seasonal life history stage migration, and changes in habitat occupancy over the years were not considered.

To select the model of best fit, variables that were not collinear (Pearson's correlation coefficient< 0.5 or >-0.5 and a Variance Inflation Factor < 2) (Dormann et al., 2013) were kept. Depth and distance from coast, and sea surface temperature and net primary production were collinear. Since most species occupied a wide latitudinal range and seasonal or yearly temperature effects were not modelled, net primary production was used instead of sea surface temperature. To identify the model of best fit a backwards stepwise model selection process was undertaken, ensuring distance from coast and depth were not in the same model. The model with the lowest deviance information criterion, which includes a penalty factor for the number of parameters, was selected (Spiegelhalter et al., 2002).

Table 2. Environmental predictors used to analyse diadromous fish distribution. For all species except *Alosa agone,* variables were extracted at a 0.1 x 0.1 decimal degree resolution. *A. agone* environmental variables were extracted at a 0.04 decimal degree resolution.

Variable	Unit	Description / Source
Depth	Meters (m)	Bathymetry data downloaded from National Aeronautics and Space Administration Earth Observations project (http://neo.sci.gsfc.nasa.gov) with a 0.1 x 0.1 grid cell resolution by using the General Bathymetric Chart of the Oceans grid.
Distance to coast	Kilometres (km)	Calculated from a continent shape file
Sediment	Categorical	Broad-scale seabed habitat types (EUSeaMap) downloaded from EMODnet. Four classes were used according to grain size: mud (≤2mm), sand (> 2 and≤4 mm), coarse grain (> 4 mm and 64 mm), and rock (> 64 mm).
Salinity	Practical Salinity Units (PSU%)	Modelled sea surface salinity at 0.1 x 0.1 decimal degree resolution downloaded from Copernicus (ERSEM, 2020).

		Modelled at 0.04 x 0.04 decimal degree resolution from Copernicus using Teruzzi et al., (2021).
Net primary production	Milligrams of carbon per meters squared (mg C/m^2)	Modelled net primary production at 0.1 x 0.1 decimal degree resolution downloaded from Copernicus (ERSEM, 2020).
		Modelled at 0.04 x 0.04 decimal degree resolution from Copernicus using Teruzzi et al., (2021).
Sea surface temperature	degrees centigrade (°C)	Modelled Sea surface temperature at 0.1 x 0.1 decimal degree resolution downloaded from Copernicus (ERSEM, 2020).
		Modelled at 0.04 x 0.04 decimal degree resolution from Copernicus using Teruzzi et al., (2021).

254
255 **Fig. 1**. Diadromous fish presence (blue dots) and absence (light grey dots) data
256 used to model their distribution (2003-2019). Dark grey lines indicate International
257 Council for the Exploration of the Sea statistical divisions.

258
2.3. Statistical analysis
A Bayesian hierarchical site occupancy Gaussian intrinsic Conditional Autoregressive
(iCAR) model was built independently for each species. Hierarchical models have
the advantages of separating the ecological process (here, the habitat suitability
model, that is the spatial distribution of the probability of presence/absence,
including spatial autocorrelation) from the observation process (here, the
imperfect detection and its variability among gears or other factors) (Dormann et
al., 2007; Isaac et al., 2020; Latimer et al., 2006; MacKenzie et al., 2002). The
fundamental concept of the model is that the true presence/absence spatial field
(including spatial autocorrelation) is modelled on a grid, each grid cell being
associated with multiple observation events with imperfect detection, with
detectability that may vary depending upon the fishing gear.
271
272   The site occupancy model integrates two processes:
273
i)   Latent ecological process (habitat suitability)
The latent field of probability of presence is modelled on a grid as a function of
environmental predictors and by explicitly accounting for spatial autocorrelation:
$logit\left(\theta_j\right) = X_j\beta + P_j$ (1)
with θ_j the latent probability of presence (the habitat suitability) within grid cell
cell j, modelled in the logit scale as a function of environmental predictors X_j, with
β a vector of fixed effect describing how much the environmental predictor
contribute to the suitability process. The spatial autocorrelation P_j is modelled at
the scale of the grid cell as an iCAR random effect (2):
iCAR: $P_j \sim Normal\left(\mu_j, \frac{V_p}{n_j}\right)$ (2)

with μ_j the mean of the random effects calculated over the n_j cells considered in
the neighbour of cell j (n_j is the number of the neighbours which is 8 in most cases,
and <8 when the cell is on the boundary of the spatial domain), V_p the variance of
the spatial random effect. A site is defined here as the area/volume covered during
a given sampling/fishing operation. Any site i within the cell j as the same latent
probability of presence θ_j.
290
ii)   Observation process (detection):
Conditionally upon the latent probability of presence/absence in each grid cell j as
describe above, the multiple detection events $y_{j,i}$ (data 0/1) associated to the
same grid cell j are modelled as mutually independent two steps Bernoulli process
with detectability $\delta_{j,i}$:
$y_{j,i} \sim Bernoulli\left(z_{j,i}\,\delta_{j,i}\right)$ (2)
with
$z_{j,i} \sim Bernoulli\left(\theta_j\right)$ (3)
and
$logit\left(\delta_{j,i}\right) = W_{j,i}\gamma$
where $\delta_{j,i}$ denotes the probability of detecting the species at site i within cell j,
modelled as a fixed effect of the gear type associated with observation i. $W_{j,i}$ is the
raw vector of the design matrix, and γ is the vector of the gear effects, so as $W_{j,i}\gamma$
is the gear affect associated with observation at site i.

305
All SDMs were undertaken using the 'hSDM' package (Vieilledent et al., 2014) using
the 'mod.hSDM.siteocc.iCAR' function in R version 3.6.3 (R Code Team 2018). The
prior distribution for the variance of the spatial random effect followed weakly
informative uniform distribution. The effect of the detectability from gear and
from the suitability process was modelled as fixed effects drawn from an weakly
informative Gaussian prior centred at zero with a fixed standard deviation of 2
(Gelman et al., 2008; Northrup and Gerber, 2018). Each model was run with 50 000
Gibbs iterations, five Markov chain Monte Carlo simulations and a burn-in phase of
50 000 iterations.
315
2.4. Grid size and spatial distribution
A critical issue in our modelling framework is the choice of the grid size. Since the
latent field of presence/absence is modelled at the scale of the grid cell (eq. 1),
the larger the grid resolution the higher the probability to get at least one
presence within a grid cell. A finer grid cell will limit the quantity of available
observations falling within each cell, which therefore increases difficulty in model
fitting. We selected the finest grid as possible that allow the model to converge.
For almost all species modelled within Eastern Atlantic waters, a grid of 0.1x0.1
decimal degrees was used. Because of the scarcity of data for Lampreys, the model
did not converge with such a resolution and a coarser resolution of 0.2x0.2 decimal
degrees was therefore used for these species. For *A. agone*, that only occurs within
the Mediterranean, a grid resolution of 0.04 x 0.04 decimal degrees was used
because of the smaller area encompassed. This spatial resolution (0.04 x 0.04
decimal degrees) was not used for other species since it was not available at the
scale required within Eastern Atlantic waters. All environmental variables were
then aggregated and modelled to their relevant spatial resolution. There were
insufficient presences within the Mediterranean for *A. anguilla*, *P. marinus*, *C.*
ramada, *P. flesus* predictions (issues with model convergence). Predictions within
the Mediterranean for these were therefore excluded (Fig. 2).
335
2.5. Model evaluation
The probability of presence was calculated from the mean of the posterior
distribution of the latent field θ_j. A threshold of 0.4 was used to bin estimated
probability of presence into binary presence/absence values. A threshold of 0.4
was used as a compromise given the rarity of the species and accuracy following
examining various thresholds and plotting positive predicted values. The model
performance was then evaluated using sensitivity (proportion of correctly classified
presences), specificity (proportion of correctly classified absences) and percent
correct classification (proportion of presences and absences correctly using the R
package 'PresenceAbsence' (Freeman and Moisen, 2008a; Fielding and Bell, 1997;
Table 3). Max sensitivity specificity which is often used for rare species distribution
models (Freeman and Moisen, 2008b; Jiménez-Valverde and Lobo, 2007), was not
used since false presences was very high when applying this metric. Additionally,
our SDM objective was to try and minimise false presences and absences within the
distribution models. The threshold independent Area under the receiver operating
curve was also used to evaluate the model performance. These four metrics were
used to better understand the accuracy of the model since, some metrics can be
biased with low prevalence data and where imperfect detection can arise (Allouche
et al., 2006; Fourcade et al., 2018; Guillera-Arroita, 2017; Leroy et al., 2018).

So that areas with probabilities of presence above the threshold could be easily identified for management purposes, all predictions below 0.4 were mapped with a uniform light blue colour (Fig. 2). Uncertainty maps for each model were also produced to gage a better idea of areas of prediction with higher uncertainty (Loiselle et al., 2003; Spiegelhalter et al., 2002). The uncertainty maps were calculated by the mean coefficient of variation from the posterior distribution of spatial parameters in the model of best fit for each species (Lambert et al., 2020; Latimer et al., 2006). Probabilities of prediction and uncertainty above 0.4 were classified into 0.2 units to highlight areas of higher and lower probability of presence more clearly (Fig. 2; Fig. S4). As a result of the low diadromous fish presence, splitting the data into learning/validation samples led to insufficient presence in the learning data set. Cross validation was therefore not undertaken.

2.6. Diadromous fish river basin presence

The predicted distributions were visually compared to the distribution of diadromous fish within their freshwater habitats using the EuroDiad database (version 4; https://data.inrae.fr/dataset.xhtml?persistentId=doi:10.15454/IVVAIC; Barber-O'Malley et al., 2022). From EuroDiad database, migrating diadromous fish population presence within the study areas our model encompassed, were extracted using 'recent' literature citing from 1951 to 2020 (see Barber-O'Malley et al., (2022) and Béguer et al., (2007) for more details on the database) and mapped with the hierarchical SDM results (Fig. 2). The EuroDiad4 database was used as opposed to IUCN red list map distribution since it has been peer reviewed and it is more up to date.

2.7. Habitat Directive MPAs of relevance to diadromous fish

'Core areas' with a probability of presence >0.4 were selected and intersected with European Union Natura 2000 Habitat Directive Sites of Community Importance (https://www.eea.europa.eu/data-and-maps/data/natura-12). A threshold of > 0.4 probability of presence was selected as a result of the low probability of presence of diadromous fish (particularly Habitat Directive protected species), and to ensure consistency with the model evaluation threshold. Since only one Site of Community Importance was submitted to the European Union from the United Kingdom, Special Areas of Conservation were taken into consideration (https://jncc.gov.uk/our-work/special-areas-of-conservation-overview/). For each species, the number of MPAs (Sites of Community Importance and Special Areas of Conservation) and the proportion of MPAs of relevance to diadromous fish (core areas within an MPA) were recorded. The extent of the core area within the MPAs was not calculated as this would be dependent of the model grid used (Table 3). Furthermore, numerous MPAs were smaller than the grid.

3. Results

3.1. Diadromous fish hierarchical SDM

For all species the Area under the receiver operating curve and Percent Correct Classification performed very well ($\geq$75%; Table 3). Correct presence classification from the confusion matrices was consistently lower (<15%) than that of correct absence classification ($\geq$64%), and false absences were lower ($\leq$3%) than false presences ($\leq$22%) because of presence absence imbalance (Table 3; Table S1). The later led to on average better specificity than sensitivity scores (Table 3).

405
P. flesus was the most prevalent species modelled (5394 presences), whereas
salmonids the least (*S. salar* = 68 presences and *S. trutta* 63 presences) (Table 3).
SDMs that had lower presences (salmonids and lampreys) did not predict as well
(lower sensitivity and higher uncertainty maps) as species with higher presence
(shads, *C. ramada*, *P. flesus* and *O. eperlanus*). In addition, species that had a
more disperse distribution (e.g., *A. anguilla* and *S. salar*) did not predict as well
(sensitivity ≤ 0.66 for both species) as those with a more aggregated distribution
(*O. eperlanus* sensitivity = 0.81) (Table 3; Fig. 1 and 2). Uncertainty maps showed
that in general, areas of high and low predicted probability of presence had lower
uncertainty (Fig. S4).
416
All species were observed to have a relatively shallow (<300m depth) coastal
distribution (~<300 km from the coast; Fig. 2 and 3). The main predictor variables
for diadromous fish habitats were depth or distance from coast and salinity (Fig. 3).
Since sufficient *C. ramada* were observed within eastern Atlantic and
Mediterranean waters to model, two salinity peaks are observed corresponding to
the different salinity ranges of the two regions (Fig. 3).
423

Table 3. Hierarchical species distribution model outputs. All models contained gear as part of the observation process. AUC = Area Under the receiver operating Curve score, PCC = Percent Correct Classification, NetPP = Net Primary Production. Max depth is the maximum depth at which the dataset was cut. Note most environmental variables (all except distance and depth) were not available within very coastal areas (e.g., within estuaries and inlets). Numerous presences and absences were therefore lost when extracting the environmental variables for modelling.

Species	Model selected	Hauls	Presence	Absence (%)	Max depth (m)	Sensitivity	Specificity	AUC	PCC	Cell size (decimal degrees)
A. alosa	Depth + Salinity + NetPP + Sediment	86 292	802	99	300	0.79	0.92	0.95	0.95	0.1x0.1
A. fallax	Distance + Salinity + NetPP + Sediment	91 135	1385	98	300	0.88	0.81	0.90	0.82	0.1x0.1
A. agone	Depth + Salinity + Sediment	5086	176	97	200	0.81	0.74	0.86	0.75	0.04x0.04
A. anguilla	Depth + Salinity + NetPP	54 124	176	99	250	0.53	0.99	0.98	0.98	0.1x0.1
L. fluviatilis	Distance + Salinity	16 536	68	99	150	0.71	0.95	0.96	0.94	0.2x0.2
P. marinus	Distance + Salinity	28 582	74	99	300	0.60	0.98	0.94	0.97	0.2x0.2
C. ramada	Distance + Salinity + Netpp + Sediment	53 083	925	98	300	0.84	0.93	0.96	0.92	0.1x0.1
P. flesus	Distance + Salinity + NetPP + Sediment	92 400	5394	94	300	0.90	0.93	0.97	0.93	0.1x0.1
O. eperlanus	Depth + Salinity + NetPP	48 877	1035	98	450	0.81	0.99	0.99	0.99	0.1x0.1
S. salar	Depth + Salinity + NetPP	36 008	68	99	150	0.66	0.98	0.95	0.98	0.1x0.1
S. trutta	Depth + Salinity + NetPP	31 754	63	99	150	0.71	0.94	0.93	0.93	0.1x0.1

Fig. 2. Diadromous fish hierarchical species distribution model probability of
presence. Darker blue shades = SDM probability of presence > 0.4, light blue shade
= SDM probability of presence ≤ 0.4. Habitat Directive Marine protected Areas
(MPAs) outlined in black. MPAs of relevance to core (probability of presence > 0.4)
diadromous fish distribution, highlighted in dark red. Dark red cells are larger than
the gridded cells to ensure smaller core areas within MPAs are visible. Darker grey
terrestrial areas represent river basins which the diadromous fish migratory
populations have been observed (EuroDiad V.4). Note, no data were available
within the EuroDiad database from Dutch river basins.

443
444 **Fig. 3.** Diadromous fish hierarchical species distribution model environmental
445 predictor effects. NetPP = Net Primary Production, C = Course grain, M = Mud, S =
446 Sand, R = Rock. The solid line represents the smooth function estimate; the shaded
447 region represents the approximate 95% credibility interval. Boxplot blue stars
448 represent means and small black dots the predictions.
449

3.2. Diadromous fish river basin connectivity

2717 references contained within the EuroDiad database were used to map the 11 diadromous fish presence within the river basin. Core distributions of diadromous fish at sea largely matched with the freshwater areas of presence (Fig. 2). The spatial extent of diadromous fish at sea was often less than that observed within river basins (e.g., lampreys, *C. ramada, O. eperlanus* and *S. trutta*; Fig. 2). It should be noted that there are gaps in the EuroDiad database (refer to Barber-O'Malley et al., (2022)). For example, no diadromous fish presence data were available within Dutch freshwater habitats, and salmon populations are known to exist within Danish waters (de Groot et al., 2012; Maes et al., 2007; Rikardsen et al., 2021). Presences of *A. alosa* and *A. fallax* within the Mediterranean from the EuroDiad database will have been due to historic classification (Bagliniere and Elie, 2000; Keith et al., 2020).

3.3. Gear type detectability

All species were caught by a range of gear categories ranging from benthic, demersal and pelagic mobile trawls, seine nets, line gear types (Fig. 4; Table S2). Some general patterns were, however, evident. Line gear types caught the least diadromous fish (2% (weighted), 17 presences over ~2887 hauls), whereas demersal mobile trawl gear types (otter beam trawl and otter twin trawl) caught the most (67%, 7409 presences over ~82 531 hauls; Fig. 4; Table S2). Shad had a higher detectability with static and demersal mobile gears. *A. anguilla* were detected by a wide range of gear types. Lamprey were mainly detected by mobile trawl gears. *P. flesus* and *O. eperlanus* had a higher detectability with demersal and benthic mobile gear, and salmonid had a higher detectability with static nets and pelagic gears. Rarer species and species that were caught by a wider range of gear types had lower detectability than more abundance species caught by fewer gear types (Fig. 4). On average static nets (e.g., trammel net and set gillnets) and pelagic mobile gears (e.g., Otter midwater trawls) caught larger diadromous fish than mobile demersal and benthic gear types (Fig. S1). Although the modelled species were not target species of the fisheries within the observer database, some species were landed (Table 4).

483
484 **Fig. 4.** Matrix of the posterior mean (marginal posterior distribution) for gear A)
485 detectability (inverse logit of γ; the probability to detect the presence of fish
486 when they are present) and B) its standard deviation. Lighter blue shades represent
487 higher detectability (A) and darker green shades higher detectability standard
488 deviation (B).

489
490 **Table 4.** Landed and discarded diadromous fish from the ObsMer data. Excluding
491 targeted catches. Proportion landed = total landed / landed + discards.

Species	Landed	Discard	Hauls	Proportion landed
A. alosa	182	515	70 378	0.26
A. fallax	152	608	64 300	0.20
A. agone	28	69	7373	0.29
A. anguilla	10	8	28 194	0.56
L. fluviatilis	0	2	828	0.00
P. marinus	2	18	10 030	0.10
C. ramada	643	101	45 699	0.86
P. flesus	856	648	66 196	0.57
O. eperlanus	6	45	31 311	0.12
S. salar	29	27	26 009	0.52
S. trutta	43	18	31 666	0.70

492
3.4. Habitat Directive MPAs relevance
A total of 482 Habitat Directive MPAs have been designated within the area in
which the species were modelled (Fig. 2). Despite the small area of core habitats
relative to the model prediction area, for most Habitat Directive listed species,
numerous MPAs overlap with species core habitats, indicating their potential for
the protection of diadromous fish at sea (Table 5; Fig. 2).
499

Table 5. Habitat Directive Marine Protected Areas of relevance to core diadromous fish distribution (probability of presence >0.4). Diadromous fish listed under the Habitat Directive in bold.

Species	Number MPAs of relevance	Proportion of MPAs of relevance
A. alosa	**35**	**0.08**
A. fallax	**60**	**0.13**
A. agone	**10**	**0.27**
A. anguilla	19	0.04
L. fluviatilis	**11**	**0.02**
P. marinus	**14**	**0.03**
C. ramada	23	0.05
P. flesus	65	0.15
O. eperlanus	13	0.03
S. salar	**17**	**0.04**
S. trutta	28	0.06

4. Discussion

To our knowledge this is the first comprehensive study to model the distribution of diadromous fish at sea across eastern Atlantic and French Mediterranean waters. Using a hierarchical model allowed us to integrate an extensive database that we collated, containing both fisheries dependent and independent data to model the distribution of eleven rare diadromous fish over a wide spatial area. The model also enabled us to evaluate the probability to detect the presence of fish by the different gear types, and thus provide key information to evaluate the risk of bycatch of diadromous fish by commercial fisheries at sea.

Diadromous fish are bound to coastal areas because of their migratory behaviour. They are known to be fragile species because of the physiological and osmoregulatory changes required to undertake migrations between fresh water and marine environments (McDowall, 2009). Yet, coastal ecosystems, particularly within eastern Atlantic and Mediterranean waters, are subject to some of the highest anthropogenic impacts (Halpern et al., 2015; Korpinen et al., 2021; Lotze et al., 2006). Even though more data were collected within coastal areas, and the species are known to migrate from freshwater habitats, our study confirms that the investigated species were found to mainly concentrate in coastal and relatively shallow waters. The strong dependency of diadromous fish species to coastal areas, as evidenced in this study, reinforces the importance of investigating the vulnerability of diadromous fish to coastal anthropogenic pressures (Halpern et al., 2015; Limburg and Waldman, 2009; Lotze et al., 2006; Worm et al., 2006).

4.1. Hierarchical Bayesian SDM

On average, the eleven SDMs performed well. All the models had lower sensitivity and positive prediction values scores, because of low presence. Areas of both high probability and very low probability of presence had lower uncertainty, giving a

more accurate picture of confidence in the distributions (e.g., higher certainty that
most species are absent within waters further from the coast). In addition to
problems with very low occurrence, the areas of higher uncertainty for certain
species (e.g., *S. salar* and *S. trutta*, *A. alosa* and *A. fallax*), may be partially
because of misidentification between these groups of diadromous fish. The core
areas of probability of presence, in combination with the confidence maps provide
a better understanding of confidence of these predictions. To improve problems of
false presence predictions which might occur with species that are difficult to
identify, Bayesian models incorporating false positive errors could be undertaken
(e.g. Diana et al., 2021; Guillera-Arroita, 2017; Royle and Link, 2006).

Separating out the spatial random field of presence absence from the observations
enabled us to explicitly consider the imperfect detection of the observation
process. Since our database contained non-diadromous fish specific survey and
catch data which sampled the same grid cell numerous times, the use of the site
occupancy model enabled imperfect detection parameters to be estimated.
Disregarding false absences would have led to biased inferences and over-stated
parameter precision (Guillera-Arroita, 2017). The spatial random field of the
probability of presence explicitly integrates spatial autocorrelation in the
probability of presence, which limits the bias in inferences due to the non-random
spatial repartition of observations.

Since there were few presences for most species, our results were sensitive to the
choice of prior on the variance of the spatial random effect. We tested both the
inverse-gamma or a uniform distribution (the two prior forms that are available
using the hSDM package; Vieilledent et al., 2014) and found that the uniform
distribution provided better result. Some alternative prior choice such as the half-
Cauchy are advocated in the literature (Gelman, 2006), and future work should
explore if other prior choice could provide more accurate inferences. Equally, for
increased confidence in predictions of rare species, several different distribution
models could be compared or an ensemble model (that uses the means of several
models) implemented to stabilise inferences (Araújo and New, 2007; Latimer et al.,
2006; Loiselle et al., 2003).

For conservation purposes low thresholds are advised for low occurring species such
as diadromous fish (Freeman and Moisen, 2008a;b; Jiménez-Valverde and Lobo,
2007). However, when lowering the threshold an increase in false positives can
occur (Freeman and Moisen, 2008b). Our objective for the distribution models was
to minimise false presence and absences where possible, to provide more accurate
predictions whilst trying to meet conservations needs and minimise potential
impacts on sea-users any spatial protection measure that might be implemented
(Domisch et al., 2018; Loiselle et al., 2003; Maxwell and Jennings, 2005). Lack of
understanding in species distribution can lead to inefficient spatial protection
measures (Wauchope et al., 2022) and resource conflict (Probst et al., 2021).

4.2.Diadromous fish river basin connectivity
The EuroDiad database is a large database that incorporates information on
diadromous fish presence and abundance during their freshwater stage (Barber-
O'Malley et al., 2022; Béguer et al., 2007). Using the Eurodiad database to provide

large scale understanding of river basin presence of diadromous fish, we provide a
first attempt at qualitatively linking diadromous fish freshwater and marine
distributions within north-eastern Atlantic waters. Although the probabilities of
presence for most species were low, and gaps within the EuroDiad database
evident (e.g., gaps in diadromous fish presence from Denmark and Netherlands),
connectivity with their freshwater habitats appears to be visible, indicating
migration pathways.
589
It is thought *A. alosa* no longer exists within the Greater North Sea (Baglinière et
al., 2003; Wilson and Veneranta, 2019). Presences observed may have been from
misidentified *A. fallax*. The high probability of presence of *A. fallax* along the east
coast of the UK may be because of suitable conditions for this species as few
spawning rivers have been recorded along the east coast (Aprahamian et al., 1998).
For *A. agone* probabilities of presence clearly match that of their watersheds.
Predicted *A. anguilla* presences closely match tagged river presences from Righton
et al., (2016) despite the gaps found in the EuroDiad database. Predicted presence
of *L. fluviatilis* was higher along the north-west coast of Germany and Holland.
Presences have been observed here, albeit in low numbers (Admiraal et al., 1993;
Pavlov et al., 2017; Thiel and Salewski, 2003). *P. marinus* predicted presence was
slightly more scattered which fits with its wide distribution including into deeper
waters (Elliott et al., 2021; Lança et al., 2014). The very low occurrences of *A.
anguilla*, and the lamprey species is also likely to be due to inadequate sampling
methods used to study these species.
605
Probabilities of presence of *C. ramada*, *P. flesus* and *O. eperlanus* matched
relatively well their known presence from the EuroDiad database. From the
EuroDiad database, there are few *S. salar* river basin presences within the North
Sea. However, *S. salar* are known to occur within the North Sea and migrate
Northwards to their feeding grounds (Mork et al., 2012; Rikardsen et al., 2021).
Occurrences observed were most likely 1 sea winter and 2 sea winters individuals
returning to their natal rivers given their size ranges. From the rivers that are
monitored, *S. trutta* distribution matched well their freshwater river occupancy
(ICES, 2020). The few salmonids observed may be as a result of the relatively few
pelagic trawls, in addition to misreporting (ICES, 2005).
616
Given the coastal habitat occupancy for most of these diadromous fish and declines
observed (Limburg and Waldman, 2009; Merg et al., 2020), detailed analysis of the
connectivity between both habitats is essential (Flitcroft et al., 2019; Lin et al.,
2017; McDowall, 2009). Unfortunately, such a model was not possible here,
because it would require more detailed knowledge on their freshwater habitat
occupancy (i.e., numerous outlets diadromous fish were observed within proximity
to, were not contained within the EuroDiad database) and their river fidelity.
Furthermore, seasonal and stage specific SDMs, which would provide more
information on their ontogenetic migration movements, were not possible because
of the low prevalence of the data and dispersion. Developing future research to
model the connectivity between both habitats is key to improving our
understanding of the pressures faced during their migrations (Flitcroft et al., 2019;
Lin et al., 2017; McDowall, 2009).
630

4.3.Gear type detectability

Even though little bycatch was observed, given a number of the diadromous fish studied here are threatened, even a small amount of bycatch may impact their populations (Dulvy et al., 2003). Furthermore, misreporting and illegal fishing is likely to remain (Elliott et al., 2020a; ICES, 2005; Stratoudakis et al., 2020; Worm et al., 2013). Bycatch detectability results from our models could help provide management advice so that gear types and areas with higher bycatch rate be avoided. Such advice can provide direct information for the MSFD and the Habitat Directive that requires the identification of gear types which may threaten protected species (1992/43/EEC; 2008/56/EC).

All species were caught by a wide range of gear types (e.g., benthic, demersal and pelagic trawls, and seine nets), despite the different gear categories being deployed at different depths. The species were probably caught by a range of gear categories because of their diel vertical migratory behaviour whilst at sea (Kristensen et al., 2018; Lança et al., 2014; Righton et al., 2016). In addition, certain gear types (e.g., trawls and static nets), can catch fish within both the demersal and pelagic water zones (He et al., 2021; Borges et al., 2008). Higher detectability from certain gear types were, however, observed. For example, shad were mainly caught by demersal mobile gear and static gear, salmonids static and pelagic mobile gear types, and *P. flesus* and *O.eperlanus* were largely captured by demersal and benthic mobile gear. These results broadly match the water column habitat occupancy these species are known to occupy and existing literature (ICES, 2005; Wilson and Veneranta, 2019). As it is thought that lampreys detach from their host upon capture (Elliott et al., 2021; Halliday, 1991), bycatch is likely to be minimal for these species.

To improve understanding of bycatch risk on a spatial level, fishing intensity by gear type should be mapped, ideally using Vessel Monitoring System data and overlaying it with the predicted distributions (Elliott et al., 2018; Quemmerais-Amice et al., 2020). Unfortunately, this information was not available to us. Additionally, since the fisheries observer data we had access to is only a small sample of the existing French fishing activity (Cornou et al., 2015), and it does not include other countries fishing effort, results would be biased.

4.4.MPA relevance

To improve protection of diadromous fish at sea, numerous mechanisms could be implemented. For example, identifying areas of high bycatch risk with measures to limit bycatch within these areas, improving diadromous fish migratory pathways, and spatial protection measures (Verhelst et al., 2021; Wilson and Veneranta, 2019). Few MPAs within European waters protect marine fish and instead focus on protecting seabed features with a few exceptions (e.g., protected seabirds and marine mammals). Therefore the only protection measure that exist for vulnerable marine fish is through the removal of targeted fishing (Dureuil et al., 2018; Probst et al., 2021; Stratoudakis et al., 2016). There is, however, increasing evidence that MPAs may have positive impacts on fish populations if appropriate management measures are implemented (Davies et al., 2021; Moland et al., 2013; Probst et al., 2021; Worm et al., 2006). The value of protected areas for highly migratory species

679 has been questioned, but there is evidence that MPAs may have positive impacts on
680 migratory species (Pendoley et al., 2014; Takashina and Mougi, 2014).
681
Our results show relatively high core area presence of protected diadromous fish
within Habitat Directive MPAs. Most marine Sites of Community Importance have
not been designated to protect Habitat Directive listed diadromous fish.
Nonetheless, our results highlight the value of these MPAs for diadromous fish
protection. Specific management measures limiting threats (e.g., limiting access to
gear types with higher probability of capture) to diadromous fish within these
areas, would be of benefit to their protection (Domisch et al., 2019; Stratoudakis
et al., 2016). Detailed analysis of Sites of Community Importance and other
categories of designated MPAs should be undertaken to evaluate their potential to
protect diadromous fish.
692
4.5. Conclusion
Understanding the spatial distribution of species and their habitats is essential for
effective management and conservation (Halpern et al., 2015; Worm et al., 2009,
2005). Much research has been carried out on the conservation of freshwater
diadromous fish (Drouineau et al., 2018; Merg et al., 2020; Verhelst et al., 2021).
Very little research has been dedicated to the value of marine and estuary habitats
for the conservation of these species and their risk of bycatch (Feunteun, 2002;
Flitcroft et al., 2019; Lin et al., 2017). Here we provide an insight into the
distribution and bycatch of diadromous fish at sea.
702
Despite diadromous fish vulnerability, targeted fishing within estuaries for
numerous species remains (Aprahamian and Walker, 2008; Castelnaud, 2000;
Feunteun, 2002; Kappel, 2005; Stratoudakis et al., 2020). Given the very coastal
distribution of these species, more detailed analysis into such catch data (small
fishing vessels targeting diadromous fish) is required. Smaller vessels and artisanal
fisheries are still largely overlooked, regardless of their potential threat to
diadromous fish (Beaulaton et al., 2008; Castelnaud, 2000; Stratoudakis et al.,
2016). Having a better understanding of fishing pressure, combined with the
outputs of the distributions and gear capture from these models, could help
improve understanding of fishing impacts on diadromous fish.
713
Finally, it has been well acknowledged that although a large impetus for the
designation of MPAs has been undertaken in the last two decades (Probst et al.,
2021; Worm et al., 2006). The implementation of management measures has been
slow to put in place (Dureuil et al., 2018; Probst et al., 2021; Stratoudakis et al.,
2016). Furthermore, little has been undertaken to protect diadromous fish within
marine ecosystems (Dureuil et al., 2018; Probst et al., 2021; Stratoudakis et al.,
2016). We highlight the value of Habitat Directive MPAs that have the jurisdiction
to protect diadromous fish. Limiting access of fisheries with a higher probability of
capture to such areas could provide additional protection for diadromous fish and
their habitats.
724

Acknowledgements
We thank the funders of the project Management of Diadromous Fish in their
Environment, OFB-INRAE-Institute Agro-UPPA. We are extremely grateful to all

those who were involved in collecting and compiling the fisheries-dependent and
fisheries-independent data and their funders. We are also grateful to IFREMER and
the French marine fisheries and aquaculture administration (DPMA) for access to
their data. Finally, thanks the reviewers for their valuable contribution.
732

References

Admiraal, W., van der Velde, G., Smit, H., Cazemier, W.G., 1993. The rivers Rhine
and Meuse in The Netherlands: present state and signs of ecological recovery.
Hydrobiologia 265, 97–128. https://doi.org/10.1007/BF00007264

Alglave, B., Rivot, E., Etienne, M.-P., Woillez, M., Thorson, J.T., Vermard, Y.,
2022. Combining scientific survey and commercial catch data to map fish
distribution. ICES J. Mar. Sci. 79, 1133–1149.
https://doi.org/10.1093/icesjms/fsac032

Allouche, O., Tsoar, A., Kadmon, R., 2006. Assessing the accuracy of species
distribution models: prevalence, kappa and the true skill statistic (TSS). J.
Appl. Ecol. 43, 1223–1232. https://doi.org/10.1111/j.1365-2664.2006.01214.x

Almeida, P.R., 1996. Estuarine movement patterns of adult thin-lipped grey mullet,
Liza ramada (Risso) (*Pisces, Mugilidae*), observed by ultrasonic tracking. J.
Exp. Mar. Bio. Ecol. 202, 137–150. https://doi.org/10.1016/0022-
0981(95)00162-X

Aprahamian, M., Walker, A., 2008. Status of eel fisheries, stocks and their
management in England and Wales. Knowl. Manag. Aquat. Ecosyst. 1–7.
https://doi.org/10.1051/kmae/2009007

Aprahmian, M.W., Lester, S.M., Aprahamian, S.D., 1998. Shad conservation in
England and Wales. R&D Technical report W110. Environment Agency:
https://www.gov.uk/government/publications/shad-conservation-in-england-
and-wales (last accessed 22/09/2022)

Araújo, M.B., New, M., 2007. Ensemble forecasting of species distributions. Trends
Ecol. Evol. 22, 42–47. https://doi.org/10.1016/j.tree.2006.09.010

Arevalo, E., Lassalle, G., Tétard, S., Maire, A., Sauquet, E., Lambert, P., Paumier,
A., Villeneuve, B., Drouineau, H., 2020. An innovative bivariate approach to
detect joint temporal trends in environmental conditions: Application to large
French rivers and diadromous fish. Sci. Total Environ. 748, 141260.
https://doi.org/10.1016/j.scitotenv.2020.141260

Barber-O'Malley, B., Lassalle, G., Lambert, P., Quinton, E., 2022. Dataset on
European diadromous species distributions from 1750 to present time in
Europe, North Africa and the Middle East. Data Br. 40, 107821.
https://doi.org/10.1016/j.dib.2022.107821

Baum, J.K., Myers, R.A., Kehler, D.G., Worm, B., Harley, S.J., Doherty, P.A., 2003.
Collapse and Conservation of Shark Populations in the Northwest Atlantic.
Science. 299, 389–392. https://doi.org/10.1126/science.1079777

Bagliniere, J.L., Elie, P., 2000. Les aloses (*Alosa alosa* et *Alosa fallax* spp.)
Ecobiologie et variabilité des populations. Institut National de la Recherche
Agronomique, Paris. ISBN 2-7380-0906-9-ISSN :0763-1707

Beaulaton, L., Taverny, C., Castelnaud, G., 2008. Fishing, abundance and life
history traits of the anadromous sea lamprey (*Petromyzon marinus*) in Europe.
Fish. Res. 92, 90–101. https://doi.org/10.1016/j.fishres.2008.01.001

Béguer, M., Beaulaton, L., Rochard, E., 2007. Distribution and richness of
diadromous fish assemblages in Western Europe: large-scale explanatory

factors. Ecol. Freshw. Fish 16, 221–237. https://doi.org/10.1111/j.1600-
0633.2006.00214.x
Belmont, J., Miller, C., Scott, M., Wilkie, C., 2022. A new statistical approach for
identifying rare species under imperfect detection. Divers. Distrib. 28, 882–
893. https://doi.org/10.1111/ddi.13495
Berg, S., Krog, C., Muus, B., Nielsen, J., Fricke, R., Berghahn, R., Neudecker, T.,
Wolff, W.J., 1996. Red List of lampreys and marine fishes of the Wadden Sea.
Helgoländer Meeresuntersuchungen 50, 101–105.
https://doi.org/10.1007/BF02366178
Berkson, J., Thorson, J.T., 2015. The determination of data-poor catch limits in
the United States: is there a better way? ICES J. Mar. Sci. 72, 237–242.
https://doi.org/10.1093/icesjms/fsu085
Bisch, A., Elliott, S.A.M., Carpentier, A., Acou, A., 2022. Modelling the distribution
of vulnerable skate from fisheries dependent data using imperfect detection.
Prog. Oceanogr. 206, 102859. https://doi.org/10.1016/j.pocean.2022.102859
Bourdaud, P., Travers-Trolet, M., Vermard, Y., Cormon, X., Marchal, P., 2017.
Inferring the annual, seasonal, and spatial distributions of marine species
from complementary research and commercial vessels' catch rates. ICES J.
Mar. Sci. 74, 2415–2426. https://doi.org/10.1093/icesjms/fsx092
Castelnaud, G., 2000. Fishing location, fishermen and production of migratory
species fisheries in the French rivers. BFPP - Bull. Fr. la Pech. la Prot. des
Milieux Aquat. 439–458. https://doi.org/10.1051/kmae/2001060
Cornou, A.S., Quinio-Scavinner, M., Delaunay, D., Dimeet, J. et al., 2015.
Observations à bord des navires de pêche professionnelle. Bilan de
l'échantillonnage 2014. Ifremer, Nantes
Costa, M.J., Duarte, G., Segurado, P., Branco, P., 2021. Major threats to European
freshwater fish species. Sci. Total Environ. 797, 149105.
https://doi.org/10.1016/j.scitotenv.2021.149105
Davies, B.F.R., Holmes, L., Rees, A., Attrill, M.J., Cartwright, A.Y., Sheehan, E. V.,
2021. Ecosystem Approach to Fisheries Management works—How switching
from mobile to static fishing gear improves populations of fished and non-
fished species inside a marine-protected area. J. Appl. Ecol. 1–16.
https://doi.org/10.1111/1365-2664.13986
de Groot, R., Brander, L., van der Ploeg, S., Costanza, R., Bernard, F., Braat, L.,
Christie, M., Crossman, N., Ghermandi, A., Hein, L., Hussain, S., Kumar, P.,
McVittie, A., Portela, R., Rodriguez, L.C., ten Brink, P., van Beukering, P.,
2012. Global estimates of the value of ecosystems and their services in
monetary units. Ecosyst. Serv. 1, 50–61.
https://doi.org/10.1016/j.ecoser.2012.07.005
Diana, A., Matechou, E., Griffin, J.E., Buxton, A.S., Griffiths, R.A., 2021. An RShiny
app for modelling environmental DNA data: accounting for false positive and
false negative observation error. Ecography (Cop.). 44, 1838–1844.
https://doi.org/10.1111/ecog.05718
Domisch, S., Friedrichs, M., Hein, T., Borgwardt, F., Wetzig, A., Jähnig, S.C.,
Langhans, S.D., 2018. Spatially explicit species distribution models: A missed
opportunity in conservation planning? Divers. Distrib. 25, 758–769.
https://doi.org/10.1111/ddi.12891
Dormann, C.F., Elith, J., Bacher, S., Buchmann, C., Carl, G., Carré, G., Marquéz,
J.R.G., Gruber, B., Lafourcade, B., Leitão, P.J., Münkemüller, T., McClean,

C., Osborne, P.E., Reineking, B., Schröder, B., Skidmore, A.K., Zurell, D.,
    Lautenbach, S., 2013. Collinearity: a review of methods to deal with it and a
    simulation study evaluating their performance. Ecography (Cop.). 36, 27–46.
    https://doi.org/10.1111/j.1600-0587.2012.07348.x

Dormann, C.F., McPherson, J.M., Araújo, M.B., Bivand, R., Bolliger, J., Carl, G.,
    Davies, R.G., Hirzel, A., Jetz, W., Daniel Kissling, W., Kühn, I., Ohlemüller,
    R., Peres-Neto, P.R., Reineking, B., Schröder, B., Schurr, F.M., Wilson, R.,
    2007. Methods to account for spatial autocorrelation in the analysis of species
    distributional data: a review. Ecography (Cop.). 30, 609–628.
    https://doi.org/10.1111/j.2007.0906-7590.05171.x

Drouineau, H., Carter, C., Rambonilaza, M., Beaufaron, G., Bouleau, G., Gassiat,
    A., Lambert, P., le Floch, S., Tétard, S., de Oliveira, E., 2018. River
    Continuity Restoration and Diadromous Fishes: Much More than an Ecological
    Issue. Environ. Manage. 61, 671–686. https://doi.org/10.1007/s00267-017-
    0992-3

Dulvy, N.K., Sadovy, Y., Reynolds, J.D., 2003. Extinction vulnerability in marine
    populations. Fish Fish. 4, 25–64. https://doi.org/10.1046/j.1467-
    2979.2003.00105.x

Dureuil, M., Boerder, K., Burnett, K.A., Froese, R., Worm, B., 2018. Elevated
    trawling inside protected areas undermines conservation outcomes in a global
    fishing hot spot. Science. 362, 1403–1407.
    https://doi.org/10.1126/science.aau0561

Elliott, M., Dewailly, F., 1995. The structure and components of European
    estuarian fish assemblages. Netherlands J. Aquat. Ecol. 29, 397–417.

Elliott, S.A.M., Bearup, D., Carpentier, A., Larivain, A., Trancart, T., Feunteun, E.,
    2020a. Evaluating the effectiveness of management measures on skates in a
    changing world. Biol. Conserv. 248, 108684.
    https://doi.org/10.1016/j.biocon.2020.108684

Elliott, S.A.M., Carpentier, A., Feunteun, E., Trancart, T., 2020b. Distribution and
    life history trait models indicate vulnerability of skates. Prog. Oceanogr. 181,
    102256. https://doi.org/10.1016/j.pocean.2019.102256

Elliott, S.A.M., Deleys, N., Rivot, E., Acou, A., Réveillac, E., Beaulaton, L., 2021.
    Shedding light on the river and sea lamprey within western European marine
    waters. Endanger. Species Res. 44, 409–419.
    https://doi.org/10.3354/esr01113

Elliott, S.A.M., Guérin, L., Pesch, R., Schmitt, P., Meakins, B., Vina-Herbon, C.,
    González-Irusta, J.M., de la Torriente, A., Serrano, A., 2018. Integrating
    benthic habitat indicators: Working towards an ecosystem approach. Mar.
    Policy 90, 88–94. https://doi.org/10.1016/j.marpol.2018.01.003

Elliott, S.A.M., Milligan, R.J., Heath, M.R., Turrell, W.R., Bailey, D.M., 2016.
    Disentangling habitat concepts for demersal marine fish mangement.
    Oceanogr. Mar. Biol. Annu. Rev. 54, 173–191.
    https://doi.org/10.1201/9781315368597

Engler, R., Guisan, A., Rechsteiner, L., 2004. An improved approach for predicting
    the distribution of rare and endangered species from occurrence and pseudo-
    absence data. J. Appl. Ecol. 41, 263–274. https://doi.org/10.1111/j.0021-
    8901.2004.00881.x

ERSEM,. 2020. NEMO-ERSEM and POLCOMS-ERSEM. Licence to use Copernicus
    Products. Reading, UK. DOI.org/10.24381/cds.dcc9295c

https://cds.climate.copernicus.eu/cdsapp#!/dataset/10.24381/cds.dcc9295c?
tab=overview.

Fauconnet, L., Trenkel, V.M., Morandeau, G., Caill-Milly, N., Rochet, M.-J., 2015.
Characterizing catches taken by different gears as a step towards evaluating
fishing pressure on fish communities. Fish. Res. 164, 238–248.
https://doi.org/10.1016/j.fishres.2014.11.019

Feunteun, E., 2002. Management and restoration of European eel population
(*Anguilla anguilla*): An impossible bargain. Ecol. Eng. 18, 575–591.
https://doi.org/10.1016/S0925-8574(02)00021-6

Fielding, A.H., Bell, J.F., 1997. A review of methods for the assessment of
prediction errors in conservation presence / absence models. Environmenal
Conserv. 24, 38–49.

Flitcroft, R.L., Arismendi, I., Santelmann, M.V., 2019. A Review of Habitat
Connectivity Research for Pacific Salmon in Marine, Estuary, and Freshwater
Environments. JAWRA J. Am. Water Resour. Assoc. 55, 430–441.
https://doi.org/10.1111/1752-1688.12708

Fourcade, Y., Besnard, A.G., Secondi, J., 2018. Paintings predict the distribution of
species, or the challenge of selecting environmental predictors and evaluation
statistics. Glob. Ecol. Biogeogr. 27, 245–256.
https://doi.org/10.1111/geb.12684

Freeman, E.A., Moisen, G., 2008a. PresenceAbsence: An R Package for Presence-
Absence Model Analysis. Journal of Statistical Software, 23(11):1-31.
http://www.jstatsoft.org/v23/i11

Freeman, E.A., Moisen, G., 2008b. PresenceAbsence: An R Package for Presence-
Absence Model Analysis. Journal of Statistical Software, 23(11):1-31.
http://www.jstatsoft.org/v23/i11

Freyhof, J., Brooks, E., 2011. European Red List of freshwater fishes. Publications
Office of the European Union, Luxembourg.
https://ec.europa.eu/environment/nature/conservation/species/redlist/dow
nloads/European_freshwater_fishes.pdf (last accessed 01/07/2020).

Gelman, A., 2006. Prior distributions for variance parameters in hierarchical
models (comment on article by Browne and Draper). Bayesian Anal. 515–534.
https://doi.org/10.1214/06-BA117A

Gelman, A., Jakulin, A., Pittau, M.G., Su, Y.-S., 2008. A weakly informative default
prior distribution for logistic and other regression models. Ann. Appl. Stat. 2,
1360–1383. https://doi.org/10.1214/08-AOAS191

Guillera-Arroita, G., 2017. Modelling of species distributions, range dynamics and
communities under imperfect detection: advances, challenges and
opportunities. Ecography (Cop.). 40, 281–295.
https://doi.org/10.1111/ecog.02445

Guisan, A., Thuiller, W., 2005. Predicting species distribution: offering more than
simple habitat models. Ecol. Lett. 8, 993–1009.
https://doi.org/10.1111/j.1461-0248.2005.00792.x

Guisan, A., Zimmermann, N.E., 2000. Predictive habitat distribution models in
ecology. Ecol. Modell. 135, 147–186. https://doi.org/10.1016/S0304-
3800(00)00354-9

Halliday, R.G., 1991. Marine distribution of the sea lamprey (*Petromyzon marinus*)
in the northwest Atlantic. Can. J. Fish. Aquat. Sci. 48, 832–842.
https://doi.org/10.1139/f91-099

Halpern, B.S., Frazier, M., Potapenko, J., Casey, K.S., Koenig, K., Longo, C., Lowndes, J.S., Rockwood, R.C., Selig, E.R., Selkoe, K.A., Walbridge, S., 2015. Spatial and temporal changes in cumulative human impacts on the world's ocean. Nat. Commun. 6, 1–7. https://doi.org/10.1038/ncomms8615

He, P., Chopin, F., Suuronen, P., Ferro, R.S.T and Lansley, J. 2021. Classification and illustrated definition of fishing gears. FAO Fisheries and Aquaculture Technical Paper No. 672. Rome, FAO. https://doi.org/10.4060/cb4966en (last accessed 22/09/2020).

Isaac, N.J.B., Jarzyna, M.A., Keil, P., Dambly, L.I., Boersch-Supan, P.H., Browning, E., Freeman, S.N., Golding, N., Guillera-Arroita, G., Henrys, P.A., Jarvis, S., Lahoz-Monfort, J., Pagel, J., Pescott, O.L., Schmucki, R., Simmonds, E.G., O'Hara, R.B., 2020. Data Integration for Large-Scale Models of Species Distributions. Trends Ecol. Evol. 35, 56–67. https://doi.org/10.1016/j.tree.2019.08.006

ICES. 2005. Report of the Study Group on the Bycatch of Salmon in Pelagic Trawl Fisheries (SGBYSAL), 8–11 February 2004, Bergen, Norway. ICES CM 2005/ACFM:13. 41 pp.

ICES. 2020. Working Group with the Aim to Develop Assessment Models and Establish Biological Reference Points for Sea Trout (Anadromous Salmo trutta) Populations (WGTRUTTA; outputs from 2019 meeting).ICES Scientific Reports. 2:59. 59 pp. http://doi.org/10.17895/ices.pub.7431

Isaac, N.J.B., Jarzyna, M.A., Keil, P., Dambly, L.I., Boersch-Supan, P.H., Browning, E., Freeman, S.N., Golding, N., Guillera-Arroita, G., Henrys, P.A., Jarvis, S., Lahoz-Monfort, J., Pagel, J., Pescott, O.L., Schmucki, R., Simmonds, E.G., O'Hara, R.B., 2020. Data Integration for Large-Scale Models of Species Distributions. Trends Ecol. Evol. 35, 56–67. https://doi.org/10.1016/j.tree.2019.08.006

Jiménez-Valverde, A., Lobo, J.M., 2007. Threshold criteria for conversion of probability of species presence to either–or presence–absence. Acta Oecologica 31, 361–369. https://doi.org/10.1016/j.actao.2007.02.001

Kappel, C. V., 2005. Losing pieces of the puzzle: Threats to marine, estuarine, and diadromous species. Front. Ecol. Environ. 3, 275–282. https://doi.org/10.1890/1540-9295(2005)003[0275:LPOTPT]2.0.CO;2

Keith, P., Poulet, N., Denys, G., Changeux, T., Feunteun, E., et Persat, H., 2020. Les poissons d'eau douce de France. Biotope, Publications scientifiques du MNHN. 703 p

Kellner, K.F., Swihart, R.K., 2014. Accounting for Imperfect Detection in Ecology: A Quantitative Review. PLoS One 9, e111436. https://doi.org/10.1371/journal.pone.0111436

Korpinen, S., Laamanen, L., Bergström, L., Nurmi, M., Andersen, J.H., Haapaniemi, J., Harvey, E.T., Murray, C.J., Peterlin, M., Kallenbach, E., Klančnik, K., Stein, U., Tunesi, L., Vaughan, D., Reker, J., 2021. Combined effects of human pressures on Europe's marine ecosystems. Ambio 50, 1325–1336. https://doi.org/10.1007/s13280-020-01482-x

Kristensen, M., Righton, D., del Villar-Guerra, D., Baktoft, H., Aarestrup, K., 2018. Temperature and depth preferences of adult sea trout *Salmo trutta* during the marine migration phase. Mar. Ecol. Prog. Ser. 599, 209–224. https://doi.org/10.3354/meps12618

Lambert, C., Authier, M., Dorémus, G., Laran, S., Panigada, S., Spitz, J., Van Canneyt, O., Ridoux, V., 2020. Setting the scene for Mediterranean litterscape management: The first basin-scale quantification and mapping of floating marine debris. Environ. Pollut. 263, 114430. https://doi.org/10.1016/j.envpol.2020.114430

Lança, M.J., Machado, M., Mateus, C.S., Lourenço, M., Ferreira, A.F., Quintella, B.R., Almeida, P.R., 2014. Investigating Population Structure of Sea Lamprey (*Petromyzon marinus*, L.) in Western Iberian Peninsula Using Morphological Characters and Heart Fatty Acid Signature Analyses. PLoS One 9, e108110. https://doi.org/10.1371/journal.pone.0108110

Lassalle, G., Béguer, M., Beaulaton, L., Rochard, E., 2008. Diadromous fish conservation plans need to consider global warming issues: An approach using biogeographical models. Biol. Conserv. 141, 1105–1118. https://doi.org/10.1016/j.biocon.2008.02.010

Lassalle, G., 2021. EuroDiad version 4.0, https://doi.org/10.15454/IVVAIC, Portail Data INRAE, V2, UNF:6:X15FgpfOWssoJvGSg0wyoA== [fileUNF]

Latimer, A.M., Wu, S., Gelfand, A.E., Silander Jr., J.A., 2006. Building Statistical Models To Analyze Species Distributions. Ecol. Appl. 16, 33–50. https://doi.org/10.1890/04-0609

Leathwick, J.R., Rowe, D., Richardson, J., Elith, J., Hastie, T., 2005. Using multivariate adaptive regression splines to predict the distributions of New Zealand's freshwater diadromous fish. Freshw. Biol. 50, 2034–2052. https://doi.org/10.1111/j.1365-2427.2005.01448.x

Legrand, M., Briand, C., Buisson, L., Artur, G., Azam, D., Baisez, A., Barracou, D., Bourré, N., Carry, L., Caudal, A.-L., Charrier, F., Corre, J., Croguennec, E., Der Mikaélian, S., Josset, Q., Le Gurun, L., Schaeffer, F., Laffaille, P., 2020. Contrasting trends between species and catchments in diadromous fish counts over the last 30 years in France. Knowl. Manag. Aquat. Ecosyst. 2020-Janua, 7. https://doi.org/10.1051/kmae/2019046

Leroy, B., Delsol, R., Hugueny, B., Meynard, C.N., Barhoumi, C., Barbet-Massin, M., Bellard, C., 2018. Without quality presence-absence data, discrimination metrics such as TSS can be misleading measures of model performance. J. Biogeogr. 45, 1994–2002. https://doi.org/10.1111/jbi.13402

Limburg, K.E., Waldman, J.R., 2009. Dramatic Declines in North Atlantic Diadromous Fishes. Bioscience 59, 955–965. https://doi.org/10.1525/bio.2009.59.11.7

Lin, H., Bush, A., Linke, S., Possingham, H.P., Brown, C.J., 2017. Climate change decouples marine and freshwater habitats of a threatened migratory fish. Divers. Distrib. 23, 751–760. https://doi.org/10.1111/ddi.12570

Loiselle, B.A., Howell, C.A., Graham, C.H., Goerck, J.M., Brooks, T., Smith, K.G., Williams, P.H., 2003. Avoiding Pitfalls of Using Species Distribution Models in Conservation Planning. Conserv. Biol. 17, 1591–1600. https://doi.org/10.1111/j.1523-1739.2003.00233.x

Lomba, A., Pellissier, L., Randin, C., Vicente, J., Moreira, F., Honrado, J., Guisan, A., 2010. Overcoming the rare species modelling paradox: A novel hierarchical framework applied to an Iberian endemic plant. Biol. Conserv. 143, 2647–2657. https://doi.org/10.1016/j.biocon.2010.07.007

Lotze, H.K., Lenihan, H.S., Bourque, B.J., Bradbury, R.H., Cooke, R.G., Kay, M.C., Kidwell, S.M., Kirby, M.X., Peterson, C.H., Jackson, J.B.C., 2006. Depletion,

Degradation, and Recovery Potential of Estuaries and Coastal Seas. Science. 312, 1806–1809. https://doi.org/10.1126/science.1128035

MacKenzie, D.I., Nichols, J.D., Lachman, G.B., Droege, S., Royle, A.A., Langtimm, C.A., 2002. Estimating site occupancy rates when detection probabilities are less than one. Ecology 83, 2248–2255. https://doi.org/10.1890/0012-9658(2002)083[2248:ESORWD]2.0.CO;2

Maes, J., Stevens, M., Breine, J., 2007. Modelling the migration opportunities of diadromous fish species along a gradient of dissolved oxygen concentration in a European tidal watershed. Estuar. Coast. Shelf Sci. 75, 151–162. https://doi.org/10.1016/j.ecss.2007.03.036

Maxwell, D., Jennings, S., 2005. Power of monitoring programmes to detect decline and recovery of rare and vulnerable fish. J. Appl. Ecol. 42, 25–37. https://doi.org/10.1111/j.1365-2664.2005.01000.x

McDowall, R.M., 2009. Making the Best of Two Worlds : Diadromy in the Evolution, Ecology, and Conservation of Aquatic Organisms. Am. Fish. Soc. Symp. 69, 1–22.

Merg, M.-L., Dézerald, O., Kreutzenberger, K., Demski, S., Reyjol, Y., Usseglio-Polatera, P., Belliard, J., 2020. Modeling diadromous fish loss from historical data: Identification of anthropogenic drivers and testing of mitigation scenarios. PLoS One 15, e0236575. https://doi.org/10.1371/journal.pone.0236575

Meyer, C.F.J., Aguiar, L.M.S., Aguirre, L.F., Baumgarten, J., Clarke, F.M., Cosson, J.F., Villegas, S.E., Fahr, J., Faria, D., Furey, N., Henry, M., Hodgkison, R., Jenkins, R.K.B., Jung, K.G., Kingston, T., Kunz, T.H., Cristina Macswiney Gonzalez, M., Moya, I., Patterson, B.D., Pons, J.M., Racey, P.A., Rex, K., Sampaio, E.M., Solari, S., Stoner, K.E., Voigt, C.C., von Staden, D., Weise, C.D., Kalko, E.K.V., 2011. Accounting for detectability improves estimates of species richness in tropical bat surveys. J. Appl. Ecol. 48, 777–787. https://doi.org/10.1111/j.1365-2664.2011.01976.x

Moland, E., Olsen, E.M., Knutsen, H., Garrigou, P., Espeland, S.H., Kleiven, A.R., Andre, C., Knutsen, J.A., 2013. Lobster and cod benefit from small-scale northern marine protected areas: inference from an empirical before-after control-impact study. Proc. R. Soc. B Biol. Sci. 280, 1–10. https://doi.org/10.1098/rspb.2012.2679

Moriarty, M., Sethi, S.A., Pedreschi, D., Smeltz, T.S., McGonigle, C., Harris, B.P., Wolf, N., Greenstreet, S.P.R., 2020. Combining fisheries surveys to inform marine species distribution modelling. ICES J. Mar. Sci. 77, 539–552. https://doi.org/10.1093/icesjms/fsz254

Nieto, A., Ralph, G.M., Comeros-Raynal, M.T., Kemp, J., Garcia Criado, M., Allen, D.J., Dulvy, N.K., Walls, R.H.L., Russell, B., Pollard, D., et al., 2015. European. Red List of marine fishes. Luxembourg: Publications Office of the European Union. https://ec.europa.eu/environment/nature/conservation/species/redlist/downloads/European_marine_fishes.pdf (last accessed 01/07/2020)

Mork, K.A., Gilbey, J., Hansen, L.P., Jensen, A.J., Jacobsen, J.A., Holm, M., Holst, J.C., Ó Maoiléidigh, N., Vikebø, F., McGinnity, P., Melle, W., Thomas, K., Verspoor, E., Wennevik, V., 2012. Modelling the migration of post-smolt Atlantic salmon (Salmo salar) in the Northeast Atlantic. ICES J. Mar. Sci. 69, 1616-1624. https://doi.org/10.1093/icesjms/fss108

Northrup, J.M., Gerber, B.D., 2018. A comment on priors for Bayesian occupancy models. PLoS One 13, e0192819. https://doi.org/10.1371/journal.pone.0192819

Pavlov, D.S., Zvezdin, A.O., Kostin, V. V., Tsimbalov, I.A., Kucheryavyy, A. V., 2017. Temporal characteristics of downstream migration of smolts of the European river lamprey *Lampetra fluviatilis* in the Chernaya River. Biol. Bull. 44, 290–295. https://doi.org/10.1134/S1062359017030062

Pendoley, K.L., Schofield, G., Whittock, P.A., Ierodiaconou, D., Hays, G.C., 2014. Protected species use of a coastal marine migratory corridor connecting marine protected areas. Mar. Biol. 161, 1455–1466. https://doi.org/10.1007/s00227-014-2433-7

Prince, J., Hordyk, A., 2019. What to do when you have almost nothing: A simple quantitative prescription for managing extremely data-poor fisheries. Fish Fish. 20, 224–238. https://doi.org/10.1111/faf.12335

Probst, W.N., Stelzenmüller, V., Rambo, H., Moriarty, M., Greenstreet, S.P.R., 2021. Identifying core areas for mobile species in space and time: A case study of the demersal fish community in the North Sea. Biol. Conserv. 254, 108946. https://doi.org/10.1016/j.biocon.2020.108946

Quemmerais-Amice, F., Barrere, J., La Rivière, M., Contin, G., Bailly, D., 2020. A Methodology and Tool for Mapping the Risk of Cumulative Effects on Benthic Habitats. Front. Mar. Sci. 7. https://doi.org/10.3389/fmars.2020.569205

Righton, D., Westerberg, H., Feunteun, E., Økland, F., Gargan, P., Amilhat, E., Metcalfe, J., Lobon-Cervia, J., Sjöberg, N., Simon, J., Acou, A., Vedor, M., Walker, A., Trancart, T., Brämick, U., Aarestrup, K., 2016. Empirical observations of the spawning migration of European eels: The long and dangerous road to the Sargasso Sea. Sci. Adv. 2, e1501694. https://doi.org/10.1126/sciadv.1501694

Rikardsen, A.H., Righton, D., Strøm, J.F., Thorstad, E.B., Gargan, P., Sheehan, T., Økland, F., Chittenden, C.M., Hedger, R.D., Næsje, T.F., Renkawitz, M., Sturlaugsson, J., Caballero, P., Baktoft, H., Davidsen, J.G., Halttunen, E., Wright, S., Finstad, B., Aarestrup, K., 2021. Redefining the oceanic distribution of Atlantic salmon. Sci. Rep. 11, 12266. https://doi.org/10.1038/s41598-021-91137-y

Royle, J.A., Link, W.A., 2006. Generalized site occupancy models allowing for false positive and false negative errors. Ecology 87, 835–841. https://doi.org/10.1890/0012-9658(2006)87[835:GSOMAF]2.0.CO;2

Simmonds, E.G., Jarvis, S.G., Henrys, P.A., Isaac, N.J.B., O'Hara, R.B., 2020. Is more data always better? A simulation study of benefits and limitations of integrated distribution models. Ecography (Cop.). 43, 1413–1422. https://doi.org/10.1111/ecog.05146

Spiegelhalter, D.J., Best, N.G., Carlin, B.P., van der Linde, A., 2002. Bayesian measures of model complexity and fit. J. R. Stat. Soc. Ser. B (Statistical Methodol. 64, 583–639. https://doi.org/10.1111/1467-9868.00353

Stratoudakis, Y., Correia, C., Belo, A.F., de Almeida, P.R., 2020. Improving participated management under poor fishers' organization: Anadromous fishing in the estuary of Mondego River, Portugal. Mar. Policy 119, 104049. https://doi.org/10.1016/j.marpol.2020.104049

Stratoudakis, Y., Mateus, C.S., Quintella, B.R., Antunes, C., Raposo de Almeida, P., 2016. Exploited anadromous fish in Portugal: Suggested direction for

conservation and management. Mar. Policy 73, 92–99.
https://doi.org/10.1016/j.marpol.2016.07.031

Takashina, N., Mougi, A., 2014. Effects of marine protected areas on overfished fishing stocks with multiple stable states. J. Theor. Biol. 341, 64–70. https://doi.org/10.1016/j.jtbi.2013.09.027

Taverny, C., Elie, P., 2001. Répartition spatio-temporelle de la grande alose *Alosa alosa* (Linne, 1766) et de l'Alose feinte *Alosa fallax* (Lacépède, 1803) dans le Golfe de Gascogne. Bull. Français la Pêche la Piscic. 803–821. https://doi.org/10.1051/kmae:2001020

Taverny, C., Lassalle, G., Ortusi, I., Roqueplo, C., Lepage, M., Lambert, P., 2012. From shallow to deep waters: habitats used by larval lampreys (genus *Petromyzon* and *Lampetra*) over a western European basin. Ecol. Freshw. Fish 21, 87–99. https://doi.org/10.1111/j.1600-0633.2011.00526.x

Teruzzi, A., Di Biagio, V., Feudale, L., Bolzon, G., Lazzari, P., Salon, S., Di Biagio, V., Coidessa, G., and Cossarini, G., 2021. Mediterranean Sea Biogeochemical Reanalysis (CMEMS MED-Biogeochemistry, MedBFM3 system) (Version 1) [Data set]. Copernicus Monitoring Environment Marine Service (CMEMS). https://doi.org/10.25423/CMCC/MEDSEA_MULTIYEAR_BGC_006_008_MEDBFM3

Thiel, R., Salewski, V., 2003. Verteilung und Wanderung von Neunaugen im Elbeästuar (Deutschland). Limnologica 33, 214–226. https://doi.org/10.1016/S0075-9511(03)80015-4

Trancart, T., Rochette, S., Acou, A., Lasne, E., Feunteun, E., 2014. Modeling marine shad distribution using data from French bycatch fishery surveys. Mar. Ecol. Prog. Ser. 511, 181–192. https://doi.org/10.3354/meps10907

Verhelst, P., Reubens, J., Buysse, D., Goethals, P., Van Wichelen, J., Moens, T., 2021. Toward a roadmap for diadromous fish conservation: the Big Five considerations. Front. Ecol. Environ. fee.2361. https://doi.org/10.1002/fee.2361

Vieilledent, G., Merow, C., Guelat, J., Latimer, A.M., Kery, M., Gelfand, A.E., Wilson, A.M., Mortier, F., Silander Jr., J.A., 2014. hSDM: Hierarchical Bayesian Species Distribution Models. R package version 1.4. https://CRAN.R-project.org/ package=hSDM

Waldman, J.R., Quinn, T.P., 2022. North American diadromous fishes: Drivers of decline and potential for recovery in the Anthropocene. Sci. Adv. 8. https://doi.org/10.1126/sciadv.abl5486

Wauchope, H.S., Jones, J.P.G., Geldmann, J., Simmons, B.I., Amano, T., Blanco, D.E., Fuller, R.A., Johnston, A., Langendoen, T., Mundkur, T., Nagy, S., Sutherland, W.J., 2022. Protected areas have a mixed impact on waterbirds, but management helps. Nature 605, 103–107. https://doi.org/10.1038/s41586-022-04617-0

Wilson, K., and Veneranta, L. (Eds). 2019. Data-limited diadromous species – review of European status. ICES Cooperative Research Report No. 348. 273 pp. http://doi.org/10.17895/ices.pub.5253

Worm, B., Barbier, E.B., Beaumont, N., Duffy, J.E., Folke, C., Halpern, B.S., Jackson, J.B.C., Lotze, H.K., Micheli, F., Palumbi, S.R., Sala, E., Selkoe, K.A., Stachowicz, J.J., Watson, R., 2006. Impacts of Biodiversity Loss on Ocean Ecosystem Services. Science (80-.). 314, 787–790. https://doi.org/10.1126/science.1132294

Worm, B., Davis, B., Kettemer, L., Ward-Paige, C.A., Chapman, D., Heithaus, M.R.,
Kessel, S.T., Gruber, S.H., 2013. Global catches, exploitation rates, and
rebuilding options for sharks. Mar. Policy 40, 194–204.
https://doi.org/10.1016/j.marpol.2012.12.034
Worm, B., Hilborn, R., Baum, J.K., Branch, T.A., Collie, J.S., Costello, C., Fogarty,
M.J., Fulton, E.A., Hutchings, J.A., Jennings, S., Jensen, O.P., Lotze, H.K.,
Mace, P.M., McClanahan, T.R., Minto, C., Palumbi, S.R., Parma, A.M., Ricard,
D., Rosenberg, A.A., Watson, R., Zeller, D., 2009. Rebuilding global fisheries.
Science. 325, 578–85. https://doi.org/10.1126/science.1173146
Worm, B., Sandow, M., Oschlies, A., Lotze, H., Myers, R., 2005. Global patterns of
predator diversity in the open oceans. Science. 309, 1365–9.
https://doi.org/10.1126/science.1113399

1179