8   Developmental staging and future research directions of the model marine tubeworm *Hydroides*

9   *elegans*

15   Katherine T. Nesbit[1], Nicholas J. Shikuma[1*]

20   [1]Molecular Biology Division, San Diego State University

21   5500 Campanile Drive, San Diego CA, 92182

22   *Corresponding author (nshikuma@sdsu.edu)

ABSTRACT

Background: The biofouling marine tube worm, *Hydroides elegans*, is an indirect developing

polychaete with significance as a model organism for questions in developmental biology and

the evolution of host-microbe interactions. However, a complete description of the life cycle

from fertilization through sexual maturity remains scattered in the literature, and lacks

standardization. **Results:** Here we present a unified staging scheme detailing the major

morphological changes that occur during the entire life cycle of the animal. These data represent

a complete record of the life cycle, and serve as a foundation for connecting molecular changes

with morphology. **Conclusions:** The present descriptions and associated staging scheme are

especially timely as this system gains traction within research communities. Characterizing the

Hydroides life cycle is essential for investigating the molecular mechanisms that drive major

developmental transitions, like metamorphosis, in response to bacteria.

INTRODUCTION

The marine tube worm *Hydroides elegans* is an emerging model Spiralian for addressing

questions in developmental biology and probing the molecular foundations of bacterial

influences on development. *Hydroides* is a simple organismal model that is highly tractable in a

laboratory setting. They release thousands of eggs and sperm at a time, and spawning is non-

lethal. Fertilization occurs externally, and the embryos/larvae will rapidly develop in a simple

dish of sea water. Metamorphosis of *Hydroides* larvae will occur at five days, and is catalyzed by

contact with a heterogeneous bacterial biofilm. In the lab, metamorphosis can be recapitulated by

contact with axenic biofilms composed of a single stimulatory bacterial strain [8, 10, 11].

Hydroides enables single animal-single microbe investigations, ultimately facilitating

characterization of the pathways through which eukaryotic organisms sense and respond to an

array of microbial products that induce major developmental transitions.

Despite its utility as a developmental model[12-17], characterizations of the growth and

development of *Hydroides* remains fragmented across the existing literature[12, 14-18]. These

works include partial descriptions of morphological and molecular features at the cleavage

stages[14] and early embryogenesis[15], as well as formation of the feeding larval stages[16],

segmentation[12], and emphasis on metamorphosis and juvenile life stages[8, 9, 17-19]. Yet, the

community has not established a standardized staging scheme that spans the entirety of

development. To this end, we provide here a complete staging of the life cycle of *Hydroides*

(Figure 1) from fertilization through sexual maturation, as well as offer perspectives on how this

model can be leveraged to advance our understanding of diverse and fundamental biological

questions.

EVOLUTION AND NATURAL HISTORY OF HYDROIDES

Annelids are a diverse and abundant group of organisms, and are part of a larger

evolutionary clade known as the Spiralia[39]. Spiralians are of particular interest in evolution

and development because of the vast diversity of bilateral body plans they exhibit throughout

ontogeny, which arise from a highly conserved embryonic cleavage pattern. The Annelids

represent an under-sampled and understudied source of Spiralian variation in the evolution of

development, despite having been shown to exhibit several important developmental differences

to molluscs[38, 40]. Existing annelid models with developmental staging schemes include the

leech – *Helobdella*[26], the ragworm – *Platynereis*[27], the parchment worm - *Chaetopterus*[28],

Capitella[12], and *Owenia*[29]. As a Spiralian developmental model, *Hydroides* can help us

understand the evolution of innovation and conservation in development at the cellular and

molecular level, as well as how this intersects with environmental input in development within

and among major metazoan lineages.

Adult *Hydroides* (Figure 1A) are protandrous hermaphrodites, and are found in intertidal

to sublittoral zones throughout the global tropical and subtropical oceans[41, 42]. They are part

of a larger group of tube-building annelids in the family Serpulidae. These animals are filter

feeders, extending branching appendages, collectively called the branchial crown, out from the

opening of their tubes to collect particles from the water. In addition, they have a special funnel-

shaped structure called an operculum that closes off the opening to the tube when the animal

retracts inside. As one of the most prolific biomineralizing annelids, this species has garnered

attention as a biofouling organism that deposits its calcareous tubes on ship hulls and other

submerged surfaces in harbors and marinas[11, 42, 43]. They are resilient, tolerating variable

temperatures (15-30°C) and salinities (15-37 PSU)[13], and able to colonize surfaces treated

with different antifouling agents[44, 45]. However, despite being seen as "pests" in harbors and

marinas, these animals are valuable research models. They, along with other members in the

genus, have been utilized for biological research since the species was first described over 100

years ago[46], with increasing interest over the last few decades. This surge in interest in

Hydroides centers around antifouling and the biphasic life history (Figure 1B) of this animal

which utilizes microbial intervention to complete development[8].

DEVELOPMENT

Unlike the gametes of other marine invertebrates, for example sea urchins, there is no

obvious morphological changes to the eggs of *Hydroides* upon fertilization (Figure 2). The early

cleavages of *Hydroides* (Figure 3) are equal-sized[14]. Unlike other members of the Spiralia and

even other Annelids[14, 37, 38], the "D" quadrant of the embryo is not morphologically distinct

from the other regions at the 4-cell stage (Figure 3B). The third cleavage, yielding an 8-cell

embryo (Figure 3C), is the first division during which the spiral pattern becomes apparent. This

division is a sinistral (left-handed) twist[14]. The cleavage spindle rotates, and as a result the

daughter cells are born on the other side of the macromere, giving the appearance that the

blastomeres on the animal side of the embryo have rotated counter-clockwise relative to the

vegetal blastomeres[14]. The 8 cells of the embryo complete the next cleavage cycle

concurrently, yielding the 16-cell stage (Figure 3D). This is in contrast to the blastomere

divisions leading from 16- to 32-cell stage (Figure 3E), and the 32- to 64-cell stage (Figure 3F)

transition which do not occur synchronously[14].

The formation of the gut (Figure 4) via invagination at the vegetal pole[16, 47] begins

shortly after the completed cleavages of the 64-cell stage embryo, now called a blastula (Figure

4A). The embryos also become ciliated (Figure 4B-C, white arrows) and hatch to begin their

free-swimming stages, forming a characteristic apical tuft (Figure 4C, yellow arrow) typical of

many marine larvae, along with a thick ciliary band that will eventually form the prototroch[48].

Prior work has demonstrated that the blastopore partially closes along the ventral midline

forming the mouth[16], and the developing gut tube traverses the blastocoel, bending towards the

aboral side of the animal[16]. Eventually the epithelium of the developing gut fuses with the

aboral ectoderm to form the anus[15, 16, 47, 48]. The complete gut is tripartite, having a

differentiated fore-, mid-, and hindgut. Feeding of the trochophore larva (Figure 4D) can then

begin at approximately 12 hours post-fertilization (hpf)[17]. The beating of the plush ciliary band

forming the prototroch, as well as the opposing accessory band, the metatroch, helps to direct

food particles towards the mouth.

Over the course of several days, the trochophore larva feeds. Two eye spots form in

sequence. The pigmentation of the first eye spot is visible as early 11 hours post-fertilization,

followed by the second which is clearly visible by 72 hpf, and the posterior portion of the body

elongates[12]. Three segments arise nearly simultaneously on the posterior portion of the body,

each segment develops chaetal sacs and parapodia[12]. The larva is now a segmented

nectochaete (Figure 4E), and continues to feed, though no additional segments are added to the

body until after metamorphosis. Upon reaching competency (Figure 4F) between 5-6 days post-

fertilization, the larva has three complete body segments and looks morphologically distinct from

the pre-competent larva. In short, competent larvae are larger in overall body size, and the body

profile from the prototroch to the tail is more cone-like in comparison to pre-competent larvae

which have a more rhomboid appearance. The head is more clearly defined from the body in

competent larvae, because there is not any tapering or gradual sloping at the anterior region of

the body, unlike in the pre-competent larvae (Figure 4E). The competent larva is then ready to

settle on a substrate and undergo metamorphosis.

138   Metamorphosis is induced by contact with heterogeneous multispecies or monospecies

biofilms[7, 8, 10, 49]. Settlement in response to a natural seawater-derived biofilm is

characterized by a slowing of swimming speed, and probing of a surface[17]. While probing the

settlement surface, the larva makes transient, reversible contact with the surface and appears to

crawl along it, while wiggling or flexing the body by contracting the muscles along the body

wall before resuming a slow swimming speed. If suitable, the larva attaches (Figure 5A) itself to

a surface and begins to secrete a primary proteinaceous tube[19] (Figure 5B, dashed line) and

shed the prototroch cilia (Figure 5C, noted with "*") which are no longer needed for swimming.

Major tissue remodeling occurs as the anterior region of the larva's body changes shape to form

a collar (Figure 5D), food groove cells between the prototroch and metatroch are shed (Figure

5E), and cilia from the metatroch are lost (Figure 5E) as lobes extended laterally (Fig 5F-G) on

the head of the animal beginning formation of the branchial crown, and a mineralized secondary

tube is deposited.

151   Post-metamorphic juveniles (Figure 5H-I) continue to elaborate the anterior portion of

the body, forming multiple radioles on the branchial crown for feeding (Figure 5I), as well as the

operculum in 10-14 days. They continue to deposit their calcareous tubes and within a week of

metamorphosis grow to several millimeters in length (Figure 5I, Figure 6). Under ideal culturing

conditions with ample food and clean, well oxygenated water, they have a generation time of

approximately three to four weeks[11]. In the wild, they settle gregariously, forming dense

communities in a matter of weeks, and can be cultured as single tubes in the lab to isolate

individual sexually mature adults (Figure 1A).

BACTERIA-STIMULATED METAMORPHOSIS OF HYDROIDES

161   The role of microbes in mediating animal development is of increasing interest. This is in

part due to the diversity of organisms that participate in host-microbe relationships, as well as the

diversity of developmental outcomes that arise from interactions with bacteria. The transition to

multicellularity[1] as well as switches between asexual to sexual reproductive strategies[2] in

choanoflagellates, maturation of the gut in zebrafish[3] and mammals[4], organogenesis in the

Hawaiian bobtail squid[5], and initiation of metamorphosis in marine invertebrates[6, 7]

including *Hydroides*[8, 9] are just some of the varied ways in which microbial interactions

influence eukaryotic development.

169   Given the remarkable impact that marine bacteria have on the life cycle of *Hydroides*, it

comes as no surprise that much attention has been given to characterizing the nature of this host-

microbe association during metamorphosis. Many different single strains of bacteria induce

Hydroides metamorphosis[8]. For example, the Gammaproteobacterium, *Pseudoalteromonas*

luteoviolacea, and the Bacteroidetes, *Cellulophaga lytica* [50]. The diverse inductive bacteria

also produce diverse bacterial products that promote the same developmental endpoints[51]. For

example, *P. luteoviolacea* produces a contractile injection system[52] that is related to the

contractile tails of bacterial viruses (bacteriophage). These syringe-like structures are assembled

into arrays and deliver a protein payload[53], the effector Mif1, to *Hydroides* larvae which then

interacts with host-derived signaling networks[54] to induce metamorphosis. Other bacteria

stimulate metamorphosis through different mechanisms[51]. For example, lipopolysaccharide

from outer membrane vesicles of some bacteria induce *Hydroides* metamorphosis[50]. These

biochemically unique products that induce metamorphosis from diverse bacteria open up a future

for discovery. The molecular mechanisms driving metamorphosis in *Hydroides* are likely to

reveal fundamental means of interaction at play between hosts and microbes across the tree of

life.

185   Working in the *Hydroides* model system raises interesting questions about the nature of

this marine tubeworm-microbe relationship: 1) do other diverse bacterial cues for metamorphosis

exist?; 2) given the heterogeneity of marine biofilms, what are the respective contributions of

individual bacteria within it to metamorphosis, effectively how potent are the various cues, are

there combinatorial effects of these products, and are they distinct enough to be differentiated by

the animal?; and 3) do these diverse bacterial cues actually yield identical developmental

outcomes, or does the mechanism of metamorphosis initiation influence the sequence of

signaling and morphogenetic events[17], and if so, how?

THE FUTURE OF HYDROIDES

195   As the *Hydroides* model continues to gain traction in the research community, it is clear

there are many areas with potential for discovery. This includes comparative evolutionary and

developmental biology, genetic tool development, biotechnology, and the discovery of bacteria-
derived natural products.

Hydroides can offer unique perspectives on the establishment of host-microbe
interactions. Developmental regulatory networks have yet to be integrated with components from
microbial symbionts. In fact, the animal sensing machinery responsible for detecting and
responding to bacteria by initiating metamorphosis is not well-resolved for *Hydroides*, or any
other animal aside from the highly conserved signaling systems (e.g., PKC and MAPK[9, 54-
57]). Thus, there exists a conspicuous gap in knowledge that this Annelid model is primed to fill.

Comparative approaches also allow for the investigation of overlap or repurposing of
molecular machinery that drives metamorphosis and facilitates host-microbe interactions. For
example, immune genes have rapidly evolved and function primarily to detect and respond to
encounters with bacteria. However, repurposing of these components could facilitate interactions
with non-pathogenic bacteria and intersect with developmental pathways like those activated
during metamorphosis[9, 58]. The *Hydroides* genome contains more genes in common with
humans than *C. elegans* does[9], placing *Hydroides* as a model system with the potential for
discoveries relating to human development, health, and disease.

The *Hydroides* model sits squarely at the junction of questions in ecology, evolution, and
development. As a broadcast spawning marine invertebrate that has a vast dispersal range, we
can gain insight into how animal ecology is influenced by environmental microbes that stimulate
metamorphosis. This would be applicable, for example, in reef environments where recruitment
and successful settlement of dispersed pelagic larvae is essential for habitat maintenance. These
eco-evo-devo perspectives[59] may illuminate the evolution of host-microbe symbioses, and also
add a new layer of context to ecosystem dynamics and management for vulnerable marine
habitats.

A clear and current bottleneck in the development of *Hydroides* as a model system for
host-microbe interactions is a lack of established techniques for genetically manipulating the
Hydroides host. There has been recent success in the genetic manipulation of related Annelids
such as *Platynereis dumerilii*[60, 61] and *Capitella teleta*[62, 63] and we are optimistic that
Hydroides will be amenable to genetic manipulation. Improvements in husbandry and long-term
culture will facilitate generation of animal lines in *Hydroides*, enabling *Hydroides* to keep pace
with advancements in other marine invertebrate larvae[64-69]. In parallel, emerging synthetic

biology tools in established model microbes also hold promise for the manipulation of more

diverse marine bacteria that promote *Hydroides* development[70]. These new tools compliment

powerful classical bacterial genetics approaches which have already helped reveal key aspects of

Hydroides interaction with stimulatory microbes[52, 53].

Future work leveraging *Hydroides* holds promise for promoting biotechnology in the

contexts of evolution, the environment, and human health. For example, insights gained from

Hydroides to understand bacteria-stimulated metamorphosis could aid in the husbandry of

invertebrates in aquaculture[71]. *Hydroides* has been studied intensively as a biofouling

organism[11] and research on antifouling technologies could provide significant economic relief

for shipping and naval sectors[43]. The products bacteria produce to promote *Hydroides*

metamorphosis and the machinery *Hydroides* uses to sense these products could lead to broader

discoveries about mechanisms and tools for host-microbe interactions, and how these

interactions evolve on a genetic and chemical level. For example, we found that gene clusters

encoding Contractile Injection Systems originally discovered to promote *Hydroides*

metamorphosis are strikingly similar to gene clusters found in *Bacteroidales* bacteria from

healthy human microbiomes[72]. Moreover, understanding the payload delivery and targeting

mechanism of the MACs produced by *P. luteoviolacea* could enable new types of therapeutic

delivery systems[73].

CONCLUSION

Environmental bacteria are becoming increasingly recognized for their significant roles in

regulating health and development[6, 74]. The utilization of the *Hydroides*–microbe model offers

opportunities in the eco-evo-devo landscape to understand the conservation and diversity of

microbial drivers of eukaryotic development and disease states.

EXPERIMENTAL PROCEDURES

Husbandry, Spawning and Fertilization

Adult worms are collected from Quivira Basin in Mission Bay, San Diego CA. Animals

are housed in a suspended basket in a recirculating tank at room temperature, supplied with

artificial filtered sea water (AFSW) at 35 PSU. Adults are fed *ad libidum* with a monoculture of

the marine algae *Isochrysis galbana* (Carolina Biological: Item #153180). To spawn animals,

individual adult tubes are selected and placed in separate glass petri dishes (60mm x 15mm)

filled with AFSW. Using forceps, the tubes are cracked, revealing the animal inside which

releases thousands of eggs or streams of sperm upon disturbance.

Eggs are collected with a trimmed plastic pipette tip and washed into fresh FASW to

minimize debris left over from tube fracture. Sperm is collected with a pipette in minimal sea

water, and then diluted further into a fresh sperm suspension for fertilization. Eggs are diluted to

~500/mL and 4-5 drops of sperm suspension is added. The dish is swirled to mix. Ten minutes

after the addition of sperm to the eggs, the eggs are collected and washed into fresh FASW three

times to eliminate excess sperm and minimize the risk of polyspermy. Eggs and embryos are

monitored over the next several hours to ensure fertilization since no obvious morphological

changes occur at fertilization. All cultures of embryos are maintained at a density of 1 larva/ml in

an incubator at 25°C, with foil covers to prevent evaporation. Larval cultures have water changes

daily starting at Day 2, and are fed *I. galbana* daily at a density of $6x10^4$ cells/mL. At 5 days

post-fertilization, the majority of larvae have reached competency (there is some asynchrony due

to variation in individual larval feeding efficiency) and are transferred to a dish that has a natural

biofilm coating the bottom of the dish. The natural biofilm is generated by filling the dish with

raw seawater from the worm's natural habitat and allowing that to sit for four to five days. The

worms metamorphose over the course of ~12 hours upon interacting with the natural biofilm,

and are then either imaged or kept for longer-term culture to adulthood. Metamorphosed worms

that are kept for culture to adulthood are fed *I. galbana* every other day *ad libidum*, and water

changes are completed on an identical schedule.

Live-cell Imaging of Development

All stages except the adult of *Hydroides elegans* were imaged live on a Zeiss Axio

Observer.Z1 using objectives ranging from 5x to 100x oil immersion, all with differential

interference contrast (DIC) optics. Images were captured using the ZEN software suite, and

measurements were added using Fiji (National Institutes of Health, Bethesda, Maryland)[16].

Composite DIC images were rendered from Z-stacks of animals using Helicon Focus Pro

Unlimited (v6.8.0, Helicon Soft Ltd.). Adult animals were imaged live on a Wild Heerbrugg

M5a stereo microscope fitted with a QImaging R3 Retiga CCD camera.

Acknowledgements

We would like to thank the members of the Shikuma Lab for constructive feedback on the

present manuscript. This work was supported by the National Science Foundation (1942251), the

Gordon and Betty Moore Foundation (GBMF9344; https://doi.org/10.37807/GBMF9344), Office

of Naval Research (N00014-20-1-2120), the National Institutes of Health, NIGMS

(R35GM146722) and the Alfred P. Sloan Foundation, Sloan Research Fellowship.

REFERENCES

1. Alegado, R.A., et al., *A bacterial sulfonolipid triggers multicellular development in the closest living relatives of animals.* elife, 2012. **1**: p. e00013.
2. Woznica, A., et al., *Mating in the closest living relatives of animals is induced by a bacterial chondroitinase.* Cell, 2017. **170**(6): p. 1175-1183. e11.
3. Bates, J.M., et al., *Distinct signals from the microbiota promote different aspects of zebrafish gut differentiation.* Developmental biology, 2006. **297**(2): p. 374-386.
4. Hooper, L.V., *Bacterial contributions to mammalian gut development.* Trends in microbiology, 2004. **12**(3): p. 129-134.
5. McFall-Ngai, M.J. and E.G. Ruby, *Symbiont recognition and subsequent morphogenesis as early events in an animal-bacterial mutualism.* Science, 1991. **254**(5037): p. 1491-1494.
6. Woznica, A. and N. King, *Lessons from simple marine models on the bacterial regulation of eukaryotic development.* Current opinion in microbiology, 2018. **43**: p. 108-116.
7. Cavalcanti, G.S., et al., *The influence of bacteria on animal metamorphosis.* Annual review of microbiology, 2020. **74**: p. 137-158.
8. Unabia, C. and M. Hadfield, *Role of bacteria in larval settlement and metamorphosis of the polychaete Hydroides elegans.* Marine Biology, 1999. **133**(1): p. 55-64.
9. Shikuma, N.J., et al., *Stepwise metamorphosis of the tubeworm Hydroides elegans is mediated by a bacterial inducer and MAPK signaling.* Proceedings of the National Academy of Sciences, 2016. **113**(36): p. 10097-10102.
10. Lau, S.C., et al., *Bioactivity of bacterial strains isolated from marine biofilms in Hong Kong waters for the induction of larval settlement in the marine polychaete Hydroides elegans.* Marine Ecology Progress Series, 2002. **226**: p. 301-310.
11. Nedved, B.T. and M.G. Hadfield, *Hydroides elegans (Annelida: Polychaeta): a model for biofouling research.* 2008, Springer.
12. Seaver, E.C., K. Thamm, and S.D. Hill, *Growth patterns during segmentation in the two polychaete annelids, Capitella sp. I and Hydroides elegans: comparisons at distinct life history stages.* Evolution & development, 2005. **7**(4): p. 312-326.
13. Qiu, J.-W. and P.-Y. Qian, *Combined effects of salinity, temperature and food on early development of the polychaete Hydroides elegans.* Marine Ecology Progress Series, 1997. **152**: p. 79-88.
14. Arenas-Mena, C., *Sinistral equal-size spiral cleavage of the indirectly developing polychaete Hydroides elegans.* Developmental dynamics: an official publication of the American Association of Anatomists, 2007. **236**(6): p. 1611-1622.
15. Arenas-Mena, C., *Brachyury, Tbx2/3 and sall expression during embryogenesis of the indirectly developing polychaete Hydroides elegans.* International Journal of Developmental Biology, 2013. **57**(1): p. 73-83.
16. Arenas-Mena, C. and A. Li, *Development of a feeding trochophore in the polychaete Hydroides elegans.* International Journal of Developmental Biology, 2015. **58**(6-7-8): p. 575-583.
17. Hadfield, M.G., M.L. Freckelton, and B.T. Nedved, *The natural sequence of events in larval settlement and metamorphosis of Hydroides elegans (Polychaeta; Serpulidae).* Plos one, 2021. **16**(5): p. e0249692.
18. Huggett, M.J., et al., *Formation and Function of the Primary Tube During Settlement and Metamorphosis of the Marine Polychaete Hydroides elegans (Haswell, 1883)(Serpulidae).* The Biological Bulletin, 2021. **240**(2): p. 82-94.

19.   Lau, S.C., T. Harder, and P.Y. Qian, *Induction of larval settlement in the serpulid polychaete*
Hydroides elegans (Haswell): role of bacterial extracellular polymers. Biofueling, 2003. **19**(3): p.
197-204.
20.   Nieuwkoop, P.D., *Normal table of Xenopus laevis (Daudin).* Normal table of Xenopus laevis
(Daudin), 1956: p. 162-203.
21.   Hamburger, V. and H.L. Hamilton, *A series of normal stages in the development of the chick*
embryo. Journal of morphology, 1951. **88**(1): p. 49-92.
22.   Kimmel, C.B., et al., *Stages of embryonic development of the zebrafish.* Developmental
dynamics, 1995. **203**(3): p. 253-310.
23.   Heyland, A. and J. Hodin, *A detailed staging scheme for late larval development in*
Strongylocentrotus purpuratus focused on readily-visible juvenile structures within the rudiment.
BMC developmental biology, 2014. **14**(1): p. 22.
24.   Smith, M.M., et al., *The larval stages of the sea urchin, Strongylocentrotus purpuratus.* Journal of
Morphology, 2008. **269**(6): p. 713-733.
25.   Nesbit, K.T. and A. Hamdoun, *Embryo, larval, and juvenile staging of Lytechinus pictus from*
fertilization through sexual maturation. Developmental Dynamics, 2020. **249**(11): p. 1334-1346.
26.   Fernández, J., *Embryonic development of the glossiphoniid leech Theromyzon rude:*
characterization of developmental stages. Developmental biology, 1980. **76**(2): p. 245-262.
27.   Fischer, A.H., T. Henrich, and D. Arendt, *The normal development of Platynereis dumerilii*
(Nereididae, Annelida). Frontiers in zoology, 2010. **7**(1): p. 1-39.
28.   Irvine, S.Q., O. Chaga, and M.Q. Martindale, *Larval ontogenetic stages of Chaetopterus:*
developmental heterochrony in the evolution of chaetopterid polychaetes. The Biological
Bulletin, 1999. **197**(3): p. 319-331.
29.   Carrillo-Baltodano, A.M., et al., *Early embryogenesis and organogenesis in the annelid Owenia*
fusiformis. EvoDevo, 2021. **12**(1): p. 1-18.
30.   Gabriel, W.N., et al., *The tardigrade Hypsibius dujardini, a new model for studying the evolution*
of development. Developmental biology, 2007. **312**(2): p. 545-559.
31.   Browne, W.E., et al., *Stages of embryonic development in the amphipod crustacean, Parhyale*
hawaiensis. genesis, 2005. **42**(3): p. 124-149.
32.   Montague, T.G., I.J. Rieth, and R. Axel, *Embryonic development of the camouflaging dwarf*
cuttlefish, Sepia bandensis. Developmental Dynamics, 2021. **250**(12): p. 1688-1703.
33.   Podrabsky, J.E., et al., *Embryonic development of the annual killifish Austrofundulus limnaeus:*
An emerging model for ecological and evolutionary developmental biology research and
instruction. Developmental Dynamics, 2017. **246**(11): p. 779-801.
34.   Griffing, A.H., et al., *Embryonic development of a parthenogenetic vertebrate, the mourning*
gecko (Lepidodactylus lugubris). Developmental Dynamics, 2019. **248**(11): p. 1070-1090.
35.   Diaz Jr, R.E., et al., *Filling in the phylogenetic gaps: induction, migration, and differentiation of*
neural crest cells in a squamate reptile, the veiled chameleon (Chamaeleo calyptratus).
Developmental Dynamics, 2019. **248**(8): p. 709-727.
36.   Adamson, C.J., N. Morrison-Welch, and C.D. Rogers, *The amazing and anomalous axolotls as*
scientific models. Developmental Dynamics, 2022.
37.   Lambert, J.D., *Developmental patterns in spiralian embryos.* Current Biology, 2010. **20**(2): p. R72-
R77.
38.   Seaver, E.C., *Variation in spiralian development: insights from polychaetes.* International Journal
of Developmental Biology, 2014. **58**(6-7-8): p. 457-467.
39.   Henry, J.Q., *Spiralian model systems.* International Journal of Developmental Biology, 2015.
58(6-7-8): p. 389-401.

40. Seaver, E.C., *Annelid models I: Capitella teleta*. Current Opinion in Genetics & Development,
2016. **39**: p. 35-41.
41. Bastida-Zavala, J.R., et al., *The fouling serpulids (Polychaeta: Serpulidae) from United States*
coastal waters: an overview. European Journal of Taxonomy, 2017(344).
42. Pettengill, J., et al., *Biofouling likely serves as a major mode of dispersal for the polychaete*
tubeworm Hydroides elegans as inferred from microsatellite loci. Biofouling, 2007. **23**(3): p. 161-
169.
43. Schultz, M., et al., *Economic impact of biofouling on a naval surface ship.* Biofouling, 2011. **27**(1):
p. 87-98.
44. Dafforn, K., T. Glasby, and E. Johnston, *Differential effects of tributyltin and copper antifoulants*
on recruitment of non-indigenous species. Biofouling, 2008. **24**(1): p. 23-33.
45. Piola, R.F., K.A. Dafforn, and E.L. Johnston, *The influence of antifouling practices on marine*
invasions. Biofouling, 2009. **25**(7): p. 633-644.
46. Haswell, W. *On some new Australian tubicolous annelids.* in *Proceedings of the Linnean Society*
of New South Wales. 1883.
47. Arenas-Mena, C. and K.S.-Y. Wong, *HeOtx expression in an indirectly developing polychaete*
correlates with gastrulation by invagination. Development genes and evolution, 2007. **217**(5): p.
373-384.
48. Arenas-Mena, C., K.S.-Y. Wong, and N. Arandi-Forosani, *Ciliary band gene expression patterns in*
the embryo and trochophore larva of an indirectly developing polychaete. Gene Expression
Patterns, 2007. **7**(5): p. 544-549.
49. Shikuma, N. and M. Hadfield, *Temporal variation of an initial marine biofilm community and its*
effects on larval settlement and metamorphosis of the tubeworm Hydroides elegans. Biofilms,
2005. **2**(4): p. 231-238.
50. Freckelton, M.L., et al., *Bacterial lipopolysaccharide induces settlement and metamorphosis in a*
marine larva. Proceedings of the National Academy of Sciences, 2022. **119**(18): p. e2200795119.
51. Freckelton, M.L., B.T. Nedved, and M.G. Hadfield, *Induction of invertebrate larval settlement;*
different bacteria, different mechanisms? Scientific Reports, 2017. **7**(1): p. 1-11.
52. Shikuma, N.J., et al., *Marine tubeworm metamorphosis induced by arrays of bacterial phage*
tail–like structures. Science, 2014. **343**(6170): p. 529-533.
53. Ericson, C.F., et al., *A contractile injection system stimulates tubeworm metamorphosis by*
translocating a proteinaceous effector. Elife, 2019. **8**: p. e46845.
54. Malter, K.E., et al., *Diacylglycerol, PKC and MAPK signaling initiate tubeworm metamorphosis in*
response to bacteria. Developmental Biology, 2022. **487**: p. 99-109.
55. Yamamoto, H., et al., *Protein kinase C (PKC) signal transduction system involved in larval*
metamorphosis of the barnacle, Balanus amphitrite. Zoological science, 1995. **12**(4): p. 391-396.
56. Amador-Cano, G., E. Carpizo-Ituarte, and D. Cristino-Jorge, *Role of protein kinase C, G-protein*
coupled receptors, and calcium flux during metamorphosis of the sea urchin Strongylocentrotus
purpuratus. The Biological Bulletin, 2006. **210**(2): p. 121-131.
57. Chambon, J.-P., et al., *ERK-and JNK-signalling regulate gene networks that stimulate*
metamorphosis and apoptosis in tail tissues of ascidian tadpoles. 2007.
58. Davidson, B. and B.J. Swalla, *A molecular analysis of ascidian metamorphosis reveals activation*
of an innate immune response. 2002.
59. Gilbert, S.F., T.C. Bosch, and C. Ledón-Rettig, *Eco-Evo-Devo: developmental symbiosis and*
developmental plasticity as evolutionary agents. Nature Reviews Genetics, 2015. **16**(10): p. 611-
622.

60.   Andreatta, G., et al., *Corazonin signaling integrates energy homeostasis and lunar phase to*
regulate aspects of growth and sexual maturation in Platynereis. Proceedings of the National
Academy of Sciences, 2020. **117**(2): p. 1097-1106.
61.   Veedin Rajan, V.B., et al., *Seasonal variation in UVA light drives hormonal and behavioural*
changes in a marine annelid via a ciliary opsin. Nature ecology & evolution, 2021. **5**(2): p. 204-
218.
62.   Neal, S., D. De Jong, and E. Seaver, *CRISPR/CAS9 mutagenesis of a single r-opsin gene blocks*
phototaxis in a marine larva. Proceedings of the Royal Society B, 2019. **286**(1904): p. 20182491.
63.   Klann, M. and E.C. Seaver, *Functional role of pax6 during eye and nervous system development*
in the annelid Capitella teleta. Developmental biology, 2019. **456**(1): p. 86-103.
64.   Nesbit, K.T., et al., *The painted sea urchin, Lytechinus pictus, as a genetically-enabled*
developmental model. Methods Cell Biol, 2019. **150**: p. 105-123.
65.   Vyas, H., et al., *Generation of a homozygous mutant drug transporter (ABCB1) knockout line in*
the sea urchin,   Lytechinus pictus. Development, 2021(*In preparation.*).
66.   Pickett, C. and R.W. Zeller, *Efficient genome editing using CRISPR-Cas-mediated homology*
directed repair in the ascidian Ciona robusta. genesis, 2018. **56**(11-12): p. e23260.
67.   Sanders, S.M., et al., *CRISPR/Cas9-mediated gene knockin in the hydroid Hydractinia*
symbiolongicarpus. BMC genomics, 2018. **19**(1): p. 1-17.
68.   Nakanishi, N. and M.Q. Martindale, *CRISPR knockouts reveal an endogenous role for ancient*
neuropeptides in regulating developmental timing in a sea anemone. Elife, 2018. **7**: p. e39742.
69.   Henry, J.Q., M.P. Lesoway, and K.J. Perry, *An automated aquatic rack system for rearing marine*
invertebrates. BMC biology, 2020. **18**(1): p. 1-22.
70.   Alker, A.T., et al., *A modular plasmid toolkit applied in marine Proteobacteria reveals functional*
insights during bacteria-stimulated metamorphosis. bioRxiv, 2023: p. 2023.01. 31.526474.
71.   Qian, P.-Y., et al., *Marine biofilms as mediators of colonization by marine macroorganisms:*
implications for antifouling and aquaculture. Marine Biotechnology, 2007. **9**(4): p. 399-410.
72.   Rojas, M.I., et al., *A distinct contractile injection system gene cluster found in a majority of*
healthy adult human microbiomes. Msystems, 2020. **5**(4): p. e00648-20.
73.   Rocchi, I., et al., *A bacterial phage tail-like structure kills eukaryotic cells by injecting a nuclease*
effector. Cell Reports, 2019. **28**(2): p. 295-301. e4.
74.   McFall-Ngai, M.J., *Unseen forces: the influence of bacteria on animal development.*
Developmental biology, 2002. **242**(1): p. 1-14.
466

467
468 **Figure 1**. *Hydroides* morphology and life cycle. A) Sexually mature adult dissected out of the
469 calcareous tube. Labelling: ab=abdomen, bc=branchial crown, c=collar, op=operculum,
470 pd=peduncle, pg=pygidium, r=radiole, th=thorax. B) Illustration of the major life history stages
471 in the biphasic life cycle of Hydroides elegans. Metamorphosis is induced by interaction with
472 single- or multi-species biofilms. Scale = 1mm.
473

474
475 **Figure 2**. Gametes and zygote of *Hydroides elegans*. A) Sperm. Scale=5µm. B) Unfertilized
476 egg. Scale=10µm. C) Zygote. No obvious morphological changes occur between the oocyte and
477 the zygote. Scale=10µm.
478

479
Figure 3. Embryonic cleavage stages in *Hydroides* elegans. All panels are depicted with the
animal pole of the embryo pointing out of the page, and the vegetal pole pointing in to the page.
Panels A-B and E-F are single equatorial focal planes. Panels C-D are composite images from
multiple focal planes. A) 2-cell embryo. B) 4-cell embryo. C) 8-cell embryo. D) 16-cell embryo.
The vegetal blastomeres are not in view. E) 32-cell embryo. F) 60-cell embryo. All panels
scale=10μm.
486

487
Figure 4. Gastrulation and larval stages in *Hydroides elegans*. A) Ciliated blastula. Dashed white
line outlines the blastocoel. B) Mid-gastrula, oral side of the animal is pointing down. White
arrows indicate ciliated cells of the developing prototroch. C) Late gastrula, hatched. White
arrows mark ciliated cells of the developing prototroch. Yellow arrow indicates cilia of the apical
tuft. D) Lateral view of a feeding trochophore larva. E) Aboral view of a segmented nectochaete
larva. F) Oral view of a metamorphically competent larva. Labelling for panels D-F: at=apical
tuft, aso=apical sensory organ, av=anal vacuole, ch=newly formed chaetae, col=collar, e=eye
spot, it=intestinal tract, m=mouth, mt=metatroch, pt=prototroch, s=larval segments with chaetae,
st=stomach. For panels A-C scale=10µm, for panels D-F scale=50µm. Panels D-F are from
composite images from multiple focal planes.
498

499
Figure 5. Morphological changes associated with bacteria-stimulated metamorphosis in
Hydroides elegans. A) Attachment to substrate. B) Formation of the primary tube (indicated by a
dashed white line on one side of the larva). C) Shedding of prototroch cilia (*). D) Collar
eversion (arrow). E) Shedding of food groove cells (arrow). F) Beginning of lobe formation
(dashed white line). G) Lateral extension and elaboration of anterior lobes. H) Juvenile at 24
hours post-metamorphosis. I) Juvenile at 1 week post metamorphosis. Inset of head and
elaboration of branchial crown. For panels A-H, scale = 50 μm. For panel I, scale = 250 μm,
inset scale = 100 μm.
508

509
Figure 6. Summary of *Hydroides* development from fertilization through sexual maturation.
Major phases of development are labelled with individual stages. Images are not to scale.
Metamorphosis stages are labelled M1-M7 and correspond to attachment to substrate (M1),
formation of the primary tube (M2), shedding of prototroch cilia (M3), collar eversion (M4),
shedding of food groove cells (M5), beginning of lobe formation (M6), lateral extension and
elaboration of anterior lobes (M7).