A High Fraction of Oral Bacteria in the Feces Indicates Gut Microbiota Depletion with Implications for Human Health

Chen Liao[1,*], Thierry Rolling[2,3,4,*], Ana Djukovic[1,*], Teng Fei[5], Vishwas Mishra[1,6], Hongbin Liu[7], Chloe Lindberg[1], Lei Dai[7], Bing Zhai[2,3,7], Jonathan U. Peled[8,9], Marcel R.M. van den Brink[8,9], Tobias M. Hohl[2,3,9,#], and Joao B. Xavier[1,#]

[1]Program for Computational and Systems Biology, Memorial Sloan-Kettering Cancer Center, New York, NY 10065, USA

[2]Infectious Disease Service, Department of Medicine, Memorial Sloan Kettering Cancer Center, New York, NY 10065, USA

[3]Immunology Program, Sloan Kettering Institute, Memorial Sloan Kettering Cancer Center, New York, NY 10065, USA

[4]Division of Infectious Diseases, First Department of Medicine, University Medical Center, Hamburg-Eppendorf, Hamburg 20251, Germany

[5]Department of Epidemiology and Biostatistics, Memorial Sloan Kettering Cancer Center, New York, NY 10065, USA

[6]Physiology, Biophysics and Systems Biology Program, Weill Cornell Medical College, New York, NY 10065, USA

[7]CAS Key Laboratory of Quantitative Engineering Biology, Shenzhen Institute of Synthetic Biology, Shenzhen Institute of Advanced Technology, Chinese Academy of Sciences, Shenzhen 518055, China

[8]Adult Bone Marrow Transplantation Service, Department of Medicine, Memorial Sloan Kettering Cancer Center, New York, NY 10065, USA

[9]Weill Cornell Medical College, New York, NY 10065, USA

*These authors contributed equally to this work

#Correspondence: xavierj@mskcc.org (J.B.X.), hohlt@mskcc.org (T.M.H.)

Abstract

The increased relative abundance of oral bacteria detected in fecal samples has been associated with intestinal diseases and digestive disorders. This observation raises two competing hypotheses: either oral bacteria invade the gut bacterial population and expand in the intestine (the *Expansion* hypothesis), or oral bacteria transit through and their relative increase in feces marks a depletion of the gut bacterial population (the *Marker* hypothesis). To address this, we conducted a comprehensive analysis of quantitative microbiome data from mouse experiments and diverse patient cohorts. Our findings consistently support the *Marker* hypothesis as the primary explanation. We further establish a robust inverse correlation between the total fraction of oral bacteria and decreased total bacterial abundance in feces. This correlation underlies the associations between the oral bacterial fraction and multiple patient outcomes consistent with a depleted gut microbiota. By distinguishing between the two hypotheses, our study guides the interpretation of microbiome compositional data and their links with human health.

Introduction

The microbiomes inhabiting different healthy human body regions have distinct bacterial populations, which reflect the unique characteristics of each ecological niche[1]. Despite occasional exchange among these sites, their resident bacterial populations remain distinctive. For example, humans swallow 1.5×10^{12} salivary bacteria per day[2], some of which can travel to the lower gastrointestinal tract[3-5] through the enteral[6,7] and hematogenous[8] routes. Gastric acids and antimicrobial peptides kill many oral travelers, but those that survive the journey must overcome colonization resistance from gut resident bacteria, which are better competitors in the gut environment[7]. Consequently, oral bacteria are typically scarce in the intestine and hardly detectable in the fecal samples of healthy individuals[9,10].

Several factors, including antibiotic use, dietary shifts, aging, and intestinal inflammation, can increase the relative abundance of oral bacteria in fecal samples[6]. This relative enrichment of oral bacteria has been associated with a number of diseases, including Crohn's disease[11], ulcerative colitis[12], irritable bowel syndrome[13], colorectal cancer[14], and liver cirrhosis[15]. However, the mechanisms underlying these associations remain unclear.

An important point to consider is that the oral bacterial enrichment in the intestine has primarily been detected through amplicon sequencing of the 16S rRNA gene using DNA extracted from fecal samples. This method produces compositional data, and we propose two competing explanations for its interpretation (Fig. 1): either the total abundance of oral bacteria expands within the intestine (the *Expansion* hypothesis), or the absolute abundance of gut resident bacteria decreases, resulting in a relatively higher representation of oral bacteria in fecal samples without a similar increase in their absolute numbers (the *Marker* hypothesis). While the *Expansion* hypothesis suggests that the gut environment has changed to favor oral residents, the *Marker* hypothesis implies that the gut resident population was damaged with reduced total bacterial load. Distinguishing between the two hypotheses is crucial for understanding the microbiome population dynamics and its implications for human health.

Here, we compared the two hypotheses using quantitative microbiome data from mice and humans. In the mouse model, we collected paired oral and fecal samples from each subject to quantify the total fraction of oral bacteria detected in the fecal samples after antibiotic treatment or chemically-induced epithelial damage. We find that antibiotic treatment, which depletes the gut bacterial population, increases the fraction of oral bacteria in feces without increasing their absolute numbers. In contrast, chemically-induced inflammation, which alters the composition of the gut microbiota without decreasing its population size, leaves unchanged the fraction of oral bacteria detected in feces. For human patients, we developed a method to estimate the oral bacterial fraction in fecal samples of individuals lacking paired oral samples. Using this method, we observed a similar increase in the relative, but not absolute, abundance of oral bacteria in feces of patients who suffered gut bacteria depletion. These results support the *Marker* hypothesis over the *Expansion* hypothesis. Finally, we show that the fraction of oral bacteria in feces not only serves as a marker for the loss of gut commensal, but is also associated with patient outcomes that involve gut bacterial depletion as a component.

81
Figure 1: The relative enrichment of oral bacteria in feces has two competing explanations.
Certain diseases and medical treatments lead to an enrichment of oral bacteria in fecal samples.
This phenomenon has two plausible explanations. In the *Expansion* hypothesis (highlighted in
red), the increased fraction of oral bacteria is caused by a similar rise in their absolute numbers.
On the other hand, the *Marker* hypothesis (highlighted in blue) attributes this phenomenon to the
depletion of gut bacteria. The *Expansion* hypothesis suggests that the gut environment has
changed to become more favorable for oral bacteria to establish, whereas the *Marker* hypothesis
proposes that perturbations have depleted the gut resident population, and the oral bacteria are
simply passing through. Distinguishing between these two hypotheses is crucial for interpreting
microbiome population dynamics through compositional analysis of fecal samples.
92
93 ## Results
94
Experiments with antibiotic-treated mice support the *Marker* hypothesis
To study the enrichment of oral bacteria in the intestine, we treated eight C57BL/6J
female mice with a cocktail of antibiotics (ampicillin, vancomycin, and neomycin) for one week
(Fig. 2a). This regimen is known to cause a significant depletion of gut-resident bacteria[16]. We
included two additional groups of (a) five untreated mice and (b) five mice treated with DSS
(dextran sulfate sodium)—a chemical used to induce epithelial damage without impacting the
total bacterial load in feces[17]—for comparison. Throughout the experiment, we collected paired
fecal and oral samples, which were subsequently analyzed by 16S amplicon sequencing to
profile bacterial composition and by 16S quantitative PCR (qPCR) to estimate total bacterial
abundance.
The composition of bacterial populations in the oral and fecal samples before antibiotic
treatment were very different, reflecting the distinct habitats of these two organs (Extended Data
Fig. 1a). Due to this niche specificity, we could define "oral resident bacteria", as opposed to
"gut resident bacteria", as those amplicon sequence variants (ASVs) highly abundant and
prevalent in pre-treatment oral samples but rarely found in their paired, pre-treatment fecal
samples (Methods). We identified 53 such oral ASVs that belong to 32 classified genera, which
further enabled us to calculate the total fraction of oral bacteria in all fecal samples (Table S1).
We subsequently compared the oral bacterial fraction that was estimated by our approach to the
oral bacterial fraction that was calculated by the established source tracking algorithm, FEAST[18],
for post-treatment samples. This comparison yielded a remarkable consistency with an R^2 value
of 0.95 (Extended Data Fig. 2) and validated our approach.

As expected, one week of antibiotic treatment resulted in a significant reduction in the
total bacterial load in feces compared to the untreated and DSS-treated mice (Extended Data Fig.
1b). In antibiotic-treated mice, the drop in the total bacterial load coincided with an increased
total fraction of oral bacteria detected in the feces (Fig. 2b). This fraction surged from an average
of 1.7e-4 in the pre-treatment samples to an average of 0.19 on day 3 and 0.21 after one week.
Remarkably, the microbiota composition of post-antibiotic treatment fecal samples was more
similar to the pre-treatment oral samples than to the pre-treatment fecal samples (Table S2,
Extended Data Fig. 1c). In contrast, there was no enrichment of oral bacteria in the untreated
mice (Fig. 2c). In DSS-treated mice, the chemically-induced gut inflammation altered gut
microbiota composition (Table S2) but did not deplete the total bacterial load. Consistent with
the *Marker* hypothesis, there was no increase in the fraction of oral bacteria detected in feces
(Fig. 2d).
We next focused on the antibiotic-treated mice to investigate whether the observed oral
bacterial enrichment aligned with the *Expansion* or *Marker* hypothesis. To determine the
absolute abundance of oral bacteria, we multiplied the total relative abundance of oral bacteria
by the total bacteria load. Notably, the average absolute load of oral bacteria detected in the feces
did not increase after antibiotic treatment (Fig. 2e, Extended Data Fig. 3). Instead, the observed
relative enrichment of oral bacteria coincided with a significant reduction in the absolute
abundance of gut bacteria, which decreased by >1,000 fold on average (Fig. 2f).
The pure *Marker* hypothesis (i.e., oral bacterial load remains constant, and relative
enrichment is solely driven by gut bacterial depletion) indicates that the derivative of $\log_{10}$(total
bacterial load) to the $\log_{10}$(oral bacterial fraction) equals -1 (Supplementary Note 1). Conversely,
a positive derivative would reject the pure *Marker* hypothesis and favor the pure *Expansion*
hypothesis (i.e., gut bacterial load remains constant, and relative enrichment is solely driven by
oral bacterial expansion). Remarkably, a linear regression between the $\log_{10}$-transformed oral
bacterial fraction and total bacterial load in mouse feces produced a slope of -0.97 (Fig. 2g).
These results, taken together, strongly support the *Marker* hypothesis as the predominant
explanation for the enrichment of oral bacteria after antibiotic treatment.
144

Figure 2. The administration of antibiotics to mice depletes gut bacteria and increases the
relative abundance of oral bacteria in fecal samples. a, Experimental design. The experiment
included three research arms: untreated controls (n=5), mice administered an antibiotic cocktail
of ampicillin, vancomycin, and neomycin (n=8), and mice treated with dextran sulfate sodium
(DSS) (n=5). Paired fecal and oral samples were collected at three-time points: the same day
before treatment initiation (pre), 3 days after treatment initiation (d3), and one week after

treatment initiation (w1). Samples with less than 1,000 reads were excluded from analysis (see
Extended Data Fig. 1a for excluded samples) and not shown in panels (b)-(g). 16S rRNA gene
amplicon sequencing and 16S quantitative PCR (qPCR) data were collected for each sample. **b-**
d, Relative abundance of oral bacteria in fecal samples from mice in the antibiotic treatment
group (b), no treatment group (c), and DSS treatment group (d). **e,f**, Absolute abundance of oral
(e) and gut (f) bacteria in fecal samples from antibiotic-treated mice. In panels (b)-(f), each circle
represents a fecal sample. Bar heights represent the means, with error bars indicating the 95%
confidence interval (CI). *P* values were calculated using a one-sided Wilcoxon signed-rank test.
g, Linear regression between oral bacterial fraction and total bacterial load in the $\log_{10}$-$\log_{10}$
scale. Pre-treatment (pre) and post-treatment (d3, w1) samples are represented by circles and
crosses, respectively. The black line represents the best linear fit and the shading of the same
color indicates the 95% CI. Samples with zero oral bacterial fraction were not shown and
excluded from the linear regression analysis. The *y*-axis unit in panels (e)-(g) is 16S copies per
gram of feces.

Method for quantifying oral bacteria in fecal samples from human subjects without paired
oral samples
We next aimed to determine whether the *Marker* hypothesis remains the major
mechanism for the reported enrichment of oral bacteria in the human intestine. Similar to the
mouse microbiome, bacterial populations residing in the human body have distinct compositions
depending on the body site[1]. We, therefore, adopted a similar strategy as used in our mouse data
analysis to identify and quantify oral bacteria in human feces (Methods). We analyzed 2,932
paired oral (collected from several oral cavity sites) and fecal samples from 223 healthy
individuals in the Human Microbiome Project (HMP)[1]. We found 178 oral ASVs that belong to
42 classified genera, including 24 *Prevotella* and 15 *Streptococcus* ASVs (Extended Data Fig. 4,
Table S3). The set of 178 reference oral ASVs obtained from the HMP cohort enabled us to
calculate the total fraction of oral bacteria in new fecal samples, even without paired oral
samples. We validated our approach by analyzing a publicly available dataset consisting of
paired fecal and saliva samples from patients with inflammatory bowel disease and their healthy
controls[19] (Extended Data Fig. 5, Supplementary Note 2).

Longitudinal analysis of patients hospitalized for allo-HCT supports the *Marker* hypothesis
Having established a method for detecting oral ASVs in human samples, we leveraged a
previously compiled human microbiome dataset[20] to test the two hypotheses. This dataset
comprises 10,433 longitudinal fecal samples collected from 1,276 patients who underwent
allogeneic hematopoietic cell transplantation (allo-HCT) over the past decade at Memorial Sloan
Kettering Cancer Center (MSKCC). Within this dataset, a nested subset of 3,108 samples
included 16S qPCR data. Since paired oral samples were unavailable, we used the above
reference set to identify oral ASVs and quantify their total fraction in fecal samples (Table S4).
Interestingly, 901 of the 10,433 fecal samples were dominated by a single oral ASV with a
relative abundance that exceeded 30% (Fig. 3a). These oral ASVs with > 30% abundance
primarily belong to three genera, *Streptococcus*, *Actinomyces*, and *Abiotrophia*, which dominated
778, 73, and 30 samples respectively. Notably, factors such as biofilm-forming capacity
(Supplementary Note 3) and sequencing depth (Fig. S1) did not account for the observed
variations in oral bacterial fraction.
As shown in Fig. 3b (top), oral bacteria became significantly enriched in the fecal
samples before allo-HCT, which temporally coincided with the onset of antibiotic prophylaxis
(Methods; n = 87, one-sided Wilcoxon signed-rank test, the same below; *P* = 1.2e-11, Cohen's *d*

= 1.09). Among the major antibiotics, piperacillin-tazobactam showed the most significant association with the enrichment (Table S5, Supplementary Note 4), which has been confirmed in a separate cohort of pediatric allo-HCT recipients (Fig. S2). Notably, the rise in the relative abundance of oral bacteria did not correlate with changes in their absolute abundance (Fig. 3b, middle and bottom, green curves), which remained stable without significant increase throughout the transplantation period ($P = 0.364$, Cohen's $d = 0.15$). As a result, the relative enrichment was attributed to the significant reduction in the absolute abundance of gut bacteria ($P = 9.9e-8$, Cohen's $d = 0.65$), as shown by the orange curve in the bottom panel of Fig. 3b.

Analogous to the analysis of our mouse experiment, we performed a linear regression analysis between the $\log_{10}$(oral bacterial fraction) and $\log_{10}$(total bacterial load). This analysis yielded a slope of -0.41 (Extended Data Fig. 6), which is higher than the theoretically expected value of -1 under the pure *Marker* hypothesis. However, by simulating the impact of interindividual variability on the regression slope (Methods), the theoretical value of the slope increased to -0.36 (Extended Data Fig. 7a), suggesting that the observed discrepancy may arise from the variability between patients. In contrast to the negative linear regression slope observed for oral bacteria, positive slopes were observed for several dominant gut bacterial genera, indicating that their relative abundance reflected their absolute abundance (Table S6).

We also analyzed metagenomic data available for a nested subset of 395 fecal samples[21] (Methods). A peak-to-trough ratio analysis revealed a slow growth rate for the most abundant oral ASV (*Streptococcus* ASV_8), suggesting that the *Streptococcus* was not actively proliferating in the intestine (Table S7, Supplementary Note 5). Collectively, our human microbiome data analysis extends our findings beyond mice and validates the *Marker* hypothesis as the primary explanation for the enrichment of oral bacteria detected in patients' feces.

Figure 3: Relative enrichment of oral bacteria in fecal samples of allo-HCT patients coincides with onset of antibiotic prophylaxis and depletion of gut bacteria. a, Bacterial population composition in all 10,433 fecal samples from our MSKCC dataset. Each thin vertical bar represents a single sample, with taxonomic composition (top), estimated oral bacterial fraction (middle), and total bacterial load (bottom) aligned for all samples. **b**, Profiles of antibiotic exposure (top) aligned with the population dynamics of relative (middle) and absolute (bottom) abundance of oral (green curves) and gut (orange curves) bacteria in feces. Lines and dots represent the means, and shadings of the same color indicate the 95% confidence interval.

233   The black dashed line marks the median day -6 when antibiotic prophylaxis was initiated. IV
234   vanco: intravenous vancomycin; FQs: fluoroquinolones; Pip-Tazo: piperacillin-tazobactam.
235
236   **The *Marker* hypothesis is generalizable to IBD patients with depleted gut microbiota**
237   The relative enrichment of oral bacteria in fecal samples has been recognized as a
characteristic feature of IBD[22]. Intriguingly, another hallmark of IBD is a depleted microbial
load in fecal samples[23,24]. If the *Marker* hypothesis holds for IBD patients, these two microbiome
signatures should exhibit an inverse relationship. To test this hypothesis, we analyzed a
previously published quantitative microbiome dataset obtained from 17 patients with Crohn's
disease and from 80 healthy controls[23]. These data included 16S amplicon sequencing and flow
cytometric enumeration of microbial cells. As expected, these patients showed a higher oral
bacterial fraction and lower total microbial load in their fecal samples compared to healthy
controls (Fig. 4, marginal distributions). As a result, these two signatures were negatively
correlated with a regression slope of -0.30, a value that matches the pure *Marker* hypothesis after
considering interindividual variability (Extended Data Fig. 7b). We, therefore, conclude that the
Marker hypothesis joins the two previously discovered IBD biomarkers and is generalizable to
patients with Crohn's disease.
250

251
Figure 4: Inverse correlation between total fraction of oral bacteria and microbial load in
fecal samples of patients with Crohn's disease. Each circle represents a fecal sample. Samples
with zero oral bacterial fraction in the original dataset were excluded from the plot and statistical
analyses. The black line represents the best linear fit, and the shading of the same color indicates
the 95% confidence interval. The plot also displays the regression slope and its standard error. *P*
values for the difference in the marginal distributions between patient and control groups were
calculated using a one-sided Mann-Whitney U test.
259
260   **Clinical implications of oral bacterial fraction as an indicator of total bacterial load**
261   Loss of gut bacteria can alter host physiology and has been linked to host health[23-26].
Having established the *Marker* hypothesis, we sought to determine whether the oral bacterial
fraction can replace the total bacterial load in assessing the impacts of gut bacterial depletion on
patient outcomes. This substitution, if validated, has important clinical applications, since the

associations between total bacterial load and patient outcomes could be effectively studied in the
absence of quantitative microbiome data. To test the hypothesis, we used the MSKCC allo-HCT
cohort which contains comprehensive patient outcome metadata. In this dataset, we showed that
the oral bacterial fraction in feces can predict the depletion of total bacteria, with a mean cross-
validation accuracy of 69% (Extended Data Fig. 8).
We considered four patient outcomes that can be influenced by gut bacterial depletion,
including stool consistency, fungal overgrowth, bloodstream infection, and overall survival. If
oral bacterial fraction could replace the total bacterial load, it should be associated with these
patient outcomes. Stool consistency is strongly correlated with gut microbiome[27]. The loss of gut
commensals reduces the total bacterial biomass and the production of short-chain fatty acids,
which can increase water absorption in the colon[28]. As expected, we found a significant
association between a high oral bacterial fraction and a low stool consistency (Extended Data
Fig. 9a). Also, gut bacterial depletion can create an ecological niche for fungi to colonize and
expand[29]. In line with our expectation, the oral bacterial fraction is significantly higher in fecal
samples with a positive fungal culture than in those with a negative fungal culture (Extended
Data Fig. 9b). Moreover, depletion of gut bacteria eliminates potentially invasive pathogens that
may translocate to bloodstream and cause infections[30,31]. Indeed, our analysis revealed that a
high oral bacterial fraction reduces the risk of total bacterial bloodstream infections (Methods;
Hazard ratio = 0.29; 95% confidence interval, 0.11-0.75; P = 0.011). This result with a high oral
bacterial fraction stands in contrast to intestinal domination states by *Enterococcus*, *Klebsiella*,
or *Escherichia*, which strongly increase the risk of subsequent bloodstream infections caused by
these organisms[30,31]. The striking contrast, in turn, supports our previous finding that the relative
enrichment of oral bacteria in feces of allo-HCT recipients was not driven by their expansion in
the gut (Fig. 3b). Finally, the oral bacterial fraction was negatively associated with all-cause and
graft-versus-host disease (GVHD)-related mortality (Table S8, Supplementary Note 6), which is
consistent with the notion that gut microbiome damage is associated with worse patient
outcomes[32].
292
Discussion
The relative enrichment of oral bacteria detected in fecal samples has been associated
with diseases[6] and changes in host gene expression[13] in various gastrointestinal disorders. To
understand these associations, it is essential to distinguish changes in relative abundance from
changes in absolute abundance of bacteria in feces[23,33]. In this study, we provide compelling
evidence for the *Marker* hypothesis: a relative enrichment of oral bacteria signifies a depletion of
commensals in the intestine rather than an expansion of the oral bacterial population with
increasing absolute numbers. This conclusion is based on population averages. On an individual
level, new oral ASVs may emerge in the gut following perturbations, leading to their increased
absolute abundances (Extended Data Fig. 3). While we demonstrate that the total numbers of oral
bacteria did not increase when gut bacteria have been depleted, it remains unclear what niche
factors prevent their growth in the gut and whether these oral bacteria are even alive. Further
research is needed to explore the growth and survival of oral bacteria in the gut niche[34].
In our mouse experiments and those carried out in a previous study[17], the administration
of DSS-induced gut inflammation, leading to a drastic alteration of the gut environment and
subsequent changes in gut microbiota composition. However, the chemically-induced gut
inflamation did not reduce the total bacterial load in mouse feces. According to the *Marker*
hypothesis, in such cases, an increase in the oral bacterial fraction should not occur due to the
lack of gut bacterial depletion, which aligns precisely with our observations. Interestingly, the

Crohn's disease patients analyzed in Fig. 4 had lower levels of total bacterial load compared to the healthy controls, which correlated with a higher fraction of oral bacteria detected in their feces. This finding suggests that the mouse model of DSS-induced gut inflammation did not reproduce the gut bacterial depletion observed in these patients. Taken together, both the mouse and human data support the *Marker* hypothesis as the primary explanation for the observed gut microbiota changes in the context of gut inflammation.

Previous studies have shown that ectopic colonization of oral bacteria in mouse gut can induce strong immune responses and gut inflammation[3,5]. These findings in mice suggest a potential causal link between the presence of translocated oral bacteria in the gut and the development of diseases. By demonstrating the *Marker* hypothesis, our study suggests that these microbiome-disease associations could be mediated—at least in part—by the loss of gut commensals. In mice, the total microbial load in the intestine has exhibited positive correlations with fecal IgA concentration[24], the proportions of mucosal RORγt[+] cells[25], and colonic lamina propria FoxP3[+] regulatory T cells[24]. Still, the *Marker* hypothesis does not exclude the possibility that oral bacteria can drive disease. Even without an absolute increase in population size, oral commensals may acquire pathogenic potential in the human gut. Further research is needed to investigate how the pathogenicity of oral bacteria and the loss of gut commensals synergistically impact host responses and exacerbate pathology.

In summary, we have validated the *Marker* hypothesis by showing a robust negative correlation between oral bacterial fraction and total bacterial load under conditions where the gut microbiota is disrupted in mice, in patients undergoing allo-HCT, and in patients with inflammatory bowel disease. However, a limitation of our approach is that it depends on a marked depletion of gut bacteria caused by perturbations such as antibiotic treatments. For example, in our mouse experiment, antibiotics depleted the gut bacterial load by over 1,000-fold. When individuals exhibit minor or no variations in the size of their gut bacterial population[35], the link between the fraction of oral bacteria detected in fecal samples and the total size of the gut bacteria population may be masked by interindividual variability. In such scenarios, the applicability of the *Marker* hypothesis and its derived results may also be limited.

Nonetheless, when subjects experience massive fluctuations in the total bacterial load, its negative correlation with oral bacterial fraction could provide a convenient method for estimating the changes in the absolute abundance of bacterial taxa from relative microbiome profiles alone. This can be achieved by calculating the ratio of their relative abundance to oral bacterial fraction in fecal samples (see Extended Data Fig. 10 for an example). This method could be applied to previously published microbiome compositional data to reevaluate previously identified bacterial taxonomic biomarkers and potentially discover new biomarkers based on estimated absolute abundances. Similarly, the oral bacterial fraction could serve as a biomarker for the outcome of fecal microbiota transplantation (FMT). A high fraction of oral bacteria in the feces of individuals being considered for FMT may signal a depleted gut microbiota in these candidates, thereby suggesting a higher chance of successful FMT engraftment. Supporting this notion, a recent study found that *Streptococcus salivarius*, a typical oral bacterial species, in the recipient's microbiota facilitates donor strain colonization[36].

Acknowledgements

Chen.L. and A.D. were supported by National Institutes of Health (NIH) grant nos. U01
AI124275 (J.B.X.), R01 AI137269 (J.B.X.) and U54 CA209975 (J.B.X.). T.R. was funded by
Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) grant no. RO-5328/1-2
(T.R.), NIH grant nos. R01 AI093808 (T.M.H.), and R21 AI156157 (T.M.H.).

Author contributions

Conceptualization, Chen.L., T.R., T.M.H., and J.B.X.; Mouse experiment, A.D., V.M., and
Chloe.L.; Microbiome data processing, Chen.L., A.D., and H.L.; Microbiome data analysis,
Chen.L., T.R. and T.F.; Writing – Original Draft, Chen.L., T.R., A.D., H.L.,T.F.; Writing –
Review and Editing, T.M.H., J.B.X., J.U.P., B.Z., L.D., and M.R.M.v.d.B.; Supervision, J.B.X.
and T.M.H.

Competing interests

J.U.P. reports research funding, intellectual property fees and travel reimbursement from Seres
Therapeutics and consulting fees from DaVolterra, CSL Behring and from Maat Pharma. He
serves on an Advisory board of and holds equity in Postbiotics Plus Research. He has filed
intellectual property applications related to the microbiome (reference nos. 62/843,849,
62/977,908 and 15/756,845). M.R.M.v.d.B. has received research support from Seres
Therapeutics; he has consulted, received honorarium from or participated in advisory boards for
Seres Therapeutics, WindMIL Therapeutics, Rheos, Frazier Healthcare Partners, Nektar
Therapeutics, Notch Therapeutics, Forty Seven Inc., Priothera, Ceramedix, Lygenesis, Pluto
Immunotherapeutics, Magenta Therapeutics, Merck & Co., Inc. and DKMS Medical Council
(Board); and he has IP Licensing with Seres Therapeutics, Juno Therapeutics and stock options
from Seres and Notch Therapeutics. T.M.H. has participated in a scientific advisory board for
Boehringer-Ingelheim Inc. T.R. is currently an employee of BioNTech SE. Memorial Sloan
Kettering Cancer Center (MSKCC) has financial interests relative to Seres Therapeutics.

Data availability

A comprehensive list of the microbiome datasets used in this study, along with an indication of
whether they contain quantitative data, is provided in Table S9. The raw 16S sequences of mouse
fecal and oral samples collected in this study have been uploaded to Sequence Read Archive
(SRA) with BioProject accession number PRJNA873058. All other microbiome datasets are
publicly available. All processed data supporting the findings of this study are available within
the article and its supplementary materials.

Code availability

Customized Python codes for reproducing the figures and tables are available on Github
(https://github.com/liaochen1988/Oral_bacterial_manuscript_revision)

Methods

Mouse experiment setup. Mice used in this study were C57BL/6J specific pathogen-free mice purchased from The Jackson Laboratories. Animals were 6-8 weeks old females. During the study, mice were single-housed in autoclaved cages with *ad libitum* access to autoclaved and acidified reverse osmosis water (pH, 2.5 to 2.8) and irradiated feed (LabDiet 5053, PMI, St Louis, MO). The animal holding room is maintained at 72 ± 2 ºF (21.5 ± 1 ºC), relative humidity between 30% and 70%, and a 12:12 hour light:dark photoperiod. Animal use is approved by MSKCC's IACUC. The institution's animal care and use program is AAALAC-accredited and operates in accordance with the recommendations provided in the *Guide*.

We performed two independent experiments. In the first one we tested the role of broad-spectrum antibiotics on the relative abundance of oral bacteria in the intestine and in the second one we repeated antibiotic cocktail treatment and included one more group that was exposed to dextran sulfate sodium (DSS) salt. Briefly, mice were treated with cocktail of ampicillin (0.5g/l), vancomycin (0.5g/l) and neomycin (1g/l) for one week in drinking water. To test the role of DSS, which is supposed to minimally affect bacterial load in the gut, we treated the mice with 3% DSS in the drinking water for a week. The water was changed once during the course of both treatments to preserve the activity of antibiotics and DSS. The third group of animals that served as a control drank regular water. Fecal pellets were collected at least immediately before and one week after the initiation of different treatments, and in some cases 3 days after the initiation of the treatment. Oral swabs were collected as per reference of Abusleme *et al.*[37] at the time fecal sampling was done. Briefly, mice were hand-held while sterile swab was introduced into mouth and swiped for at least 30 seconds. After, the swab was put into 150 µl of TE buffer, and the tip was cut off so that the Eppendorf can be closed. Samples were put immediately to dry ice. One negative control swab was taken by pulling out the swab from the pouch and swirling through air for at least 30 seconds, after which it was put in TE and on dry ice. Fecal samples and oral swabs were kept at -80ºC until further processing.

DNA extraction and sequencing. DNA extraction and library preparation of the fecal samples collected during the first mouse experiment was done in the sequencing core facility at MSKCC. However, since the DSS treatment results in difficult amplification of the DNA, all fecal samples from the second experiment (control, antibiotic-treatment, and DSS-treatment samples) were subjected to DNA extraction and further processing in the Xavier lab with protocols adapted to ensure the proper DNA amplification and library preparation.

1) First experiment fecal samples processing. Fecal DNA was extracted and 16S rRNA gene was amplified using the previously described protocol[38]. QIAseq 1-step Amplicon Library Kit was used for generating libraries that were later quantified, normalized, and sequenced using MiSeq Reagent Kit V3.
2) Second experiment fecal samples processing. Bacterial DNA from fecal samples was extracted using the QIAamp® DNA Fast Stool Mini kit with introduction of a mechanic disruption step with bead-beating as per reference of Djukovic *et al.*[39]. The V4-V5 region of the 16S rRNA gene was amplified with Q5 High-Fidelity DNA Polymerase following the manufacturer's recommendations. Briefly, for each sample, 25 µl of the PCR was prepared containing 5 µl of the Q5 buffer, 5 µl of the Q5 enhancer, 0.5 µl of the 10 mM dNTPs and 0.625 µl of the Q5 polymerase. 2.5 µl of the forward and 2.5 µl of the reverse primer at concentrations of 10 µM were added. Forward and reverse primers used for each sample contained unique sequences that would allow demultiplexing during

439 sequence processing step. 1 ng of the DNA extracted from fecal samples collected from
440 control and antibiotic-treated groups was added to reaction, while the DNA obtained
441 from DSS-treated animals was diluted 1:20 and 5 ul was amplified. This was done since
442 the carryover DSS prevented amplification of the DNA from some fecal samples, while
443 in diluted samples the amplification worked as expected. Where necessary, the volume of
444 the reaction was completed with water. Cycling conditions of the PCR were 96 ºC for 10
445 minutes, and 35 cycles of 96 ºC for 10 seconds, 51 ºC for 30 seconds and 72 ºC for 30
446 seconds. The final elongation step was performed at 72 ºC for 5 minutes. Amplification
447 was confirmed with agarose gel electrophoresis and PCR products were purified with
448 AxyPrep Magnetic Beads, quantified with Quant-iT PicoGreen dsDNA kit, normalized
449 and pooled. KAPA LTP Library Preparation Kit was used to generate sequencing
450 libraries that were later quantified with Quant-iT PicoGreen dsDNA kit, normalized, and
451 sequenced using MiSeq Reagent Kit V3.

 3) Oral samples processing. Oral DNA was extracted by using modified DNeasy Blood and
453 Tissue Kit protocol as described in Abusleme et al.[37]. After extraction V4-V5 region of
454 the 16S rRNA gene was amplified with Q5 High-Fidelity DNA Polymerase by cycling 1
455 ng of the extracted DNA. Volumes and concentrations of PCR reagents were the same as
456 described above. The following steps that include agarose gel electrophoresis, PCR
457 product purification, normalization, pooling, library preparation, library quantification,
458 and sequencing were performed as described in the previous section.

460 **Quantitative PCR (qPCR) for determining bacterial load.** For assessing the bacterial load in
461 fecal and oral samples, qPCR against standard curve was used to determine 16S rRNA copy
462 number. For this purpose, the PowerUP qPCR Kit was used. Briefly, for each sample, 20 μl PCR
463 triplicates were prepared with each containing 2 μl of the DNA used as template, 10 μl of mix
464 provided by the manufacturer, and 1 μl of forward and reverse primers at the final concentration
465 of 0.5 μM. We used the primer pair 27F/338R to amplify the V1-V2 region of the 16S rRNA
466 gene (F- AGAGTTTGATCMTGGCTCAG; R- TGCTGCCTCCCGTAGGAGT). In order to
467 complete the volume of the reaction, 6 μl of water was added. A PCR product of the 16S rRNA
468 gene from *Enterococcus faecium* ATCC 700221 strain was used for obtaining a standard curve
469 by amplifying its 16S rRNA gene and purifying the product. The copy number of the PCR
470 product was determined based on its concentration and 16S rRNA sequence. A standard curve
471 was obtained by using 10-fold dilutions.

472 Cycling conditions of the qPCR were 50 ºC for 2 minutes, 95 ºC for 2 minutes, and 40
473 cycles of 95 ºC for 15 seconds, 56 ºC for 15 seconds and 72 ºC for 60 seconds. By extrapolating
474 results by looking the ones obtained from standard curve samples, the number of 16S rRNA
475 genes was determined for each sample. The final number of 16S rRNA genes per 1 g of fecal
476 sample was calculated by multiplying the number of 16S rRNA molecules obtained by qPCR
477 with DNA elution volume after DNA extraction and dividing this number with the weight of the
478 fecal pellet from which DNA extraction was performed.

480 **16S amplicon sequencing data processing.** Samples collected from the first mouse experiment
481 were sequenced once. Fecal and oral samples from the second mouse experiment were
482 sequenced twice using the MiSeq sequencers at the Joao Xavier and Thomas Norman labs at
483 MSKCC. Since the two sequencing runs yielded very similar read coverage and bacterial
484 profiles, we combined the read counts from both runs for samples in the second mouse
485 experiment. Sequence processing was done following DADA2 (Divisive Amplicon Denoising
486 Algorithm) tutorial with an in-house script. Briefly, after demultiplexing, reads were trimmed to

the first 180 bp or the first point with a quality score Q < 2, and removed if they contained ambiguous nucleotides (N) or if two or more errors were expected based on the quality of the trimmed reads. Paired reads were merged, and chimeras were removed. ASVs were identified using DADA2 and classified against the SILVA v138 database[40].

The bacterial profiles of > 10,000 fecal samples from MSKCC allo-HCT recipients were previously analyzed by an in-house pipeline and compiled in a recent study[20]. Similarly, reads were trimmed to the first 180 bp or the first point with a quality score Q < 2, and removed if they contained ambiguous nucleotides (N) or if two or more errors were expected based on the quality of the trimmed reads. ASVs were identified using DADA2[41] and classified against the SILVA v138 database[40].

The demultiplexed and primer-trimmed HMP[1] 16S sequences (V3-V5 region) were obtained from the Qitta repository[42] and processed using QIIME (Quantitative Insights Into Microbial Ecology) 2[43]. DADA2 was applied to denoise the data and generate an ASV per sample count table, using the QIIME denoise-pyro plugin[43]. To remove low-quality tails, we used parameter --p-trunc-len 395 to generate ASVs that cover the entire high-quality V4-V5 region. Taxonomy classification of the ASV sequences was performed using the QIIME plugin "feature-classifier"[44] and the SILVA v138 database[40]. The classification took three steps. We first extracted the V3-V5 region of the SILVA reference sequences using the extract-reads method. Then we created a classifier by using the fit-classifier-naïve-bayes method with extracted reads and the SILVA reference taxonomy. Finally, we ran the classifier on the ASV sequences using the classify-sklearn method to get their taxonomy.

All other microbiome datasets in this study were processed using QIIME 2[43]. Demultiplexed short reads were trimmed using the QIIME cutadapt plugin[45] with parameters "--p-error-rate 0.1" and "--p-overlap 3". The trimmed reads were then denoised using the QIIME dada2 plugin with truncation lengths determined by per-base quality scores to generate ASV-level feature tables. Taxonomic classification was performed using the QIIME plugin "feature-classifier classify-sklearn"[46] against the SILVA v138 database[40].

For all microbiome datasets analyzed in this study, samples with sequencing coverage below 1,000 reads were excluded. Additionally, we excluded non-bacterial ASVs and ASVs with "Chloroplast" or "Mitochondria" in their taxonomy labels.

Identification of oral ASVs in fecal samples. We leveraged paired oral and fecal samples to identify bacterial ASVs typically colonizing the oral cavity. An ASV is considered of oral origin if it satisfies all four of the following criteria simultaneously: (1) its relative abundance, averaged across all oral cavity samples, is above θ_a; (2) its relative abundance, averaged across all fecal samples, does not exceed θ_a; (3) its prevalence, across all oral cavity samples, is above θ_p; (4) its prevalence, across all fecal samples, does not exceed θ_p. We identified a total of 53 such oral ASVs in all mice using θ_a = 1e-3 and θ_p = 10%. Mouse-specific oral ASVs were selected by excluding any ASV from this set if it was undetectable in the pre-treatment oral samples. Exceptions were made for three mice (Control_2D, DSS_2A, DSS_2C) lacking pre-treatment oral samples, for which no ASVs were excluded. To identify oral ASVs in humans, we employed the HMP[1] dataset and applied θ_a = 1e-4 and θ_p = 5%. The cutoff values are consistent with those used in a previous study[47] for identifying oral species (not ASVs) from metagenomic data. For both mice and humans, the prevalence of an ASV was calculated as the proportion of samples containing the ASV at a relative abundance above 1e-3[48]. Unless otherwise specified, we calculated the absolute abundance of oral and gut bacteria by multiplying their relative

533   abundances by the total bacterial loads, which can be quantified by either 16S qPCR or flow
534   cytometry.
535

Impact of interindividual variability on regression slope between oral bacterial fraction
and total bacterial load. In the absence of interindividual variability, the derivative of the log-
transformed total bacterial load with respect to the log-transformed oral bacterial fraction equals
to -1 under the pure *Marker* hypothesis and remains positive under the pure *Expansion*
hypothesis (see Supplementary Note 1). To understand the impact of interindividual variability
on this derivative, we simulated the oral bacterial fraction (f_{oral}) and total bacterial load (F_{total})
in fecal samples from both a control and a case group. The control group samples are collected
from individuals with low oral bacterial fraction. These individuals, for example, include healthy
individuals or patients before a medicial treatment. The case group samples are collected after a
perturbation that induces relative enrichment of oral bacteria, such as antibiotic treatment and
certain intestinal disorders. It is important to note that the control and case group samples do not
necessarily come from the same longitudinal study and may be collected from different
populations in a cross-sectional study.
For each control group sample i, we assume that the $\log_{10}$-transformed values of the oral
bacterial fraction ($f_{oral,i,ctr}$) and total bacterial load ($F_{total,i,ctr}$) follow separate Gaussian
distributions:

$$p\left(\log_{10} f_{oral,i,ctr} = x\right) = \frac{1}{\sqrt{2\pi}\sigma_f} \exp\left(-\frac{1}{2\sigma_f^2}(x - \mu_f)^2\right)$$ Eq. 1

$$p\left(\log_{10} F_{total,i,ctr} = x\right) = \frac{1}{\sqrt{2\pi}\sigma_F} \exp\left(-\frac{1}{2\sigma_F^2}(x - \mu_F)^2\right)$$ Eq. 2

Here, μ_f and μ_F represent the means of the respective distributions, and σ_f and σ_F represent the
standard deviations.
For each case group sample j, we first generate its baseline values of oral bacterial
fraction ($f_{oral,j,bl}$) and total bacterial load ($F_{total,j,bl}$) from the same Gaussian distributions.
These baseline values characterize the microbiome state prior to any perturation that enriches
oral bacteria in this sample. The $\log_{10}$-transformed value of oral bacterial fraction after the
perturbation ($f_{oral,j,case}$) is drawn from a uniform distribution between $\log_{10} f_{oral,j,bl}$ and
$\log_{10} f_{oral,max}$. Here, $f_{oral,max}$ represents the maximum possible oral bacterial fraction, which
may vary based on the strength and type of the perturbation. After sampling $f_{oral,j,case}$, the total
bacterial load ($F_{total,j,case}$) in this sample can be computed in two different ways. Under the pure
Marker hypothesis (i.e., oral bacterial load remains constant), we have $F_{total,j,case} =$
$f_{oral,j,bl}F_{total,j,bl}/f_{oral,j,case}$. Under the pure *Expansion* hypothesis (i.e., gut bacterial load
remains constant), we have $F_{total,j,case} = F_{total,j,bl}(1 - f_{oral,j,bl})/(1 - f_{oral,j,case})$.
The simulation parameters (μ_f, μ_F, σ_f, σ_F) specific to the MSKCC allo-HCT cohort
presented in Fig. 3 and the Crohn's disease cohort presented in Fig. 4 were determined as
follows. For the allo-HCT recipients, their fecal samples collected between day -20 and 40
relative to the day of transplantation were divided into two sets: 185 samples obtained prior to
antibiotic prophylaxis constitute the control group, and 2,339 samples after the initiation of
prophylaxis constitute the case group. Using the control group samples, we estimated the values
of the following parameters: μ_f = -1.72, μ_F = 7.79, σ_f = 0.70, and σ_F = 0.99. On the other hand,
the Crohn's disease cohort consists of 80 healthy individuals in the control group and 17 patients

with Crohn's disease in the case group. We calculated the following parameter values from the control group samples: μ_f = -2.71, μ_F = 11.01, σ_f = 0.48, and σ_F = 0.33.

Cox proportional hazard model. We used a time-varying Cox proportional hazard model to evaluate the association between antibiotic exposure and intestinal domination by oral ASVs in allo-HCT recipients. A total of 291 patients with at least 10 samples between day -10 and 40 relative to transplantation were included. The event of interest was intestinal domination of any oral bacterial ASV that exceeds 30%. Binary indicators of antibiotic exposure were used as covariates (see Table S5 for included antibiotics). Patients without any oral bacterial domination at day 40 were censored. The model outputs the hazard ratio, which compares the likelihood of oral ASV domination occurring among patients with antibiotic exposure and patients without.

Using data from the same 291 patients, we performed the Cox proportional hazard regression analysis to assess the association between oral bacterial domination and total bloodstream infections (BSIs). Oral bacterial domination, as defined above, was analyzed as a time-varying predictor. The events of interest were BSIs caused by any bacterial species from *Citrobacter, Enterobacter, Enterococcus*, *Klebsiella*, *Escherichia*, *Pseudomonas*, *Stenotrophomonas*, and *Streptococcus*. Patients without any infection of interest at day 40 were censored. The hazard ratio indicates the relative risk of developing bacterial BSIs among patients with intestinal dominations of oral bacteria compared to those without dominations.

For both associations, we fit the Cox models using the CoxTimeVaryingFitter method from the Python package lifelines.

Survival analysis of allo-HCT recipients. Following our previous approach[32,49], we employed multivariable survival models with time-varying covariates to examine the associations between oral bacterial fraction in feces and all-cause or GVHD-related mortality among allo-HCT recipients. A total of 1,268 patients with fecal samples collected on or after day 0 relative to transplantation were included. The endpoint of interest was all-cause mortality or GVHD-related mortality until 2 years after allo-HCT, where patients who survived beyond 2 years were censored. Here, GVHD-related mortality was defined as death due to GVHD or after GVHD onset, without cancer relapse[50] or other-cause mortality. Therefore, the GVHD-related mortality is the endpoint of interest with relapse and other-cause mortality as competition risks. Due to the different endpoint characteristics, we applied the Cox model and Fine-Gray model for all-cause mortality and GVHD-related mortality, respectively. Both models included oral bacterial fraction as the predictor, along with other covariates including log-transformed *Enterococcus* absolute abundance (a risk factor for both all-cause and GVHD-related mortality[51]), age, underlying disease, graft source, and conditioning intensity. Since qPCR data was available for fewer than a third of the samples, we used the *Enterococcus*-to-oral bacteria relative abundance ratio to estimate the absolute abundance of *Enterococcus* across all 10,433 samples. We have validated the estimation in Extended Data Fig. 10. The theoretical underpinning for this estimation is provided by the inverse correlation between oral bacterial fraction and total bacterial load: dividing by the oral bacterial fraction is equivalent to multiplying by the total bacterial load. The two survival analyses were performed using functions coxph and finegray from the R package survival, respectively.

Shotgun metagenomic sequencing data processing. We adapted a recently published bioinformatic pipeline[52] to assemble bacterial genomes from metagenomic data. The pipeline uses MEGAHIT[53] to assemble contigs from short reads. It subsequently employs Metabat2[54] and

CONCOCT[55] to bin these contigs into Metagenome-Assembled Genomes (MAGs). In the last
step, it uses DAS Tool[56] to generate an optimized, non-redundant set of MAGs. High-quality
Streptococcus MAGs ($\geq$ 75% complete, $\leq$ 175 fragments/Mbp sequence, and $\leq$ 2%
contamination) classified by Kraken2[57] were further analyzed by iRep[58]. The iRep value of a
MAG represents the average number of replication events over different subpopulations of the
MAGs weighted by their relative abundances.
628
Statistical testing. Detailed descriptions of statistical analyses, including sample size, hypothesis
test name, *P* value, and effect size, are provided. All *P* values were corrected for multiple testing
using false discovery rate (FDR) adjustment (the Benjamini-Hochberg procedure). Statistical
significance was defined as (adjusted) *P* value < 0.05. The pairwise Adonis test was performed
using the R package pairwiseAdonis. The Mann-Whitney U test, Wilcoxon signed-rank test, and
linear regression analysis were performed using the corresponding functions in the Python Scipy
package. Specifically, we used a one-sided Wilcoxon signed-rank test to assess the differences in
the oral bacterial fraction, oral bacterial load, and gut bacterial load before and after the
administration of antibiotic prophylaxis among MSKCC allo-HCT recipients. We defined the
pre-treatment period as spanning from 20 days before transplantation until the onset of antibiotic
prophylaxis. The post-treatment period extended from the initiation of antibiotic prophylaxis to
the day of neutrophil engraftment. We computed their average values across all samples
collected during each of the pre-treatment and post-treatment periods.
642
643   **Plots.** For all main and supplementary figures displaying a boxplot, the central line represents the
644   median. The box limits correspond to the first and third quartiles (25th and 75th percentiles). The
645   whiskers extend to the smallest and largest values or at most to 1.5 times the interquartile range,
646   whichever is smaller. All plots were generated using Python Matplotlib library and the Seaborn
647   package.

References

1. The Human Microbiome Project Consortium. Structure, function and diversity of the healthy human microbiome. *Nature* **486**, 207-214 (2012).
2. Segata, N*., et al.* Composition of the adult digestive tract bacterial microbiome based on seven mouth surfaces, tonsils, throat and stool samples. *Genome Biol.* **13**, 1-18 (2012).
3. Atarashi, K*., et al.* Ectopic colonization of oral bacteria in the intestine drives T$_H$1 cell induction and inflammation. *Science* **358**, 359-365 (2017).
4. Li, B*., et al.* Oral bacteria colonize and compete with gut microbiota in gnotobiotic mice. *Int. J. Oral Sci.* **11**, 1-9 (2019).
5. Kitamoto, S*., et al.* The intermucosal connection between the mouth and gut in commensal pathobiont-driven colitis. *Cell* **182**, 447-462. e414 (2020).
6. Kitamoto, S., Nagao-Kitamoto, H., Hein, R., Schmidt, T. & Kamada, N. The bacterial connection between the oral cavity and the gut diseases. *J. Dent. Res.* **99**, 1021-1029 (2020).
7. Jin, S., Wetzel, D. & Schirmer, M. Deciphering mechanisms and implications of bacterial translocation in human health and disease. *Curr. Opin. Microbiol.* **67**, 102147 (2022).
8. Abed, J*., et al.* Colon cancer-associated *Fusobacterium nucleatum* may originate from the oral cavity and reach colon tumors via the circulatory system. *Front. Cell. Infect. Microbiol.* **10**, 400 (2020).
9. Schmidt, T.S*., et al.* Extensive transmission of microbes along the gastrointestinal tract. *Elife* **8**, e42693 (2019).
10. Rashidi, A., Ebadi, M., Weisdorf, D.J., Costalonga, M. & Staley, C. No evidence for colonization of oral bacteria in the distal gut in healthy adults. *Proc. Natl Acad. Sci. USA* **118**, e2114152118 (2021).
11. Gevers, D*., et al.* The treatment-naive microbiome in new-onset Crohn's disease. *Cell Host Microbe* **15**, 382-392 (2014).
12. Schirmer, M*., et al.* Compositional and temporal changes in the gut microbiome of pediatric ulcerative colitis patients are linked to disease course. *Cell Host Microbe* **24**, 600-610.e4 (2018).
13. Priya, S*., et al.* Identification of shared and disease-specific host gene–microbiome associations across human diseases using multi-omic integration. *Nat. Microbiol.*, 1-16 (2022).
14. Kostic, A.D*., et al. Fusobacterium nucleatum* potentiates intestinal tumorigenesis and modulates the tumor-immune microenvironment. *Cell Host Microbe* **14**, 207-215 (2013).
15. Qin, N*., et al.* Alterations of the human gut microbiome in liver cirrhosis. *Nature* **513**, 59-64 (2014).
16. Reikvam, D.H*., et al.* Depletion of murine intestinal microbiota: effects on gut mucosa and epithelial gene expression. *PLoS One* **6**, e17996 (2011).
17. Taguer, M*., et al.* Changes in gut bacterial translation occur before symptom onset and dysbiosis in dextran sodium sulfate-induced murine colitis. *mSystems* **6**, e00507-00521 (2021).
18. Shenhav, L*., et al.* FEAST: fast expectation-maximization for microbial source tracking. *Nat. Methods* **16**, 627-632 (2019).
19. Imai, J*., et al.* A potential pathogenic association between periodontal disease and Crohn's disease. *JCI Insight* **6**, e148543 (2021).

20.   Liao, C., *et al.* Compilation of longitudinal microbiota data and hospitalome from
hematopoietic cell transplantation patients. *Sci. Data* **8**, 1-12 (2021).
21.   Yan, J., *et al.* A compilation of fecal microbiome shotgun metagenomics from
hematopoietic cell transplantation patients. *Sci. Data* **9**, 1-11 (2022).
22.   Read, E., Curtis, M.A. & Neves, J.F. The role of oral bacteria in inflammatory bowel
disease. *Nat. Rev. Gastroenterol. Hepatol.* **18**, 731-742 (2021).
23.   Vandeputte, D., *et al.* Quantitative microbiome profiling links gut community variation to
microbial load. *Nature* **551**, 507-511 (2017).
24.   Contijoch, E.J., *et al.* Gut microbiota density influences host physiology and is shaped by
host and microbial factors. *Elife* **8**, e40553 (2019).
25.   Britton, G.J., *et al.* Defined microbiota transplant restores Th17/RORγt[+] regulatory T cell
balance in mice colonized with inflammatory bowel disease microbiotas. *Proc. Natl*
Acad. Sci. USA **117**, 21536-21545 (2020).
26.   Zarrinpar, A., *et al.* Antibiotic-induced microbiome depletion alters metabolic
homeostasis by affecting gut signaling and colonic metabolism. *Nat. Commun.* **9**, 1-13
(2018).
27.   Vandeputte, D., *et al.* Stool consistency is strongly associated with gut microbiota
richness and composition, enterotypes and bacterial growth rates. *Gut* **65**, 57-62 (2016).
28.   Bedford, A. & Gong, J. Implications of butyrate and its derivatives for gut health and
animal production. *Anim. Nutr.* **4**, 151-159 (2018).
29.   Zhai, B., *et al.* High-resolution mycobiota analysis reveals dynamic intestinal
translocation preceding invasive candidiasis. *Nat. Med.* **26**, 59-64 (2020).
30.   Stoma, I., *et al.* Compositional flux within the intestinal microbiota and risk for
bloodstream infection with Gram-negative bacteria. *Clin. Infect. Dis.* **73**, ciaa068 (2020).
31.   Taur, Y., *et al.* Intestinal domination and the risk of bacteremia in patients undergoing
allogeneic hematopoietic stem cell transplantation. *Clin. Infect. Dis.* **55**, 905-914 (2012).
32.   Peled, J.U., *et al.* Microbiota as predictor of mortality in allogeneic hematopoietic-cell
transplantation. *N. Engl. J. Med.* **382**, 822-834 (2020).
33.   Maghini, D.G., *et al.* Quantifying bias introduced by sample collection in relative and
absolute microbiome measurements. *Nat. Biotechnol.*, 1-11 (2023).
34.   Rojas-Tapias, D.F., *et al.* Inflammation-associated nitrate facilitates ectopic colonization
of oral bacterium *Veillonella parvula* in the intestine. *Nat. Microbiol.*, 1-13 (2022).
35.   Sarrabayrouse, G., *et al.* Fungal and bacterial loads: noninvasive inflammatory bowel
disease biomarkers for the clinical setting. *mSystems* **6**, e01277-20 (2021).
36.   Schmidt, T.S., *et al.* Drivers and determinants of strain dynamics following fecal
microbiota transplantation. *Nat. Med.* **28**, 1902-1912 (2022).
37.   Abusleme, L., *et al.* Oral microbiome characterization in murine models. *Bio Protoc.* **7**,
pii:e2655 (2017).
38.   Taur, Y., *et al.* Reconstitution of the gut microbiota of antibiotic-treated patients by
autologous fecal microbiota transplant. *Sci. Transl. Med.* **10**, eaap9489 (2018).
39.   Djukovic, A., *et al. Lactobacillus* supports *Clostridiales* to restrict gut colonization by
multidrug-resistant *Enterobacteriaceae*. *Nat. Commun.* **13**, 5617 (2022).
40.   Quast, C., *et al.* The SILVA ribosomal RNA gene database project: improved data
processing and web-based tools. *Nucleic Acids Res.* **41**, D590-D596 (2012).
41.   Callahan, B.J., *et al.* DADA2: High-resolution sample inference from Illumina amplicon
data. *Nat. Methods* **13**, 581-583 (2016).
42.   Gonzalez, A., *et al.* Qiita: rapid, web-enabled microbiome meta-analysis. *Nat. Methods*
15, 796-798 (2018).

43. Bolyen, E*., et al.* Reproducible, interactive, scalable and extensible microbiome data
science using QIIME 2. *Nat. Biotechnol.* **37**, 852-857 (2019).
44. Bokulich, N.A*., et al.* Optimizing taxonomic classification of marker-gene amplicon
sequences with QIIME 2's q2-feature-classifier plugin. *Microbiome* **6**, 1-17 (2018).
45. Martin, M. Cutadapt removes adapter sequences from high-throughput sequencing reads.
EMBnetJ. **17**, 10-12 (2011).
46. Pedregosa, F*., et al.* Scikit-learn: machine learning in Python. *J. Mach. Learn. Res.* **12**,
2825-2830 (2011).
47. Thomas, A.M*., et al.* Metagenomic analysis of colorectal cancer datasets identifies cross-
cohort microbial diagnostic signatures and a link with choline degradation. *Nat. Med.* **25**,
667-678 (2019).
48. Machado, D*., et al.* Polarization of microbial communities between competitive and
cooperative metabolism. *Nat. Ecol. Evol.* **5**, 195-203 (2021).
49. Nguyen, C.L*., et al.* High-resolution analyses of associations between medications,
microbiome, and mortality in cancer patients. *Cell* **186**, 2705-2718. e2717 (2023).
50. Copelan, E*., et al.* A scheme for defining cause of death and its application in the T cell
depletion trial. *Biol. Blood Marrow Transplant.* **13**, 1469-1476 (2007).
51. Stein-Thoeringer, C*., et al.* Lactose drives *Enterococcus* expansion to promote graft-
versus-host disease. *Science* **366**, 1143-1149 (2019).
52. Siranosian, B.A*., et al.* Rare transmission of commensal and pathogenic bacteria in the
gut microbiome of hospitalized adults. *Nat. Commun.* **13**, 1-17 (2022).
53. Li, D., Liu, C.-M., Luo, R., Sadakane, K. & Lam, T.-W. MEGAHIT: an ultra-fast single-
node solution for large and complex metagenomics assembly via succinct *de Bruijn*
graph. *Bioinformatics* **31**, 1674-1676 (2015).
54. Kang, D.D*., et al.* MetaBAT 2: an adaptive binning algorithm for robust and efficient
genome reconstruction from metagenome assemblies. *PeerJ* **7**, e7359 (2019).
55. Alneberg, J*., et al.* Binning metagenomic contigs by coverage and composition. *Nat.*
Methods **11**, 1144-1146 (2014).
56. Sieber, C.M*., et al.* Recovery of genomes from metagenomes via a dereplication,
aggregation and scoring strategy. *Nat. Microbiol.* **3**, 836-843 (2018).
57. Wood, D.E., Lu, J. & Langmead, B. Improved metagenomic analysis with Kraken 2.
Genome Biol. **20**, 1-13 (2019).
58. Brown, C.T., Olm, M.R., Thomas, B.C. & Banfield, J.F. Measurement of bacterial
replication rates in microbial communities. *Nat. Biotechnol.* **34**, 1256-1263 (2016).
776

**Extended Data Figure 1: Paired fecal and oral microbiome samples in the mouse
experiment. a,** Bacterial ASV profiles displayed using stacked bar plots. Samples were collected
at three time points: pre (before treatment), d3 (3 days after treatment initiation), and w1 (one
week after treatment initiation). Missing samples in the second experiment were due to low
sequence coverage (samples with less than 1,000 reads were excluded). **b,** Total bacterial load in
fecal samples (circles). Bar heights represent the means, with error bars indicating the 95%
confidence interval. *P* values were calculated using a one-sided Wilcoxon signed-rank test. **c**,
Principal coordinate analysis plot of these samples. PC1 and PC2 represent principal component
1 and 2, respectively. The unit of total bacterial load is 16S copies per gram of feces for fecal
samples and 16S copies per swab for oral samples. DSS: Dextran Sulfate Sodium.

788
789 **Extended Data Figure 2:** Quantitative agreement between our method and FEAST[18] for
790 estimating the oral bacterial fraction in post-treatment fecal samples. DSS: Dextran Sulfate
791 Sodium.

793
Extended Data Figure 3: Emergence of new oral ASVs in mouse feces after antibiotic
treatment. Each panel plots data from one mouse. Mouse Abx_2B was excluded due to the
absence of its pre-treatment fecal sample. In each panel, individual oral ASVs are shown by
lines. ASVs that increased in absolute abundance after treatment (i.e., post-treatment average >
pre-treatment) are shown in red, while those that decreased or remained unchanged (post-
treatment average ≤ pre-treatment) are shown in blue. The black line in each panel represents the
total absolute abundance, which is the sum of all red and blue lines. Except for one oral ASV in
mouse Abx_1B (shown as a red dashed line), all ASVs displaying increased absolute abundance
(solid red lines) were not present in the pre-treatment fecal samples; they represent new ASVs
emerging from the oral cavity. Samples were collected at three time points: pre (before
treatment), d3 (3 days after treatment initiation), and w1 (one week after treatment initiation).
The unit for absolute abundance of oral ASVs in mouse feces is 16S copies per gram of feces.

806
Extended Data Figure 4: Oral bacterial ASVs identified from healthy human individuals
involved in the HMP dataset. a,b, Mean relative abundance (a) and prevalence (b) of ASVs
across 2,641 oral cavity samples (*x*-axis) and 291 paired fecal samples (*y*-axis). Black dashed
lines represent the cutoffs used to identify oral ASVs (see Methods for details). Each dot
represents an ASV. There is a total of 23,411 ASVs, where 178 ASVs identified as of oral origin
are highlighted in orange. **c,** Distribution of the 178 oral ASVs at the genus level.

Extended Data Figure 5: Validation of oral bacterial ASVs identified from healthy
individuals in patients with inflammatory bowel disease. This cohort consists of 16 patients
with Crohn's disease (CD), 42 patients with ulcerative colitis (UC), and 43 healthy controls
(HC). Both microbiome data and clinical metadata were obtained from a literature study[19].
Notably, none of the participants had taken antibiotics within the last three months prior to the
study entry. **a,** Impact of cutoff parameters on the estimated oral bacterial fraction in feces of
these participants. We systematically varied cutoffs for the mean relative abundance (0.01%,
0.1%), prevalence (5%, 10%), and the definition of ASV presence (0.01%, 0.1%) in the
computation of prevalence to establish the reference set of oral ASVs. For each reference set,
oral ASVs in their feces were inferred through exact sequence matching. The default parameter
combination used throughout the study is outlined in the red box. *P* values were calculated using
a one-sided Mann-Whitney U test. **b,** Percentage of inferred oral ASVs in feces that are also
found in paired saliva samples. **c,** Percentage of the total relative abundance of inferred oral
ASVs in feces contributed by those found in paired saliva samples. Two HC samples with a zero
oral bacterial fraction are not shown in all box plots.

Slope = -0.41 ± 0.03

829
Extended Data Figure 6: Inverse correlation between oral bacterial fraction and total
bacterial load in fecal samples from MSKCC allo-HCT recipients. Each dot in the plot
represents a fecal sample. Samples with zero oral bacterial fraction, or with total bacterial load
less than 1,000 16S copies per gram of feces, or collected 20 days before or 40 days after
transplantation were excluded from the plot and the linear regression analysis. The number of
remaining samples is 2,524. The red line represents the best linear fit, and the shading of the
same color indicates the 95% confidence interval. The plot also displays the estimated regression
slope and its standard error.

838
Extended Data Figure 7: Impact of interindividual variability on the regression slope
between oral bacterial fraction and total bacterial load. We generated synthetic datasets
using parameters estimated from the MSKCC allo-HCT cohort analyzed in Fig. 3 (**a**) and the
Crohn's disease cohort analyzed in Fig. 4 (**b**). In both panels, we systematically varied σ_F (Eq. 2)
from 0 to 2 while co-varying σ_f (Eq. 1) to maintain a constant ratio of σ_f/σ_F. For each synthetic
dataset, the regression slope was determined through linear regression between oral bacterial
fraction and total bacterial load in the log-log space. The red and blue lines represent the mean
slopes over 100 simulation runs, with the shaded regions of the same color indicating the
standard deviations. Vertical dashed lines in the panel (a) and (b) mark the σ_F values estimated
from the pre-antibiotic-prophylaxis samples in the allo-HCT cohort and from the healthy
individuals in the Crohn's disease cohort, respectively. At these σ_F values, the pure *Marker*
hypothesis predicted that the regression slopes are -0.36 and -0.30. For detailed information on
our simulation approach, please refer to the Methods section.

852
Extended Data Figure 8: Cross-validation accuracy for classifying total bacterial load in
fecal samples from MSKCC allo-HCT recipients. A schematic illustration of our classification
model is enclosed in the dashed box. Our model uses two threshold parameters, θ_o and θ_t, to
convert the oral bacterial fraction and total bacterial load into binary categories, respectively.
Given training data, we optimized the two cutoff parameters by minimizing the P value of
Fisher's exact test of independence, subject to the constraints $\theta_o \in [10^{-4}, 1]$ and $\theta_t \in$
$[10^3, 10^{10}]$. The optimized θ_o was subsequently applied to predict high or low bacterial loads in
the test set by comparing the oral bacterial fractions to θ_o. Simultaneously, we binarized the
observed bacterial loads by comparing them to θ_t. Accuracy was assessed by comparing the
predicted bacterial load categories to the observed bacterial load categories. In the boxplot, each
dot corresponds to a single 5-fold cross-validation split and the random train-test split was
repeated 50 times.

866
867 **Extended Data Figure 9: Distribution of oral bacterial fraction in fecal samples from**
868 **MSKCC allo-HCT recipients, categorized by stool consistency (a) and fungal cultivability**
869 **(b).** Each dot represents a fecal sample. FDR-corrected *P* values were calculated using a two-
870 sided Mann-Whitney U test.

871
Extended Data Figure 10: Estimating *Enterococcus* absolute abundance through the
Enterococcus-to-oral bacteria relative abundance ratio. Each data point represents a fecal
sample from MSKCC allo-HCT recipients. We excluded samples with zero relative abundance
of *Enterococcus* or zero oral bacterial fraction from both the plot and the Pearson's correlation
analysis, resulting in a subset of 2,765 samples. The red line represents the best linear fit, with
the shading in the same color indicating the 95% confidence interval. The absolute abundance of
Enterococcus on the *y*-axis was calculated by multiplying its relative abundance by the total
bacterial load measured by 16S qPCR, and thus has a unit of 16S copies per gram of feces.