1

Evaluation of metabolic and functional properties of cholesterol-reducing and GABA-producer *Limosilactobacillus fermentum* strain AGA52 isolated from lactic acid fermented Shalgam by using *in vitro* and *in silico* probiogenomic approaches

Ahmet Yetiman[1]*, Mehmet Horzum[2], Mikail Akbulut[3]

[1]Erciyes University, Faculty of Engineering, Food Engineering Department, 38030, Kayseri, Turkey

[2]Erciyes University, Graduate School of Natural and Applied Sciences, Food Engineering Department, 38030, Kayseri, Turkey

[3]Erciyes University, Faculty of Science, Department of Biology, 38030, Kayseri, Turkey

*Corresponding author: ahmetyetiman@erciyes.edu.tr

Abstract

The objective of this study is characterization of the genomic and metabolic properties of a novel *Lb. fermentum* strain AGA52 which was isolated from a lactic acid fermented beverage called "Shalgam". The genome size of AGA52 was 2,001,084 bp, which is predicted to carry 2,024 genes, including 50 tRNAs, 3 rRNAs, 3ncRNAs, 15 CRISPR repeats, 14 CRISPR spacers, and 1 CRISPR array. The genome has a GC content of 51.82% including 95 predicted pseudogenes, 56 complete or partial transposases and 2 intact prophages. The similarity of the clusters of orthologous groups (COG) was analysed by comparison with the other *Lb. fermentum* strains. The detected resistome on the genome of AGA52 was found to be intrinsically originated. Based on genome analyses many CAZYmes were identified that are responsible for carbohydrate degradation and adaptation to gastrointestinal conditions. Besides, it has been determined that AGA52 has an obligate heterofermentative carbohydrate metabolism due to the absence of the 1-phosphofructokinase (*pfK*) enzyme. Additionally, it is determined that the strain has a higher antioxidant capacity and is adaptable to gastrointestinal simulated environments. It was also observed that the AGA52 has antimicrobial activity against *Y. Enterocolitica* ATCC9610, *B. cereus* ATCC33019, *S. enterica* sv. Typhimurium, *E.coli* O157:h7 ATCC43897, *L.monocytogenes* ATCC7644, *K.pneumoniae* ATCC13883, and *P. vulgaris* ATCC8427. Cholesterol assimilation (33.9±0.005%) and GABA production capacities were also confirmed by "*in silico*" and "*in vitro*". Overall, investigation of genomic and metabolic features of the AGA52 revealed that is a potential psychobiotic and probiotic dietary supplement candidate and can bring functional benefits to the host.

Keywords: *Limosilactobacillus fermentum*, comparative-genomics, probiotic, cholesterol, GABA, psychobiotic

Introduction

The lactobacilli are a bacterial genus that is a member of Lactic acid bacteria (LAB) which can be found in fermented foods and the gastrointestinal microbiome of humans and animals [1]. They are widely used in the food biotech industry because of their well-documented beneficial and functional effects on health [2]. The members of the lactobacilli can produce different kinds of metabolic products such as bacteriocins and various organic acids, which generate an unsuitable environment for the growth of pathogenic and saprophytic microorganisms [3, 4]. Furthermore, metabolites or bacterial cells by themselves can show antiviral activity [5, 6]. Many species from the *Lactobacillus*, have been recognized as GRAS (Generally Recognized As Safe) and/or QPS (Presumably Qualified as Safe), due to their potential probiotic properties. Probiotics are defined as nonpathogenic living microorganisms that confer health benefits to the host when administered in adequate amounts by the International Scientific Association for Probiotics and Prebiotics (ISAPP) [7].

Lately, the genus *Lactobacillus* has been reclassified into 25 different genera, 23 of which are novel. In accordance with this new taxonomic order, *Lactobacillus fermentum* has been renamed *Limosilactobacillus fermentum* and qualified as gram-positive, coccoid, or rod-shaped, aerotolerant or anaerobic, and heterofermentative. *L. fermentum* has been generally isolated from fermented plant materials including cereals, dairy products, feces, manure, sewage, oral cavity, human vagina, and breast milk [8, 9]. Moreover, it has been reported that *L fermentum* exhibits health-promoting effects on the host when administered time-dependently. These effects are an increment of the amount of short-chain fatty acid in the colon, advancement of the gut microbiome composition, enhancement of the antioxidant activities in the colon, liver, kidney, and heart tissues of the host, and decrement of dyslipidemia and blood pressure. However, the type and intensity of the aforementioned effects vary from strain to strain [10-13].

Each strain of these microorganisms that have been recognized as probiotics has unique and

unusual properties which vindicates the studies for new probiotics. The biological activities of

probiotics are strain-specific and these properties cannot be hypothesized to other strains of

the same species. Therefore, it is important to reveal these properties through physiological

and metabolic tests, and probiogenomic approaches when an attention-grabbing new strain is

recovered from any niche [14-16].

The Shalgam is a traditional lactic acid fermented Turkish beverage that is characterized by

reddish color, sour-soft taste, and cloudy appearance [17]. It has been reported that Shalgam

exhibits presumptive health benefits against a variety of health risks and diseases due to the

high content of lactic acid bacteria present in its microbiome [14, 18, 19]. In its microbial

distribution, *Lactiplantibacillus plantarum* and *Lactocaseibacillus paracasei* are reported as

prevalent strains by several authors [19, 20]. Except those, *Lb. casei*, *Lb. brevis*, *Lb.*

parabrevis, *Lb. pentosus*, *Lb. buchneri*, *Lb. delbrueckii* subsp. *delbrueckii*, *Lb. fermentum*, *Lb.*

reuteri, *Lb.helveticus, Lb. gasseri*, *Lb. acidophilus*, *Pediococcus* spp., *Leu. mesenteroides*

subsp. *mesenteroides/ dextranicum*, *Leu. mesenteroides* subsp. *cremoris*, *Leu. mesenteroides*

subsp. *mesenteroides* species were also reported to be found in the shalgam microbiome [18,

21].

This study has aimed to evaluate the genomic, physiological, and metabolic characteristics of

Limosilactobacillus fermentum strain AGA52 by using "*in vitro*" and "*in silico*" approaches.

The strain AGA52 has been isolated from shalgam and as far as we know, there have been no

studies regarding the genomic and phenotypic characterization of a *Limosilactobacillus*

fermentum strain sourced from traditional lactic acid fermented beverage called "shalgam",

although several kinds of research have been done to enlighten its microbiota, chemical, and

volatile composition relationship. For this purpose, the strain's physiological properties were

examined under the screening and selection criteria of probiotics. Further, the whole genome

of the AGA52 was sequenced and analyzed with bioinformatic tools. This is the first report

clarifying genomic and metabolic characteristics of cholesterol-reducing and GABA-producer

Limosilactobacillus fermentum strain AGA52 isolated from lactic acid fermented Shalgam

beverage.

Materials and Methods

Materials and Methods

Growth Conditions of Bacterial Strain

Limosilactobacillus fermentum strain AGA52 was isolated from a Turkish fermented shalgam

juice (pH: 3.29) that was purchased from a local producer (Ali Göde: Hot) in Adana, Turkey.

A 10 mL of sample was diluted with 90mL of sterile physiological saline solution (0.85%)

and homogenized for 1 minute with a highspeed vortex (MS-3 Basic, IKA-Werke GmbH,

Staufen, Germany). Later, serial decimal dilutions were get set from the suspension and

100µL of each diluent was spread on MRS agar (Merck GmbH, Darmstadt, Germany). The

plates were incubated at 30°C for five days in an anaerobic atmosphere. The AGA52 isolate

has been chosen from the dilution of 10^{-5} and subjected to colony purification twice and,

catalase test and gram staining were applied to the pure isolate of the AGA 52. The cryo

stocks of AGA52 were prepared in MRS broth (Merck) with 25% glycerol and were stored at

-80°C.

DNA extraction, identification, whole-genome sequencing, and assembly

First, *Lb. fermentum* AGA52 cryo culture was subcultured twice in MRS broth (Merck)

followed by incubation anaerobically at 37 °C for 24h. A 1 mL fresh culture was pipetted onto

a sterile microcentrifuge tube (2mL) and centrifuged at 6000 x g for 10 min at 4 °C. Later,

Total DNA was harvested from the cell pellet using the PureLink Genomic DNA Mini Kit

(Invitrogen, Thermo-Fisher Scientific, Carlsbad, CA, USA) in accord with the manufacturer's

instructions for gram-positive bacteria. The quality and concentration of genomic DNA were

checked by a Qubit 3.0 fluorometer (Invitrogen, Thermo-Fisher Scientific, Carlsbad, CA,

USA) and agarose gel (1.5 %). The test strain was determined by full-length nucleotide

sequencing of the 16S rRNA gene. PCR amplification and further steps were carried out as

per Yetiman et al. [14]. Following the verification of the strain AGA52 using 16S rDNA, the

whole genome sequencing libraries were constructed using Nextera XT DNA Library

Preparation Kit (Illumina, San Diego, CA, USA) and sequencing was carried out by Illumina

Novaseq platform as paired-end (PE) 2x150 bases read. The adapter sequences were trimmed

with the JGI-RQC Filter pipeline (BBTools v38.22) by preserving the reads belonging to the

strain AGA52 and trimmed raw data were assembled with SPAdes v. 3.15.3 [22].

Bioinformatic analyses

Genome annotation was performed by using NCBI Prokaryotic Genome Annotation Pipeline

(PGAP) [23]. Next, RASTtk and BV-BRC annotations have been conducted for comparison

with PGAP [24, 25]. A BLAST Ring Image of the genome of the AGA52 strain and other

compared *Lb. fermentum* strains were generated using BRIG v0.95 [26]. The calculation of

orthologous average nucleotide identity values (OrthoANI) of the AGA52 and others

compared *Lb. fermentum, Lb. gastricus,* and *Lb. reuterii* strains were executed by OrthoANI

tool v0.93.1 [27]. Hierarchical clustering analysis of the strain AGA52 and other 45 *Lb.*

fermentum genomes were done via the genome clustering service of IMG/M [28]. Prediction

of metabolic pathways of *Lb. fermentum* AGA52 was carried out using BlastKOALA for

scanning against the KEGG database [29]. Prediction of CAZymes (Carbohydrate active

enzymes) was implemented by use of dbCAN2 meta server [30]. A cluster of Orthologous

Groups (COG) prediction based on protein FASTA sequences was conducted via webMGA with default options [31]. The upSet plot has been visualized by an intervene web-basedsed intersection tool for the determination of the shared COG between the AGA52 and other reference genomes [32].

The prophage elements on the genome of AGA52 were identified with the PHASTER-Phage Search Tool Enhanced Release [33]. All protein-coding sequences obtained from PHASTER have been screened against the non-redundant protein (NR) database by performing protein-BLAST to identify the horizontally transferred genes. If a gene's homologous protein was got to match a microorganism other than *Lb. fermentum* $\geq$ 80%, that gene was noted as horizontally transferred [14]. A resistome screening was conducted by scanning the complete genome sequences of the AGA52 strain versus the ResFinder 4.1, CARD, BV-BRC and KEGG databases, respectively [24, 29, 34, 35]. As with the phage elements, horizontal gene transfer screening was performed within the detected resistome. Prediction of CRISPR-Cas regions were implemented via the CRISPR finder online tool (http://crispr.i2bc.paris-saclay.fr/Server/). The complete genome sequence of *Lb. fermentum* AGA52 was submitted to NCBI under accession number (CP091132.1).

Carbohydrate fermentation

According to the manufacturer's protocols, the carbohydrate fermentation patterns of the AGA52 strain were determined by API 50 CHL kit (BioMérieux, Marcy l'Etoile, France).

Determination of antibiotic susceptibility

Antibiogram assays were performed to find out the resistance or sensitivity of the strain

AGA52 in return for commonly used antibiotics. Ready to use commercial antibiotic disks

[azithromycin, methicillin, vancomycin, amikacin, kanamycin, tetracycline, penicillin G

(Bioanalyse, Yenimahalle, Ankara, Turkey); amoxicillin, amoxicillin, ampicillin,

erythromycin, oxacillin, carbenicillin, streptomycin, rifampicin (Oxoid, Basingstoke,

Hampshire, UK)] were used for antibiotic susceptibility testing of *Lb. fermentum* AGA52.

The application of disk diffusion assay was carried out considering the Kirby-Bauer method

[36] Interpretation of inhibition zone (mm) was carried out as described by Clinical and

Laboratory Standards Institute's performance standards for antimicrobial testing [37].

Determination of probiotic properties

To determine probiotic properties of the AGA52: β-haemolysis, *in vitro* simulation of gastric

digestion, cell surface hydrophobicity, cellular auto-aggregation tests, and antibacterial

activity assay against several pathogens were performed, respectively. β-hemolytic activity of

the AGA52 was evaluated by using 5% sheep blood containing Columbia agar plate. The

isolate was streaked on the Columbia agar followed by incubation at 37°C for 48 h under

anaerobic conditions [38]. To check the response of the AGA52 against gastric digestion, the

gastric environment was simulated as described by Zhang et al. [39] with a slight

modification (1% bile salt was used) for preparation of artificial gastroenteric juice [14].

Cell surface hydrophobicity and auto-aggregation assays were applied as for that Krausova et

al. [40]. The bactericidal activity of the strain AGA52 has been tested by using the agar well

diffusion method [41]. The supernatant of 48h grown bacterial cells was assayed against

Escherichia coli O157:H7 (ATCC 43895), *Staphylococcus aureus* (ATCC 25923), *Bacillus*

cereus (ATCC 33019), *Salmonella enterica* sv. *Typhimurium* (ATCC 14028), *Proteus*

vulgaris (ATCC 8427), *Listeria monocytogenes* (ATCC 7644), *Yersinia enterocolitica*

(ATCC 9610) and *Klebsiella pneumoniae* (ATCC 13883).

Antioxidant Activity Assays

The antioxidant activity of the AGA52 was determined by using DDPH (2,2-diphenyl-1-

picrylhydrazyl) and ABTS (2,2 azino-bis(3-ethylbenzothiazoline-6-sulfonic acid)) methods.

For this purpose, cell-free supernatant (CFS) was collected from MRS broth via centrifugation

at 3500rpm for 15 min. DPPH scavenging activity was assessed according to Ozturk et al.

[42] with some modifications. The reaction mixture was prepared by adding 2 mL freshly

prepared 0.2mM DPPH onto 1 mL four times diluted CFS. The mixture was incubated in the

dark for 30 min at 25°C. Later, the mixture was centrifuged at 5000 rpm before measuring the

absorbance at 520 nm with five replicates. The blank control includes dH_2O and DPPH

solution. The DPPH scavenging activity has been computed as scavenging activity $\% = (1 -$

$A_{test}/A_{blank}) \times 100$.

ABTS assay was performed as for that Pieniz et al. [43] with some modifications. ABTS was

dissolved in water (7 mM). Later, ABTS stock solution was mixed with 2.45 mM potassium

persulfate (final concentration) for the formation of radical cation ($ABTS^+$) and the mixture

was incubated in dark for 16h at 25°C before usage. The stock solution can be used for only 3

days. Before application, $ABTS^+$ solution was diluted with phosphate buffer (pH: 7.2) to get

an absorbance of 0.700±0.03 at 734 nm. Likewise, The CFS was also diluted four times with

phosphate buffer (pH: 7.2). Next, 40 mL of CFS was added to 4mL of $ABTS^+$ solution, the

mixture was incubated in dark for 5 min and the absorbance was measured with five

replicates. Phosphate buffer (pH: 7.2) was used as blank and control. The percentage of

inhibition was predicted according to the ascorbic acid standard curve (0-9 µg/mL).

Cholesterol assimilation assay

The cholesterol utilization by the AGA52 in MRS broth was analysed by O-phthaldehyde (OPA) method as explained by Rudel, Morris [44] with some modifications. The AGA52 was grown in MRS broth (0.25% dextrose + 0.3% ox bile containing) supplemented with 100ppm cholesterol (5-cholesten-3β-ol (Sigma, Merck GmbH, Darmstadt, Germany), dissolved in 2-propanol) at 37 °C for 24 and 48h, respectively. After incubation, the cultures were centrifuged (3500rpm, 15min, 25°C) and CFS were used for determination of residual cholesterol. Next, 0.5mL of CFS was mixed with 2mL of KOH (50% wt/vol) and 3mL of absolute ethanol, vortexed for 1 min and later heated at 60°C for 15 min. After chilling, 3mL dH_2O and 5mL of hexane were added to the mixture and it vortexed for 1 min. Afterwards, 2.5 mL of hexane phase was transferred to another glass tube and then was kept for evaporation at 80°C. The residue was directly dissolved in 4 mL of OPA (0.5 mg/mL in glacial acetic acid) and incubated at 25°C for 10 min. Later, 2 mL of H_2SO_4 (98%) was added gently and vortexed for 1min and the mixtures incubated again at 25°C for 10min, and absorbance was measured at 550nm (Shimadzu UV-1800 UV/VIS spectrophotometer, Tokyo, Japan). The cholesterol assimilation was calculated according to the difference between the control (uninoculated MRS broth) and test samples.

Confirmation of GABA production by ^{1}H-nuclear magnetic resonance

The monosodium-glutamate (MSG) standard, γ-aminobutyric acid (GABA) standard and deuterium oxide were purchased from Merck GmbH (Darmstadt, Germany). The two-times subcultured AGA52 culture was inoculated in 1% MSG containing MRS broth (0.25% dextrose) and incubated at 37 °C for 48h. Next, the culture media was centrifuged at 3500 rpm for 15 min and cells were discarded. The CFS was lyophilised by using a freeze-dryer (Christ,

Alpha 2–4 LSCplus, Germany). Next, aprroximately 100mg of CFS, MSG, and GABA were

transferred into a cylindiric glass NMR tube and dissolved with deuterium oxide, respectively.

Each sample was subjected to ^{1}H-nuclear magnetic resonance by using an NMR spectrometer

(Bruker, Model 400, MA, USA) at 400mhz for 5 min. Afterwards, NMR spectra were

analysed by comparing standards via data analysis software (Bruker, TopSpin, MA, USA) of

the device.

Results and Discussion

Functional genomic characterization

The genome of the *L. fermentum* strain AGA52 comprised a circular chromosome of

2,001,184 bp with a 51.82% GC ratio, and it contains 2.024 genes, 1.873 protein-coding

sequences, 50 tRNA, 3 rRNA, 3 non-coding RNA, 1 CRISPR array region and 95

Pseudogenes. The BLAST atlas of the genome of AGA52 and other compared well-known *L.*

fermentum genomes were displayed in Fig 1, and differences between each other can be

noticed clearly due to the gaps in the genomes. Among these gaps, the detection of genomic

islands is possible due to the corresponding decrease in GC content and the absence of these

islands in other strains. Generally, integrases, insertion sequences, and transposases are

positioned in and/or around these islands which is an indicator of potentially horizontal

transferred genes [45, 46].

The orthoANI comparison results amongst the AGA52 and the aforecited genomes were

presented in Fig 2. Based on the OrthoANI values, the genome of the *L. fermentum* AGA52

bears a resemblance to the genomes of SK152 (99.18%), ATCC 14931 (99.15%), DSM20052

(99.11%), YLF016 (99.11%), IFO 3956 (99.04%), CECT 5716 (98.92%), and B44(98.92%),

respectively. It has been informed that the recognition of the genomes of the two species as

the same depends on the ANI value above 96% [27]. Besides, It has been known that SK152,

ATCC14931 and DSM 20052 originated from fermented plant materials [45, 47]. On the

other side, YLF016 (Yak's GIT), CECT5716 (human breast milk), and IFO3956 (cheese) are

derived from non-plant sources [5, 39, 48]. The genetic distance between the genome of

AGA52 and the mentioned genomes can be explained by differences or similarities in the

strain's isolation niche, evolutionary histories, and the conditions exposed [14, 49]. In

addition, assembly details and annotated genome properties of *L. fermentum* AGA52 and

higher orthoANI similarity shown genomes were summarized in Table 1.

Moreover, the protein-coding sequences of the genome of the AGA52 have been screened for

determination of the clusters of orthologous groups (COG) and categorized into 23 different

function classes in Fig 3. A detailed demonstration of the number of shared orthologous

proteins by the UpSet plot can be seen in Fig 4. According to the hierarchical clustering

results (Fig 5.) which are based on clusters of orthologous groups (COG) profiles of 46

different *L. fermentum* strains, the AGA52 shared the highest similarity with KMB612 (0.95),

KMB613 (0.95), AF15-40LB (0.95), AF16-22LB (0.95), 279 (0.94), 779_LFER (0.94), and

VRI-003 (0.94). The isolates of KMB 612 and KMB 613 were isolated from the Bryndza

cheese (Fig S1). The characteristic of cheese is that it is produced from sheep's milk. Based on

past studies on *L. fermentum*, It is possible to make certain assumptions to explain this genetic

relationship between AGA52, KMB 612, and KMB 613. This strain may have been

transferred to sheep's milk through the consumption of plant-based fermented feed (e.g.,

silage) or may itself be an autochthonous member of the existing sheep microbiome [45, 49].

In the IMG/M database, AF15-40LB, AF16-22LB and 279 genomes were reported as faecal

originating, while 779_LFER has been expressed to originate from the respiratory system. It

was also described in the IMG/M that VRI-003 was isolated from a lyophilized commercial probiotic formulation from Australia. Nonetheless, It has been reported that VRI-003 has an increasing effect on interferon-gamma (IFNγ) production in whole blood culture [50]. As above, a similar analogy can be drawn between AGA52 and these strains.

Predicted CRISPR structures in the genome of the AGA52 consist of 1 copy of CRISPR-associated endonuclease Cas9, 1 copy of type II CRISPR-associated endonuclease Cas1, 1 copy of type II-A CRISPR-associated protein Csn2, 1 copy of CRISPR-associated endonuclease Cas2, 15 repeat regions, 14 spacer regions, and 1 array region (Table S1). CRISPR constructs and related (Cas) proteins serve as adaptive immune systems against prophages and foreign plasmids in prokaryotes and have been used for genotyping and genetic engineering purposes since the day they were discovered [51, 52]. Especially, from the point of view of the development of next-generation probiotics with enhanced functionalities, this technology has uncovered new opportunities. In this sense, It is possible to improve the therapeutic potential of the probiotics for vaccine delivery or modulation of the host immune response [53]. Therefore, the prediction and characterization of the CRISPR-Cas structures in the genome of the *L. fermentum* strain AGA52 could enable the development of novel biotherapeutics by regulating the expression and production of existing functional metabolites or acquiring novel biosynthetic gene clusters (e.g bacteriocins, non-bacteriocin antimicrobial compounds, metabolic products) by using itself.

Prophages, related horizontal gene transfer, and mobilome

The phage scan results revealed that AGA52 had three phage sites in its genome, two intact and the other questionable. The intact phage regions (regions 2 and 3) were displayed similarity with Lactob_LfeSau_NC_029068 (32.2 kb) and Lactob_521B_NC_048752 (11.7

kb), respectively. It has been observed that Lactob_LfeSau presented the highest matching protein (16) among the identified phages which is also shown in Table S2. This phage is the most common one found to be among the *L. fermentum* strains. Only, Lactob_LfeSau has attL/attR sequences and integrase (PP_00944) except for region 1 (Klebsi_phiKO2) and 3 (Lactob_521B). In a bacterial genome, the presence of the integrase is a functional identifier for pathogenicity islands, phages, and integrative plasmids [46, 54]. Besides, GC% content of the phages is quite lower than the average GC% content of the genome of the AGA52. All phage regions have been located around or in the genomic islands which are seen in BLAST ring alignment (Fig 1). Both attachment sites are positioned forward strand and upstream of the integrase (attL: 922221-922233; attR:947491-947503). Merely, Lactob_LfeSau was found to comprise all packaging/head/tail gene clusters, and DNA packaging genes. Unlike, previous studies no endolysin gene has been observed among all of the identified phages [55, 56]. Endolysins are phage proteins that rapidly break the bacterial cell wall down and release new viral particles [57]. This means that identified phages are temperate for the strain AGA52 and each member of identified phage regions was listed in Table S3-S5.

Horizontal gene transfer (HGT) amidst bacteria generally appears in so far as bacteriophage infection or natural competence [14, 58]. Based on protein BLAST results for screening HGT from phage-related proteins, 0.79% (16 genes) of the genome of AGA52 have been acquired by phage-associated HGT from other bacteria and viruses, eg. *Ligilactobacillus animalis*, *Pediococcus acidilactici*, *Lactobacillus* sp., *Siphoviridae* sp. (bacteriophage), *Lentilactobacillus* sp., *Limosilactobacillus reuteri* (Table S6). On the other side, all of the detected phage-associated transposases were only found in the Lactob_521B phage. Among the phage elements of Lactob_521B, 11 transposase genes (PP_01962, PP_01963, PP_01965, PP_01966, PP_01968, PP_01970, PP_01972, PP_01973, PP_01975, PP_01976, and PP_01977) that may have been acquired by recombination, repair and replication. In addition,

342 as for the whole genome-based transposase search result, IS3, IS4, IS6, ISL3, IS30,

343 IS200/IS605, IS256, and IS982 families' members of transposases have been identified by

344 using IS Finder and were summarised in Table S7.

Phenotypic antibiotic resistance and Safety-related genes assessment

In agreement with the European Food Safety Authority (EFSA) suggestions, a probiotic strain

should not hold gained or transferable resistome to clinically used drugs [59]. Therefore, it is

important to determine antibiotic resistance phenotypically and genotypically. Assessment of

the antibiotic susceptibility of the AGA52 has been performed following The Clinical and

Laboratory Standards Institute's performance standards. The value of inhibition zones (ZOI)

that belongs to tested antibiotics against the AGA52 and resistome search results were

presented in Table S8. Based on the antibiogram results, AGA52 was found to be sensitive

(ZOI ≥ 20mm) to Ampicillin (10 μg), Penicillin G (10 U), Carbenicillin (100 μg),

Amoxycillin (25 μg), and Rifampicin (5 μg), while it displayed resistance (ZOI ≤14mm) to

Methicillin (5 μg), Oxacillin (1 μg), Streptomycin (10 μg), Vancomycin (30 μg), Amikacin

(30 μg), and Kanamycin (30 μg). Unlike these, intermediate sensitivity (ZOI ~ 15-19mm) to

Azithromycin (15 μg) and Erythromycin (10 μg) was also observed.

Based on Resfinder and CARD analyses, there are no antibiotic-resistance genes have been

found in the genome of the AGA52. Nonetheless, sixteen antibiotic resistance genes have

been found by using KEGG and BV-BRC databases which were displayed in Table S8. The

detected genes were found to be related to β-Lactams (4), aminoglycosides (7), peptide-

antibiotics (3) and multidrug efflux transporters (2). It is well known that lactobacilli show a

higher resistance profile to aminoglycosides. Especially, *Lactobacillus* species have been

acknowledged as resistant to vancomycin under their intrinsic peptidoglycan precursors

containing D-lactate instead of D-alanine at the C-terminus [60]. The vancomycin resistance

genes were detected in the genome of the AGA52 comprises of *mraY, alr*, *ddl, murF, murG*,

and *VanS* which were presented in Table S9. Interestingly, detected vancomycin resistance

genes showed differences from previous studies [14, 39]. Especially, the *VanX* gene could not

be found which is highly specific for hydrolyzation of D-ala-D-ala dipeptides and is an

important precursor of the cell wall. This might be caused by variations in tolerance or gene

expression [61]. Additionally, as an aminoglycoside, streptomycin resistance responsible *gidB*

gene has also been observed. However, no specific resistance genes were identified for

kanamycin and amikacin even if resistance subsists. This could be caused by phenotype do

not completely represent genotype [62]. Aside from these, *blaI, mcrA, pbp2A*, and *penP* genes

responsible for β-Lactam resistance have been determined which can be seen in Table S8. It is

well-known that oxacillin and methicillin resistance is related to penicillin-binding proteins

(*mcrA, pbp2A*). Despite the presence of beta-lactamase (*penP*) and penicillinase repressor

(*blaI*), AGA52 exhibited a sensitive profile to ampicillin, penicillin, carbenicillin, and

amoxycillin. This would be explained by the aforementioned case for aminoglycosides. On

the other hand, there are no resistance genes were found for the macrolides, tetracyclines, and

rifamycins tested in this study. However, strain AGA52 exhibited an intermediate and

sensitive pattern to these antibiotics (Table S8). This situation could be derived from

membrane impermeability and/or multidrug efflux transporters (the presence of *efrA* and *efrB*

is confirmed in Table S9 and S12) in reduced susceptibility to these drugs [63]. In addition,

daptomycin-specific *pgsA*, *mprF*, and *gdpD* genes were also identified in the genome of the

AGA52. Due to increasing concerns regarding various food niches and/or common bacteria

that might serve as potential reservoirs for resistome, the presence of transferable

antimicrobial resistance genes in probiotics is undesirable which is used for humans or

animals. Lactobacilli have intrinsically originated resistance to various antimicrobial

compounds, and it is well-known that such resistance has not been related to any particular

safety concerns. But, the intrinsic resistome on the bacterial chromosome must not be flanked

by transposases or integrases. According to the protein BLAST results for the resistome

detected in this study, no horizontally transferred gene was found (Table S9).

Carbohydrate fermentation patterns and active enzymes

Limosilactobacillus fermentum acquires ATP through heterofermentative carbohydrate

fermentation based on strain-specific sugar preference. Uncovering strain-specific metabolic

tendencies is important due to the prediction of industrial applicability and their probiotic

potential. Since, carbohydrate-active enzymes (CAZYmes) also have a probiotic function, e.g.

xylose-isomerase stimulation of gut persistence [64]. According to API 50 CHL test results,

the AGA52 can utilize 13 different carbohydrates out of 49 being tested. At first glance, the

utilization of D-ribose, D-xylose, D-galactose, D-glucose, D-fructose, D-cellobiose, D-

maltose, D-lactose, D-melibiose, D-sucrose, and D-raffinose can be comprehended from

Table S10. *L. fermentum* is a cohesive species and could ferment various kinds of sugars. This

status is directly involved in the presence of the genes responsible for the CAZYmes and

transporters. Most of the transporters associated with carbohydrate metabolism are positioned

in the phosphoenolpyruvate-dependent sugar phosphotransferase system (PTS) [65]. For

strain AGA52, the entire PTS enzymes were encoded by its genome and it consists of PTS

System Enzyme I (general enzyme gene, *ptsI*, K08483), phosphocarrier protein HPr gene

(*ptsH*, K02784), and 13 complete or incomplete substrate-specific enzyme II (EII) complexes

genes (Table S13). These complex genes seem cellobiose, fructose, mannose, sucrose, beta-

glucoside, and ascorbate specific and their substrate specificity is unidentified. However, it is

well-known that various sugar transporters can bring in more than one substrate [58]. In

addition, several carbohydrate-specific common ABC transporters were listed in Table S12.

It has been observed that the AGA52 has not encoded 1-phosphofructokinase (*pfK*) enzyme

gene on its genome. Based on previous studies, it is commonly accepted that species lacking

this enzyme have an obligate heterofermentative carbohydrate metabolism and they also

produce CO_2, lactate, and ethanol via a heterofermentative pathway [66, 67]. The same case is

valid for AGA52. Moreover, the presence of l-arabinose isomerase, l-ribulose kinase, and

ribulose phosphate epimerase enzymes are confirmatory to a lesser extent that a lactobacilli

strain has an obligate heterofermentative pathway [2, 68]. In this study, both L-arabinose

isomerase [EC:5.3.1.4] and ribulose-phosphate 3-epimerase [EC:5.1.3.1] could only be found

in the genome of the AGA52 (Table S11). Nevertheless, it could be implied that AGA52 has

an obligate heterofermentative pathway. Meanwhile, the key enzymes responsible for the

EMP, PK and Leloir pathways and their copy numbers are presented in Table S14 by being

compared with well-known *L.fermentum* strains. Among the compared strains, all key

enzymes are available for the aforecited pathways apart from phosphofructokinase 1,

fructose-bisphosphate aldolase, and L-ribulose kinase. These findings are also confirmatory

for the mentioned situation above. The CAZYmes encoding genes connected with the intact

EMP (Fig S1) and PK (Fig S2) pathways have been identified in the genome of the AGA52

and presented in Table S11.

The AGA52 can synthesize both isomers of lactate and has got four copies of L-lactate

dehydrogenase (*ldh*), three copies of D-lactate dehydrogenase (*ldhA*), and one copy of D-

lactate dehydrogenase (*dld*; quinone) enzymes encoding genes on its genome. The difference

between *dld* and others is that it is a peripheral membrane-specific dehydrogenase which

plays a role in respiration. The electrons originating from D-lactate oxidation are turned over

to membrane soluble quinone pool [69]. Apart from these, the AGA52 also possesses two

copies of alcohol dehydrogenase (*adhP*) enzyme encoding genes. The reaction products of

this enzyme differ depending on the substrate catalyzed. The enzyme produces aldehydes as a

product in reactions where primary alcohols are substrates, while ketones are released as

products when secondary alcohols are substrates [70]. The effect of the reaction products on

the specific flavour of the shalgam cannot be ignored.

Following KEGG mapper results, the AGA52 has a total of 158 carbohydrate metabolism-

related genes which are encoded in its genome. The distribution of those genes is as follows;

18 Embden–Meyerhof–Parnas pathway genes, 14 phosphoketolase pathway genes, 6 TCA

cycle-related genes (Fig S3), 6 pentose and glucuronate interconversions associated genes, 13

galactose metabolism genes, 11 fructose and mannose metabolism genes, 3 ascorbate and

aldarate metabolism genes, 13 starch and sucrose metabolism genes, 19 amino sugar and

nucleotide sugar metabolism genes, 20 pyruvate metabolism genes, 6 glyoxylate and

dicarboxylate metabolism genes, 9 propanoate metabolism genes, 13 butanoate metabolism

genes, 3 inositol phosphate metabolism genes, and 5 C5-branched dibasic acid metabolism

genes. Moreover, the number of identified CAZYmes encoded in the genome of AGA52 is as

follows; 32 Glycosyl transferases, 1 auxiliary activity responsible enzyme, 3 carbohydrate-

binding modules, and 38 Glycoside Hydrolases. There is no existence of carbohydrate

esterases has been observed in CAZY annotation results. Further, it can be seen in Fig 6 that

the CAZYmes patterns of ATCC 14931, DSM 20052 and SK152 are similar to AGA52.

Overall, the characterization results revealed that the AGA52 has an acceptable potential for

carbohydrate degradation and adaptive nature specific to the gastrointestinal tract.

Probiotic properties and antioxidant activities

The probiotic characterization tests were carried out for confirmation of the presence or

absence of the features responsible for becoming probiotics. First, the result of the β-

haemolysis assay revealed that AGA52 does not have β-hemolytic activity. The cell surface

hydrophobicity (CSH) and autoaggregation capacity (AC) were determined as 73,6% (with

xylene) and 44%, respectively. The CSH is essential for understanding probiotics' overall

adhesion capacity, which is generally measured by assessing the bacterial affinity to a

hydrocarbon (e.g. xylene, n-hexadecane, hexane) solvent [40, 71]. The hydrocarbon affinity

method does not reveal exact CSH, instead, it expresses the presence of van der Waals and

electrostatic interactions which are significant factors influencing the overall adhesion

capacity [72]. Recently, it has been reported that bacteria which has a higher CSH can bind

better to intestinal epithelial cells [73]. Similar to the present study, approximate CSH values

were reported for *L.fermentum* SJRP40 (79.69%), SJRP46 (80.57%), and *L. reuteri* K18

(76%) by different studies [40, 73]. CSH value of *L. fermentum* YLF016 (≥65%) has been

reported as low as than AGA52. Moreover, the lipoteichoic acid (LTA) production

responsible genes (*MurJ*, *LtaS, LtaS, LafC, LafB, LafA, dltD, dltA*, and *dltB*) that can be

associated with CSH were identified and their putative functions were explained in Table S15.

Autoaggregation ability for probiotic bacteria is crucial due to its direct relationship to the

adhesion of epithelial cells and mucosal surfaces and is also a phenotypic marker for adhesion

[74]. Due to the AC, bacteria can generate cellular aggregates which promote persistence in

the intestine [75]. Bacterial cell wall components and other structures, like mucus-binding

proteins, adhesins, surface layer proteins, fibronectin-binding proteins, exopolysaccharides,

and lipoteichoic acids give an advantage to bacteria for colonization and adhesion to host

epithelial cells [74, 76, 77]. In this study, the presence of some of these structures has been

confirmed in the genome of AGA52 and summarised in Table S15. The AC of the AGA52

shows a similar pattern to previously studied strains such as colostrum-originated *L. reuteri*

K14 (44.4%) and infant feces-originated *L. reuteri* E1M1C (46.3%). Besides, Li et al. [78]

reported between 0.86 to 65.15% autoaggregation values for a variety of *L. fermentum* strains

which were isolated from different Chinese fermented foods.

Resistance to stress factors of gastrointestinal systems, such as long-term survival and

colonization are uniquity of Lactobacilli. Uncovering host-probiotic interaction regulation

mechanisms have become more significant due to the gradually increasing number of

microbes used as probiotics. The survival behaviour of AGA52 was tested against

physiological pH, and bile salt environment and mimicked the gastrointestinal fluid. Broadly,

survivability at low pH of most Lactobacilli differs between 2.5 to 3.5 which is a crucial

selection criterion for potential acid-tolerant probiotic strains. During gastric transit,

probiotics must survive at pH 3.0 as low and remain alive for at least four 4 hours before

reaching the lower tract [79]. It has been observed that the AGA52 has a tolerance to an acidic

environment with a survival rate of 20±0.3% in pH 3.0 (with an initial load of 8.74 log

CFU/mL). Besides, the viability goes down to 0.09±0.01% at pH 2.0 (Fig 7B) which overlaps

the previous studies [14, 39]. Furthermore, the survival profile to bile salts (0.2-1% (w/v)) was

checked and the AGA52 presented a moderate survival pattern (38±0.1%) after 4h (with an

initial load of 9.19 log CFU/mL). Commonly, it is strongly suggested that approximately

0.3% bile salts concentration is proper for probiotic selection purposes [80]. Based on the

survival pattern at 0.3% bile salt concentration (64±0.1%), the AGA52 appears to partially

meet the requirements of this criterion (Fig 7A). Apart from these, survivability of the

AGA52 in the gastrointestinal simulated environment was determined for understanding

whether the strain shows tolerance or not. Because, gastrointestinal tolerance is significant for

the colonization and metabolic activity of probiotics to stimulate a positive effect on the host

[39]. The survival patterns were determined as 7.40 and 8.18 log CFU/mL for gastric and

intestinal fluids, respectively (Fig 7C and D). The resistance of the mentioned stress factors is

associated with the existence of the putative genes related to stress resistance, DNA and

protein protection/repair and these are listed in Table S15. Especially, cation-proton

antiporters (*NapA*, *NhaC*, *NhaP*), ABC ATPases, F_0F_1-type ATP synthase, and chaperones

(Clp ATPases) are responsible for homeostasis and intracellular pH regulation which have

contributed to acid and bile resistance. Similarly, the same acid bile resistance mechanisms

have been reported in *L. plantarum* Y44 [81], *L. amyloticus* L6 [82], and *L.fermentum*

YLF016 [39]. On the other side, choloylglycine hydrolase (*cbh*) and inorganic

pyrophosphatases (e.g. *PpaC*) were supposed to be prominent in acid bile resistance.

Differently, based on antimicrobial activity assay results, CFS of the AGA52 showed a zone

of inhibition (> 5 mm) versus all tested pathogen strains except for *S.aureus* (Table S16). As

far as we are concerned, inhibition of test pathogens by CFS of the AGA52 has originated

from the produced organic acids by itself. Because this strain does not encode any bacteriocin

production-responsible gene cluster in its genome.

Antioxidant activity is one of the most important probiotic functions that are directly related

to health benefits. Following antioxidant assays, DPPH and ABTS inhibitions were

determined as $92.9\pm0.01\%$ and $85.3\pm0.065\%$, respectively. DPPH and ABTS are well-known

free radicals which are generally used for testing the free radical scavenging activity of

antioxidants and can be analyzed colourimetrically. These strong inhibition values are related

to the antioxidant enzymes such as *trxA* (Thioredoxin), *trxB* (Thioredoxin reductase), *tpx*

(Thiol peroxidase), *dsbA* (Thiol disulfide oxidoreductase), *arsC* (Arsenate reductase), *Gor*

(Glutathione reductase), *nrdH* (Glutaredoxin) and other putative antioxidant genes. The

identifiers and putative functions of aforecited genes were summarized in Table S15. The "*in*

vitro" and "*in silico*" findings regarding the antioxidant activity of AGA52 are compatible and

further demonstrate that it can adapt to higher oxidative stress environments.

Cholesterol assimilation

Cardiovascular and coronary artery disease risks are directly correlated with dyslipidemia and are important public health concerns. Dyslipidemia generally occurs due to unbalanced concentrations of cholesterol, low-density lipoprotein cholesterol, (LDL-C), triglycerides, and high-density lipoprotein (HDL). Different researchers have reported that various *Lb. fermentum* strains showed blood cholesterol-lowering effects [10, 11]. In this regard, similar cholesterol-decreasing effects have been observed "*in vitro*" for this study. Based on cholesterol assimilation assay results, the AGA52 was assimilated 28.8±0.003% of the 100ppm cholesterol for 24h, while it has been utilized the 33.9±0.005% of total cholesterol added to MRS broth for 48h. In previous studies, 11.51±1.44%, 37.19±4.99%, and 40.84±1.90% cholesterol assimilations were reported for *Lb. fermentum* strains NCIMB 5221, NCIMB 8829, and NCIMB 2797, respectively [83]. In another study, cholesterol assimilation ranges from 28.3-88% were described for different *Lb. fermentum* strains by Pan et al. [84]. The molecular mechanism of hypocholesterolemic effects is generally explained by the presence of bile salt hydrolase (*bsh* or *cbh*) enzymes, or the production of short-chained-fatty acids that inhibit enzymes like 3-hydroxy-3-methylglutaryl coenzyme A which is component of ketogenesis pathways. In this study, the presence of choloylglycine hydrolase (*cbh*) enzyme which has bile salt hydrolase like activity was confirmed "*in silico*". The presence of organic acids, like propionic, acetic, butyric, and lactic acids was also confirmed by GC-MS (method and data not shown) which is another factor related with cholesterol assimilation.

GABA production as a psychobiotic feature

Gamma-aminobutyric acid (GABA) is a free amino acid which is produced from L-glutamic acid or MSG by Glutamate decarboxylase (GAD, EC 4.1.1.15) and is extensively found in animals, plants and microbes [85]. GABA has multiple physiological functions and acts as

major inhibitory neurotransmitter which delivers chemical message in the central nervous

systems of mammalians and has a significant role on stress response, cognition and behaviour

[86]. Besides, GABA supplementation is positively affects depression, sleeplessness,

immunity, blood pressure regulation, visual cortical function, anxiety, and menopausal

syndrome [87-91]. Various lactobacilli strains such as *Lb. fermentum*, *Lb. plantarum*, *Lb.*

helveticus, *Lb. paracasei* and *Lb. delbrueckii* sp. *bulgaricus* are able to produce GABA that

originated from cheese, pickles, yoghurt, fermented soybean etc. Recently, a great number of

Lactic acid bacteria strains have been isolated from various fermented foods and utilized for

the enrichment of functionality of foods by supplementation of GABA [85]. In microbes,

biosynthesis of GABA is carried out by pyridoxal-5'-phosphate-dependent-glutamic acid

decarboxylase (GAD) system which is comprised of the GAD enzyme (encoded by *gadA* or

gadB) and glutamate/GABA antiporter (*gadC*) [92-94]. In this study, the existence of these

genes and GABA production ability from MSG were confirmed via "*in silico*" and "*in vitro*"

analyses and the members of GAD system were summarized in Table S15. The NMR spectra

were shown in FigS4 describes that the AGA52 can convert the MSG to GABA by using its

related enzyme system. It is clearly seen in FigS4 that G1 (CH_2: 1.76 ppm), G2 (CH_2: 2.16

ppm), and G3 (CH_2: 2.87 ppm) peaks of MRS broth are showing highest similartiy with the

G1(CH_2: 1.75 ppm), G2(CH_2:2.15 ppm), and G3 (CH_2:2.85 ppm) peaks of GABA standard

which is the evidence of the AGA52 is able to produce GABA from MSG.

Conclusion

585   Briefly, it has been determined that *Lb. fermentum* AGA52 is a novel strain that possesses

586   crucial probiotic and psychobiotic features such as higher antioxidant activiy, cholesterol

587   assimilation capacity, GABA production ability, auto-aggregation ability, cell surface

588   hydrophobicity, gut persistance, adaptation to gastointestinal simulated environment, acid-bile

tolerance. It has obligate heterofermentative carbohydrate metabolism due to the absence of the phosphofructokinase enzyme which is confirmed from its genome. The AGA52 can also produce CO_2, lactate, and ethanol by heterofermentative pathway. Three phage regions (2 intact and 1 questionable) were identified and were detected as temperate due to the absence of the endolysins and 0.79% of the genome of the AGA52 has orginated from phage related horizontal gene transfer. On the other side, all detected resistome were found as intrinsically originated. The detected CRISPR structures revealed that the AGA52 can be used as model organism for CRISPR studies or can be used for development of next-generation probiotics by acquiring novel functional gene clusters. Besides, the survival behaviour of the strain proved that it has an adaptable nature to gastrointestinal mimicked environment and the diversity of the CAZYmes has also confirmed this case. When desired, survivability of this strain to gastrointestinal conditions can be enhanced more by encapsulation or gaining high copy number of relavant genes (e.g. *bsh*, Na+/H+ antiporters, F_0F_1 ATPases). The AGA52 has displayed the highest antioxidant capacity according to DPPH and ABTS+ assays which is also validated by presence of relevant genes. Overall, all findings revealed that the AGA52 has promising bacterial strain which has a potential for use as dietary supplement that might provide functional and therapeutic benefits to the host.

Acknowledgements

Mr. Mehmet Horzum would like to thank the Council of Higher Education of Türkiye (YÖK) due to the 100/2000 PhD scholarship program.

Funding

This study has been financially supported by Erciyes University Scientific Research Projects Coordination Unit under grant number FDK-2017-7281.

Declarations

Author Contribution

All authors contributed to the study's conception and design. Material preparation, data collection and analysis were performed by Ahmet E. Yetiman and Mehmet Horzum. The first draft of the manuscript was written by Ahmet E. Yetiman and was proofread by Mikail Akbulut. All authors read and approved the final manuscript.

Conflict of Interest

The authors declare no competing interests.

References

1. Seddik HA, Bendali F, Gancel F, Fliss I, Spano G, Drider D. Lactobacillus plantarum and Its Probiotic and Food Potentialities. Probiotics Antimicrob Proteins. 2017;9(2):111-22. doi: 10.1007/s12602-017-9264-z.

2. Buron-Moles G, Chailyan A, Dolejs I, Forster J, Miks MH. Uncovering carbohydrate metabolism through a genotype-phenotype association study of 56 lactic acid bacteria genomes. Appl Microbiol Biotechnol. 2019;103(7):3135-52. doi: 10.1007/s00253-019-09701-6.

3. Bangar SP, Suri S, Trif M, Ozogul F. Organic acids production from lactic acid bacteria: A preservation approach. Food Bioscience. 2022;46:101615.

4. Tamang JP, Shin D-H, Jung S-J, Chae S-W. Functional properties of microorganisms in fermented foods. Frontiers in microbiology. 2016;7:578.

5. Jiménez E, Langa S, Martín V, Arroyo R, Martín R, Fernández L, et al. Complete genome sequence of Lactobacillus fermentum CECT 5716, a probiotic strain isolated from human milk. Journal of bacteriology. 2010;192(18):4800-.

6. Erol I, Kotil SE, Fidan O, Yetiman AE, Durdagi S, Ortakci F. In silico analysis of bacteriocins from lactic acid bacteria against SARS-CoV-2. Probiotics and antimicrobial proteins. 2021:1-13.

7. Sanders ME, Akkermans LM, Haller D, Hammerman C, Heimbach JT, Hörmannsperger G, et al. Safety assessment of probiotics for human use. Gut microbes. 2010;1(3):164-85.

8. Zheng J, Wittouck S, Salvetti E, Franz CM, Harris H, Mattarelli P, et al. A taxonomic note on the genus Lactobacillus: Description of 23 novel genera, emended description of the genus Lactobacillus Beijerinck 1901, and union of Lactobacillaceae and Leuconostocaceae. International Journal of Systematic and Evolutionary Microbiology. 2020;70(4):2782-858.

9. D'ambrosio S, Ventrone M, Fusco A, Casillo A, Dabous A, Cammarota M, et al. Limosilactobacillus fermentum from buffalo milk is suitable for potential biotechnological process development and inhibits Helicobacter pylori in a gastric epithelial cell model. Biotechnology Reports. 2022;34:e00732.

10. de Araújo Henriques Ferreira G, Magnani M, Cabral L, Brandão LR, Noronha MF, de Campos Cruz J, et al. Potentially probiotic limosilactobacillus fermentum Fruit-Derived strains alleviate cardiometabolic disorders and gut microbiota impairment in male rats fed a high-Fat diet. Probiotics and Antimicrobial Proteins. 2022;14(2):349-59.

11. de Luna Freire MO, do Nascimento LCP, de Oliveira KÁR, de Oliveira AM, dos Santos Lima M, Napoleão TH, et al. Limosilactobacillus fermentum strains with claimed probiotic properties exert anti-oxidant and anti-inflammatory properties and prevent cardiometabolic disorder in female rats fed a high-fat diet. Probiotics and Antimicrobial Proteins. 2021:1-13.

12. Bhathena J, Martoni C, Kulamarva A, Tomaro-Duchesneau C, Malhotra M, Paul A, et al. Oral
probiotic microcapsule formulation ameliorates non-alcoholic fatty liver disease in Bio F1B Golden
Syrian hamsters. PLoS One. 2013;8(3):e58394.
13. Yamamoto N, Shoji M, Hoshigami H, Watanabe K, Takatsuzu T, Yasuda S, et al. Antioxidant
capacity of soymilk yogurt and exopolysaccharides produced by lactic acid bacteria. Bioscience of
Microbiota, Food and Health. 2019:18-017.
14. Yetiman AE, Keskin A, Darendeli BN, Kotil SE, Ortakci F, Dogan M. Characterization of
genomic, physiological, and probiotic features Lactiplantibacillus plantarum DY46 strain isolated
from traditional lactic acid fermented shalgam beverage. Food Bioscience. 2022;46:101499.
15. Paulino do Nascimento LC, Lacerda DC, Ferreira DJS, de Souza EL, de Brito Alves JL.
Limosilactobacillus fermentum, Current Evidence on the Antioxidant Properties and Opportunities to
be Exploited as a Probiotic Microorganism. Probiotics and Antimicrobial Proteins. 2022:1-20.
16. Feng H, Yuan Y, Yang Z, Xing X-h, Zhang C. Genome-wide genotype-phenotype associations in
microbes. Journal of bioscience and bioengineering. 2021;132(1):1-8.
17. Coskun F. A traditional Turkish fermented non-alcoholic beverage,"Shalgam". Beverages.
2017;3(4):49.
18. Ekinci FY, Baser GM, Özcan E, Üstündağ ÖG, Korachi M, Sofu A, et al. Characterization of
chemical, biological, and antiproliferative properties of fermented black carrot juice, shalgam.
European Food Research and Technology. 2016;242(8):1355-68.
19. Tanguler H, Erten H. Occurrence and growth of lactic acid bacteria species during the
fermentation of shalgam (salgam), a traditional Turkish fermented beverage. LWT-Food Science and
Technology. 2012;46(1):36-41.
20. Agirman B, Settanni L, Erten H. Effect of different mineral salt mixtures and dough extraction
procedure on the physical, chemical and microbiological composition of Şalgam: A black carrot
fermented beverage. Food Chemistry. 2021;344:128618.
21. Tanguler H, Erten H. Chemical and microbiological characteristics of shalgam (şalgam): a
traditional turkish lactic acid fermented beverage. Journal of Food Quality. 2012;35(4):298-306.
22. Prjibelski A, Antipov D, Meleshko D, Lapidus A, Korobeynikov A. Using SPAdes de novo
assembler. Current protocols in bioinformatics. 2020;70(1):e102.
23. Tatusova T, DiCuccio M, Badretdin A, Chetvernin V, Nawrocki EP, Zaslavsky L, et al. NCBI
prokaryotic genome annotation pipeline. Nucleic acids research. 2016;44(14):6614-24.
24. Wattam AR, Davis JJ, Assaf R, Boisvert S, Brettin T, Bun C, et al. Improvements to PATRIC, the
all-bacterial bioinformatics database and analysis resource center. Nucleic acids research.
2017;45(D1):D535-D42.
25. Brettin T, Davis JJ, Disz T, Edwards RA, Gerdes S, Olsen GJ, et al. RASTtk: a modular and
extensible implementation of the RAST algorithm for building custom annotation pipelines and
annotating batches of genomes. Scientific reports. 2015;5(1):1-6.
26. Alikhan NF, Petty NK, Ben Zakour NL, Beatson SA. BLAST Ring Image Generator (BRIG):
simple prokaryote genome comparisons. BMC Genomics. 2011;12:402. doi: 10.1186/1471-2164-12-
402.
27. Lee I, Kim YO, Park S-C, Chun J. OrthoANI: an improved algorithm and software for calculating
average nucleotide identity. International journal of systematic and evolutionary microbiology.
2016;66(2):1100-3.
28. Chen I-MA, Chu K, Palaniappan K, Ratner A, Huang J, Huntemann M, et al. The IMG/M data
management and analysis system v. 6.0: new tools and advanced capabilities. Nucleic acids research.
2021;49(D1):D751-D63.
29. Kanehisa M, Sato Y, Morishima K. BlastKOALA and GhostKOALA: KEGG tools for functional
characterization of genome and metagenome sequences. Journal of molecular biology.
2016;428(4):726-31.
30. Zhang H, Yohe T, Huang L, Entwistle S, Wu P, Yang Z, et al. dbCAN2: a meta server for
automated carbohydrate-active enzyme annotation. Nucleic acids research. 2018;46(W1):W95-W101.
31. Wu S, Zhu Z, Fu L, Niu B, Li W. WebMGA: a customizable web server for fast metagenomic
sequence analysis. BMC genomics. 2011;12(1):1-9.
32. Khan A, Mathelier A. Intervene: a tool for intersection and visualization of multiple gene or
genomic region sets. BMC bioinformatics. 2017;18(1):1-8.

33. Arndt D, Grant JR, Marcu A, Sajed T, Pon A, Liang Y, et al. PHASTER: a better, faster version of
the PHAST phage search tool. Nucleic acids research. 2016;44(W1):W16-W21.
34. Bortolaia V, Kaas RS, Ruppe E, Roberts MC, Schwarz S, Cattoir V, et al. ResFinder 4.0 for
predictions of phenotypes from genotypes. Journal of Antimicrobial Chemotherapy.
2020;75(12):3491-500.
35. Alcock BP, Raphenya AR, Lau TT, Tsang KK, Bouchard M, Edalatmand A, et al. CARD 2020:
antibiotic resistome surveillance with the comprehensive antibiotic resistance database. Nucleic acids
research. 2020;48(D1):D517-D25.
36. Khan ZA, Siddiqui MF, Park S. Current and emerging methods of antibiotic susceptibility testing.
Diagnostics. 2019;9(2):49.
37. CLSI C. Performance Standards for Antimicrobial Susceptibility Testing; Twenty-Second
Informational Supplement. 2012.
38. Angmo K, Kumari A, Bhalla TC. Probiotic characterization of lactic acid bacteria isolated from
fermented foods and beverage of Ladakh. LWT-food Science and Technology. 2016;66:428-35.
39. Zhang L, Ma H, Kulyar MF-e-A, Pan H, Li K, Li A, et al. Complete genome analysis of
Lactobacillus fermentum YLF016 and its probiotic characteristics. Microbial Pathogenesis.
2022;162:105212.
40. Krausova G, Hyrslova I, Hynstova I. In vitro evaluation of adhesion capacity, hydrophobicity, and
auto-aggregation of newly isolated potential probiotic strains. Fermentation. 2019;5(4):100.
41. Mishra V, Prasad D. Application of in vitro methods for selection of Lactobacillus casei strains as
potential probiotics. International journal of food microbiology. 2005;103(1):109-15.
42. Ozturk G, Yetiman AE, Dogan M. The bioactive efficiency of ultrasonic extracts from acorn
leaves and green walnut husks against Bacillus cereus: a hybrid approach to PCA with the Taguchi
method. Journal of Food Measurement and Characterization. 2019;13(2):1257-68.
43. Pieniz S, Andreazza R, Anghinoni T, Camargo F, Brandelli A. Probiotic potential, antimicrobial
and antioxidant activities of Enterococcus durans strain LAB18s. Food Control. 2014;37:251-6.
44. Rudel LL, Morris M. Determination of cholesterol using o-phthalaldehyde. Journal of Lipid
Research. 1973;14(3):364-6.
45. Brandt K, Nethery MA, O'Flaherty S, Barrangou R. Genomic characterization of Lactobacillus
fermentum DSM 20052. BMC genomics. 2020;21(1):1-13.
46. Juhas M, Van Der Meer JR, Gaillard M, Harding RM, Hood DW, Crook DW. Genomic islands:
tools of bacterial horizontal gene transfer and evolution. FEMS microbiology reviews.
2009;33(2):376-93.
47. Yoo D, Bagon BB, Valeriano VDV, Oh JK, Kim H, Cho S, et al. Complete genome analysis of
Lactobacillus fermentum SK152 from kimchi reveals genes associated with its antimicrobial activity.
FEMS Microbiology Letters. 2017;364(18).
48. El-Ghaish S, Dalgalarrondo M, Choiset Y, Sitohy M, Ivanova I, Haertlé T, et al. Characterization
of a new isolate of Lactobacillus fermentum IFO 3956 from Egyptian Ras cheese with proteolytic
activity. European food research and technology. 2010;230(4):635-43.
49. Duar RM, Lin XB, Zheng J, Martino ME, Grenier T, Pérez-Muñoz ME, et al. Lifestyles in
transition: evolution and natural history of the genus Lactobacillus. FEMS microbiology reviews.
2017;41(Supp_1):S27-S48.
50. Cox AJ, Pyne DB, Saunders PU, Fricker PA. Oral administration of the probiotic Lactobacillus
fermentum VRI-003 and mucosal immunity in endurance athletes. British Journal of Sports Medicine.
2010;44(4):222-6.
51. Goh YJ, Barrangou R. Harnessing CRISPR-Cas systems for precision engineering of designer
probiotic lactobacilli. Current opinion in biotechnology. 2019;56:163-71.
52. Roberts A, Barrangou R. Applications of CRISPR-Cas systems in lactic acid bacteria. FEMS
Microbiology Reviews. 2020;44(5):523-37.
53. Hidalgo-Cantabrana C, O'Flaherty S, Barrangou R. CRISPR-based engineering of next-generation
lactic acid bacteria. Current opinion in microbiology. 2017;37:79-87.
54. Mahony J, Van Sinderen D. Current taxonomy of phages infecting lactic acid bacteria. Frontiers in
microbiology. 2014;5:7.
55. Oechslin F, Zhu X, Dion MB, Shi R, Moineau S. Phage endolysins are adapted to specific hosts
and are evolutionarily dynamic. PLoS biology. 2022;20(8):e3001740.

56. Miller-Ensminger T, Mormando R, Maskeri L, Shapiro JW, Wolfe AJ, Putonti C. Introducing Lu-
1, a novel lactobacillus jensenii phage abundant in the urogenital tract. Plos one.
2020;15(6):e0234159.
57. Cahill J, Young R. Phage lysis: multiple genes for multiple barriers. Advances in virus research.
2019;103:33-70.
58. Kleerebezem M, Boekhorst J, van Kranenburg R, Molenaar D, Kuipers OP, Leer R, et al.
Complete genome sequence of Lactobacillus plantarum WCFS1. Proceedings of the National
Academy of Sciences. 2003;100(4):1990-5.
59. EFSA. Guidance on the assessment of bacterial susceptibility to antimicrobials of human and
veterinary importance. EFSA Journal. 2012;10(6):2740.
60. Deghorain M, Goffin P, Fontaine L, Mainardi J-L, Daniel R, Errington J, et al. Selectivity for D-
lactate incorporation into the peptidoglycan precursors of Lactobacillus plantarum: role of Aad, a
VanX-like D-alanyl-D-alanine dipeptidase. Journal of bacteriology. 2007;189(11):4332-7.
61. Campedelli I, Mathur H, Salvetti E, Clarke S, Rea MC, Torriani S, et al. Genus-wide assessment
of antibiotic resistance in Lactobacillus spp. Applied and environmental microbiology.
2019;85(1):e01738-18.
62. Zhang Z-Y, Liu C, Zhu Y-Z, Wei Y-X, Tian F, Zhao G-P, et al. Safety assessment of Lactobacillus
plantarum JDM1 based on the complete genome. International Journal of Food Microbiology.
2012;153(1-2):166-70.
63. Rozman V, Lorbeg PM, Accetto T, Matijašić BB. Characterization of antimicrobial resistance in
lactobacilli and bifidobacteria used as probiotics or starter cultures based on integration of phenotypic
and in silico data. International journal of food microbiology. 2020;314:108388.
64. Walter J, Heng NC, Hammes WP, Loach DM, Tannock GW, Hertel C. Identification of
Lactobacillus reuteri genes specifically induced in the mouse gastrointestinal tract. Applied and
environmental microbiology. 2003;69(4):2044-51.
65. Gänzle MG, Follador R. Metabolism of oligosaccharides and starch in lactobacilli: a review.
Frontiers in microbiology. 2012;3:340.
66. Salvetti E, Harris HM, Felis GE, O'Toole PW. Comparative genomics of the genus Lactobacillus
reveals robust phylogroups that provide the basis for reclassification. Applied and Environmental
Microbiology. 2018;84(17):e00993-18.
67. Morita H, Toh H, Fukuda S, Horikawa H, Oshima K, Suzuki T, et al. Comparative genome
analysis of Lactobacillus reuteri and Lactobacillus fermentum reveal a genomic island for reuterin and
cobalamin production. DNA research. 2008;15(3):151-61.
68. Schleif R. AraC protein, regulation of the l-arabinose operon in Escherichia coli, and the light
switch mechanism of AraC action. FEMS microbiology reviews. 2010;34(5):779-96.
69. Jiang T, Guo X, Yan J, Zhang Y, Wang Y, Zhang M, et al. A bacterial multidomain NAD-
independent d-lactate dehydrogenase utilizes flavin adenine dinucleotide and Fe-S clusters as cofactors
and quinone as an electron acceptor for d-lactate oxidization. Journal of bacteriology.
2017;199(22):e00342-17.
70. Thomas LM, Harper AR, Miner WA, Ajufo HO, Branscum KM, Kao L, et al. Structure of
Escherichia coli AdhP (ethanol-inducible dehydrogenase) with bound NAD. Acta Crystallographica
Section F: Structural Biology and Crystallization Communications. 2013;69(7):730-2.
71. Goel A, Halami PM, Tamang JP. Genome analysis of Lactobacillus plantarum isolated from some
Indian fermented foods for bacteriocin production and probiotic marker genes. Frontiers in
microbiology. 2020;40.
72. Martienssen M, O R, U K. Surface properties of bacteria from different wastewater treatment
plants. Acta biotechnologica. 2001;21(3):207-25.
73. de Souza BMS, Borgonovi TF, Casarotti SN, Todorov SD, Penna ALB. Lactobacillus casei and
Lactobacillus fermentum strains isolated from mozzarella cheese: probiotic potential, safety,
acidifying kinetic parameters and viability under gastrointestinal tract conditions. Probiotics and
antimicrobial proteins. 2019;11(2):382-96.
74. Chaffanel F, Charron-Bourgoin F, Soligot C, Kebouchi M, Bertin S, Payot S, et al. Surface
proteins involved in the adhesion of Streptococcus salivarius to human intestinal epithelial cells.
Applied microbiology and biotechnology. 2018;102(6):2851-65.

75. Pan M, Kumaree KK, Shah NP. Physiological changes of surface membrane in Lactobacillus with
prebiotics. Journal of food science. 2017;82(3):744-50.
76. Haddaji N, Mahdhi AK, Krifi B, Ismail MB, Bakhrouf A. Change in cell surface properties of
Lactobacillus casei under heat shock treatment. FEMS microbiology letters. 2015;362(9).
77. Lesuffleur T, Barbat A, Dussaulx E, Zweibaum A. Growth adaptation to methotrexate of HT-29
human colon carcinoma cells is associated with their ability to differentiate into columnar absorptive
and mucus-secreting cells. Cancer research. 1990;50(19):6334-43.
78. Li Q, Liu X, Dong M, Zhou J, Wang Y. Aggregation and adhesion abilities of 18 lactic acid
bacteria strains isolated from traditional fermented food. Int J Agric Policy Res. 2015;3(2):84-92.
79. Ouwehand AC, Kirjavainen PV, Shortt C, Salminen S. Probiotics: mechanisms and established
effects. International dairy journal. 1999;9(1):43-52.
80. Saarela M, Mogensen G, Fonden R, Mättö J, Mattila-Sandholm T. Probiotic bacteria: safety,
functional and technological properties. Journal of biotechnology. 2000;84(3):197-215.
81. Gao Y, Liu Y, Sun M, Zhang H, Mu G, Tuo Y. Physiological function analysis of Lactobacillus
plantarum Y44 based on genotypic and phenotypic characteristics. Journal of dairy science.
2020;103(7):5916-30.
82. Fei Y, Li L, Zheng Y, Liu D, Zhou Q, Fu L. Characterization of Lactobacillus amylolyticus L6 as
potential probiotics based on genome sequence and corresponding phenotypes. LWT. 2018;90:460-8.
83. Tomaro-Duchesneau C, Jones ML, Shah D, Jain P, Saha S, Prakash S. Cholesterol assimilation by
Lactobacillus probiotic bacteria: an in vitro investigation. BioMed research international. 2014;2014.
84. Pan DD, Zeng XQ, Yan YT. Characterisation of Lactobacillus fermentum SM-7 isolated from
koumiss, a potential probiotic bacterium with cholesterol-lowering effects. Journal of the Science of
Food and Agriculture. 2011;91(3):512-8.
85. Cui Y, Miao K, Niyaphorn S, Qu X. Production of gamma-aminobutyric acid from lactic acid
bacteria: A systematic review. International Journal of Molecular Sciences. 2020;21(3):995.
86. Wu J-Y, Matsuda T, Roberts E. Purification and characterization of glutamate decarboxylase from
mouse brain. Journal of Biological Chemistry. 1973;248(9):3029-34.
87. Okada T, Sugishita T, Murakami T, Murai H, Saikusa T, Horino T, et al. Effect of the defatted rice
germ enriched with GABA for sleeplessness, depression, autonomic disorder by oral administration.
Nippon Shokuhin Kagaku Kogaku Kaishi= Journal of the Japanese Society for Food Science and
Technology. 2000;47(8):596-603.
88. Abdou AM, Higashiguchi S, Horie K, Kim M, Hatta H, Yokogoshi H. Relaxation and immunity
enhancement effects of γ-aminobutyric acid (GABA) administration in humans. Biofactors.
2006;26(3):201-8.
89. Ting Wong CG, Bottiglieri T, Snead III OC. Gaba, γ-hydroxybutyric acid, and neurological
disease. Annals of Neurology: Official Journal of the American Neurological Association and the
Child Neurology Society. 2003;54(S6):S3-S12.
90. Hayakawa K, Kimura M, Kasaha K, Matsumoto K, Sansawa H, Yamori Y. Effect of a γ-
aminobutyric acid-enriched dairy product on the blood pressure of spontaneously hypertensive and
normotensive Wistar–Kyoto rats. British Journal of Nutrition. 2004;92(3):411-7.
91. Leventhal AG, Wang Y, Pu M, Zhou Y, Ma Y. GABA and its agonists improved visual cortical
function in senescent monkeys. Science. 2003;300(5620):812-5.
92. Lyu C, Zhao W, Peng C, Hu S, Fang H, Hua Y, et al. Exploring the contributions of two glutamate
decarboxylase isozymes in Lactobacillus brevis to acid resistance and γ-aminobutyric acid production.
Microbial cell factories. 2018;17(1):1-14.
93. Wu Q, Tun HM, Law Y-S, Khafipour E, Shah NP. Common distribution of gad operon in
Lactobacillus brevis and its GadA contributes to efficient GABA synthesis toward cytosolic near-
neutral pH. Frontiers in Microbiology. 2017;8:206.
94. Yunes R, Poluektova E, Dyachkova M, Klimina K, Kovtun A, Averina O, et al. GABA production
and structure of gadB/gadC genes in Lactobacillus and Bifidobacterium strains from human
microbiota. Anaerobe. 2016;42:197-204.

Fig 1. The BLAST Ring alignment of *Limosilactobacillus fermentum* strain AGA52 versus other compared well-known *Lb. fermentum* genomes which were aligned from the red ring to the inside as CECT 5716, IFO3956, B44, YLF016, DSM 20052, ATCC 14931, SK152, and AGA52, respectively. The GC content was illustrated in the third inner circle before the genome of AGA52. The GC skew (+/-) of the genome of AGA52 was also shown in the 2nd inner circle and in the first inner circle, the genome size has been demonstrated. On the other hand, prophage regions of the strain AGA52 were depicted as navy blue colored arcs in the outer circle, the numbers in parentheses show the number of matched proteins of the phages. Besides, at the end of the phage names, (IN) means intact and (Q) means questionable.

Fig 2. Orthologous Average Nucleotide Identity (OrthoANI) phylogram of the genomes of the *Limosilactobacillus fermentum* strain AGA52 and other well-known *Limosilactobacillus* strains.

Fig 3. Graphical presentation of the identified numbers of clusters of orthologous groups (COG) of protein-coding sequences in the genome of *Limosilactobacillus fermentum* AGA52.

Fig 4. The UpSet plot demonstrates the number of shared orthologous proteins amongst AGA52 and the other well-known strains with bar charts. On the other hand, the number of total identified clusters of orthologous groups (COG) via the webMGA engine has been shown on the left of the strain names.

Fig 5. Pairwise intersection with hierarchical clustering in accordance with the existence of clusters of orthologous groups (COG) of 48 different *Lb. fermentum*, *Lb. reuteri* DSM 20016, and *Lb. gastricus* DSM 16045 genomes which were available on IMG/M. The correlation coefficient was calculated according to Pearson correlation and the color bar on the bottom of the heatmap indicates the degree of the correlation coefficient. The AGA52 shows a higher correlation with KMB612 (0.95), KMB613 (0.95), AF15-40LB (0.95), AF16-22LB (0.95), 279 (0.94), 779_LFER (0.94), and VRI-003 (0.94), respectively. Interestingly, it is apprently seen that MTCC 8711 inversely correlated with the all genomes clustered.

Fig 6. Heatmap of carbohydrate-active enzymes (CAZYmes) and their distribution amongst the compared *L. fermentum* strains. The color gradient bar indicates the exact number of identified enzymes from the CAZY (http://www.cazy.org/) database by using DIAMOND (E-value <1e-102). CE: Carbohydrate Esterases, AA: Auxiliary Activities, CBM: Carbohydrate-Binding Modules, GT: Glycosyl Transferases, GH: Glycoside Hydrolases.

Fig 7. Survival rate and growth characteristics under different conditions. A. Survival at
different bile salts concentrations, B. Growth at different pH values, C. Survival tendency of
AGA52 at simulated gastric juice environment, D. Survival behaviour of AGA52 at simulated
intestinal juice environment.

985 **Table 1.** Summary of assembly details and annotated genome properties of
986 *Limosilactobacillus fermentum* AGA52 and other compared strains which have higher
987 orthoANI similarity to itself.

Item	AGA52	SK152	ATCC 14931 (B1 28)	YLF016
Contigs	128	1	1	1
Genome length (bp)	2,001,084	2,092,273	1,905,587	2,094,354
GC content (%)	51,82	51,94	52,3	51,46
Sequencing platform	Illumina Novaseq	PacBio	Oxford Nanopore GridION	PacBio
Sequencing depth	296.0x	500.0x	342x	200.0x
Genes (total)	2,024	2,121	1,916	2,136
Protein coding sequences	1,873	1,950	1,744	1,950
tRNA	50	58	58	58
rRNA	3	15	15	15
Non-coding RNA	3	3	3	3
CRISPR_repeat	15	100	13	10
CRISPR_spacer	14	98	12	9
CRISPR_array	1	2	1	1
Pseudogenes (total)	95	95	96	110
Pseudogeens (frameshifted)	19 of 95	28 of 95	36 of 96	35 of 110
Pseudogenes (incomplete)	68 of 95	50 of 95	35 of 96	60 of 110
Pseudogenes (internal stop)	24 of 95	29 of 95	33 of 96	30 of 110
OrthoANI similarity to AGA52	-	99.18	99.15	99.11

989