1 **Seedless fruit in *Annona squamosa* L. is monogenic and conferred by *INO***
2 **locus deletion in multiple accessions**
3

4 Bruno Rafael Alves Rodrigues[1], Charles S. Gasser[2], Samy Pimenta[1], Marlon Cristian Toledo
5 Pereira[1] and Silvia Nietsche[3]*
6
[1] State University of Montes Claros, Department of Agricultural Sciences, CEP 39401-369,
Janaúba, MG, Brazil
[2] Dept. of Molecular and Cellular Biology, University of California, Davis, Davis, CA 95616
USA
[3] Federal University of Minas Gerais, Institute of Agricultural Science, CEP 39404-547,
Montes Claros, Minas Gerais, Brazil
*Author for correspondence: silvia.nietsche@gmail.com

14
15
16 **Running head: Inheritance of seedless fruit in *A. squamosa***
17

Summary. Understanding the genetic basis and inheritance of a trait facilitates the planning
of breeding and development programs of new cultivars. In the sugar apple tree (*Annona*
squamosa L.), the mechanism of the desirable seedless trait in the Thai seedless (Ts) and
Brazilian seedless (Bs) accessions was associated with a deletion of the *INNER NO OUTER*
(*INO*) locus. Genetic analysis of F_1, F_2 and backcross descendants of crosses of Bs to fertile
wild-type varieties showed that seedlessness was recessive and monogenic. Whole genome
sequencing of a third accession, Hawaiian seedless (Hs), identified a 16 kilobase deletion
including *INO*. The finding of an identical deletion in Ts and Bs indicated a common origin
among genotypes, from a single deletion event. Analysis of microsatellite markers could not
preclude the possibility that all three accessions are vegetatively propagated clones. The
sequence of the deletion site enabled formulation of a codominant assay for the wild-type and
mutant genes that validated the *INO* gene deletion as the cause of seedless trait, and can be
used in the selection of new seedless varieties. The study findings and obtained progenies
should be useful in breeding and introgression programs of the trait into elite sugar apple lines
and into other *Annonas* by means of interspecific crossings.

Introduction

The global fruit market has grown appreciably in the past decade. The absence of seeds from fruit that is consumed has become one of the most appreciated traits by consumers. Reducing seed content without changing the size of the fruit is one of the main objectives of the cultivation of many fruit trees. The goal is to enhance the experience of fruit consumption by consumers and improve the quality of fruits for food processing (Varoquaux et al., 2000).

Sugar apple (*Annona squamosa* L.) ($2n = 14$) is a tropical fruit species in the family Annonaceae. Seedless fruits have been described for some spontaneous mutants of *A. squamosa* whose origin is still unknown (Pinto et al., 2005; Wongs-Aree and Noichinda, 2011; Lora et al., 2018; Pereira and Borém, 2021). They include the Cuban cultivar Cuban seedless (Cs), with good fruit characteristics but lower productivity than fertile cultivars (Araújo et al., 1999); Brazilian seedless (Bs) originally identified in northeast Brazil, which produces small, asymmetric fruits that frequently perish (Santos, 2014); the Thai seedless (Ts) mutant that produces normal size fruits (Lora et al. 2011) among other types with apparently similar fruits, such as in the Philippines and Hawaii (Pinto et al., 2005; this work).

Lora et al. (2011) were among the first researchers to examine details of the seedless trait in *A. squamosa* in Ts. This variety produces fruit following pollination and fertilization. The authors demonstrated that seedlessness results from a defect in ovule development where the outer of the two integuments sheathing the nucellus fails to form. This defect directly mirrors the effect of *INNER NO OUTER* (*INO*) loss of function mutants in *Arabidopsis thaliana* (Villanueva et al., 1999). *INO* encodes a putative transcription factor belonging to the *YABBY* family. Members of this family are involved in the determination of abaxial identity in a variety of plant organs (Sawa et al., 1999; Siegfried et al., 1999; Bowman, 2000). Lora et al. (2011) isolated an *A. squamosa INO* ortholog and demonstrated the association of

the Ts mutant with an apparent deletion of the *INO* gene, indicating a candidate gene for the

seedless trait.

The Bs variety was also evaluated in a breeding program undertaken at the State

University of Montes Claros. Results showed that the absence of seeds was also associated

with failure in the development of the external integument of the ovule and in the

development of seeds, similar to that observed in Ts (Mendes et al., 2012; Santos et al., 2014).

In addition, preliminary studies of inheritance of the stenospermocarpic absence of seeds

involving F_1 progenies in Bs (*A. squamosa* wild type x Bs) also indicated the same expected

recessive nature for the mutation as in Ts (Nassau et al., 2021). However, it was not known

whether the molecular basis of the *INO* deletion described is widespread in *A. squamosa* and

may be responsible for the case of aspermia described in Bs.

The present study addressed the following questions: (i) Is the inheritance of the

presence/absence of seeds in *A. squamosa* monogenic?; (ii) What are the molecular details of

the *INO* gene deletion?; (iii) Is there a complete correspondence of homozygosity for the

deletion with the seedless phenotype; (iv) What is the relationship between the known

seedless accessions, and (v) Can codominant molecular markers specific to *INO* be designed

for use in assisted selection?

95
96
97   **Materials and Methods**
98
99
100   **Plant material**
101
Wild-types (fertile) M_1, M_2 and M_3 and seedless Bs lines were previously described by

de Souza et al. (2010), and Ts was described by Lora et al. (2011). For whole genome

sequencing a wild-type sugar apple fruit was purchased from a retail source in the United

States, seeds from the fruit were planted, and one plant grown in the UC Davis Conservatory

was sampled for sequencing with voucher herbarium samples stored as DAV225058 and

DAV225059. The Hawaiian seedless (Hs) line was obtained as budwood from Frankie's

Nursery (Waimanalo, HI), grafted to a wild-type *A. squamosa* rootstock and grown in the UC

Davis Conservatory with a voucher sample stored as DAV225060.

For genetic inheritance studies, plants (M_1, M_2, M_3 and Bs) were grown at the

Experimental Farm and molecular analysis were performed at Molecular Biology laboratory

of the State University of Montes Claros, latitude 15º48'09''S, longitude 43°18'32''W and

altitude 516 m. For phenotypic characterization of seedless versus seeded (fertile) two

strategies were applied: (a) fruits were harvested, pulped, and examined for the presence or

absence of seeds (Supplemental Figure S1); or (b) flowers either fresh or fixed in formalin

acetic acid-alcohol (FAA) were dissected to separate the ovules from the carpel tissue.

Isolated ovules were observed with a stereomicroscope (Nikon, SMZ800) and photographed

by a digital camera (Sony, Dsc-W350). The wild-type ovules present a domed shape opposite

the funiculs, while the mutant ovules come to a point at this position (Supplemental Figure

S2).

Filial generations (F_1), self-fertilization (F_2), backcrosses with wild-types parents M_1,

M_2, M_3 (BC_M), and backcrosses with mutant parent Bs (BC_{Bs}) were obtained. Segregations

were evaluated for conformity to predicted ratios with the Chi-square test (χ^2) using the Genes

statistical software (Cruz, 2016).

125
126
127   **Polymerase chain reaction (PCR) analysis**
128
DNA was extracted from young leaf samples with hexadecyltrimethylammonium

bromide buffer (CTAB) as described by Doyle and Doyle (1990) and separated from

polysaccharides as described by Cheung et al., (1993). Primers used in PCR are listed in

Supplemental Table S1. PCR was performed with DreamTaq (ThermoFisher, Waltham, MA)

and the included reagents with an initial denaturation at 94ºC for 3 min; 35 cycles with

denaturation at 94ºC for 30 sec, annealing at 56°C for 30 sec, and extension at 72ºC for 1.5

min; and a final extension of 72ºC for 4 min. For reactions using the AsINODel primers a 60°

annealing temperature was used.

PCR products were electrophoresed on 1.2 % (w/v) agarose buffered with 1× TBE (89

mM Tris-base, 89 mM boric acid, 2 mM EDTA, pH 8.0) or SB (10 mM Sodium Borate pH

8.0) and DNA visualized by staining with ethidium bromide (2 µg/ml) an illumination with

ultraviolet light.

For sequencing, PCR products were processed with ExoSAP-IT (ThermoFisher,

Waltham, MA) or Quiapure (Quiagen, Germantown, MD) and sequenced using amplification

primers on an ABI 3500 or 3730 genetic analyzer (ThermoFisher, Waltham, MA) at Análises

Moleculares Ltda. (Centro de Biotecnologia, UFRGS, Porto Alegre, RS) or the University of

California Davis CBS DNA Sequencing Facility (Davis, CA).

146
147
148 **Whole genome shotgun sequencing**
149
DNA for whole genome sequencing was isolated from young leaves by grinding in

100 mM TRIS-Cl, 20 mM EDTA, 1.4 M NaCl, 2% (w/v) CTAB, 1% each

polyvinylpyrrolidone and sodium metabisulfite pH 8.0. Samples were treated with 70 µg/ml

RNAaseA (ThermoFisher, Waltham, MA), extracted with 1:24 (v/v) mixture of isoamyl

alcohol and chloroform, and precipitated with isopropanol. Samples were dissolved in 10 mM

TRIS pH 8.0, 1 mM EDTA, adjusted to 0.3 M Na Acetate, pH 4.8, precipitated with 2

volumes of ethanol and dissolved in 10 mM TRIS pH 8. Wild-type *A. squamosa* DNA was

processed and sequenced at the University of California, Davis Genome Center (Davis, CA).

For PacBio (Menlo Park, CA) sequencing, DNA fragments greater than 10 kb were selected

by BluePippin (Sage Sciences, Beverly, MA) electrophoresis and were sequenced on a

PacBio RSII or Sequel Single Molecule, Real-time device. This resulted in 2.46 million reads

with an average read length of 8 kb comprising more than 29 Gbases, or approximately 37 X

genome representation. For Illumina (San Diego, CA) sequencing the DNA was sheared and

fragments of an average size of 400 bp were selected and sequenced on a HiSeq 4000

apparatus by the paired-end 150 bp method (PE150) resulting in approximately 390 million

sequences. The sequences were trimmed of poor quality regions and primer sequences with

Sickle (Joshi and Foss, 2011) and Scythe (Buffalo, 2011), respectively resulting in 229

Gbases or approximately 124 X genome representation. Hs DNA was similarly processed and

sequenced by QuickBiology (Pasadena, CA) resulting in 408 million sequences and

approximately 130 X genome representation.

170
171
172 **Genome sequence assembly and analysis**
173
The PacBio reads of wild-type DNA were assembled using Canu (Koren et al., 2017)

with default settings, producing 3519 contigs. The wild-type Illumina reads were aligned with

the assembly using BWA MEM (Li, 2013) with default settings, and Pilon (Walker et al.,

2014) used the alignment to correct the contigs, changing 148k single nucleotide errors and

adding a net of more than 1.8 Mbases of insertions for a final assembly of 707.7 Mbases with

an average contig length of 201 kb and an N90 of 93.9 kb. A BLAST (Altschul et al., 1990)

search with the known *A. squamosa INO* gene sequence (Lora et al., 2011) identified a 587 kb

contig (tig00001115) containing the *INO* gene (Supplemental Figure S3). BWA MEM aligned

the Hs Illumina reads with the assembly and Tablet (Milne et al., 2013) was used to examine

the alignment with the 587 kb contig containing INO. BLAST was used to search one half of

the set of Hs Illumina sequence reads for those extending across a detected deletion and the

resulting sequences were aligned and assembled using Sequencer 5.4.1 (Gene Codes, Ann

Arbor, MI).

187
188

Genetic diversity using SSR markers

Genetic diversity was assessed among varieties of seedless sugar apple Bs, Ts, and Hs, with the fertile parent M_2 as a contrasting control. Sixty-seven pairs of SSR microsatellite markers, described for *A. cherimola* were used, with fifteen having been described by Escribano et al. (2004) and fifty-two by Escribano et al. (2008).

DNA extraction was performed as described above for the markers association of seedless trait with INO deletion. Amplification utilized an initial denaturation at 94ºC for 1 min; 35 cycles at 94ºC denaturation for 30 sec, annealing at 48 to 57°C depending on the primer; and extension at 72ºC for 1 min; and a final extension of 72ºC for 7 min. The amplification products were separated by 3.0% agarose gel electrophoresis buffered, stained and visualized as above.

To calculate diversity, the amplification data of the SSR primers were converted into numerical code per locus for each allele. The presence of a band was designated by 1 and the absence by 0. Although the microsatellite markers can be codominant, molecular analyses of the locus were performed based on the presence/absence of each amplified fragment. The established binary matrix was used to obtain estimates of genetic similarities between genotype pairs, based on the Jaccard coefficient. The Genes statistical program (Cruz, 2016) was used for data processing.

Procedures adopted chronologically

The present study was developed according the following chronological events: (1) obtaining the segregating generations (F_2, BC_M and BC_{Bs}) from the crossing of the fertile parents M_1, M_2 and M_3 with the mutant Bs (2018-2019); (2) application of the dominant marker LMINO primer-set in order to associate the seedless trait with *INO* deletion in the segregating generations (F_1, F_2, BC_M and BC_{Bs}) (2019-2020); (3) sequencing and

217 identification of the *INO* gene deletion region and design the codominant primer AsINODel

218 (2021); (4) AsINODel marker validation and phenotypic characterization of plants (F_2, BC_M

219 and BC_{Bs}) in the field conditions (2021-2022).

220
221
222 # Results

223
224
225 **Seedless trait inheritance**

226
The results of the phenotypic analysis of the parents M_2 and Bs, F_1, F_2, backcrosses

with the wild-type parent M_2 (BC_M) and with mutant parent Bs (BC_{Bs}) are displayed in Table

1. In generation F_1, all individuals presented fruits with seeds. In the F_2 population, among the

plants in reproductive stage during the evaluation period, 48 formed fruits with fully

developed seeds and ten presented only seed rudiments, characterized by the absence of seeds

(Supplemental Figure S1). Considering segregation hypotheses expected for one, two and

three genes (Supplemental Table S2), the Chi-square test revealed that the trait under study

segregated at a 3:1 ratio (presence/absence), consistent with a monogenic inheritance. These

results were corroborated by data from backcrosses with the parent M_2 (BC_M), where all

plants that produced fruits had seeds, consistent with the 1:0 ratio, while the plants evaluated

through backcrossing with the mutant parent Bs (BC_{Bs}), showed segregation of 1:1 for

presence and absence of seeds. Taken together, these results corroborate the monogenic

inheritance found in the analyses of F_2 generations, indicating that a single recessive locus

controls the seedless trait in Bs *A. squamosa*.

Table 1. Phenotypic analysis of the segregation of the seed presence/absence in fruits of F_1 and F_2 generations, BC_M and BC_{Bs} backcrosses of the M_2 family in *A. squamosa*. Janaúba-MG, Brazil.

Parents and Generations	Observed proportion		Expected ratio	χ^2	*p*-value
	Presence	Absence			
Parents					
M_2	1				
Bs		1			
Generation F_1					
$M_2 \times$ Bs	17	0	1:0		
Generation F_2					
$F_1(M_2 \times$ Bs$) \times F_1(M_2 \times$ Bs$)$	48	10	3:1	1.86	0.172[ns]
Backcrossing					
BC_M					
$F_1(M_2 \times$ Bs$) \times M_2$	12	0	1:0		
BC_{Bs}					
$F_1(M_2 \times$ Bs$) \times$ Bs	8	6	1:1	0.286	0.592[ns]

[ns] Non-significant values at the 5% significance level, χ^2 Chi-square test estimate.

Association of seedless trait with *INO* deletion

Previously described molecular markers for the presence of the *INO* gene (Lora et al., 2011) were tested on parents M_1, M_2, M_3, and Bs and displayed the expected band patterns. These markers generated amplification products only in the three wild-type parents, with no amplification of any fragment in Bs for any of the primer pairs used (Supplemental Figure S4A).

The dominant marker LMINO primer-set were also used to amplify DNA from F_1 plants obtained from crosses between genotypes of *A. squamosa* (M_1, M_2, and M_3) with the mutant Bs (Table 1). All evaluated F_1 individuals produced fruits with seeds in the field and amplified the products with all primer pairs, as shown in the Supplemental Figure S4B.

The same procedure was applied in order to genotype segregating generations (F_2, BC_M and BC_{Bs}) in seedling stage in the nursery (Table 2). Figure 1 shows a sample of individuals amplified with the LMINO1/2 primers and the results confirm the discriminatory capacity of those genetic markers (Figure 1A, B and C). The field confirmation of presence/

absence of seeds in the fruits in these generations F_2, BC_M and BC_{Bs} were obtained later

(Table 1).

In the F_2 generations of the three crosses ($M_1 \times Bs$, $M_2 \times Bs$ and $M_3 \times Bs$), there was a

segregation of the products of the amplification of the LMINO markers that correlated exactly

with the presence/absence of seeds. Fertile plants in this generation uniformly produced an

amplification product with the LMINO1/2 primer set, while plants producing no product

(indicating homozygosity for the *INO* gene deletion) produced only seedless fruit (Figure

1A). The same complete cosegregation pattern seen in F_2 individuals for the presence/absence

of seeds and PCR product was also observed in backcross populations of BC_{Bs} (Figure 1B).

For BC_M backcross plants, the formation of *INO* amplification products was observed in all

DNA samples tested (Figure 1C) for these uniformly fertile/seed bearing plants.

Figure 1. Amplification products for primer pair LMINO1/2 obtained from DNA samples from segregating generations: A) F_2, B) BC_{Bs} C) BC_M in 1.2% agarose gels in 1× TBE buffer. M bp: molecular weight marker in base pairs. Lanes 1 to 20 contain the respective amplified fragments from the 20 genotypes. S: seedless; F: fertile (S and F are phenotypic determinations).

The χ^2 test was performed with the data generated in the molecular analysis to confirm the segregation of the dominant amplification (Table 2). F_1 plants displayed the expected genotypic ratio of 1:0 (presence/absence of amplified band) that had been linked to the trait of seeded fruits. In F_2 generations, six segregation hypotheses expected for one, two and three genes were tested (Supplemental Table S3). Considering a significance of 5% probability, the frequencies of genotypes fit a ratio of 3:1, but allowed rejection of the other predicted ratios, confirming the hypothesis that a single locus confers the phenotype for the trait under study, with the dominant allele responsible for the presence and the recessive allele for the absence of the amplification product.

To identify the homogeneity between the F_2 crossings ($M_1 \times$ Bs, $M_2 \times$ Bs, and $M_3 \times$

Bs), statistical techniques were applied to verify whether the differences observed in the

results could be explained by chance or not. The heterogeneity test was not significant and

indicated, with a 55% likelihood, that the results of the χ^2 were consistent for the populations

of the three families studied, confirming the expected segregation (Table 2).

To further support the hypothesis of segregation in F_2 generations, BC_M and BC_{Bs}

backcrosses were used. Similarly, the heterogeneity of segregation between the families of the

BC_{Bs} backcrossing was not significant (Table 2). BC_M and BC_{Bs} progenies analysed separately

displayed segregation in a manner consistent with the hypothesis of a single gene. In BC_{Bs}

backcrossing, carried out between generations F_1 and the parent Bs, the proportion was close

to 1:1 presence/absence of seeds in the fruits. χ^2 test was applied and the deviations between

the observed and expected frequencies were not significant. In the BC_M backcrossing between

generations F_1 and parents (M_1, M_2, and M_3), the proportion was 1:0 presence/absence of

seeds in the fruits. These results confirmed the monogenic inheritance found in the analyses of

F_2 generations consistent with a single recessive allele being responsible for the seedless trait

in *A. squamosa* considering the 3:1 segregation hypothesis.

321 **Table 2.** Molecular analysis of segregation of dominant primers in parents (M_1, M_2, M_3 and

322 Bs), generations F_1, F_2, BC_M and BC_{Bs} in *A. squamosa*, Janaúba-MG, Brazil.

Parents and Generations	Observed proportion		Expected ratio	χ^2	*p*-value
	Presence	**Absence**			
Parents					
M_1	1				
M_2	1				
M_3	1				
Bs		1			
Generation F_1					
$M_1 \times$ Bs	9	0	1:0		
$M_2 \times$ Bs	17	0	1:0		
$M_3 \times$ Bs	6	0	1:0		
Generation F_2					
$F_1(M_1 \times Bs) \times F_1(M_1 \times Bs)$	99	43	3:1	2.113	0.146[ns]
$F_1(M_2 \times Bs) \times F_1(M_2 \times Bs)$	190	71	3:1	0.676	0.411[ns]
$F_1(M_3 \times Bs) \times F_1(M_3 \times Bs)$	69	33	3:1	2.941	0.086[ns]
Heterogeneity				1.182	0.553[ns]
Backcrosses					
BC_M					
$F_1(M_1 \times Bs) \times M_1$	63	0	1:0		
$F_1(M_2 \times Bs) \times M_2$	67	0	1:0		
$F_1(M_3 \times Bs) \times M_3$	61	0	1:0		
BC_{Bs}					
$F_1(M_1 \times Bs) \times$ Bs	48	61	1:1	1.550	0.213[ns]
$F_1(M_2 \times Bs) \times$ Bs	57	59	1:1	0.034	0.852[ns]
$F_1(M_3 \times Bs) \times$ Bs	26	30	1:1	0.286	0.593[ns]
Heterogeneity				0.586	0.746[ns]

[ns] Non-significant values at the 5% significance level, χ^2 Chi-square test estimate.

The nature of the *INO* gene deletion event

Whole genome shotgun sequencing was used to determine the characteristics of the

INO gene deletion event. A draft wild-type *A. squamosa* genome was assembled through

sequencing of total DNA isolated from a plant grown from seed derived from commercially

available *A. squamosa* fruit. Genomic DNA was sequenced by both long-read (PacBio (Menlo

Park, CA) Single Molecule, Real-Time sequencing) and short read (Illumina (San Diego,

CA)) paired-end 150 base (PE150) methods. The long reads were assembled into a draft

sequence that was corrected with the higher coverage short read sequences. The resulting

assembly comprised 707 Mbases of DNA in 3,519 contigs, with average contig length of 201

kb. A BLAST (Altschul et al., 1990) search with a previously published *A. squamosa INO*

gene sequence (GenBank GU828033.1) was used to identify a 587 kb contig that included the

INO gene (Supplemental Figure S3). Total Hs *A. squamosa* DNA was used to produce a

second short-read sequence set (~140-fold redundant) and this was aligned with the

assembled wild-type sequence. Visualization of the alignment of the Hs sequences with the

587 kb contig including *INO* revealed a clear absence of reads over a region of 16,020 bp

indicating a 16 kb deletion that included the *INO* gene (Figure 2A). The alignment program

truncates read sequences where they do not align with the reference sequence, so a deletion or

a deletion with a heterologous insertion would appear similar in this visualization. Truncated

sequence reads at the immediate borders of the putative deletion region were used to perform

a BLAST search of the Hs sequence reads to identify untruncated read sequences. Identified

sequence reads from the two sides of the deletion region overlapped allowing us to assemble

the Hs sequence in this region (Figure 3). This showed that the lesion was in fact a clean

deletion of the region deficient in read sequences with no inserted DNA and no sequence

duplication at the deletion borders (Figure 2B). The deleted region included the entire *INO*

gene and part of the first coding region of another putative gene encoding a possible UDP-N-

acetylglucosamine-N-acetylmuramyl-pyrophosphoryl-undecaprenol N-acetylglucosamine

protein. The genomic region containing *INO* in wild-type and the corresponding region from

Hs were deposited in GenBank as accessions ON248606 and ON248607, respectively.

A

B

Wild-type

...AGCGCTAGATTGACTAAAAT...(16 kb including *INO*) ...GAGGGCAGGTTTTGTTCGCA...

...AGCGCTAGATTGAGGTTTTGTTCGCA...
Hawaiian seedless

Figure 2. Identification of the *INO* deletion region. A) Screen capture of Tablet (Milne et al., 2013) visualization of the alignment of Hs Illumina sequence reads with the 587 kb contig of *A. squamosa* wild-type sequence that contains the *INO* gene. At top is a view of 500 kb of the alignment with each of the aqua or blue colored dots representing one 150 base read of the Hs sequence. A region including a deficiency of aligning reads is visible and is highlighted by a translucent purple box. The lower portion of the panel is an expanded view of the highlighted region where each small colored line again depicts a 150 base Hs read aligning with the wild-type reference sequence. The near complete absence of aligning reads in the center of this region (barring a small number of reads aligning with repeated sequences within this region) shows a clear deletion in the Hs sequence relative to the wild-type reference sequence. The black arrow edited onto the figure depicts the location of the *INO* gene transcribed region in the wild-type reference sequence. B) Depiction of the wild-type sequence compared to the sequence of the corresponding region in Hs as determined by assembling Hs Illumina reads that overlapped the deletion junction. A deletion of 16,020 bp was observed with no duplication of sequence at the junction site and no additional inserted sequence.

Primers were designed that would amplify the region spanning the deletion in mutant

plant DNA (Figure 3). These primers (AsINODel F and AsINODel R) were used for PCR on

DNA from Hs and Ts lines. Both lines produced a fragment with migration consistent with a

length consistent with the 456 bp expected for the deletion (Figure 4A). Further, because the

region between the primers in wild type is too large to be amplified in standard procedures,

this procedure provides a positive assay for the presence of the deletion. In combination with

primers that will amplify a region of the wild-type *INO* gene, this would enable a codominant

test for the wild-type and seedless alleles of the gene in a single reaction. To test this, we

modeled amplification from a "heterozygote" by mixing DNA from Hs and wild-type *A.*

squamosa. We found that the amplification with the deletion-specific primers and LMINO1/2

primers in a single reaction could clearly differentiate the homozygous seedless, homozygous

wild-type and heterozygous lines (Figure 4B).

Subsequent work showed that AsINODel primer PCR on all three seedless isolates

produced comigrating fragments (Figure 4A and Figure 5A). The sequences of the PCR

products were determined for all three seedless lines and were aligned with the assembled

deletion region sequence (Figure 3). The sequence of the deletion junction was confirmed in

the sequence from Hs, and an identical deletion was present in the same position in the

sequences from mutants Bs and Ts (Figure 3), consistent with all lines producing PCR

products of the same size. Further, the presence of identical deletions in all three lines

indicates a single origin for the *INO* deletion among the isolates.

```
AssembledDelRegion    GCTGTTTCCTTACCTGTTTAGTTGTCTATGTTCCACATGATCAATGATCGACGCCCAAGC
HsDelPCR              ------------------------------------------------------------
TsDelPCR              ------------------------------------------------------------
BsDelPCR              ------------------------------------------------------------

                                     ____AsINODelF_______  >
AssembledDelRegion    TATGCATGCATGTACGGGAACATCATTCTTTAATTGCTGAACCTTTTTTTTAAAAAAAAA
HsDelPCR              --------------------------TTTAATTGCTGAACCTTTTTTTTAAAAAAAAA
TsDelPCR              --------------------------TTTAATTGCTGAACCTTTTTTTTAAAAAAAAA
BsDelPCR              -------------------------------------GAACCTTTTTTTTAAAAAAAAA

AssembledDelRegion    AAAATTCACACGTGTGATGAATTTAATATCTAATTGTGGAAGTGGGCCCCAATTCAACGG
HsDelPCR              AAAATTCACACGTGTGATGAATTTAATATCTAATTGTGGAAGTGGGCCCCAATTCAACGG
TsDelPCR              AAAATTCACACGTGTGATGAATTTAATATCTAATTGTGGAAGTGGGCCCCAATTCAACGG
BsDelPCR              AAAATTCACACGTGTGATGAATTTAATATCTAATTGTGGAAGTGGGCCCCAATTCAACGG

AssembledDelRegion    ACTTATGCATGTTTTGACTTTTGACTCTATATATTCGTTGTTGTTTTTTCAGACAGAACA
HsDelPCR              ACTTATGCATGTTTTGACTTTTGACTCTATATATTCGTTGTTGTTTTTTCAGACAGAACA
TsDelPCR              ACTTATGCATGTTTTGACTTTTGACTCTATATATTCGTTGTTGTTTTTTCAGACAGAACA
BsDelPCR              ACTTATGCATGTTTTGACTTTTGACTCTATATATTCGTTGTTGTTTTTTCAGACAGAACA

AssembledDelRegion    GGTCAAAAGGCTTCGAAGTTTGGAAGAAGGAAGCCATGGAAATAATGTCTCATTTATTCA
HsDelPCR              GGTCAAAAGGCTTCGAAGTTTGGAAGAAGGAAGCCATGGAAATAATGTCTCATTTATTCA
TsDelPCR              GGTCAAAAGGCTTCGAAGTTTGGAAGAAGGAAGCCATGGAAATAATGTCTCATTTATTCA
BsDelPCR              GGTCAAAAGGCTTCGAAGTTTGGAAGAAGGAAGCCATGGAAATAATGTCTCATTTATTCA

AssembledDelRegion    TATTCATGTGATAATTTAGAAAGCATAGCGCTAGATTGagGTTTTGTTCGCAATGAAGAG
HsDelPCR              TATTCATGTGATAATTTAGAAAGCATAGCGCTAGATTGagGTTTTGTTCGCAATGAAGAG
TsDelPCR              TATTCATGTGATAATTTAGAAAGCATAGCGCTAGATTGagGTTTTGTTCGCAATGAAGAG
BsDelPCR              TATTCATGTGATAATTTAGAAAGCATAGCGCTAGATTGagGTTTTGTTCGCAATGAAGAG

AssembledDelRegion    TGTTGGAGGCCTTGTTTGAAAGAAATCCATCCTGCCTGGCTTCTGATTTTTCGCATTGTG
HsDelPCR              TGTTGGAGGCCTTGTTTGAAAGAAATCCATCCTGCCTGGCTTCTGATTTTTCGCATTGTG
TsDelPCR              TGTTGGAGGCCTTGTTTGAAAGAAATCCATCCTGCCTGGCTTCTGATTTTTCGCATTGTG
BsDelPCR              TGTTGGAGGCCTTGTTTGAAAGAAATCCATCCTGCCTGGCTTCTGATTTTTCGCATTGTG

AssembledDelRegion    GCCTTTTGTGTGCTTGTGCCATTGCTTGTTGTCTATGCAGTTGTTGATGGAAGCAGCATA
HsDelPCR              GCCTTTTGTGTGCTTGTGCCATTGCTTGTTGTCTATGCAGTTGTTGA-------------
TsDelPCR              GCCTTTTGTGTGCTTGTGCCATTGCTTGTTGTCTATGCAGTTGTTGA-------------
BsDelPCR              GCCTTTTGTGTGCTTGTGCCATTGCTTGTTGTCTATG----------------------

                                            <___AsINODelR________
AssembledDelRegion    TTTCACTACTATACTCAGTAAGCTGCTTCCAGCTCTTGGTTTGATTGAACAAAATACTCA
HsDelPCR              ------------------------------------------------------------
TsDelPCR              ------------------------------------------------------------
BsDelPCR              ------------------------------------------------------------

AssembledDelRegion    TGATTGATGAAGTTTAGAAAAGTAACTTTAGAAACTATATGTTTGAAACTCAGTCGAGTT
HsDelPCR              ------------------------------------------------------------
TsDelPCR              ------------------------------------------------------------
BsDelPCR              ------------------------------------------------------------

AssembledDelRegion    CTTGAAATATGCTAGCTTCAATGTTCCATATCCATCTTCATGAAGTGTCGAATGAACATG
HsDelPCR              ------------------------------------------------------------
TsDelPCR              ------------------------------------------------------------
BsDelPCR              ------------------------------------------------------------

AssembledDelRegion    TGGAACATAACTGGCA
HsDelPCR              ----------------
TsDelPCR              ----------------
BsDelPCR              ----------------
```

Figure 3. Aligned sequences of the deletion region junction. "AssembledDelRegion" is the sequence assembled from the Illumina reads selected from among all reads by BLAST searches with the reads flanking the deletion. The deletion falls between the "ag" nucleotides shown in lowercase light blue. The other sequences are those of PCR products produced using the AsINODel primers (indicated by arrows above the sequence) on DNA from Hs, Ts, or Bs as indicated, with sequence determined using the AsINODelR primer. All sequences show an identical deletion.

407
408
Figure 4. PCR of *INO* deletion region. A) AsINODel primers designed to amplify a fragment
spanning the deletion junction were used on Hs and Ts DNA and the products were
electrophoresed on an agarose gel. The Hs DNA produced the expected 457 bp fragment, and
a comigrating fragment was produced from Ts. B) A combination of the AsINODel and
LMINO1/2 primer pairs was used in single reactions to amplify DNA from Ts, wild-type and
a mixture of wild-type and Ts ("Mix"). Only the expected 457 and 350 bp fragments were
amplified from the Ts and wild-type DNA, respectively. Both bands were amplified from the
mixture demonstrating effective detection of a heterozygous state.
417
418
419 **Codominant marker analysis**
420
To evaluate use of the AsINODel primers together with the LMINO1/2 pair as

codominant markers for plant breeding, lines M_1, M_2, and M_3 were used as wild-type for *INO*

gene and the presence of seeds, and the mutant Bs for the *INO* deletion and the absence of

seeds. The genotype Hs was used as a positive control for the marker AsINODel. After PCR

reactions, the combined primer pairs amplified a single fragment of 350 bp from the wild-type

parents, representing the presence of the *INO* gene, and a 456 bp fragment from the mutants

Bs and Hs, corresponding to the deletion junction fragment (Figure 5A).

429
Figure 5. Amplifications products for combined AsINODel and LMINO1/2 primer pairs
obtained from DNA samples of populations indicated below the agarose gel lanes (A). M bp:
molecular weight marker in base pairs; Hs: Hawaiian seedless; Bs: Brazilian seedless; M_1,
M_2, M_3: wild-types; F_1: first filial generation (lanes 6 to 11); F_2: self-fertilization (lanes 12 to
20). Generations BC_{Bs} (B) and BC_M (C) lanes 1 to 12 contain the amplified fragments from
the 12 individuals each. S: seedless; F: fertile (phenotypic determinations of the source
plants).
437
As expected, the wild-type fertile parents (M_1, M_2 and M_3) were homozygous for the

presence, and the mutant seedless parent Bs was homozygous for the deletion of the *INO*

gene. In generation F_1 both *INO* gene and deletion region bands were present in all 32

genotypes evaluated, consistent with the expected heterozygous state (Table 3 and Figure 5A).

The codominant markers were also used in genotyping of individuals of the F_2 and

backcross generations (BC_M and BC_{Bs} from $M_2 \times Bs$) (Table 3) with some individuals selected

to illustrate the discriminatory capacity of markers in Figure 5 A, B and C. In generation F_2,

analysis of 145 plants gave results consistent with the expected ratio of 1:2:1 for homozygous

INO: heterozygous: homozygous *ino* deletion (Table 3). The expected ratios of genotypes

were also observed for the sixty-one BC_{Bs} plants and the fifty-six BC_M plants, where ratios

close to 1:1:0 and 0:1:1 were observed respectively (Figure 5B, 5C and Table 3). The χ^2 test

on each of these generations showed probabilities consistent with a monogenic segregation of

the wild-type and mutant alleles.

451
452   **Table 3.** Molecular analysis of codominant segregation of parents (M_1, M_2, M_3 and Bs), F_1
453   generations and generation F_2, BC_M and BC_{Bs} backcrosses of the M_2 family in *A. squamosa*,
454   Janaúba-MG, Brazil.

Parents and Generations	Observed proportion			Expected ratio	χ^2	p-value
	INO INO	*INO ino*	*ino ino*			
Parents						
M_1	1					
M_2	1					
M_3	1					
Bs			1			
Generation F_1						
$M_1 \times$ Bs	0	9	0	0:1:0		
$M_2 \times$ Bs	0	17	0	0:1:0		
$M_3 \times$ Bs	0	6	0	0:1:0		
Generation F_2						
$(M_2 \times$ Bs$) \times (M_2 \times$ Bs$)$	34	70	41	1:2:1	0.848	0.654[ns]
Backcrosses						
BC_M						
$F_1(M_2 \times$ Bs$) \times M_2$	30	26	0	1:1	0.286	0.593[ns]
BC_{Bs}						
$F_1(M_2 \times$ Bs$) \times$ Bs	0	30	31	1:1	0.016	0.898[ns]

455   [ns] Non-significant values at the 5% significance level, χ^2 Chi-square test estimate.
456
Fruit was obtained from sixty-three of the genotyped segregating plants. In every case

where a band corresponding to the wild-type *INO* gene was present (11 homozygous *INO* and

37 heterozygous plants) the seeded phenotype was observed. Similarly, all of the fifteen

plants homozygous for the deletion allele exhibited the Bs mutant seedless phenotype. Thus,

there was 100% correlation of the seeded and seedless phenotypes with the wild-type

containing and homozygous *ino* deletion genotypes, respectively. To obtain more information

to further characterize the linkage between the molecular and visible phenotypes we examined

the ovules of plants that were flowering but not yet producing fruit where the *ino* mutant

ovules could be differentiated from wild-type (Supplemental Figure S2). This determination

was done on ninety-five plants and once again there was a complete correspondence between

the molecular genotype and the presence of wild-type ovules or aberrant ovules incapable of

forming seeds (Supplemental data file 1).

If the *ino* mutant gene is only linked to the seedless mutant allele, and not causally

responsible for it, then recombination between the molecular and phenotypic markers would

be possible. Among segregating F_2 plants, a single recombination event could be observed in

the wild-type *INO* containing chromosome of heterozygous progeny, or in either chromosome

of homozygous *ino* progeny. The same would be true of chromosomes deriving from the F_1

plants in the BC Bs progeny. In these cases, recombination would lead to a switch of the

seedless/seeded phenotype and breakage of the 100% cosegregation (Supplemental Methods).

Seed or ovule phenotypic data were produced for 57 informative heterozygous plants and 36

homozygous mutant plants (allowing examination of a total of 114 chromosomes)

(Supplemental data file 1). Thus, no recombination events were observed between the *INO*

locus and the Bs gene in 114 chromosomes. This allows an initial limit on the maximum

genetic distance between *ino* and Bs. A genetic distance of 3.5 cM would lead to a prediction

of 3.5% recombination, where 0% recombination was observed. The comparison of these

observed and expected result using the χ^2 test produces $\chi^2 = 3.99$ and a corresponding P-value

of 0.046. Thus, a genetic distance of 3 cM (or greater) can be rejected at the 5% probability

level (Supplemental data file 1). In addition, within the 587 kb of sequence surrounding the

INO gene, we find that the 16 kb deletion is the only significant difference between wild-type

and Hs in this region (the only other being a single base change in an intron of a predicted

gene). Together these observations are consistent with the *INO* gene deletion being the lesion

responsible for the seedless phenotype.

489
490   **Genetic diversity using SSR markers**
491
Of the 67 SSR primers used, 41 amplified under the tested conditions, 12 presented a

null amplification pattern with no bands evident, and the remaining 14 did not present a clear

band pattern in the tested samples (Supplemental Table S4). Among the primers that

amplified, 38 generated a total of 63 monomorphic alleles; only three primers (LMCH 3, 39,

and 137) generated specific polymorphisms among the genotypes evaluated.

The LMCH 3 primer generated two alleles between 200 and 300 bp for the accession

M_2 genotype and the seedless genotype, respectively. The primer LMCH 39 amplified five

alleles in total. Three generated a band only in genotype M_2, distinguishing it from seedless

genotypes. Similarly, the primer LMCH 137 amplified five alleles. However, in this marker

the seedless genotypes presented two specific bands differing from genotype M_2, displaying

amplification of two monomorphic bands common to all individuals. The genetic distance

between genotypes, estimated by the complement of the similarity matrix generated by the

Jaccard index, ranged from 0 among genotypes Bs, Ts, and Hs, and 0.0933 between the M_2

with Bs, Ts, and Hs. Thus, these markers differentiated between the mutant lines and the

wild-type line, but could not demonstrate any divergence between the three mutant

accessions.

508
509
510   **Discussion**
511
512
Three seedless stenospermocarpous sugar apple varieties, Bs, Ts, and Hs, from

differing global locales have been described [Lora et al. (2011), Santos et al. (2014) and this

work]. The seedless phenotype results from an ovule defect (Lora et al., 2011, Santos et al.,

2014) indistinguishable from the ovule effects of the *inner no outer* mutation in *Arabidopsis*

(Villanueva et al., 1999), and was associated with a deletion of this locus in the Ts accession

(Lora et al., 2011). On this basis, Lora et al. (2011) hypothesized that the deletion was the

cause of the seedless trait, but segregation data were not available. We show that the three

seedless accessions all carry identical 16 kb deletions of the region including *INO*. The

identical sequence of the deletion region in the three accessions indicates that the mutant

allele arose as a single deletion event in a common ancestor and propagated globally. The

deletion was not associated with any apparent repeated sequence at the deletion junction, but

was a precise excision of the deleted region. We tested the causal relationship between the

deletion and seedlessness by examining segregation of the seedless trait and the deletion. We

demonstrated that seedlessness is a single locus recessive trait and that the fertile/seedless

phenotypes cosegregated 100% with the presence/absence of the *INO* gene among progeny

plants (114 evaluated chromosomes). Our data demonstrate that the *ino* deletion and the

seedless trait are separated by less than 3.5 cM. We have determined the sequence of 587 kb

surrounding the deletion region, and based on prior measures of the relationship between

genetic and physical mapping distances (Puchta and Hohn, 1996) this would constitute from

three to thirty centimorgans and so would include the seedless mutation. Outside of the

deletion that includes the *INO* gene, we found no other significant differences from wild type

in the sequenced region surrounding the *INO* locus. Together with the clear expectation of an

ino gene deletion causing the observed ovule phenotype, these data support the *ino* deletion

being the sole cause of the seedless trait.

It is possible that the deletion and seedless trait was retained in multiple lines by

selection despite significant outcrossing, or alternatively that all lines with this deletion are

vegetatively propagated clones. Genotyping through the use of SSR markers failed to detect

polymorphism among our available seedless accessions (Bs, Hs, Ts) but did show a low level

of polymorphism (4.7%) when compared to the wild-type M_2 (indicating low genetic

diversity among genotypes). These data therefore do not rule out a possible spread by

vegetative propagation. The utilized SSR markers were developed for *A. cherimolia* and no

specific SSR primers were available for *A. squamosa* and some outbreeding could have been

missed in this analysis. But with the current data the simplest explanation is that a single

mutational event was dispersed to different continents through vegetative propagation.

Tracing a temporal and geographical route of this dispersal is difficult since the species has

been widely developed and naturalized.

It is believed that the primary center of origin of *A. squamosa* is in the lowlands of

Central America. Historical data indicate accessions to Mexico by the natives of the region

(Pinto et al., 2005). According to these authors, after their arrival, the Spaniards were

responsible for spreading the seeds to the Philippines. By 1590, the species had been

introduced into India. In Brazil, 1626 is the first record of introduction of the species. By the

early 17th century the species was already widespread in Indonesia, China, Australia,

Polynesia and Hawaii. In 1955, the Cuban seedless variety had been introduced in the state of

Florida in the United States. The mutant Brazilian accession Bs was first described in 1940 in

the state of São Paulo (Cunha, 1953). We obtained Hs from a commercial nursery in Hawaii.

The nursery reports obtaining the line from the Philippines. Their source in the Philippines

reports that the line was brought to the Philippines from an unknown location in the 1920s,

but documentary evidence of this was not found. Information on the origin of Ts was not

available.

Lora et al. (2011) designed primers within, or closely flanking the transcribed region

of *INO* for detection of the wild-type locus, and demonstrated amplification from wild-type *A.*

squamosa, A. cherimola, and four other species that cover the phylogenetic range of the genus

Annona. These primers were confirmed to function as expected in our *A. squamosa* varieties

(Nassau et al., 2021 and this work). Notably the primer pairs only allow detection of the wild-

type locus and a homozygous mutant could only be detected by the absence of amplification.

According to Singh and Singh (2015), markers that can directly identify the traits of interests

should be more efficient in incorporating and monitoring genes in breeding programs. Our

determination of the sequence of the region from which *INO* has been deleted enabled

formulation of primers (AsINODel) for direct detection of the mutant allele allowing

codominant detection of both alleles in wild-type and seedless mutant parents and to

differentiate fertile heterozygotes from homozygous fertile plants in a single reaction.

The improvement of perennial plants, especially fruit species, is a challenging task due

to a long juvenile period and seed dormancy, among other factors, which translates into a

relatively long generation period (McClure et al., 2014). In *A. squamosa*, for example, the

beginning of flowering of adult plants, originated from seeds, under specific cultivation

conditions, would take at least three years (Pereira and Borém, 2021) until the

characterization of the fruits, limiting the number of genotypes that can be studied, which

makes the breeding process slow and expensive. The PCR assay using a combination of the

primers detecting the wild-type and mutant alleles allows plants to be genotyped in the

seedling stage, accelerating the breeding program. Given the generality of the *INO* gene

detecting primers for use in multiple species of *Annona* (Lora et al., 2011), and the fertility of

interspecific crosses in this genus, our codominant genotyping is applicable to introgression

of the seedless trait into elite sugar apple varieties and into other cultivated *Annona* species.

586
587
588   **Conclusions**
589
590
The presence/absence of seeds in *A. squamosa* displays a monogenic inheritance. The

INO allele*,* whether homozygous or heterozygosis (*INO__*), conditions the presence of seeds

and has complete dominance over the recessive seedless condition (*ino ino*). The Bs, Ts and

Hs mutant accessions all display the same deletion sequence for the *INO* locus, demonstrating

a common origin of genotypes. The deletion is simple, not resulting in insertion or duplication

of any sequence at the deletion junction. Determination of the deletion sequence allowed

formulation of primers (AsINODel) that specifically detect the deletion. The AsINODel

primers together with the marker LMINO1/2 display co-dominance and can distinguish

dominant homozygotes from heterozygotes. The validated LMINO and AsINODel primers

can facilitate introgression of the seedless trait in *A. squamosa* and other members of this

genus.

602
603
604 **Acknowledgments**
605
606

607   We thank Trang Dang for technical support and Debra Skinner for help with planning

608 and analysis for sequence determination.

609
610
611
612 **Funding**
613
614

We are grateful for the support of the Coordination for the Improvement of Higher Education

Personnel - Brazil (*Coordenação de Aperfeiçoamento de Pessoal de Nível Superior* - CAPES)

- Financing Code 001, the National Council for Scientific and Technological Development

(*Conselho Nacional de Desenvolvimento Científico e Tecnológico* - CNPq), the U. S. National

Science Foundation grant IOS1354014 (to CSG) and the Fundação de Amparo à Pesquisa do

Estado de Minas Gerais - FAPEMIG) for granting scholarships.

621
Author Contributions

Experiments were conceived by S.N., S.P and C.S.G. Crossing and genetic analyses were

performed by B.R.A.R. with help from M.C.T.P. Whole genome sequence analysis was

performed by C.S.G. and B.R.A.R. S.N., B.R.A.R and C.S.G. wrote the manuscript.

The author responsible for distribution of materials integral to the findings presented in this

article in accordance with the policy described in the Instructions for Authors

(https://academic.oup.com/plphys/pages/general-instructions) is: Silvia Nietsche

(silvia.nietsche@gmail.com)

References

Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ (1990) Basic local alignment search tool. J Mol Biol **215**: 403–410. https://doi.org/10.1016/S0022-2836(05)80360-2.

Araújo JF, Araújo JF, Alves AAC (1999) Instruções técnicas para o cultivo da pinha (*Annona squamosa* L.). Salvador, BA: EBDA

Bowman JL (2000) The YABBY gene family and abaxial cell fate. Curr Opin Plant Biol **3**: 17–22. https://doi.org/10.1016/S1369-5266(99)00035-7.

Buffalo V (2011) Scythe—A very simple adapter trimmer. GitHub, San Fr Available https://github.com/najoshi/scythe Accessed Oct **17**: 2013

Cheung WY, Hubert N, Landry BS (1993) A simple and rapid DNA microextraction method for plant, animal, and insect suitable for RAPD and other PCR analyses. Genome Res **3**: 69–70. https://doi.org/10.1101/gr.3.1.69.

Cruz CD (2016) Genes Software-extended and integrated with the R, Matlab and Selegen. Acta Sci Agron **38**: 547–552. https://doi.org/10.4025/actasciagron.v38i4.32629.

Cunha JC (1953) Pinha sem semente. Chácara e Quintais **88**: 839–841

Doyle JJ, Doyle JL (1990) Isolation of plant DNA from fresh tissue. Focus **12**: 13–15

Escribano P, Viruel MA, Hormaza JI (2008) Development of 52 new polymorphic SSR markers from cherimoya (*Annona cherimola* Mill.): transferability to related taxa and selection of a reduced set for DNA fingerprinting and diversity studies. Mol Ecol Resour **8**: 317–321. https://doi.org/10.1111/j.1471-8286.2007.01941.x.

Escribano P, Viruel MA, Hormaza JI (2004) Characterization and cross-species amplification of microsatellite markers in cherimoya (*Annona cherimola* Mill., Annonaceae). Mol Ecol Notes **4**: 746–748. https://doi.org/10.1111/j.1471-8286.2004.00809.x.

Joshi NA, Fass JN (2011) Sickle: a sliding-window, adaptive, quality-based trimming tool for FastQ files. (Version 1.33) [Software]. Available at https://github.com/najoshi/sickle. Accessed month day: year (Oct 17: 2013)

Koren S, Walenz BP, Berlin K, Miller JR, Bergman NH, Phillippy AM (2017) Canu: scalable and accurate long-read assembly via adaptive k-mer weighting and repeat separation. Genome Res **27**: 722–736. https://doi.org/10.1101/gr.215087.116.

Li H (2013) Aligning sequence reads, clone sequences and assembly contigs with BWA-MEM. arXiv Prepr. arXiv1303.3997. https://doi.org/10.48550/arXiv.1303.3997.

Lora J, Hormaza JI, Herrero M, Gasser CS (2011) Seedless fruits and the disruption of a

conserved genetic pathway in angiosperm ovule development. Proc Natl Acad Sci
108: 5461–5465. https://doi.org/10.1073/pnas.1014514108.

Lora J, Larranaga N, Hormaza JI (2018) Genetics and Breeding of Fruit Crops in the
Annonaceae Family: *Annona* spp. and *Asimina* spp. *In* JM Al-Khayri, SM Jain, D V
Johnson, eds, Springer International Publishing, Cham, pp 651–672.
https://doi.org/10.1007/978-3-319-91944-7_16.

McClure KA, Sawler J, Gardner KM, Money D, Myles S (2014) Genomics: a potential
panacea for the perennial problem. Am J Bot **101**: 1780–1790.
https://doi.org/10.3732/ajb.1400143.

Mendes HTA, Costa MR, Nietsche S, Oliveira JAA, Pereira MCT (2012) Pollen grain
germination and fruit set in 'Brazilian seedless' sugar apple (*Annona squamosa* L.).
Crop Breed Appl Biotechnol **12**: 277–280. https://doi.org/10.1590/S1984-
70332012000400007.

Milne I, Stephen G, Bayer M, Cock PJA, Pritchard L, Cardle L, Shaw PD, Marshall D
(2013) Using Tablet for visual exploration of second-generation sequencing data.
Brief Bioinform **14**: 193–202. https://doi.org/10.1093/bib/bbs012.

Nassau B, Mascarenhas PSC, Guimarães AG, Feitosa FM, Ferreira HM, Castro BMC,
Zanuncio JC, Costa MR, Nietsche S (2021) Inheritance of seedlessness and the
molecular characterization of the INO gene in Annonaceae. Brazilian J. Biol. 83:
e246455. https://doi.org/10.1590/1519-6984.246455.

Pereira MCT, Borém A (2021) Anonáceas: do plantio a colheita. Viçosa, MG: UFV

Pinto AC de Q, Cordeiro MCR, De Andrade SRM, Ferreira FR, Filgueiras HA de C,
Alves RE, Kinpara DI (2005) *Annona* species. Southampton, UK: International
Centre for Underutilised Crops. https://www.embrapa.br/en/busca-de-publicacoes/-
/publicacao/890581/annona-species.

Puchta H, Hohn B (1996) From centiMorgans to base pairs homologous recombination in
plants. Trends Plant Sci. **1**: 340-348. https://doi.org/10.1016/S1360-1385(96)82595-0.

Santos RC, Ribeiro LM, Mercadante-Simoes MO, Costa MR, Nietsche S, Pereira MCT
(2014) Stenospermy and seed development in the "Brazilian seedless" variety of sugar
apple (*Annona squamosa*). An Acad Bras Cienc **86**: 2101–2108.
https://doi.org/10.1590/0001-3765201420140206.

Sawa S, Watanabe K, Goto K, Kanaya E, Morita EH, Okada K (1999) FILAMENTOUS
FLOWER, a meristem and organ identity gene of Arabidopsis, encodes a protein with
a zinc finger and HMG-related domains. Genes Dev **13**: 1079–1088.
https://doi.org/10.1101/gad.13.9.1079.

Siegfried KR, Eshed Y, Baum SF, Otsuga D, Drews GN, Bowman JL (1999) Members of
the YABBY gene family specify abaxial cell fate in Arabidopsis. Development **126**:
4117–4128. https://doi.org/10.1242/dev.126.18.4117.

Singh BD, Singh AK (2015) Marker-Assisted Selection. *In* Marker-Assisted Plant Breeding:
Principles and Practices. New Delhi: Springer. https://doi.org/10.1007/978-81-322-
2316-0_9.
734
735   **de Souza DA, de Melo LC, Librelon SS, Costa MR, Nietsche S, Pereira MCT** (2010)
736   Identification of hybrids of intra and interspecific crosses in Annonaceae by RAPD
737   markers. Crop Breed. Appl. Biotechnol. **10**: 110–115.
738   http://dx.doi.org/10.12702/1984-7033.v10n02a02.
739
740   **Varoquaux F, Blanvillain R, Delseny M, Gallois P** (2000) Less is better: new approaches
741   for seedless fruit production. Trends Biotechnol **18**: 233–242.
742   https://doi.org/10.1016/s0167-7799(00)01448-7.
743
744   **Villanueva JM, Broadhvest J, Hauser BA, Meister RJ, Schneitz K, Gasser CS** (1999)
745   INNER NO OUTER regulates abaxial–adaxial patterning in Arabidopsis ovules.
746   Genes Dev **13**: 3160–3169. https://doi.org/10.1101%2Fgad.13.23.3160.
747
748   **Walker BJ, Abeel T, Shea T, Priest M, Abouelliel A, Sakthikumar S, Cuomo CA, Zeng**
749   **Q, Wortman J, Young SK** (2014) Pilon: an integrated tool for comprehensive
750   microbial variant detection and genome assembly improvement. PLoS One **9**:
751   e112963. https://doi.org/10.1371/journal.pone.0112963.
752
753   **Wongs-Aree C, Noichinda S** (2011) Sugar apple (*Annona squamosa* L.) and atemoya (*A.*
754   *cherimola* Mill.× *A. squamosa* L.). Postharvest Biol. Technol. Trop. Subtrop. Fruits.
755   Elsevier, pp 399-427e. http://dx.doi.org/10.1533/9780857092618.399.
756

Parsed Citations

Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ (1990) Basic local alignment search tool. J Mol Biol 215: 403–410. https://doi.org/10.1016/S0022-2836(05)80360-2.
  Google Scholar: Author Only Title Only Author and Title

Araújo JF, Araújo JF, Alves AAC (1999) Instruções técnicas para o cultivo da pinha (Annona squamosa L.). Salvador, BA: EBDA
  Google Scholar: Author Only Title Only Author and Title

Bowman JL (2000) The YABBY gene family and abaxial cell fate. Curr Opin Plant Biol 3: 17–22. https://doi.org/10.1016/S1369-5266(99)00035-7.
  Google Scholar: Author Only Title Only Author and Title

Buffalo V (2011) Scythe-A very simple adapter trimmer. GitHub, San Fr Available https://github.com/najoshi/scythe Accessed Oct 17: 2013
  Google Scholar: Author Only Title Only Author and Title

Cheung WY, Hubert N, Landry BS (1993) A simple and rapid DNA microextraction method for plant, animal, and insect suitable for RAPD and other PCR analyses. Genome Res 3: 69–70. https://doi.org/10.1101/gr.3.1.69.
  Google Scholar: Author Only Title Only Author and Title

Cruz CD (2016) Genes Software-extended and integrated with the R, Matlab and Selegen. Acta Sci Agron 38: 547–552. https://doi.org/10.4025/actasciagron.v38i4.32629.
  Google Scholar: Author Only Title Only Author and Title

Cunha JC (1953) Pinha sem semente. Chácara e Quintais 88: 839–841
  Google Scholar: Author Only Title Only Author and Title

Doyle JJ, Doyle JL (1990) Isolation of plant DNA from fresh tissue. Focus 12: 13–15
  Google Scholar: Author Only Title Only Author and Title

Escribano P, Viruel MA, Hormaza JI (2008) Development of 52 new polymorphic SSR markers from cherimoya (Annona cherimola Mill.): transferability to related taxa and selection of a reduced set for DNA fingerprinting and diversity studies. Mol Ecol Resour 8: 317–321. https://doi.org/10.1111/j.1471-8286.2007.01941.x.
  Google Scholar: Author Only Title Only Author and Title

Escribano P, Viruel MA, Hormaza JI (2004) Characterization and cross-species amplification of microsatellite markers in cherimoya (Annona cherimola Mill., Annonaceae). Mol Ecol Notes 4: 746–748. https://doi.org/10.1111/j.1471-8286.2004.00809.x.
  Google Scholar: Author Only Title Only Author and Title

Joshi NA, Fass JN (2011) Sickle: a sliding-window, adaptive, quality-based trimming tool for FastQ files. (Version 1.33) [Software]. Available at https://github.com/najoshi/sickle. Accessed month day: year (Oct 17: 2013)
  Google Scholar: Author Only Title Only Author and Title

Koren S, Walenz BP, Berlin K, Miller JR, Bergman NH, Phillippy AM (2017) Canu: scalable and accurate long-read assembly via adaptive k-mer weighting and repeat separation. Genome Res 27: 722–736. https://doi.org/10.1101/gr.215087.116.
  Google Scholar: Author Only Title Only Author and Title

Li H (2013) Aligning sequence reads, clone sequences and assembly contigs with BWA-MEM. arXiv Prepr. arXiv1303.3997. https://doi.org/10.48550/arXiv.1303.3997.
  Google Scholar: Author Only Title Only Author and Title

Lora J, Hormaza JI, Herrero M, Gasser CS (2011) Seedless fruits and the disruption of a conserved genetic pathway in angiosperm ovule development. Proc Natl Acad Sci 108: 5461–5465. https://doi.org/10.1073/pnas.1014514108.
  Google Scholar: Author Only Title Only Author and Title

Lora J, Larranaga N, Hormaza JI (2018) Genetics and Breeding of Fruit Crops in the Annonaceae Family: Annona spp. and Asimina spp. In JM Al-Khayri, SM Jain, D V Johnson, eds, Springer International Publishing, Cham, pp 651–672. https://doi.org/10.1007/978-3-319-91944-7_16.
  Google Scholar: Author Only Title Only Author and Title

McClure KA, Sawler J, Gardner KM, Money D, Myles S (2014) Genomics: a potential panacea for the perennial problem. Am J Bot 101: 1780–1790. https://doi.org/10.3732/ajb.1400143.
  Google Scholar: Author Only Title Only Author and Title

Mendes HTA, Costa MR, Nietsche S, Oliveira JAA, Pereira MCT (2012) Pollen grain germination and fruit set in 'Brazilian seedless' sugar apple (Annona squamosa L.). Crop Breed Appl Biotechnol 12: 277–280. https://doi.org/10.1590/S1984-70332012000400007.
  Google Scholar: Author Only Title Only Author and Title

Milne I, Stephen G, Bayer M, Cock PJA, Pritchard L, Cardle L, Shaw PD, Marshall D (2013) Using Tablet for visual exploration of second-generation sequencing data. Brief Bioinform 14: 193–202. https://doi.org/10.1093/bib/bbs012.
Google Scholar: Author Only Title Only Author and Title

Nassau B, Mascarenhas PSC, Guimarães AG, Feitosa FM, Ferreira HM, Castro BMC, Zanuncio JC, Costa MR, Nietsche S (2021) Inheritance of seedlessness and the molecular characterization of the INO gene in Annonaceae. Brazilian J. Biol. 83: e246455. https://doi.org/10.1590/1519-6984.246455.
Google Scholar: Author Only Title Only Author and Title

Pereira MCT, Borém A (2021) Anonáceas: do plantio a colheita. Viçosa, MG: UFV
Google Scholar: Author Only Title Only Author and Title

Pinto AC de Q, Cordeiro MCR, De Andrade SRM, Ferreira FR, Filgueiras HA de C, Alves RE, Kinpara DI (2005) Annona species. Southampton, UK: International Centre for Underutilised Crops. https://www.embrapa.br/en/busca-de-publicacoes/-/publicacao/890581/annona-species.
Google Scholar: Author Only Title Only Author and Title

Puchta H, Hohn B (1996) From centiMorgans to base pairs homologous recombination in plants. Trends Plant Sci. 1: 340-348. https://doi.org/10.1016/S1360-1385(96)82595-0.
Google Scholar: Author Only Title Only Author and Title

Santos RC, Ribeiro LM, Mercadante-Simoes MO, Costa MR, Nietsche S, Pereira MCT (2014) Stenospermy and seed development in the "Brazilian seedless" variety of sugar apple (Annona squamosa). An Acad Bras Cienc 86: 2101–2108. https://doi.org/10.1590/0001-3765201420140206.
Google Scholar: Author Only Title Only Author and Title

Sawa S, Watanabe K, Goto K, Kanaya E, Morita EH, Okada K (1999) FILAMENTOUS FLOWER, a meristem and organ identity gene of Arabidopsis, encodes a protein with a zinc finger and HMG-related domains. Genes Dev 13: 1079–1088. https://doi.org/10.1101/gad.13.9.1079.
Google Scholar: Author Only Title Only Author and Title

Siegfried KR, Eshed Y, Baum SF, Otsuga D, Drews GN, Bowman JL (1999) Members of the YABBY gene family specify abaxial cell fate in Arabidopsis. Development 126: 4117–4128. https://doi.org/10.1242/dev.126.18.4117.
Google Scholar: Author Only Title Only Author and Title

Singh BD, Singh AK (2015) Marker-Assisted Selection. In Marker-Assisted Plant Breeding: Principles and Practices. New Delhi: Springer. https://doi.org/10.1007/978-81-322-2316-0_9.
Google Scholar: Author Only Title Only Author and Title

de Souza DA, de Melo LC, Librelon SS, Costa MR, Nietsche S, Pereira MCT (2010) Identification of hybrids of intra and interspecific crosses in Annonaceae by RAPD markers. Crop Breed. Appl. Biotechnol. 10: 110–115. http://dx.doi.org/10.12702/1984-7033.v10n02a02.
Google Scholar: Author Only Title Only Author and Title

Varoquaux F, Blanvillain R, Delseny M, Gallois P (2000) Less is better: new approaches for seedless fruit production. Trends Biotechnol 18: 233–242. https://doi.org/10.1016/s0167-7799(00)01448-7.
Google Scholar: Author Only Title Only Author and Title

Villanueva JM, Broadhvest J, Hauser BA, Meister RJ, Schneitz K, Gasser CS (1999) INNER NO OUTER regulates abaxial–adaxial patterning in Arabidopsis ovules. Genes Dev 13: 3160–3169. https://doi.org/10.1101%2Fgad.13.23.3160.
Google Scholar: Author Only Title Only Author and Title

Walker BJ, Abeel T, Shea T, Priest M, Abouelliel A, Sakthikumar S, Cuomo CA, Zeng Q, Wortman J, Young SK (2014) Pilon: an integrated tool for comprehensive microbial variant detection and genome assembly improvement. PLoS One 9: e112963. https://doi.org/10.1371/journal.pone.0112963.
Google Scholar: Author Only Title Only Author and Title

Wongs-Aree C, Noichinda S (2011) Sugar apple (Annona squamosa L.) and atemoya (A. cherimola Mill.× A. squamosa L.). Postharvest Biol. Technol. Trop. Subtrop. Fruits. Elsevier, pp 399-427e. http://dx.doi.org/10.1533/9780857092618.399.
Google Scholar: Author Only Title Only Author and Title