A cellular and transcriptomic dissection of the human breast for studying mechanisms of cell and tissue function

Katelyn Del Toro[1,3], Rosalyn W. Sayaman[2,3,4], Kate Thi[2], Yamhilette Licon-Munoz[1], and William C. Hines[1]

[1]*Department of Biochemistry and Molecular Biology; University of New Mexico School of Medicine, 1 University of New Mexico MSC08 4670, Albuquerque NM 87131, USA*

[2]*Life Sciences Division, Lawrence Berkeley National Laboratory; Mailstop 977R225A, 1 Cyclotron Road; Berkeley, CA 94720, USA.*

[3]Equal contributions to work

Current affiliations:

[4]Present address: Department of Laboratory Medicine, Helen Diller Family Comprehensive Cancer Center, University of California, San Francisco, San Francisco, CA 94143, USA.

Correspondence and contact for reagent and resource sharing should be addressed to Curt Hines (e-mail: wchines@salud.unm.edu), www.breastcancerlab.com.

Keywords: Breast, flow cytometry, FACS, RNA-sequencing (RNA-seq), human mammary epithelial cells (HMEC), Gene Set Enrichment Analysis (GSEA), Cytoscape, Enrichment Map, luminal epithelial cells, myoepithelial, vascular smooth muscle, pericytes, endothelial, lymphatic, fibroblasts, adipocyte-derived mesenchymal cells, adipocytes, leukocytes, erythrocytes.

Abstract

A fundamental question in biology, central to our understanding of cancer and other pathologies, is determining how different cell types coordinate to form and maintain tissues. Recognizing the distinct features and capabilities of the cells that compose these tissues is critical. Unfortunately, the complexity of tissues often hinders our ability to distinguish between neighboring cell types and, in turn, scrutinize their transcriptomes and generate reliable and tractable cell models for studying their inherently different biologies. In a companion article, we introduced a novel method that permits the identification and purification of the twelve cell types that compose the human breast—nearly all of which could be reliably propagated in the laboratory. Here, we explore the nature of these cell types. We sequence mRNAs from each purified population and investigate transcriptional patterns that reveal their distinguishing features. We describe the differentially expressed genes and enriched biological pathways that capture the essence of each cell type, and we highlight transcripts that display intriguing expression patterns. These data, analytic tools, and transcriptional analyses form a rich resource whose exploration provides remarkable insights into the inner workings of the cell types composing the breast, thus furthering our understanding of the rules governing normal cell and tissue function.

Introduction

Understanding the biology of tissues and ultimately developing effective cancer therapies requires us to recognize the symphony of interactions in tissues that occur among its many different cell types—and the myriad of molecules they synthesize. Determining how cells communicate with each other and their microenvironments is critical, as is defining the roles of stromal cells in regulating epithelial cell behavior, shaping tissues, and promoting or preventing overt cancer[1-4]. To this puzzle, there are many pieces. Clarifying and appreciating the roles of each cell type requires a substantial foundation in our ability to identify and distinguish among these distinct tissue elements so we may learn how they fit together to form a functioning organ. Tumors are caricatures of normal tissues[5], yet the intricate cellular composition and arrangement of cells in many tissues, including the breast, have yet to be adequately dissected, leaving many unanswered questions about these cells' unique and essential qualities.

The human dissections and inspiring observations made by Andreas Vesalius, William Harvey, and other Renaissance anatomist-physicians produced astonishing breakthroughs in medical science[6,7]. In a companion article and with a similar passion for anatomical dissection, we applied modern tools of histology and cell and molecular biology to examine the fabric of the human breast—to learn its cellular composition and study its mechanistic underpinnings. Using immunofluorescence and advanced flow cytometry, we developed methods to resolve, isolate, quantify, and culture each breast cell type, reconciling every cell in the tissue. The product of this labor was a fluorescence-activated cell sorting (FACS) strategy that used redundant markers and a rigorous gating strategy to identify the twelve distinct breast cell types (Figure 1a, Figure 1—figure supplement 1)]. These populations included two separate luminal epithelial cell types, myoepithelial cells, adipocytes, leukocytes, pericytes, vascular smooth muscle cells, erythrocytes, adipose-derived mesenchymal stem cells (fibroblasts), lymphatic and vascular endothelial cells, and a newly identified rare epithelial cell population with unknown function.

Although this FACS method resolved leukocytes as a single CD45[Pos] fraction, the strategy is
adaptable and can be further refined to scrutinize the array of leukocyte populations in the tissue.
We now routinely sort and culture nine of the twelve breast cell types (excluded are red blood
cells, leukocytes, and vascular smooth muscle cells). Some of these primary models are the first
of their kind from this tissue, including breast pericytes, lymphatic endothelial cells, and our
newly defined epithelial cell population. Their isolation and culture add to the repertoire of
available models and permit functional studies that were not previously possible[8].

Here, we asked the critical question of what defines each breast cell type—and sought to identify
their characteristic features and functions. Unsurprisingly, hierarchical clustering of
transcriptional profiles echoed FACS results by cleanly separating the eleven sequenced cell
types—revealing remarkable correlations among biological replicates (of each cell type, isolated
from random individuals). Principal component analysis (PCA) established genes that heavily
contribute to cellular diversity. We used PCA-derived gene lists to identify the distinguishing
features of epithelial, mesenchymal, endothelial, and leukocyte cell lineages—each being a
group of cell types with similar origins and characteristics. These included well-recognized
features, such as keratinization of epithelial cells, muscle contraction by mesenchymal cells,
vasculogenesis by endothelial cells, and more. While these results validated our analytic
approach, they revealed many other processes we had not previously associated with a specific
cell lineage. Shifting to a more gene-centric approach, we investigated what families of genes
contributed to cell-type-specific functions by identifying the gene families that varied most
across the spectrum of cell types. Many gene families showed little to no change in expression
(e.g., exportins), whereas others were extensively differentially expressed (e.g., laminins,
integrins, Rho GTPase activating proteins, and more). Differences in these genes' expression,
and the biological processes in which they are involved, were further solidified—and others
identified—via comparisons between cell-type pairs and application of gene set enrichment and
pathway analyses to reveal the 'essence' of each cell type. We highlighted six comparisons that
we believed to be of the highest interest as they reveal features of every major cell type and
identify those distinguishing closely related cell types. Moreover, we have performed all 55
possible pairwise cell-type comparisons and provided the results in a user-friendly analysis tool
(Pathway Analyzer[9]) and web application (Breast Cell Atlas: Transcript Finder[10]). These tools
allow one to thoroughly engage with the dataset and actively explore each cell type's expressed
genes and enriched biological pathways.

Results and Discussion

Identification of cell lineages and analytic pipeline

The breast is a compound tubuloalveolar gland whose subcutaneous tissue comprises, at a
minimum, 12 major cell types. In an adjoining article, we describe the development and
validation of an antibody panel and gating strategy capable of isolating these cell types[8], cell
populations we numbered from one to twelve (which refers to their respective FACS gate using
the isolation strategy outlined in Figure 1a and Figure 1—figure supplement 1). These cell
populations are described as follows: ER[Pos] (Pop1/Lum1) and ER[Neg] (Pop2/Lum2) luminal
epithelial cells, myoepithelial cells (Pop3), a rare epithelial cell population (Pop4), vascular

smooth muscle cells (vSMCs, Pop5), pericytes (Pop6), lymphatic endothelial cells (Pop7),
fibroblasts (Pop8), vascular endothelial cells (Pop9), CD45[Pos] leukocytes (Pop10), erythrocytes
(Pop11), and adipocytes (Pop12). These different cell types are often broadly classified as
members of epithelial, endothelial, mesenchymal, or hematopoietic lineages (Figure 2b).

In the early days of this study, we had not yet completed identifying all populations revealed by
FACS, and several remained a mystery even after finalizing our isolation strategy. Establishing
their identities was particularly challenging. We were aided by several helpful clues, however.
One was the distinct morphologies the cell types displayed when cultured. These observations
allowed us to determine whether a particular population appeared epithelial-like, e.g., Pop4
epithelial cells, or presented with more mesenchymal-like characteristics, e.g., Pop6 pericytes.
We also backtracked and stained tissues using the same antibody combinations used in FACS
and hunted for cell types in the tissue that shared identical staining patterns. Ultimately, the
transcriptome information from RNA-sequencing provided definitive evidence that solidified
each cell type's identity.

With their identities and histological locations now solved (Figure 2a-f, Figure 2—figure
supplement 1-2)[11], we shifted our attention to exploring the individual biologies of these
different cell types by diligently dissecting their transcriptomes through a comparative lens
provided by the comprehensive collection of cell types we had isolated and sequenced. Our
comprehensive analytic pipeline is described in Figure 3 (Figure 3, Figure 3—figure supplement
1).

mRNA sequencing and validation of FACS purified cell types

To investigate the unique biology of the breast cell types, we FACS isolated each population to
sequence and analyze their transcriptomes. This process entailed processing and collagenase-
treating breast tissues (reduction mammoplasty tissues free of proliferative disease), followed by
low speed (<100g) centrifugation. The first population collected was the lipid layer that had
phase-separated under centrifugal force (Pop12 adipocytes). All other cell types were FACS-
purified from the pelleted tissue 'organoids' following their trypsinization to single-cell
suspensions (Figure 1 and Figure 1- figure supplement 1). While developing and testing our
FACS antibody panel, we typically stained between 100,000 to 500,000 cells[8]. However, many
more were needed for mRNA library construction and sequencing here, so we scaled up our
operation by increasing the number of cells stained by roughly 1000-fold.

FACS purifying cells for RNA-sequencing was performed on freshly collected tissues. This
process required relatively large amounts of tissue (>30 grams) and sort times (14 to 18
continuous hours for each tissue) to produce cell yields that could support deep RNA-
sequencing. In addition to the above lipid layer, the sorting provided 11 FACS-purified cell
populations from each specimen. This exercise yielded hundreds of thousands to millions of cells
for the more abundant cell types (Figure 3-table supplement 1). In contrast, yields were much
lower for the rarer types, which produced a range of only 24,000-130,000 cells. After cell
purification, the RNA isolated from each sample correlated with cell counts, and in pilot
experiments, we found it essential to isolate RNA immediately following FACS to preserve its
integrity. In two independent sequencing runs, we sequenced 36 samples representing eleven of

the twelve identified breast cell types (red blood cells did not produce measurable RNA and thus were not sequenced). All cell populations were isolated from four random individuals (aged 16, 23, 27, and 29 years/old), and nearly all were sequenced in triplicate (from three of the four individuals, Figure 3—table supplement 2). In only two instances did we not have enough RNA of sufficient quality to sequence in triplicate –and both were rare populations with low FACS yields (Pop5 vascular smooth muscle cells and Pop7 lymphatic endothelial cells, Figure 1b, Figure 4a). We also isolated and sequenced RNAs from snap-frozen tissues and unsorted cell suspensions, which we used as mixed cell controls. After sequencing mRNAs from the samples and aligning the FASTQ sequencing data to the human genome, we calculated the average mRNA counts across all samples to be 39.5 million paired-end reads—all with Phred[12] quality scores of 25 or more, indicating the sequencing was of high depth and quality. We found these early sequencing results encouraging and moved forward with quality control analyses.

Nearly all sequenced samples demonstrated excellent read counts. However, covariance analysis, assessment of Cook's distances, and distribution of regularized log values[13] identified five samples with reduced transcript complexity and lower correlations. All were the rarer cell types, i.e., Pop4 epithelial cells, Pop5 vSMCs, and lymphatic endothelial cells (Figure 4b,c). Nevertheless, we determined that only Pop4 epithelial cells from individual N239 needed to be removed from critical downstream analyses. We retained the four other low-complexity samples because they clustered well with their cell-type counterparts (hierarchical clustering and PCA) and provided meaningful insights for many transcripts (Figure 4b, Figure 3—table supplement 2). These samples were: Pop4 epithelial cells (from individuals N225 and N228), Pop5 vSMCs (from N228), and Pop7 lymphatic endothelial cells (from individual N228). Although outlier genes existed in these samples, concordance between replicates was high for most others. Despite these minor issues, the sequencing data across the board was impressively robust. We did not detect sequencing artifacts, such as correlations with sequencing run or individuals from which the cells were derived, as all samples clustered tightly based on their inherent cell type (Figure 4b,d). The substantial number of reads detected per sample, the highly correlated replicates, and the large number of unique transcripts detected in these specimens (21,559) indicated that the dataset was rich and of excellent quality. Indeed, these data were immediately helpful in determining the identities of several obscure cell populations.

Before we had sequenced the mRNAs from the cell types, there were several FACS populations whose identities were a mystery (as discussed in the adjoining article)[8]. Most notably were populations #4, 5, and 6, which we ultimately identified as a rare epithelial cell type, vascular smooth muscle cells, and pericytes. RNA-sequencing provided conclusive evidence that helped identify these cell types by revealing their relationships to other populations and supplying information on expressed markers. After identifying these three populations, we explored all remaining cell types. Reassuringly, transcript levels of genes encoding cell surface markers used to sort each cell type tracked nicely with their respective FACS-staining levels (Figure 4—figure supplement 1a,b; Figure 1a). And GAPDH, often used as a uniformly-expressed internal control gene, was indeed expressed equally among the cell types, as were the transcripts of other widely-used internal control genes (Figure 4—figure supplement 1c, Figure 3—table supplement 2). In addition, analysis of hundreds of other genes, such as those uniquely expressed by each cell type

165 or those used as cell type markers, i.e., genes encoding CD antigens, validated and helped us
166 build confidence in our dataset (Figure 5, Figure 5—figure supplement 1). To facilitate the
167 exploration of these RNA-sequencing data, we have created a publicly available tool for
168 visualizing transcript levels in each cell type (http://breastcancerlab.com/breast-atlas/).

Identification of cell-type specific transcriptional patterns

One of our early goals was to identify the distinct transcriptional patterns of the different breast
cell types. To address this question and further explore the dataset, we clustered the samples'
transcriptomes using a standard Euclidean distance metric. Hierarchical clustering recapitulated
the distinct nature of the different cell types and mixed cell controls (Figure 6). Clustering was
robust, especially for the two luminal cell types, myoepithelial cells, pericytes, fibroblasts,
vascular endothelial cells, and adipocyte samples (all of which displayed approximately unbiased
(AU) p-values >0.95 when evaluated by multiscale bootstrap resampling[14]). Notably, biological
replicates of each cell type clustered tightly together, and similar cell types formed adjacent
clusters. These related pairs included: a) vascular and lymphatic endothelial cells (FACS Pops
7&9), b) vSMCs and pericytes (Pops5&6), c) ER[Pos] and ER[Neg] luminal epithelial cells (Pops
1&2), and d) myoepithelial cells and related Pop4 epithelial cells (Pops3&4). As expected,
fibroblasts (Pop8) clustered with the other mesenchymal cell types (vSMCs and pericytes)—
albeit at a greater distance— reflecting their inherently distinct features. In contrast, the most
distant samples that stood far from the others were the CD45[Pos] leukocytes (Pop10)—a
separation that we anticipated, given their unique biology. The clustering confirmed that the
eleven sorted cell types were indeed distinct and captured the underlying differences—and
similarities— in their gene expression patterns.

Several features emerged after closely examining the hierarchical clustering of genes. Gene
expression trends, in particular, were apparent across populations, and many clusters contained
genes encoding prominent cell markers, which helped us identify or confirm each cell type's
identity (Figure 6). For example, p-selectin (SELP) and HOXD1 were genes in a cluster that
differentiated endothelial cells. Epithelial cell adhesion molecule (EPCAM), zona occludens 3
(TJP3), and epidermal growth factor (EGF) were among the many genes in a cluster highly
expressed by luminal epithelial cells. Calmodulin-like 3 (CALML3) and G-protein-coupled
receptor 87 (GPR87) were highly expressed by Pop4 epithelial and myoepithelial cells, along
with the chloride channel regulator CLCA2, which is regulated by the myoepithelial lineage
marker p63 and is involved in basal cell adhesion. Finally, the perivascular cell cluster contained
angiopoietin 1 (ANGPT1) and connexin 40 (GJA5), which are essential for blood vessel
development and maintenance. The clustering thus demonstrated high agreement among the
biological replicates and identified clusters of shared genes that provided substantial evidence
confirming the identity of each cell population (or pointing to its identity).

The hierarchical clustering used to compare cell types was also notably stable. Whereas Figure 6
shows the clustering of the 2000 most differentially expressed genes, the pattern did not
significantly change when we repeated the analysis with different inclusive thresholds, such as
when we used the top 100, 200, 500, 1000, or 3000 genes that were most differentially
expressed. Although the cellular relationships that emerged from these analyses were largely

confirmatory, they helped us identify and characterize several populations early on. For example,
before we had ascertained the identity of FACS Pop5 as vSMCs, its close clustering with related
Pop6 pericytes here provided a critical clue[8]. The clustering also revealed similarities between
the rare Pop4 epithelial cells and myoepithelial cells. However, it also showed that Pop4's
expression of some genes more resembled those found in luminal epithelial cells that
distinguished this cell population from myoepithelial cells[8]. The molecular patterns that emerged
provided information on each population and objective evidence of the relatedness among cell
types. To explore these relationships—and features fundamental to each cell type—we needed to
decipher their transcriptomes in finer detail.

Distinguishing the features of cell lineage

Differentiated cells are often classified by their developmental history and the lineage from
which they are derived. The different breast cells, for example, are often described as being
mesenchymal (fibroblasts, pericytes, vSMC, and adipocytes), hematopoietic/blood (leukocytes,
erythrocytes), endothelial (vascular and lymphatic), or epithelial derived (ER^{Pos} and ER^{Neg}
luminal, myoepithelial, and Pop4 epithelial, Figure 2b). Although our primary goal was to
identify the qualities that make each cell type unique, our sequencing of every breast cell type
presented an unprecedented opportunity to explore the broad attributes of cell lineage through
the comparative lens afforded by our dataset's comprehensive nature. We accomplished this
through principal component and pathway analysis, as this approach allowed us to analyze the
samples en masse—and not focus on any particular pairwise comparison prematurely. As a
result, these analyses produced an astonishing amount of data. Unfortunately, we can present
only a fraction of it here. However, we have provided additional tools and tables to aid the
exploration of all data and results presented herein (Figure 3—table supplements, Pathway
Analyzer[9], and the Breast Cell Atlas: Transcript Finder App[10]). Below are our analyses of the
first four principal components that revealed the distinguishing features of the four broad cell
lineages composing breast tissues.

Principal component analysis (PCA) of the transcripts from all cell types showed gene
expression variances were distributed across multiple principal components (PCs), with the bulk
of the variation (71%) being captured by the first four components (Figure 7—figure supplement
1). The projection of principal component 1 (PC1) and principal component 2 (PC2) markedly
separated the four cell lineages in two-dimensional space. Genes contributing to PC1 were
primarily responsible for distinguishing the mesenchymal and epithelial cell types, as these
lineages projected at PC1's opposing ends (Figure 7a). Interestingly, endothelial cells—which
derive from mesenchyme but are classically regarded as a specialized epithelial cell type[15,16]—
projected as an independent group that aligned more closely with the mesenchymal cell types
along PC1. In contrast, the variance explained by genes composing the second principal
component (PC2) emerged from leukocytes, as these cells formed a single group at its distal end,
far apart from the other cell types (Figure 7a). A similar analysis of principal components 3 and 4
resolved the endothelial cells and two subtypes of the mesenchymal and epithelial lineages, i.e.,
perivascular cells and myoepithelial/Pop4 epithelial cells (Figure 8a).

To gain further insights into the distinguishing attributes shared by members of each cell lineage,
we examined the genes impacting their PCA groupings, i.e., the loading factors with the highest
influence on each principal component (Figure 3—table supplement 3). We then used these
unique gene lists to probe functionally grouped Gene Ontology[17] and Reactome[18] pathway
annotation networks[19].

Epithelial Lineage *(ER^Pos & ER^Neg Luminal cells, Myoepithelial cells, Pop4 Epithelial cells).*
We first analyzed the cluster defined by the epithelial lineages: the ER^{Pos} and ER^{Neg} luminal myoepithelial
and Pop4 epithelial cell types (Figure 2b). Based on their positioning on PC1, we expected the genes with
the highest contributions to PC1 to reflect those that distinguish the epithelial and mesenchymal lineages,
and in particular, highlight genes associated with luminal epithelial cells—as the ER^{Pos} and ER^{Neg} luminal
cells projected the farthest along on PC1. Analysis of gene expression in each cell type showed this was
indeed the case (Figure 7b). Many of PC1's top genes (at the negative epithelial end) were recognizable,
as they are often used as markers of the epithelial lineage. They included keratins 6b, 8, 18, and 19, P-
and E- cadherins, claudin 4, and Her2 (Figure 7b, Figure 9a-e; Figure 3—table supplement 3). Other
familiar genes included integrin beta 6, the transcription factors ELF3, EHF, and TFAP2A; peptidase
inhibitors SERPINA3 and SPINT2; serine protease ST14, cytoskeletal and associated proteins
SHROOM3, MYO5D, and MYH14; the calcium-binding protein S100A14, among others (Figure 3—
table supplement 3). These genes were among the 200 extracted from the negative epithelial end of
PC1—a gene list we used to identify the biological pathways enriched within the epithelial cell lineage.

Using the epithelial gene set, we performed a two-sided hypergeometric (enrichment/depletion)
test to determine if particular molecular functions or biological pathways were disproportionately
represented (Gene Ontology[20], Reactome Pathway database[18]). We found many were, and we
aggregated the results to form a functionally organized pathway network, which grouped similar
biological processes based on their shared genes (ClueGO[21]). The 124 enriched gene sets
aggregated into twenty-four distinct groups related to cell adhesion, peptidase activity,
keratinization, estrogen signaling, AP-2 transcription factor regulation, mucins, and innate
immune system factors (Figure 10a, Figure 3—table supplement 4: PC1 Epi tab). The most
significant group (*p*-value =1.4×10^{-8}) contained the *cell adhesion molecule binding* and *cadherin*
binding Gene Ontology terms. These gene sets represent different levels of the same ontology
hierarchy and are highly correlated. For example, they both contain components of adherens
junctions, i.e., epithelial- and placental-cadherins (CDH1 and CDH3), cingulin (CGN), and 348
other genes. By comparison, the *cell adhesion* gene set contains an additional 196 genes related
to other aspects of cellular adhesion, such as desmosome-associated desmoplakin (DSP) and
desmoglein 2 (DSG2, Figure 3—table supplement 3, sorted by PC1). The algorithm's ability to
cluster overlapping gene sets like these, and assemble them into related clusters, allowed us to
extract the common biological themes obscured within the volume and complexity of the data.
Reassuringly, when we examined the expression of the genes in these pathways, we found they
were indeed elevated in the four epithelial cell types (Figure 10b-e, (Figure 3—table supplement
4: PC1 Epi tab). These enriched pathways and themes describe general features of the epithelial
lineage. After exploring the remaining cell lineages' shared attributes (defined by each PC), we
will revisit these cells and explore their individual traits.

Leukocytes *(CD45^Pos Cells).* The second principal component separated the leukocytes (CD45^{Pos}
FACS population), and analysis of PC2's loading factors revealed many well-known leukocyte

markers (Figure 7a,c). For example, 'protein tyrosine phosphatase receptor type C' (PTPRC), the gene encoding the pan-leukocyte marker CD45—which we used to FACS isolate the leukocytes—was notably the first gene on the list. Other top genes included the CD antigens: CD2, CD3E, CD6, CD7, CD8A, and others. Also present was the transcription factor RUNX3, which is critical for T-cell differentiation, as was selectin P ligand (SELPLG), which is crucial for leukocyte trafficking. Using the same technique described above (for PC1 and the epithelial cells), we extracted the top genes from the leukocyte-defining end of PC2 and performed pathway enrichment analysis. We identified 453 enriched gene sets that clustered into 62 distinct thematic groups (Figure 3—table supplement 4: PC2 Leukocyte tab). Representative pathways from the ten most significant groups were: *adaptive immune response, activation of immune response, leukocyte homeostasis, T-cell activation, leukocyte chemotaxis, phagocytosis, IL-6 production, acute inflammatory response, regulation of GTPase activity, mast cell degranulation, and monocyte differentiation* (p-values ranged between $6.07 \times 10^{-46} - 9.53 \times 10^{-11}$, Figure 10—figure supplement 1a). Although most leukocytes in the breast are macrophages[8], the sequenced mRNAs from the CD45^{Pos} FACS gate represent a diverse population of white blood cells. Therefore, finding a broad range of enriched pathways attributable to different leukocyte subpopulations was expected. When we compared the expression of enriched genes in each cell type, we found they were indeed highly expressed by the Pop10 leukocytes (Figure 10—figure supplement 1b-d).

Mesenchymal Lineage *(vSMCs, Pericytes, Fibroblasts, Adipocytes).* Similar gene expression patterns within the four mesenchymal cell types positioned this cell group toward the positive end of PC1's axis in the 2D projection of PC1 and PC2 (Figure 7a). However, endothelial cells were also located in this region, indicating they and mesenchymal cells share the distinct expression of many genes. The relatively similar projection of endothelial and mesenchymal cells along PC1 prevented extraction of loading factors that could be solely attributable to either cell class. As we were primarily interested in identifying the defining characteristics of mesenchymal cells at this stage, we digitally removed the endothelial cell data and repeated the principal component analysis. The new PCA projection resolved the mesenchymal cells, situating them at the positive end of the first principal component in this ancillary graph (PC1^{No} ^{ENDO}, Figure 7—figure supplement 2a). Exploring the loading factors of this new principal component, we confirmed that these genes were indeed highly expressed by the mesenchymal cells. For example, vimentin (VIM), an intermediate filament protein and classical mesenchymal marker (Figure 9f), was near the top of the list, at position 192 (out of 21,560 genes, Figure 7—figure supplement 2b). The top loading factor was the non-selective receptor for endothelins (EDNRB). Endothelins are potent vasoactive peptides, and EDNRB expression was notably highest in the perivascular cell types (vSMC and pericytes) –these cells surround blood vessels and regulate vessel diameter by influencing vasoconstriction and dilation. Genes highly contributing to PC1 included many collagen family members (e.g., COL4A1, COL1A2, COL6A3, COL4A2, COL3A1, COL15A1, COL5A3, and others). Also present were ADAM metalloproteinases, the transcription factors ZEB1and ZEB2, platelet-derived growth factor receptor beta (PDGFRB), insulin-like growth factor II, laminin subunit alpha-4 (LAMA4), interleukin 24 (IL24), chondroitin sulfate proteoglycan 4 (CSPG4), along with many other notable genes (Figure 7— figure supplement 2b, Figure 3—table supplement 3: PC1^{NoEndo}). To

understand how these genes collectively contribute to mesenchymal cells, we performed pathway analysis using the loading factors from the mesenchymal-defining end of $PC1^{No\ ENDO}$.

Pathway enrichment using 200 of PC1's most critical mesenchymal genes identified 337 enriched gene sets from the Reactome and Gene Ontology databases (p-values< 0.05). These pathways assembled into a network of 96 distinct thematic groups (Figure 3—table supplement 4: PC1 Mesenchymal tab). The most significant group (p-value =$5x10^{-33}$) contained 94 different aggregated gene sets/pathways that dealt mainly with: cardiac development (chamber, muscle, septum), cartilage morphogenesis, smooth muscle proliferation, bone development, and chondrocyte differentiation—pathways that are consistent with our primary cultures showing the isolated breast fibroblasts have multi-lineage differentiation potential[8]. The defining genes in this group included TBX2 and HEYL transcription factors, the type III TGFβ receptor TGFBR3, fibroblast growth factor FGF2, ADAM metallopeptidase with thrombospondin type 1 motif 1a and 9 (ADAMTS1, ADAMTS9); platelet-derived growth factor receptor beta (PDGFRB), and others (Figure 11a). Representative pathways from the other top significant groups included: *Fibroblast migration, GAG metabolic process, chondrocyte development, bone development, smooth muscle contraction, negative regulation of calcium ion transport, regulation of fatty acid biosynthetic process, and dermatan sulfate biosynthesis,* with p-values ranging between $5.00x10^{-33}$ to $4.01x10^{-08}$ (Figure 11a, Figure 3—table supplement 4: PC1 Mesenchymal tab).

While the four mesenchymal cell types highly expressed genes in the enriched pathways, transcript levels were not always uniform across these cell types (Figure 11b-d). For example, in the *cardiac chamber development* gene set, expression of the pathway's genes was consistently higher in the perivascular cell types. The heart is composed of mesenchymal cardiac muscle cells and is a highly vascularized tissue that requires perivascular support, which may explain the frequent enrichment of cardiac-related pathways within the mesenchymal gene set (Figure 11—cardiac chamber development). The enrichment with these and other tissue-related gene sets (cartilage, bone, muscle, and adipose) indicates a deeply-rooted bond among mesenchymal cells and common dependence on a core set of genes. Later, we will use specific pairwise comparisons to explore the distinguishing features of the different mesenchymal cell types.

Endothelial Lineage *(Lymphatic and Vascular Endothelial Cells).* To identify the genes and processes responsible for shaping the unique characteristics of endothelial cells, we explored the third and fourth principal components. These components resolved the vascular and lymphatic endothelial cells as a distinct group near the positive end of PC3 (Figure 8a). Many genes contributing to the endothelial end of PC3 were familiar, such as vascular-endothelial cadherin (CDH5)—a prominent endothelial adhesion molecule (Figure 8b). We also observed von Willebrand factor (VWF); the angiopoietin regulator tie1 (TIE1); the endothelial marker CD31 (PECAM1); endothelial selectin (SELE); angiopoietin 2 (ANGPT2); as well as vascular endothelial growth factor receptors KDR and FLT4, and many other notable genes. The top gene on the list was 'podocalyxin like' (PODXL), which binds leukocyte selectin and is vital to lymphocyte-endothelial adhesion. Also present was Claudin 5, which is crucial for endothelial cell junctions. Finding so many genes with known importance to endothelial cells corroborated these genes' notable contributions to endothelial cells and showed they are characteristics of both vascular and lymphatic cell types. At the same time, the analysis identified other influential

genes that are perhaps less appreciated or whose importance to endothelial cell function are not
as well characterized (listed in Figure 9, Figure 3—table supplement 3). To explore the
biological contribution of these genes to endothelial cell biology, we performed pathway
enrichment analysis.

Hypergeometric analysis of the endothelial genes from PC3 identified 650 enriched biological
pathways (p-values< 0.05, Figure 3—table supplement 4: PC3 Endothelial tab). These pathways
assembled into 61 groups based on the genes they shared. Not surprisingly, most groups
pertained to blood vessel and circulatory system biology (Figure 12a). The most significant
group (p-value =6.27×10^{-19}) involved cell motility and chemotaxis and contained *ameboidal-type*
cell migration, *angiogenesis*, and *signaling by VEGF* gene sets (Figure 12a,b). As expected,
PC3-identified genes within these enriched pathways were highly and differentially expressed by
the lymphatic and endothelial cell types (Figure 12b-d). Representative pathways from these
groups included: *cell-matrix adhesion, endothelial- and muscle- cell differentiation, lymph vessel*
morphogenesis, regulation of establishment of endothelial barrier, regulation of coagulation,
vasculogenesis, and monocyte chemotaxis. Many notable genes were present in these pathways,
including podoplanin (PDPN) —a marker we used to sort lymphatic endothelial cells— ephrins
A1 and B2 (EFNA1 and EFNB2), activin a receptor like type 1 (ACVRL1), platelet-derived
growth factor subunit b (PDGFB), along with the aforementioned cadherin 5, claudin 5 and the
VEGF receptors (KDR and FLT4). Finding vascular-related pathways enriched in the endothelial
data set was not surprising, but identifying these pathways was reassuring, as it validated the
approach and indicated these processes were occurring in both lymphatic and vascular
endothelial cell types found in the breast.

Outside of the anticipated pathways, enrichment analysis of the endothelial genes also
uncovered: *Signaling by SCF-KIT* (stem cell factor/c-kit)*, adherens junction organization,*
regulation of peptidyl-tyrosine phosphorylation (JAK/STAT)*, semaphorin receptor binding,*
interleukin-6 secretion (and IL-10, IL-12 cytokines)*, aminoglycan biosynthetic process,*
hyaluronic acid binding, along with roughly 500 other enriched processes and pathways (p-
values ranging between 1.64E-06 - 2.08E-02, Figure 3—table supplement 4: PC3 Endothelial
tab). Enrichment of '*Signaling by SCF-KIT'* pathway was driven by elevated endothelial
expression of the kit tyrosine kinase receptor (KIT), the GRB2-Related Adapter Protein (GRAP),
and LYN Proto-Oncogene (LYN). Notably, Kit's ligand, stem cell factor (KITLG), was
differentially expressed between endothelial cell types, as vascular endothelial cells expressed it
at a much higher level (40-fold higher) than lymphatic endothelial cells (Figure 12—figure
supplement 1).

Endothelial semaphorin expression was intriguing. Semaphorins are a large family of signaling
ligands that exist as either secreted or membrane-associated proteins which provide movement
and guidance cues[22]. The *Semaphorin receptor binding* pathway was enriched predominantly
because of endothelial expression of semaphorin 3a, 3d and 6b (SEMA3A, SEMA3D, SEMA6B,
p-value=1.47×10^{-3}; (Figure 12—figure supplement 2, Figure 3—table supplement 4: PC3
Endothelial tab). The class 3 semaphorins are the only semaphorins in humans that are secreted,
and we found that both SEMA3A and SEMA3D were expressed rather exclusively by lymphatic
endothelial cells (Figure 12—figure supplement 2a). Interestingly, the co-receptor for sema3

proteins, neuropilin1 and 2 (NRP1 and NRP2), which bind VEGF family members, were highly expressed by the endothelial cell types (*p*-values of 3.13×10^{-27} and 1.23×10^{-53}, respectively), with vascular endothelial cells (and pericytes) differentially expressing the highest levels of NRP1. In contrast, the highest levels of NRP2 occurred in lymphatic endothelial cells, which was 2-fold higher than in their vascular counterparts (Figure 12—figure supplement 2b,c). Others have implicated NRP2 as a critical factor in lymphangiogenesis[23,24], and our results are consistent with it fulfilling such a role for breast lymphatic cells.

The above analysis has identified many features shared between lymphatic and vascular endothelial cells. Later, we will directly compare these two endothelial cell types and explore their individual traits.

Perivascular sub-lineage *(vSMC and Pericytes).* After analyzing the endothelial cells, we were left with two other cell groups to explore that were resolved by PC3 and PC4. Perivascular cells were the first of these (Figure 8a). This mesenchymal cell subclass includes pericytes and vascular smooth muscle cells (vSMCs), and the group projected toward the negative end of the third principal component, opposite to the endothelial cells. Like fibroblasts and adipocytes, perivascular cells are mesenchymal in nature; but perform critical functions to the blood vessels they envelop. PC3's loading factors (at the perivascular end) included genes with known pericyte and vSMC significance (Figure 8c), such as smooth muscle actin (ACTA2, Figure 9g), smooth muscle calponin (CNN1), glutamyl aminopeptidase (ENPEP), regulator of G-protein signaling 5 (RGS5), and platelet-derived growth factor receptor beta (PDGFRB). At the top of the list was the atypical cadherin FAT1, along with phosphodiesterase 3A (PDE3A), peptidase inhibitor 15 (PI15), and carboxypeptidase E (CPE, Figure 8c). Both perivascular cell types differentially expressed AXL receptor tyrosine kinase, which was sixth on the list and, in addition to its other physiological functions, serves as a receptor/co-receptor for many viruses, including Ebola, Lassa, Marburg[25], SARS-CoV-2[26], and Zika viruses[27] –which may help explain the vessel involvement often observed in infected individuals. The perivascular cells also notably and differentially expressed members of the Notch and Hedgehog signaling pathways, such as NOTCH3, HEYL, GLI2, and GLI3, suggesting these cell types fulfill fundamental roles during tissue morphogenesis and maintenance. As before, we used the top 200 genes from the defining principal component for pathway enrichment analysis.
Hypergeometric analysis of the perivascular genes identified 456 enriched processes and pathways from the Gene Ontology and Reactome databases (*p*-values< 0.05, Figure 3—table supplement 4: PC3 Perivascular tab). These aggregated into 24 distinct groups (Figure 13a). Many groups reflected vascular-related biology in line with known perivascular cell functions. Among these were *regulation of blood vessel diameter, regulation of systemic arterial blood pressure by hormone,* and *smooth muscle contraction.* Enriched pathways from other significant groups included: *Epithelial tube morphogenesis, blood vessel-* and *vascular- development, ECM proteoglycans, artery-* and *cardiac tissue- development, cell fate commitment, syndecan interactions,* and *mesenchymal cell differentiation* (Figure 13 b-e). These analyses identify characteristic processes of both perivascular cell types.

Myoepithelial sub-lineage *(Myoepithelial Cells and Pop4 Epithelial Cells).* The final cell group separated by PC3 and PC4 was an epithelial subclass comprised of myoepithelial cells and

related Pop4 epithelial cells (a rare cell type we identified during FACS panel development[8]).
The fourth principal component resolved these cells, and when we investigated PC4's loading
factors, we found that the most significant gene was none other than keratin 5 (KRT5). Finding
keratin 5 was reassuring because it has a long and storied history as a marker of basal epithelial
cells[28-30]. Another top gene was alpha-6 integrin (ITGA6/CD49f), which we anticipated because
CD49f formed the backbone of our FACS panel –and myoepithelial cells consistently exhibited
the highest CD49f staining intensities (Figure 1a, Figure 9f,h). Among the many genes
contributing significantly to PC4 (Figure 8d), included the well-known myoepithelial markers:
keratins 14 (KRT14, Figure 8d) and 17 (KRT17), laminin subunit alpha-3 (LAMA3), the alpha-1
chain of collagen type XVII (COL17A1), smooth muscle actin (ACTA2, Figure 9g), p63 (TP63,
Figure 9i), E-cadherin (CDH1, Figure 9c,h), P-cadherin (CDH3), the transcriptional regulatory
factor TRIM29, and integrin subunit beta 4 (ITGB4, Figure 8d, Figure 3—table supplement 3).
Many of these myoepithelial genes notably contribute to the molecular cluster defining the basal-
like breast cancer subtype—which provided the rationale for its 'basal-like' designation[31].

As with the preceding cell lineages, we extracted the top 200 loading factors from the defining
end of the principal component (PC4, in this case) and used the gene list to explore potentially
enriched pathways and processes. Pathway enrichment identified 333 gene sets, which
aggregated into 24 groups of related biological functions (Figure 3—table supplement 4: PC4
Myoepithelial (MEP) tab). Processes related to the basement membrane, extracellular matrix
organization, and laminin interactions led the way, forming the most significantly enriched group
(p-value $=3.46 \times 10^{-16}$, Figure 14a). These gene sets were dominated by integrins (ITGA3, ITGA6,
ITGB4, ITGB6, ITGB8) and laminin chains (LAMA3, LAMB3, LAMC2), which were highly
expressed by myoepithelial cells (Figure 14b-c). Notably, these three laminin genes encode the
a3, b3, and g2 chains that form the heterotrimeric laminin 332 molecule—an essential epithelial
basement-membrane-specific variant that reinforces the cellular adhesion to the basal lamina[32].

Representative pathways from the other significantly enriched groups included: *cell junction*
organization, keratinization and skin development, prostate glandular acinus development,
morphogenesis of a branching epithelium, epithelial tube formation, actin filament bundle
assembly, smooth muscle contraction, intermediate filament organization, regulation of Wnt
signaling pathway, and others. (Figure 14a-e, Figure 3—table supplement 4: PC4 Myoepithelial
tab). Among the many integrin, laminin, keratin, and collagen genes represented in these
enriched pathways were cadherins CDH1 and CDH3, desmosome-related desmocollin 3 (DSC3)
and desmogleins DSG2 and DSG3, stratafin (SFN, 14-3-3 sigma); wnt-modulators 'secreted
frizzled-related protein 1' (SFPR1) and 'Dickkopf wnt signaling pathway inhibitor 3' (DKK3);
epidermal growth factor receptor (EGFR), fibroblast growth factor 1 (FGF1), and others (Figure
3—table supplement 4: PC4 Myoepithelial tab).

Myoepithelial cells have historically been the most studied breast cell type, likely because of
their prevalence in the tissue and ease of culture. Our results are consistent with prior knowledge
of myoepithelial function: that these cells are basally located, help shape the basement membrane
to which they attach via hemidesmosomes and are rich with smooth muscle actin and myosin
that provide them with contractile ability. Furthermore, the mammary gland is formed through
invagination of the skin, and the myoepithelial gene list reflects this connection, as it contains

genes associated with epidermal development. This list includes basal keratins (KRT-14, -17, -5, -6b, -6c, and -7), the laminin 332 chains, desmosomal genes, vitamin D receptor (VDR), and transcription factors SOX9 and p63. These attributes distinguish myoepithelial cells from all others in the breast. Our results pinpoint the specific factors produced by myoepithelial cells, affirming their pivotal role in ECM-mediated cellular functions.

This section highlights the broad characteristics of cell lineages composing the breast. These characteristics emerged from pathway enrichment analyses using gene lists defined by the principal components aligning and separating each cell lineage. Hundreds of pathways were revealed. Some we anticipated, whereas many others were not. Although we have here highlighted the most statistically significant pathways emerging from this analysis (Figures 10-14), many more exist and are listed in the supplementary tables (Figure 3—table supplement 4). A full review of these results will undoubtedly provide a more comprehensive perspective.

Gene families contributing to cellular identity

Above, we explored the distinguishing qualities of different cell lineages by determining the pathways enriched in PCA-defined gene lists. These analyses relied on curated gene sets from the Gene Ontology (GO) and Reactome databases—but not every human gene is represented in these archives. For example, the Reactome knowledgebase currently contains 11,291 human proteins (including known isoforms) that, by our count, are encoded by just over 9,300 genes. Because we measured over 20,000 genes in each cell type, many are not captured in the above analysis. Therefore, we have complemented the prior results by shifting to a more 'gene-centric' approach. This method entailed identifying all differentially expressed (DE) genes in the breast cell dataset (DESeq2[13], adj. p-value ≤ 0.1, Figure 3—table supplement 5). We then used this information to identify gene families with the most variably expressed genes. Due to their differential expression, these gene types must contribute to cell-specific processes and the unique features of each cell type. Their identification and patterns of expression provided countless insights.

The HUGO Gene Nomenclature Committee has organized 25,320 human genes into 1,519 manually curated gene families (HGNC gene group index[33,34]). When we explored the breast cells' transcriptomes, we identified 1,421 gene families in our dataset (Figure 3—table supplement 6). These included *vacuolar-type ATPases*, *matrix metallopeptidases*, *mucins*, *collagens*, *glutathione transferases*, *erb-B2 tyrosine kinases*, *histones*, *interleukins*, and more (Figure 3— Table supplement 6). The sizes of the families varied widely, from 1 to 831 genes ($\bar{x} = 14, \mathrm{sd} = 42.2$) with a median family size of six genes per family (Fig.15—figure supplement 1).

To determine the families with the most variably expressed genes, we examined the expression of each gene, from each family, within every breast cell type—and focused on families that had six or more genes (≥ median size). We discovered 187 gene families composed entirely of differentially expressed genes (Figure 15a). Twenty of these families remained once we excluded those containing fewer than six genes. These included *Rho GTPase activating proteins*, *fibulins*, *histocompatibility complex genes*, *laminin subunits, integrin beta subunits, and more* (Figure 15b, Figure 3—Table supplement 6). The cellular processes affected by genes in these twenty

families are wide-ranging, as they are involved in Rho signaling, cAMP regulation, interactions
with the immune system, and cellular adhesion—between other cells, the extracellular matrix,
and basement membranes. Below, we highlight nine families that we believe are most intriguing,
but we have also provided a full accounting of each gene family in a table supplement—
indicating the number and percentage of genes differentially expressed among them (Figure 3—
Table supplement 6)

Laminins. Each gene family told a story. The family of laminins, for example, is composed of 12
genes that code for laminin a, b, and g chains. These chains aggregate to form an array of
heterotrimeric laminin molecules with distinct properties that polymerize in the extracellular
matrix and help shape basement membranes. Above, we found that myoepithelial cells highly
expressed the chains composing laminin 332 (Figure 14). Inspection of the entire laminin family
revealed that fibroblasts highly express four laminins (LAMA1, LAMA2, LAMB1, and
LAMC3); endothelial cells express the highest levels of three (LAMA4, LAMB1, and LAMC1);
whereas ER^{Pos} luminal, vascular endothelial, and pericytes significantly expressed LAMA5
(Figure 15b- laminin subunits). There were also several intriguing instances where closely
related cell types expressed widely disparate levels of a particular laminin chain. Notable
examples were the differential expression of LAMA1 between ER^{Pos} and ER^{Neg} luminal cells,
LAMB1 between the two endothelial cell types, and LAMC3 between pericytes and vSMCs.
Furthermore, no cell type meaningfully expressed LAMB4, consistent with its consideration as a
pseudogene[35].

The relationship between cells and their basement membranes is dynamic and reciprocal. Each
cell type contributes to the basal lamina's structural diversity by secreting basement membrane
components like the laminins and other extracellular molecules. These integral basement
membrane proteins anchor the cells via membrane-tethered receptors and transmit signals to the
cells, where they become integrated and influence multiple cellular processes.

Fibulins. One such related and intriguing ECM-related gene family was the fibulins. Fibulins are
RGD-containing extracellular matrix glycoproteins that bind basement membrane proteins, such
as tropoelastin, fibrillin, and fibronectin[36]. Whereas we found that fibroblasts highly expressed
most fibulin family members, fibulin 5 (FBLN5) levels were highest in the ER^{Neg} luminal
epithelial cells (Figure 15b- fibulins). FBLN5 was also notably expressed by lymphatic
endothelial cells at substantially higher levels than their vascular counterparts. Fibulin 5 is one of
the smallest fibulins, is secreted, has unique calcium and integrin binding properties, and is
essential for elastic fiber formation[36]—and thus likely contributes to the elasticity of the
lymphatic endothelium. We also found fibulin 3 (EFEMP1) was also differentially expressed by
lymphatic endothelial cells, presumably providing similar contributions.

The fibulins and laminins are just two examples of the 187 gene families whose members were
entirely differentially expressed. Of the remaining families (having fewer than 100% DE genes),
we identified 211 containing between 75 to 99% DE genes (Figure 15a). Over half of these
families were composed of six or more genes. These 126 families included *integrin alpha*
subunits, *toll-like receptors, kruppel like factors, adenylate cyclases, low-density lipoprotein*

receptors, glypicans, interleukin receptors, ephrins, and others (Figure 3—Table supplement 6).
One of the larger and more familiar families was the *collagens*.

Collagens. The family of collagens consists of 45 individual genes, and, as expected, we found
mesenchymal and myoepithelial cell types highly expressed most of them. There were some
notable exceptions, however. For example, the vascular endothelial cells highly expressed
COL15A1 and COL9A3, whereas COL22A1 was expressed uniquely by luminal epithelial cells
and adipocytes (Figure 15c). Particularly enlightening were the cell types expressing the collagen
IV genes.

Collagen IV is a sheet forming collagen and the principal constituent of basement membranes. It
provides cellular integrin and non-integrin attachment sites through its triple-helical and non-
collagenous domains[37]. There are six collagen IV isoforms, which assemble to form one of three
triple-helical heterotrimers, α1α1α2, α3α4α5, or α5α5α6[37,38]. We found the expression of
collagen IV chains in the breast cells quite revealing, as the most robust levels occurred in cells
associated with the basal lamina, but patterns of isoform expression differed by cell type. For
example, pericytes and vSMCs, which are embedded within, and help maintain the vascular
basement membrane, expressed the highest levels of COL4A1, COL4A2, and COL4A4 (Figure
15c, Figure 15—figure supplement 2a-c). In contrast, myoepithelial cells, which contribute
significantly to epithelial basement membranes, exhibited the highest levels of COL4A5 and
COL4A6. From these results, we would predict that the dominant collagen IV heterotrimer in
epithelial basement membranes to be α5α5α6— whereas our data suggests it would be α1α1α2
within the basal lamina of the vasculature.

Integrins. The family of integrins was highly differentially expressed across the breast cell types.
Integrins connect the extracellular matrix (ECM) to the cytoskeleton and act as receptors that
integrate extracellular signals, relaying information to the cell's interior. Each integrin
heterodimer forms from one α and one β subunit, from families of 18 α and 8 β isoforms,
generating the 24 observed αβ combinations. When we investigated the family of integrins in the
breast cell types, we found they were all differentially expressed—except for ITGAD, which was
essentially absent in every cell type (Figure 15b- integrin beta subunits, Figure 15—figure
supplement 2d). Due to their diversity and cell-surface expression, sixteen integrins serve as
'cluster of differentiation' (CD) antigens. Above, we introduced that a critical marker in our
FACS was CD49f, which is integrin α6 (ITGA6) —and that myoepithelial and endothelial cell
types highly expressed it (Figure 4—figure supplement 2a, Figure 14b). Integrin a6 binds
laminins through heterodimers formed with either b1 or b4 subunits. The a6b1 and a6b4
integrins, along with a3b1 and a7b1, are the four main laminin-binding integrins. Thus, along
with α6 integrin, the main laminin-binding integrin heterodimers use a3 and a7 integrins.
Investigating these transcripts' expression in breast cells, we found that myoepithelial cells and
ER[Pos] luminal epithelial cells expressed the highest levels of ITGA3. In contrast, pericytes and
vSMCs expressed the highest levels of ITGA7, and fairly specifically, which is consistent with
prior reports of this subunit's expression in vascular smooth muscle cells[39] (Fig.15—figure
supplement 2d-e). These data suggest that myoepithelial and perivascular cells use distinct
integrin heterodimers to bind laminins within their respective basement membranes.

Another CD antigen is ITGAM (CD11b), which is frequently used to identify myeloid cells. We
indeed found that within the breast cell populations, ITGAM was highly expressed by
leukocytes—although it was also strongly expressed by adipocytes and luminal epithelial cells
(Fig.15—figure supplement 2d). Moreover, leukocytes also highly expressed six other integrins:
ITGAE, ITGAL, ITGAX, ITGA4, ITGB7, and ITGB2 (Fig.15—figure supplement 2d). ITGB7
and ITGB2 were exclusively expressed by leukocytes, consistent with their recognized role in
trafficking leukocytes to sites of injury and infection[40]. Our data reflect this cell-type specificity,
and transcript levels within the different breast cell types provide a detailed accounting of these
and other expressed integrin isoforms, offering valuable insights into potential dimer
combinations used by the different cell types.

Along with the collagens, alpha integrins, and 206 additional gene families, families containing
between 75 to 99% DE genes included the *plakins* (88% genes DE), *spectrins* (86%), and
receptor tyrosine kinases (80%). By comparison, we identified 209 families whose members
were expressed relatively equally across all breast cell types (0-24%), such as the *exportins* (0%,
Figure 15e). Furthermore, we also investigated families formed by a more moderate makeup of
DE genes, i.e., between 25 to 74% (Figure 3—table supplement 6). Among these 814 families
were the *semaphorins* (70%), *proteoglycans* (71%), *mucins* (71%), *prostaglandin receptors*
(67%), *nuclear hormone receptors* (66%), and *claudins* (64%, Figure 3—table supplement 6).
Also included were the *matrix metalloproteinases* (61% DE), which are a family of proteinases
that degrade all extracellular matrix components—and thus help regulate the homeostatic
balance of ECM molecules in the breast microenvironment (Figure 15d).

Matrix Metalloproteinases (MMPs). Of the 23 MMP isoforms encoded within the human
genome, we identified 14 differentially expressed among the breast cell types. Fibroblasts highly
expressed over half of these MMPs, which included gelatinase A (MMP2), stromelysin 1 and 2
(MMP3, MMP10), macrophage elastase (MMP12), membrane type 1,3, and 4 (MMP14,
MMP16, MMP17), and MMP19. Predominant substrates for these MMPs include fibronectin,
laminins, type III, IV, and V collagens, elastin, fibrin, pro-MMP2, pro-gelatinase A, aggrecan,
and cartilage oligomeric matrix protein[41]. In addition, we found leukocytes (and adipocytes)
expressed gelatinase B (MMP9) at its highest levels, consistent with reports of it being absent in
fibroblasts and playing an essential role in ECM proteolysis and leukocyte migration[42,43].
Arguably the most notable deviation from the near-universal fibroblast expression of MMPs was
the differential expression of matrilysin (MMP7) by ER^{Neg} luminal epithelial cells, where we
found levels to be 18-fold higher than in their ER^{Pos} counterparts –and 237-fold higher than in
fibroblasts (Figure 15d).

Tight Junctions. Essential to multicellular life are the mechanisms in which cells join to form
the fabric of tissues and guide their shape and functions. Above, we introduced the family of
integrins that serve as central receptors binding cells to the ECM. Different cell lineages have
distinct requirements. For example, breast fibroblasts are embedded within a collagen-rich
matrix, whereas perivascular cells, which are also mesenchymal in nature, encircle vessels, help
generate the endothelial basement membrane, and collaborate with endothelial cells to form the
vasculature. The endothelial cells, in turn, must create tight junctions to generate a patent and
sealed network for transporting liquid blood—especially important for forming the blood-brain

barrier of the central nervous system. Epithelial cells are also unique because their glandular role requires them to develop functional sheets while maintaining a semi-permeable barrier to the luminal space, where they may secrete and collect milk. The lumen is also essentially connected to the outside environment through the ductal network and nipple opening, so epithelial cells must also establish a barrier to protect against bacteria and other pathogens—working in concert with adjacent leukocytes that closely associate with the epithelium. Thus, the formation of tight junctions is essential to the gland.

Tight junctions are the prominent cell-to-cell junctions of epithelia and endothelia, establishing the selectively permeable barrier between neighboring cells[44]. Claudin and occludin proteins are vital components of this cell adhesion complex. Investigating their expression in the breast cells, we found the vast majority of claudins (11 of the 19 expressed) were predominantly limited to the two luminal epithelial cell types, although there were some notable exceptions (Figure 16a,b). One was the specific expression of claudins 5 and 14 (CLDN5, CLDN14) by endothelial cells—and to a lesser extent, by adipocytes. Whereas the vascular and lymphatic endothelial cells both highly expressed CLDN5, CLDN14-expression was limited to the vascular endothelial cells. Another exception was myoepithelial CLDN11 expression. Also, of the eleven claudins expressed by the luminal epithelial cells, five were differentially expressed between the two luminal types, with CLDN9, 22, and 23 transcripts being more highly expressed by the ER^{Pos} cells and CLDN16 and 10 being more highly expressed by the ER^{Neg} cells (Figure 16a,b). Furthermore, claudin 1 was one of the six claudins expressed by the luminal epithelial cells, and tissue staining confirmed luminal cell expression, as the staining was confined to the apical surface of the paracellular space, circumscribing the luminal surface (Figure 16—figure supplement 1). These data are consistent with the epithelial and endothelial use of tight junctions, and results pinpoint the differential isoform specificity among the five claudin-expressing cell types.

The other critical components of tight junctions are the occludins (tight junction proteins encoded by TJP1, TJP2, and TJP3 genes). When we explored their expression among the breast cell types, we found that TJP1 and TJP2 (encoding zona occludens -1 and -2) shared a similar pattern, as they were both expressed relatively equally among all cell types. The only exceptions were that for TJP1, leukocyte levels were roughly 50-fold lower than the other cell types, and for TJP2, levels were lower in both fibroblasts and leukocytes. In contrast, TJP3 expression (ZO-3) was predominantly limited to the luminal epithelial cells (Figure16c-e).

Gap Junctions. In contrast to the occluding tight junctions that seal the paracellular space between neighboring cells, gap junctions create channels between them, allowing the exchange of ions and low molecular weight metabolites between neighboring cells[45]. We found that the breast cell types expressed eleven of the 20 genes in the gap junction family (Figure 17 a,b). Myoepithelial cells expressed two of these genes (GJA1 and GJB5, i.e., connexin 43 and 31.1), whereas luminal epithelial cells expressed six (GJB2, B3, B4, B5, C3, and D3), sharing only with myoepithelial cells the expression of GJB5. Furthermore, ER^{Pos} and ER^{Neg} luminal cells differentially expressed three gap junction genes, with GJB3 and GJB4 being higher in ER^{Pos} cells—and GJB2 being highest in the ER^{Neg} luminal cells. Another intriguing feature was the fibroblast and endothelial expression of GJA1, which they shared with Pop4 epithelial and

myoepithelial cells. We also found that leukocytes uniquely expressed the highest levels of
GJB7, although we are uncertain what role it would fulfill in these cells. One final observation
was the perivascular expression of GJA4, GJA5, and GJC1; and the differential expression of
GJA5 by vSMCs, at a 5.5-fold higher level than pericytes (Figure 17 a,b). From these data, we
can identify the potential homomeric and heteromeric gap junction complexes used by the
perivascular and other cell types.

Whereas gap junctions facilitate the direct connection of cytoplasm between cell types and are
essential mediators of cell-to-cell communication, other forms of communication occur through
autocrine, paracrine, and endocrine signaling mechanisms. Multicellular organisms—and
tissues— rely on efficient and responsive communication between different cells, sometimes
separated by great distances. One of the most evolutionary-protected mechanisms facilitating cell
communication is the production of proteins, peptides, fatty acids, glycans, cholesterol
derivatives, gasses, and other metabolites by one cell, which are detected by another and elicit a
response[46]. Unfortunately, the vast number of ligand and receptor families preclude us from
exploring each of them here. Instead, we chose to inspect two different receptor families that
exemplify the heterogeneous expression patterns observed among breast cell types: *receptor*
tyrosine kinases and *nuclear hormone receptors*. The most studied and best-understood family of
receptors possessing inherent kinase activity are the receptor tyrosine kinases (RTKs).

Receptor Tyrosine Kinases. The human genome encodes 58 different receptor tyrosine kinases
(RTK) whose ligands span across twenty subfamilies. These ligands are wide-ranging and
include the ephrins, glial-cell derived neurotrophic factor, and: epidermal-, nerve-, hepatocyte-,
fibroblast-, vascular endothelial-, and insulin-like growth factors, among others. RTKs are vital
regulators of critical cellular processes and are frequently involved in many cancer types. Among
the breast cell types, most RTKs displayed a cell-type-specific expression pattern (Figure 18a-
m). Notably among these was the robust mesenchymal expression of platelet-derived growth
factors alpha and beta (PDGFRA, PDGFRB), fibroblast growth factor receptor 1 (FGFR1), AXL
tyrosine kinase receptor (AXL), and orphan receptor 2 (ROR2, Figure 18a-m). In contrast, most
cells equally expressed insulin receptor (INSR), insulin-like growth factor 1 receptor (IGF1R),
receptor-like tyrosine kinase (RYK), and lemur tyrosine kinase 2 (LMTK2). As expected, the
endothelial cells differentially expressed angiopoietin receptors tie2 and tie1 (TEK and TIE1)
and VEGF receptors 1 and 3 (FLT1 and FLT4), which are known to regulate many facets of
vascular biology (Figure 18a,i). We have previously described, consistent with prior reports[47-49],
how stem cell growth factor receptor (KIT) was highly expressed by ER[Neg] luminal cells[8], but we
also found it to be appreciably expressed by vascular endothelial cells and vSMCs (Figure 18a,
Figure 12—figure supplement 1). Her2 (ERBB2) is amplified in a subset of breast cancers and
has been successfully targeted for therapeutic intervention (Herceptin/trastuzumab). We found
that Her2, along with two other members of the ErbB family (ERBB3 and ERBB4), were highly
expressed by luminal epithelial cells. In contrast, the remaining family member, epidermal
growth factor receptor (EGFR), was expressed across a broader range of cell types, with
fibroblasts and myoepithelial cells displaying the highest transcript levels (Figure 18j-m).
Overall, RTKs display a complex pattern of expression. Most, 80%, were expressed

differentially across breast cell types, sometimes even within the same lineage, e.g., PDGFRB,
KIT, ERBB3.

Nuclear Hormone Receptors. Nuclear hormone receptors were another receptor family
demonstrating high cell type specificity. These receptors act as ligand-induced transcription
factors and are activated upon binding to small lipophilic molecules, such as thyroid or steroid
hormones, vitamin D3, retinoic acids (vitamin A), ecdysone, oxysterols, bile acids, fatty acids,
leukotrienes, and prostaglandins[50]. The human genome has 49 nuclear hormone receptors, and
their regulatory effects are wide-ranging. Of this family's many prominent members, the most
notable for the breast field is estrogen receptor (ERα).

Interestingly, much of what we know about nuclear hormone receptors is rooted in the early
studies of estrogen and how it regulates female reproductive organs, including the breast[51]. We
found 44 nuclear hormone receptors expressed in breast cells—31 differentially. The androgen,
progesterone, and estrogen α receptors were all strongly expressed by luminal epithelial cells—
more so in ER^{Pos} cells than their ER^{Neg} counterparts (Figure 19a-d). Both thyroid hormone
receptors were similarly expressed among the cell types, except perivascular cells, which
expressed 5.3-fold lower levels of THRA—and endothelial cells and leukocytes, which
expressed 8.5-fold lower levels of THRB (Figure 19e). The cell-type specific expression
extended to many other nuclear hormone receptors, such as vitamin D receptor (epithelial cell
types), estrogen-related receptor beta (ER^{Pos} luminal cells), hepatocyte nuclear factor 4 gamma
(both luminal cell types), estrogen receptor beta (Pop4 epithelial and vascular endothelial cells),
and nuclear receptor subfamily members –5A2 (both endothelial cell types), –4A1(mesenchymal
cells), –4A2 and –4A3 (perivascular cells, Figure 19f-m). Nuclear hormone receptors possess
broad influence over many cell processes. The cell-specific expression patterns detailed here
identify the cell types in which these receptors likely play a prominent role.

Analyzing gene families has provided us with an unconventional look at how different types of
genes contribute to the cell types in the breast, and the results identified the kinds of genes that
vary most across the spectrum of cell types. Above, we meticulously described nine gene
families whose functions extend from the nucleus (hormone receptors) to the exterior cell
microenvironment (laminins). However, these represent only a limited percentage of families we
identified as having a high proportion of highly variable genes. Many more are listed in the
supplementary tables (Figure 15—Table supplement 1, Figure 3—Table supplement 6). To
further understand how these different genes collectively contribute to the individuality of each
breast cell type, we needed to identify and evaluate the cellular pathways in which these genes
are collectively involved—and the higher-order processes they bestow.

Defining the nature of each breast cell type

To acquire a broad perspective of each breast cell type's fundamental properties and individual
functions, we performed a wholesale assessment of the biological pathways and processes
disproportionately employed by these different cell types. Gene set enrichment analysis
(GSEA)[52-54] accounted for each of the 20,000+ expressed genes in these cells' transcriptomes.
The algorithm determined whether a particular pathway's genes were enriched within a
bidirectionally ranked list of genes constructed from two contrasted cell types. One advantage of

GSEA is that it doesn't require large transcriptional variations to pinpoint enriched pathways, as
it can identify situations where many genes in a pathway differ in a small but coordinated
fashion.

Several databases organize functionally related genes into distinct pathways. Here, we present
our GSEA results that used gene sets defined by the Reactome Knowledgebase[18]. This database
contains a manually curated list of 2,585 pathways cross-referenced to other scholarly resources,
such as NCBI[55], ENSEMBL[56], KEGG[57], and GO[17,20]. Each Reactome pathway represents a
collection of genes whose products coordinate to perform a specialized biological process. To
determine which of these pathways are potentially active—or differentially enhanced—in a
particular cell type relative to another, we analyzed every transcript, in each breast cell type,
across every cell-cell comparison, which required 55 separate differential gene expression
contrasts (Figure 20, Figure 3—Table supplement 7 [1v2]).). For each pairwise comparison, we
calculated: a) the log fold change for each measured transcript, b) the p-value for the calculated
difference (Wald test), and c) the Benjamini-Hochberg adjusted p-value. Using these values, we
ranked all expressed genes bidirectionally, placing the most statistically significant genes at the
extreme ends of the list, with one cell type represented at the top (most positive) —and the other
at the bottom (most negative). After ranking the genes, we performed GSEA, which extracted
pathways from the Reactome database, identified the position of each pathway's genes within
our ranked list, and used a running sum algorithm to calculate a normalized enrichment score for
each pathway. For each pairwise comparison, hundreds of enriched pathways were typically
identified, each consisting of 15 to 200 genes.

To help us navigate the overwhelming amount of data generated from these combined analyses,
we integrated our GSEA results and gene expression values into a single pathway analysis tool
that we programmed within Microsoft Excel (Pathway Analyzer[9]). This facile tool efficiently
summarizes results according to user inputs (e.g., cell types to contrast, FDR cutoffs to use,
biological themes to investigate, specific pathways to examine, and genes to explore).

With the aid of the Pathway Analyzer tool, we found that the number of pathways enriched in
each cell type varied widely, ranging between 7 to 462 ($\bar{x} = 194, \text{sd} = 120$). On average,
fibroblasts had the fewest enriched pathways across all comparisons ($\bar{x} = 85 \pm 59$), whereas
those having the most were the vSMCs ($\bar{x} = 315 \pm 86$), pericytes ($\bar{x} = 272 \pm 121$), vascular
endothelial cells ($\bar{x} = 287 \pm 126$), and lymphatic endothelial cells ($\bar{x} = 229 \pm 114$). When we
compared cell types across different lineages, akin to the PCA analysis performed above, many
of the same enriched pathways emerged, e.g., keratinization within the epithelial cell types,
angiogenesis by endothelial cells, and more. Unfortunately, the magnitude and comprehensive
nature of the GSEA results preclude us from presenting here every identified aspect. Instead, we
will focus on the biology of closely related cell types since the broad features of different cell
lineages were examined above. However, the results from all analyses are accessible, and one
can thoroughly explore any of the 55 possible comparisons using the Pathway Analyzer tool.
Below, we present the most thought-provoking findings from six cell-type comparisons; these
are: a) the two luminal epithelial cell types, b) ER[Neg] luminal epithelial cells and myoepithelial
cells, c) ER[Neg] luminal epithelial cells and vascular endothelial cells, d) the two endothelial cell
types, e) fibroblasts and adipocytes, and finally f) fibroblasts and pericytes. Some enriched

pathways we anticipated, such as finding the enrichment of 'estrogen signaling' in ER[Pos] luminal
cells. These results validated the analytic approach and helped us establish confidence in the
pathway enrichments.

Estrogen signaling in luminal epithelial cells: a validating example

Luminal epithelial cells help shape the breast's glandular structure by forming the lumen of
mammary ducts. The breast's two major luminal cell types are commonly classified by their
divergent staining for estrogen receptor alpha (ERα), and we were able to isolate each of these
cell types based on their mutually exclusive staining for α6 integrin (ITGA6/CD49f; Figure1,
Figure 1—figure supplement 1)[8]. To determine the distinguishing characteristics of these two
cell types, we contrasted their expressed transcripts, exposing 1,301 differentially expressed
genes (BH/FDR adj. p-value < 0.1, Figure 21). ER[Pos] luminal cells expressed higher levels of 546
of these, including progesterone receptor (PGR), epiregulin (EREG), fibronectin (FN1), and
FOXA1. In contrast, transcripts differentially expressed by the ER[Neg] cells included epidermal
growth factor (EGF), matrix metallopeptidase 7 (MMP7), KIT, and 752 other genes (Figure 21,
Figure 3—table supplement 7 [1v2]). We ranked all expressed genes by their differential
expression (based on their fold-change and statistical significance) and performed pathway
enrichment analysis to identify the cellular functions impacted by these cell types' transcriptional
differences.

Gene set enrichment analysis (GSEA) assessed where each pathway's genes fell along our
ranked gene list, and the algorithm calculated a normalized enrichment score, nominal *p*-value,
and false discovery rate (FDR) for each Reactome pathway. The *estrogen-dependent gene*
expression set was among the top pathways enriched in the ER[Pos] luminal cells (*p*-value <0.001,
q-value =4.84×10^{-4}). Identifying this estrogen signaling pathway was validating and served as a
fine example of the enrichment algorithm (Pathway Analyzer [1v2]). Differentially expressed
genes contributing to this pathway's enrichment included progesterone receptor (PGR), FOXA1,
erb-b2 receptor tyrosine kinase 4 (ERBB4), GATA binding protein 3 (GATA3), estrogen
receptor a (ESR1), MYC, and others (Figure 22a). These genes were among the 65% in the
pathway that fell within the leading edge of the analysis (the leading-edge genes are the core
genes that contributed most to the GSEA enrichment score, Figure 22a). Furthermore, when we
analyzed transcripts from this estrogen-related pathway, we confirmed that the leading-edge
genes were indeed differentially expressed between the ER[Pos] and ER[Neg] luminal epithelial cells
(Figure 22b, Pathway Analyzer).

Apart from *estrogen-dependent gene expression,* we identified an additional 324 enriched
pathways between the two luminal cell types, 211 within the ER[Pos] cells and 113 within the
ER[Neg] cells (FDR ≤ 10%, Pathway Analyzer). We used a pathway enrichment and visualization
method to extract information rooted within these GSEA results (Enrichment Map[58]). This tool
analyzed the pathway's gene sets and, based on the proportion of genes they shared, formed a
complex network of interconnected pathways, placing redundant and biologically related
pathways close to each other. We connected pathways when they shared 50% or more genes, and
this caused distinct clusters, or 'biological themes,' to emerge in the network (Figure 23,
Supplemental Cytoscape file [Enrichment Map 1v2]). We circled these grouped pathways and

named their overarching theme using Reactome's hierarchy nomenclature as a guide. For
example, the *estrogen-dependent gene expression* gene set clustered with the two other sets
related to estrogen signaling: *estrogen-receptor mediated signaling* and *extra-nuclear estrogen*
signaling. We encircled all three on the map and labeled this overall theme 'Estrogen Signaling.'
(Figure 23). We adhered to this strategy for each theme in the network and often arranged related
themes together. For example, we placed the enriched 'FOXO mediated transcription' theme
adjacent to 'Estrogen Signaling' on the enrichment map because FOXO3 is an established ERα
regulator [59,60].

To evaluate the GSEA results and biological rationale for each pathway's enrichment, we studied
the genes from each pathway—and their expression levels in the different breast cell types.
These details often provided valuable insights (Pathway Analyzer). For example, high-scoring
genes within the *ER mediated signaling* pathway included: HSBP1, which binds to ERα and
promotes membrane localization; JUND, which interacts with ERα and occupies estrogen
response element sites within the promoters of target genes; progesterone receptor, which is a
direct downstream target of ERα; and others, like FOXA1, AKT1, epiregulin (EREG),
amphiregulin (AREG), and SRC proto-oncogene. These genes were all highly expressed by
ER^{Pos} luminal cells, and their transcript levels in these cells stood in stark contrast to those we
observed within ER^{Neg} luminal cells –and the same was observed for FOXO mediated
transcription (Figure 23—figure supplement 1a,b). In some cases, a pathway's genes were
uniquely expressed (or nearly so) by one of the contrasted cell types. We indicated these genes
on the enrichment map under each theme's name. For example, the genes encoding estrogen
receptor a (ESR1), amphiregulin growth factor (AREG), transcriptional factors FOXA1 and
MYB; tyrosine kinase receptor ErbB4, and the trefoil factor mucosal proteins TFF1 and TFF3
were all highly and differentially expressed by ER^{Pos} luminal cells (Figure 22b, Figure 23).
Collectively, these data provided strong evidence that estrogen signaling is active in this cell
type—as one would anticipate in cells expressing estrogen receptor.

Identifying the association between estrogen signaling pathways and ER^{Pos} cells was
validating—as was later finding additional anticipated themes in other cell-type contrasts.
Moreover, these correlations assured us that the pathway analysis had indeed identified processes
used disproportionately between cell types. Below, we move on from the detailed example of
estrogen signaling in these luminal cells and sift through the complexity and sheer volume of
remaining results.

ER^{Pos} vs. ER^{Neg} luminal epithelial cells (Pop1 vs. Pop2)

When we compared transcripts expressed by ER^{Pos} and ER^{Neg} luminal cell types (FACS
populations #1 and #2, Figure 1a), we identified 325 enriched pathways at a false discovery rate
of 10%. The majority of these pathways were associated with cell signaling (83 pathways),
metabolism (55 pathways), and immune-related functions (42 pathways). Others involved
extracellular matrix organization, developmental biology, nervous system development, cell
signaling, muscle contraction, and vesicle-mediated transport (Figure 23, Pathway Analyzer).
Further exploration of expressed transcripts defined by each pathway showed that the GSEA
algorithm functioned well at identifying significant, concordant differences in the gene levels

between cell types—even if individual transcript levels only modestly diverged. There were,
however, pathways that contained a significant share of highly differentially expressed genes—
some of these being principal players in the pathway. For example, fibronectin (FN1) and
TGFB1 are members of the *ECM proteoglycans* gene set and were expressed at over 35- and 10-
fold higher levels in ER^{Pos} luminal cells than their ER^{Neg} cell counterparts (Adj. *p*-value = $3.2x10^{-8}$
and $3.8x10^{-7}$, Figure 23, Figure 3—table supplement 7 [1v2]). Because implications of these
stark differences were often patently evident, we primarily focus on pathways that contain these,
but details of all enriched pathways can be thoroughly examined with the provided analysis tool
(Pathway Analyzer).

Cell Signaling & Signal Transduction (1v2). The PI3K/Akt, ErbB4, VEGF, and HGF/MET
pathways are all notable receptor tyrosine kinase pathways that we found enriched within ER^{Pos}
cells (Figure 23, Pathway Analyzer, Supplemental Cytoscape file [Enrichment Map 1v2]). Along
with p53, Phosphatidylinositol 3-kinase (PIK3CA) is the most mutated gene in breast tumors
(36% of tumors), and genes in the phosphoinositide 3-kinase (PI3K/AKT) pathway are the most
frequently altered in human cancers[61]. Across the span of normal breast cell types, we found
PIK3CA to be evenly expressed (Figure 23—figure supplement 1c). The PI3K/AKT pathway's
enrichment in ER^{Pos} luminal cells was instead primarily driven by the differential expression of
the serine-threonine kinase AKT1, the EGFR ligand epiregulin (EREG), and the receptor
tyrosine kinases: fibroblast growth factor receptor 1 (FGFR1) and platelet-derived growth factor
receptor beta (PDGFRB). Other genes contributing to the pathway's enrichment included the
EGFR receptors ERRB2, ERBB3, and ERBB4 (Figure 23—figure supplement 1c). These data
show that the ER^{Pos} luminal cells contain different signaling elements than ER^{Neg} cells and
possess the potential for stimulating themselves and neighboring cell types.

Another prominent member of the epidermal growth factor family of receptors is ERBB4. This
receptor dimerizes with ERα and is involved in many facets of downstream ER signaling[62]. Not
surprisingly, we found ER^{Pos} luminal cells enriched with the ERBB4 pathways: *signaling by*
ERBB4 and *nuclear signaling by ERBB4* (FDR=3.9% and 2.9%, Figure 23). This enrichment
was driven by the cell's prominent expression of ERBB4 and its ligands betacellulin (BTC) and
epiregulin (EREG), although ER^{Neg} luminal cells differentially expressed several pathway
members as well, including neuregulins 1 & 2 (NRG1, NRG2). In addition, glial fibrillary acidic
protein (GFAP), another pathway member, was also expressed—nearly exclusively—by ER^{Neg}
luminal cells. In CNS astrocytes (cells that express high levels of GFAP), GFAP promoter
activity is inhibited when bound by an erbb4-containing complex[63]. This inhibition perhaps
explains the reduced GFAP levels in ER^{Pos} cells, as these cells have high ERBB4 levels (Figure
23—figure supplement 1d).

Another theme emerging from the analysis of the two luminal cell types was VEGF signaling, as
we found the pathways *signaling by VEGF* and *VEGFR2 mediated vascular permeability*
enriched within ER^{Pos} luminal cells (FDR=0.17% and 4.2%, respectively). This result is,
however, an excellent example of how comprehensive information on all the breast cell types
provided substantial perspective. Between the two luminal cell types, the VEGF pathways were
indeed enriched in the ER^{Pos} cells—predominantly due to their differential expression of the
serine/threonine kinase AKT1, amyloid beta precursor like protein 2 (APLP2), neuropilin 1

(NRP1), and other pathway members. However, when we explored these transcript levels across all cell types, we found their expression in ER^{Pos} cells was not dramatically different from most other breast cell types (Figure 23—figure supplement 1e). It thus became clear that the pathway's enrichment was not a feature of its overexpression in ER^{Pos} cells but rather its relative repression in ER^{Neg} luminal epithelial cells (Pathway Analyzer). Because of the importance of these comparative insights, we similarly evaluated gene expression across all cell types for each pathway, using heat maps and the Pathway Analyzer tool.

Another pathway significantly enriched in ER^{Pos} luminal cells was HGF/MET. Hepatocyte growth factor (HGF) was first identified as a fibroblast-secreted factor that caused dramatic epithelial cell migration[64]. Its cognate receptor is MET receptor tyrosine kinase[65]. We found three HGF/MET signaling pathways enriched within ER^{Pos} luminal cells, including *signaling by MET* (FDR=0.6%, Figure 23, Pathway Analyzer). Enrichment was mainly driven by the differential expression of fibronectin, integrin alpha 3 (ITGA3), and several laminin subunits (LAMA1, -A5, and -B2). These proteins are involved in focal adhesion signaling that acts through focal adhesion kinase (PTK2), which was also modestly elevated in ER^{Pos} cells (Figure 23—figure supplement 1f). Interestingly, when we examined HGF expression across all cell types, we found, consistent with its initial discovery, that it was indeed highly expressed by breast fibroblasts, although pericytes and adipocytes also expressed it. HGF's receptor, MET, was expressed by vascular endothelial cells, adipocytes, and the breast's four epithelial cell types (Figure 22—figure supplement 1f).

Metabolism (1v2). To evaluate metabolism-related pathways, we coded a filter in the Pathway Analyzer tool, allowing us to extract pathways involved in specific biological themes. At a 10% FDR threshold, we identified fifty-five enriched pathways related to metabolism—thirty within ER^{Pos} luminal cells and twenty-five within ER^{Neg} luminal cells. Those related to energy metabolism were enriched in ER^{Pos} luminal cells (Figure 23). Notable among these were: *integration of energy metabolism* (FDR=0.550%), *glycolysis* (FDR=7.94%), *regulation of insulin secretion* (FDR=5.51%), *PKA activation in glucagon signaling* (FDR=4.87%), and *glucuronidation* (FDR=3.25%, Pathway Analyzer). Genes contributing to these pathways' enrichment included: the inositol 1,4,5-trisphosphate receptor (ITPR1), which is a PKA-activated calcium channel that acts downstream of glucagon and cAMP signaling; fatty acid synthase (FASN), which catalyzes the synthesis of palmitic acid from acetyl-CoA and malonyl-CoA; and cAMP-dependent protein kinase (PRKAR2B), which transduces cAMP signals (Figure 23—figure supplement 1g). Elevated transcript levels of several glycolytic and regulatory enzymes within ER^{Pos} cells also led to the enrichment of the core glycolysis pathway. These were: hexokinase-1 (HK1), phosphofructokinase (PFKP), the bifunctional enzyme 6-phosphofructo-2-Kinase/Fructose-2,6-Biphosphatase 1 (PFKFB1), enolase gamma (ENO2), and pyruvate kinase M1/2 (PKM). These enzymes catalyze the first, last, and an intermediate step of glycolysis (HK1, PKM, ENO2) –and regulate other key regulatory metabolites in the pathway (PFKFB1),Figure 23, Figure 23—figure supplement 1h, Pathway Analyzer). Thus, significant evidence suggests ER^{Pos} cells have increased bioenergetic demands. As such, it was not surprising that we also found enrichment of pathways involved in lipid metabolism.

Metabolism of phospholipids, sphingolipids, and fatty acids were three themes that emerged
from the comparison of the two luminal cell types (Figure 23). The ER^{Pos} cells' disproportional
expression of many of these pathways' genes led to the enrichment *sphingolipid metabolism*
(FDR=0.04%), *phosphoinositol metabolism* (FDR=0.06%), *fatty acid metabolism* (FDR=9.53%),
and related pathways (Pathway Analyzer). Analysis of the leading-edge genes from these
pathways demonstrated their elevated expression in the ER^{Pos} cells (Figure 23—figure
supplement 1i-k). The most notable genes involved in phospholipid metabolism included the
solute carriers SLC44A4 and SLC44A5. These membrane-embedded choline transporters
provide a critical metabolic precursor for phosphatidylcholine synthesis. ER^{Pos} cells also highly
expressed pleckstrin homology domain protein (PLEKHA6) and phosphatidylinositol 4,5-
bisphosphate 5-phosphatase A (INPP5J). PLEKHA6 interacts with phosphatidylinositol 3,4,5-
trisphosphate (PIP3), whereas INPP5J dephosphorylates PIP3. Fatty acid elongase (ELOVL5)
and fatty acid synthase (FASN) are both involved in synthesizing fatty acids, were highly ranked,
and contributed to the pathways' enrichment.

The set of pathways involving protein metabolism covers the range of reactions that proteins
encounter over their lifespan, from synthesis to degradation (including protein folding, post-
translational modifications, and trafficking). We found both luminal cell types enriched with
distinct protein metabolic pathways (Figure 23). In ER^{Pos} cells, two of the eleven significant
pathways were *N-glycan antennae elongation in the medial trans golgi* (FDR=0.79%) and
sumoylation of intracellular receptors (FDR=0.95%, Pathway Analyzer). Differentially
expressed genes in the former pathway included carbohydrate (N-acetylgalactosamine 4-0)
sulfotransferases (CHST8, CHST10), ST8 alpha-N-acetyl-neuraminide alpha-2,8-
sialyltransferases (ST8SIA2, ST8SIA6), and mannosyl (alpha-1,3-)-glycoprotein beta-1,4-N-
acetylglucosaminyltransferase isoenzymes (MGAT4A, MGAT4B). These enzymes are active in
the medial Golgi, where N-glycan remodeling occurs. When we explored the second pathway,
sumoylation of *intracellular receptors,* we found that the small ubiquitin-like modifiers
(SUMO1, 2, 3) were surprisingly expressed evenly across cell types. The pathway's enrichment
was instead driven by elevated expression of the retinoic acid, retinoid X, progesterone,
androgen, and estrogen receptors, indicating the receptors that could be modified by sumoylation
(Pathway Analyzer).

The protein metabolism pathways enriched in ER^{Neg} cells included *amyloid fiber formation*
(FDR=5.95%) and *O-linked glycosylation of mucins* (FDR=9.28%, Pathway Analyzer). We also
observed that many, but not all, large ribosomal protein components were elevated in ER^{Neg}
cells, leading to the enrichment of ribosome-involved pathways, *eukaryotic translation initiation*
(FDR≤0.001%) and *eukaryotic translation elongation* (FDR≤0.001%). Historically, these genes
have been considered invariant, but more recent evidence suggests otherwise, that levels may be
linked to cell specification[66]—as we observed among the breast cell types here.

Extracellular Matrix Organization (1v2). The extracellular matrix (ECM) is a dynamic
structure consisting of a heterogeneous composition of scaffolding proteins, growth factors,
proteoglycans, and remodeling enzymes. It undergoes persistent remodeling, primarily through
reciprocal interactions with nearby cells in the immediate microenvironment. We found eleven
pathways enriched between the two luminal cell types, with ER^{NEG} cells overexpressing genes

related to *collagen synthesis* and assembly (FDR=1.58%), whereas the ER^Pos^ cells were enriched in degradative pathways *degradation of the extracellular matrix* and *collagen degradation* (FDR 4.08% and 5.55%, Figure 23, Pathway Analyzer). Driving enrichment of the synthetic pathway in ER^Neg^ cells were collagens (COL4A2, 6A2, 6A1, 22A1), lysyl oxidases (LOX, LOXL4), and matrix metallopeptidases (MMP3, MMP7). In contrast, the degradative pathway in ER^Pos^ cells was driven by differential expression of tissue inhibitor of metalloproteinase 2 (TIMP2), calpains (CAPN8, CAPN2, CAPN13), ADAM metallopeptidase 10 (ADAM10), cathepsin L (CTSL), elastin (ELN), matrix metalloproteinases (MMP12, MMP13), and collagens (COL7A1, 12A1, 4A5, 4A6). These data suggest that these two related—but distinct— luminal cell types have contrasting roles in the synthesis and degradation of specific ECM proteins (Figure 23—figure supplement 1 l,m).

Additional enriched pathways in ER^Pos^ cell relating to the ECM included: *laminin interactions* (FDR=4.1%), *non-integrin ECM interactions* (FDR=0.18%), *integrin cell-surface interactions* (FDR=0.47%), *syndecan interactions* (FDR=6.47%), and *ECM proteoglycans* (FDR=0.31%; Figure 23—figure supplement 1 n-r). Differentially expressed genes within these pathways included collagen IV isoforms A5 and A6 (COL4A5, COL4A6), laminin chains (LAMA5, LAMB2), fibronectin (FN1), transforming growth factor-beta -1 and -2 (TGFB1, TGFB2), platelet-derived growth factor beta (PDGFB), agrin (AGRN), synedecan 3 (SDC3), and integrins α5 and β3 (ITGA5, ITGB3), all of which had higher transcript levels in ER^Pos^ luminal cells compared to their ER^Neg^ counterparts.

Tight junctions (1v2). Claudins are a family of genes we explored in the previous section, where we found, with few exceptions, that their transcription was mainly limited to endothelial and epithelial cells—predominantly the two luminal cell types (Figure 16a,b). Here, using GSEA, we identified significant differences between the ER^Pos^ and ER^Neg^ luminal cells (FDR = 9.1%). Specifically, the ER^Neg^ cells expressed significantly higher levels of CLDN10, CLDN16, CLDN11, CLDN8, and CLDN1, which led to enrichment of the *tight junction interactions* pathway (Figure 23, Figure 16a). This was notably the only Reactome 'Cell-cell Communication' pathway enriched in either cell type (other pathways, for example, involve *adherens junctions*, *cell-ECM interactions*, and *hemidesmosome assembly*, etc.). Despite the elevated levels of five claudins in ER^Neg^ cells, CLDNs 1, 7, 3, and 12 were expressed at roughly the same level in both cell types, whereas ER^Pos^ cells had higher levels of CLDNs 23, 6, and 9. Thus, the enrichment indicated that the two luminal cell types possess unique tight junction components to support their individual physiologies. These tight junctions are critical to the cells' ability to form semi-permeable barriers, which helps retain fluid and protect against microorganisms in the outside environment –to which the ductal system is connected via the nipple opening.

Immune System (1v2).

Using GSEA, we identified forty-two enriched pathways related to immune function. The vast majority of these were in the ER^Neg^ luminal cells (33 pathways), which overlapped with the two innate immune system pathways previously identified by the PCA/overrepresentation analysis (Figure 10a). One of the larger themes was interleukin signaling, where we found the enrichment

of nine pathways related to: *interleukin-4 and interleukin-13 signaling, interleukin-10 signaling,*
interleukin-1 family signaling, interleukin-2 family signaling, and *interleukin-12 family*
signaling (FDR= 0.181%, 0.792%, 2.33%, 4.79%, 1.096%, respectively; Figure 23; Pathway
Analyzer). Enrichment of these pathways was driven by the ER[Neg] cell's differential expression
of the interleukins 33, 1a, and 18 (IL33, IL1A, IL18), chemokine ligands 4 and 5 (CCL4, CCL5),
tumor necrosis factor receptor TNFRSF1B, serum amyloid-A1 (SAA1), interleukin-2 receptor
gamma (IL2RG), interleukin-1 receptor type II (IL1R2), interleukin-27 receptor alpha (IL27RA),
colony-stimulated factor 2RB (CSF2RB), and lipocalin 2 (LCN2, Figure 23- figure supplement
1s,t; figure supplement 2a-c). We notably found breast macrophages concentrated near the
epithelial structures and within the ducts and epithelial compartment[8], and thus were not
surprised to find luminal cell enrichment of interleukin pathways that involve macrophage
activation. This indicates that the ER[Neg] cells are disproportionately responsible for regulating
macrophage physiology –a crosstalk that is subverted in breast tumors[67].

Among the other enriched immune-related pathways, we discovered ER[Neg] luminal cells
expressed higher levels of genes involved in *toll-like receptor cascades* and *antimicrobial*
peptides (FDR=8.208%, 4.1067%, respectively). The ER[Neg] cells differentially expressed toll-
like receptors 1 and 2 (TLR1 and TLR2), defensins B4A and B1 (DEFB4A and DEFB1),
peptidase inhibitor 3 (PI3), and clusterin (CLU, Figure 23- figure supplement 2d,e). Furthermore,
all ten toll-like receptors were highly expressed by luminal cells, with half being the highest in
these cells compared to all other cell types (TLR1, TLR2, TLR2, TLR5, and TLR6, Figure 23—
figure supplement 2d ; Figure 23—figure supplement 3a-f). Toll-like receptors play a vital role
in the innate immune system by triggering an immune response to microbial agents. For
example, TLR1 and TLR2 coordinate to mediate the response to bacterial lipoproteins. Although
the toll-like receptors are well studied in the colonic epithelium, their contributions to the
homeostasis of the breast epithelium are not as well characterized. Because luminal epithelial
cells have the potential for elevated microorganism exposure, it is reasonable to expect TLR and
antimicrobial peptide expression, as they are needed to provide innate immune protection.

Milk proteins (1v2). Although not part of the Reactome gene sets, we noticed several genes
encoding prominent milk proteins were highly differentially expressed by ER[Neg] luminal cells,
such as lactotransferrin (LTF) and casein kappa (CSN3). Lactotransferrin is the major iron-
binding protein in milk, whereas casein kappa is critical for stabilizing casein micelle formation
in milk. To explore the cellular expression of other milk constituents, we created a list of genes
from prior reports of human milk composition [68-70]. This list comprised 48 genes and included
lactotransferrin, caseins, lactalbumin, and other milk proteins (Figure 23—figure supplement 4a).
Most of these were expressed by at least one breast cell type (41/48), and nearly all were
differentially expressed (37/41; BH/FDR adj. p-value < 0.05), with one of the two luminal cells
most often being the cell type with the highest transcript level (25/37). After comparing
expression between the luminal cell types, we found 12 genes whose expression significantly
differed between the two (adj.*p*-val <0.05), with ER[Neg] cells most often having the higher
transcript levels (8/12). These eight genes were: lactotransferrin (LTF), secreted phosphoprotein
1 (SPP1), polymeric immunoglobulin receptor (PIGR), xanthine dehydrogenase (XDH), clusterin
(CLU), complement component 3 (C3), haptoglobin (HP), and milk fat globule-EGF factor 8

protein (MFGE8, Figure 23—figure supplement 4b-j). These results demonstrate that the luminal cells are primed for milk production despite being derived from the resting breast tissues of non-lactating women and, at rest, transcribe the genes encoding major milk proteins–with the highest levels often being in the ER^{Neg} luminal cells.

ER^{Neg} luminal epithelial vs. Myoepithelial cells (Pop2 vs. Pop3)

Whereas the luminal cells shape the lumen of the gland's ductal structure, myoepithelial cells form the basal layer, anchoring the epithelium to the basal lamina (Figure 9 a,d,e,g,h). The ER^{Neg} luminal and myoepithelial cells (FACS populations #2 and #3) are two of the most abundant cell types in the breast (Figure 1); they share the same epithelial lineage and spatial proximity—and physically interact. Most clinical breast tumors have a luminal phenotype and presumably originate from luminal cells, with the ER^{Neg} luminal cells being the most-likely cell of origin for basal-like breast tumors[49,71], despite these tumors' basal-like features. For these reasons, we were interested in—and explored— the differences between ER^{Neg} luminal and myoepithelial cells, focusing on the biochemical and physiological processes distinguishing these two cell types.

Analysis of transcripts expressed by the myoepithelial and ER^{Neg} luminal cells revealed 2,239 differentially expressed genes (BH/FDR adj. p-value < 0.1, Figure 24, Figure 3—table supplement 7 [2v3]). ER^{Neg} luminal cells expressed the higher level of roughly half of these genes (51.6%), which included CD24, keratin 19, prominin 1 (PROM1), epidermal growth factor (EGF), and peptidase inhibitor 3 (PI3). Myoepithelial cells, in contrast, differentially expressed 1,083 genes; among these were genes previously associated with this cell type, including smooth muscle actin (ACTA2), keratin 5 (KRT5), p63 (TP63), and oxytocin receptor (OXTR, Figure 24). Using the p-values from this differential expression analysis, we created a ranked list of all expressed genes and used this list to perform pathway enrichment analysis. GSEA revealed enrichment of 426 Reactome pathways (FDR≤10%), which clustered into distinct biological themes (Figure 25). The most significant of these involved cell-cell communication, extracellular matrix organization, and metabolism (Pathway Analyzer).

Cell-cell communication (2v3). Proper intercellular communication is essential for maintaining homeostatic balance and normal tissue function. When we contrasted the ER^{Neg} luminal and myoepithelial cells, we found enrichment of four cell-cell communication pathways: one in the ER^{Neg} luminal cells (tight junctions) and three in the myoepithelial cells (cell-junctions, cell communication, and ECM interactions). Given the previous identification of claudin expression in the luminal epithelial cells, it was not surprising to find the *tight junction interactions* pathway enriched in the ER^{Neg} luminal cells, as they—in comparison to myoepithelial cells— differentially expressed most epithelial claudins, such as CLDN1, 3, 4, 7, 8, 10, 12, and 16 (FDR =5.57%). The only significant exception was the myoepithelial expression of CLDN 11 (Figure 16 a,b). These results show that luminal cells express significantly more claudins than myoepithelial cells and thus rely on the tight junction features that these different claudins provide.

Although luminal cells differentially expressed claudins and were enriched with the *tight junction interactions* pathway, we found other cell communication pathways enriched in myoepithelial cells. These were the *cell-cell communication* pathway (FDR = 0.0919%) and its

underlying pathways: *cell junction organization* (FDR =0.0076%) and *cell-extracellular matrix*
interactions (FDR =0.1682%). Notable genes shared among these three pathways include actinin
and filamin proteins that promote the branching of actin filaments and anchor the filaments to
proteins in the cell's membrane. Shared in the enriched pathways were also the cell adhesion
molecule 3 (CADM3), keratins 5 and 14 (KRT5, KRT14), and cadherins 3, 4, and 11 (CDH3,
CDH4, CDH11, Figure 25—figure supplement 1a). As we found above with the PCA-derived
gene list for myoepithelial cells (myoepithelial and Pop4 epithelial cells), a general feature of
myoepithelial cells is their expression of ECM and basement membrane components and
formation of desmosomes involved in cell junction organization (Figure 14 a-c). Thus, we were
not surprised to find myoepithelial enrichment of the *cell-extracellular matrix interactions*
pathway because these cells form and help shape the epithelial basement membrane. For the
same reason, we anticipated other pathways involved in extracellular matrix organization would
be enriched within myoepithelial cells. We found this was indeed the case.

ECM organization (2v3). As the non-cellular component of tissues, the extracellular matrix is
formed by a network of proteins, proteoglycans, growth factors, and minerals that provides
structural support, elasticity, and tensile and compressive strength to the tissue[72]. The ECM and
basement membrane composition is dynamic and is constantly remodeled by cells in their
immediate vicinity. Notably, we found pathways associated with extracellular matrix
organization as some of the most significantly enriched pathways in myoepithelial cells, as nine
of the ten most statistically significant pathways in the myoepithelial cells involved ECM and
basement membrane organization (Pathway analyzer). Myoepithelial cells were enriched in
nearly every aspect of ECM organization, including *Elastic fiber formation* (FDR= 0.01%),
Laminin interactions (FDR<0.001%), *non-integrin membrane ECM interactions* (FDR<0.001%),
ECM Proteoglycans (FDR<0.001%); *activation of matrix metalloproteinases* (FDR=4.38%),
collagen formation and *collagen degradation* pathways (FDR<0.001%). These cells
differentially expressed many collagen and laminin encoding genes, including COL2A1, 4A5,
4A6, 8A2, 11A1, and 17A1; and laminin chains LAMA1, LAMA3, and LAMB1). Other genes
contributing to these pathways' enrichment included elastin microfibril interfacer 1 (EMILIN1),
metalloproteinase 3 (MMP3), tissue inhibitor of metalloproteinases 2 (TIMP2), actinin alpha
1(ACTN1), thrombospondin 1 (THBS1), fibromodulin (FMOD), heparan sulfate proteoglycan 2
(HSPG2), and syndecan 3 (SDC3), among others (Figure 25- figure supplement 1b-e). The
myoepithelial cells—which reside on the basement membrane at the edge of the epithelial
compartment— are thus a major contributor and remodeler of the basement membrane and
ECM. This is a significant role that diverges from that of the luminal epithelial cells. However,
the luminal cells were a major producer of matrilysin (MMP7) that, in addition to several ECM
proteins, cleaves the milk protein casein, which we also found differentially expressed by ER[Neg]
luminal cells.

Muscle Contraction (2v3). A defining feature of myoepithelial cells has long been their
expression of smooth muscle actin[73], which allows them to contract when stimulated by
oxytocin, e.g., during lactation. It was thus reassuring to find *muscle contraction* (FDR =
0.0085%) and associated pathways enriched in myoepithelial cells, including *smooth muscle*
contraction, phase 0 rapid depolarization, and *cardiac conduction—*the latter sharing similar

genes despite being a pathway in the heart (FDR= 0.0707%, 2.011%, 0.148% respectively, Figure 25, Pathway Analyzer). Myoepithelial cells highly expressed smooth muscle actins alpha-1 and gamma-2 (ACTA1, ACTG2), the voltage-gated sodium channel alpha subunit 3 (SCN3A), the calcium voltage-gated channel subunit L (CACNA1C), myosin heavy chain 11 (MYH11), myosin light chain kinase (MYLK), and leiomodin 1 (LMOD1, Figure 25- figure supplement 1f) that contribute to the contractile ability of this cell type.

Ion channel transport (2v3). Calcium is essential for myoepithelial contractility; Its intracellular availability is regulated by the transport of ions between the extracellular and intracellular spaces. This process was significantly enriched in myoepithelial cells, as we found *ion channel transport* and associated pathways enriched in myoepithelial cells (FDR=5.99%, Figure 25). The enrichment was driven predominantly by these cell's differential expression of anoctamin-1 (ANO1), chloride channel accessory-2 (CLCA2), and the intracellular calcium channels ryanodine receptors 1 and 3 (RYR1, RYR3, Figure 25- figure supplement 1g). Furthermore, myoepithelial cells differentially expressed the transient receptor potential (TRP) cation channels TRPV6, TRPC1, TRPC4, TRPM3, TRPC6, and TRPV2 (*TRP channels,* FDR=1.59%)*.* TRP channels are calcium permeable and convert external stimuli into electrical or chemical signals[74] (Figure 25- figure supplement 1g). These pathways likely contribute to regulating myoepithelial cells' intracellular hyperpolarization/depolarization and contractility.

Cell signaling (2v3). We identified the enrichment of eighty-one cell signaling pathways between the ER[Neg] and myoepithelial cells. Most of these (sixty-two) were enriched within the myoepithelial cells. The bulk of these pathways involved receptor tyrosine kinases (23/62), G-protein coupled receptors (13/62), Notch (9/62), Wnt (8/62), and Hedgehog pathways (6/62). Although the ER[Neg] luminal cells differentially expressed KIT and were enriched with both *signaling by SCF Kit* (FDR=5.7%) and ERBB2 signaling pathways, the myoepithelial cells were enriched with twenty receptor tyrosine kinase pathways, most at a conservative false discovery rate of less than 1% (13/20, Pathway Analyzer). These pathways involved MET, neurotrophic tyrosine receptor kinases, platelet-derived growth factor, vascular endothelial growth factor, fibroblast growth factor, and insulin-like growth factor pathways (Figure 25). For example, the enrichment of the FGFR signaling pathways *signaling by FGFR, signaling by FGFR1,* and *signaling by FGFR2* (FDR= 0.328%, 0.822%, and 4.00%,) was driven by their differential expression of fibroblast growth factors 1, 2, and 7 (FGF1, FGF2, FGF7); FGF receptors 1, 2, and 4 (FGFR1, FGFR2, FGFR4); and the ras proto-oncogene (HRAS, Figure 25- figure supplement 1h).

The enrichment of Notch signaling pathways split between the luminal and myoepithelial cells. *Signaling by notch* and *signaling by notch4* were enriched in myoepithelial cells (FDR= 0.0268%, 0.059%), as these cells differentially expressed smooth muscle actin (ACTA2) and notch ligands, JAG1 and JAG2 (Figure 25- figure supplement 1i). They also had slightly elevated levels of many proteosome subunits that help compose this pathway. In contrast, ER[Neg] luminal cells were enriched with Notch 1, 2, and 3 signaling pathways (FDR= 3.38%, 7.75%, 1.99%) due to their differential expression of the Notch receptors, NOTCH2 and NOTCH3; the downstream effector HEY2, delta/notch-like EGF repeat containing (DNER) and epidermal growth factor (EGF), and other pathway members (Figure 25- figure supplement 1j).

Other myoepithelial enriched pathways involved G-protein coupled receptor-, Wnt-, and
Hedgehog-signaling. Differentially expressed genes from the twelve enriched G-protein coupled
receptor pathways included G-protein coupled receptor 176 (GPR176), dopamine receptor (DR),
and adenylate cyclase 5 (ADCY5). Enrichment of the Wnt and Hedgehog signaling pathways
was driven by elevated transcript levels of these pathways' downstream effectors, which
included transcription factor 7 (TCF7), lymphoid enhancer-binding factor-1 (LEF1), GLI1, and
GLI3 (Pathway Analyzer).

Metabolism (2v3). Within the eighty-seven metabolism-related pathways enriched between
luminal and myoepithelial cells, one of the larger emergent themes involved the metabolism of
glycosaminoglycans (Figure 25). These pathways were enriched in myoepithelial cells and
included *glycosaminoglycan metabolism, chondroitin sulfate dermatan sulfate metabolism,* and
heparan sulfate heparin HS-GAG metabolism pathways (FDR= 3.76%, 5.50%, and 3.04%,
Figure 25). Among the genes contributing to their enrichment included key enzymes in GAG
metabolism, e.g., a) hyaluronan synthase 3 (HAS3), which catalyzes the addition of
acetylglucosamine and glucuronic acid monosaccharides to hyaluronan polymers during
hyaluronic acid synthesis; b) chondroitin-6 sulfotransferase-3 (CHST3), which catalyzes the
sulfation of chondroitin; and c) the heparan sulfate proteoglycan glypican-3 (GPC3), an ECM
glycoprotein that binds to, and modulates, several signaling pathways (Figure 25—figure
supplement 1k).

A characteristic feature of epithelial cells is their secretion and release of mucins from their
apical surfaces[75]. Not surprisingly, we found the *O-linked glycosylation of mucins* pathway
enriched in ER^{Neg} luminal epithelial cells (Figure 25). Luminal cells notably expressed mucin-1
(MUC1), which is a marker in our FACS panel and was used to sort the luminal cells. We found
many other mucins expressed by the luminal cells, including MUC15 and MUC16 (Figure 25-
figure supplement 2 a-f). Mucins are highly glycosylated proteins and a component of milk,
where they provide antimicrobial resistance in the GI tract of breastfed infants[76]. These mucins
are derived from the luminal epithelial cells.

The *metabolism of steroids* was another large theme in the network, which was enriched within
ER^{Neg} luminal cells (FDR=0.998%, Figure 25, Pathway Analyzer). These cells differentially
expressed hydroxysteroid dehydrogenases 17B2 and 17B4 (HSD17B2 and HSD17B4)—
essential enzymes involved in the production of the androgen and estrogen sex hormones[77].
Additionally, luminal cells highly expressed other steroid hormone regulators, HSD11B1 and
steroid 5-alpha-reductase (SRD5A3), as well as the cytochrome p450s (CYP24A1 and
CYP24B1), which function in Vitamin D metabolism (Figure 25- figure supplement 1l). We
found these genes were highly expressed by both luminal cell types, indicating steroid
metabolism is a general feature of the luminal lineage.

The ER^{Neg} luminal cells were also enriched with other metabolic pathways, such as *fatty acid*
metabolism (FDR=6.04%, Figure 25, Pathway Analyzer), as they differentially expressed acyl
synthetase long-chain family members 1, 3, 5, and 6 (ACSL1, ACSL3, ACSL5, ACSL6), aldo-
keto reductase family member C3 (AKR1C3), glutathione peroxidase 2 (GPX2), and carnitine O-
acetyltransferase (CRAT, Figure 25- figure supplement 1m). The acyl synthetase enzymes

catalyze long chain fatty acids into active fatty acyl-CoA esters, which is a critical step in beta-
oxidation and lipid biosynthesis. The roughly 10-fold difference in acyl synthetase transcripts
between luminal and myoepithelial cells indicates that luminal cells have different metabolic
demands requiring the processing of long-chain fatty acids.

In our analysis of the citric acid cycle pathway, we found most genes in this pathway were
uniformly expressed across the entire spectrum of cell types. However, there were three notable
exceptions. The first, which occurred in both luminal cell types, was their pronounced
differential expression of both isoforms of isocitrate dehydrogenase (IDH1, IDH2), which
operate in the mitochondria to regulate 2-hydroxyglutarate, alpha-ketoglutarate, and isocitrate
(adj. p-values = 5.2e-5 and 2.1e-13). Another differentially expressed TCA cycle gene was
pyruvate decarboxylase (PC), which was repressed in both endothelial cell types. Finally, we
found phosphoenolpyruvate carboxykinase-1 (PCK1) expression was limited to pericytes and
adipocytes (Figure 25- figure supplement 2f).

ER^{Neg} *luminal epithelial vs. Vascular endothelial cells (Pop2 vs. Pop9)*

Histologists consider endothelial cells specialized epithelial cells because they exhibit classical
epithelial features[16]. For example, vascular endothelial cells and epithelial cells both form the
lumen of their respective structures (e.g., vessels, ducts), use tight junctions for cell-cell
adhesion, and rest on a basal lamina. However, when we analyzed the breast cell transcriptomes
by PCA (Figure 7a) we found that endothelial cells projected more closely to mesenchymal cell
types than to the epithelial cells, which may reflect their mesodermal origin. We found this
curious and decided to compare the features of vascular endothelial cells to the abundant ER^{Neg}
luminal epithelial cell type (FACS populations #9 and #2).

Comparing transcripts between vascular endothelial cells and ER^{Neg} luminal cells, we identified
4,167 differentially expressed genes (Figure 26). Of those, 1,652 were differentially expressed by
the endothelial cells, which included endothelial markers: platelet/endothelial cell adhesion
molecule (PECAM1/CD31), CD93, TIE1, and TEK tyrosine kinase. In contrast, luminal cells
differentially expressed the transcriptional regulator, SMARCD3, keratin 15 (KRT15), CD24,
cadherin 1 (CDH1), and transcriptional repressor GATA binding-1 (TRPS1, Figure 26, Figure
3—table supplement 7 [2v9]). We ranked all expressed genes by their degree of differential
expression and used this list for pathway analysis. GSEA revealed five-hundred pathways
enriched between ER^{Neg} luminal and vascular endothelial cells (FDR≤10%), which provided
insights into the different biological pathways and systems used by each cell type (Pathway
Analyzer).

Hemostasis (2v9). One of the most significantly enriched pathways, *cell surface interactions at*
the vascular wall (FDR<0.001%), was found in vascular endothelial cells—cells that indeed
form the vascular wall (Figure 27). Endothelial cells differentially expressed selectins E, L, and
P (SELE, SELL, SELP)—which mediate cell-to-cell adhesion; endothelial cell adhesion
molecule (ESAM); junctional adhesion molecule 2 (JAM3); platelet/endothelial cell adhesion
molecule (PECAM1); and the angiopoietin receptor TEK tyrosine kinase (TEK, Figure 27-
figure supplement 1a). These gene products regulate leukocyte adhesion to vessel walls during
an inflammatory response and facilitate leukocyte migration. In addition, endothelial cells

differentially expressed von Willebrand factor (VWF)— a historical marker of endothelial cells, thrombomodulin (THBD), and coagulation factors F2R and F3 (Figure 27- figure supplement 1b). These contributed to the enrichment of *formation of fibrin clot (clotting cascade,* FDR = 3.04%). This is an essential step for blood clotting after injury, indicating that endothelial cells supply critical factors for this process, thus distinguishing them from other epithelial cell types. Notably, several of these pathway's genes are shared by mesenchymal perivascular cells (e.g., ITGA5, ESAM, ANGPT); this shared expression contributes to the mesenchymal alignment of endothelial cells in the PCA projection (Figure 7a).

Keratinization (2v9). Another divergent feature of endothelial and epithelial cells was the keratinization that occurred in ER^{Neg} luminal cells (*keratinization,* FDR<0.001%, Figure 27, Pathway Analyzer). Keratin filaments are heteropolymers of different Type I and II keratins that confer distinct biophysical properties to cells, connecting the cells' cytoskeleton to other cells and the basement membrane via desmosomes and hemidesmosomes[78]. We found ER^{Neg} luminal cells differentially expressed fifteen keratins, including keratins 7, 8, 19, 6b, 23, 81, 15, 16, 17, and 18. In addition, they also differentially expressed caspase-14 (CASP14)—which is involved in the cornification of epithelia— and desmoglein-1 (DSG1), a component of desmosomes (Figure 27- figure supplement 1c).

Cell Communication (2v9). Among the pathways associated with cell communication, we found that ER^{Neg} luminal cells were, compared to the vascular endothelial cells, enriched with the broad *cell-cell communication* pathway (FDR= 0.954%) and others involving tight junctions (Figure 27- figure supplement 1d). These were *cell junction organization* (FDR=1.8%) and *tight junction interactions* pathways (FDR= 0.835%, Figure 27- figure supplement 1e). The enrichment resulted from ER^{Neg} luminal cells' differential expression of laminin beta 3 (LAMB3), cadherins 1 and 3 (CDH1, CDH3), as well as the crumbs-cell-polarity complex component 3 (CRB3), which helps form tight junctions. Although the vascular endothelial cell types expressed claudins 5 and 14 rather exclusively, the ER^{Neg} luminal cells differentially expressed an overwhelming number of claudins (CLDN1, -3, -4, -7, -8, -10, -11, and -12), leading to the pathway's luminal cell enrichment (Figure 27- figure supplement 1e, Figure 16). Another cell-communication pathway is the *cell-extracellular matrix interactions* pathway, which we found enriched in vascular endothelial cells (FDR<0.001%). This pathway includes integrins that link ECM proteins to the cell's cytoskeleton –with the help of scaffolding and focal adhesion proteins. Genes in this pathway differentially expressed by the vascular endothelial cells included filamin C (FLNC)—which was notably repressed by both luminal cell types (compared to all others), integrin-linked kinase (ILK), and parvin beta (PARVB, Figure 27- figure supplement 1f, Pathway Analyzer [2v9]).

Metabolism (2v9). We identified eighty-two enriched pathways that involve the metabolism of nucleic acids, lipids, proteins, and other metabolites. Some of the more statistically significant pathways within ER^{Neg} luminal cells included the *metabolism of vitamins and cofactors* (FDR= 0.618%), *metabolism of water-soluble vitamins and cofactors* (FDR=0.71%)*,* and *nicotinamide salvaging* (FDR=4.63%, Figure 27, Pathway Analyzer). Genes contributing to their enrichment included a) vanins 1 and 2 (VNN1 and VNN2), which are involved in vitamin B5 metabolism, b) transcobalamin 1 (TCN1)—a vitamin 12 binding protein, and c) pyridoxal kinase (PDXK),

which phosphorylates vitamin B6 (Figure 27- figure supplement 1g). We also found *sphingolipid metabolism* enriched in luminal cells (FDR= 4.37%), as they differentially expressed sphingosine-1-phosphate phosphatase 2 (SGPP2), sphingomyelin synthase 1 (SGMS1), and aldehyde dehydrogenase 3 family member B2 (ALDH3B2, Figure 27- figure supplement 1h). Additionally, *fatty acid metabolism* was enriched in ER[Neg] luminal cells (FDR= 0.636). Enrichment resulted from the luminal cells' differential expression of epoxide hydrolase 2 (EPHX2), ELOVL fatty acid elongase 6 (ELOVL6), acyl-CoA synthetase long-chain family members 1, 3, and 6 (ACSL1, ACSL3, ACSL6); fatty acid-2 hydrolase (FA2H); and the mitochondrial isocitrate dehydrogenase IDH2 (Figure 27- figure supplement 1i). *Glycosaminoglycan metabolism* was another enriched pathway that contained substantial transcriptional differences (FDR=0.95%, Figure 27, Pathway Analyzer). Leading to its enrichment in vascular endothelial cells was their differential expression of carbohydrate sulfotransferase 1 (CHST1), versican (VCAN), biglycan (BGN), decorin (DCN), and heparan sulfate proteoglycan 2 (HSPG2/perlecan, Figure 27- figure supplement 1j). This is notable because endothelial surfaces and their surrounding ECM contain large quantities of heparan sulfate proteoglycans[79] that modulate vascular functions, such as sequestering growth factors (FGF, VEGF), regulating lipoprotein metabolism, and inhibiting clotting[80].

Immune system (2v9). Pathways associated with the immune system were enriched in both luminal and endothelial cells (Figure 27). Nine were enriched in luminal cells and 39 in endothelial cells. *Antimicrobial peptides* (FDR=0.058%), *defensins* (FDR=0.349%), and *beta-defensins* (FDR=1.19%) were enriched in luminal cells, which differentially expressed defensin beta 1 (DEFB1), S100 calcium-binding proteins A7A, A7, A8, and A9 (S100A7A, S100A7, S100A8, and S100A9), toll-like receptors 1 and 2 (TLR1 and TLR2), lipocalin 2 (LCN2), and bactericidal/permeability-increasing protein (BPI, Figure 27- figure supplement 1k). Luminal cells also differentially expressed protein tyrosine phosphatase 2 and 6 (PTPN1 and PTPN6), interferon-inducible guanylate binding protein 1, 2, 3, and 5 (GBP1, GBP2, GBP3, and GBP5), and interferon regulatory factor 6 (IRF6)—which define the enrichment of *interferon-gamma signaling* in luminal cells (Figure 27, Figure 27- figure supplement 1l).

Interferon signaling was enriched in vascular endothelial cells, which differentially expressed intercellular adhesion molecule 1 (ICAM1) and interferon alpha-inducible protein 27 (IFI27, FDR= 0.512%, Figure 27, Figure 27- figure supplement 1m, Pathway Analyzer). ICAM1 encodes a receptor for leukocyte adhesion protein LFA-1 (ITGAL) and is essential for leukocyte trans-endothelial migration[81]. Additionally, *toll-like receptor cascades* (FDR=1.36%) were enriched in vascular endothelial cells, that differentially expressed suppressor of cytokine signaling (SOCS1), interleukin receptor-associated kinase 2 (IRAK2), and TLR4 (Figure 27, Figure 27- figure supplement 1n). Two pathways associated with TLR cascades were enriched in luminal cells: *IRAK4 deficiency TLR2/4* and *regulation of TLR by endogenous ligand* (FDR=0.926% and 4.94%, respectively, Figure 27). Notably, we discovered that enrichment of many pathways associated immune system was due to the expression of proteasome subunits. For example, enrichment of *interferon signaling*, *TCR signaling, BCR signaling,* and *Interleukin-1 signaling* were all affected by overexpression of the proteosome subunits PSMD2, PSMB6, and PSMB5, among others (FDRs < 0.001%, Figure 27, Figure 27- figure supplement 1o-q).

Proteosomes function in peptide processing for many pathways, including those involved in antigenic peptide presentation during immune responses. However, protein modification is required for many other pathways like cell cycle progression, apoptosis, and DNA repair.

Cell signaling (2v9). As expected, *signaling by VEGF* and *VEGFR2 mediated vascular permeability* were enriched in vascular endothelial cells (FDR=0.005% and 6.81%, respectively), with differential expression of placental growth factor (PGF), neuropilin 1 and 2 (NRP1 and NRP2), nitric oxide synthase 3 (NOS3), and vascular endothelial growth factors B and C (VEGFB and VEGFC)—all of which are involved in VEGF-regulated angiogenesis (Figure 27, Figure 27- figure supplement 1r)

Interestingly, *signaling by ERBB2* was enriched in endothelial cells (FDR=8.00%, Figure 27). The differentially expressed genes contributing to the enrichment of this pathway include FYN proto-oncogene (FYN), YES proto-oncogene (YES1), and heparin-binding EGF-like growth factor (HBEGF, Figure 27, figure supplement 1s). However, the luminal cells differentially expressed ERBB2, ERBB3, ERBB4, phosphoinositide-3-kinase regulatory subunit 1 (PIK3R1), epidermal growth factor (EGF), and EGF receptor (EGFR), which define the enrichment of other pathways associated with ERBB2 signaling, like *PI3K events in ERBB2 signaling* and *ERBB2 regulates cell motility* (Figure 27).

One other surprising finding was, despite the higher levels of estrogen receptor α mRNA in ER[Neg] luminal cells (so named because of their undetectable protein expression and lower mRNA levels compared to the ER[Pos] luminal cells), we found *estrogen receptor-mediated signaling* and associated pathways enriched in vascular endothelial cells (FDR = 0.008%). After investigating other members of this pathway, we found vascular endothelial cells differentially expressed estrogen receptor β (ESR2), calmodulin-1 (CALM1), and the G protein subunits GNAI2, GNG2, GNG5, GNG11, GNB1, and GNB2, and endothelial nitric oxide synthetase (NOS3) that compose this pathway. (Figure 27, Figure 27, figure supplement 1t). This is noteworthy because endothelial cells respond to estrogen for nitric oxide and prostacyclin production[82]. Our data thus indicate that the bulk of estrogen's effects on breast endothelium likely results through ERβ signaling.

Lymphatic endothelial vs. Vascular endothelial cells (Pop7 vs. Pop9)

Whereas the blood's vascular system circulates blood throughout the body, the lymphatic system collects and filters interstitial fluid, ultimately returning it to the bloodstream—a process that helps maintain fluid homeostasis. Much like capillaries of the blood system, lymphatic vessels are also formed by endothelial cells, although the two vascular architectures are distinct (Figure 2 d,f). We found lymphatic and vascular endothelial cells expressed similar markers, like CD31 (PECAM1) and von-Willebrand factor (VWF, Figure 8b) but diverge and expressed other genes that contribute to their individual functions. To explore the biological processes shaping these two endothelial cell types, we investigated their transcriptional differences.

Lymphatic endothelial cells are a rare cell type in the breast. Because of this, acquiring enough cells required for mRNA-sequencing proved challenging. However, by scaling up the number of cells stained—and through sheer brute force (~15 hours of continuous FACS sorting for each sample), we succeeded in purifying a sufficient number of cells from two samples (24,713 cells

from N228 and 29,111 cells from N239, Figure 3—table supplement 1). We detected roughly 21
million and 43 million paired end reads, respectively from these purified lymphatic endothelial
cells (N228 and N239). During quality control assessment of the RNA-sequencing data, we
however found outliers and conflicts between the lymphatic cell replicates (Figure 4c).
Nonetheless, the pair of lymphatic endothelial cell types clustered together well— in the
hierarchical clustering dendrogram (Figure 6) and projection of principal components (Figure
7a). We thus retained the lymphatic endothelial samples for further analyses, yet, we did expect
to find discordance between replicates for some genes.

When we compared transcript levels between lymphatic and vascular endothelial cell types, we
identified 3,739 differentially expressed genes (Figure 28, Figure 3—table supplement 7 [7v9]).
Lymphatic endothelial cells expressed the higher level of 1,462 of these, including podoplanin
(PDPN), the marker we used to differentiate lymphatic cells in our FACS strategy (adj.p-value of
1.14E-7). We also found other lymphatic markers differentially expressed, such as 'lymphatic
vessel endothelial hyaluronan receptor1' (LYVE1), 'vascular endothelial growth factor receptor-
3' (FLT4), and prospero homeobox 1 (PROX1, Figure 28). In contrast, the vascular endothelial
cells differentially expressed 2,277 genes. Among these were the vascular endothelial growth
factor C (VEGFC), cytokine-like 1 (CYTL1), lipocalin 1 (LCN1), endothelial selectin (SELE),
and the endothelial marker CD93 (Figure 28). After ranking the expressed genes by degree of
differential expression, we performed GSEA, and discovered 240 pathways enriched between the
lymphatic and endothelial cell types (FDR≤10%, Pathway Analyzer). Of these enriched
pathways, we've concentrated only on those established from genes where there was
concordance between lymphatic endothelial cell replicates.

Immune system (7v9). Both lymphatic and blood vascular systems are essential and active
components of immune responses. Lymphatic vessels serve as conduits for lymphocyte
migration from peripheral tissues to the lymph nodes, making them crucial to adaptive immune
response. In contrast, vascular vessels are first responders to pathogen detection and danger-
signal-sensing mechanisms in the bloodstream. Indeed, when we inspected the enrichment
results, both cell types were enriched with immunity-related pathways. For instance, we found
interleukin-4 and interleukin-13 signaling pathways enriched in lymphatic endothelial cells
(FDR= 1.54%). Contributing to this enrichment was the lymphatic cells' differential expression
of transforming growth factor-beta 1 (TGFB1) and IL13 receptors A1 and A2 (IL13RA1 and
IL13RA2, Figure 29a). In contrast, the vascular endothelial cells were enriched with *the TCR*
signaling and *interleukin-1 family signaling pathways* (FDR= 0.009% and 0.394%, Pathway
Analyzer). The TCR signaling pathway's most highly differentially expressed genes included
major histocompatibility complexes HLA-DRB1 and HLA-DQA1, vasodilatory-stimulated
phosphoprotein (VASP), and the TCR subunit CD247 (Figure 29b), whereas differentially
expressed genes of the IL-1 signaling pathway included HMGB1, which encodes the 'high
mobility group box 1' protein that promotes inflammatory responses to infectious signals, and
protein tyrosine phosphatase 7 (PTPN7) which regulates TCR signaling (Figure 29c).

Other immune-related genes that we identified as being differentially expressed by lymphatic
endothelial cells but were not associated with the above pathways included MRC1, TLR4,
CCL21, and ACKR2 (Figure 29—figure supplement d-f, Figure 23- figure supplement 3g).

MRC1 encodes mannose receptor 1, a cognate binding partner of L-selectin—that is expressed by lymphocytes and facilitates their binding to endothelial cells. Toll-like receptor 4 (TLR4) plays a crucial role in mediating the innate immune response to bacterial lipopolysaccharide (LPS), where it engages other levers of the immune response upon binding to bacterial LPS[83]. chemokine ligand 21 (CCL21) mediates homing of lymphocytes to lymph organs, whereas the atypical chemokine receptor 2 (ACKR2) regulates cellular interactions with inflammatory leukocytes[84]. Interestingly, we found lymphatic and vascular endothelial cells both expressed the pattern recognition receptor NOD1, which detects bacterial peptidoglycan fragments and initiates immune responses to gram-negative bacteria[85] (Figure 29g). There is thus ample evidence showing the respective endothelial cell types are interacting with cells of the immune system and actively participating in immune surveillance –in distince ways.

ECM Organization (7v9). While vascular endothelial cells are surrounded by a continuous basement membrane, the basement membrane surrounding lymphatic endothelial cells is thin and discontinuous (Figure 2d, Figure 2—figure supplement 2)[8]. We thus expected to find differences in their respective expression of extracellular matrix components. After performing pathway analysis, we indeed found enrichment of *ECM proteoglycans* in the lymphatic cells (FDR=1.09%). Contributing to the pathway's enrichment was the strong lymphatic expression of all three TGFβ ligands (TGFB1, -B2 and -B3), fibromodulin (FMOD), biglycan (BGN), fibronectin (FN1), tenascin XB (TNXB), integrin alpha 8 (ITGA8), collagen 6A2 (COL6A2), and laminin beta 2 (LAMB2, Figure 29h). Elastic fibers appear to be strongly associated with lymphatics, as we found the *elastic fiber formation* pathway highly enriched in this cell type (FDR=0.5%). Twenty-four genes in the elastic fiber pathway were elevated in lymphatic endothelial cells, including 'elastin microfibril interfacer 1' (EMILIN1) and latent transforming growth factor beta (LTBP1). In a prior section, we introduced the fibulin gene family, where we discussed their importance to elastic fibers and identified several family members (EFEMP1, FBLN1, FBLN5) differentially expressed by lymphatic endothelial cells –these genes are also members of the *elastic fiber formation* pathway (Figure 29i).

The only ECM-related pathway enriched in vascular endothelial cells was c*ollagen biosynthesis and modifying enzymes* (FDR = 6.65%). Differentially expressed genes contributing to its enrichment included the collagens COL9A3, COL4A1, COL4A2, COL18A1, COL21A1, COL13A1, COL8A1, COL15A1; procollagen-lysine 2-oxoglutarate 5-dioxygenase 1 (PLOD1); and collagen beta (1-O) galactosyltransferase 1 (COLGALT1, Figure 29j). The type IV collagens are a major structural component of basement membranes, indicating that endothelial cells collaborate with pericytes to produce the vascular basement membrane.

Cell signaling (7v9). VEGF signaling is a critical regulator of vascular development. Interestingly, we found *signaling by VEGF* enriched in lymphatic endothelial cells (FDR= 0.563%) due to their elevated expression of 55 of the 105 genes in the pathway, despite the vascular endothelial cells' differential expression of several key pathway members, like vascular endothelial growth factor c (VEGFC), placental growth factor (PGF), and VEGF Receptor-1 (VEGFR1/FLT1). The lymphatic endothelial cells, in contrast, expressed higher levels of other pathway members, like: 'p21 (Cdc42/Rac)-activating protein 3' (PAK3), cytochrome b245 (CYBA), the mTORC2 target PRR5, and VEGFR3 (FLT4, Figure 29k).

Distinct Notch signaling pathways were enriched in both endothelial cell types. We found the
broader *Signaling by notch* pathway enriched in vascular endothelial cells (FDR=0.434%),
whereas the *signaling by notch2* pathway was enriched in lymphatic endothelial cells—at a false
discovery rate of 9.97% (near the 10% cut-off). Of the 174 genes in the *signaling by notch*
pathway, 103 were differentially expressed by vascular endothelial cells, including genes
encoding the Notch ligands jagged 1 (JAG1), jagged 2 (JAG2), and delta-like 4 (DLL4, Figure
29l). Notch2 transcript levels were higher in the lymphatic cells, which, compared to all other
breast cell types, appeared to stem from its repression within vascular endothelial cells (Figure
29m). This contributed to *signaling by notch2* pathway's enrichment in lymphatics, along with
robust lymphatic expression of other pathway members, like ADAM metallopeptidase domain
10 (ADAM10) and mastermind-like transcription coactivator (MAML2).

Muscle contraction (7v9). We found the *muscle contraction* pathway enriched in vascular
endothelial cells (FDR= 4.30%, Pathway Analyzer). However, both endothelial cell types did not
appreciably express skeletal actin (ACTA1) or smooth muscle actins (ACTA2 or ACTG2), so
they do not appear to be contractile. Of course, this is consistent with contractile smooth muscle
cells being the primary regulator of vessel diameter. Instead, we found the vascular endothelial
cells' enrichment of the *muscle contraction* pathway primarily resulted from the differential
expression of a) the calcium-permeable transient receptor potential cation channel (TRPC1), b)
the calcium-dependent membrane protein annexin A6 (ANXA6), and c) several potassium
channels, including KCNE3, KCNK5, and KCNK6. These intermediate and small conductance
channels play an essential role by regulating the release of endothelial cell autacoids, like nitric
oxide[86], and were noticeably absent in lymphatic endothelial cells (Figure 29n).

Fibroblasts (ADMSCs) vs. Adipocytes (Pop8 vs. Pop12)

As epithelial cells form the parenchyma of the breast, the bulk of the tissue stroma is formed by
the mesenchymal cell types (adipocytes, fibroblasts, vascular smooth muscle cells, and
pericytes), with endothelial cells and a spectrum of leukocytes providing key elements[8]. We did
not quantify adipocytes by FACS, but we could deduce from the RNA-sequencing data that
adipocytes are one of, if not the most, abundant breast cell type. This was inferred from the
covariance analysis and hierarchical clustering that grouped adipocytes with the whole tissue
control samples, indicating they were disproportionately comprised of adipocytes (Figure 4b and
Figure 6). The remaining cell types were assessed by FACS, and it came as no surprise that
fibroblasts were also abundant (Figure 1b).

Fibroblasts are spindle-shaped cells that predominantly provides the tissue's foundation—a
collagenous-rich matrix of fibrillar extracellular proteins, which fibroblasts shape through the
secretion of growth factors, proteoglycans, and matrix-bound proteases. Fibroblasts are the
archetypical mesenchymal cell type, existing within a developmental lineage that is shared with
adipocytes. In our adjoining article, we described how breast fibroblasts express mesenchymal
stem cell markers (CD10, CD34, CD73, Thy1, CD44, CD105) and displayed a functional multi-
lineage potential when cultured under osteogenic or adipogenic conditions[8]. For these reasons,
we designated Pop8 fibroblasts as adipocyte-derived mesenchymal stem cells (ADMSCs) but
commonly refer to them as fibroblasts. In our cultures, and especially within the tissue,

adipocytes appeared as large, spherical cells containing a single large lipid droplet (unilocular adipocytes) that stained for neutral lipids[8] (Figure 2a,c; Figure 2—figure supplement 1a, 2a). Adipocytes notably serve as the body's chief storage depot of triglycerides and other acylglycerols and regulate energy metabolism through secretion of adipokines[87], e.g, leptin hormone (LEP) and adipsin (CFD), which we found highly expressed in the pop12 lipid (adipocyte) samples (Figure 31a,b). To establish the different cellular processes shaping adipocytes and fibroblasts, we examined their expressed transcripts.

In contrast to the other breast cell types, freshly-isolated adipocytes were not FACS purified prior to RNA-sequencing, and therefore results should be interpreted differently. Adipocytes were instead isolated from the phase-separated lipid layer of the collagenase tissues digests, which we had visually inspected for contaminating cellular content (Hoechst staining). The number of paired-end mRNA fragments measured from the RNA isolated from the lipid layers was excellent (34 to 45 million fragments), and the replicates nicely clustered together (Figure 4b; Figure 6; Figure 7—figure supplement 2a, Figure 8a). However, one sample did stray in the projection of PC1 and PC2 (Figure 7a), indicating some discordance between samples. We also noticed, in all three adipocyte replicates, low transcript levels for many genes (e.g., this can be observed in the many heatmaps presented herein, such as those containing the claudins and gap junction proteins, Figures 16,17). These adipocyte samples thus have a higher noise level than that observed in the FACS-purified samples, which is to be expected given the purity and excellent quality of those samples. The noise is likely derived from a fraction of contaminating cells in the samples, such as leukocytes. Therefore, we have prioritized genes and pathways in our analyses that are not heavily associated with immune response or where there is not evidence of robust adipocyte mRNA levels. With the caveat that the adipocyte samples should be interpreted differently, we explored the nature of adipocytes by comparing them with fibroblasts.

Pairwise comparison of RNA transcripts from adipocytes and fibroblasts revealed 1,171 differentially expressed genes. This was notably the lowest number of differentially expressed genes between any cell-type comparison, similar to the number differentially expressed between the two luminal epithelial cell types (1,301, figure 20). We anticipated fewer DE genes given the similarity and shared lineage of adipocytes and fibroblasts. Of the 1,171 differentially expressed genes we identified, 207 were more highly expressed by fibroblasts, and 964 were more highly expressed by adipocytes (Figure 30, Figure 3—table supplement 7 [8v12]).

When we investigated the identity of the differentially expressed genes, we found fibroblasts expressed the ECM protein biglycan, delta-like non-canonical Notch ligand 1 (DLK1), matrix metalloproteinase 3 (MMP3), and others. In contrast, adipocytes differentially expressed leptin (LEP), thyroid hormone-responsive protein (THRSP), and glycerol-3-phosphate dehydrogenase (GPD1), along with 961 other genes (Figure 30). After performing GSEA, we identified 387 pathways enriched between the two cell types (FDR $\leq$ 20%), with 69 pathways in fibroblasts and 318 in adipocytes (Pathway Analyzer). Due to the large number of of immune-related genes and pathways in adipocyte samples, we increased the false discovery rate cutoff to 20% for this comparison, which resulted in the enrichment of more pathways related to fat storage and metabolism.

Extracellular matrix organization (8v12). Twelve pathways associated with ECM organization were enriched in fibroblasts, including *degradation of the extracellular matrix*, *ECM proteoglycans*, and *elastic fiber formation* (FDR< 0.001%, < 0.001%, and =0.090%, respectively, Pathway Analyzer). Fibroblasts differentially expressed genes contributing to these pathways, including TIMP metallopeptidase inhibitor 1 and 2 (TIMP1 and TIMP2); metallopeptidases 2, 3, 13, 14, and 16 (MMP2, MMP3, MMP13, MMP14, and MMP16); fibulin-1 (FN1); fibrillin-1 (FBN1); and elastin (ELN, Figure 31c-e). There were two pathways associated with ECM organization enriched in adipocytes, including *laminin interactions* and *anchoring fibril formation* (FDR=3.23% and 9.97%, Figure 31f). Adipocytes differentially expressed integrins A6, B4, A3, and A2; laminins B3, A3, and C3; and collagens 4A5 and 4A6, which contributed to the enrichment of these pathways (Figure 31f).

Metabolism (8v12). Included in the enriched metabolic pathways was *Glycosaminoglycan metabolism* that was associated with fibroblasts (FDR<0.001%). This pathway's genes included heparanase-2 (HPSE2), hyaluronan synthase-2 (HAS2), and glypican-3 (GPC3, Figure 31g). In contrast, we found adipocytes enriched in pathways associated with lipid metabolism, that included *triglyceride metabolism, triglyceride catabolism*, and *arachidonic acid metabolism* (FDR= 15.5%, 15.5%, and 4.13%, Pathway Analyzer, Figure 31h). Differential expression of fatty acid binding proteins (FABP4, FABP5, FABP7, and FABP9) were among those genes contributing to enrichment of pathways associated with triglyceride metabolism and catabolism (Figure 31h). These intracellular fatty acid-binding proteins (FABPs) exhibit high affinity for a single long-chain fatty acids and are essential for triglyceride storage; they mediate intracellular fatty acid transport and interact with PPAR nuclear receptors[88]. Enrichment of *arachidonic acid metabolism* in adipocytes was defined by differential expression of arachidonate 5-lipoxygenase (ALOX5) and its activating protein (ALOX5AP, Figure 31i), which catalyzes the first two steps of the biosynthesis of leukotrienes, which are potent mediators of inflammation.

We also found adipocytes uniquely expressed cell death inducing DFFA like effector C (CIDEC, Figure 31j), which binds to lipid droplets and regulates their enlargement, thereby restricting lipolysis and favoring storage[89]. The adipocytes also uniquely expressed aquaporin AQP7 and the aquaporin pseudogenes AQP7P1 and AQP7P3 (Figure 31k-m), which control water and glycerol transport and promote the catabolism of lipids in adipocytes[90].

Cell signaling (8v12). Accumulation of adipose tissue is regulated in part by estrogen signaling[91]. Therefore, it is unsurprising that *ESR-mediated signaling* and its associated pathways were enriched in adipocytes (FDR= 0.110%), which differentially express the contributing genes GATA binding protein 3 (GATA3), caveolin 2 (CAV2), epidermal growth factor (EGF), and epiregulin (EREG, Figure 31n). Adipocytes also expressed estrogen receptors 1 and 2 (ESR1 and ESR2), albeit at 10x lower levels lower than ER[Pos] luminal epithelial cells but at a 4.3x higher level than fibroblasts (Figure31n, Pathway Analyzer).

Pericytes vs. Fibroblasts (ADMSCs, Pop6 vs. Pop8)

Whereas mesenchymal adipocytes contribute to the storage and mobilization of energy-rich metabolites; and mesenchymal fibroblasts secrete ECM proteins and enzymes that shape the extracellular stromal environment, the mesenchymal perivascular cells influence and modulate

blood vessels. In a companion article[8], we described how we identified vascular smooth muscle
cells and pericytes while devising methods to isolate and culture all breast cell types, finding
these perivascular cells to be an elusive set of cell types. Pericytes have long been
overlooked[92,93], largely stemming from their heterogeneity that, in turn, required a combinatoric
assessment of markers to identify and purify them. After discovering how to reliably isolate and
propagate breast pericytes, we found their microscopic appearance fascinating, as they had a
stellate astrocyte-like morphology that was entirely distinct from fibroblasts and adipocytes[8].
However, it was difficult to tell the pericytes and fibroblasts apart at higher confluency. We were
thus interested in exploring the biological processes defining these two similar mesenchymal cell
types (i.e., FACS populations #6-pericytes and #8-ADMSC/fibroblasts).

Analysis of transcripts expressed by freshly isolated pericytes and fibroblasts revealed 2,932
differentially expressed genes (BH/FDR adj. p-value < 0.1, Figure 32). Interestingly, these were
not split evenly between the two cell types, as fibroblasts had roughly three times the number of
DE genes than pericytes. Among these 2,187 genes, were platelet-derived growth factor receptor
alpha (PDGFRA), matrix metallopeptidase 2 (MMP2), and hydroxyacid oxidase 2 (HAO2). In
contrast, pericytes differentially expressed 748 genes, including pyruvate dehydrogenase kinase
4 (PDK4), CD36, and desmin (DES, Figure 32). Along with desmin, we found the pericytes
differentially expressed other genes previously associated with pericytes, such as melanoma cell
adhesion molecule (MCAM), G protein receptor RGS5, and the potassium channel KCNJ8. We
also found pericytes uniquely expressed the serotonin receptor isoform HTR1F, which was
distinct from the HTR1B isoform we discovered uniquely expressed by vascular smooth muscle
cells that, coincidently, helped us identify the vSMC FACS population[8] (serotonin is a
vasoactive peptide[94]). We ranked all expressed genes by their degree of differential expression
and used this list for pathway analysis.

We discovered 477 pathways enriched between pericytes and fibroblasts at a false discovery rate
of 10%. We found the enrichment of many pathways resulted from small concordant expression
differences between the cell types, such as those involved in transcriptional regulation, cell cycle,
and immunity. Here, we focus on pathways containing significantly differentially expressed
genes.

Muscle Contraction (6v8). As one would predict, we found perivascular pericytes enriched with
pathways involved with *muscle contraction* and *smooth muscle contraction* (FDR =0.361% and
0.026%, respectively, Figure 33). Pericytes differentially expressed myosin light chain kinase
(MYLK), myosin heavy chain 11 (MYH11), actin gamma 2 (ACTG2), and the voltage-gated
sodium channel type IV (SCN4A). Also differentially expressed by pericytes was dystrophin
(DMD), which anchors ECM proteins to cytoplasmic filamentous actin (and whose defects lead
to Duchenne and Becker muscular dystrophies). Tropomodulin 1 (TMOD1) was also highly
expressed by pericytes, which is a protein that regulates tropomyosin via blocking the elongation
and depolymerization of actin filaments (Figure 33- figure supplement 1a). Interestingly, *cardiac*
conduction was enriched in fibroblasts (FDR=5.64%), which differentially expressed T-box 5
(TBX5) transcription factor; potassium channels KCNK1, KCNK2, KCNIP1, and KCNIP4; and
the sodium channel SCN1B (Figure 33, Figure 33- figure supplement 1b). Fibroblasts share
similar expression of voltage-gated channels to cardiac myocytes, which are responsible for

cardiac conduction in the heart. Additionally, the dramatic difference in gene expression could be due to the low expression of many of these channels in pericytes, which may instead rely on endothelial cells for depolarization signals (as we found similar channels were differentially expressed by vascular endothelial cells when compared to their lymphatic counterparts, Figure 29n).

Hemostasis (6v8). We found fourteen pathways associated with hemostasis enriched in pericytes, including *platelet homeostasis* (FDR= 0.732%, Figure 33). Its enrichment was defined by differential expression of phosphodiesterases 1A, 1B, and 5A (PDE1A, PDE1B, and PDE5A), transient receptor cation channel C6 (TRPC6), cGMP-dependent protein kinase 1 (PRKG1), and the prostaglandin receptor PTGIR (Figure 33- figure supplement 1c). The phosphodiesterases are responsible for hydrolyzing cGMP and modulating the effects of nitric oxide signaling. Also related to the vasculature was the enrichment, in pericytes, of *cell surface interactions at the vascular wall* (FDR=1.14%), resulting from their differential expression of MER proto-oncogene (MER), thrombomodulin (THBD), and angiopoietin 1 (ANGTP1, Figure 33, Figure 33- figure supplement 1d).

Intracellular Transport (6v8). Pericytes differentially expressed kinesins (KIF4A, KIF2A, and KIF26B) and tubulins (TUBB2A, 3, and 4B), leading to their enrichment of *kinesins* pathway (FDR<0.001%, Figure 33, Figure 33- figure supplement 1e). *Clathrin-mediated endocytosis* was also enriched in pericytes (FDR<0.001%), due to their differential expression of the endocytic adapter protein SGIP1, WNT5A ligand, the wnt receptor 'frizzled class receptor 4' (FZD4), angiotensin II receptor (AGTR1), and the endocytosis inhibitor synaptotagmin XI (SYT11, Figure 33, Figure 33- figure supplement 1f). This pathway regulates the cell surface expression and signaling of RTKs and GPCRs[95,96], and we found that most of the enriched pathways associated with RTKs or GPCRs were enriched in pericytes (Pathway Analyzer).

Cell signaling (6v8). We identified thirty enriched pathways associated with RTK signaling in pericytes, including *signaling by NTRKs*, *signaling by VEGF*, and *signaling by PDGF* (FDR=0.011%, 0.365%, and 6.67%, respectively, Figure 33). Neurotropic tyrosine kinase receptors are activated by neurotrophins like NGF and NTF3—which were highly and differentially expressed by pericytes (Figure 33- figure supplement 1g). Pericytes also differentially express AXL receptor tyrosine kinase (AXL), caveolin-1 (CAV1), and integrin beta 3 (ITGB3), which define the enrichment of *signaling by VEGF* (Figure 33- figure supplement 1h). *Signaling by PDGF* was enriched in pericytes due to the differential expression of collagens COL4A1, COL4A2, COL4A4, COL5A3; platelet-derived growth factor alpha (PDGFA); and thrombospondin 4 (THBS4, Figure 33- figure supplement 1i).

The enrichment of thirteen Notch signaling pathways in pericytes indicate their importance to pericyte function. The broader *signaling by notch* pathway (FDR<0.001%) was enriched largely because of the differential expression of NOTCH receptors 1 and 3 (NOTCH1, NOTCH3), the downstream transcription factors HEYL and HEY2, and SMAD family member 3 (SMAD3, Figure 33- figure supplement 1s).

Cell-Cell Communication (6v8). Communication between pericytes and endothelial cells is critical to vessel development and function. We found pericytes enriched with the broad *cell-cell*

communication and cell junction pathways (FDR=0.10% and 0.06%, respectively, Figure 33).
These pathways involve tight junctions, adherins junctions, gap junctions, and signal regulatory
proteins, and pericytes differentially expressed a) cadherins 3 and 6 (CDH3 and CDH6), classical
calcium adhesion molecules; b) plectin (PLEC), which interacts and organizes connections to
intermediate filaments and microtubules and are essential components of desmosomes and
hemidesmosomes; and c) laminin subunit alpha-3 (LAMA3), which assembles into laminin
heterotrimers that are essential components of the vascular basement membrane. (Figure 33-
figure supplement 1j).

Extracellular Matrix Organization (6v8). Pericytes and fibroblasts have different functional
roles requiring distinct interactions with different ECM components. This was reflected in the
different ECM organization pathways we found in either cell type. For example, acting in
opposition, the *collagen formation* pathway was enriched in pericytes, whereas the *collagen
degradation* and *degradation of the extracellular matrix* pathways were enriched in fibroblasts.
Enrichment of the *Collagen formation* pathway in pericytes (FDR=4.28%) resulted from their
differential expression of COL4A1, COL4A2, COL4A3, COL4A4, COL4A5, COL5A3,
COL17A1, COL23A1, COL25A1, and COL27A1 (Figure 33, Figure 33- figure supplement 1k).
As introduced in the discussion of the collagen family above, the perivascular cells expressed the
highest levels of type IV collagens (basement membrane collagens, Figure 15—figure
supplement 2a-c), consistent with their spatial location in the tissues—embedded within the
vascular basement membranes—and role of maintaining these basement membranes. In contrast,
fibroblasts exist in the extracellular matrix, which they are responsible for forming and shaping.
We found fibroblasts differentially expressed collagens COL1A1, COL3A1, COL6A6,
COL8A1, COL8A2, COL14A1, and COL16A1, contributing to the *degradation of the
extracellular matrix* pathway's enrichment (FDR=0.705%, Figure 33, Figure 33- figure
supplement 1l). Additional differentially expressed genes included matrix metalloproteinases
MMP2, MMP3, MMP10, MMP12, MMP13, and MMP16; TIMP metallopeptidase inhibitor 1
(TIMP1); calpains 5 and 6 (CAPN 5 and CAPN6); and the bone remodeling protein cathepsin
K—which cleaves collagen I fibers (Figure 33- figure supplement 1l). We also found enrichment
of *ECM proteoglycans* in fibroblasts (FDR=4.68%), as they differentially expressed versican
(VCAN), asporin (ASPN), lumican (LUM), heparan sulfate proteoglycan 2 (HSPG2), decorin
(DCN), and biglycan (BGN) proteoglycans (Figure 33, Figure 33- figure supplement 1m). Other
genes in this pathway highly expressed by fibroblasts were tenascins N and C (TNN, TNC),
TGFB3, and laminin alpha 1 (LAMA1, Figure 33- figure supplement 1m).

Elastic Fibers (6v8). Elastic fibers are essential components of connective tissue and contribute
to tissue elasticity. We discovered pathways related to elastic fibers were among the most
significantly enriched pathways in fibroblasts, i.e., *molecules associated with elastic fibers*
(FDR=0.29%) and *elastic fiber formation* (FDR=0.90%). Leading to these pathways' enrichment
were the fibroblasts' robust expression of elastin (ELN), elastic microfibril interfacer 1
(EMILIN1), fibrillins 1 and 2 (FBN1 and FBN2), bone morphogenic protein 7 (BMP7),
fibronectin (FN1), and lysyl oxidase 1 (LOXL1, Figure 33, Figure 33- figure supplement 1n).

Metabolism (6v8). Pericytes and fibroblasts were enriched with different metabolic pathways.
There were fifteen pathways enriched in pericytes and thirteen in fibroblasts (Pathway analyzer).

One of the most significant pathways enriched in pericytes was *Integration of energy metabolism* (FDR=0.016%), stemming from the pericytes' differential expression of adrenoreceptor alpha 2A and 2C (ADRA2A and ADRA2C), thrombospondin receptor (CD36), adenylate cyclases 6 and 9 (ADCY6 and ADCY9), cysteinyl leukotriene receptor 2 (CYSLTR2), and the voltage-dependent calcium channel CACNB2 (Figure 33, Figure 33- figure supplement 1o). In contrast, fibroblasts were enriched in *fatty acid metabolism* (FDR=0.889%), owing to their differential expression of prostaglandin synthases PTGES, PTGS2, PTGIS, and PTGDS; the C2 domain-containing phospholipase PLA2G4A; and their unique expression of hydroxyacid oxidase 2 (HAO2, Figure 33, Figure 33—figure supplement 1p).

White adipocyte differentiation (6v8). We found forty-five of the eighty-four genes composing the *transcriptional regulation of white adipocyte differentiation* pathway were elevated in pericytes, leading to its enrichment (FDR=0.007% Figure 33). Genes differentially expressed by pericytes included NR2F2, which regulates apolipoprotein A-I transcription; CD36 (fatty acid translocase), which regulates the transport and uptake of long-chain fatty acids, and fatty acid binding protein-4 (FABP4), which mediates intracellular fatty acid transport (Figure 33—figure supplement 1q). One of the most notable genes in this pathway is phosphoenolpyruvate carboxykinase 1 (PCK1), which catalyzes the production of phosphoenol pyruvate from oxaloacetate. The activity of PCK1, which is regulated at the transcriptional level, commits mitochondrial pyruvate to gluconeogenesis or glyceroneogenesis[97]. This step is irreversible under physiological conditions and is the rate-limiting step controlling gluconeogenesis flux. We found PCK1 expression in breast cells was limited to the pericytes and adipocytes (Figure 33—figure supplement 1r), indicating, at the very least, that pericytes have an increased demand for cytoplasmic pyruvate.

Summary

Resolving the types and properties of cells in a tissue is vital to understanding the underlying symphony of interactions that make life possible. It is essential to overcome diseases such as cancer. Here, and in the adjoining article, we explored the fabric of the breast through a cytometric and molecular dissection of the tissue, which has provided us with insights into its inner workings. From the outset, we had questions regarding the number and types of cells composing the breast, their spatial location, their functions, and how they contributed to the tissue as a whole.

To address these questions, we revisited breast tissues to explore their architecture and cellular composition through an exhaustive cytometric dissection of the tissue, which led to our development of a FACS procedure that reconciled every cell in the tissue (Figure 1). We isolated twelve cell populations from four lineages and developed techniques to culture nearly every one (Figure 2)[8]. Our system of defining breast cell types provides a comprehensive resource that can be used to explore interactions in more complex and physiologically relevant culture systems aimed at mimicking the tissue microenvironment.

Here, we explored the nature of each breast cell type by dissecting their transcriptomes. These were acquired by scaling up our FACS procedure and painstakingly sorting each cell type to purity. RNA was then isolated and sequenced, leading to the acquisition of transcriptomes that were of high depth and quality. Although the rare cell types yielded fewer cells and transcripts, concordance of transcriptomic profiles between all cell replicates illustrated the reproducibility of our methods (Figure 4). Analysis of transcripts identified uniquely expressed genes that have already helped us decipher spatially resolved transcripts from breast tumors. Bioinformatic analyses of transcriptional profiles further helped validate the identity of several cell types by providing information on expressed markers and revealing their functional commonalities to related cell types (Figure 6).

Over 20,000 genes were measured in each cell type, providing a wealth of information. As we explored the transcriptomes from every major cell type in the breast, there were so many exciting features that we had to come to terms with the fact that we couldn't describe all that we encountered. Our solution was to build resource-sharing tools that aid exploration of our data[9,10]. To demonstrate the utility of these resources, we addressed three pressing questions: a) What features distinguish each cell lineage? b) What gene types confer cell-specific functions? and c) What biological processes define each cell type?

To identify the cell lineages' distinguishing features, we investigated the cellular pathways impacted by lineage-specific genes, which we defined through principal component analysis. The pathway enrichment analyses affirmed established attributes of each cell lineage while identifying others we had not previously associated with these lineages (Figures 10-14).

Focusing on the genes families that varied most among the different cell types provided us with an unconventional look at the types of genes contributing to the essential qualities of the breast cell types. We examined the expression of each gene, from each gene family, within every breast cell type, revealing the gene types that vary across cell types. We provide a full accounting of

each gene family in a supplemental table and have meticulously described nine gene families
whose functions extend from the nucleus (e.g., hormone receptors) to the exterior cell
microenvironment (e.g., laminins, Figures 15-19).

To address our last question and develop a broad perspective of each breast cell type's
fundamental properties and individual functions, we performed a comprehensive assessment of
the biological pathways and processes that are differentially enriched in each cell population. We
performed fifty-five independent analyses identifying differentially expressed genes and
enriched pathways between every possible cell-type comparison. These results are available as a
downloadable interactive application (Pathway Analyzer[9]). We comprehensively analyzed six
comparisons we believed were of the highest interest, as they revealed features of every major
cell type and identified those distinguishing closely related cell types (Figures 21-33). As the
field moves forward with single-cell and spatial transcriptomic approaches to explore and define
the significance of cellular heterogeneity within tissues and tumors[98,99], we predict that the
abundant number of transcripts measured here, within the full spectrum of FACS-purified breast
cell types, will provide a strong foundation for identifying cell populations and interpreting
results.

A broad wealth of information touching on many scientific disciplines is presented herein, and
we recognize that developing deeper insights requires engagement and a focused exploration of
the associated datasets. We offer novel tools to aid this exploration[9,10] and look forward to
creating and sharing others. This body of work, presented here and in the adjoining article,
presents a systematic approach for identifying, isolating, and culturing nearly every major breast
cell type of epithelial, endothelial, and mesenchymal origin. We spent many years refining our
FACS strategy, used multiple rounds of sorting to ensure purity, exhaustively stained tissues and
primary cultures to validate and characterize cell types, and comprehensively and painstakingly
analyzed each transcriptome. This rich resource, which we submit here, has provided countless
insights and has already changed our perception of the cells composing this elaborate and
beautiful tissue.

Methods

Tissue acquisition and processing. Breast tissues from reduction mammoplasties were obtained
from the Cooperative Human Tissue Network (CHTN), a program funded by the National
Cancer Institute. All specimens were collected with patient consent and were reported negative
for proliferative breast disease by board-certified pathologists. The University of California at
Berkeley Institutional Review Board and University of New Mexico Human Research
Protections Office granted use of anonymous samples through exemption status, according to the
Code of Federal Regulations 45 CFR 46.101. Upon receipt, several fragments (roughly 2cm^2)
were embedded in optimal cutting temperature compound (OCT) in tissue cassettes, flash-frozen
in nitrogen, and archived at -80ºC for later cryosectioning and staining. The remaining tissue,
typically 20-100g, was thoroughly rinsed with phosphate-buffered saline and processed to
organoids as previously described (gentle agitation method)[100]. Briefly, this included manually
mincing and scoring tissues with a scalpel and incubating them overnight (12-18 hrs, 37°C) with
0.1% collagenase I (Gibco/Invitrogen) in Dulbecco's Modified Eagle Medium containing 100

U/ml penicillin, 100 mg/ml streptomycin, and 100 mg/ml Normocin™ (Invivogen, San Diego,
CA). The resulting divested tissue fragments (organoids) were collected by centrifugation (100g
x 2 min) and either archived in liquid nitrogen (90%FBS+10% DMSO) or immediately
processed to cell suspensions for FACS analysis. Tissues used for RNA-sequencing were
immediately processed upon receipt and kept at 4°C during processing and FACS sorting, except
for the overnight collagenase digestion at 37°C. RNA-sequenced cells were from fresh
specimens and were not previously archived in nitrogen.

Antibodies. A list of antibodies and reagents used in this study is provided in Figure 3—Table
Supplement 8. The table includes antibody clone designations, conjugations, isotype, supplier
product numbers, dilution factors used.

Cryosectioning and Immunostaining. Immunofluorescence was performed on 10-80 mm
cryosectioned tissue sections as described in the adjoining article[8]. Briefly, specimens were fixed
with 4% paraformaldehyde for 5 minutes at 23°C, which was followed by 4% formaldehyde /
0.1% saponin for 5 minutes at 23°C. Tissues were rinsed for 20 minutes in wash buffer (0.1%
saponin/10% goat serum in PBS) and then incubated with primary antibodies, typically overnight
at 4°C. Following the primary incubation period, samples were washed and incubated with
secondary antibodies conjugated with Alexafluor 488, 568, 594, or 647 (Thermo). After 1 hour
incubation at 23°C, samples were rinsed in PBS and their nuclei counterstained with 300nM
DAPI (4',6-Diamidino-2-Phenylindole, Dihydrochloride, Thermo). Coverslips were mounted
with Fluoromount G (Southern Biotech). Images were captured using a Zeiss LSM710 confocal
microscope, Zeiss Axioscope, or EVOS FL Auto imaging station. If needed, image contrast was
applied to the entire image using Photoshop and were annotated with the antibodies used to stain
the tissue.

Cell Preparation for Flow Cytometry and FACS analysis. To prepare cell suspensions for
FACS and flow cytometry analysis, we thoroughly rinsed the organoids in PBS, then pelleted
them via centrifugation at 100g x 2 min. After removing PBS by aspiration, the organoids were
suspended in 1 ml of 'Cell Dissociation Reagent' (Sigma# C5914; or Thermo# 13150016),
incubated for two minutes at room temperature, upon which 3ml trypsin (used for RNA-
sequenced samples) was added (0.25%, Thermo 25200072; Note that we have more recently
found TrypLE better preserves CD49f staining, although it dampens CD34 resolution). Samples
were incubated by hand/body temperature, thus permitting visual inspection and brief (1-3sec)
pulse vortexing every 30-60 seconds. After the mixture became cloudy from the dissociating
cells (about 8-10 min), the organoids/cells were gently and repeatedly pipetted through a 16-
gauge needle until clumps dissipated. Afterward, we filtered cell suspensions through a 100mm
cell strainer and added 3ml 0.1% w/v soybean trypsin inhibitor to stop the digest (Sigma#
T9128). The suspensions were then filtered through a 40mm cell strainer, rinsed in 10-20ml
PBS, and pelleted by centrifugation at high centrifugal force (400g x 5 min.—or longer, until all
cells had pelleted, which we confirmed by microscopic examination of the supernatant). The
cells were then rinsed in 10 ml Hanks balanced salt solution/1% BSA(w/v), counted, and again
pelleted by centrifugation. After centrifugation, nearly all the Hanks/BSA was aspirated, leaving
the cell pellet with roughly 60ml of Hanks/BSA. The pellet was resuspended in this small
volume, and then we added FACS antibodies to the samples and incubated the samples on ice,

covered, for 30 minutes (See Figure 3- table supplement 8 for antibodies). Following the
incubation, the cells were rinsed in Hanks/1% BSA, centrifuged (400g x 5 min), and resuspended
in Hanks/BSA with To-Pro-3 viability marker (diluted 1:4000, Thermo T3605), then filtered
again through a 40mm cell strainer cap (into the FACS tube). We did not use DNase enzyme or
hypotonic RBC lysis solutions on the cell preparations because we found they were unnecessary.
We also wanted to avoid any potential deleterious effect these treatments may have on the cells,
such as reducing cell viability or altering gene expression levels. We, again, also conscientiously
chilled samples during cell preparation and the sorting procedure for similar concerns.

Flow Cytometry. The FACS panel and gating strategy outlined herein was developed using a
BD FACS Vantage 8-channel cytometer (FACSDIVA software) and has since been adapted and
reproduced using a 14-channel SONY SY3200. PMT voltages were optimized on both
instruments by analyzing individually stained compensation beads (AbC™ Anti-Mouse Bead
Kit, Thermo # A10344) at increasing voltage intervals for each detector. Because the FACS
Vantage has only eight available channels, it was necessary to use two markers (podoplanin and
CD45) in a single channel (A488), which we experimentally validated. To-Pro-3 was used as the
viability marker, so the corresponding 405nm channel (455/50) could be used for BV421. This
channel was dedicated to CD49f to provide a maximum resolution of cell populations, which is
essential to this sorting strategy. The considerable compensation typically required between the
heavily overlapping dyes PE/Cy5 and To-Pro-3 was not needed, as the To-Pro-3 negative
(viable) cells were used for downstream analyses, allowing us to use the PE/Cy5 channel for
Thy1. Negative controls consisted of unlabeled beads and cells incubated with isotype control
antibodies conjugated to each fluorophore. Sort times typically ranged 14-18 hours (50-120
million cells stained— needed to yield enough cells for RNA-sequencing). Cells were chilled at
the sample intake and collection chambers, and all samples were chilled on ice during the entire
experiment. All media and PBS were filtered through a 0.2mm filters to prevent the FACS
machine from triggering events from crystals and debris. Staining profiles of cells collected at
the beginning and end of the sorts (sometimes separated by as many as 18 hours) were identical
and did not show any signal loss or appreciable decrease in viability. Because our machines
could sort cells into a maximum of 4 tubes, we first sorted cells into four groups, containing 2-4
populations each. These mixtures were: Tube1 (CD34Pos) and contained Pops 7,8,9; Tube 2
(Thy1[Pos]) contained Pops 3,6; Tube 3 (Luminal) contained Pops 1&2; and Tube 4 (Null gate)
contained Pops 4,5,10,11. During the first round of sorting, cells were typically sorted at
approximately 8,000 cell/sec. We followed this initial enrichment with purity sorts for each cell
type, using stringent sort masks, sorting each population into individual tubes at a rate of
approximately 2,000 cells/sec. FACS data were analyzed using Flowjo software (version v7.6.3
– v10.8), Tree Star Inc./ Becton, Dickinson & Company).

RNA Isolation and Sequencing. Following the overnight digestion of tissues with collagenase,
we centrifuged the samples at 100g x 2 min. This produced aqueous and lipid phases that
separated under centrifugal force. The top lipid phase was collected, visualized microscopically
for cellular content, and then processed immediately for RNA extraction using Qiagen RNA
Lipid Tissue Mini Kit (Qiagen 74804), according to the manufacturer's instructions. Total RNA
from the FACS-purified cells was isolated immediately after FACS sorting using Qiagen RNA

Mini (>100,000 cells, Qiagen 74104) or RNA Micro kits (<100,000 cells, Qiagen 74004)
according to the manufacturer's instructions. Total RNA was treated with DNAse I for 30
minutes at 37°C (Ambion DNA-*free* DNA Removal kit, Thermo AM1906*)*, then treated with
DNAse Inactivation beads. RNA was then concentrated with RNA Clean & Concentrator™-5
columns (Zymo R1015), quantified by Nanodrop and Agilent Bioanalyzer assays, and stored at -
80°C.

RNA Library construction and sequencing. mRNA was captured using poly-T coated dishes
(Clontech) and used for library construction using SMARTer technology (SMARTer Stranded
RNA-seq Kit, Takara Biotech) following the manufacturer's guidelines. For low abundance
samples, mRNA was captured on biotinylated LNA coated beads, magnetically captured and
rinsed 3x in binding and wash buffer (10 mM Tris-HCl (pH 7.5), 1 mM EDTA, 2 M NaCl).
mRNA was eluted from Dynabeads in 10ml volume of elution buffer, following by a 5 min.
incubation at 68°C. It was then used for first-strand cDNA synthesis (SMARTer Stranded RNA-
seq Kit). Sample clean-up and removal of adapter-dimers) was performed using Ampure XP
beads (Beckman Coulter), according to the manufacturer's protocol. Specific attention was given
to the samples to prevent under- or over-drying the beads, which we found to reduce PCR
efficiency. PCR amplification of the RNA-Seq library was performed using SeqAmp DNA
Polymerase (Takara Bio, cat# 638509). Each library was then purified prior to sequencing using
SPRI AMPure Beads (Beckman Coulter). Sequencing was performed using Illumina Trueseq v.2
technology, targeting approximately 30 million paired-end reads for each cell population
(Illumina HiSeq 2x100bp).

RNA-Seq pre-processing and quality control assessment. We processed raw count data from
aligned fragments (Figure 3—table supplement 2) using R programming language (RStudio) and
associated BioConductor packages. Quality control analyses (Cook's distance box plot,
covariance matrix, sample correlation matrix) were performed using the DESeq2 package[13]. For
data visualization (donut and heatmaps), the count matrix data was transformed by two methods
used in the DESeq2 package: (i) regularized log transformation (rlog), and (ii) variance-
stabilizing transformation (vst), both normalized to library size. Quality control assessment of
expressed transcripts across sorted cell samples was conducted, and one sample was removed
from statistical analyses as it was an extreme outlier (Epithelial Population 4 Sample N239). This
was determined by: (i) distance matrix, (ii) Principal Component Analysis, and (iii)
Kolmogorov-Smirnov test that compares each sample to the reference probability distribution of
the pooled data. We included this sample in downstream visualizations for reference purposes,
for example, the bar and donut graphs used throughout the manuscript, which display rlog
transformed values on both log2 (bar graphs) and linear (donut graph) scales.

Differential Gene Expression Analysis. All differential expression analyses were performed
using DESeq2[13]. The DESEQ2 likelihood ratio test (LRT) was used to identify genes
differentially expressed genes across all cell types (Figure 3—table supplement 5); These genes
were then used to calculate the proportion of differentially expressed genes within each HGNC
families/group in our expression dataset (https://www.genenames.org/data/genegroup).
Differential expression p-values were corrected for multiple hypothesis testing using Benjamini-
Hochberg method across all genes and all 55 possible pairwise contrasts. Volcano plots of the

DESeq2 contrasted cell types were generated using the EnhancedVolcano R/Bioconductor
package[101]

Principal component analysis. Principal component analysis was performed on VST
transformed data (subsetting without tissue and bulk cell controls) using the pcaExplorer
R/Bioconductor package[102]. Loading factors from the ends of principal components 1, 2, 3, and 4
were further analyzed using the ClueGo[21] Cytoscape package, referencing the Gene Ontology
biological processes (GOBP) and Reactome pathway databases[18,20]. Subsequent enriched
pathways were grouped automatically by likeness to obtain a condensed list of defining features
for each cell class.

Gene Set Enrichment and Pathway Analysis. Pathway enrichment analysis was performed
using Broad Institute's GSEA software (version 4.3.2)[52]. We pre-ranked our individual gene lists
using the DESeq2 calculated Fold changes and Benjamini-Hochberg adjusted p-values from all
55 possible pairwise contrasts (Figure 3—table supplement 7). The rank metric was calculated
by: (sign[Fold Change]$*$ -$\log_{10}\{p$-value$\}$). GSEA analysis was performed using a classic,
unweighted enrichment statistic (p=0)[52] and was used to evaluate the enrichment of Reactome
pathways[18] Pathways were obtained via the Molecular Signatures Database[54], and we included
pathways with sizes between 15-200 genes. The resulting analysis produced a collection of
enriched pathways and calculated statistics (1000 permutations to correct for multiple hypotheses
testing) for each of the 55 pairwise comparisons. These data became the foundation for the
Excel-based Pathway Analyzer tool.

Enrichment Map/Cytoscape. Visualization of enriched gene sets was performed using the
Enrichment Map Cytoscape plugin[52,58], with applied edge cut-offs of 0.5 and FDR cutoffs of 0.1
(10%) for most comparisons, with the exception of the 8v12 comparison between fibroblasts and
adipocytes, which had a FDR of 0.2 (20%). The produced nodes (biological pathways) were
manually adjusted/grouped into themes for presentation purposes using the Reactome hierarchy
as a reference. Annotations of select biological themes and notable genes were added manually.

Supplemental Resources

Pathway Analyzer: This supplemental tool integrates regularized log transformed transcript
levels (mRNA-sequencing values) and enriched biological pathways identified by Gene Set
Enrichment Analysis (GSEA). Data from every possible pairwise comparison of cell types is
embedded in the tool, which is supplied as a MS Excel file. The file is approximately 300Mb and
contains computationally intensive algorithms (e.g., we run the file on both Windows and MAC-
based computers (Surface Laptop Studio [Core i7/3.3GHz with 32Gb RAM], Surface Book 2
[Corei7/4.2GHz with 16Gb RAM], Dell Inspiron Desktop [Corei5/3.2GHz with 8Gb RAM],
MacBook Pro [Corei7/2.1GHz with 16Gb RAM]), and we recommend using it on computers
with ≥16Gb RAM).

Breast Cell Atlas: Transcript Finder: http://breastcancerlab.com/breast-atlas/ . This online tool
facilitates exploration of mRNAs expressed by the breast cells. When provided individual gene
names (official gene symbol), the tool will display the transcript levels measured in each cell
type. Regularized log transformed values are provided on a log2 and linear scale (bar and donut

graph, respectively), similar to those presented in the many figure elements contained herein. This application was coded using the R/Shiny package[103].

Author Contributions

Conceptualization, WCH; FACS & Methodology, WCH; RNA isolation, WCH; mRNA Library production, ZW; RNA-seq alignment, quality control, and count matrix formation, AC; Bioinformatics, KDT, RS, AC, and WCH; Pathway Analyzer: KDT and WCH; Breast Cell Atlas: Transcript Finder: KDT, WCH; Immunostaining, WCH, KT, KDT, YLM; Writing – Original draft, WCH and KDT; Writing – Review & Editing, WCH, KDT, RS; Funding, WCH.

Acknowledgments

We thank Mina Bissell, Irene Kuhn, Alex Bazarov, Ritu Mukhopadhyay (Lawrence Berkeley Laboratory), Sandy Borowsky (U.C. Davis), Kornelia Polyak (Dana Farber Cancer Institute), Eric Prossnitz (University of New Mexico Health Sciences Center), Curtis Hines (Sandia National Laboratory, retired) for thoughtful scientific discussions and/or for critical review of the manuscript. We thank Tristan Oper, Alice Rindestig, Kremena Karagyozova, Alyssa Cozzo, Gaelen Stanford-Moore, Maria Rojec, Anya Afasisheva, and Sun-Young Lee for their technical assistance. We also thank Dana Pe'er, Ambrose Carr, Zhenmao Wan, and the staff at Columbia University's Genomic Sequencing Core for Illumina RNA-sequencing. Much of the analyses presented herein would not have been possible without Reactome's curation of genes, pathways, and publicly available resources. We thank Reactome's Principal Investigators and the team of dedicated scientists that have contributed to the Reactome Project. We also thank Kerry Wiles and Erik Brooks of the Cooperative Human Tissue Network, and the anonymous women who graciously donated their remnant surgical tissue to science. This project depended on flow cytometry instruments provided at the University of New Mexico Flow Cytometry and High Throughput Screening Resource and we thank Cecelia Rietz and Wade Johnson for their assistance. Finally, a very special thank you to Michelle Scott, Lawrence Berkeley National Laboratory Advanced Microscopy and Flow Cytometry facility, for her expert technical guidance and support.

Funding

Research reported in this publication was supported by the University of New Mexico Cancer Center Support Grant (P30CA118100), an Institutional Development Award (IDeA) from the National Institute of General Medical Sciences of the National Institutes of Health under grant number P20GM103451, the U.S. Department of Defense (W81XWH12M9532), and the Breast Cancer Research Foundation.

References

1. Cox, T.R. The matrix in cancer. *Nature reviews. Cancer* **21**, 217-238 (2021).
2. Maman, S. & Witz, I.P. A history of exploring cancer in context. *Nature Reviews Cancer* **18**, 359-376 (2018).
3. Quail, D.F. & Joyce, J.A. Microenvironmental regulation of tumor progression and metastasis. *Nature medicine* **19**, 1423-1437 (2013).
4. Bissell, M.J. & Hines, W.C. Why don't we get more cancer? A proposed role of the microenvironment in restraining cancer progression. *Nature medicine* **17**, 320-329 (2011).
5. Pierce, G.B. Neoplasms, differentiations and mutations. *The American journal of pathology* **77**, 103-118 (1974).
6. Zampieri, F., ElMaghawry, M., Zanatta, A. & Thiene, G. Andreas Vesalius: celebrating 500 years of dissecting nature. *Global cardiology science & practice* **2015**(2015).
7. Androutsos, G., Karamanou, M. & Stefanadis, C. William Harvey (1578-1657): discoverer of blood circulation. *Hellenic J Cardiol* **53**, 6-9 (2012).
8. Thi, K., Del Toro, K., Licon-Munoz, Y., Sayaman, R.W. & Hines, W.C. Comprehensive identification, isolation, and culture of human breast cell types. *bioRxiv* (2022).
9. Hines, W.C. & Del Toro, K. Pathway Analyzer. Supplementary Software for: A cellular and transcriptomic dissection of the human breast for studying mechanisms of cell and tissue function. (2022).
10. Del Toro, K. & Hines, W.C. Breast Cell Atlas: Transcript Finder (R/Shiny App). Supplementary Web Application for: A cellular and transcriptomic dissection of the human breast for studying mechanisms of cell and tissue function. (2022).
11. Hines, W.C., Thi, K., Rojec, M., Stanford-Moore, G. & Bissell, M.J. Abstract B86: A cytometric atlas of the human breast: Comprehensive characterization reveals 12 distinct cell populations. *Cancer research* **75**, B86-B86 (2015).
12. Ewing, B., Hillier, L., Wendl, M.C. & Green, P. Base-calling of automated sequencer traces usingPhred. I. Accuracy assessment. *Genome research* **8**, 175-185 (1998).
13. Love, M.I., Huber, W. & Anders, S. Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. *Genome Biol* **15**, 550 (2014).
14. Suzuki, R. & Shimodaira, H. Pvclust: an R package for assessing the uncertainty in hierarchical clustering. *Bioinformatics* **22**, 1540-1542 (2006).
15. Dyer, L.A. & Patterson, C. Development of the endothelium: an emphasis on heterogeneity. in *Seminars in thrombosis and hemostasis*, Vol. 36 227-235 (© Thieme Medical Publishers, 2010).
16. Mescher, A.L. Author. in *Junqueira's Basic Histology: Text and Atlas, 15e* (McGraw-Hill Education, New York, NY, 2018).
17. Gene Ontology, C. The Gene Ontology resource: enriching a GOld mine. *Nucleic acids research* **49**, D325-D334 (2021).
18. Jassal, B.*, et al.* The reactome pathway knowledgebase. *Nucleic acids research* **48**, D498-D503 (2020).
19. Bindea, G.*, et al.* ClueGO: a Cytoscape plug-in to decipher functionally grouped gene ontology and pathway annotation networks. *Bioinformatics* **25**, 1091-1093 (2009).
20. Ashburner, M.*, et al.* Gene ontology: tool for the unification of biology. The Gene Ontology Consortium. *Nat Genet* **25**, 25-29 (2000).

21. Bindea, G.*, et al.* ClueGO: a Cytoscape plug-in to decipher functionally grouped gene ontology and pathway annotation networks. *Bioinformatics* **25**, 1091-1093 (2009).

22. Kruger, R.P., Aurandt, J. & Guan, K.L. Semaphorins command cells to move. *Nat Rev Mol Cell Biol* **6**, 789-800 (2005).

23. Yuan, L.*, et al.* Abnormal lymphatic vessel development in neuropilin 2 mutant mice. *Development* **129**, 4797-4806 (2002).

24. Wang, J.*, et al.* NRP-2 in tumor lymphangiogenesis and lymphatic metastasis. *Cancer letters* **418**, 176-184 (2018).

25. Neil, S. & Bieniasz, P. Human immunodeficiency virus, restriction factors, and interferon. *J Interferon Cytokine Res* **29**, 569-580 (2009).

26. Wang, S.*, et al.* AXL is a candidate receptor for SARS-CoV-2 that promotes infection of pulmonary and bronchial epithelial cells. *Cell research* **31**, 126-140 (2021).

27. Liu, S., DeLalio, L.J., Isakson, B.E. & Wang, T.T. AXL-mediated productive infection of human endothelial cells by Zika virus. *Circulation research* **119**, 1183-1189 (2016).

28. Purkis, P.E.*, et al.* Antibody markers of basal cells in complex epithelia. *Journal of cell science* **97 (Pt 1)**, 39-50 (1990).

29. Dairkee, S. & Heid, H.W. Cytokeratin profile of immunomagnetically separated epithelial subsets of the human mammary gland. *In Vitro Cell Dev Biol Anim* **29A**, 427-432 (1993).

30. Taylor-Papadimitriou, J.*, et al.* Keratin expression in human mammary epithelial cells cultured from normal and malignant tissue: relation to in vivo phenotypes and influence of medium. *Journal of cell science* **94 (Pt 3)**, 403-413 (1989).

31. Perou, C.M.*, et al.* Molecular portraits of human breast tumours. *Nature* **406**, 747-752 (2000).

32. Miner, J.H. & Yurchenco, P.D. Laminin functions in tissue morphogenesis. *Annual review of cell and developmental biology* **20**, 255-284 (2004).

33. Yates, B.*, et al.* Genenames.org: the HGNC and VGNC resources in 2017. *Nucleic acids research* **45**, D619-D625 (2017).

34. Gray, K.A., Seal, R.L., Tweedie, S., Wright, M.W. & Bruford, E.A. A review of the new HGNC gene family resource. *Hum Genomics* **10**, 6 (2016).

35. Aumailley, M. The laminin family. *Cell adhesion & migration* **7**, 48-55 (2013).

36. Timpl, R., Sasaki, T., Kostka, G. & Chu, M.L. Fibulins: a versatile family of extracellular matrix proteins. *Nat Rev Mol Cell Biol* **4**, 479-489 (2003).

37. Khoshnoodi, J., Pedchenko, V. & Hudson, B.G. Mammalian collagen IV. *Microscopy research and technique* **71**, 357-370 (2008).

38. Borza, D.-B.*, et al.* The NC1 domain of collagen IV encodes a novel network composed of the α1, α2, α5, and α6 chains in smooth muscle basement membranes. *Journal of Biological Chemistry* **276**, 28532-28540 (2001).

39. Yao, C.C., Breuss, J., Pytela, R. & Kramer, R.H. Functional expression of the alpha 7 integrin receptor in differentiated smooth muscle cells. *Journal of cell science* **110 (Pt 13)**, 1477-1487 (1997).

40. Hyun, Y.-M., Lefort, C.T. & Kim, M. Leukocyte integrins and their ligand interactions. *Immunologic research* **45**, 195-208 (2009).

41. Visse, R. & Nagase, H. Matrix metalloproteinases and tissue inhibitors of metalloproteinases: structure, function, and biochemistry. *Circulation research* **92**, 827-839 (2003).

42. Welgus, H.G.*, et al.* Neutral metalloproteinases produced by human mononuclear phagocytes. Enzyme profile, regulation, and expression during cellular development. *The Journal of clinical investigation* **86**, 1496-1502 (1990).

43. Lepidi, S.*, et al.* MMP9 production by human monocyte-derived macrophages is decreased on polymerized type I collagen. *Journal of vascular surgery* **34**, 1111-1118 (2001).

44. Zihni, C., Mills, C., Matter, K. & Balda, M.S. Tight junctions: from simple barriers to
multifunctional molecular gates. *Nat Rev Mol Cell Biol* **17**, 564-580 (2016).
45. Nielsen, M.S*., et al.* Gap junctions. *Comprehensive Physiology* **2**(2012).
46. Heldin, C.H., Lu, B., Evans, R. & Gutkind, J.S. Signals and Receptors. *Cold Spring Harb Perspect*
Biol **8**, a005900 (2016).
47. Westbury, C.B*., et al.* Genome-wide transcriptomic profiling of microdissected human breast
tissue reveals differential expression of KIT (c-Kit, CD117) and oestrogen receptor-alpha
(ERalpha) in response to therapeutic radiation. *The Journal of pathology* **219**, 131-140 (2009).
48. Moreira, J.M*., et al.* Tissue proteomics of the human mammary gland: towards an abridged
definition of the molecular phenotypes underlying epithelial normalcy. *Mol Oncol* **4**, 539-561
(2010).
49. Lim, E*., et al.* Aberrant luminal progenitors as the candidate target population for basal tumor
development in BRCA1 mutation carriers. *Nature medicine* **15**, 907-913 (2009).
50. Cato, A. & Shatkina, L. Nuclear Hormone Receptors: Regulators of Gene Transcription and
Cellular Signaling. *Encyclopedic Reference of Genomics and Proteomics in Molecular Medicine*,
1314-1320 (2006).
51. Frigo, D.E., Bondesson, M. & Williams, C. Nuclear receptors: From molecular mechanisms to
therapeutics. Vol. 65 847-856 (Portland Press Ltd., 2021).
52. Subramanian, A*., et al.* Gene set enrichment analysis: a knowledge-based approach for
interpreting genome-wide expression profiles. *Proc Natl Acad Sci U S A* **102**, 15545-15550
(2005).
53. Mootha, V.K*., et al.* PGC-1alpha-responsive genes involved in oxidative phosphorylation are
coordinately downregulated in human diabetes. *Nat Genet* **34**, 267-273 (2003).
54. Liberzon, A*., et al.* Molecular signatures database (MSigDB) 3.0. *Bioinformatics* **27**, 1739-1740
(2011).
55. Coordinators, N.R. Database resources of the National Center for Biotechnology Information.
Nucleic acids research **44**, D7-19 (2016).
56. Cunningham, F*., et al.* Ensembl 2022. *Nucleic acids research* **50**, D988-D995 (2022).
57. Kanehisa, M. & Goto, S. KEGG: kyoto encyclopedia of genes and genomes. *Nucleic acids research*
28, 27-30 (2000).
58. Merico, D., Isserlin, R., Stueker, O., Emili, A. & Bader, G.D. Enrichment map: a network-based
method for gene-set enrichment visualization and interpretation. *PLoS One* **5**, e13984 (2010).
59. Giamas, G*., et al.* Kinome screening for regulators of the estrogen receptor identifies LMTK3 as a
new therapeutic target in breast cancer. *Nature medicine* **17**, 715-719 (2011).
60. Guo, S. & Sonenshein, G.E. Forkhead box transcription factor FOXO3a regulates estrogen
receptor alpha expression and is repressed by the Her-2/neu/phosphatidylinositol 3-kinase/Akt
signaling pathway. *Molecular and cellular biology* **24**, 8681-8690 (2004).
61. Cancer Genome Atlas, N. Comprehensive molecular portraits of human breast tumours. *Nature*
490, 61-70 (2012).
62. Zhu, Y*., et al.* Coregulation of estrogen receptor by ERBB4/HER4 establishes a growth-promoting
autocrine signal in breast tumor cells. *Cancer research* **66**, 7991-7998 (2006).
63. Sardi, S.P., Murtie, J., Koirala, S., Patten, B.A. & Corfas, G. Presenilin-dependent ErbB4 nuclear
signaling regulates the timing of astrogenesis in the developing brain. *Cell* **127**, 185-197 (2006).
64. Stoker, M., Gherardi, E., Perryman, M. & Gray, J. Scatter factor is a fibroblast-derived modulator
of epithelial cell mobility. *Nature* **327**, 239-242 (1987).
65. Weidner, K.M*., et al.* Evidence for the identity of human scatter factor and human hepatocyte
growth factor. *Proc Natl Acad Sci U S A* **88**, 7001-7005 (1991).

66.   Guimaraes, J.C. & Zavolan, M. Patterns of ribosomal protein expression specify normal and
malignant human cells. *Genome Biol* **17**, 236 (2016).
67.   Lin, E.Y., Gouon-Evans, V., Nguyen, A.V. & Pollard, J.W. The macrophage growth factor CSF-1 in
mammary gland development and tumor progression. *Journal of mammary gland biology and*
neoplasia **7**, 147-162 (2002).
68.   Andreas, N.J., Kampmann, B. & Mehring Le-Doare, K. Human breast milk: A review on its
composition and bioactivity. *Early Hum Dev* **91**, 629-635 (2015).
69.   Lonnerdal, B. Nutritional and physiologic significance of human milk proteins. *Am J Clin Nutr* **77**,
1537S-1543S (2003).
70.   Bawden, W.S., Passey, R.J. & Mackinlay, A.G. The genes encoding the major milk-specific
proteins and their use in transgenic studies and protein engineering. *Biotechnology and Genetic*
Engineering Reviews **12**, 89-138 (1994).
71.   Visvader, J.E. Cells of origin in cancer. *Nature* **469**, 314-322 (2011).
72.   Frantz, C., Stewart, K.M. & Weaver, V.M. The extracellular matrix at a glance. *Journal of cell*
science **123**, 4195-4200 (2010).
73.   Skalli, O.*, et al.* A monoclonal antibody against alpha-smooth muscle actin: a new probe for
smooth muscle differentiation. *The Journal of cell biology* **103**, 2787-2796 (1986).
74.   Venkatachalam, K. & Montell, C. TRP channels. *Annu Rev Biochem* **76**, 387-417 (2007).
75.   Kufe, D.W. Mucins in cancer: function, prognosis and therapy. *Nature reviews. Cancer* **9**, 874-
885 (2009).
76.   Schroten, H.*, et al.* Inhibition of adhesion of S-fimbriated Escherichia coli to buccal epithelial
cells by human milk fat globule membrane components: a novel aspect of the protective
function of mucins in the nonimmunoglobulin fraction. *Infect Immun* **60**, 2893-2899 (1992).
77.   Labrie, F.*, et al.* The key role of 17β-hydroxysteroid dehydrogenases in sex steroid biology.
Steroids **62**, 148-158 (1997).
78.   Fuchs, E. & Weber, K. Intermediate filaments: structure, dynamics, function, and disease. *Annu*
Rev Biochem **63**, 345-382 (1994).
79.   Simionescu, M., Simionescu, N., Silbert, J.E. & Palade, G.E. Differentiated microdomains on the
luminal surface of the capillary endothelium. II. Partial characterization of their anionic sites. *The*
Journal of cell biology **90**, 614-621 (1981).
80.   Rosenberg, R.D., Shworak, N.W., Liu, J., Schwartz, J.J. & Zhang, L. Heparan sulfate proteoglycans
of the cardiovascular system. Specific structures emerge but how is synthesis regulated? *The*
Journal of clinical investigation **99**, 2062-2070 (1997).
81.   van Buul, J.D.*, et al.* RhoG regulates endothelial apical cup assembly downstream from ICAM1
engagement and is involved in leukocyte trans-endothelial migration. *The Journal of cell biology*
178, 1279-1293 (2007).
82.   Arnal, J.-F.*, et al.* Estrogen receptors and endothelium. *Arteriosclerosis, Thrombosis, and*
Vascular Biology **30**, 1506-1512 (2010).
83.   Tatematsu, M.*, et al.* Raftlin controls lipopolysaccharide-induced TLR4 internalization and
TICAM-1 signaling in a cell type–specific manner. *The Journal of Immunology* **196**, 3865-3876
(2016).
84.   Hauser, M.A. & Legler, D.F. Common and biased signaling pathways of the chemokine receptor
CCR7 elicited by its ligands CCL19 and CCL21 in leukocytes. *Journal of leukocyte biology* **99**, 869-
882 (2016).
85.   Moreira, L.O. & Zamboni, D.S. NOD1 and NOD2 signaling in infection and inflammation. *Frontiers*
in immunology **3**, 328 (2012).
86.   Jackson, W.F. Potassium channels in the peripheral microcirculation. *Microcirculation* **12**, 113-
127 (2005).

87. Gupta, R.K. Adipocytes. *Curr Biol* **24**, R988-993 (2014).

88. Storch, J. & Thumser, A.E. Tissue-specific functions in the fatty acid-binding protein family. *The*
Journal of biological chemistry **285**, 32679-32683 (2010).

89. Li, Y.*, et al.* Cidec differentially regulates lipid deposition and secretion through two tissue-
specific isoforms. *Gene* **641**, 265-271 (2018).

90. Chen, L.*, et al.* AQP7 mediates post-menopausal lipogenesis in adipocytes through FSH-induced
transcriptional crosstalk with AP-1 sites. *Reproductive BioMedicine Online* **41**, 1122-1132 (2020).

91. Blüher, M. Importance of estrogen receptors in adipose tissue function. *Molecular metabolism*
2, 130 (2013).

92. Winkler, E.A., Sagare, A.P. & Zlokovic, B.V. The Pericyte: A Forgotten Cell Type with Important
Implications for A lzheimer's Disease? *Brain pathology* **24**, 371-386 (2014).

93. Armulik, A., Genové, G. & Betsholtz, C. Pericytes: developmental, physiological, and pathological
perspectives, problems, and promises. *Developmental cell* **21**, 193-215 (2011).

94. Rapport, M.M. Serum vasoconstrictor (serotonin) the presence of creatinine in the complex; a
proposed structure of the vasoconstrictor principle. *The Journal of biological chemistry* **180**, 961-
969 (1949).

95. Goh, L.K. & Sorkin, A. Endocytosis of receptor tyrosine kinases. *Cold Spring Harb Perspect Biol* **5**,
a017459 (2013).

96. Wolfe, B.L. & Trejo, J. Clathrin-dependent mechanisms of G protein-coupled receptor
endocytosis. *Traffic* **8**, 462-470 (2007).

97. Beale, E.G., Harvey, B.J. & Forest, C. PCK1 and PCK2 as candidate diabetes and obesity genes.
Cell Biochem Biophys **48**, 89-95 (2007).

98. Nguyen, Q.H.*, et al.* Profiling human breast epithelial cells using single cell RNA sequencing
identifies cell diversity. *Nature Communications* **9**, 1-12 (2018).

99. Wu, S.Z.*, et al.* A single-cell and spatially resolved atlas of human breast cancers. *Nature genetics*
53, 1334-1347 (2021).

100. Hines, W.C., Su, Y., Kuhn, I., Polyak, K. & Bissell, M.J. Sorting Out the FACS: A Devil in the Details.
Cell reports **6**, 779-781 (2014).

101. Blighe, K., Rana, S. & Lewis, M. EnhancedVolcano: Publication-ready volcano plots with
enhanced colouring and labeling. *R package version* **1**(2019).

102. Marini, F. & Binder, H. pcaExplorer: an R/Bioconductor package for interacting with RNA-seq
principal components. *BMC Bioinformatics* **20**, 331 (2019).

103. Chang, W.*, et al.* shiny: Web Application Framework for R. R package version 1.6. 0. 2021. *URL*
R-project. org/package= shiny (2021).

2310

2311

Figure 3—Table supplement 8. Antibodies

Tissue Staining (Immunofluorescence)				
Antibody specificity	**Clone**	**Conjugated**	**Source**	**Dilution**
Alpha-Smooth Muscle Actin	1A4	No	Mouse IgG2a, κ, Lab Vision/Neomarkers MS-113-P0ǁ	1:100
CD29	TS2/16	A488	Mouse IgG1, κ; Biolegend 303015	½ assay¶
CD49f	GoH3	No	Rat IgG2a, κ; Biolegend 313602	½ assay¶
E-Cadherin (CD324)	24E10	No	Rabbit mAb; Cell Signaling 3195S	1:100
E-Cadherin (CD324)	67A4	A488	Mouse IgG1, κ; BioLegend 324110	½ assay¶
erbB2/Her2 (CD340)	24D2	PE	Mouse IgG1, κ; Biolegend 324406	½ assay¶
Keratin 14	Polyclonal	No	Rabbit Polyclonal; Thermo/Lab Vision RB-9020-P1	1:400
Keratin 14	Polyclonal	No	Chicken Polyclonal; Biolegend 906001	1:400
Keratin 18	DC10	No	Mouse IgG1; Thermo/Lab Vision MS142P0	1:400
Keratin 19	A53-B/A2.26; same as Ks19.1	No	IgG2a/λ; Lab Vision/Neomarkers MS-198-PABX	1:400
Keratin 8		No	Mouse IgG2a κ	1:400
P63	4A4	None	Abcam ab32353	1:100
Pan-Laminin	Polyclonal	No	Rabbit Polyclonal; L9393	1:400
Pan-Laminin	Polyclonal	No	Rabbit Polyclonal; L9393	1:400
Podoplanin	NC-08	No	Rat IgG2a, λ	½ assay¶
Facs Panel Antibodies / Viability Dyes				
CD49f (alpha-6 integrin)	GoH3	BV421*	BioLegend 313624, 313612, 313607	2.5μl§
CD24	ML5	BV605*	BioLegend 311124, 311106, 311108	2.5μl§
CD227 (Muc1)	16A	PE	BioLegend 355604,	4μl§
Podoplanin	NC-08	A488†	BioLegend 337006, 337004	2.5μl§
CD45	HI30	A488†	BioLegend 304052, 304032	3μl§

CD90 (Thy1)	5E10	PE-Cy5	BioLegend 328112, 328134	3μl§
CD34	581	PE-Cy7	BioLegend 343516	1-2μl§
CD10	HI10a	APC-Cy7	BioLegend 312212, 312204	3μl§
TO-PRO™-3 Iodide	N/A	(642/661)	Thermo (T3605)	1:4000

Brilliant Violet; † Alexa Fluor; (Biolegend); § 1x volume per 6×10^6 cells stained in FACS experiments (up to 36×10^6, after which cells received 6x volumes; max stained cells $= 120 \times 10^6$); titration required and must be matched to cytometer. ¶ Percentage of manufacturer's suggested assay volume per 100ul volume (antibody is prediluted).

List of Table supplements and Data	
Figure 3—table supplement 1	Sample Treatment.xlsx
Figure 3—table supplement 2	CountMatrix_rLog_VST.xlsx
Figure 3—table supplement 3	PCA_Loading_Factors.xlsx
Figure 3—table supplement 4	GO from PCAs.xlsx
Figure 3—table supplement 5	DESeq2_LRT.xlsx
Figure 3—table supplement 6	Gene Families_DE Genes.xlsx
Figure 3—table supplement 7	DESeq2_files.xlsx
OSF; https://osf.io/gfhc3/	Cytoscape/Enrichment Map files (1v2, 2v3, 2v9, 6v8)
OSF; https://osf.io/hpb9t/	Pathway Analyzer[9]
http://breastcancerlab.com/breast-atlas	Breast Cell Atlas: Transcript Finder (R/Shiny App)[10]

Figure Legends

Figure 1. FACS gating strategy for comprehensive cell type isolation. (a) Prior to FACS isolation, breast tissues were enzymatically digested with collagenase. This was followed by low-speed centrifugation, which yielded an organoid pellet, supernatant, and an adipocyte-enriched lipid layer. RNA was isolated from the lipid layer (adipocytes, Pop12) and stored for later sequencing. The remaining cell types were FACS purified from the trypsin-dissociated organoid cell pellet, according to the outlined gating scheme. Each cell type was assigned a population number that corresponds to each population's isolation strategy (Pops 1-12). **(b)** Relative abundance of each cell type from 30 different individuals (proportion of total cell counts, as measured by FACS analysis of 44 independent FACS experiments). Outliers (*) are defined as points falling (1.5 x IQR) above/below the 75^{th} and 25^{th} quartile.

Figure 1—figure supplement 1. FACS strategy designed for an 8 channel FACS machine (BD FACS Vantage). (a) FACS gating scheme used to isolate the RNA-sequenced cell types (antibody panel designed for an 8 channel FACS machine- BD FACS Vantage). Representative FACS data are of cells derived from disease-free reduction mammoplasty tissue surgically resected from a 22-year-old female (sample #N239).

Figure 2. IHC staining of cell types in the breast. (a) Representative 4x scan of normal breast tissue (reduction mammoplasty tissue from 23-year-old female). The general location of the 12 cell types are indicated by the their FACS population identification numbers, Pops1-12. **(b)** Chart of the 12 breast cell types and their assigned colors (used throughout this article). Cell types are grouped by their common histological classification: epithelial, mesenchymal, and blood cell types. Epithelial types are further subdivided into glandular and endothelial types, whereas the mesenchymal perivascular types are indicated (pericytes and vascular smooth muscle cells). **(c-f)** spatial position of the cell types in normal tissue immunostained with: **(c)** keratin 18 (green; luminal epithelial cells) and pan-laminin (red; epithelial and vascular basement membrane, and adipocytes). Staining reveals a collecting duct within several breast lobules (center) composed of epithelial (Pops 1-4), fibroblasts (Pop8), and capillaries that consist of pericytes (Pop6) and vascular endothelial cells (Pop9). Stromal adipocytes are also present (Pop12, upper-right). Scale = 1000μm. **(d)** podoplanin (green) and pan-laminin (red) staining reveal podoplanin-staining lymphatic vessels (Pop7) and vascular blood vessels (Pop9). Laminin embedded pericytes (Pop6) are also discernible on the capillaries; Scale = 200μm; **(e)** CD34 (red) and pan-laminin (green) staining reveals a cross-section of three-and a-half breast lobules within a bed of CD34^{Pos} fibroblasts. Endothelial cells forming capillaries and larger blood vessels also stain for CD34, both inside and outside the lobular areas, e.g., larger vessels are visible at lower right, along with pan-laminin staining adipocytes (Pop12); Scale = 200μm **(f)** CD49f (green) and pan-laminin (red) reveals a large blood vessel (center) surrounded by vascular smooth muscle cells (Pop5). Endothelial cells (Pop9) and pericytes (Pop6) of smaller capillaries are present, as are lymphatic vessels (Pop7) and adipocytes (Pop12); Scale = 200μm.

Figure 2—figure supplement 1. (a) High-resolution image of the immunostained breast tissues provided in Fig. 2a. **(b)** magnified area indicated by white box in a (on left) containing several breast lobules (a.k.a, terminal ductal lobular units). **(c)** magnified area indicated by white box in

a (on right) containing the cross section of a large lactiferous duct. The duct is surrounded by
numerous blood vessels running parallel to the duct and is embedded in a collagenous-rich
matrix.

Figure 2- Figure supplement 2. (a,b) High-resolution (label-free) images of the immunostained
breast tissues provided in Figs 2c,d.

Figure 2- Figure supplement 3. (a,b) High-resolution (label-free) images of the immunostained
breast tissues provided in Figs 2e.f.

Figure 3. Distinguishing the traits of each breast cell type. Shown is a sketch of the RNA-
sequencing workflow used for analyzing the eleven sequenced cell types. This began with **(a)**
developing a FACS isolation method for separating each breast cell type (as detailed in Fig.1 and
adjoining article). This procedure was scaled up and cells were sorted and RNA-sequenced.
Analysis of the RNA-sequencing data involved: **(b)** validation and exploration of cellular
relationships by hierarchical clustering, **(c)** covariance matrix and quality control analyses,
identification of unique cell markers, and principal component analysis (PCA). PCA-derived
gene lists were used to perform gene ontology enrichment analyses that revealed the shared
characteristics of cell lineage. **(d)** Differential gene expression analysis was performed and used
to identify gene families that varied most among breast cell types; Pairwise comparisons
identified transcripts whose levels varied between sample pairs. Differentially expressed genes
between cell pairs were ranked and analyzed by **(e)** Gene Set Enrichment Analysis, leading to the
identification of key pathways and processes that differ between cell types. GSEA results were
further analyzed by **(f)** Constructing and analyzing enrichment maps that identified the central
biological themes and distinguishing features of each cell type. Analyzing the data required the
development of software code and packages that are available for download: Pathway Analyzer
(Open Science Framework, https://osf.io/), and online Transcript Finder
(http://breastcancerlab.com/breast-atlas).

Figure 3—figure supplement 1. Project workflow. Diagram of the sequential processes and
analyses presented in the adjoining and current articles. The adjoining article (in green) describes
the development and validation of a FACS isolation strategy for separating all twelve major
breast cell types, culminating with the development of primary cell models for nine major breast
cell types. The current article (in blue) explores the nature of cell type through bulk-mRNA-
sequencing and analysis of transcript levels measured in each FACS-purified population.
Analysis included principal component analysis, differential gene expression, and gene set
enrichment analysis. Red-outlined boxes indicate the major products from each publication.

Figure 3—table supplement 1. Sample information/Meta Data. Table of sample information,
containing age/sex/race information supplied with the acquired tissues (B=black; W=white),
Patient number, FACS gate, cell type, cell yields, RNA-isolation strategy/kit, elution volumes,
nanodrop concentrations, DNase treatment used, Bioanalyzer results (concentration and integrity
number), RNA-sequencing batch, and calculated reads in each sample.

Figure 3—table supplement 2. RNA-sequencing count matrix.

Figure 3—table supplement 3. PCA loading factors. Sheet 1, 'All Pops,' contains loading factors calculated from the analysis of all cell types (Pops 1-12). Sheet 2, 'Endothelial Removed,' contains loading factors calculated from the analysis of non-endothelial cell types (Pops 1-7,8,10-12).

Figure 3—table supplement 4. Tabulated Pathway Enrichment Results (ClueGo/Reactome/Gene Ontology Biological Processes) that used the 200 loading factors associated with the four cell lineages: a) Epithelial end of PC1, b) Leukocyte end of PC2, c) Mesenchymal end of $PC1^{No\ Endo}$, d) Endothelial end of PC3, and the perivascular and myoepithelial lineage subtypes e) Perivascular cell end of PC3, and f) myoepithelial end of PC4.

Figure 3—table supplement 5. RNA-Seq DESeq2 Likelihood Ratio Test (LRT) statistics, p-values, and Benjamini Hochberg adjusted *p*-values for differentially expressed genes with LRT adj.*p*-val<0.05. These results were used to calculate the proportion of differentially expressed genes in each HGNC gene family.

Figure 3—table supplement 6. Table of the 1,421 HGNC Gene families, number of genes in each family, and the calculated number of differentially expressed genes in each family (using the DESeq2 Benjamini Hochberg Adj. *p*-value threshold of ≤ 0.1).

Figure 3—table supplement 7. Excel workbook containing 55 DESeq results (on separate tabulated worksheets) from each possible pairwise contrast of the 11 seqenced breast cell types; e.g. Pop1v2 tab contains results from the contrast of Pop1 (ER^{Pos} luminal epithelial cells) and Pop2 (ER^{Neg} luminal epithelial cells). Tab 1v2 also contains the calculated rank metric used for GSEA analysis; i.e., *sign([Fold Change] * -log₁₀ {p-value}).*

Figure 3—table supplement 8. Antibodies. Antibody clones used for immunostaining and FACS.

Figure 4. RNA-seq quality assessment. (a) Sample metadata for the mRNA-sequenced cell populations. **(b)** Covariance matrix displaying the correlation of gene expression for all pairwise combinations of samples. Samples N228-Pop4, N225-Pop4, N228-Pop5, N228-Pop7 have been flagged for lower correlations between replicates (*). These cell populations are the least abundant types in the breast and are among those that produced the lowest FACS cell yields **(c)** distribution of regularized log transcript levels in each sample. **(d)** Sample correlation matrix demonstrates absence of sample/batch effects.

Figure 4—figure supplement 1. mRNA levels correspond to FACS staining. (a) Normalized mRNA values for genes encoding FACS markers used to purify cell types from tissues (and CD36, which was later used to purify cultured adipocytes). Transcript levels are provided on a) a log2 scale (bar graphs containing each biological replicate) and b) a linear scale (donut graphs of the replicate's median value), which are both color-coded by cell type. **(b)** Schematic indicating relative FACS staining levels of each marker in the twelve breast cell types. "Neg" indicates relative negative staining (i.e., compared to other populations, not absolute negativity), "-/+" indicates near background level staining, and the increasing number of "+" indicates greater staining intensity. [†] CD36 is a marker used to purify cultured adipocytes but was not used to

purify adipocytes prior to RNA-sequencing in this analysis. **(c)** Normalized transcript levels for the commonly used internal control gene GAPDH (rlog, DEseq2) are presented on a log2 scale (bar graph) and linear scale (donut graph of median value).

Figure 5. Uniquely expressed genes. A sliding threshold (fold-change) was used to identify the top 20 most uniquely expressed genes in each of the eleven sequenced cell types. Notable genes included: **a)** albumin (ALB) and connexin 30.3 (GJB4) in ER-positive luminal cells, **(b)** lactotransferrin (LTF) in ER-negative luminal cells, **(c)** alpha-1 chain of collagen XI (COL11A1) and notch ligand delta-like 3 (DLL3) in myoepithelial cells, **(d)** protocadherin 20 (PCDH20) in Pop4 epithelial cells, **(e)** lymphatic vessel endothelial hyaluronan receptor 1 (LYVE1) and prospero homeobox 1 (PROX1) in lymphatic endothelial cells, **(f)** endothelial selectin (SELE) and delta like canonical notch ligand 4 (DLL4) in vascular endothelial cells, **(g)** calcitonin related polypeptide alpha (CALCA) and 5-hydroxytryptamine receptor 1b (HTR1B) in vascular smooth muscle cells, **(h)** desmin (DES) and 5-hydroxytryptamine receptor 1f (HTR1F) in pericytes, **(i)** alpha-1 chain of collagen I (COL1A1) and wnt ligand family member 2 (WNT2) in fibroblasts, **(j)** the lipid-binding protein 'cell death inducing dffa-like effector c' (CIDEC) and glycerol-3-phosphate dehydrogenase 1 (GPD1) in adipocytes, and **(k)** CD7 and fas ligand (FASLG) in leukocytes.

Figure 5—figure supplement 1. CD antigen genes. Median mRNA expression levels of genes encoding CD antigens (VST, log2 scale)**.** The genes are arranged by the cell type that expressed them at their highest level, as indicated by the background shading specific to each cell type. Genes differentially expressed across cell populations are annotated with (*). CD antigen genes in the uncolored panel were not expressed by any breast cell type.

Figure 6. Transcript levels of top 2000 most differentially expressed genes. Heatmap with dendrogram of the top 2000 most differentially expressed genes across all FACS sorted populations, as well as unsorted cells and matched breast tissues (mixed cell controls). Clustering was performed using Euclidean distance measures and Ward clustering method on variance stabilized expression values (DESeq2; Variance Stabilized Transformation, VST). Adjusted p-values for these 2000 differentially expressed genes ranged between 4.9e-97 to 0.0017. All 2000 genes symbols can be viewed in Figure 3—table supplement 5 (when the data is ordered by adj. *p*-value).

Figure 7. Principal Component Analysis (a) projection of breast cell types on principal components 1 and 2 (PC1, PC2). PC1 separates the luminal ER^{Pos} and ER^{Neg}, Pop4 epithelial, and myoepithelial cell types (epithelial lineages) from the mesenchymal (fibroblasts, adipocytes, pericytes and vSMCs) and endothelial (lymphatic and vascular) lineages. PC2 separates the $CD45^{Pos}$ leukocytes (mesodermal lineage) from all other cell types. The cell types composing each of the epithelial, mesenchymal, endothelial, and mesodermal cell-lineages are indicated (dashed circles). **(b)** Median transcript levels of the 50 loading factors (genes) that contribute to the negative end of PC1(bottom end of the list, rlog/log2 scale). **(c)** Median mRNA expression of the 50 loading factors (genes) contributing to the negative end of PC2 (bottom end of the list) that separates the leukocyte samples (rlog, log2 scale). Stars (★) denote commonly used markers for luminal cells (blue), myoepithelial cells (red), and leukocytes (pink).

Figure 7—figure supplement 1. Variance captured by principal components. A screeplot shows the sample variances captured by each principal component. 71% of the total variance is captured by the first four components.

Figure 7—figure supplement 2. PCA with endothelial cells removed. (a) PCA projection of breast cell types, after removal of the lymphatic and vascular endothelial cells, resolves the mesenchymal cell types on PC1^{NoEndo}. **(b)** Median mRNA expression of the 50 loading factors (genes) contributing to the positive end of PC2 (top end of the list; rlog, log2 scale). Stars (★) denote common markers of pericytes (purple) and fibroblasts (yellow).

Figure 8. PCA plot of PC3 and PC4. (a) PCA projection of breast cell types on principal components 3 and 4 (PC3, PC4). PC3 separates the endothelial (most positive) and perivascular (most negative) cell types, whereas PC4 separates the myoepithelial and myoepithelial-related cell types (myoepithelial and Pop4 epithelial cells). Median mRNA expression of the 50 loading factors contributing to the **(b)** positive end of PC3 (top of the list; rlog, log2 scale), **(c)** negative end of PC3 (bottom of the list; rlog, log2 scale), and **(d)** the positive end of PC4 (top of the list; rlog, log2 scale). Stars (★) denote common markers of endothelial cells (orange), myoepithelial cell (red), fibroblasts (yellow), pericytes (purple), and luminal cells (blue).

Figure 9. IHC validation of PCA markers. Tissue immunostaining of normal human breast tissue with markers contained in the principal components. **(a)** keratin 19 (green) and pan-laminin (red); **(b)** keratin 8 (red); **(c)** e-cadherin (green) and alpha smooth muscle actin (red); **(d)** Her2 (green) and keratin 14 red); **(e)** CD49f (green) and vimentin (red); **(f)** alpha smooth muscle actin (green) and pan-laminin (red); **(g)** CD49f (green) and e-cadherin (red); and **(h)** p63 (green) and keratin 8 (red). Loading factors at the negative end of PC1 (epithelial end) included: keratin 19/Krt19 (#5 on list), keratin 8/KRT8 (#6), E-cadherin/CDH1 (#22), Her2/ERBB2 (#85), and keratin 18/KRT18 (#138), and CD49f/ITGA6 (#5574). Loading factors at the positive end of PC1$^{No\ Endo}$ (mesenchymal end) included: smooth muscle actin/ACTA2 (#155 on the list), and Vimentin/VIM (#192). Loading factors at the negative end of PC3 (perivascular end) included: smooth muscle actin/ACTA2 (#7 on the list). Loading factors at the positive end of PC3 (endothelial end) included: CD49f/ITGA6 (#344). Loading factors at the positive end of PC4 (myoepithelial end) included: smooth muscle actin/ACTA2 (#7 on the list), P63/TP63 (#25) and CD49f/ITGA6 (#68).

Figure 10. Pathway analysis of epithelial genes (PC1). (a) Grouped GO/Reactome terms overrepresented within the gene list 200 genes contributing to the negative/epithelial end of PC1 (bottom of the list) are ordered by adjusted p-value. Grouping was done by the similarity of terms to evaluate broad differences in cell class. Associated genes for each group are listed in the 4th column. Median mRNA expression (rlog, log2 scale) for PC1 genes associated with select pathways including **(b)** c*ell adhesion and binding*, **(c)** *peptidase activity*, **(d)** *keratinization*, and **(e)** *cytoskeletal binding*.

Figure 10—figure supplement 1. Pathway analysis of leukocyte genes (PC2). (a) Grouped GO/Reactome terms associated with the 200 genes contributing to the positive end (bottom of list) of PC1 ordered by adjusted p-value. Median mRNA expression (rlog transformed) for PC2

genes associated with select pathways, including **(b)** a*daptive immune response*, **(c)** *immune*
response activation, and **(d)** *leukocyte homeostasis*.

Figure 11. Pathway analysis of mesenchymal genes (PC1$^{No\ Endo}$). (a) Grouped GO/Reactome
terms overrepresented within the gene list 200 genes contributing to the positive/mesenchymal
end of PC1^{NoEndo}, ordered by adjusted p-value. Median mRNA expression (rlog, log2 scale) for
PC1^{NoEndo} genes associated with select pathways, including **(b)** *cardiac tissue development*, **(c)**
fibroblast migration, **(d)** *regulation of peptidyl-tyrosine phosphorylation*, and **(e)**
chondrocyte/skeletal development.

Figure 12. Pathway analysis of Endothelial genes (PC3). (a) Grouped GO/Reactome terms
associated with the 200 genes contributing to the positive end of PC3 (top of list), ordered by
adjusted p-value. Median mRNA expression (rlog, log2 scale) for PC3 genes associated with
select pathways including **(b)** *angiogenesis*, **(c)** *cell motility*, **(d)** c*ell-matrix adhesion*, and **(e)**
endothelial cell differentiation.

Figure 12—figure supplement 1. Transcript levels of genes composing the *signaling by*
***SCF-KIT* pathway.** Transcript levels for each gene in the Reactome pathway s*ignaling by SCF-*
KIT. VST gene levels are relative (normalized within each row of the heatmap). Boxed areas
highlight expression of select genes in lymphatic and vascular endothelial, and ER^{Neg} luminal
epithelial cell types.

Figure 12—figure supplement 2. Transcript levels of genes composing the *Semaphorins*
gene family. (a) Transcript levels of genes in Semaphorin gene family. VST gene levels are
relative (normalized within each row of the heatmap). Boxed areas highlight select genes highly
expressed by the lymphatic and vascular endothelial cell types. Normalized mRNA values of **(b)**
NRP1 and **(c)** NRP2 (rlog, DEseq2) are provided on log2 scale (bar graph of each biological
replicate) and linear scale (donut graph of median value), which are both color-coded by cell
type.

Figure 13. Pathway analysis of perivascular genes (PC3). (a) Grouped GO/Reactome terms
associated with the 200 genes contributing to the negative end of PC3 (bottom of the list),
ordered by adjusted p-value. Median mRNA expression (rlog, log2 scale) for PC3 genes
associated with select pathways, including **(b)** *epithelial tube morphogenesis*, **(c)** *blood vessel*
development, **(d)** *muscle contraction*, and **(e)** *extracellular matrix organization*.

Figure 14. Pathway analysis of myoepithelial-related genes (PC4). (a) Grouped
GO/Reactome terms associated with the 200 genes contributing to the positive end of PC4 (top
of the list), ordered by adjusted p-value. Median mRNA expression (rlog, log2 scale) for PC4
genes associated with select pathways, including **(b)** *extracellular matrix organization*, **(c)** *cell*
junction organization, **(d)** *keratinization*, and **(e)** *morphogenesis of branching epithelium*.

Figure 15—table 1. Variably expressed gene types. Genes composing each gene family
(HGNC) were assessed by DESeq2 (Adj.*p*-value ≤ 1, across all cell types). The number of genes
in each family, along with the numer of those found to be differentially expressed are provided
(for gene families ≥ 6 genes).

Figure 15. Gene types contributing to cell-specific features. Of the 1,519 types of genes (gene families) defined by HGNC, we found 1,421 gene families (of all sizes) expressed within the breast cells transcriptomes. **(a)** Distribution of the proportions of differentially expressed (DE) genes in each gene family (BH Adj. p-value ≤ 0.1). The proportion of DE genes in these families ranged between 0-100%, which we grouped into different bins: 0-24% (blue, n=209), 25%-74% (green, n=814), 75% to 99% (orange, n=211), and 100% (red, n=187). **(b-e)** Median transcript levels (rlog, log2 scale) across all cell types of genes composing select families with **(b)** 100% DE genes, **(c)** 75%-99% DE genes, **(d)** 25-74% DE genes, and **(e)** 0-24% DE genes.

Fig.15—figure supplement 1. Distribution of gene family size. The sizes of the HGNC families range from 1 to 831 genes. The distribution of these family sizes is provided, along with a box & whisker plot (top). The median family size is six genes per family.

Fig.15—figure supplement 2. Transcript levels of DE genes. Transcript levels for select Type IV collagen chains: **(a)** COL4A1, **(b)** COL4A2, and **(c)** COL4A4. **(d)** Median mRNA expression (rlog, log2 scale) for integrin alpha subunits. The dashed circles identify notable differentially expressed genes, ITGA6, ITGAE, and ITGAM. **(e)** Transcript levels of the laminin binding integrins. The predominant laminin-binding integrins are αβ heterodimers of: α3β1, α7β1, α7β1 and α6β4. These are integrins are encoded by ITGB1, ITGA3, ITGA7, ITGA6, and ITGB4. Collagen-binding integrins that also show laminin binding properties include heterodimers of: α1β1, α2β1, and α10β1 (encoded by ITGA1, ITGA2, ITGA10, and ITGB1). ITGA7 is predominantly limited to the perivascular cell types, whereas ITGA3, ITGA6, and ITGB4 are largely expressed by the epithelial cell types.

Figure 16. Tight Junctions (a) Heatmap displaying transcript levels of genes belonging to the claudin gene family (VST, normalized by row). Dashed box indicates claudins that were not appreciably expressed by any cell type. Dark boxes highlight genes differentially expressed by luminal epithelial cells, endothelial cells, and myoepithelial cell types. **(b)** median rlog levels of the claudin family genes. Dashed circles identify genes outlined in the heat map. **(c-e)** Transcript levels for the zona occludens tight junction proteins: **(c)** TJP1, **(d)** TGP2, and **(e)** TJP3 (rlog values on log2 (bar graph) and linear (donut graph) scales).

Figure 16—figure supplement 1. IHC staining. (a) Claudin-1 staining of normal breast tissue (reduction mammoplasty of 28-year-old female). (▼) Claudin-1 staining is found confined to the apical surface of the paracellular space of luminal epithelial cells, circumscribing the luminal surface of the cells. Scale = 50 μm.

Figure 17. Gap junction proteins. (a) Heatmap displaying transcript levels of genes encoding gap junction proteins (VST, normalized by row). Boxed areas highlight the cell types expressing select gap junction proteins. Dashed boxes identify genes not appreciably expressed by any cell type, as evidenced by the rlog values shown in b. **(b)** Median rlog transcript levels of gap junction genes on a log2 scale; differentially expressed genes are encircled.

Figure 18. Receptor tyrosine kinases. (a) Heatmap displaying transcript levels of genes belonging to the tyrosine kinase receptor gene family (VST, normalize by row). Boxed areas

indicate cell types expressing highest levels of select kinases. **(b-m)** Transcript levels of the
select RTKs (rlog values on log2 (bar graph) and linear (donut graph) scales).

Figure 19. Transcript levels of nuclear hormone receptors. (a) Heatmap displaying transcript
levels of genes belonging to the nuclear hormone receptor gene family (VST, normalized by
row). Boxed areas indicate high/low gene expression in different cell types, which are
highlighted below by the donut plots. Select genes include **(b-m)** mRNA values (rlog) are
provided on log2 scale (bar graph of each biological replicate) and linear scale (donut graph of
median value) which are both color-coded by cell type.

Figure 20. Differentially expressed genes across all cell type pairs. The number of
differentially expressed genes between cell-type pairs, as reported by DESeq2; Benjamini-
Hochberg adjusted *p*-values ≤ 0.1.

Figure 21. Differences between ER^{Pos} and ER^{Neg} luminal epithelial cell types. A contrast of
expressed transcripts between the two luminal epithelial cell types revealed 1,301 differentially
expressed genes (DeSeq2/Benjamini-Hochberg adjusted p-value ≤ 0.1). All expressed genes are
displayed. The horizontal dashed lines (separating red and green dots) represents an Adj. *p*-value
of 0.1. Established cell type markers (of each cell type) are denoted. mRNA expression values
for ten genes (indicated by filled, color-coordinated boxes) are shown at left and right of the
volcano plot. Color-coded stars in the bar graphs indicate the cell type being contrasted that had
the higher level of that gene; i.e., ER^{Pos}(light blue) and ER^{Neg} luminal cells (dark blue).

Figure 22. Inspection of GSEA results and the validating example of the *estrogen dependent
***gene expression* pathway. (a)** Gene Set Enrichment Analysis (GSEA) was performed using a
list of genes expressed by ER^{Pos} and ER^{Neg} luminal epithelial cells, ranked by the statistical
significance of their differential expression, *-log$_{10}$ {p-value} * sign([Fold Change])*. The ranked
list, depicted by the shaded bar, has the most statistically significant genes located at its extreme
ends, with the ER^{Pos} cells' most differentially expressed genes represented at the left end and the
ER^{Neg} cells' most differentially expressed genes on the right. GSEA identified the Reactome
pathway *Estrogen dependent gene expression* as one of the the most statistically significant
enriched pathways identified (FDR<0.001). The position of the 89 genes composing the *estrogen*
dependent gene expression pathway within the ranked list of expressed genes are depicted by
vertical black lines. Names and positions of several familiar genes in the pathway are indicated,
e.g., progesterone receptor (PGR) is found at position 337 (out of 21,085 expressed genes). The
GSEA algorithm assessed the distribution of the pathway's 89 genes within the ranked list using
a running sum score (green trace), calculating statistics by comparing results to analysis of a
randomized ranked list (1000 permutations). **(b)** Heat map depicting transcript levels within
ER^{Pos} and ER^{Neg} luminal epithelial cells of the 89 genes belonging to *estrogen dependent gene*
expression pathway, ordered by their relative position in the ranked list (VST values normalized
by row). Positions of the familiar genes are indicated, including SP1 (which was near the bottom
of the list and was not differentially expressed). Transcript levels of PGR, ESR1, and SP1 in
each cell type (rlog values on log2 (bar graph) and linear (donut graph) scales).

Figure 23. ER^{Pos} and ER^{Neg} luminal cells' enriched pathways. Visualization of pathways
enriched among ER^{Pos} and ER^{Neg} luminal cells (Enrichment Map/Cytoscape). All enriched
pathways with a FDR≤0.1 appear on the map (nodes, color coded by cell type). Pathways(nodes)
are connected when they share 50% of their genes (or greater). Node size corresponds to the
number of genes in the pathway, as indicated (bottom left). Biologically related pathways
sharing genes formed clusters in the network (biological themes), and different themes with a
common biological function (e.g., metabolism-, immune -, and ECM-related, etc.) were typically
placed adjacent to each other (when possible). Select themes were encircled and annotated.
Highly differentially expressed genes are indicated (above the theme when it coincided with the
attributed cell type—and below if it was expressed by the contrasted cell type). The names of
each pathway are available in the supplemental Cytoscape file and Pathway Analyzer tool.

Figure 23—figure supplement 1. Transcript levels from enriched pathways. (a-t) Transcript
levels of leading-edge genes from select pathways enriched in ER^{Pos} or ER^{Neg} luminal cells (from
325 enriched pathways with a FDR of ≤ 10%, GSEA). VST gene levels are relative (normalized
by row). Arrowheads under the heatmaps identify the cell type either over- or under-expressing
the pathway's genes. Samples are color-coded. Due to space constraints and the large number of
presented genes, their names (rows of the heatmap) must be inspected digitally (magnified in the
PDF).

Figure 23—figure supplement 2. Transcript levels from enriched pathways. (a-e) Transcript
levels of leading-edge genes from select pathways enriched in ER^{Pos} or ER^{Neg} luminal cells (from
325 enriched pathways with a FDR of ≤ 10%, GSEA). VST gene levels are relative (normalized
by row). Arrowheads under the heatmaps identify the cell type either over- or under-expressing
the pathway's genes. Samples are color-coded. Due to space constraints and the large number of
presented genes, their names (rows of the heatmap) must be inspected digitally (magnified in the
PDF).

Figure 23—figure supplement 3. Transcript levels of toll-like receptors (a) Transcript levels
of TLR gene family. **(b-f)** Normalized mRNA values (rlog, DEseq2) provided on log2 scale (bar
graph of each biological replicate) and linear scale (donut graph of median value) which are both
color-coded by cell type.

Figure 23—figure supplement 4. Transcript levels of milk proteins. (a) Transcript levels of
milk proteins. VST gene values are relative (normalized by row). **(b-j)** Normalized mRNA
values (rlog, DEseq2) provided on log2 scale (bar graph of each biological replicate) and linear
scale (donut graph of median value), which are both color-coded by cell type.

Figure 24. Differences between ER^{Neg} luminal and myoepithelial cell types. A contrast of
expressed transcripts between ER^{Neg} luminal and myoepithelial cells revealed 2,239
differentially expressed genes (DeSeq2/Benjamini-Hochberg adjusted p-value ≤ 0.1). All
expressed genes are displayed. The horizontal dashed lines (separating red and green dots)
represents an Adj. *p*-value of 0.1. Established cell type markers (of each cell type) are denoted.
mRNA expression values for ten genes (indicated by filled, color-coordinated boxes) are shown
at left and right of the volcano plot. Color-coded stars in the bar graphs indicate the cell type

being contrasted that had the higher transcript level, i.e., ER^{Neg}(blue) and myoepithelial cells

(red).

Figure 25. The ER^{Neg} luminal and myoepithelial cells' enriched pathways. Visualization of

pathways enriched among ER^{Pos} and myoepithelial cell types (Enrichment Map/Cytoscape). All

enriched pathways with a FDR≤0.1 appear on the map (nodes, color coded by cell type).

Pathways(nodes) are connected when they share 50% of their genes (or greater). Node size

corresponds to the number of genes in the pathway, as indicated (bottom left). Select biological

themes were encircled and annotated. Highly differentially expressed genes are indicated (above

the theme when it coincided with the attributed cell type—and below if it was expressed by the

contrasted cell type). The names of each pathway are available in the supplemental Cytoscape

file and Pathway Analyzer tool.

Figure 25—figure supplement 1. Transcript levels from enriched pathways. Transcript

levels of leading-edge genes from select pathways enriched in ER^{Pos} or myoepithelial cells (from

426 enriched pathways with a FDR of ≤ 10%, GSEA). VST gene levels are relative (normalized

by row). Arrowheads under the heatmaps identify the cell type, either over- or under-expressing

the pathway's genes. Samples are color-coded. Due to space constraints and the large number of

presented genes, their names (rows of the heatmap) must be inspected digitally (magnified in the

PDF).

Figure 26. Differences between ER^{Neg} luminal and vascular endothelial cell types. A contrast

of expressed transcripts between ER^{Neg} luminal and vascular endothelial cells revealed 4,167

differentially expressed genes (DeSeq2/Benjamini-Hochberg adjusted p-value ≤ 0.1). All

expressed genes are displayed. The horizontal dashed lines (separating red and green dots)

represent an adj. *p*-value of 0.1. Established cell type markers (of each cell type) are denoted.

mRNA expression values for ten genes (indicated by filled, color-coordinated boxes) are shown

at the left and right of the volcano plot. Color-coded stars in the bar graphs indicate the cell type

being contrasted that had the higher transcript level, i.e., ER^{Neg}(blue) and vascular endothelial

cells (orange).

Figure 27. The ER^{Neg} luminal and vascular endothelial cells' enriched pathways.

Visualization of pathways enriched among ER^{Pos} luminal and vascular endothelial cell types

(Enrichment Map/Cytoscape). All enriched pathways with a FDR≤0.1 appear on the map (nodes,

color-coded by cell type). Pathways(nodes) are connected when they share 50% of their genes

(or greater). Node size corresponds to the number of genes in the pathway, as indicated (bottom

left). Select biological themes were encircled and annotated. Highly differentially expressed

genes are indicated (above the theme when it coincided with the attributed cell type—and below

if it was expressed by the contrasted cell type). Samples are color-coded. Due to space

constraints and a large number of presented genes, their names (rows of the heatmap) must be

inspected digitally (magnified in the PDF). The names of each pathway are also available in the

supplemental Cytoscape file and Pathway Analyzer tool.

Figure 27—figure supplement 1. Transcript levels from enriched pathways. Transcript

levels of leading-edge genes from select pathways enriched in ER^{Pos} luminal and vascular

endothelial cells (from 500 enriched pathways with a FDR of ≤ 10%, GSEA). VST gene levels

are relative (normalized by row). Arrowheads under the heatmaps identify the cell type, either
over- or under-expressing the pathway's genes.

Figure 28. Differences between lymphatic and vascular endothelial cell types. A contrast of
expressed transcripts between lymphatic and vascular endothelial cells revealed 3,739
differentially expressed genes (DeSeq2/Benjamini-Hochberg adjusted p-value ≤ 0.1). All
expressed genes are displayed. The horizontal dashed lines (separating red and green dots)
represent an adj. *p*-value of 0.1. Established cell type markers (of each cell type) are denoted.
mRNA expression values for ten genes (indicated by filled, color-coordinated boxes) are shown
at left and right of the volcano plot. Color-coded stars in the bar graphs indicate the cell type
being contrasted that had the higher transcript level, i.e., lymphatic (light orange) and vascular
(dark orange) endothelial cells.

Figure 29. The lymphatic and vascular endothelial cells' enriched pathways. Visualization
of pathways enriched among lymphatic and vascular endothelial cell types (Enrichment
Map/Cytoscape). All enriched pathways with a FDR≤0.1 appear on the map (nodes, color-coded
by cell type). Pathways(nodes) are connected when they share 50% of their genes (or greater).
Node size corresponds to the number of genes in the pathway, as indicated (bottom left). Select
biological themes were encircled and annotated. Highly differentially expressed genes are
indicated (above the theme when it coincided with the attributed cell type—and below if it was
expressed by the contrasted cell type). Samples are color-coded. Due to space constraints and a
large number of presented genes, their names (rows of the heatmap) must be inspected digitally
(magnified in the PDF). The names of each pathway are also available in the supplemental
Cytoscape file and Pathway Analyzer tool.

Figure 29—figure supplement 1. Transcript levels from enriched pathways. Transcript
levels of leading-edge genes from select pathways enriched in lymphatic and vascular
endothelial cell (from 240 enriched pathways with a FDR of ≤ 10%, GSEA). VST gene levels
are relative (normalized by row). Arrowheads under the heatmaps identify the cell type, either
over- or under-expressing the pathway's genes.

Figure 30. Differences between fibroblasts and adipocytes. A contrast of expressed transcripts
between fibroblasts and adipocytes revealed 1,171 differentially expressed genes
(DeSeq2/Benjamini-Hochberg adjusted p-value ≤ 0.1). All expressed genes are displayed. The
horizontal dashed lines (separating red and green dots) represent an adj. *p*-value of 0.1.
Established cell type markers (of each cell type) are denoted. mRNA expression values for ten
genes (indicated by filled, color-coordinated boxes) are shown at left and right of the volcano
plot. Color-coded stars in the bar graphs identify the position of the fibroblasts (yellow) and
adipocytes (brown).

Figure 31. The fibroblasts and adipocytes' enriched pathways. Visualization of pathways
enriched among fibroblasts and adipocytes (Enrichment Map/Cytoscape). All enriched pathways
with a FDR≤0.1 appear on the map (nodes, color-coded by cell type). Pathways(nodes) are
connected when they share 50% of their genes (or greater). Node size corresponds to the number
of genes in the pathway, as indicated (bottom left). Select biological themes were encircled and
annotated. Highly differentially expressed genes are indicated (above the theme when it

coincided with the attributed cell type—and below if it was expressed by the contrasted cell
type). The names of each pathway are available in the supplemental Cytoscape file and Pathway
Analyzer tool. Samples are color-coded. Due to space constraints and a large number of
presented genes, their names (rows of the heatmap) must be inspected digitally (magnified in the
PDF).

Figure 31—figure supplement 1. Transcript levels from enriched pathways. Transcript
levels of leading-edge genes from select pathways enriched in fibroblasts and adipocytes (from
387 enriched pathways with a FDR of $\leq 10\%$, GSEA). VST gene levels are relative (normalized
by row). Arrowheads under the heatmaps identify the cell type, either over- or under-expressing
the pathway's genes.

Figure 32. Differences between pericytes and fibroblasts. A contrast of expressed transcripts
between pericytes and fibroblasts revealed 2,932 differentially expressed genes
(DeSeq2/Benjamini-Hochberg adjusted p-value ≤ 0.1). All expressed genes are displayed. The
horizontal dashed lines (separating red and green dots) represent an adj. *p*-value of 0.1.
Established cell type markers (of each cell type) are denoted. mRNA expression values for ten
genes (indicated by filled, color-coordinated boxes) are shown at the left and right of the volcano
plot. Color-coded stars in the bar graphs identify the position of the pericytes (purple) and
fibroblasts (yellow).

Figure 33. The pericytes and fibroblasts' enriched pathways. Visualization of pathways
enriched among pericytes and fibroblasts (Enrichment Map/Cytoscape). All enriched pathways
with a FDR≤ 0.1 appear on the map (nodes, color-coded by cell type). Pathways(nodes) are
connected when they share 50% of their genes (or greater). Node size corresponds to the number
of genes in the pathway, as indicated (bottom left). Select biological themes were encircled and
annotated. Highly differentially expressed genes are indicated (above the theme when it
coincided with the attributed cell type—and below if it was expressed by the contrasted cell
type). Samples are color-coded. Due to space constraints and a large number of presented genes,
their names (rows of the heatmap) must be inspected digitally (magnified in the PDF). The
names of each pathway are also available in the supplemental Cytoscape file and Pathway
Analyzer tool.

Figure 33—figure supplement 1. Transcript levels from enriched pathways. Transcript
levels of leading-edge genes from select pathways enriched in pericytes and fibroblasts (from
477 enriched pathways with a FDR of $\leq 10\%$, GSEA). VST gene levels are relative (normalized
by row). Arrowheads under the heatmaps identify the cell type, either over- or under-expressing
the pathway's genes.

A cellular and transcriptomic dissection of the human breast for studying mechanisms of cell and tissue function

Katelyn Del Toro, Rosalyn Sayaman, Kate Thi, Yamhilette Licon-Munoz, and William C. Hines

Figures

Figure 1

a

Breast Tissue → All Cells

All Cells —(Lipid Layer)→ Adipocytes (Pop12)

FACS (Organoid fraction)

CD34 + → CD49f − → Podoplanin + → Lymphatic Endothelial (Pop7)

Podoplanin − → Endothelial (Pop9)

CD34 − → MSC/Fibroblast (Pop8)

Thy1 + → CD49f + → CD10 + → Myoepithelial (Pop3)

CD49f − → Pericytes (Pop6)

CD24 ∧ Muc1 + → CD49f + → Luminal #2 (Pop2)

CD49f Low → Luminal #1 (Pop1)

CD49f + → Epithelial (Pop4)

CD45 + → Leukocytes (Pop10)

Podoplanin Low → vSMC (Pop5)

RBCs (Pop11)

b

Relative Abundance vs Cell Type

Legend:
1. Luminal 1 (ER^Pos)
2. Luminal 2 (ER^Neg)
3. Myoepithelial
4. Epithelial
5. vSMCs
6. Pericytes
7. Lymphatic Endothelial
8. Fibroblasts
9. Vascular Endothelial
10. CD45+ Leukocytes
11. Erythrocytes
12. Adipocytes

Figure 1—figure supplement 1

BD FACS VANTAGE
8/8 channels used

Figure 2

a

Arteriolar vSMCs **5**

Capillary pericytes **6**

Lymphatic Vessels **7**

Stromal Fibroblasts **8**

Blood Vessels
Endothelial Cells, Leukocytes, RBCs
9,10,11

Lobules (TDLU)
Epithelium, fibroblasts, capillaries
1,2,3,4, 6,8,10,11

Lactiferous Duct
1,2,3,10

Adipose **12**

b

Epithelial   **Mesenchymal**   **Blood**

Glandular   Endothelial

Luminal   ME   Novel   Peri-endothelial

ER+ Luminal Epithelial	ER^Neg Luminal Epithelial	Myoepithelial	Rare Epithelial	Lymphatic Endothelial	Vascular Endothelial	vSMCs	Pericytes	Fibroblasts	Adipocytes	Leukocytes (combined)	Erythrocytes
1	**2**	**3**	**4**	**7**	**9**	**5**	**6**	**8**	**12**	**10**	**11**

Cell Population ID#s

c
8 6,9 12
5,9
1,2,3,4, 6,8,9
1
2 3
Keratin18 Pan-Laminin DAPI

d
7 9
6
7
9
6
Podoplanin Pan-Laminin DAPI

e
8
12 6,9 8
6,9 8
12
3 8 12
1,2 12 5,9
Pan-Laminin CD34 DAPI

f
12 12
5 9 5
5
9 7 6
6
8 8 6
7 9 6
7 12
CD49f Pan-Laminin DAPI

Figure 2—figure supplement 1

Figure 2—figure supplement 2

Figure 3

Distinguishing the traits of each breast cell type

a Identification and purification of 12 cell types composing the breast

b RNA-sequencing & Validation

c Covariance, Clustering, PCA, and identification of lineage differences

PCA/Pathway Enrichment Analysis

d Differential Gene Expression Analysis

e Gene Set Enrichment Analysis

f Distinguishing features (of each cell type)

g Pathway Analyzer & Online Tool

Figure 3—figure supplement 1

Figure 3—table supplements (supplemental files)

List of Table supplements and Data	
Figure 3—table supplement 1	Sample Treatment.xlsx
Figure 3—table supplement 2	CountMatrix_rLog_VST.xlsx
Figure 3—table supplement 3	PCA_Loading_Factors.xlsx
Figure 3—table supplement 4	GO from PCAs.xlsx
Figure 3—table supplement 5	DESeq2_LRT.xlsx
Figure 3—table supplement 6	Gene Families_DE Genes.xlsx
Figure 3—table supplement 7	DESeq2_files.xlsx
OSF; https://osf.io/gfhc3/	Cytoscape/Enrichment Map files (1v2, 2v3, 2v9, 6v8)
OSF; https://osf.io/hpb9t/	Pathway Analyzer[9]
http://breastcancerlab.com/breast-atlas	Breast Cell Atlas: Transcript Finder (R/Shiny App)[10]

Figure 4

a

		N222	N225	N228	N239
	Cell Populations sequenced	29 y/o ♀	16 y/o ♀	27 y/o ♀	23 y/o ♀
Pop1	Luminal 1 (ER+)	1		1	1
Pop2	Luminal 2 (ER-)	2		2	2
Pop3	Myoepithelial	3		3	3
Pop4	Epithelial		4	4	4
Pop5	Vascular Smooth Muscle Cells			5	5
Pop6	Pericytes		6	6	6
Pop7	Lymphatic Endothelial Cells			7	7
Pop8	Fibroblasts	8		8	8
Pop9	Vascular Endothelial Cells	9		9	9
Pop10	Leukocytes		10	10	10
Pop11	Red Blood Cells *no RNA / NS	-	-	-	-
Pop12	Adipocytes	12		12	12
Pop13	Unsorted Cells (dissociated, pre-FACS)	13		13	13
Pop14	Tissue			14	14

b

Covariance Matrix

c

Regularized log values

rlog values

Reads per Sample

d

Sample Correlations (Ordered by Individual)

Individual-Cell Population

a

Legend:
1. Luminal 1 (ER^Pos)
2. Luminal 2 (ER^Neg)
3. Myoepithelial
4. Epithelial
5. vSMCs
6. Pericytes
7. Lymphatic Endothelial
8. Fibroblasts
9. Vascular Endothelial
10. CD45+ Leukocytes
11. Erythrocytes
12. Adipocytes
13. Cells (unsorted)

b

		ANTIBODY STAINING IN SORTING SCHEME								
Pop ID	Cell Type	CD36†	CD34	CD49f (ITGA6)	PDPN	CD90 (Thy1)	CD10 (MME)	CD24	CD227 (MUC1)	CD45 (PTPRC)
1	ER^Pos Luminal Epithelial		Neg	-/+	Neg	Neg	Neg	+++	+++	Neg
2	ER^Neg Luminal Epithelial		Neg	++	Neg	Neg	Neg	+++	+++	Neg
3	Myoepithelial		Neg	++++	++	+++	+++	Neg	Neg	Neg
4	Pop4 Epithelial		Neg	++	+	Neg	-/+	Neg	Neg	Neg
5	Vascular Smooth Muscle Cells		Neg	Neg	-/+	Neg	Neg	Neg	Neg	Neg
6	Pericytes		Neg	Neg	Neg	+++	Neg	Neg	Neg	Neg
7	Lymphatic Endothelial Cells		+++	++	+++	-/+	Neg	Neg	Neg	Neg
8	Adipocyte Derived Mesenchymal Stem Cells (Fibroblasts)		+++	Neg	+	++	+	Neg	Neg	Neg
9	Vascular Endothelial Cells		+++	++	Neg	-/+	Neg	Neg	Neg	Neg
10	Leukocytes (CD45^Pos)		Neg	Neg	Neg	Neg	-/+	Neg	Neg	+++
11	Red Blood Cells		Neg	Neg	Neg	Neg	Neg	Neg	Neg	Neg
12	Adipocytes	+++								

c

glyceraldehyde-3-phosphate dehydrogenase

Figure 5

Figure 5-figure supplement 1

HGNC Gene Family 471: CD Molecules

Mean VST Expression Value

1. Luminal 1
2. Luminal 2
3. Myoepithelial
4. Epithelial
5. vSMCs
6. Pericytes
8. Fibroblasts
12. Adipocytes
7. Lymphatic Endothelial
9. Vascular Endothelial
10. CD45+ Leukocytes

Figure 6

Top 2000 Differentially Expressed Genes

Density

Row Z-Score

-3 -1 1 3

1. Luminal 1 (ER^Pos)
2. Luminal 2 (ER^Neg)
3. Myoepithelial
4. Epithelial
5. vSMCs
6. Pericytes
7. Lymphatic Endothelial
8. Fibroblasts
9. Vascular Endothelial
10. CD45+ Leukocytes
11. Erythrocytes
12. Adipocytes
13. Cells (unsorted)
14. Tissue

10 7 9 8 5 6 2 1 4 3 13 12 14

Leuk Endo Mesen. Epithelial Adip/Tissue

Figure 7

Figure 7-figure supplement 1

Figure 7-figure supplement 2

Figure 8

Figure 9

Figure 10

a

Principal Component 1: Epithelial Cell Types

ER+ Luminal	ER- Luminal	Myoepithelial	Epithelial

(# pathways in group)	Representative Pathways	Adj. *p*-value	Associated Genes (from PC1 BOTTOM 200) Epithelial
2	cell adhesion molecule binding & cadherin binding	1.40E-08	[ANK3, BAIAP2L1, CDH1, CDH3, CGN, CLDN7, DSG2, DSP, EPCAM, EPS8L2, ITGB6, ITGB8, KRT18, LAD1, LAMB3, NECTIN4, OLFM4, PKP2, PKP3, PPL, PROM1, PTPRF, SFN, SHTN1, SPP1]
63	peptidase activity, hydrolase activity, Regulation of kinase activity…	6.05E-05	[ARHGEF5, ATP1B1, CLDN3, CLDN4, DHCR24, EFNA1, ERBB2, ESR1, LTF, PERP, PPP1R1B, PROM2, RARRES1, S100A8, S100A9, SERPINA1, SERPINA3, SERPINB3, SERPINB4, SERPINB7, SFN, SLPI, SPINT1, SPINT2, SYTL2, TFAP2B, TNFSF10, TPM1, VAV3, WFDC2, XDH]
9	keratinization, programmed cell death…	8.32E-05	[DSC2, DSG2, DSP, KRT15, KRT16, KRT18, KRT19, KRT23, KRT6B, KRT7, KRT8, PERP, PKP2, PKP3, PPL, PRSS8, ST14]
3	actin filament binding, cytoskeletal protein binding, & actin binding	2.68E-03	[ANK3, BAIAP2L1, CDH1, CDH3, CGN, CLU, COBL, EPB41L4B, EPS8L2, HOMER2, HOOK1, LIMCH1, MYH14, MYO5B, PLS1, PPP1R9A, PROM1, PRSS8, PSTPIP2, RAB27B, S100A8, S100A9, SHROOM3, SHTN1, TPM1]
34	RTK signaling, ESR signaling, regulation by TFAP (AP-2) …	5.58E-03	[CYP27B1, EHF, ELF3, ESR1, FOXA1, GATA3, GRHL1, MUC1, RORC, SOX10, SOX9, TFAP2A, TFAP2B, TFAP2C, VDR]
1	calcium ion binding	6.13E-03	[ANXA3, CDH1, CDH3, CELSR2, CLU, DCHS2, DSC2, DSG2, GALNT3, LRP2, MGP, PADI2, PLS1, RASEF, S100A14, S100A8, S100A9, SYT7, TPD52]
5	Metabolism of vitamins and cofactors, ECM organization, glycosylation, vascular wall interactions, non-integrin ECM interactions	6.91E-03	[CDH1, DDR1, ITGB6, ITGB8, LAMB3, MMP7, SDC1, SDC4, SPP1]
2	Innate immune system	7.77E-03	[CFB, CHI3L1, CLU, DSP, FRK, LCN2, LTF, MGST1, MUC1, MUC4, OLFM4, PADI2, PIGR, S100A8, S100A9, SAA1, SERPINA1, SERPINA3, SERPINB, SLPI, TLR2, VAMP8, VAV3, VNN1]

Principal Component 2: Leukocytes (Mesodermal)

Leukocytes

a

(# pathways in group)	Representative Pathways	*Adj. p*-value	Associated Genes (from PC2 <u>BOTTOM</u> 200) Leukocytes
6	adaptive immune response, lymphocyte activation, regulation of immune response …	6.07E-46	[BTK, CAMK4, CD247, CD28, CD3D, CD3E, CD48, CD6, CD7, CD74, CD79A, CD8A, CD8B, GPR183, HLADRA, HLA-DRB1, HLA-DRB5, IGLL5, IL7R, INPP5D, IRF4, ITK, JCHAIN, KLRK1, LY9, MYO1G, PIK3CG, PRF1, PRKCB, PRKCQ, PTK2B, PTPRC, SAMSN1, SASH3, SKAP1, SLA2, SLAMF6, SLAMF7, SPN, TARP, THEMIS, TNFRSF13C, WAS, ZAP70]
15	activation of immune response, T cell receptor signaling pathway, B cell receptor signaling pathway…	1.78E-41	[CARD11, CCR7, CD247, CD28, CD3D, CD3E, FYB1, HLADRA, HLA-DRB1, HLA-DRB5, INPP5D, ITK, LCK, LCP2, PRKCQ, PTPN22, PTPRC, SKAP1, SLA2, TARP, TESPA1, THEMIS, WAS, ZAP70] ; [BLK, BTK, CD22, CD79A, IGLL5, ITK, LCK, LPXN, MS4A1, NCKAP1L, PAX5, PRKCB, PTPN22, PTPRC]
141	leukocyte homeostasis, cytokine production, hemopoiesis, interleukin-10, -4, -2, -8 production, cell-cell adhesion…	2.42E-41	[CD74, CORO1A, DOCK10, DOCK11, GPR183, IL2RA, NCKAP1L, TNFRSF13C, TSC22D3]
25	T cell activation involved in immune response, Interleukin-17 production, T-helper 17 type immune response…	1.94E-26	[ADAM8, APBB1IP, ARHGAP45, ARHGAP9, ATP8A1, BIN2, BTK, CD28, CD53, CD74, CORO1A, DOCK10, DOCK11, DOCK2, GPR183, IQGAP2, IRF4, ITGAL, ITGB2, LCP1, LRMP, LY9, NCKAP1L, PIK3CG, PTGER4, PTK2B, PTPRC, RAC2, RASGRP1, SELL, SLAMF6, SPN]
53	leukocyte chemotaxis and migration, T cell chemotaxis, cellular extravasation…	5.88E-23	[ADAM8, CCL5, CCR5, CCR7, CD74, CORO1A, CXCR4, GPR183, IL16, ITGB2, JAML, KLRK1, NCKAP1L, PIK3CG, PREX1, PTK2B, RAC2, RIPOR, STK4, VAV1]
6	phagocytosis, Fc-gamma receptor signaling pathway involved in phagocytosis…	8.85E-22	[BIN2, CD247, CORO1A, CYFIP2, DOCK2, IGLL5, IL2RB, IL2RG, ITGAL, ITGB2, MYO1G, NCKAP1L, PTPRC, RAC2, TARP, VAV1, WAS, WIPF1]
36	Interleukin-6 production, interleukin-1 secretion…	5.64E-17	[BANK1, CD74, IL16, INPP5D, LCP1, NCKAP1L, PTPN22]
10	acute inflammatory response…	3.69E-14	[ADAM8, ALOX5AP, BTK, CCR7, CD6, PIK3CG]
9	regulation of GTPase activity, Ras - and Rho- guanyl-nucleotide exchange factor activity…	1.29E-11	[ACAP1, ARAP2, ARHGAP15, ARHGAP30, ARHGAP45, ARHGAP9, ARHGDIB, CCL5, CCR7, DOCK10, DOCK11, DOCK2, DOCK8, FGD3, IQGAP2, KLRK1, NCKAP1L, PREX1, PTK2B, RASAL3, RASGRP1, RASGRP2, RGS1, RGS2, RHOH, SMAP2, TAGAP, VAV1]
19	mast cell degranulation, inflammatory response to antigenic stimulus…	3.44E-11	[BTK, CCR7, CD28, CD6, HLADRB1, IL2RA, RASGRP1]
6	monocyte differentiation…	9.53E-11	[EVI2B, HCLS1, INPP5D]

Figure 11

a. Principal Component 1b: Mesenchymal Cell Types

vSMCs	Pericytes	Fibroblasts	Adipocytes

(# pathways in group)	Representative Pathways	Adj. p-value	Associated Genes (from PC1 TOP 200) Mesenchymal
94	cardiac muscle tissue development, septum development, cartilage morphogenesis, regulation of vSMC proliferation …	5.00E-33	[ADAMTS1, CPE, HEG1, HEYL, MEF2C, MYOCD, NRP2, SHOX2, SLIT3, TBX2, TGFBR3, ZFPM2]
85	Fibroblast migration, endothelial migration, epithelial migration, regulation of angiogenesis…	5.37E-31	[ADAMTS1, ADAMTS9, ADGRA2, ANGPT2, COL4A2, CXCL8, DCN, EMILIN1, ETS1, FGF2, GPR4, HGF, HSPB6, IL6, ITGA5, PTGS2, SERPINE1, SPARC, TCF4, THBS1, THBS2, THBS4, TWIST1]
77	positive regulation of peptidyl-tyrosine phosphorylation, cardiac epithelial to mesenchymal transition, GAG metabolic process, regulation of chemokine production, …	7.40E-21	[BMP6, CSPG4, FGF7, FYN, HGF, IGF2, IL24, IL6, ITGA5, ITGB3, LIF, NR4A3, THBS4]
35	chondrocyte, cartilage, skeletal system development; regulation of mesenchymal cell proliferation …	1.67E-18	[ADAMTS12, ADAMTS4, BGN, BMP6, CDH11, CHST11, COL1A2, COL3A1, COL5A2, COL6A1, COL6A2, COL6A3, FGF2, FGFR1, GLI2, IGF2, MEF2C, MMP2, PDGFRB, PRRX1, SATB2, SFRP4, SHOX2, SPARC, TBX15, TWIST1, ZEB1]
31	bone development, bone morphogenesis, regulation of histone acetylation…	4.53E-18	[BGN, BMP6, COL6A1, COL6A2, COL6A3, MEF2C, SFRP4, SHOX2, SPARC, TWIST1]
27	positive regulation of blood vessel endothelial cell migration…	1.55E-19	[ETS1, FGF2, FGFR1, PTGS2, THBS1]
25	smooth muscle contraction, vascular smooth muscle contraction…	4.58E-12	[ACTA2, CAV1, EDNRA, EDNRB, MYH11, MYOCD, PDE4D, PTGS2, SSTR2]
19	negative regulation of calcium ion transport, muscle contraction…	1.29E-11	[CAV1, GEM, KCNE4, PLN, PTGS2, SFRP4, SYT11, THBS1, TWIST1]
18	regulation of fatty acid biosynthetic process, prostaglandin metabolic process…	3.84E-12	[AVPR1A, PDK4, PTGS2]
17	glycosaminoglycan catabolic process, dermatan sulfate biosynthesis, positive chemotaxis…	4.01E-08	[ANGPT2, CXCL8, FGF2, FGF7, HGF, IL6]

b. Genes associated with Cardiac Chamber Development

c. Genes associated with Fibroblast Migration

d. Genes associated with Regulation of Peptidyl-Tyrosine Phosphorylation

e. Genes associated with Chondrocyte/Skeletal Development

Figure 12

a   Principal Component 3: Endothelial Cell Types

Lymphatic Endothelial   **Vascular**

(# pathways in group)	Representative Pathways	Adj. *p*-value	Associated Genes (from PC3 TOP 200) Endothelial
65	Signaling by VEGF, endothelial cell proliferation, angiogenesis, positive regulation of vasculogenesis…	6.27E-19	[ACVRL1, ANGPT2, ARHGEF15, CALCRL, CARD10, CD34, CDH13, CDH5, CLDN5, CTSH, CX3CL1, DYSF, ECSCR, EFNA1, EFNB2, EGFL7, FLT4, HSPB1, JCAD, KDR, KIT, NDST1, NEDD4, PDGFB, PDPN, PECAM1, PPP1R16B, PRCP, PROX1, PTPRB, RAMP2, ROBO4, S1PR1, SHB, SMAD1, SOX18, STAB1, TAL1, TBX1, TIE1]
8	regulation of cell motility, regulation of cell migration…	1.33E-16	[ABI3, ACVRL1, AFDN, ANGPT2, CARD10, CD200, CDH13, CDH5, CLDN5, CTSH, CX3CL1, DMTN, DOCK4, DYSF, EFNA1, FLT4, HSPB1, HYAL2, JCAD, KDR, KIT, LDB2, LGALS9, LIMCH1, LRRC15, LYN, MCTP1, PDGFB, PDPN, PECAM1, PODXL, PPM1F, PRCP, PROX1, RELN, ROBO4, S1PR1, SELE, SEMA3A, SEMA3D, SEMA6B, SSH2, STC1, TIE1, TNFRSF14]
18	cell-matrix adhesion, regulation of chemotaxis, positive regulation of blood vessel endothelial cell migration…	5.83E-14	[ACVRL1, CD34, CDH13, CX3CL1, DMTN, KDR, LDB2, LIMCH1, LYPD5, LYVE1, PECAM1, PPM1F]
16	Endothelial-, muscle- cell differentiation, lymph vessel morphogenesis, lymphangiogenesis , aorta development…	9.18E-11	[ACVRL1, AFDN, CDH5, CLDN5, F11R, KDR, PDPN, PECAM1, PPP1R16B, PROX1, ROBO4, S1PR1, SOX18, STC1]
41	Cell-cell junction assembly, regulation of vascular permeability, tight junction organization, regulation of establishment of endothelial barrier, endothelial cell apoptotic process…	1.61E-12	[ACVRL1, AFDN, CDH13, CDH5, CLDN5, DCHS1, DMTN, EFNB2, F11R, FSCN1, KDR, LDB2, LIMCH1, MPP7, PECAM1, PPM1F, PRKCH, RAMP2, SHANK3]
12	regulation of endothelial cell migration, negative regulation of chemotaxis, negative regulation of blood vessel endothelial cell migration…	1.38E-11	[ACVRL1, ANGPT2, CARD10, EFNA1, FLT4, HSPB1, JCAD, KDR, PDGFB, PRCP, PROX1, STC1]
40	regulation of coagulation, regulation of hemostasis, regulation of platelet activation, positive regulation of phosphatidylinositol 3 -kinase activity…	9.22E-09	[CD34, DMTN, F11R, LYN, PDGFB, PDPN, TLR4] [AFDN, CNKSR3, CX3CL1, DMTN, EFNB2, F11R, LGALS9, LYN, MAGI1, PDPN, PODXL, SELE, TNFRSF14]
9	regulation of blood vessel endothelial cell migration, regulation of cell - matrix adhesion, negative regulation of angiogenesis…	1.27E-08	[ACVRL1, ANGPT2, CARD10, EFNA1, HSPB1, JCAD, KDR, PDGFB, PRCP]
15	Signaling by VEGF, regulation of vascular permeability, vasculogenesis…	3.53E-08	[CDH5, FLT4, HSPB1, KDR, MAPK11, SHB]
37	peptidyl-tyrosine modification, Signaling by SCF -KIT, regulation of lipid kinase activity, monocyte chemotaxis…	4.93E-08	[BMX, CSF2, CSF3, DMTN, EFNA1, FLT4, GPRC5B, HYAL2, KDR, KIT, LYN, PDGFB, PECAM1, RELN, STAP2, TAL1, TIE1, TNFRSF14]

b Genes associated with Angiogenesis

c Genes associated with Cell Motility

d Genes associated with Cell-Matrix Adhesion

e Genes associated with Endothelial Cell Differentiation

Figure 12—figure supplement 1

Figure 12—figure supplement 2

Semaphorins

HGNC: SEMA

The heatmap rows (top to bottom): SEMA4F, SEMA3E, SEMA4A, SEMA3F, SEMA4D, SEMA7A, SEMA6B, SEMA3G, SEMA4G, SEMA5B, SEMA6C, SEMA3B, SEMA3D, SEMA3A, SEMA4B, SEMA4C, SEMA6A, SEMA5A, SEMA6D, SEMA3C

1. Luminal 1
2. Luminal 2
3. Myoepithelial
4. Epithelial
5. vSMCs
6. Pericytes
8. Fibroblasts
12. Adipocytes
7. Lymphatic Endothelial
9. Vascular Endothelial
10. CD45+ Leukocytes

Figure 13

Principal Component 3: Perivascular Cell Types

a

| vSMCs | Pericytes |

(# pathways in group)	Representative Pathways	Adj. *p*-value	Associated Genes (from PC3 <u>BOTTOM</u> 200) Perivascular
39	Epithelial tube morphogenesis, neuron fate commitment, branching morphogenesis of an epithelial tube, cell fate commitment…	2.33E-17	[DLG5, EDNRA, FGF2, GLI2, GLI3, IRX1, LMO4, NFATC4, SPRY2, STARD13, TBX2, TBX3, TNC, TWIST1, ZEB2]
5	Blood vessel development, vasculature development, urogenital system development, tissue morphogenesis, tube development	4.25E-16	[ACTA2, ADAMTS1, AGTR1, ANTXR1, APOE, APOLD1, COL1A2, CSPG4, EDNRA, ENPEP, FGF2, GJA5, GLI3, HSPB6, ITGB3, MMP19, MYOCD, NFATC4, NOTCH3, NR4A1, NTRK2, PDGFRB, PRRX1, RUNX1, SEMA3C, SMAD7, SPRY2, STARD13, TBX2, TBX3, TGFBI, TWIST1]
9	Muscle contraction, regulation of blood vessel diameter, regulation of systemic arterial blood pressure by hormone, regulation of blood pressure…	1.92E-14	[ACTA2, ACTG2, ADRA2A, CACNA1C, CACNA1G, CACNA1H, CNN1, CRYAB, EDNRA, EDNRB, GJA5, HSPB6, KCNE4, KCNJ8, LMOD1, MYH11, MYL9, MYOCD, PLN, SLMAP, SMAD7, SSTR2, TPM2]
21	Extracellular matrix organization, ECM proteoglycans, NCAM1 interactions, Degradation of the extracellular matrix…	5.74E-14	[ADAMTS1, ADAMTS5, CD44, COL14A1, COL1A2, COL5A3, COL6A1, COL6A2, FGF2, ITGA7, ITGB3, MMP19, PTPRS, SDC2, SH3PXD2A, TNC]
13	Artery development, cardiac muscle tissue development, muscle tissue development mesenchyme morphogenesis…	1.17E-10	[APOE, GJA5, GLI3, MYOCD, NOTCH3, PDGFRB, PRRX1, SMAD7, TBX2]
13	Cardiac muscle cell action potential, NCAM signaling for neurite out - growth, muscle contraction…	2.29E-10	[CACNA1C, CACNA1G, GJA5, KCNE4, KCNJ8, SLMAP]
9	Smooth Muscle Contraction, muscle contraction, regulation of muscle system process…	1.27E-10	[ACTA2, ADRA2A, CNN1, EDNRA, EDNRB, MYH11, MYOCD, SSTR2]
4	Cell fate commitment, cartilage development, neuron fate commitment	5.22E-07	[FGF2, GLI2, GLI3, LMO4, NOTCH3, PRRX1, SATB2, SPRY2, TBX15, TBX18, TBX2, TBX3, TEAD3]
7	Syndecan interactions, ECM proteoglycans, Integrin cell surface interactions…	1.06E-06	[COL1A2, COL5A3, FGF2, ITGB3, SDC2, TNC]
6	Mesenchymal cell differentiation, mesenchymal cell development, atrioventricular valve development…	3.16E-06	[DLG5, EDNRA, EDNRB, HEYL, SEMA3C, SMAD7, SPRY2, TWIST1, VASN, ZEB2]

b Genes associated with Epithelial Tube Morphogenesis

c Genes associated with Blood Vessel Development

d Genes associated with Muscle Contraction

e Genes associated with Extracellular Matrix Organization

Principal Component 4: Myoepithelial Cells

Myoepithelial Cells	Pop4 Epithelial

(# pathways in group)	Representative Pathways	Adj. *p*-value	Associated Genes (from PC4 TOP 200) Myoepithelial / Epithelial
24	Extracellular matrix organization, Laminin interactions, hemidesmosome assembly, Cell-Cell communication, Degradation of the extracellular matrix, …	3.46E-16	[A2M, ADTRP, ANTXR1, CDH1, COL17A1, ELF3, ITGA3, ITGA6, ITGB4, ITGB6, ITGB8, LAMA3, LAMB3, LAMC2, LCP1, MYH11, NTN4, SERPINB5, SOX9, SPINT2, TNC, TPSAB1]
12	Cell junction organization, desmosome organization, apoptosis,…	1.00E-14	[ACTN1, AMIGO2, CDH1, CDH3, COL17A1, DSG2, DSP, DST, FZD1, ITGA3, ITGA6, ITGB4, KRT14, KRT5, LAMA3, LAMB3, LAMC2, MPP7, NECTIN1, NFASC, NTRK2, PARD3, PATJ, PERP, PLXNB1, PTPRF, PTPRT, SFRP1, SORBS1, TLR2, TNC]
9	Keratinization, skin development, formation of the cornified envelope, …	8.47E-14	[CDH3, DSC3, DSG2, DSG3, DSP, KRT14, KRT17, KRT5, KRT6B, KRT6C, KRT7, PERP, PRSS8, SFN, ST14]
6	Epidermal cell differentiation, prostate glandular acinus development, hair follicle development, …	6.03E-11	[CDH3, DSC3, DSG2, DSG3, DSP, ESRP1, IRF6, KRT14, KRT17, KRT5, KRT6B, KRT6C, KRT7, PALLD, PERP, PRSS8, SFN, ST14, TP63, VDR]
18	Morphogenesis of a branching epithelium, stem cell differentiation, tissue morphogenesis, prostate glandular acinus development, …	2.27E-10	[FGF1, FGFR2, LGR4, NTN4, SEMA3C, SFRP1, SOX10, SOX9, SPINT2, ST14, TACSTD2, TNC, TP63, VDR]
6	Cell-cell junction organization, cell-substrate junction assembly, cell junction organization, …	8.52E-10	[CDH1, CDH3, DSG2, DSP, MPP7, NECTIN1, NFASC, PARD3, PATJ, PERP]
7	Epithelial tube formation, branching involved in labyrinthine layer morphogenesis, epithelial cell development,…	1.32E-07	[FGFR2, FZD1, IRX1, SFRP1, SHROOM3, SOX9, SPINT2, ST14]
10	Actin filament bundle assembly,…	2.07E-06	[ACTN1, CGNL1, EZR, LCP1, PIK3R1, SFRP1, SORBS1, SYNPO2, TACSTD2]
2	Intermediate filament cytoskeleton organization, intermediate filament organization	1.63E-05	[DSP, DST, EPPK1, KRT14, KRT17, KRT6C, SYNM]
1	Regulation of Wnt signaling pathway	3.16E-05	[CDH3, DIXDC1, DKK3, EGFR, FGFR2, FZD1, ITGA3, LGR4, PLEKHA4, RNF43, SFRP1, SLC25A23, SOX10, SOX9, TLR2, WIF1]
2	Smooth muscle contraction, muscle contraction	4.59E-05	[ACTA2, ACTG2, LMOD1, MYH11, MYLK, SORBS1, TPM2]

b Genes associated with Extracellular Matrix Organization

c Genes associated with Cell Junction Organization

d Genes associated with Keratinization

e Genes associated with Morphogenesis of a Branching Epithelium

Legend: 1. Luminal 1, 2. Luminal 2, 3. Myoepithelial, 4. Epithelial, 5. vSMC, 6. Pericytes, 7. Lymph.Endo, 8. Fibroblasts, 9. Endothelial, 10. Leukocytes, 12. Adipocytes

Figure 15

Figure 15-table 1

Gene Family	DE Genes	Genes in Family
Adenylate cyclases	9	10
Aldehyde dehydrogenases	16	19
Collagens	38	45
Death inducing signaling complex	6	6
DOCK family Rho GEFs	11	11
Ephrins	6	8
Erb-b2 receptor tyrosine kinases	4	4
Fibulins	7	7
Glutathione S-transferases	15	23
GTPases, IMAP	7	7
Histocompatibility complex	14	14
Histones	37	89
Integrin alpha subunits	17	18
Integrin beta subunits	9	9
Interleukins	18	37
Interleukin receptors	34	42
Keratins, Type I	7	26
Keratins, Type II	10	26
Kruppel like factors	14	17
Laminin subunits	12	12
Lysophosphatidic acid receptors	6	6
MAP kinase phosphatases	10	11
Matrix metallopeptidases	14	23
Mucins	10	14
Plexins	9	9
Ras association domain family	10	10
Rho GTPase activating proteins	46	48
Suppressors of cytokine signaling	8	8
Tetraspanins	22	33
Toll like receptors	9	10
V-type ATPases	10	23

Fig.15—figure supplement 1

HGNC Gene Family Size

Median Size   6
Mean Size   14.2
sdev   42.2

Figure 15—figure supplement 2

Figure 16

Claudins
HGNC: CLDN

Figure 16—figure supplement 1

CD49f   Claudin-1   DAPI

Figure 17

Figure 18

Receptor Tyrosine Kinases
HGNC: RTKs

Color Key and Histogram

1. Luminal 1
2. Luminal 2
3. Myoepithelial
4. Epithelial
5. vSMCs
6. Pericytes
7. Lymphatic Endothelial
8. Fibroblasts
9. Vascular Endothelial
10. Leukocytes
12. Adipocytes
13. Cells (unsorted)

b. platelet-derived growth factor — PDGFRA
c. platelet-derived growth factor — PDGFRB
d. fibroblast growth factor receptor 1 — FGFR1
e. AXL receptor tyrosine kinase — AXL
f. insulin receptor — INSR
g. TEK tyrosine kinase, endothelial — TEK
h. tyrosine kinase with immunoglobulin- — TIE1
i. fms-related tyrosine kinase 1 — FLT1
j. epidermal growth factor receptor — EGFR
k. erb-b2 receptor tyrosine kinase 2 — ERBB2
l. erb-b2 receptor tyrosine kinase 3 — ERBB3
m. erb-b2 receptor tyrosine kinase 4 — ERBB4

Figure 19

Figure 20

Figure 21

Figure 22

Biological Themes
Select Leading Edge Genes
ER^Pos vs. ER^Neg Luminal Epithelial Cells

Figure 23

Enriched Gene Sets & Biological Themes
Luminal Epithelial Cells:
ER^Pos Luminal (Pop1, ◯) vs. ER^Neg Luminal (Pop2, ●)

Ion Channel Transport
(WNK4)
(ATP13A5)

Keratinization
(KRT83)

Rho-GTPase
Signaling
(ARHGAP40)

Semaphorin
Interactions

MAPK Signaling

Muscle Contraction
(MYBPC1)

FOXO Mediated
Transcription

NCAM
Signaling

Sumoylation
(TFAP2b)

Hemostasis
(KIF12, ATP1B1, PSG9, CEACAM6,
ALB, ORM1, ORM2)

IGN-gamma
Signaling

MTOR Signaling

Antimicrobial Peptides
(DEFB1, EPPIN-WFDC6
BPIFB1, LTF, S100A7,
PI3, LCN2)

Estrogen Signaling
(FOXA1, MYB, TFF1, TFF3,
ERBB4, Amphiregulin, ESR1)

Chemokine
Signaling
(CXCL16, CCL28, CCL4)

(PGLYRP2)

FGFR Signaling

MHC Class I

MHC Class II

Integration of
Metabolism
(PRKAR2B, ABCC8)

Autophagy

Translation
(RPL genes)

Other Interleukin
Signaling
(CSF3R)

Glycolysis
(6-Phosphofructo-2-Kinase/
Fructose-2,6-Biphosphatase 4)

Fatty Acid
Metabolism
(Cytochrome p450 4F22)

(EEF1A2)

Interleukin Signaling
(IL1R2)

Dectin-1 Signaling

IL10   IL2
IL1
IL12
IL4/13

Non-canonical
NFKB Signaling

TCR Signaling
(LBP, S100A9, S100A1, BPi)

Metabolism of Proteins

VEGF Signaling
(VAV3)

TLR Cascades
(Toll-like Receptor 2,6)

FCERI Activation

Vesicle Mediated
Signaling

Tight Junctions
(Claudin 8)

Respiratory Electron Transport
(Isocitrate Dehydrogenase 2)

Phospholipid Metabolism
(SLC44A4, PLEKHA6, INPP5J, SLC44A5)

ERBB4 Signaling
(ERBB4, EREG, PGR)

Sphingolipids

PTK6 Signaling

(NRG1, NRG2, GFAP)

Phospholipids

HGF/MET Signaling
Focal Adhesions

PI3K AKT Signaling
(Amphiregulin, ERBB4, ERBB2
ERBB3, ESR1)

O-linked
Glycosylation
(MUC15)

BCR Signaling

Neurotrophic Tyrosine
Receptor Kinase
Signaling

(GCNT3)

Non-integrin
ECM Interactions

ECM Proteoglycans
(APP, TGFB1)

Collagen
Formation
(LOXL4, COL22A1,
MMP7)

Collagen / ECM
Degradation
(ADAM9, Calpain-8)

Integrin Cell
Surface Interactions

Elastic Fiber
Formation

Laminin
Interactions
(LAMA5)

◯ 212 Gene Sets, FDR ≤ 0.1

● 113 Gene Sets, FDR ≤ 0.1

Gene Set Size
◯ 180
◯ 100
○ 30

—— 50% shared genes

Figure 23—Supplementary Figure 1

Biological Themes
Select Leading Edge Genes
ER^{Pos} vs. ER^{Neg} Luminal Epithelial Cells

Figure 23—figure supplement 2

Biological Themes contd.
Select Leading Edge Genes
ER^Pos vs. ER^Neg Luminal Epithelial Cells

1. Luminal 1
2. Luminal 2
3. Myoepithelial
4. Epithelial
5. vSMCs
6. Pericytes
7. Lymphatic Endothelial
8. Fibroblasts
9. Vascular Endothelial
10. Leukocytes
12. Adipocytes

IL1 Signaling

a

SAA1
RPS27A
UBA52
S100B
IL1R2
IL1A
NOD2
PELI1

IL2 Signaling

b

PIK3R1
IL2RG
CSF2RB
CSF2
JAK2
CSF2RA
JAK3
IL15RA
SOS2
INPP5D
IL5RA
STAT4
IL3RA
SOS1
IL21R
PTK2B

IL12 Signaling

c

RPLP0
IL27RA
TCP1
HNRNPDL
GSTA2
SERPINB2
JAK2
MSN
IL12A
PPIA
EBI3
IL12B

Toll Like Receptor

d

SAA1
RPS27A
S100A8
TANK
MEF2C
DUSP7
S100A1
S100A9
BIRC3
UBA52
S100B
BPI
MAP2K3
PLCG2
LBP
NOD2
ITGB2
PELI1
CTSV
DUSP6
RIPK2
CTSS
CD180
MAPK14
TLR7
TLR2
PELI2
LY96
TLR6
BIRC2
NFKB1
TLR4

Defensins & Antimicrobial Peptides

e

LTF
S100A8
PI3
CLU
S100A9
RNASE6
BPI
S100A7
TLR2
LCN2
S100A7A
DEFB1
RNASE7
PGLYRP4
CCR6
TLR1
EPPIN- WFD
6

Fig.23—figure supplement 3

Figure 23—figure supplement 4

Milk Proteins

Figure 24

Myoepithelial vs. ER^Neg Luminal Cells

2,239 Differentially Expressed Genes

Legend:
1. Luminal 1
2. Luminal 2
3. Myoepithelial
4. Epithelial
5. vSMCs
6. Pericytes
7. Lymphatic Endothelial
8. Fibroblasts
9. Vascular Endothelial
10. Leukocytes
12. Adipocytes
13. Cells (unsorted)

Figure 25

Enriched Gene Sets
Epithelial Cells:
ER^Neg Luminal (Pop2, ●) vs. Myoepithelial (Pop3, ●)

TLR Cascades
(S100A8, Toll-like receptor 1,2)

NCAM1 Interactions
(Calcium channel subunit 4, ST8SIA2)

Notch Signaling
(NOTCH2, NOTCH3)
(JAG1, JAG2)

O-linked glycosylation
(MUC5B, MUC16, MUC15, MUC7, MUC1, B3GNT3)

Notch4

Integrin Cell
Surface Interactions

RUNX1 Regulation
(TP73)

ECM Proteoglycans

VEGF
Signaling

Respiratory Electron Transport

Fatty Acid
Metabolism
(Glutathione Peroxidase -2)

Glucose
Metabolism

Glycolysis

FGFR Signaling
(FGFR2, FGF1)

L1CAM
Interactions

(CD24)

Negative regulation of
PI3K/Akt Signaling
(IL1RAP, FGF1, FGFR2, Neuregulin-2)

Negative regulation of
ERBB2 Signaling
(ERBB2/HER2, ERBB3, EGF)

Myogenesis
(MAP2K6)

Interactions at the
Vascular Wall

Interferon
Signaling
(TRIM29, GBP3)

Muscle
Contraction
(Voltage-gated sodium channel)

Metabolism of
Steroids
(HSD11B1, HSD17B2, CYP24A1)

Glycosaminoglycan
Metabolism
(HAS3, Thrombospondin 7B, CHST3)

Cell-cell
Communication
(Keratin 5, 14, Cadherin 3)

Tight Junction
Organization
(Claudin 1,3,4,7,8,10)

P53 Regulation
(BCL2L14, TP63)

Interleukin-10
Signaling
(Interleukin-1 Receptor-2)

MET Signaling
(Tensin-4)

Ion Channel
Transport
(Chloride Channel Accessory -2 ,
Anoctamin-1)

Non-integrin
Membrane Interactions
(COL11A1, COL4A6, COL17A1)

Collagen Regulation

(Laminin alpha-3, Integrin alpha-2,
COL17A1, COL8A2, COL2A1,
COL4A5, Dystonin)

Defensins
(Lactotransferrin, DEFB1, CLU,
LCN2, S100A7, S100A9,
S100A7A, BPI, EPPIN-WFDC6)

Keratinization
(PI3, Caspase-14, ST14, KLK5,
Keratin 8, 18, 19, 23, 80, 81, 83)

ATP13A5, ATP6V0A4

ECM Degradation

Activation of MMPs

(MMP7)

(RNASE7)

(Keratin 5, 6ABC, 14, 17)

● 110 Gene Sets, FDR ≤ 0.1

● 316 Gene Sets, FDR ≤ 0.1

Gene Set Size
○ 180
○ 100
○ 30

— 50% shared genes

Figure 25—figure supplement 1

Biological Themes
Select Leading Edge Genes
ER^Neg Luminal Cells vs Myoepithelial Cells

Fig.25—figure supplement 2

Figure 26

ER^Neg Luminal Cells vs. Vascular Endothelial Cells

4,167 Differentially Expressed Genes

Figure 27

Enriched Gene Sets & Biological Themes
Luminal Epithelial Cells:
ER^Pos Luminal (Pop2, ⬤) vs. ER^Neg Luminal (Pop9, ⬤)

Biological Themes
Select Leading Edge Genes
ER^Neg Luminal Epithelial Cells vs. Vascular Endothelial Cells

Figure 28

Lymphatic vs. Vascular Endothelial Cells

3,739 Differentially Expressed Genes

Figure 29

Biological Themes
Select Leading Edge Genes
Lymphatic vs. Vascular Endothelial Cells

Figure 30

Fibroblasts vs. Adipocytes
1,171 Differentially Expressed Genes

Figure 31

Biological Themes
Select Leading Edge Genes
Fibroblasts vs. Adipocytes

Figure 32

Fibroblasts vs. Pericytes

2,932 Differentially Expressed Genes

Fibroblasts ← → Pericytes

Log₂ fold change

1. Luminal 1 2. Luminal 2 3. Myoepithelial 4. Epithelial 5. vSMCs 6. Pericytes 7. Lymphatic Endothelial 8. Fibroblasts 9. Vascular Endothelial 10. Leukocytes 12. Adipocytes 13. Cells (unsorted)

Figure 33

Enriched Gene Sets & Biological Themes
Mesenchymal Cells:
Pericytes (Pop6, ●) vs. Fibroblasts (Pop8, ○)

Biological Themes
Select Leading Edge Genes
Pericytes vs. Fibroblasts