A community genomics approach to natural hybridization

Zachery D. Zbinden[1*], Marlis R. Douglas[1], Tyler K. Chafin[2], Michael E. Douglas[1]

[1]Department of Biological Sciences, University of Arkansas, Fayetteville, AR, USA

[2]Biomathematics and Statistics Scotland, Edinburgh, Scotland, UK

[*]Corresponding Author: zdzbinde@uark.edu

Abstract

Hybridization is a complicated, oft-misunderstood process. Once deemed unnatural and uncommon, hybridization is now recognized as ubiquitous among species. But hybridization rates within and among communities are poorly understood despite the relevance to ecology, evolution, and conservation. To clarify, we examined hybridization across 75 freshwater fish communities within the Ozarks of the North American Interior Highlands (USA) by SNP genotyping 33 species (N=2,865 individuals; ddRAD). We found evidence of hybridization (70 putative hybrids; 2.4% of individuals) among 18 species-pairs involving 73% (24/33) of study species, with the majority being concentrated within one family (Leuciscidae/minnows; 15 species; 66 individuals). Interspecific genetic exchange— or introgression— was evident from 24 backcrossed individuals (10/18 species-pairs). Hybrids occurred within 42 of 75 communities (56%). Four selected environmental variables (species richness, protected area extent, precipitation [May and annually]) exhibited 73–78% accuracy in predicting hybrid occurrence via random forest classification. Our community-level assessment identified hybridization as spatially widespread and environmentally dependent (albeit predominantly within one large, diverse family). Our approach provides a more holistic survey of natural hybridization by testing a wide range of species-pairs, thus contrasting with more conventional evaluations.

Keywords:

environmental predictors, flooding, hybridity, introgression, Leuciscidae, minnows

1. Introduction

Hybridization was once considered "exceedingly rare" [1], but it is now acknowledged as relatively common, primarily because of better detection using modern DNA sequencing [2,3]. A concurrent recognition is the significant role of hybridization and introgression in ecology and evolution [4–6], which diverges sharply from the historical perspective rooted in the biological species concept [7]. But while a growing catalog of species known to hybridize has helped to draw attention to hybridization (per-species documentation), the prevalence of hybrids within and among communities is poorly understood (per-individual documentation) [2].

The more frequently we encounter hybridization, the more evidence for its influential role in nature [2,8]. Hybridization and introgression are often considered maladaptive threats to biodiversity [6,9]. However, it is becoming more appreciated that the genetic novelty injected into a lineage through introgressive hybridization can be adaptive [8] and even provide evolutionary rescue [10]. Moreover, hybridization is expected to increase in lockstep with global environmental change [11,12]. Therefore, baseline estimates are needed to precisely gauge increases in hybridization and detect where it impacts ecosystems [10,13]. Attempts to understand the geographic patterns and processes of hybridization have not often considered an array of taxa. However, doing so will help identify where it occurs more generally and what environmental factors might drive it. These insights are essential because these places/environments may disproportionately impact biodiversity and ecosystems [14].

Broad comparative examinations often employ meta-analyses of numerous single-species-pair studies to quantify hybridization rates [15] or assess biogeographic relationships [14,16,17]. Several generalizations have emerged. Hybridization tends to be unevenly distributed across taxonomic groups and occurs more in plants than animals [2]. It has been documented more so in fish than other vertebrates [18], likely due to external fertilization within the aquatic environment [19]. Among fish, evidence of hybridization is more prevalent in freshwaters [1], especially between carps and minnows (Cypriniformes) [19], owing to their high diversity, sympatry, and breeding behaviors [20].

The less time since two species have diverged, the less time for pre- and post-zygotic barriers to

46   evolve [20–22]. Similarly, the geographic overlap between two closely-related species also controls the

pressure to evolve reproductive barriers [23], with most hybridizing lineages first evolving in allopatry

48   [24]. Pre-zygotic barriers are often keyed (directly or indirectly) with the environment [25,26], weakening

concurrently with environmental fluctuations, thus facilitating hybridization. For example, many fish rely

50   on coloration to identify conspecific mates, and any environmental alterations impacting visual acuity

(e.g., turbidity, siltation) can promote hybridization [27]. Much work has focused on human impacts due

52   to environmental homogenization, sensory cue changes, and creating novel environments where hybrids

may have advantages [28].

54   We quantified hybridization across a North American freshwater fish metacommunity by

genotyping genome-wide single nucleotide polymorphisms (SNPs) that allowed us to detect hybrids

56   among the individuals we sampled. We addressed the following questions: (i) How many hybrids occur

within and among localities, and what is their frequency within species/families? (ii) Is genetic exchange

58   among species (i.e., introgression) occurring as evinced by the presence of backcrossed individuals (e.g.,

F_1 x Parental)? (iii) Is the incidence of hybrid individuals within communities predicated on

60   environmental factors which allow the prediction of hybrid occurrence?

62   **2. Methods**

(a) Data generation and processing

64   Our sampling area spanned the White River Basin (71,911 km^2; Ozark Plateau, North American Interior

Highlands). It represents an ideal study location because: (i) It is an unglaciated refugium with elevated

66   fish diversity and limited anthropogenic impacts [29–31]; and (ii) Central to and representative of the

Mississippi system, it is an excellent region from which generalizable patterns can be interpreted [32].

68   Sampling procedures were approved by the University of Arkansas Institutional Animal Care and Use

Committee (IACUC #17077) and appropriate permitting agencies (Supplement S1). Fishes were seined

70   (June 2017 to September 2018) and euthanized via tricaine methanesulfonate (MS-222) and 95% ethanol.

Species diagnoses occurred in the laboratory.

72   Genomic DNA was isolated from fin clips (Qiagen Fast kits; Qiagen Inc.) and quantified by

fluorometry (Qubit; Thermo-Fisher Scientific). Individuals were SNP genotyped using double-digest

74   restriction site-associated DNA sequencing (ddRAD), using modified procedures [33,34], and sequenced

as pooled batches (N=144/1x100 lane) on the Illumina HiSeq 4000 (Supplement S1).

76   Raw Illumina reads were demultiplexed (IPYRAD; [35]), and family-level phylogenies were

generated to verify individual species IDs via an alignment and assembly-free method (PHYLORAD; [36];

78   Supplement S1). Individuals (N=3,042) were grouped by family (N=6) and processed de novo in IPYRAD to

generate separate family-level assemblies. Adapters/primers were removed, and reads with >5 low-quality

80   bases (Phred<20) were discarded. Clusters were assembled using an 85% identity threshold, with loci

subsequently removed via conditional criteria to ensure high-quality data (Supplement S1). Biallelic

82   SNPs were further filtered and visualized using RADIATOR [37] (Supplement S1).

84   **(b) Hybrid detection**

Our initial objective was to detect putative hybrids and determine hybridization rates. We focused only

86   within families for two reasons: No evidence implicates hybridization among these divergent families

within six different orders; Also, a high number of species in RADseq alignments can reduce the power to

88   detect hybrids by diminishing recovery of homologous loci (via allelic dropout from accumulating

mutations in restriction enzyme cut-sites) [38]. All analyses were conducted using R 4.1.3 [39]. Individual

90   genetic variation was first visualized using separate principal components analyses (PCAs) on each

family-level SNP alignment (ADE4; [40]).

92   Admixture analysis employing sparse non-negative matrix factorization (sNMF) was used to

estimate ancestry coefficients for individuals within families (25 repetitions/K value (1–25) with

94   regularization parameter (α)=100 (LEA; [41]). The best K from each run (via cross-validation) was used to

impute missing data (*impute* function, method='mode' in LEA), with sNMF repeated using imputed data (as above). An individual was flagged as a putative hybrid if the assignment probability for the majority ancestry cluster was <0.9.

Results were contrasted with a maximum-likelihood clustering approach considering hybrid categories per expected allele frequencies (SNAPCLUST; [42]). Three models were tested: (i) F_1 only; (ii) F_1+ first-generation backcross; and (iii) F_1+first- and second-generation backcrosses, with AIC used to determine the best fit [42]. Pairwise analyses were among species within families.

We assessed the power of sNMF and SNAPCLUST to detect hybrids. We first removed putative hybrids following the above analyses. Then, we sampled alleles from the remaining pure individuals for each within-family species-pair to simulate hybridization between parentals (P_1 & P_2). We simulated 10 each of F_1, F_2, and backcross (5 P_1 x F_1 & 5 P_2 x F_1) (function *hybridize*; ADEGENET; [43]). SNPs for simulated hybrids were combined with parentals before running sNMF and SNAPCLUST (as above). We then quantified the proportion of hybrids detected by each analysis.

(c) Hybrid classification

Hybrids detected with sNMF and SNAPCLUST were classified by assignment as either F_1, F_2, or backcross (NEWHYBRIDS; [44]). Backcrosses underscore hybrid viability and gene exchange among species (i.e., introgression). We implemented HYBRIDDETECTIVE [45] to confirm sufficient statistical power for classification (F_1, F_2, or BXs). All NEWHYBRIDS analyses were conducted pairwise among species from which hybrids were detected. We first simulated known class hybrids by randomly sampling two alleles/locus from appropriate parental pools, with convergence assessed via three replicates. We then determined an optimal posterior probability threshold (=0.70) from which hybrid classes could be reliably assigned. The final MCMC was 1,000,000 iterations (250,000 burn-in). Assignments were made using a reduced panel of 200 SNPs exhibiting the greatest among-species differentiation (Weir and Cockerham's

F_{ST} among target species-pair) and lowest linkage disequilibrium (r^2<0.2) via *getTopLoc*, HYBRIDDETECTIVE

120 [45].

122 **(d) Correlates of hybridization**

We tested whether hybridizing/non-hybridizing species-pairs differed significantly with respect to genetic

124 differentiation and basin-wide co-occurrence. Weir and Cockerham's F_{ST} (DIVERSITY; [46]) and the number

of co-occurrences across communities (COOCCUR; [47]) were calculated among *N*=137 species-pairs within

126 families. Differences among hybridizing/non-hybridizing pairs were assessed using the Wilcoxon rank

sum test.

128   We evaluated the influence of environmental factors on the incidence of hybrids within

communities to determine their predictability. Each community (*N*=75) was classified as with/without

130 hybrids, with potential predictor variables being: Species richness at each site; Drainage identity (8-digit

USGS hydrologic unit); and high-resolution hydro-environmental descriptors (*N*=281, RiverATLAS;

132 [48]). These encompassed factors broadly related to hydrology, physiography, climate, land cover,

geology, and anthropogenic impact. Associations between hybrid incidence and predictive variables were

134 assessed using random forest classification (i.e., supervised machine-learning employing linear/non-linear

relationships among mixed data types [49]). Bootstrapped decision trees (via predictor variables) are

136 aggregated, with each trained on a random 2/3 of samples (="in-bag"), with validation via the remaining

1/3 (="out-of-bag").

138   We implemented the random forest quantile-classifier approach [50] in RANDOMFORESTSRC [51]

using 100,000 bootstrapped decision trees (=ntree), each of which randomly sampled 1/3 of total

140 predictors (=mtry). Variable Importance (VIMP; Increase in prediction error when a variable is

randomized) was quantified via permutation for each predictive variable. VIMPs significantly >0 were

142 assessed by subsampling 1,000 trees. Random forest variable selection was executed using minimal depth

and a conservative threshold set to "high," employing only that subset of predictive variables with

144   significant VIMP [52]. A final random forest model was built from this selected subset of predictors.

Accuracy was evaluated using G-mean (0–1; the geometric mean of the true negative and positive rates),

146   which replaces the misclassification rate in imbalanced data settings [50] and the normalized Brier score

(0–1; mean square difference between true classes and predicted probabilities).

3. Results

150   Fish collections across 75 communities/sites yielded 72 species (N=3,605 individuals), averaging 10.8

species/site—typical for streams across the Mississippi Basin [53]. We genotyped 33 species to represent

152   the fish metacommunity [54] (i.e., 84% of total individuals collected, averaging 9.44 spp./site). Of N=137

intra-familial species-pairs examined, 13 (9.5%) did not co-occur at our sites. Four do have non-

154   overlapping, parapatric distributions, each with one occurring in the upper White and the other in the

Black River: one ictalurid (*Noturus albater/ maydeni*); one leuciscid (*Luxilus pilsbryi/ zonatus*); and two

156   percids (*Etheostoma juliae/ uniporum*; *E. spectabile/ uniporum*). Despite range overlap, nine other pairs

did not co-occur (six leuciscids and three percids), which may be due to chance/sampling, given that pairs

158   included at least one uncommon species (≤10 occurrences).

We examined N=2,865 individuals across six families post-filtering using SNP genotypes (Table

160   1; mean missing data=21%; mean coverage=56x). SNPs varied by family and were inversely related to

the number of species (Table 1). Power analyses verified the robustness of panels for detecting/classifying

162   hybrids based on genotype frequencies (Supplement S2–S19). Among N=30 simulated hybrids for each

unique species-pair assessed (N=137), sNMF and SNAPCLUST detected 93% and 100%, respectively.

164   Simulated hybrids among pairs targeted for classification via HYBRIDDETECTIVE and NEWHYBRIDS showed 90-

100% assignment accuracy across genotype frequency classes and species-pairs for posterior probability

166   thresholds ≥ 0.70.

(a) Hybrid detection

The number of species-pairs examined varied considerably among the six families/orders, with just one pair tested for two families versus 105 pairs within Leuciscidae (Table 2). Hybrids (N=70; 2.4%) were detected within four families (Table 2), with hybrid proportions ranging from 0–4.4% of individuals. All but four hybrids were minnows (Leuciscidae), with exceptions being: (i) *Micropterus dolomieu* x *salmoides* (Centrarchidae); (ii) *Etheostoma juliae* x *zonale* (Percidae); (iii) *Etheostoma spectabile* x *caeruleum* (Percidae); and (iv) a putative multi-specific hybrid: *Noturus maydeni* x *albater* x *exilis* (Ictaluridae). Thus, we recognized N=18 hybridizing species-pairs and N=8 species-triplets, i.e., multi-species hybrids. We did not detect hybrids within Cottidae or Fundulidae.

Within Leuciscidae, we identified 66 hybrids and 15 hybridizing species-pairs (Supplement S20). The most remarkable (N=29 individuals; 41%) involved *Campostoma anomalum* x *oligolepis*. All minnow species herein (N=15) appear to hybridize with at least one other species in the study basin. The number of hybrids per species was significantly related to the number of community occurrences (R^2=0.16, F=6.1, p=0.02).

PCA, sNMF, and SNAPCLUST were largely congruent in detecting hybrids (Figures 1-2). Both sNMF and SNAPCLUST detected the same 36 hybrids, each separately identifying additional (26 and 8, respectively). SNAPCLUST showed excellent power in detecting hybrids based on allele frequencies (100%). However, it comes with a cost: the same individual was often suggested as a hybrid in multiple pairwise tests, regardless of the second parental species. Fortunately, we could adjudicate these using PCA and sNMF. A list of hybrid individuals and inferences for each are available in Supplementary Material (S21).

(b) Hybrid classification and introgression

We also found evidence of introgression based on the occurrence of backcrossed hybrid individuals. NEWHYBRIDS identified 39% of hybrids as backcrossed (24/62, excluding eight multi-specific hybrids); this corresponds to 55% (10/18) of all putatively hybridizing species-pairs showing evidence of introgressive

hybridization (Table 3). Approximately 24% of hybrids (15/62) were seemingly F_1 or F_2. Finally, NEWHYBRIDS designated 37% (23/62) of putative hybrids as pure parentals, although possibly introgressed individuals poorly classified, i.e., late-generation hybrids [42].

(c) Correlates of hybridization

Genetic differentiation among species (Table 4; Supplement S22) was significantly lower for hybridizing versus non-hybridizing pairs (mean F_{ST} = 0.86 vs. mean F_{ST} = 0.93; W=1521; p=0.002). The number of co-occurrences between hybridizing (mean=8.4) and non-hybridizing pairs (mean=7.1) did not differ significantly (W=960; p=0.76).

At least one hybrid individual occurred within 42/75 communities (56%). We identified 12 environmental variables with a significant ability to reduce predictive error (i.e., VIMP) based on random permutation of predictors in the random forest (Supplement S23-S24). From these, just four were used to maximize the predictive capacity of the final model based on minimal depth selection (Figure 3), including species richness (VIMP=0.09), protected area extent (pac_pc_use; VIMP=0.04), mean precipitation in May (pre_mm_c05; VIMP=0.03), and mean annual precipitation (pre_mm_cyr; VIMP=0.03). The accuracy of the model was satisfactory: G-mean=0.73 and normalized Brier score=0.78. A similar misclassification error occurred for communities with and without hybrids (0.26 vs. 0.27, respectively).

Neither species richness nor protect area extent were strongly correlated with other predictor variables (r >0.70; Pearson correlation). The correlation coefficient between mean May and annual precipitation was r=0.61. Both precipitation variables were strongly and positively correlated with other climatic variables related to temperature, evapotranspiration, and soil water content. Annual precipitation was also strongly but negatively correlated with river network position (distance from network outlet; DIST_DN_KM). Although not selected for the final model, eight additional variables were significantly associated with hybridization based on VIMP, including indices of human impact (N=2), road density (N=1), annual snow cover (N=1), and vegetative cover (N=4) (Supplement S23-24).

4. Discussion

Hybridization and introgression can impact fitness, facilitate gene exchange among species, or generate new lineages [5,6,55–57]. While hybridization has been widely documented at the per-species level, its occurrence within communities is expected to be rare; otherwise, species boundaries would seriously deteriorate [2]. Despite these expectations, hybrid prevalence at the community level is largely unknown, given that most studies to date focus on single species-pairs (or but a few closely related pairs) [55].

This study quantified hybridization at the community level without *a priori* assumptions of putative hybridization, thus serving as an appropriate broad-scale model without species-specific biases. It provided a more holistic survey of hybridization across the riverscape and thus stood in contrast to more conventional studies based on single species-pair evaluations.

(a) Frequency of natural hybridization

The relevance of hybridization in ecology and evolution is underscored in our study by the detection of hybrids within four (of six) families (67%), involving 24 (of 33) species (73%). We documented hybridization among 18 unique species-pairs, 10 of which have been reported previously [20,58–62]. Hybridization across our study region was overwhelmingly within leuciscids, a family oft-dominating within Ozark stream communities [63]. The unbalanced number of representatives from each family reflects that and should not necessarily be interpreted as differences in the propensity to hybridize among the families more generally. Surprisingly, few hybrids were detected within Centrarchidae and Percidae, despite an extensively-documented presence [19,64,65].

Hybridization was encountered more frequently than anticipated, whether viewed per-individual (2.4%) or per-species (73%). Previous estimates in plants and animals are lower than found herein (0.002–0.06%) [15,66,67]. If rates among our study species were comparable to the previous (e.g., 0.1%), we would have expected but 2–3 hybrids. Earlier estimates relied on morphological identification, a methodology less sensitive in detecting later-generation hybrids [19,62]. Additionally, rates herein also

246   reflect both our breadth of individuals evaluated, as well as potential comparisons so tested. While our

study is but a subsample of individuals from the region and lacks representatives from every extant

248   species, we nevertheless attempted to reduce unsampled lineages that could bias inferences and allow

undetected or misclassified hybrids [68,69].

250   Hybridization is seemingly greater in fishes than in other vertebrates due to their external

fertilization within aqueous environs [1,19,70]. Similar per-individual rates (0–4%) were identified in

252   marine fishes [71], with even higher rates (22.5%) among invasive Mississippi River carp [72]. Published

per-species estimates varied from 1–10% across animals and ~25% for plants [2,73]. Although

254   differences in the former are apparent from literature and museum records, one could ask if it reflects an

actual biological signal or variance in diversity and/or research effort. For example, once adjustments

256   were made for diversity and research intensity, rates were mostly homogenous among taxonomic groups

compiled in a meta-analysis, but with the caveat that rates for fishes were still demonstrably higher than

258   expected [18].

Our per-species rate was highest within our most specious group (Leuciscidae; Table 2). This

260   parallels previous meta-analyses identifying disproportionately high hybridization rates within minnows

[1,19,20]. Interestingly, leuciscid breeding behaviors—especially nest building and association—

262   significantly predicted hybridization rates across the clade [20]. Many minnows broadcast gametes widely

or affix them onto substrates, often shared among species [20,74], with as many as six simultaneously

264   employing the same gravel substrate at once [1]. In addition, minnows comprise the most diverse and

widely distributed North American stream fish family and numerically dominate stream communities

266   [75]. They thus encounter broad environmental heterogeneity, demonstrate a high degree of sympatry,

and exhibit uneven species abundances, which promote hybridization [19].

(b) Introgression

270   The evidence of backcrossing between hybrids and parentals suggests that hybrids can be viable and

fertile, thus facilitating introgression [67]. This genetic exchange can provide a source of novelty for

272 evolutionary forces to act upon [73], and the prevalence found here provides some indication as to how probable an extraneous genetic contribution can be. We identified 24 individuals seemingly resulting

274 from parental/hybrid backcrosses spanning ten species-pairs and two families (Leuciscidae, Percidae). It is intriguing that the most diverse group of North American freshwater fish—leuciscids—has an

276 established reputation for frequent hybridization and is also a prolific contributor to gene exchange among species in our study. Introgressive hybridization has played a crucial role in cyprinid evolutionary history

278 [67], and the evidence gathered here suggests it may continue to do so.

Questions remain regarding whether the introgression seen in our region is evolutionarily

280 adaptive, maladaptive, or neutral. This may ultimately hinge upon differences among species and environments [76]. Answers would facilitate our capacity to predict and mitigate adverse outcomes of

282 hybridization [77]. Climate-mediated environmental shifts will predictably exacerbate hybridization, especially for fishes whose life histories are sensitive to spawning temperatures and streamflow [12]. Yet

284 we recognize that hybridization is not necessarily negative. For example, adaptive introgression between generalist and specialist rainbow fishes has seemingly diminished climate change vulnerability in

286 admixed individuals compared to pure populations [10]. It is thus viewed as a valuable component of "evolutionary rescue" and an underappreciated conservation tool [10].

(c) Correlates of hybridization

290 Our data concur with the premise that hybridization is strongly influenced by both divergences among species and also environmental factors. We focus our discussion on the latter, per the novelty of our data

292 and the previously established support for the former [20–22]. In synopsis, communities were more likely to harbor hybrids when: (i) Greater species richness existed; (ii) Protected area within the catchment was

294 limited; and (iii) Habitats were prone to more precipitation.

While hybridizing and non-hybridizing species-pairs did not differ significantly in their co-

296 occurrence, we did note that communities where more species occur together (greater richness) also possessed more hybrids (Figure 3). Although seemingly intuitive, the opposite is suggested at larger

298 spatial and temporal scales where species-richness diminishes with greater niche availability at

biogeographic borders [16,23]. Empirical results are few, and conclusions varied: For example, the

300 number of plant hybrids across US counties was significantly related to species richness [17]. However,

hybridization and species richness were not significantly related among coral reefs [14,16].

302 The extent of protected areas within the total watershed upstream of the reach was negatively

associated with basin-level hybridization (Figure 3). Similarly, anthropogenic impacts were positively

304 associated with hybridization (Supplement S24). Environmental perturbations in general, and specifically

those anthropogenic, loom large in the hybridization literature [19,28]. Stable, more pristine environments

306 are expected to harbor fewer hybrids than those perturbed due mainly to a breakdown of reproductive

isolating mechanisms facilitated by translocations, habitat modifications, and ongoing climate change

308 [28,78].

Greater mean precipitation within the reach catchment (annually and in May) was also associated

310 with elevated hybridization (Figure 3). More precipitation is associated with warmer and more

downstream communities in our system (i.e., more flow). Elevated levels of hybridization in these

312 communities are potentially driven by flooding magnitude, particularly given the combination of higher

precipitation and lower network position. Horton stream order and catchment size promote the magnitude

314 of flooding [79], as compounded by the precipitation gradient of our study system [80]. Additionally, the

Hydrologic Disturbance Index (a compendium of several anthropogenic impacts) is more significant in

316 streams with larger drainage areas and lower gradients (i.e., further downstream), which in turn has been

shown to promote variance in fish community composition [31].

318 Flooding can affect fish spawning in several ways: promotion of spawning activity [80];

concentration of fishes within refugia [53,81]; disturbance of spawning habitats and nests [82];

320 displacement of oviposited eggs [53]; and elevation of discharge/turbidity, which weakens those sensory

cues (e.g., visual, olfactory, environmental) that sustain breeding isolation [25–27]. The inclusion of May

322 precipitation in our most predictive models supports the above, in that study species (save two cottids)

spawn in May/June [83].

5. Conclusion

326   Hybridization occurs more frequently than expected in the White River Basin and is predictable based on

specifics of the environment. Although recognized as a creative evolutionary force, it is also considered a

328   maladaptive threat. Further research may blueprint an even more complicated future and potentially

demonstrate that groups (such as minnows) thrive proportional to their prolific gene exchange. Moreover,

330   hybridization is predicted to increase in frequency with global environmental change [11,12], a

recognition consistent with our finding of hybrid occurrence in lockstep with climate-related variables.

332   Therefore, baseline estimates are required to gauge the increase in hybridization, predict which

ecosystems will be so impacted (and how severely), and promote a more robust conservation/management

334   strategy that allows those impacts to be understood and adjudicated (if so needed) [10]. Future studies like

ours will be performed at the whole-genome level, with a resolution more robust for detecting and

336   untangling hybridization and its genomic consequences [24].

Acknowledgments

We thank M. Flurry, M. George, T. Goodhart, K. Hollar, and M. Reed, who assisted with DNA extractions. The Arkansas High-Performance Computing Center provided analytical resources.

Funding

Funding was provided by the University of Arkansas Distinguished Doctoral Fellowship and Harry and Jo Leggett Chancellor's Fellowship (ZDZ), the Bruker Professorship in Life Sciences (MRD), the Twenty-First Century Chair in Global Change Biology (MED), and an NSF Postdoctoral Research Fellowship in Biology (TKC) [DBI: 2010774]. The findings, conclusions, and opinions expressed in this article represent those of the authors and do not necessarily represent the views of the NSF or other affiliated or contributing organizations.

Conflict of interest

The authors declare that they have no competing interests.

Author contributions

ZDZ conceived the research with input from all authors. Specimen collection was done by ZDZ & TKC. Laboratory work, bioinformatics, data analysis, and manuscript drafting were done by ZDZ. All authors contributed to the analysis interpretation and critically revising the manuscript. MRD and MED administered funding through their endowments.

Data availability statement

Raw sequence files are accessioned in the NCBI GenBank Sequence Read Archive (SRA) BioProject: PRJNA809538. [84]. SNP alignments and R code are archived on Open Science Framework [85] .

REFERENCES

1. Hubbs CL. 1955 Hybridization between fish species in nature. *Syst. Zool.* **4**, 1–20. (doi:10.2307/sysbio/4.1.1)

2. Mallet J. 2005 Hybridization as an invasion of the genome. *Trends Ecol. Evol.* **20**, 229–237. (doi:10.1016/j.tree.2005.02.010)

3. Edelman NB, Mallet J. 2021 Prevalence and adaptive impact of introgression. *Annu. Rev. Genet.* **55**, 265–283. (doi:10.1146/annurev-genet-021821-020805)

4. Pfennig KS. 2021 Biased hybridization and its impact on adaptive introgression. *Trends Ecol. Evol.* **36**, 488–497. (doi:10.1016/j.tree.2021.02.010)

5. Lowe WH, Muhlfeld CC, Allendorf FW. 2015 Spatial sorting promotes the spread of maladaptive hybridization. *Trends Ecol. Evol.* **30**, 456–462. (doi:10.1016/j.tree.2015.05.008)

6. Rhymer JM, Simberloff D. 1996 Extinction by hybridization and introgression. *Annu. Rev. Ecol. Syst.* **27**, 83–109. (doi:10.1146/annurev.ecolsys.27.1.83)

7. Harrison RG, Larson EL. 2014 Hybridization, introgression, and the nature of species boundaries. *J. Hered.* **105**, 795–809. (doi:10.1093/jhered/esu033)

8. Taylor SA, Larson EL. 2019 Insights from genomes into the evolutionary importance and prevalence of hybridization in nature. *Nat. Ecol. Evol.* **3**, 170–177. (doi:10.1038/s41559-018-0777-y)

9. Todesco M *et al.* 2016 Hybridization and extinction. *Evol. Appl.* **9**, 892–908. (doi:10.1111/EVA.12367)

10. Brauer CJ, Sandoval-Castillo J, Gates K, Hammer MP, Unmack PJ, Bernatchez L, Beheregaray LB. 2023 Natural hybridization reduces vulnerability to climate change. *Nat. Clim. Chang.* , early access. (doi:10.1038/s41558-022-01585-1)

11. Cordonnier M, Gayet T, Escarguel G, Kaufmann B. 2019 From hybridization to introgression between two closely related sympatric ant species. *J. Zool. Syst. Evol. Res.* **57**, 778–788. (doi:10.1111/jzs.12297)

12. Muhlfeld CC, Kovach RP, Jones LA, Al-Chokhachy R, Boyer MC, Leary RF, Lowe WH, Luikart G, Allendorf FW. 2014 Invasive hybridization in a threatened species is accelerated by climate change. *Nat. Clim. Chang.* **4**, 620–624. (doi:10.1038/nclimate2252)

13. Becker M *et al.* 2013 Hybridization may facilitate in situ survival of endemic species through periods of climate change. *Nat. Clim. Chang.* **3**, 1039–1043. (doi:10.1038/nclimate2027)

14. Hobbs JPA, Richards ZT, Popovic I, Lei C, Staeudle TM, Montanari SR, DiBattista JD. 2022 Hybridisation and the evolution of coral reef biodiversity. *Coral Reefs* **41**, 535–549. (doi:10.1007/s00338-021-02193-9)

15. Justyn NM, Callaghan CT, Hill GE. 2020 Birds rarely hybridize: A citizen science approach to estimating rates of hybridization in the wild. *Evolution* **74**, 1216–1223. (doi:10.1111/evo.13943)

16. Ng I, Bellwood DR, Siqueira AC. 2022 Do currents shape global patterns of hybrid richness in coral reef fishes? *Glob. Ecol. Biogeogr.* **31**, 2524–2540. (doi:10.1111/geb.13592)

17. Guo Q. 2014 Plant hybridization: The role of human disturbance and biological invasion. *Divers. Distrib.* **20**, 1345–1354. (doi:10.1111/ddi.12245)

18. Schwenk K, Brede N, Streit B. 2008 Introduction. Extent, processes and evolutionary impact of

interspecific hybridization in animals. *Philos. Trans. R. Soc. B Biol. Sci.* **363**, 2805–2811. (doi:10.1098/rstb.2008.0055)

19. Scribner KT, Page KS, Bartron ML. 2000 Hybridization in freshwater fishes: A review of case studies and cytonuclear methods of biological inference. *Rev. Fish Biol. Fish.* **10**, 293–323. (doi:10.1023/A:1016642723238)

20. Corush JB, Fitzpatrick BM, Wolfe EL, Keck BP. 2021 Breeding behaviour predicts patterns of natural hybridization in North American minnows (Cyprinidae). *J. Evol. Biol.* **34**, 486–500. (doi:10.1111/jeb.13751)

21. Coyne JA, Orr HA. 1998 The evolutionary genetics of speciation. *Philos. Trans. R. Soc. B Biol. Sci.* **353**, 287–305. (doi:10.1098/rstb.1998.0210)

22. Fitzpatrick BM. 2004 Rates of evolution of hybrid inviability in birds and mammals. *Evolution* **58**, 1865–70. (doi:10.1111/j.0014-3820.2004.tb00471.x)

23. Seehausen O. 2004 Hybridization and adaptive radiation. *Trends Ecol. Evol.* **19**, 198–207. (doi:10.1016/j.tree.2004.01.003)

24. Payseur BA, Rieseberg LH. 2016 A genomic perspective on hybridization and speciation. *Mol. Ecol.* **25**, 2337–2360. (doi:10.1111/mec.13557)

25. Takemura A, Rahman MS, Park YJ. 2010 External and internal controls of lunar-related reproductive rhythms in fishes. *J. Fish Biol.* **76**, 7–26. (doi:10.1111/j.1095-8649.2009.02481.x)

26. Engström-Öst J, Candolin U. 2007 Human-induced water turbidity alters selection on sexual displays in sticklebacks. *Behav. Ecol.* **18**, 393–398. (doi:10.1093/beheco/arl097)

27. Moran RL, Zhou M, Catchen JM, Fuller RC. 2017 Male and female contributions to behavioral isolation in darters as a function of genetic distance and color distance. *Evolution* **71**, 2428–2444. (doi:10.1111/evo.13321)

28. Grabenstein KC, Taylor SA. 2018 Breaking barriers: Causes, consequences, and experimental utility of human-mediated hybridization. *Trends Ecol. Evol.* **33**, 198–212. (doi:10.1016/j.tree.2017.12.008)

29. Wiley EO, Mayden RL. 1985 Species and speciation in phylogenetic systematics, with examples from the North American fish fauna. *Ann. Missouri Bot. Gard.* **72**, 596. (doi:10.2307/2399217)

30. Robison HW, Buchanan TM. 2020 *Fishes of Arkansas*. 2nd edn. Fayetteville, AR: University of Arkansas Press.

31. Fox JT, Magoulick DD. 2019 Predicting hydrologic disturbance of streams using species occurrence data. *Sci. Total Environ.* **686**, 254–263. (doi:10.1016/j.scitotenv.2019.05.156)

32. Warren ML *et al.* 2000 Diversity, distribution, and conservation status of the native freshwater fishes of the southern United States. *Fisheries* **25**, 7–31. (doi:10.1577/1548-8446(2000)025<0007:DDACSO>2.0.CO;2)

33. Peterson BK, Weber JN, Kay EH, Fisher HS, Hoekstra HE. 2012 Double digest RADseq: An inexpensive method for de novo SNP discovery and genotyping in model and non-model species. *PLoS One* **7**, e37135. (doi:10.1371/journal.pone.0037135)

34. Chafin TK, Douglas MR, Martin BT, Douglas ME. 2019 Hybridization drives genetic erosion in sympatric desert fishes of western North America. *Heredity* **123**, 759–773. (doi:10.1038/s41437-019-0259-2)

35. Eaton DAR, Overcast I. 2020 Ipyrad: Interactive assembly and analysis of RADseq datasets. *Bioinformatics* **36**, 2592–2594. (doi:10.1093/bioinformatics/btz966)

36. Fan H, Ives AR, Surget-Groba Y. 2018 Reconstructing phylogeny from reduced-representation
genome sequencing data without assembly or alignment. *Mol. Ecol. Resour.* **18**, 1482–1491.
(doi:10.1111/1755-0998.12921)

37. Gosselin T. 2020 R package radiator: RADseq data exploration, manipulation and visualization
using R. See https://thierrygosselin.github.io/radiator/.

38. Cerca J *et al.* 2021 Removing the bad apples: A simple bioinformatic method to improve loci-
recovery in de novo RADseq data for non-model organisms. *Methods Ecol Evol* **12**, 805–817.
(doi:10.1111/2041-210X.13562)

39. R Core Team. 2022 R: A language and environment for statistical computing.
(doi:10.1038/sj.hdy.6800737)

40. Dray S, Dufour A. 2007 The ade4 Package: Implementing the duality diagram for ecologists. *J.
Stat. Softw.* **22**, 1–20. (doi:10.18637/jss.v022.i04)

41. Frichot E, François O. 2015 LEA: An R package for landscape and ecological association studies.
Methods Ecol. Evol. **6**, 925–929. (doi:10.1111/2041-210X.12382)

42. Beugin MP, Gayet T, Pontier D, Devillard S, Jombart T. 2018 A fast likelihood solution to the
genetic clustering problem. *Methods Ecol. Evol.* **9**, 1006–1016. (doi:10.1111/2041-210X.12968)

43. Jombart T. 2008 Adegenet: A R package for the multivariate analysis of genetic markers.
Bioinformatics **24**, 1403–1405. (doi:10.1093/bioinformatics/btn129)

44. Anderson EC, Thompson EA. 2002 A model-based method for identifying species hybrids using
multilocus genetic data. *Genetics* **160**, 1217–1229. (doi:10.1093/genetics/160.3.1217)

45. Wringe BF, Stanley RRE, Jeffery NW, Anderson EC, Bradbury IR. 2017 hybriddetective: A
workflow and package to facilitate the detection of hybridization using genomic data in r. *Mol.
Ecol. Resour.* **17**, e275–e284. (doi:10.1111/1755-0998.12704)

46. Keenan K, Mcginnity P, Cross TF, Crozier WW, Prodöhl PA. 2013 diveRsity: An R package for
the estimation and exploration of population genetics parameters and their associated errors.
Methods Ecol. Evol. **4**, 782–788. (doi:10.1111/2041-210X.12067)

47. Griffith DM, Veech JA, Marsh CJ. 2016 cooccur : Probabilistic species co-cccurrence analysis in
R. *J. Stat. Softw.* **69**, 1–17. (doi:10.18637/jss.v069.c02)

48. Linke S *et al.* 2019 Global hydro-environmental sub-basin and river reach characteristics at high
spatial resolution. *Sci. data* **6**, 1–15. (doi:10.1038/s41597-019-0300-6)

49. Manco F, Lang SDJ, Trathan PN. 2022 Predicting foraging dive outcomes in chinstrap penguins
using biologging and animal-borne cameras. *Behav. Ecol.* **33**, 989–998.
(doi:10.1093/beheco/arac066)

50. O'Brien R, Ishwaran H. 2019 A random forests quantile classifier for class imbalanced data.
Pattern Recognit. **90**, 232–249. (doi:10.1016/J.PATCOG.2019.01.036)

51. Ishwaran H, Kogalur UB. 2023 Fast unified random forests for survival, regression, and
classification (RF-SRC). See https://cran.r-project.org/package=randomForestSRC.

52. Ishwaran H, Kogalur U, Gorodeski E, Minn A, Lauer M. 2010 High-dimensional variable
selection for survival data. *J. Am. Stat. Assoc.* **105**, 205–217. (doi:10.1198/jasa.2009.tm08622)

53. Matthews WJ. 1998 *Patterns in freshwater fish ecology*. New York: Chapman and Hall.

54. Matthews WJ, Robison HW. 1988 The distribution of the fishes of Arkansas: A multivariate
analysis. *Copeia* **1988**, 358. (doi:10.2307/1445876)

55.   Bangs MR, Douglas MR, Brunner PC, Douglas ME. 2020 Reticulate evolution as a management
490   challenge: Patterns of admixture with phylogenetic distance in endemic fishes of western North
491   America. *Evol. Appl.* **13**, 1400–1419. (doi:10.1111/eva.13042)

492   56.   Arnold ML. 1992 Natural hybridization as an evolutionary process. *Annu. Rev. Ecol. Syst.* **23**,
493   237–261. (doi:10.1146/annurev.es.23.110192.001321)

494   57.   Meng C, Kubatko LS. 2009 Detecting hybrid speciation in the presence of incomplete lineage
495   sorting using gene tree incongruence: A model. *Theor. Popul. Biol.* **75**, 35–45.
496   (doi:10.1016/j.tpb.2008.10.004)

497   58.   Rakocinski CF. 1980 Hybridization and introgression between *Campostoma oligolepis and C.
498   anomalum pullum* (Cypriniformes: Cyprinidae). *Copeia* **1980**, 584. (doi:10.2307/1444433)

499   59.   Grady JM, Cashner RC. 1988 Evidence of extensive intergeneric hybridization among the cyprinid
500   fauna of Clark Creek, Wilkinson Co., Mississippi. *Southwest. Nat.* **33**, 137. (doi:10.2307/3671888)

501   60.   Cross FB. 1954 Fishes of Cedar Creek and the South Fork of the Cottonwood River, Chase
502   County, Kansas. *Trans. Kansas Acad. Sci.* **57**, 303. (doi:10.2307/3626057)

503   61.   Barthel BL, Dorothy OM, Philipp DP. 2010 Molecular genetic confirmation of hybridization
504   between Largemouth and Smallmouth Bass (*Micropterus*) in the wild. *Copeia* **2010**, 671–675.
505   (doi:10.1643/CG-09-186)

506   62.   Keck BP, Near TJ. 2009 Patterns of natural hybridization in darters (Percidae: Etheostomatinae).
507   *Copeia* **2009**, 758–773. (doi:10.1643/CI-09-008)

508   63.   Matthews W. 1982 Small fish community structure in Ozark streams: Structured assembly patterns
509   or random abundance of species? *Am. Midl. Nat.* **107**, 42–54. (doi:Doi 10.2307/2425187)

510   64.   Ray JM, Lang NJ, Wood RM, Mayden RL. 2008 History repeated: recent and historical
511   mitochondrial introgression between the Current Darter *Etheostoma uniporum* and Rainbow
512   Darter *Etheostoma caeruleum* (Teleostei: Percidae). *J. Fish Biol.* **72**, 418–434.
513   (doi:10.1111/J.1095-8649.2007.01732.X)

514   65.   Kim D, Taylor AT, Near TJ. 2022 Phylogenomics and species delimitation of the economically
515   important Black Basses (*Micropterus*). *Sci. Rep.* **12**, 9113. (doi:10.1038/s41598-022-11743-2)

516   66.   Mayr E. 1963 *Animal species and evolution*. Cambridge: Harvard University Press.

517   67.   Dowling TE, Demarais BD. 1993 Evolutionary significance of introgressive hybridization in
518   cyprinid fishes. *Nature* **362**, 444–446. (doi:10.1038/362444a0)

519   68.   Tricou T, Tannier E, De Vienne DM. 2022 Ghost lineages highly influence the interpretation of
520   introgression tests. *Syst. Biol.* **71**, 1147–1158. (doi:10.1093/SYSBIO/SYAC011)

521   69.   Hersch-Green EI, Cronn R. 2009 Tangled trios?: Characterizing a hybrid zone in Castilleja
522   (Orobanchaceae). *Am. J. Bot.* **96**, 1519–1531. (doi:10.3732/AJB.0800357)

523   70.   Wallis GP, Cameron-Christie SR, Kennedy HL, Palmer G, Sanders TR, Winter DJ. 2017
524   Interspecific hybridization causes long-term phylogenetic discordance between nuclear and
525   mitochondrial genomes in freshwater fishes. *Mol. Ecol.* **26**, 3116–3127. (doi:10.1111/mec.14096)

526   71.   Burford MO, Bernardi G, Carr MH. 2011 Analysis of individual year-classes of a marine fish
527   reveals little evidence of first-generation hybrids between cryptic species in sympatric regions.
528   *Mar. Biol.* **158**, 1815–1827. (doi:10.1007/s00227-011-1694-7)

529   72.   Lamer JT, Dolan CR, Petersen JL, Chick JH, Epifanio JM. 2010 Introgressive hybridization
530   between Bighead Carp and Silver Carp in the Mississippi and Illinois Rivers. *North Am. J. Fish.
531   Manag.* **30**, 1452–1461. (doi:10.1577/m10-053.1)

73. Twyford AD, Ennos RA. 2012 Next-generation hybridization and introgression. *Heredity* **108**, 179–189. (doi:10.1038/hdy.2011.68)

74. Peoples BK, Floyd SP, Frimpong EA. 2016 Nesting microhabitat comparison of Central Stoneroller and Bluehead Chub: potential inference for host-switching by nest associates. *J. Freshw. Ecol.* **31**, 251–259. (doi:10.1080/02705060.2015.1091390)

75. Matthews WJ, Marsh-Matthews E. 2017 *Stream fish community dynamics*. Baltimore, MD: Johns Hopkins University Press.

76. De Santis V, Quadroni S, Britton RJ, Carosi A, Gutmann Roberts C, Lorenzoni M, Crosa G, Zaccara S. 2021 Biological and trophic consequences of genetic introgression between endemic and invasive Barbus fishes. *Biol. Invasions* **23**, 3351–3368. (doi:10.1007/s10530-021-02577-6)

77. Gibson I, Welsh AB, Welsh SA, Cincotta DA. 2019 Genetic swamping and possible species collapse: tracking introgression between the native Candy Darter and introduced Variegate Darter. *Conserv. Genet.* **20**, 287–298. (doi:10.1007/s10592-018-1131-2)

78. Ottenburghs J. 2021 The genic view of hybridization in the Anthropocene. *Evol. Appl.* **14**, 2342–2360. (doi:10.1111/eva.13223)

79. Gupta VK, Ayalew TB, Mantilla R, Krajewski WF. 2015 Classical and generalized Horton laws for peak flows in rainfall-runoff events. *Chaos* **25**, 75408. (doi:10.1063/1.4922177)

80. King AJ, Tonkin Z, Mahoney J. 2009 Environmental flow enhances native fish spawning and recruitment in the Murray River, Australia. *River Res. Appl.* **25**, 1205–1218. (doi:10.1002/RRA.1209)

81. Fitzsimons JM, Nishimoto RT. 1995 Use of fish behavior in assessing the effects of Hurricane Iniki on the Hawaiian island of Kaua'i. *Environ. Biol. Fishes* **43**, 39–50. (doi:10.1007/BF00001816/METRICS)

82. Sear DA. 1993 Fine sediment infiltration into gravel spawning beds within a regulated river experiencing floods: Ecological implications for salmonids. *Regul. Rivers Res. Manag.* **8**, 373–390. (doi:10.1002/RRR.3450080407)

83. Frimpong EA, Angermeier PL. 2009 Fish Traits: A database of ecological and life-history traits of freshwater fishes of the United States. *Fisheries* **34**, 487–495. (doi:10.1577/1548-8446-34.10.487)

84. Zbinden ZD, Douglas MR, Chafin TK, Douglas ME. 2022 Data from: Ozark fish community genomics: ddRAD seq (BioProject: PRJNA809538). *NCBI sequence read archive.* See https://www.ncbi.nlm.nih.gov/bioproject/PRJNA809538

85. Zbinden ZD, Douglas MR, Chafin TK, Douglas ME. 2022 Data from: White river fish hybridization. *Open Science Framework*. (doi:10.17605/OSF.IO/PSBDH)

565   **TABLE 1** Samples (N=33 species) from the White River Basin, USA. N_I=number of individuals/species
566   (post-filtering), N_S=number of collection sites/species (post-filtering), Miss=mean proportion of missing
567   data, Depth=mean sequencing depth, Ho=observed heterozygosity, N_{FA}=number of individuals/family,
568   SNPs=single nucleotide polymorphisms in family panel (post-filtering).

Family	Common Name	Scientific Name	N_I	N_S	Miss	Depth	Ho	N_{FA}	SNPs
Centrarchidae	Bluegill Sunfish	*Lepomis macrochirus*	69	23	0.15	46	0.028		
	Longear Sunfish	*Lepomis megalotis*	239	54	0.09	47	0.036	375	1926
	Smallmouth Bass	*Micropterus dolomieu*	43	30	0.47	39	0.011		
	Largemouth Bass	*Micropterus salmoides*	24	17	0.46	41	0.013		
Cottidae	Banded Sculpin	*Cottus carolinae* [†]	33	18	0.08	64	0.043	75	5344
	Ozark Sculpin	*Cottus hypselurus* [‡]	42	10	0.16	63	0.057		
Fundulidae	Northern Studfish	*Fundulus catenatus*	105	28	0.12	37	0.008	226	2366
	Blackspotted Topminnow	*Fundulus olivaceus*	121	34	0.20	33	0.024		
Ictaluridae	Ozark Madtom	*Noturus albater*	10	5	0.12	65	0.033		
	Slender Madtom	*Noturus exilis*	16	12	0.09	51	0.042	31	2744
	Black River Madtom	*Noturus maydeni*	5	3	0.28	34	0.034		
Leuciscidae	Central Stoneroller	*Campostoma anomalum*	128	44	0.27	58	0.010		
	Largescale Stoneroller	*Campostoma oligolepis*	135	41	0.29	52	0.008		
	Southern Redbelly Dace	*Chrosomus erythrogaster*	38	10	0.41	38	0.001		
	Whitetail Shiner	*Cyprinella galactura*	75	16	0.19	59	0.012		
	Steelcolor Shiner	*Cyprinella whipplei*	30	8	0.27	65	0.010		
	Striped Shiner	*Luxilus chrysocephalus*	63	18	0.18	59	0.013		
	Duskystrip Shiner	*Luxilus pilsbryi*	258	33	0.09	76	0.023		
	Bleeding Shiner	*Luxilus zonatus*	100	17	0.09	85	0.013	1507	343
	Redfin Shiner	*Lythrurus umbratilis*	24	5	0.26	47	0.004		
	Bigeye Shiner	*Notropis boops*	226	31	0.10	89	0.019		
	Ozark Minnow	*Notropis nubilus*	193	35	0.11	76	0.016		
	Carmine Shiner	*Notropis percobromus*	67	15	0.20	66	0.010		
	Telescope Shiner	*Notropis telescopus*	83	15	0.14	75	0.006		
	Bluntnose Minnow	*Pimephales notatus*	55	24	0.25	73	0.007		
	Creek Chub	*Semotilus atromaculatus*	32	14	0.32	51	0.001		
Percidae	Greenside Darter	*Etheostoma blennioides*	62	26	0.24	55	0.006		
	Rainbow Darter	*Etheostoma caeruleum*	348	53	0.09	54	0.026		
	Fantail Darter	*Etheostoma flabellare*	26	11	0.33	38	0.001		
	Yoke Darter	*Etheostoma juliae* [§]	63	15	0.24	58	0.011	651	687
	Orangethroat Darter	*Etheostoma spectabile* [¶]	50	10	0.24	48	0.023		
	Current Darter	*Etheostoma uniporum*	18	7	0.25	41	0.005		
	Banded Darter	*Etheostoma zonale*	84	26	0.25	51	0.018		

Note: Several taxonomic changes have been adopted in the most recent *Fishes of Arkansas* [30] based on the literature but have
yet to be recognized by the American Fisheries Society or Eschmeyers's Catalog of Fishes.
[†] *Uranidea carolinae*
[‡] *Uranidea immaculata* (Knobfin Sculpin)
[§] *Nothonotus juliae*
[¶] *Etheostoma* sp. *cf. spectabile* (Ozark Darter)

 TABLE 2 Hybridization within six families sampled across the White River Basin, USA. Columns
include: N Species=number of species analyzed; Unique pairs=number of unique intra-familial; Pairs Not
Co-occurring=number of pairs within a family not found together at the same site; N Indiv=number of
individuals examined; N Hybrids=number of hybrids detected; Percent Indiv=Hybrid percentage. Unique
species-pairs with hybrids are only considered for ancestry between two species (i.e., species triplets were
not counted).

Family	N Species	Unique Pairs	Pairs Not Co-Occurring	N Indiv.	N Hybrids	Percent Indiv.	Species w/ Hybrids	Percent Species w/ Hybrid	Unique Pairs w/ Hybrid	Percent Pairs w/ Hybrid
Fundulidae	2	1	0	226	0	0.0%	0	0%	0	0%
Cottidae	2	1	0	75	0	0.0%	0	0%	0	0%
Ictaluridae	3	3	1	31	1	3.2%	3	100%	0	0%
Centrarchidae	4	6	0	375	1	0.3%	2	50%	1	17%
Percidae	7	21	5	651	2	0.3%	4	57%	2	10%
Leuciscidae	15	105	7	1507	66	4.4%	15	100%	15	14%
Overall	33	137	13	2865	70	2.4%	24	73%	18	13%

TABLE 3 Observed genotypic frequency classes of hybrid individuals inferred from NewHybrids analysis for 18 hybridizing fish species-pairs collected across the White River basin, USA. Note that 8 multi-specific hybrids were not included. Putative hybrids were assigned to a genotype frequency class (F_1, F_2, Bx[=backcross], pure) based on Bayesian posterior probability >0.70.

No.	Species A	Species B	N Indiv.	F_1	F_2	Bx	Pure
1	*Campostoma anomalum*	*Campostoma oligolepis*	29	5	-	11	13
2	*Campostoma anomalum*	*Chrosomus erythrogaster*	1	1	-	0	-
3	*Campostoma anomalum*	*Luxilus pilsbryi*	1	-	1	0	-
4	*Campostoma oligolepis*	*Notropis telescopus*	1	1	-	0	-
5	*Cyprinella galactura*	*Cyprinella whipplei*	2	1	-	1	-
6	*Cyprinella whipplei*	*Lythrurus umbratilis*	1	-	-	1	-
7	*Luxilus chrysocephalus*	*Luxilus zonatus*	2	1	-	0	1
8	*Luxilus chrysocephalus*	*Semotilus atromaculatus*	1	-	-	1	-
9	*Luxilus pilsbryi*	*Lythrurus umbratilis*	6	-	1	5	-
10	*Luxilus pilsbryi*	*Notropis percobromus*	2	1	-	1	-
11	*Luxilus pilsbryi*	*Luxilus zonatus*	8	-	-	0	8
12	*Luxilus pilsbryi*	*Luxilus chrysocephalus*	1	-	-	1	-
13	*Luxilus zonatus*	*Pimephales notatus*	1	1	-	0	-
14	*Notropis boops*	*Notropis nubilus*	2	-	-	1	1
15	*Pimephales notatus*	*Semotilus atromaculatus*	1	-	1	0	-
16	*Micropterus salmoides*	*Micropterus dolomieu*	1	1	-	0	-
17	*Etheostoma spectabile*	*Etheostoma caeruleum*	1	-	-	1	-
18	*Etheostoma juliae*	*Etheostoma zonale*	1	-	-	1	-
		Totals	62	12	3	24	23

587 **TABLE 4** Mean, minimum, and maximum values of Weir and Cockerham's pairwise F_{ST} calculated
588 among species within families collected across the White River Basin, USA. Families Cottidae and
589 Fundulidae were represented by only two species each, hence one value of F_{ST}.

Family	Mean F_{ST}	Min F_{ST}	Max F_{ST}
Centrarchidae	0.94	0.85	0.98
Cottidae	-	0.82	0.82
Fundulidae	-	0.98	0.98
Ictaluridae	0.81	0.66	0.91
Leuciscidae	0.92	0.65	0.99
Percidae	0.93	0.78	0.98

FIGURE 1 First two principal components (PCs) derived from SNP genotypes in six stream fish families collected across the White River Basin, USA. The variance explained by each component is in the bottom left or right corner of each plot. *N*=numbers of individuals/family.

595
FIGURE 2 Calculated ancestral proportions for *N*=2,865 individuals from *N*=33 fish species grouped by
family collected from the White River Basin, USA. *N*=number of individuals, *K*=optimum number of
clusters.

FIGURE 3 Significant predictors of hybridization across the White River Basin, USA. Maps depict
collection sites with circles (red indicates presence of a hybrid). In the first map, site-diameter is scaled by
species richness, while in the remaining maps, it is scaled by the probability of hybrid occurrence based
on the model. Basin backgrounds differ for enhanced visibility. Below the maps are partial marginal
effects plots showing relationships between probability of hybrid individuals occurring per four predictive
variables. The mean (black) and 95% confidence intervals (red) are shown, along with distribution of
variable values across sites (black tick marks above x-axis).