The dispensability of 14-3-3 proteins for the regulation of human cardiac sodium channel Na$_v$1.5

Short running title: Role of 14-3-3 in Na$_v$1.5 regulation

Oksana Iamshanova[1], Anne-Flore Hämmerli[1], Elise Ramaye[1], Arbresh Seljmani[1,2], Daniela Ross-Kaschitza[1], Noëlia Schärz[1,2], Maria Essers[1], Sabrina Guichard[1], Jean-Sébastien Rougier[1], Hugues Abriel[1]

[1] - Institute of Biochemistry and Molecular Medicine and Swiss National Center of Competence in Research (NCCR) TransCure, University of Bern, Bern, Switzerland

[2] - medi - Center for Medical Education, Bern, Switzerland

Correspondence should be addressed either to:
Oksana Iamshanova, PhD (ORCID:0000-0002-1984-0013) or Abriel Hugues, MD, PhD (ORCID: 0000-0003-0465-5138)
Institute of Biochemistry and Molecular Medicine, University of Bern
Bühlstrasse 28, CH-3012 Bern, Switzerland
e-mail: oksana.iamshanova@unibe.ch
e-mail: hugues.abriel@unibe.ch

Summary

This work shows that only 14-3-3η, exhibits weak/transient interaction with Na$_v$1.5, and does not modify its total protein expression, plasmalemmal trafficking, and basal biophysical properties of the whole-cell current. Furthermore, inhibition of endogenous 14-3-3/ligand interactions with difopein does not affect the dimerization of Na$_v$1.5. Therefore, 14-3-3 proteins are suggested to be dispensable for the Na$_v$1.5 regulation in a heterologous expression system.

Abstract

Background: 14-3-3 proteins are ubiquitous proteins that play a role in cardiac physiology (*e.g.*, metabolism, development, and cell cycle). Furthermore, 14-3-3 proteins were proposed to regulate the electrical function of the heart by interacting with several cardiac ion channels, including the voltage-gated sodium channel Na$_v$1.5. Given the many cardiac arrhythmias associated with Na$_v$1.5 dysfunction, understanding its regulation by the protein partners is crucial.
Aims: In this study, we aimed to investigate the role of 14-3-3 proteins in the regulation of the human cardiac sodium channel Na$_v$1.5.
Methods and Results: Amongst the seven 14-3-3 isoforms, only 14-3-3η (encoded by *YWHAH* gene) weakly co-immunoprecipitated with Na$_v$1.5 when heterologously co-expressed in tsA201 cells. Total and cell surface expression of Na$_v$1.5 was however not modified by 14-3-3η overexpression or inhibition with difopein, and 14-3-3η did not affect physical interaction between Na$_v$1.5 α-α subunits. The current-voltage relationship and the amplitude of Na$_v$1.5-mediated sodium peak current density were also not changed.
Conclusions: Our findings illustrate that the direct implication of 14-3-3 proteins in regulating Na$_v$1.5 is not evident in a transformed human kidney cell line tsA201.

Keywords

Introduction

In the heart voltage-gated sodium channel, Na$_v$1.5 (encoded by *SCN5A* gene), is responsible for the genesis and propagation of the action potential in the contractile myocardium. The function of Na$_v$1.5 largely relies on the interaction with its protein partners that regulate proper localization, turnover, and biophysical properties of the channel (Dong et al., 2020). Dysfunction of Na$_v$1.5 is associated with various lethal diseases such as cardiac arrhythmias and cardiomyopathy. Therefore, understanding the relationship between Na$_v$1.5 and its regulatory protein partners may uncover novel therapeutic strategies for patients suffering from *SCN5A*-related diseases.

One of the protein partners of Na$_v$1.5 was reported to be 14-3-3 proteins, which are a family of conserved regulatory molecules expressed in all eukaryotic cells. 14-3-3 can bind target proteins, modulating protein-protein interactions and the activity of their ligands due to inducing structural rearrangements, masking or unmasking of the functional sites, and changing the ligand's intracellular localization (Obsil and Obsilova, 2011). *YWHAB*, *YWHAG*, *YWHAE*, *SFN*, *YWHAH*, *YWHAQ*, and *YWHAZ* genes encode the mammalian 14-3-3 isoforms β/α, γ, ε, σ, η, θ, and ζ/δ, respectively, where α and δ represent the phosphorylated forms. All 14-3-3 isoforms exhibit a high level of structural homology and consist of 9 antiparallel α-helices arranged in an L-shape (Obsil and Obsilova, 2011; Yang et al., 2006). Through their N-terminal region, 14-3-3 proteins form homo- and heterodimers (Yang et al., 2006). Each ~30-kDa monomer binds to a variety of the target proteins at the consensus sequences R(S/X)XpSXP, RXXXpSXP, and pS/pTX1–2-COOH of the phosphorylated motifs (Thompson and Goldspink, 2022). The best-studied examples of the unphosphorylated targets are P*seudomonas aeruginosa* virulence factor exoenzyme S and R18, a peptide identified from a phage display library (Henriksson et al., 2002; Wang et al., 1999). R18 is widely used as a potent antagonist of 14-3-3 protein/ligand interactions due to its high affinity with all 14-3-3 isoforms (K_d = 80 nM) followed by occlusion of their ligand-binding groove (Wang et al., 1999). Accordingly, dimer of R18, difopein, is also a competitive non-isoform specific inhibitor of 14-3-3 protein/ligand interactions (Masters and Fu, 2001).

14-3-3 proteins target a wide variety of ligands and regulate many cellular processes (Obsil and Obsilova, 2011). Accordingly, 14-3-3 proteins were shown to be involved in cardiac physiology, including cardiomyocyte development, cell cycle, and Ca^{2+} signaling (Gittenberger-de Groot et al., 2016; Kosaka et al., 2012; Martens et al., 2021; Menzel et al., 2020). Importantly, 14-3-3 was proposed to play a role in cardiac electrophysiology due to their interactions with cardiac ion channels, exchangers, and pumps (Thompson and Goldspink, 2022). Specifically, the α-subunit of Na$_v$1.5 channel was shown to coimmunoprecipitate with 14-3-3 proteins in mouse heart tissue as well as in heterologous expression systems (Allouis et al., 2006). Although 14-3-3η, -θ, and -ζ were identified by yeast 2-hybrid screen to directly interact with Na$_v$1.5, only 14-3-3η was investigated for functional consequences on the channel activity (Allouis et al., 2006). In the presence of exogenous 14-3-3η, the density of sodium current and activation curve were unchanged, while the inactivation curve was negatively shifted (Allouis et al., 2006). Based on this data, computer simulations suggested proarrhythmic effects of 14-3-3η (Allouis et al., 2006). Nevertheless, transgenic mice

expressing a cardiac-specific loss-of-function 14-3-3η mutant exhibited normal cardiac morphology, cardiomyocyte appearance, and ventricular systolic function under basal conditions, while nothing was reported about its cardiac electrical activity (Xing et al., 2000). Interestingly, immunostaining analysis revealed colocalization between exogenous 14-3-3η and Na$_v$1.5 in the intercalated discs of adult rabbit cardiomyocytes, suggesting a role in clusterization (Allouis et al., 2006). In line with these findings, 14-3-3 proteins were reported to mediate the coupled gating of Na$_v$1.5 (Clatot et al., 2017). Although two 14-3-3 binding sites were identified in the intracellular loop between domains I and II of Na$_v$1.5 by co-immunoprecipitation analysis, it was unclear which isoform mediated the channel gating properties through its direct interaction (Clatot et al., 2017). Therefore, our goal was to investigate the direct role of all 14-3-3 isoforms on Na$_v$1.5 in a heterologous expression system. Specifically, we addressed their interaction, expression, trafficking, and sodium current.

Materials and Methods

cDNA constructs and cloning

Template cDNA constructs and their derivatives are listed in Table 1.

cDNA construct	Transcript (Genbank)	Source	
pcDNA3.1	Empty vector	V86020, Invitrogen	
pcDNA3.1-*SCN5A*-WT	NM_000335.5	GenScript, NJ, USA	
pcDNA3.1-3X-FLAG-*SCN5A*-WT			
pcDNA3.1-HA-*SCN5A*-WT			
pFN217K-LgBiT(Nter)-*SCN5A*-WT		Cloned from pcDNA3.1-*SCN5A*-WT into NanoBiT® CMV Flexi BiBit ready vectors (ID# CS1603B33) according to Flexi® vector systems technical manual #TM254 from Promega (Switzerland)	
pFC219K-*SCN5A*-WT-LgBiT(Cter)			
pFN218K-SmBiT(Nter)-*SCN5A*-WT			
pFC220K-*SCN5A*-WT-SmBiT(Cter)			
CMV/SmBiT-CA/BlastR	NM_002730.4	NanoBiT® CMV PPI control vectors (ID# CS1603B54) from Promega (Switzerland)	
CMV/LgBiT-R2A/HygR	NM_004157.4		
BiBiT-RI/SmBiT-CA/LgBiT-R2A/BlastR	NM_002730.4 and NM_004157.4		
CMV/HaloTag®-SmBiT	HM157289.1		
pcDNA3-HA-*YWHAB*-WT	NM_003404.5	a gift from Michael Yaffe (Yaffe, 2002)	Addgene #13270
pcDNA3-HA-*YWHAG*-WT	NM_012479.4		Addgene #13274
pcDNA3-HA-*YWHAE*-WT	NM_006761.5		Addgene #13273
pcDNA3-HA-*SFN*-WT	NM_006142.5		Addgene #11946
pcDNA3-HA-*YWHAH*-WT	NM_003405.4	GenScript, NJ, USA	
pcDNA3-HA-*YWHAQ*-WT	NM_006826.4		
pcDNA3-HA-*YWHAZ*-WT	NM_003406.4		
pcDNA3-FLAG-*YWHAB*-WT	NM_003404.5	a gift from Dario Diviani (University of Lausanne, Switzerland)	
pcDNA3-FLAG-*YWHAZ*-WT	NM_003406.4		
pSCM YFP-difopein	Peptide sequence: SADGAPHCVPRDLSWLD LEANMCLPGAAGLDSAD GAPHCVPRDLSWLDLEA NMCLPGAAGLE	a gift from Haian Fu (Emory University School of Medicine, GA, USA)	
pSCM YFP-mutant difopein R18(D12K,E14K)	Peptide sequence: PHCVPRDLSWLKLKANM CLP		
pIRES-*CD8*-WT	NM_001768.7	a gift from David Kass (Johns Hopkins University, MD, USA)	
pIRES-*CD8*-WT-*SCN1B*-WT	NM_001768.7 and NM_001037.5		

Table 1. Description of cDNA constructs used in this study.

To clone *SCN5A*-WT into NanoBiT® CMV Flexi BiBit ready vectors, SgfI and PmeI restriction sites were introduced, for which the primers were designed by the Flexi® Vector Primer Design Tool (https://ch.promega.com/techserv/tools/flexivectortool/) and purchased from Eurofins Genomics (Germany) (forward primer 5'-3': GTC GGC GAT CGC CAT GGC AAA CTT CCT ATT ACC TCG G; reverse primer 5'-3': GCG AGT TTA AAC CAC GAT GGA CTC ACG GTC CC).

Cell culture and transfection

Human embryonic kidney cells tsA201 (ECACC Cat# 96121229, RRID:CVCL_2737) were cultured up to 20 passages at 37°C with 5% CO_2 in Dulbecco's modified Eagle's culture medium (41965, Gibco™, Thermo Fisher Scientific, USA), supplemented with 2 mM L-glutamine (G7513, Sigma), 50 U/mL penicillin-streptomycin (15140122, Gibco™) and 10% heat-inactivated fetal bovine serum (10270-106, Lot: 2440045, Gibco™). When needed, cells were split using phosphate-buffered saline (PBS, 10010-015, Gibco™) and 0.05% Trypsin-EDTA (25300-054, Gibco™). Mycoplasma contamination status was tested weekly with PCR Mycoplasma Test Kit I/C (PK-CA91-1096, Promokine, PromoCell GmbH, Germany). cDNA constructs were introduced into cells by transfection with LipoD293™ (SL100668, SignaGen® Laboratories). Cells were harvested 48 hours after transfection unless stated otherwise. Stable cell lines tsA201-*SCN5A*-WT and tsA201-FLAG-*SCN5A*-WT were established by polyclonal selection and were maintained in culture growth media with 200 µg/mL Zeocin™ Selection Reagent (R25005, Gibco™).

RNA extraction, reverse transcription polymerase chain reaction, and agarose gel

electrophoresis

RNA extraction was performed on PBS-prewashed cell pellets using ReliaPrep™ RNA cell Miniprep Systems according to the manufacturer's instruction (Z6010, Promega). The concentration of extracted RNA was determined by NanoDrop™ One/One^C Microvolume UV-Vis Spectrophotometer (ND-ONE-W, Thermo Fischer Scientific™). Reverse transcription (RT) was performed using a High-Capacity cDNA Reverse Transcription kit (4375222, Applied Biosystems) followed by polymerase chain reaction (PCR) with GoTaq® G2 Master Mix (M7822, Promega) on Biometra TRIO Touch Thermocycler (207072X, LabGene Scientific, Switzerland). The PCR protocol was as follows: initial denaturation at 95°C for 2 minutes, 35 cycles of denaturation at 95°C for 1 minute, annealing at 60°C for 45 seconds and elongation at 72°C for 45 seconds per kb, and a final extension of 72°C for 5 minutes. PCR products were stored at 4°C. Primers with their expected amplicon sizes are listed in Table 2.

Target gene	Forward primer 5'-3'	Reverse primer 5'-3'	Amplicon size, bp
GAPDH	CCTTCATTGACCTCAACTACATGG	GGGATCTCGCTCCTGGAAGATGG	140
PSMB4	GCACTTTACAGAGGTCCAATCAC	CTCGGTAGTACAGCACTCGC	565
TBP	CTTGACCTAAAGACCATTGCACTTC	GAAACTTCACATCACAGCTCCC	266
YWHAB	CTGCTCTCTGTTGCCTACAAGAATGT	CTTCGTCTCCCTGGTTTTCCGAT	583
YWHAG	GCGAGCAACTGGTGCAGAAAGC	GTACTGGGTCTCGCTGCAATTCTT	341
YWHAE	GACAAGCTAAAAATGATTCGGGAATATCG	CTGAAGTCCATAGTGTCAGATTATCACG	472
SFN	GAGAGAGCCAGTCTGATCCAGAA	GTAGTAGTCACCCTTCATCTTCAGGTAG	381
YWHAH	TACGACGACATGGCCTCCG	CAAACTGTCTCCAGCTCCTTCTCA	233
YWHAQ	GCTCTCCGTGGCCTACAAGAAC	CGATCATCACCACACGCAACTTCA	285
YWHAZ	GATGACAAGAAAGGGATTGTCGATCAGT	CACTTAATGTATCAAGTTCAGCAATGGCT	217

Table 2. Description of primers used in this study.

Protein lysate extraction

Pre-washed (PBS) adherent monolayers of cells were scraped in 2 mL of cold PBS (pH 7,4) and pelleted by centrifugation for 5 minutes at 200 g at 4°C. Cell pellets were lysed in lysis buffer (50 mM NaCl, 50 mM imidazole/HCl, 2 mM 6-aminohexanoic acid, 1 mM EDTA, pH 7) with addition of cOmplete tablets EDTA-free (04693132001, Roche, Switzerland), 0.5 mM Na_3VO_4, 0.5 mM NaFl, 10 µg/mL aprotinin, 10 µg/mL leupeptin, 1 mM phenylmethylsulfonyl fluoride and 1% digitonin (D141, Sigma) for 1 hour at 4°C. Lysates were centrifuged at 16,100 g for 15 minutes at 4°C. The obtained supernatant was taken for further analysis. Protein concentration was determined by Coo Assay Protein Dosage Reagent (Uptima, UPF86420).

Co-immunoprecipitation

For immunoprecipitation nProtein A Sepharose 4 Fast Flow (17528001, Cytiva) or anti-FLAG® M2 Magnetic Beads (M8823, Sigma) were used in the proportion of 1 µL of beads per 20 µg of total protein lysate. Protein lysates were diluted in Tris-buffered saline (TBS, 20 M Tris, 0.1 mM NaCl, pH 7.6, HCl-adjusted) and were mixed with the TBS-prewashed antibody-coupled beads at 4°C overnight. After thoroughly washing the beads with TBS, the co-immunoprecipitated protein complex was eluted with 4X LDS sample buffer (NP0007, Invitrogen). The samples were analyzed with the immunoblotting technique (see below).

Cell surface biotinylation assay

Adherent cell monolayers were gently washed three times with cold PBS (pH 8) and incubated, while slowly rocking, with 1 mg/mL EZlink™ Sulfo-NHS-SS-Biotin (21331, Thermo Fisher Scientific) in PBS (pH 8) for 45 minutes at 4°C. Afterward, the excess biotin was quenched by washing with 50 mM Tris-HCl (pH 8), followed by several washes with PBS (pH 7,4). Extracted protein lysates were mixed with Streptavidin Sepharose High-Performance beads (GE Healthcare, USA) at 4°C overnight. After thoroughly washing the beads with washing buffer (0.1% BSA, 0.001% Tween20, PBS pH 7,4), the biotinylated fraction was eluted with 4X LDS sample buffer and analyzed with the immunoblotting technique (see below).

Sodium dodecyl sulfate–polyacrylamide gel electrophoresis (SDS-PAGE) and

immunoblotting analysis

Protein lysates containing LDS sample buffer and 100 mM dithiothreitol (DTT) were loaded onto 4-12% Tris-acetate acrylamide gels and let to run at 60 mV in Tris-acetate running buffer (50 mM Tris, 50 mM Tricine, 0.1% SDS, pH 8.3). Afterward, proteins were transferred to the nitrocellulose membrane by using the Trans-Blot Turbo system (1704158, Bio-Rad, USA). After validation of successful protein transfer with Ponceau S solution (0.1% Ponceau S, 12.5% acetic acid), the membranes were blocked with 5% bovine serum albumin (BSA) in TBS-T (TBS, 0.1% Tween20) for 1 hour at room temperature and then incubated with primary antibody dilutions at 4°C overnight. After incubating membranes with secondary antibodies for 1 hour at room temperature, the infrared fluorescent signals were revealed with LiCor Odyssey Infrared imaging

system (LI-COR Biosciences) and quantified with ImageJ software (Rasband, W.S., U. S. National Institutes of Health, Bethesda, Maryland, USA).

Primary and secondary antibodies used in this study are listed in Table 3.

Antigen	Dilution	Host	Epitope	RRID	Source
FLAG	1:1000	mouse	DYKDDDDK	AB_262044	F1804, Sigma-Aldrich
GFP	1:1000	mouse	n/a	AB_390913	11814460001, Roche (Switzerland)
GFP	1:1000	rabbit	Full length denatured and non-denatured TurboGFP and CopGFP	n/a	AB514, Lot#51402250467, Evrogen
HA	1:500	mouse	YPYDVPDYA	n/a	ENZ-ABS118-0500, Enzo Life Sciences
HA	1:1000	rabbit	YPYDVPDYA	AB_2810986	ab137838, Abcam (UK)
$Na_v1.5$	1:1000	rabbit	DTVSRSSLEMSPLAPV	n/a	generated by Pineda (Germany)
α1-Na$^+$/K$^+$ ATPase	1:500	mouse	Full length native α1-Na$^+$/K$^+$ ATPase	AB_306023	ab7671, Abcam (UK)
α1-syntrophin	1:1000	rabbit	SGRRAPRTGLLELRAC	n/a	generated by Pineda (Germany)
α-actin	1:1000	rabbit	SGPSIVHRKCF	AB_476693	A2066, Sigma-Aldrich
IgG mouse	1:20,000	goat	Reacts with the heavy and light chains of mouse IgG$_1$, IgG$_{2a}$, IgG$_{2b}$, and IgG$_3$, and with the light chains of mouse IgM and IgA.	AB_10956588	926-68070, LI-COR Biosciences
IgG rabbit	1:20,000	goat	Reacts with the heavy and light chains of rabbit IgG, and with the light chains of rabbit IgM and IgA.	AB_621843	926-32211, LI-COR Biosciences

Table 3. Description of primary and secondary antibodies used in this study.

Patch clamp electrophysiology

To test data replicability, whole-cell sodium current recordings from two researchers (one of whom was blinded) using different setups were pooled together: either with an Axopatch 200B amplifier (Molecular Devices, Wokingham, United Kingdom) or with a VE-2 amplifier (Alembic Instrument, USA). Cells, solutions, micropipettes, and protocols were identical.

All cells were transiently co-transfected with *CD8*. Only successfully transfected cells, visualized by brief pre-wash with 1:1000 dilution of Dynabeads^{TM}CD8 (Thermo Fisher Scientific), were chosen for electrophysiological recordings. Thin-wall capillaries (TW150F-3, World Precision Instruments, Germany) were pulled with a DMZ-universal puller (Zeitz, Germany) with a tip resistance of 1.0 to 3.0 MΩ. The intracellular pipette solution contained (in mM): CsAsp 70, CsCl 60, EGTA 11, CaCl$_2$ 1, MgCl$_2$ 1, HEPES 10, and 5 Na$_2$ATP. The extracellular solution contained (in mM): NaCl 20, NMDG-Cl 110, CaCl$_2$ 2, MgCl$_2$ 1.2, HEPES 10, CsCl 5, D-glucose 5. pH was adjusted to 7.2-7.4 with CsOH or HCl. Osmolarity was measured by Osmometer type OM 806 (Löser) in mOsmol/kg H$_2$O range. All recordings were performed at a stable temperature of 25°C using Axon™ pCLAMP™ 10 Electrophysiology Data Acquisition & Analysis Software, Version 10.2 (Axon Instruments, CA, USA). The recorded sodium current I_{Na} was filtered with a low-pass Bessel 5 kHz filter at a sampling rate of 20 kHz per signal. Recordings without leak subtraction were selected for analysis. Raw traces were not corrected for the liquid junction potential.

Exclusion criteria for individual traces were: absence of I_{Na}; current at holding potential (-100 mV) <= -100 pA; slope of the activation (A) curve <=4; and slope of the steady-state inactivation (SSI) curve >=8. Sodium current density (pA/pF) was calculated by dividing peak current by cell capacitance. Current-voltage (I–V) curves were fitted with the equation $y = g(V_m - V_{rev})/((1 + \exp[(V_m - V_{1/2})/k]))$. SSI and A curves were fitted with the Boltzmann equation $y = 1/(1 + \exp[(V_m - V_{1/2})/k])$, where y is the normalized current or conductance at a given holding potential, V_m is the membrane potential, V_{rev} is the reversal potential, $V_{1/2}$ is the potential at which half of the channels are activated, and k is the slope factor. Data analysis was not blinded.

Protein-protein interaction assay

The NanoLuc® Binary Technology, NanoBiT® (Promega) allowed us to monitor protein-protein interactions between two $Na_v1.5$-WT subunits in live cells using an aqueous, cell-permeable furimazine substrate. For this purpose, *SCN5A*-WT was cloned into different vectors that encoded $Na_v1.5$-WT tagged with Large BiT (LgBiT) and Small BiT (SmBiT) on N- or C-termini when transfected into tsA201 cells. In the case of physical interaction between LgBiT and SmBiT, the bright luminescent signal was formed. Since Nano-Glo® Live Cell Assay (N2012, Promega) was not ratiometric and depended on the cell quantity, we normalized the luminescent signal to the fluorescent signal obtained with CellTiter-Fluor® Cell Viability assay (G6082, Promega). The signals were detected with Spark® multimode microplate (Tecan) and GloMax® Explorer Multimode Microplate Reader (GM3500, Promega).

Data and statistical analysis

Data are represented as means ± SEM. Data normality was tested by using Shapiro-Wilk test. Statistical significance for normally distributed data was calculated with ordinary one-way ANOVA. In case data were normalized to control condition statistical significance was calculated with one-sample two-tailed t-test with hypothetical mean value=1. Statistical significance for non-normally distributed data was calculated with Kruskal-Wallis test (Prism version 7.04; GraphPad, CA, USA). Exact *p*-values are indicated in the figures.

Supplemental material

Suppl. fig. S1-S2 show blots for co-immunoprecipitation between each of 14-3-3 isoforms and $Na_v1.5$ in tsA201 cell line. Suppl. fig. S3 shows endogenous expression of all 14-3-3 genes in tsA201 cell line. Suppl. fig. S4 shows validation of difopein as an efficient competitor for 14-3-3/ligand interaction in tsA201 cells: difopein, but not the mutant of difopein, interacts with 14-3-3 and stabilizes its oligomerization.

Results

Identification of 14-3-3 isoforms interacting with $Na_v1.5$

To identify which 14-3-3 isoform interacts with $Na_v1.5$, we performed co-immunoprecipitation analysis. Individual overexpression of all seven human 14-3-3 isoforms tagged with HA revealed that by themselves 14-3-3 proteins have a certain degree of non-specific binding with the anti-FLAG magnetic beads (Suppl. fig. S1-2).

Accordingly, the control condition with non-tagged Na$_v$1.5 was run along with FLAG-Na$_v$1.5 and compared for the intensity of co-immunoprecipitated 14-3-3 protein. As an indicator of successfully co-immunoprecipitated Na$_v$1-5-specific fraction, we used α1-syntrophin, a known partner of Na$_v$1.5 macromolecular complex and endogenously present in tsA201. From all the isoforms, only 14-3-3η exhibited reproducibly brighter band intensity in the FLAG-immunoprecipitated fraction compared to the non-tagged fraction; hence, we concluded that only this isoform reliably interacts with Na$_v$1.5 (Suppl. fig. S1-2). Furthermore, sepharose beads coupled with anti-Na$_v$1.5 antibody successfully co-immunoprecipitated α1-syntrophin and 14-3-3η, but not 14-3-3σ (Fig. 1). However, due to the overall weakness of Na$_v$1.5-immunoprecipitated 14-3-3η protein, we suggest that the interaction is rather weak and/or transient.

14-3-3 proteins do not affect expression level of Na$_v$1.5 at the cell surface

Next, we assessed the role of 14-3-3 proteins on the cell surface density of Na$_v$1.5. This was motivated by findings from Pohl *et al.*, who suggested that 14-3-3 proteins modulate cell surface expression of membrane receptors and ion channels upon binding with the E3 ubiquitin ligase Nedd4-2 (Pohl et al., 2021). Specifically, they showed that the high affinity of 14-3-3 with phosphorylated Ser342 and Ser448 sites of Nedd4-2 induces its structural rearrangement with plausible exposure of its tryptophan-rich WW-domain (Pohl et al., 2021). Our group previously reported the interaction between the WW-domain of Nedd4-2 with the PY-motif of Na$_v$1.5 and its importance for Na$_v$1.5 cell surface localization (Van Bemmelen et al., 2004; Rougier et al., 2005). According to our current results, the levels of the total and cell surface Na$_v$1.5 expression, represented by the biotinylated fraction, did not change upon overexpression of each individual 14-3-3 isoform when compared to the sodium channel alone (Fig. 2A-B).

Of note, in tsA201, we detected endogenous mRNAs of all 14-3-3 isoforms (Suppl. fig. S3). In line with our results, transcripts of all 14-3-3 isoforms were previously reported for HEK293, the cell line from which tsA201 was originally derived (Mathew et al., 2016). Therefore, to inhibit endogenous 14-3-3/ligand interactions, we used difopein, a dimer of R18, the unphosphorylated peptide antagonist of 14-3-3 (Masters and Fu, 2001). Difopein strongly interacts with two 14-3-3 proteins, stabilizing them in dimeric conformation, occupying their binding grooves, and preventing their interaction with other ligands (Masters and Fu, 2001). We performed co-immunoprecipitation analysis to validate that difopein is an effective competitor for 14-3-3/ligand interactions in our heterologous expression system. First, we confirmed that 14-3-3 strongly interacts with difopein but not with its mutant (Suppl. fig. S4a). As expected, difopein-YFP exhibited higher molecular weight than the mutant difopein-YFP, represented by the monomeric R18-D12K,E14K peptide (Suppl. fig. S4a). Further, we demonstrated that 14-3-3 dimers were stabilized by difopein but not by its mutant (Suppl. fig. S4b). Last, we confirmed that such an effect did not depend on any specific 14-3-3 isoform (Suppl. fig. S4b). Of note, difopein and its mutant did not affect the total and cell surface expression of Na$_v$1.5 (Fig. 2c-d).

Therefore, we concluded that 14-3-3 proteins do not regulate Na$_v$1.5 expression and cell surface density in the tested heterologous expression system.

Impact of 14-3-3 proteins onto Na$_v$1.5-mediated sodium current

Since of all 14-3-3 isoforms, only 14-3-3η weakly but reproducibly interacted with Na$_v$1.5 (Fig. 1, Suppl. fig. S1-S2), we assessed the whole cell recordings of the sodium current (I_{Na}) when transiently overexpressing *YWHAH*, encoding 14-3-3η. Moreover, to establish comparable conditions with the previous study that described a 14-3-3η-induced negative shift in I_{Na} inactivation curve, we similarly co-expressed *SCN1B*, encoding the Na$_v$β1 subunit of the voltage-gated sodium channel (Allouis et al., 2006). In our heterologous expression system, neither 14-3-3η or Na$_v$β1 alone nor their combination modified the baseline whole-cell properties of I_{Na}, as described by the current-voltage (*I-V*) relationship, peak current density, reversal potential (V_{rev}) and half-maximal potentials ($V_{1/2}$) of activation and inactivation curves (Fig. 3, Table 4). Similarly, disruption of endogenous 14-3-3/ligand interactions with difopein did not modify basal current properties of Na$_v$1.5 (Fig. 4, Table 5).

In summary, the isoform non-specific antagonist of 14-3-3, difopein, and 14-3-3η, the only isoform interacting with Na$_v$1.5, left the channel function unmodified.

	Peak I_{Na} density (pA/pF)	V_{rev} (mV)	Steady-state activation		Steady-state inactivation	
			k slope	$V_{1/2}$ (mV)	*k* slope	$V_{1/2}$ (mV)
+ CD8 + empty vector (n=32)	-140.18 ± 14.11	38.15 ± 1.68	6.38 ± 0.17	-27.01 ± 0.89	5.25 ± 0.12	-78.93 ± 0.90
+ CD8 + 14-3-3η (n=32)	-120.69 ± 19.06	35.62 ± 1.13	6.45 ± 0.18	-26.02 ± 1.38	5.22 ± 0.16	-78.98 ± 0.97
+ CD8 + hNa$_v$β1 + empty vector (n=31)	-129.51 ± 17.83	35.59 ± 1.35	6.49 ± 0.13	-27.61 ± 1.22	4.97 ± 0.14	-79.86 ± 1.10
+ CD8 + hNa$_v$β1 + 14-3-3η (n=28)	-143.15 ± 16.60	36.88 ± 1.57	6.39 ± 0.17	-25.05 ± 1.18	5.35 ± 0.17	-77.75 ± 0.90

Table 4. Biophysical properties of sodium currents conducted by Na$_v$1.5 stably present in tsA201 cells in combination with transiently expressed 14-3-3η and/or Na$_v$β1. CD8 was co-expressed to visualize successfully transfected cells. Peak sodium current (I_{Na}) densities are extracted from the respective I-V curves. Data are presented as mean ± SEM and are compared to control values (i.e., "+CD8 + empty vector"). n, number of cells analyzed.

	Peak I_{Na} density (pA/pF)	V_{rev} (mV)	Steady-state activation		Steady-state inactivation	
			k slope	$V_{1/2}$ (mV)	*k* slope	$V_{1/2}$ (mV)
+ CD8 + empty vector (n=31)	-125.89 ± 15.65	35.30 ± 1.66	6.40 ± 0.14	-26.14 ± 1.21	5.32 ± 0.17	-78.66 ± 1.28
+ CD8 + difopein (n=31)	-155.07 ± 17.48	39.89 ± 1.31	6.32 ± 0.17	-24.51 ± 0.72	5.30 ± 0.14	-75.80 ± 0.92
+ CD8 + mut. difopein (n=30)	-133.26 ± 18.63	37.40 ± 1.17	6.51 ± 0.13	-24.13 ± 0.84	5.54 ± 0.18	-77.79 ± 0.94

Table 5. Biophysical properties of sodium currents conducted by Na$_v$1.5 stably present in tsA201 cells in combination with transiently expressed difopein and its mutant. CD8 was co-expressed to visualize successfully transfected cells. Peak sodium current (I_{Na}) densities are extracted from the respective I-V curves. Data are presented as mean ±

SEM and are compared to control values (i.e., "+CD8 + empty vector"). n, number of cells analyzed.

Inhibition of 14-3-3 dimerization does not affect the interaction between α-subunits of Na$_v$1.5.

One role of the 14-3-3 family of adapter proteins is to stabilize multiprotein complexes (Obsil and Obsilova, 2011). In particular, 14-3-3η was proposed to mediate the coupled gating of Na$_v$1.5 dimers by bridging two α-subunits through their DI-DII loops (Clatot et al., 2017). We evaluated whether the 14-3-3 antagonist, difopein, affected Na$_v$1.5 dimerization. Our co-immunoprecipation analysis confirmed the existence of Na$_v$1.5-Na$_v$1.5 interactions in a heterologous expression system, but these were not affected by difopein overexpression (Fig. 5a-b). To confirm this finding in a more physiological and dynamic environment, we performed a NanoBiT assay. In brief, N- and C-termini of Na$_v$1.5 were tagged with LgBiT and SmBiT, respectively, which are parts of the bright and stable Nanoluciferase. Accordingly, when two Na$_v$1.5 physically interact, LgBiT and SmBiT would reconstitute Nanoluciferase, subsequently leading to luminescence emission (Fig. 5c). Even though intrinsic interaction between LgBiT and SmBiT is rather weak (K_d = 190 µM) (Dixon et al., 2016), all experiments were accompanied by a simultaneous co-expression of Na$_v$1.5-LgBiT with the non-interacting HaloTag-SmBiT as a readout of the background luminescence (Fig. 5c). In live tsA201 cells, the level of interaction between two Na$_v$1.5 was not modified by difopein (Fig. 5d-g). Interestingly, when comparing various locations of LgBiT and SmBiT tags on Na$_v$1.5, the brightest luminescent signal was detected when N-N and C-C termini were combined, compared to the N-C and C-N conditions (Fig. 5d-g). It might indicate a preferred conformational orientation of α-subunits within Na$_v$1.5 dimers.

Discussion

We reported that amongst seven 14-3-3 isoforms, only η (encoded by *YWHAH* gene) weakly co-immunoprecipitated with the major cardiac voltage-gated sodium channel Na$_v$1.5 when heterologously expressed in tsA201 cells. Nevertheless, neither 14-3-3η overexpression nor its inhibition with difopein affected the total protein level of Na$_v$1.5, its cell surface localization, dimerization, and basal biophysical properties of I_{Na}.

Our finding that 14-3-3η interacted with Na$_v$1.5 is in line with previous report of Allouis *et al.*, who demonstrated that 14-3-3 interacted with Na$_v$1.5 in Cos-7 and native mouse cardiac tissue and identified by yeast 2-hybrid screen DI-DII loop as a specific binding region of 14-3-3η (Allouis et al., 2006). Of note, when all cardiac voltage-gated sodium channels, including Na$_v$1.5, were immunoprecipitated from the mouse left ventricles using pan-Na$_v$ antibody, the most abundant peptides in the fraction were corresponding to 14-3-3ζ/δ, 14-3-3ε, 14-3-3γ, 14-3-3β/α, and 14-3-3σ, but not to 14-3-3η (Lorenzini et al., 2021).

Furthermore, we reported that difopein and 14-3-3η did not modify I_{Na} density and gating kinetics. Likewise, I_{Na} density was not modified by exogenous 14-3-3η, by inhibition of endogenous 14-3-3η, and by isoform-unspecific antagonist, difopein (Allouis et al., 2006; Utrilla et al., 2017; Zheng et al., 2021). However, regarding the activation and inactivation kinetics of Na$_v$1.5, several discrepancies were described. Similarly with our study, 14-3-3η overexpression did not modify conductance-voltage relationship (Allouis et al., 2006). But in contrast with our study, Zheng *et al.* demonstrated that difopein induced negative shift of the voltage-dependent conductance curve for the wild-type Na$_v$1.5 as well as for its truncated mutant Gly1642X with His558Arg polymorphism (Zheng et al., 2021). Furthermore, unlike our study, Allouis *et al.* showed that 14-3-3η shifted the inactivation curve to more negative potentials and decelerated the recovery from the inactivation of Na$_v$1.5 (Allouis et al., 2006). By co-expressing *SCN1B*, we eliminated the possibility that such discrepancy could be due to the presence of the Na$_v$β1 subunit. In fact, we did not observe any significant effects of *SCN1B* expression onto biophysical properties of the heterologous Na$_v$1.5. Indeed, previous studies reported conflicting results regarding the impact of Na$_v$β1 on the function of Na$_v$1.5. While some groups detected changes of the peak current density and/or differences in the channel kinetics and gating properties, others like us did not observe any Na$_v$β1-mediated changes of I_{Na} (Malhotra et al., 2001; Maroni et al., 2019; Martinez-Moreno et al., 2020; Valdivia et al., 2002; Zhu et al., 2021). Unlike the tsA201 cells used in our study, Allouis *et al.* used Cos-7 cells derived from monkey kidney fibroblasts as their heterologous expression system (Allouis et al., 2006). Like tsA201 cells, Cos-7 endogenously express all seven 14-3-3 isoforms, too (Abdrabou et al., 2020). However, their expression pattern, representing subcellular localization of 14-3-3 proteins, differed noticeably from human embryonic kidney (HEK293) cells (Abdrabou et al., 2020). Furthermore, other protein partners that regulate the function of Na$_v$1.5 may also vary between cellular models and may explain the disparity of our outcomes.

Our observation that 14-3-3 proteins do not participate in the dimerization of heterologously expressed Na$_v$1.5 was also corroborated by previous reports (Clatot et al., 2017; Utrilla et al., 2017). Both studies highlighted that instead of directly mediating the formation of Na$_v$1.5 homo- and heterocomplexes with other ion channels (*e.g.*, Na$_v$1.5-Na$_v$1.5 and Na$_v$1.5-K$_{ir2.1}$), 14-3-3η played a significant role in their biophysical coupling (Clatot et al., 2017; Utrilla et al., 2017).

Zheng *et al.* also suggested additional indirect roles for 14-3-3-dependent regulation of Na$_v$1.5 (Zheng et al., 2021). Indeed, since Na$_v$1.5 is regulated by many proteins, including kinases, of which many are 14-3-3 substrates, too (Abriel, 2010; Iqbal and Lemmens-Gruber, 2019), it is plausible that 14-3-3 proteins could alter the function of Na$_v$1.5 indirectly through other partner proteins (Fig. 6). Indeed, monomeric peptide R18, of which difopein is composed, disrupts 14-3-3γ/ε from protein kinase A (PKA), unmasking PKA's catalytic activity and promoting PKA-mediated phosphorylation of its downstream targets (Kent et al., 2010). Na$_v$1.5 is among PKA's targets, and PKA stimulation in HEK293 cells has been shown to enhance I_{Na} (Aiba et al., 2014). Furthermore, 14-3-3γ was shown to slow dephosphorylation of calcium/calmodulin-dependent protein kinase (CaMKII) (Psenakova et al., 2018). CaMKII is reported to

mediate both gain- and loss-of-function effects of I_{Na}, highlighting its complex relationship with Na$_v$1.5 (Iqbal and Lemmens-Gruber, 2019). Moreover, 14-3-3ζ has been shown to regulate protein kinase C (PKC), which, in turn, also modulates Na$_v$1.5 activity through phosphorylation (Gannon-Murakami and Murakami, 2002). Nevertheless, in our heterologous expression system the effect of isoform-unspecific antagonist, difopein, on the regulation of Na$_v$1.5 was not evident.

In summary, ubiquitous presence of homo- and hetero-oligomerizing 14-3-3 proteins makes investigation of their direct role onto Na$_v$1.5 rather challenging. Furthermore, expression of 14-3-3 proteins and their downstream targets might drastically vary between different models, potentially providing for distinct direct and indirect modes of Na$_v$1.5 regulation.

Limitations

To eliminate possible clonal specificity of our cell model, we developed polyclonal tsA201 stably expressing Na$_v$1.5. This led to high variability in whole-cell Na$_v$1.5 expression, which made single-cell electrophysiological analyses challenging, we strengthened the reliability of our data by testing a relatively large number of cells for each condition.

One of the limitations of this study is the presence of all endogenous 14-3-3 isoforms, including 14-3-3η, that could saturate Na$_v$1.5-binding sites and hence preclude interaction with the overexpressed proteins. In addition to homodimerization 14-3-3 isoforms are also capable to form heterodimers (Jones et al., 1995; Obsil and Obsilova, 2011). Thus, it is possible that even when overexpressed 14-3-3η-comprised heterodimers may interact with substrates other than Na$_v$1.5, hindering its direct role on the channel. Therefore, using gene-editing techniques to silence specific isoforms would be advantageous to unravel the complexity of 14-3-3 signaling.

Another limitation of heterologously expressed proteins is the difficulty in controlling post-translational modifications, such as phosphorylation, that impact Na$_v$1.5 function and 14-3-3 signaling. A recent study reported a successful semisynthetic approach to stabilize Na$_v$1.5 phosphorylation *in vitro* (Galleano et al., 2021). Thus, it would be interesting to test 14-3-3-specific regulation under stably phosphorylated Na$_v$1.5.

While 14-3-3 might not act directly onto Na$_v$1.5, it was shown to act onto its macromolecular complex with other protein partners (*e.g.*, PKA, PKC, CaMKII) and cardiac ion channels (*e.g.*, Kir2.1) (Pohl et al., 2021; Utrilla et al., 2017). Because of the vast spectrum of 14-3-3 targets, one should be aware of the possible indirect effects measured in any experiment targeting 14-3-3 function. As such, any results obtained with difopein should be interpreted carefully.

In summary, while direct effects of 14-3-3, specifically isoform η, on the cardiac sodium channel are not evident in the tested heterologous expression system, it still may regulate Na$_v$1.5 through proteins of its macromolecular complex.

Conclusions

We confirmed that 14-3-3η forms a macromolecular complex with human Na$_v$1.5 in a heterologous expression system. Difopein is a potent tool to antagonize 14-3-3-ligand interactions but not the direct physical interaction between two Na$_v$1.5 α-subunits. Overexpression of 14-3-3η and difopein did not affect I_{Na} density, gating kinetics as well as total and cell surface expression of Na$_v$1.5. Overall, the role of 14-3-3 proteins on the functionality of Na$_v$1.5 is dispensable in tsA201 cell line.

Author contributions

O.I, **D.R.K**., **J.S.R.**, and **H.A.** designed this research project. **O.I.**, **A.F.H.**, **E.R.**, **A.S.**, **D.R.K.**, **N.S.**, **M.E.**, **S.G.**, and **J.S.R.** performed the experiments. **O.I.**, **A.F.H.**, and **J.S.R.** analyzed the data. **O.I.** and **A.F.H.** made figures and tables. **O.I.** drafted and edited the manuscript. **O.I.**, **J.S.R.**, and **H.A.** critically reviewed the manuscript.

Disclosure statement

The authors declare no potential conflict of interest.

Funding

This work was funded by the Swiss National Science Foundation [SNF 310030_184783] to H.A.

Acknowledgments

We thank Dr Zoja Selimi (University of Bern) and Prof Jan Kucera (University of Bern) for their invaluable insights and fruitful discussions. We thank Dr Sarah Vermij for proofreading of the article.

References

Abdrabou, A., Brandwein, D., and Wang, Z. (2020). Differential subcellular distribution and translocation of seven 14-3-3 isoforms in response to EGF and during the cell cycle. Int. J. Mol. Sci. *21*, 1–24. https://doi.org/10.3390/ijms21010318.

Abriel, H. (2010). Cardiac sodium channel Nav1.5 and interacting proteins: Physiology and pathophysiology. J. Mol. Cell. Cardiol. *48*, 2–11. https://doi.org/10.1016/j.yjmcc.2009.08.025.

Aiba, T., Farinelli, F., Kostecki, G., Hesketh, G.G., Edwards, D., Biswas, S., Tung, L., and Tomaselli, G.F. (2014). A mutation causing brugada syndrome identifies a mechanism for altered autonomic and oxidant regulation of cardiac sodium currents. Circ. Cardiovasc. Genet. *7*, 249–256. https://doi.org/10.1161/CIRCGENETICS.113.000480.

Allouis, M., Le Bouffant, F., Wilders, R., Péroz, D., Schott, J.J., Noireaud, J., Le Marec, H., Mérot, J., Escande, D., and Baró, I. (2006). 14-3-3 Is a regulator of the cardiac voltage-gated sodium channel Nav1.5. Circ. Res. *98*, 1538–1546. https://doi.org/10.1161/01.RES.0000229244.97497.2c.

Van Bemmelen, M.X., Rougier, J.S., Gavillet, B., Apothéloz, F., Daidié, D., Tateyama, M., Rivolta, I., Thomas, M.A., Kass, R.S., Staub, O., et al. (2004). Cardiac voltage-gated sodium channel Nav1.5 is regulated by Nedd4-2 mediated ubiquitination. Circ. Res. *95*, 284–291. https://doi.org/10.1161/01.RES.0000136816.05109.89.

Clatot, J., Hoshi, M., Wan, X., Liu, H., Jain, A., Shinlapawittayatorn, K., Marionneau, C., Ficker, E., Ha, T., and Deschênes, I. (2017). Voltage-gated sodium channels assemble and gate as dimers. Nat. Commun. *8*, 1–14. https://doi.org/10.1038/s41467-017-02262-0.

Dixon, A.S., Schwinn, M.K., Hall, M.P., Zimmerman, K., Otto, P., Lubben, T.H., Butler, B.L., Binkowski, B.F., MacHleidt, T., Kirkland, T.A., et al. (2016). NanoLuc Complementation Reporter Optimized for Accurate Measurement of Protein Interactions in Cells. ACS Chem. Biol. *11*, 400–408. https://doi.org/10.1021/acschembio.5b00753.

Dong, C., Wang, Y., Ma, A., and Wang, T. (2020). Life Cycle of the Cardiac Voltage-Gated Sodium Channel NaV1.5. Front. Physiol. *11*, 1–11. https://doi.org/10.3389/fphys.2020.609733.

Galleano, I., Harms, H., Choudhury, K., Khoo, K., Delemotte, L., and Pless, S.A. (2021). Functional cross-talk between phosphorylation and disease-causing mutations in the cardiac sodium channel Nav1.5. Proc. Natl. Acad. Sci. U. S. A. *118*. https://doi.org/10.1073/pnas.2025320118.

Gannon-Murakami, L., and Murakami, K. (2002). Selective association of protein kinase C with 14-3-3 ζ in neuronally differentiated PC12 cells: Stimulatory and inhibitory effect of 14-3-3 ζ in vivo. J. Biol. Chem. *277*, 23116–23122. https://doi.org/10.1074/jbc.M201478200.

Gavillet, B., Rougier, J.S., Domenighetti, A.A., Behar, R., Boixel, C., Ruchat, P., Lehr, H.A., Pedrazzini, T., and Abriel, H. (2006). Cardiac sodium channel Nav1.5 is regulated by a multiprotein complex composed of syntrophins and dystrophin. Circ. Res. *99*, 407–414. https://doi.org/10.1161/01.RES.0000237466.13252.5e.

Gittenberger-de Groot, A.C., Hoppenbrouwers, T., Miquerol, L., Kosaka, Y., Poelmann, R.E., Wisse, L.J., Yost, H.J., Jongbloed, M.R.M., Deruiter, M.C., and Brunelli, L. (2016). 14-3-3Epsilon Controls Multiple Developmental Processes in the Mouse Heart. Dev. Dyn. *245*, 1107–1123. https://doi.org/10.1002/dvdy.24440.

Henriksson, M.L., Francis, M.S., Peden, A., Aili, M., Stefansson, K., Palmer, R., Aitken, A., and Hallberg, B. (2002). A nonphosphorylated 14-3-3 binding motif on exoenzyme S that is functional in vivo. Eur. J. Biochem. *269*, 4921–4929. https://doi.org/10.1046/j.1432-1033.2002.03191.x.

Iqbal, S.M., and Lemmens-Gruber, R. (2019). Phosphorylation of cardiac voltage-gated sodium channel: Potential players with multiple dimensions. Acta Physiol. *225*, 1–18. https://doi.org/10.1111/apha.13210.

Jones, D.H., Ley, S., and Aitken, A. (1995). Isoforms of 14-3-3 protein can form homo- and heterodimers in vivo and in vitro: implications for function as adapter proteins. FEBS Lett. *368*, 55–58. https://doi.org/10.1016/0014-5793(95)00598-4.

Kent, C.B., Shimada, T., Ferraro, G.B., Ritter, B., Yam, P.T., McPherson, P.S.,

Charron, F., Kennedy, T.E., and Fournier, A.E. (2010). 14-3-3 proteins regulate protein kinase A activity to modulate growth cone turning responses. J. Neurosci. *30*, 14059–14067. https://doi.org/10.1523/JNEUROSCI.3883-10.2010.

Kosaka, Y., Cieslik, K.A., Li, L., Lezin, G., Maguire, C.T., Saijoh, Y., Toyo-oka, K., Gambello, M.J., Vatta, M., Wynshaw-Boris, A., et al. (2012). 14-3-3ε Plays a Role in Cardiac Ventricular Compaction by Regulating the Cardiomyocyte Cell Cycle. Mol. Cell. Biol. *32*, 5089–5102. https://doi.org/10.1128/mcb.00829-12.

Lorenzini, M., Burel, S., Lesage, A., Wagner, E., Charriere, C., Chevillard, P.M., Evrard, B., Maloney, D., Ruff, K.M., Pappu, R. V., et al. (2021). Proteomic and functional mapping of cardiac NaV1.5 channel phosphorylation sites. J. Gen. Physiol. *153*. https://doi.org/10.1085/JGP.202012646.

Malhotra, J.D., Chen, C., Rivolta, I., Abriel, H., Malhotra, R., Mattei, L.N., Brosius, F.C., Kass, R.S., and Isom, L.L. (2001). Characterization of Sodium Channel α- and β-Subunits in Rat and Mouse Cardiac Myocytes. Circulation *103*, 1303–1310. https://doi.org/10.1161/01.CIR.103.9.1303.

Maroni, M., Körner, J., Schüttler, J., Winner, B., Lampert, A., and Eberhardt, E. (2019). β1 and β3 subunits amplify mechanosensitivity of the cardiac voltage-gated sodium channel Nav1.5. Pflugers Arch. Eur. J. Physiol. *471*, 1481–1492. https://doi.org/10.1007/s00424-019-02324-w.

Martens, M.D., Seshadri, N., Nguyen, L., Chapman, D., Henson, E.S., Xiang, B., Falk, L., Mendoza, A., Rattan, S., Field, J.T., et al. (2021). Misoprostol treatment prevents hypoxia-induced cardiac dysfunction through a 14-3-3 and PKA regulatory motif on Bnip3. Cell Death Dis. *12*. https://doi.org/10.1038/s41419-021-04402-3.

Martinez-Moreno, R., Selga, E., Riuró, H., Carreras, D., Parnes, M., Srinivasan, C., Wangler, M.F., Pérez, G.J., Scornik, F.S., and Brugada, R. (2020). An SCN1B Variant Affects Both Cardiac-Type (NaV1.5) and Brain-Type (NaV1.1) Sodium Currents and Contributes to Complex Concomitant Brain and Cardiac Disorders. Front. Cell Dev. Biol. *8*. https://doi.org/10.3389/fcell.2020.528742.

Masters, S.C., and Fu, H. (2001). 14-3-3 Proteins Mediate an Essential Anti-apoptotic Signal. J. Biol. Chem. *276*, 45193–45200. https://doi.org/10.1074/jbc.M105971200.

Mathew, D.E., Larsen, K., Janeczek, P., and Lewohl, J.M. (2016). Expression of 14-3-3 transcript isoforms in response to ethanol exposure and their regulation by miRNAs. Mol. Cell. Neurosci. *75*, 44–49. https://doi.org/10.1016/j.mcn.2016.06.006.

Menzel, J., Kownatzki-danger, D., Tokar, S., Ballone, A., Unthan-fechner, K., Kilisch, M., Lenz, C., Urlaub, H., Mori, M., Ottmann, C., et al. (2020). 14-3-3 binding creates a memory of kinase action by stabilizing the modified state of phospholamban. *1436*. .

Obsil, T., and Obsilova, V. (2011). Structural basis of 14-3-3 protein functions. Semin. Cell Dev. Biol. *22*, 663–672. https://doi.org/10.1016/j.semcdb.2011.09.001.

Pohl, P., Joshi, R., Petrvalska, O., Obsil, T., and Obsilova, V. (2021). 14-3-3-protein regulates Nedd4-2 by modulating interactions between HECT and WW domains. Commun. Biol. *4*. https://doi.org/10.1038/s42003-021-02419-0.

Psenakova, K., Petrvalska, O., Kylarova, S., Lentini Santo, D., Kalabova, D., Herman, P., Obsilova, V., and Obsil, T. (2018). 14-3-3 protein directly interacts with the kinase domain of calcium/calmodulin-dependent protein kinase kinase (CaMKK2). Biochim. Biophys. Acta - Gen. Subj. *1862*, 1612–1625. https://doi.org/10.1016/j.bbagen.2018.04.006.

Rougier, J.S., Van Bemmelen, M.X., Bruce, M.C., Jespersen, T., Gavillet, B., Apothéloz, F., Cordonier, S., Staub, O., Rotin, D., and Abriel, H. (2005). Molecular determinants of voltage-gated sodium channel regulation by the Nedd4/Nedd4-like proteins. Am. J. Physiol. - Cell Physiol. *288*, 692–701. https://doi.org/10.1152/ajpcell.00460.2004.

Thompson, W.C., and Goldspink, P.H. (2022). 14–3-3 Protein Regulation of Excitation–Contraction Coupling. Pflugers Arch. Eur. J. Physiol. 267–279. https://doi.org/10.1007/s00424-021-02635-x.

Utrilla, R.G., Nieto-Marín, P., Alfayate, S., Tinaquero, D., Matamoros, M., Pérez-Hernández, M., Sacristán, S., Ondo, L., de Andrés, R., Javier Díez-Guerra, F., et al. (2017). Kir2.1-Nav1.5 channel complexes are differently regulated than Kir2.1 and Nav1.5 channels alone. Front. Physiol. *8*, 1–16. https://doi.org/10.3389/fphys.2017.00903.

Valdivia, C.R., Nagatomo, T., and Makielski, J.C. (2002). Late Na Currents Affected by α Subunit Isoform and β1 Subunit Co-expression in HEK293 Cells. J. Mol. Cell. Cardiol. *34*, 1029–1039. https://doi.org/10.1006/jmcc.2002.2040.

Wang, B., Yang, H., Liu, Y.C., Jelinek, T., Zhang, L., Ruoslahti, E., and Fu, H. (1999). Isolation of high-affinity peptide antagonists of 14-3-3 proteins by phage display. Biochemistry *38*, 12499–12504. https://doi.org/10.1021/bi991353h.

Xing, H., Zhang, S., Weinheimer, C., Kovacs, A., and Muslin, A.J. (2000). 14-3-3 Proteins block apoptosis and differentially regulate MAPK cascades. EMBO J. *19*, 349–358. https://doi.org/10.1093/emboj/19.3.349.

Yaffe, M.B. (2002). How do 14-3-3 proteins work? - Gatekeeper phosphorylation and the molecular anvil hypothesis. FEBS Lett. *513*, 53–57. https://doi.org/10.1016/S0014-5793(01)03288-4.

Yang, X., Lee, W.H., Sobott, F., Papagrigoriou, E., Robinson, C. V., Grossmann, J.G., Sundström, M., Doyle, D.A., and Elkins, J.M. (2006). Structural basis for protein-protein interactions in the 14-3-3 protein family. Proc. Natl. Acad. Sci. U. S. A. *103*, 17237–17242. https://doi.org/10.1073/pnas.0605779103.

Zheng, Y., Wan, X., Yang, D., Ramirez-Navarro, A., Liu, H., Fu, J.D., and Deschênes, I. (2021). A Heart Failure-Associated SCN5A Splice Variant Leads to a Reduction in Sodium Current Through Coupled-Gating With the Wild-Type Channel. Front. Physiol. *12*. https://doi.org/10.3389/fphys.2021.661429.

Zhu, W., Wang, W., Angsutararux, P., Mellor, R.L., Isom, L.L., Nerbonne, J.M., and Silva, J.R. (2021). Modulation of the effects of Class-Ib antiarrhythmics on cardiac NaV1.5-encoded channels by accessory NaVβ subunits. JCI Insight *6*. https://doi.org/10.1172/jci.insight.143092.

Figure 1. Na$_v$1.5 co-immunoprecipitated with 14-3-3η. Immunoblot of total lysate and Na$_v$1.5-specific immunoprecipitated fraction of tsA201 cells 48 hours after transient co-expression of an empty vector, 14-3-3σ, 14-3-3η and Na$_v$1.5. Immunoprecipitation was performed with Sepharose beads pre-coupled with anti-Na$_v$1.5 antibody. HA-tagged 14-3-3σ and 14-3-3η were revealed with an anti-HA antibody. Endogenous α1-syntrophin was used as a positive control for co-immunoprecipitation with Na$_v$1.5, and α-actin as a negative control. Yellow triangle indicates on the band corresponding to the co-immunoprecipitated 14-3-3η.

Figure 2. 14-3-3 proteins do not affect the total expression level of Na$_v$1.5 and cell surface density. Representative immunoblots of the total lysate and cell surface fraction of tsA201 stably expressing Na$_v$1.5 after 48 hours of transient overexpression of: **(a)** HA-tagged 14-3-3 proteins; **(c)** YFP-tagged difopein and its mutant.

Endogenous Na$^+$/K$^+$-ATPase was used as a positive control of cell surface fraction, while α-actin was used as a negative control. **(b** and **d)** Intensities of Na$_v$1.5 in total protein lysate and at the cell surface were normalized to Na$^+$/K$^+$-ATPase and α-actin and to the control condition ("+ empty vector"). Data are presented as mean ± SEM from three-five biological replicates. Individual *p*-values, calculated with one-sample two-tailed t-test with hypothetical mean value=1, are indicated in each panel.

Figure 3. 14-3-3η expressed alone or with Na$_v$β1 does not affect Na$_v$1.5-conducted sodium current. (a) Current density-voltage (IV) relationships from tsA201-Na$_v$1.5 transiently expressing CD8 with empty vector, or 14-3-3η, and/or

Na$_v$β1. **(b)** Representative whole-cell I_{Na} traces recorded from the listed conditions. **(c)** Peak current densities of each group. **(d)** Activation (A) and steady-state inactivation (SSI) curves. **(e)** Reversal potential (V_{rev}). **(f)** Half-voltage of SSI ($V_{1/2}$ SSI). **(g)** Half-voltage of A ($V_{1/2}$ A). Values pertaining to the biophysical properties are shown in Table 4. Data are presented as mean ± SEM. *n*, number of cells taken for analysis. Kruskal-Wallis has been performed in **(c, e, f,** and **g)** and the multiplicity adjusted *p*-values are indicated in each panel.

Figure 4. Difopein, an antagonist of 14-3-3/ligand interactions, does not affect Na$_v$1.5-conducted sodium current. (a) Current density-voltage (IV) relationships from Na$_v$1.5-expressing tsA201 cells transiently expressing CD8 with empty vector, difopein, or mutant of difopein. **(b)** Representative whole-cell I_{Na} traces recorded from the listed conditions. **(c)** Peak current densities of each group. **(d)** Activation (A) and

steady-state inactivation (SSI) curves. **(e)** Reversal potential (V_{rev}). **(f)** Half-voltage of SSI ($V_{1/2}$ SSI). **(g)** Half-voltage of A ($V_{1/2}$ A). Values pertaining to the biophysical properties are shown in Table 5. Data are presented as mean ± SEM. *n*, number of cells taken for analysis. Kruskal-Wallis has been performed in **(c, e, f,** and **g)** and the multiplicity adjusted *p*-values are indicated in each panel.

Figure 5. Inhibition of 14-3-3/ligand interaction does not affect oligomerization between α-subunits of Na$_v$1.5. Difopein and its mutant did not affect co-immunoprecipitation between α subunits of Na$_v$1.5. **(a)** Representative immunoblot of the total lysate ("INPUT") and FLAG-specific immunoprecipitated fraction ("IP: FLAG") of tsA201 cells transiently expressing Na$_v$1.5 with empty vector, difopein, or mutant of

difopein. Overexpressed YFP-tagged difopein and its mutant were revealed with an anti-GFP antibody. Endogenous α-actin was used as a negative control for co-immunoprecipitation with Na$_v$1.5. **(b)** Intensity of co-immunoprecipitated Na$_v$1.5-HA was normalized to the intensity of the total Na$_v$1.5-HA divided by the intensity of α-actin. Data are presented as mean ± SEM from three biological replicates and are normalized to the control condition ("+ empty vector"). Individual *p*-values, calculated with one-sample two-tailed t-test with hypothetical mean value=1, are indicated in the panel. **(c)** Schematic illustration of NanoBiT assay. Background signal was determined by co-expression of N- or C-termini LgBiT-Na$_v$1.5 with non-interacting control, SmBiT-HaloTag. **(d, e, f,** and **g)** In living cells difopein did not modify the level of Na$_v$1.5 dimers. Results of the NanoBiT assay are presented as relative intensity of luminescence (indicating the level of protein-protein interactions) normalized to fluorescence (indicating the number of cells) in tsA201 cells transiently transfected with *SCN5A* constructs. Each dataset was normalized to its relevant background. Different variations between Na$_v$1.5 N- and C-termini interactions were tested. Since difopein and its mutant were tagged with YFP, we also included control by co-transfection with GFP. NT, non-transfected. Data are presented as mean ± SEM from three biological replicates. The multiplicity adjusted *p*-values, calculated with one-way ANOVA, are indicated in each panel.