Survey of the Phytophthora species diversity reveals *P. europaea*-like isolates as

Phytophthora root rot pathogens in Québec Christmas tree plantations

Guillaume Charron[1,2], Julie Yergeau[1,3], Hervé Van der Heyden[2], Guillaume J. Bilodeau[4],

Carole Beaulieu[3], Philippe Tanguay[1,*]

[1]Natural Resources Canada, Laurentian Forestry Centre, 1055 rue Du P.E.P.S., P.O. Box
10380, Québec (Québec), Canada, G1V 4C7

[2]Phytodata Inc. research company, 291 rue de la Coopérative, Sherrington (Québec),
Canada J0L 2N0.

[3]Département de Biologie, 2500, boulevard de l'Université, Université de Sherbrooke,
Sherbrooke (Québec), Canada, J1K 2R1

[4]Canadian Food Inspection Agency, 3851 Fallowfield Road, Ottawa, ON, Canada, K2J
4S1

[*]Corresponding author: P. Tanguay; E-mail: philippe.tanguay@NRCan-RNCan.gc.ca

Funding: This study was supported by a grant in the Cellule d'innovation des

méthodologies de diagnostic des ennemis de cultures from the Prime-Vert funding

program of the Québec Ministry of Agriculture, Fisheries and Food.

Abstract

Christmas trees are an economically and culturally important ornamental plant in North America. Many microorganisms are pathogens of firs cultivated as Christmas trees. Among those, *Phytophthora* causes millions of dollars in damage to plantations every year. In Canada, it is still not known which species or group of species is responsible for Phytophthora root rot (PRR) of the cultivated *Abies* species. Between 2019 and 2021, soil and root samples were collected from 40 Christmas tree plantations in Québec province. We used soil baiting and direct root isolation to assess the diversity of culturable *Phytophthora* spp. The obtained isolates were identified with a multi-locus sequencing approach and we used the sequencing data to place them along the *Phytophthora* phylogeny. A total of 44 isolates from 6 different *Phytophthora* species were identified, one of which was fitting the undescribed *P. sp kelmania*. There was also a group of 10 isolates which could not be assigned to a known Phytophthora species based on DNA sequences. Among the known species, a *Phytophthora europaea*-like was the prevalent isolated species associated with PRR symptomatic trees. The fulfillment of Koch's postulates confirmed the pathogenicity of *P. europaea*-like on both Fraser and balsam fir seedlings. Our studies provide a first snapshot of the *Phytophthora* diversity in Québec's Christmas tree productions and describe multiple first associations between *Phytophthora* species and *Abies balsamea* and *A. fraseri*.

Keywords: Christmas trees, *Abies fraseri*, *Abies balsamea*, *Phytophthora* root rot, ITS1-5.8S-ITS2, *Phytophthora europaea*, *Phytophthora abietivora*, Phytophthora diversity, Phytophthora isolation, Koch's posulates.

Introduction

With a little less than 1000 km^2 in cultivated surface, Christmas trees are an

economically important ornamental plant for the USA and Canada (Statistics Canada

2021; USDA National Agricultural Statistics Service 2021). In eastern North America,

Fraser fir is the preferred species cultivated as Christmas trees. Reasons for this

preference include shape and growth, combined with its postharvest qualities such as

needle retention and durability (Bates et al. 2004; Mitcham-Butler et al. 1988).

Unfortunately, Fraser fir is also one of the most susceptible species to *Phytophthora* root-

rot (PRR) (Benson and Grand 2000; Chastagner and Benson 2000). Hence, PRR is

responsible for millions of yearly losses (Grand and Lapp 1974; Kuhlman and Hendrix

1963; Pettersson et al. 2017). The first symptom of PRR is a reddish-brown discoloration

of the cambium region of the roots, which is difficult to detect unless roots are unearthed.

As the infection progresses through the root system, symptoms such as chlorosis of

foliage, wilting of new growth, branch flagging and stunting begin to show in the above-

ground parts of the tree (Chastagner and Benson 2000). Unfortunately, it is usually too

late to take actions at the onset of visible symptoms and infected trees will die within

months.

In the last 40 years, most of the data concerning *Phytophthora* species associated

with Christmas tree productions came from the US (Benson and Grand 2000; Chastagner

et al. 1990; Grand and Lapp 1974; McKeever and Chastagner 2016). More than 20

Phytophthora species have been associated with more than 20 different species of true

firs (Farr 2022), with at least 12 Phytophthora species known to induce PRR (McKeever

and Chastagner 2016). Since the mid-2000s, data available for European countries

overlap with what is found in the US (Pettersson et al. 2019; Shafizadeh and Kavanagh

2005; Talgø et al. 2006; Talgø et al. 2007; Talgø et al. 2017), suggesting that either the

association between these Phytophthora species and firs have been longstanding or that,

as for other *Phytophthora* species, worldwide plant trade has contributed to their global

dissemination (Bienapfl and Balci 2014; Brasier 2013; Jung et al. 2016). While the

information currently available connects *Phytophthora* species to PRR symptomatic

trees, few studies have shown their pathogenic potential with Koch's postulates

(Frampton and Benson 2012; Hamm and Hansen 1982; Hoover and Bates 2013;

McKeever and Chastagner 2016). Moreover, when Koch postulate are tested, the

experiments are usually performed by injuring the tree and directly putting the pathogen

at the wounded site (Li et al. 2019; McKeever and Chastagner 2016; Pettersson et al.

2019; Quesada-Ocampo et al. 2009), which does not capture all infection patterns

(Hardham 2001; Judelson and Blanco 2005).

The province of Québec represents more than 35% of the total Canadian cultivated

area used for Christmas tree plantations. Eighty-five percent of this area can be found in

two regions; Estrie and Chaudière-Appalaches (Statistics Canada 2016). The two most

cultivated species used as Christmas trees in Québec's province are Fraser fir (*Abies*

fraseri) and balsam fir (*Abies balsamea*). While Canadian producers had past PRR

occurrences in their plantations, the data available on Phytophthora species diversity is

insufficient to tackle tasks such as systematic detection and prevention of disease

spreading for this host.

In this study, we sampled Christmas tree plantations in Estrie and Chaudière-

Appalaches to get an overview of the *Phytophthora* species diversity associated with PRR

symptomatic trees in Quebec main producing regions. We also demonstrated the

pathogenic potential of *P. europaea*-like isolates obtained from sampling and found in

association with both PRR symptomatic *A. balsamea* and *A. fraseri*. Finally, this study

constitutes a first documented report for *P. europaea* in Canada.

Material and Methods

Christmas tree plantations sampling. Soils were sampled during the fall of 2019,

2020, and the spring of 2021 from 40 different sites in Chaudière-Appalaches and Estrie

(20 sites each, Supplementary Table S1). All of the sites were harboring trees showing

Phytophthora root rot symptoms. Twelve soil samples were taken in each site from trees

aged 4-12 years: four showing signs of PRR and four healthy in appearance, and four

from apparently healthy trees from a natural forest near the production site. Using a 5 cm

diameter soil probe, three soil sub-samples, 120° apart, were taken at the crown limit of

the trees up to 17 cm in depth. After removing the organic horizon, the three soil cores

were cut in half and the three halves were pooled together in a plastic bag (38x19 cm

Whirl-Pak). The probe was rinsed with water and heat-disinfected with a propane hand

torch between samples. Samples were kept at 4°C until being processed.

Soil samples processing and baiting. Each soil sample was mixed and sieved

through a 4 mm mesh basket. Root fragments retained in the sieve were transferred at

4°C for oomycetes isolation. Two hundred and fifty milligrams of homogenized soil was

transferred to a microtube for subsequent DNA extraction. Two hundred and fifty milliliters

of homogenized soil were put into a clean 1 L plastic container (IPL plastics, Canada) to

which 500 mL of distilled water was added. Four to five healthy rhododendron

'Cunningham's White' leaves without any visible injury were rinsed with sterile distilled

water and laid upon the water surface of each soil container. Five rhododendron leaves

were also placed in a container filled with 500 mL of distilled water to serve as a control.

The containers were incubated in the dark at 20°C until the apparition of necrosis on the

rhododendron leaf tissue (4-7 days).

Following the emergence of necrosis, leaves were washed with running tap water,

dried for 10 minutes on absorbing paper, and taken under a laminar flow hood. Necrotic

tissues were sampled using a paper punch sterilized with ethanol and flame to obtain leaf

disks (6 mm in diameter). Ten disks were taken at the margin of leaf necrotic tissues for

each soil sample: five were frozen in 2 mL safe-lock tubes (Eppendorf, Hamburg,

Germany) for subsequent DNA extraction and the other five were inserted into solid V8-

PARPNH (clarified V8 50 mL/L, agar 15 g/L, PCNB 25 µg/mL, Ampicillin 250 µg/mL,

Rifampicin 10 µg/mL, Pimaricin 10 µg/mL, Nystatin 25 µg/mL, and Hymexazol 25 µg/mL)

medium (100 mm × 15 mm Petri dishes). Leaf disks were inserted at a 45° angle keeping

half of the disc above the agar. Petri dishes were incubated in the dark at 20°C and were

examined daily for growth. After 2 to 10 days, morphologically distinct colonies were

purified by transferring hyphal tips to fresh V8-PARPH. The cultures were incubated in

the dark at 20°C. All cultures not resembling a *Phytophthora* colony were excluded from

further investigation. Grown cultures were preserved at 12°C.

Oomycetes isolation from roots. Lateral roots gathered from soil samples were

transferred to a fine mesh strainer that was first immersed in a beaker containing distilled

water and agitated to remove most of the soil. The strainer was then transferred to a

beaker containing a 0.5% sodium hypochlorite solution and soaked for 30 seconds,

followed by two rinses in distilled water. Surface-sterilized root segments were dried into

a laminar flow cabinet for 5-10 minutes and cut into 5-10 mm sections with a sterile

scalpel. Five root sections were inserted halfway into a V8-PARPNH agar plate. The

remaining roots were put into a 2 mL safe-lock tube and stored at −20°C. Media were

incubated in the dark at 20°C. Morphotype colonies were purified as described above.

DNA extractions. Purified cultures were grown on V8-agar (1.5% agar, clarified

V8 50 mL/L) overlayed with a sterile cellophane membrane. Mycelium was scraped with

a scalpel blade and transferred to a 2 mL safelock tube (Eppendorf, Hamburg, Germany).

A tungsten carbide bead (Qiagen, Hilden, Germany) was added to each tube and the

bottom of the tubes was immersed in liquid nitrogen for 1 minute. The tubes were then

quickly transferred to a Mixer mill adapter (Retsch, Haan, Germany) previously cooled at

−80°C. Tubes were shaken for 1 minute at a frequency of 26 Hz after which a second

cycle of freezing and shaking was performed. DNA was extracted following the protocol

developed in Penouilh-Suzette et al. (2020) with minor modifications. In short, 800µL of

Lysis buffer (0.1 M Tris-HCl pH 8, 10 mM EDTA pH 8, 1 M NaCl and 1% SDS) were

added to up to 50 mg of ground mycelium along with 2 µL of Proteinase K (20 mg/mL).

Samples were mixed by inversion and incubated at 50℃ for 10 minutes, while mixing by

inversion twice during the incubation. 270 µL of potassium acetate 5 M was added, and

tubes were mixed by inversion before incubating on ice for 10 minutes. Samples were

centrifuged for 10 minutes (5,000 g), the supernatant was transferred to a new 2 mL tube,

and 2 µL of RNaseA (100 mg/mL) was added. The samples were incubated for 30 minutes

at 37℃ and mixed by inversion every 10 minutes. Next, 1 mL of Isopropanol was added

and tubes were mixed by inversion. The samples were incubated at RT for 30 minutes.

Following incubation, samples were centrifuged for 2 minutes at 10,000 g, the

supernatant was removed, and the DNA pellets were rinsed with fresh 70% ethanol.

Samples were centrifuged for 1 minute (10,000 g) and the supernatant was removed.

Pellets were air dried for 1 to 5 minutes under a lamp and were resuspended with 50 µL

PCR grade water.

Molecular identification. To identify the isolated organisms we amplified five loci:

the ITS1-5.8S-ITS2 region of the 45S ribosomal RNA (ITS), the Beta tubulin, Heat Shock

Protein 90 (HSP90), and the mitochondrial genes Cytochrome oxidase I (cox1) and

NADH dehydrogenase (nad1). We used primer pairs already described in the literature

(Supplementary Table S2). The PCR cycles were: 94℃ 2 min; 35 cycles of 94℃ 30s;

Primer specific Tm 30s; 72℃ with an amplicon specific time; 72℃ 10 min. First, PCR

products of expected length for the ITS (Supplementary Table S2) were sent to Sanger

sequencing at the CHUL sequencing platform on an ABI 3730xl Data Analyzer (Applied

Biosystems, Waltham, USA). Then, for every strain confirmed as *Phytophthora*, the other

four loci were amplified and sent for sequencing. For each of the isolated strains, the

sequences were used as a query against the nt/nr collection of the NCBI using BLAST

(Altschul et al. 1990). All of the sequences generated for this study were deposited into

Genbank under accession numbers ON008003-ON008178 (for cox1, nad1, Beta-tubulin

and HSP90), OM984685-OM984728 (ITS) and OP676073 (Cox1 sequence for the

RC2017-1 *P. abietivora* isolate).

Phylogenetic analysis. For each of the sequenced locus, sequences identical for

more than one isolate were collapsed into one sequence. Sequences of the 5 loci from

strains of 69 of *Phytophthora* species belonging to the different kown clades were

gathered from GenBank and used as references (Supplementary Table S3). Sequences

obtained from our isolates and GenBank-downloaded sequences of the reference strains

were aligned with Clustal W and trimmed using the MEGA X software (Kumar et al. 2018).

These alignments were concatenated using a custom script in R v4.1.1 with the *seqinr*

v1.0-2 package (Charif and Lobry 2007; R Core Team 2013). Phylogenetic analysis

employed maximum-likelihood and the Tamura-Nei model (Tamura and Nei 1993) in

MEGA X. The dataset comprised 80 concatenated sequences, 69 from the reference

strains of the *Phytophthora* species and 11 distinct concatenated sequences covering the

44 isolates of *Phytophthora* recovered in this study. Positions with less than 90% site

coverage were eliminated resulting in a total of 3372 positions in the final dataset.

Elongisporangium undulatum was designated as an outgroup. Node confidence was

estimated with 1000 bootstrap replicates.

Inoculum and seedlings preparation. Three *Phytophthora europaea*-like

isolates obtained from the sampling were selected to test Koch's postulates. Isolates were

inoculated on a V8-agar medium and incubated for one week at 20℃. Ten mycelium plugs

(5 mm $\times$ 5 mm) were taken at the margin of the actively growing cultures and inoculated

into a sterile mixture of vermiculite, overnight water-soaked steel-cut oat, and clarified V8

vegetable juice in a 1 : 1 : 1 ratio. Controls were prepared the same way using plugs from

an uninoculated V8-agar medium. These inocula were incubated for two weeks at 20°C

in the dark.

Dormant two-year-old *Abies balsamea* and *Abies fraseri* seedlings were obtained

from a local nursery (Productions Resinex inc., La Durantaye, Canada) and kept in

dormancy at −2℃ for two months. They were prepared for the experiment by transferring

them at 4°C for a week, followed by a week at 10°C. Fraser and balsam fir seedlings were

respectively re-potted in individual 1.5L and 1L containers filled with an autoclaved

soilless mix containing Pro-mix LP15 peat moss (Premier Tech, Rivière-du-Loup,

Canada), vermiculite (Perlite Canada Inc., Lachine, Canada) and silica sand (Atlantic

silica inc., Poodiac, Canada) in a 2:1:1 ratio amended with 4 kg/m^3 of 18-6-8 NPK

Nutricote™ fertilizer (Chisso-Asahi Fertilizer Co., Tokyo, Japan). Seedlings were

maintained in a greenhouse with a 16-hour photoperiod under high-pressure sodium

lighting (OSRAM Sylvania Inc., Wilmington, USA), with temperatures of 20°C during the

day and 18°C during the night. Seedlings were irrigated for 10 minutes periods, four times

a week (≈400 mL per irrigation period). These conditions were maintained until bud burst.

Koch's postulates. After bud burst, 100 mL of the top layer of the potting medium

was replaced with a fresh potting mix inoculated with 0.1% (V/V) of the *P. europaea*

inoculum. Trials followed a randomized block design where three blocks containing five

replicates of each inoculum/tree essence combinations were placed in different areas of

the greenhouse (three blocks ✕ two essence ✕ (three isolates + one control) ✕ five

replicates = 120 seedlings). Growth conditions were the same as stated above except

that every two weeks, the seedlings were flooded (1 cm of water above the potting

medium) for a period of 48 hours to create conditions favorable to zoospores production

and dissemination. The height (soil surface to the tip of the seedling rounded to the closest

5 mm) and the collar diameter of the seedlings were recorded right before inoculation

(time point "0"), every other week through the first month (time points 2 and 4), and every

week between week five and twelve (time points 5-12). Symptoms of PRR were assessed

by visual examination of the seedlings looking for stem and branch wilting or chlorotic and

dead needles. An estimated percentage of the seedling tissues affected by these

symptoms was recorded for each seedling at each time point. Seedlings were considered

dead once symptoms affected 100% of their tissues.

After 12 weeks, all seedlings were carefully removed from their containers and the

root systems were sampled by separating them from the stem at the height of the root

collar. The root systems were washed to remove potting media, taking care to retain as

much root tissue as possible. The clean root systems were successively plunged into a

beaker containing clean distilled water for 1 minute, then surface sterilized into a beaker

containing a 0.5% sodium hypochlorite solution for 30 seconds. Finally, the roots were

rinsed twice in clean distilled water for 30 seconds. Surface-sterilized root systems were

dried for 10-15 minutes in a laminar flow cabinet and then 5 to 10 roots were sampled

and cut into 5 to 10-mm segments. Two aliquots of 10 root segments were taken for each

seedling. The remainder of the surface sterilized root systems were oven-dried for 48

hours at 70℃. The dried root systems were weighed to assess the severity of root

degradation. To satisfy the fourth postulate, we used the two aliquots of root segments.

One was used for oomycete isolation as previously described. The other was used for

DNA extraction using the DNeasy Powersoil Pro Kit (QIAGEN, Hilden, Germany) and

molecular detection of Phytophthora using the nad1 PCR also as previously described.

Statistical analyses. All statistical analyses were performed with the R software

v4.1.1 (R Core Team 2013) using custom scripts and the FSA package v0.9.3 for the

Dunn's Test function (Ogle 2017). Unless stated otherwise, the significance level used for

all statistical tests is $\alpha = 0.05$. The area under the disease progression curves (AUDPC)

was calculated as described (Madden et al. 2007). Data representations, survival

analyses and statistical comparisons were made using the beanplot v1.3.1 (Kampstra

2008), beeswarm v0.4.0 (Eklund 2021), survival v3.2-11 (Therneau and Grambsch 2000)

and survminer v0.4.9 (Kassambra 2021) packages.

Results

Phytophthora isolations. In 2019, 41 *Phytophthora* isolates were recovered

from sampling, while the 2020 sampling yielded three isolates (Supplementary Table

S4). Among them, 40 were recovered by baiting soil samples with Rhododendron

leaves. These isolates were obtained from 22 unique soil samples with some samples

yielding up to four isolates. DNA sequencing of the Beta-tubulin, ITS, HSP90, cox1 and

nad1 genes coupled with BLAST analyses revealed that the isolates recovered belong

to at least seven different species. Based on the ITS sequences, we obtained 18 *P.*

chlamydospora isolates, 8 *P. europaea*-like isolates, 7 *P. gonapodyides* isolates, 3 *P.*

gregata isolates, 6 *P. megasperma* isolates, and 2 *P. sp. Kelmania* isolates. Among the

P. chlamydospora isolates, 10 isolates had BLAST hit results for the sequence of the

other loci that did not correspond to *P. chlamydospora*. Excepted for 6 isolates of *P.*

gonapodyides and 2 isolates of *P. chlamydospora* that were obtained from soil samples

taken in natural fir stands in the vicinity of christmas tree plantations, all other isolates

were from soil samples under firs showing PRR symptoms. Despite the fact that 160

soil samples from healthy trees in plantations were analysed, no *Phytophthora* was

recovered from these samples. The fact that few isolates were obtained from healthy

trees coupled with the low numbers of *Phytophthora* isolates recovered from the 2019,

2020 and 2021 samplings precluded statistical analysis. However, we can observe that

P. europaea-like isolates were recovered from the highest number of independent

samples (8 isolates from independent samples) which could imply a role for this

organism in the development of PRR.

Phylogenetics analysis. Our phylogenetic analysis revealed that the isolates

obtained from Christmas tree plantations primarily come from clades 6 and 7 (Figure 1).

Most of the isolates obtained from our survey are related to known *Phytophthora* species,

except for a group of 10 isolates (from 7 independent soil samples) that seem to be related

among themselves but not to any other species used as reference. For those sequences,

BLAST analyses would not return consistent identity results. Our phylogeny placed those

isolates in clade 6b as a monophyletic taxa sister to the *P. chlamydospora* and *P.*

gonapodyides group.

While we identified *P. europaea*-like isolates as potential pathogen candidates, our

phylogeny indicates that these isolates would be more closely related to *P. abietivora*.

From our concatenated sequence data (3884 bp), we identified 19 SNPs between our *P.*

europaea reference and the *P. abietivora* type strain. Half of these SNP (9) were located

in the coxI mitochondrial locus. From those 19 SNPs, 17 were shared between our *P.*

europaea-like isolates and *P. abietivora*, only one was shared between our *P. europaea*-

like isolates and *P. europaea* (in the ITS locus) and one seemed to be private to our *P.*

europaea-like isolates while *P. europaea* and *P. abietivora* shared the same nucleotide

(in the coxI locus).

Progression of PRR and survival in seedlings. Visual examination of the

seedlings allowed the observation of a rapid onset of PRR symptoms after the second

flooding period (four weeks post-inoculation, Figure 2a, b). The statistical comparison of

AUDPC values indicates a difference between treatments (Kruskal-Wallis test, $H(7) =$

118.53, $P < 0.001$). Post-hoc tests showed that, in Fraser firs, the AUDPC of inoculated

seedlings was significantly different from the uninoculated controls for all the tested

isolates (Two-sided Dunn's test $P < 0.001$, Benjamini-Hochberg corrected). In balsam firs,

the results show that two out of three seedling groups are significantly different from the

uninoculated controls (Two-sided Dunn's test P: 170_B = 0.049, 62_E < 0.001, and 97_A

= 0.301, Benjamini-Hochberg corrected). The comparison between inoculated Fraser and

balsam seedlings also shows a significant difference in AUDPC (Wilcoxon rank sum test,

W=595, $P < 0.001$).

First seedling deaths were recorded as soon as five weeks post-inoculation. While

uninoculated controls showed only little losses (Fraser 13.333%, 2 replicates out of 15

and balsam 6.666%, 1 replicate out of 15), the inoculated seedlings showed a large

proportion of dead seedlings (Fraser 97.7%, 44 replicates out of 45 and balsam 60%, 27

replicates out of 45) (Figure 2c). Accordingly, there is a significant difference in survival

probability between treatments (log-rank test, $\chi^2(7) = 95$, $P < 0.001$). Pairwise

comparisons show that all isolates inoculated onto Fraser fir seedlings lead to a

significantly different survival probability compared to the uninoculated control (Figure 2c,

log-rank test $P < 0.001$ for all isolates, Benjamini-Hochberg corrected). In balsam fir

survival probability was significantly different in two isolates when compared to the

uninoculated control (Figure 2c, log-rank test P: 170_B = 0.012, 62_E < 0.001, and 97_A

= 0.054 Benjamini-Hochberg corrected). The comparison between inoculation Fraser and

balsam seedlings shows a significant difference in survival probability. (log-rank test, $\chi^2(1)$

= 37.3, $P < 0.001$).

Impacts of *P. europaea*-like isolates on fir seedlings growth. We used the

height, stem diameter and root system mass collected during the experiment to assess

the effect of the inoculation with *P. europaea*-like isolates on the growth of balsam and

Fraser firs seedlings.

We calculated height increment as the difference in height between the final (t_{12})

and initial (t_0) time points measurements. Statistical comparisons indicate significant

differences in height growth between treatments (Kruskal-Wallis test, $H(7) = 67.182$, $P <$

0.001). Post-hoc tests show that inoculated Fraser firs had height increment significantly

different than the uninoculated trees (Figure 3a, Two-sided Dunn's test P: 170_B = 0.040,

62_E = 0.013, and 97_A = 0.004, Benjamini-Hochberg corrected) while such observation

could only be made with isolate 62_E in balsam firs (Figure 3b, Two-sided Dunn's test P:

170_B = 0.310, 62_E = 0.04, and 97_A = 0.989, Benjamini-Hochberg corrected).

Root collar diameter increment of the seedling was calculated the same way as

height increment by subtracting the final and initial time points measurements. Statistical

comparisons indicate significant differences in root collar diameter increment between

treatments (Kruskal-Wallis test, $H(7) = 60.946$, $P < 0.001$). Post-hoc tests show that

inoculated Fraser firs had significantly lower diameter increment than the uninoculated

trees (Figure 3c, Two-sided Dunn's test P: 170_B = 0.003, 62_E < 0.001, and 97_A <

0.001, Benjamini-Hochberg corrected). In balsam fir, only the seedlings inoculated with

the 170_B and 62_E *P. europaea*-like isolates had significantly different root collar

diameter increments than the controls. (Figure 3d, Two-sided Dunn's test P: 170_B =

0.0233, 62_E < 0.001, and 97_A = 0.3189, Benjamini-Hochberg corrected).

Dry-root weight at the final time point also showed significant differences between

treatments Kruskal-Wallis test, H(7) = 61.244, P < 0.001). Post-hoc tests show that both

inoculated Fraser firs (Figure 3e, Two-sided Dunn's test Ps: 170_B < 0.001, 62_E < 0.001,

and 97_A = 0.002, Benjamini-Hochberg corrected) and inoculated balsam firs (Figure 3f,

Two-sided Dunn's test P: 170_B = 0.003, 62_E < 0.001, and 97_A = 0.004, Benjamini-

Hochberg corrected) had significantly lower dry-root weight than their respective

uninoculated control.

Re-isolation of the *P. europaea*-like isolates on diseased seedlings. None of

the necrotic root sections used for reisolation of the *P. europaea*-like isolates yielded live

oomycetes after inoculation on a fresh medium. Using a PCR approach based on the

amplification and sequencing of the nad1 locus allowed the detection of the *P. europaea*-

like isolates in DNA extracted from root samples. While the isolates could be detected in

more than the third of our symptomatic or dead seedlings (16/45 for Fraser fir and 18/45

for balsam fir), they were not detected in any of our uninoculated controls regardless of

their health status. Statistical analyses indicate a significant relationship between

positivity of the nad1 PCR and both the status (healthy or symptomatic/dead) of the

seedling at the end of the experiment (Pearson's Chi-squared test with Yates' continuity

correction, χ^2(1, 120) = 16.860, P = < 0.001) and its inoculation status (Pearson's Chi-

squared test with Yates' continuity correction, χ^2(1, 120) = 14.008, P < 0.001).

Discussion

Phytophthora root rot has been reported throughout the Christmas fir cultivation

range. The problem has been well documented in the USA where both the diversity of the

Phytophthora species (Chastagner and Benson 2000; McKeever and Chastagner 2016)

and the severity (Benson and Grand 2000) of the PRR were described. Surveys were

also conducted in European countries to identify Phytophthora species associated with

PRR on Christmas trees (Pettersson et al. 2019; Shafizadeh and Kavanagh 2005; Talgø

et al. 2006; Talgø et al. 2007). While PRR is present in Canadian Christmas tree

plantations, details about the cohorts *Phytophthora* species and the intensity of the

disease are lacking in Canada. To fulfill this gap, we surveyed PRR in Christmas tree

plantations in Quebec, the province with the largest production in Canada.

Despite extensive surveying over three years, only a small number of

Phytophthora isolates were recovered. In total, 44 *Phytophthora* isolates belonging to

seven different taxa were recovered. Of the 480 soil samples processed, 22 (4.58%)

yielded at least one *Phytophthora* isolates. This is low compared to other studies that

reported *Phytophthora* recovery rates of 21% from forest soil samples of the Eastern and

North-Central U. S. oak ecosystems (Balci et al. 2007), 50% for the Fraser fir plantations

in the Southern Appalachians (Pettersson et al. 2017), 9% from the total sampled trees

in North Carolina (Benson and Grand 2000). Our *Phytophthora* spp. recovery rate is still

lower when isolation from PRR symptomatic trees is considered. Fifteen soil samples

(12.55%) yielded *Phytophthora* isolates compared to 54% and 20% respectively reported

by Pettersson et al. (2017) and Benson and Grand (2000). Direct isolation from roots was

generally less successful than baiting, with less than 10% (4 out of 44) of the isolates

obtained from root samples. Similar results were obtained by (Aghighi et al. 2016), where

most of the *Phytophthora* isolates associated with declining *Rubus anglocandicans* were

recovered by baiting rather than direct isolation from roots.

Phytophthora chlamydospora, P. *gonapodyides*, and *P. gregata* belong to clade 6,

and are considered riparian-associated species (Brasier et al. 1993). These three species

have been reported worldwide in various natural ecosystems (Hansen et al. 2015; Wan

et al. 2020). *Phytophthora chlamydospora* and *P. gonapodyides*, are believed to mainly

act as saprophytes, with episodic fine root infections on weakened hosts (Hansen et al.

2015). Associations between *Phytophthora gonapodyides* and true fir species were

previously reported (Chastagner et al. 1995; McKeever and Chastagner 2016;

Pettersson et al. 2019), but to our knowledge, this study is the first to report the

association between *P. gonapodyides* balsam fir (*Abies balsamea*). *Phytophthora*

chlamydospora was previously found to be associated with *Abies concolor* and *Abies*

procera (Brasier et al. 1993). Still, the present study is the first to report associations of

this species with *Abies balsamea* and *A. fraseri*. *Phytophthora gregata* was identified as

an aggressive pathogen of Australian wetland plants, being able to infect through the

roots and the stem of susceptible hosts (Wan et al. 2020). Other studies have reported

associations of *P. gregata* with other broadleaf forest species (see the summary table of

global *Phytophthora gregata* records in Wan et al. 2020). Still, to our knowledge, this

study is the first report of this species in Canada, and also the first to report the association

of this species with a conifer species (*Abies balsamea*). Further studies are required to

investigate the aggressivity of *P. gregata* on true firs.

Among the isolated *Phytophthora chlamydospora* there was a group of 10 isolates

that did not group with *P. chlamydospora* in our phylogeny. Besides the ITS sequences,

the other loci had higher nucleotide identities with other species. These strains may

represent a new species or a hybridization event which could be confirmed with more in

depth genomic and phenotypic studies.

Phytophthora megasperma is a known aggressive pathogen of fir species and can

affect many other plant hosts. It has been isolated from *Abies fraseri*, *A. grandis*, and *A.*

procera in the USA (Chastagner et al. 1995; McKeever and Chastagner 2016), from *A.*

procera in Ireland (Shafizadeh and Kavanagh 2005), from *A. lasiocarpa* in Norway (Talgø

et al. 2007), and *A. nordmanniana* in Sweden (Pettersson et al. 2019). This study is the

first to report associations of *P. megasperma* with *Abies balsamea* and *A. fraseri*.

True firs were shown to be susceptible to *Phytophthora sp. 'kelmania'*, but this

study is the first to report the association of this *Phytophthora* species with balsam fir (*A.*

balsamea). Previously, McKeever and Chastagner (2016) have isolated *P. sp. 'kelmania'*

from Fraser fir (*Abies fraseri*), Canaan fir (*Abies balsamea var. phanerolepis*) and Turkish

fir (*Abies bornmuelleriana*). This was the second most commonly isolated *Phytophthora*

species in North Carolina on Fraser fir showing PRR symptoms, and artificial inoculations

confirmed that Fraser fir was highly susceptible to *P. sp. 'kelmania'*. More recently, Kondo

et al. (2022) performed artificial inoculations and showed that *P. sp. 'kelmania'* induced

PRR symptoms on Sakhalin fir seedlings (*Abies sachalinensis*).

Phytophthora europaea was initially isolated from soil associated with declining

oak forests (*Quercus spp.* and *Q. robur)* in France and Germany (Jung et al. 2002). It was

since then isolated in Europe and associated with *Quercus* species (Balci and

Halmschlager 2003; Tkaczyk et al. 2017), in North America associated with diverse

deciduous tree species and riparian ecosystems (Balci et al. 2006; Reed et al. 2019;

Reeser et al. 2011; Sims et al. 2015), and in Australia associated with *Dacrydium*

cupressinum and *Iris sibirica* (International Collection Of Microorganisms (ICMP) 2022).

More recently, surveys of Christmas tree plantations in the US recovered anecdotal *P.*

europaea isolates on *Abies fraseri* (McKeever and Chastagner 2016; Pettersson et al.

2017). *P. europaea* has already been previously reported twice in Canada, once from a

diseased balsam fir imported from North Carolina (Robideau et al. 2011), and a second

time from a diseased balsam fir reported in the Annual 2017 Canadian Plant Disease

Survey (Elmhirst 2017). This study constitutes the first official and well documented report

of its presence in Canada's Christmas tree plantations and its association with PRR

symptomatic *Abies balsamea* and *A. fraseri*.

Our phylogenetic analysis revealed that our isolates were more closely related to

P. abietivora than *P. europaea*. However, our phylogeny and the phylogeny of Li et al.

(2019), which formally described *P. abietivora*, show some discordances. They show *P.*

abietivora as a sister species to *P. flexuosa* and those two as a sister group to *P.*

europaea (Li et al. 2019) while we show *P. europaea* and *P. abietivora* in a group sister

to *P. flexuosa*. This may be the result of a difference in the resolution of both phylogenies

as ours covers most of the *Phytophthora* clades but lacks multiple sequences per species

ans Li's phylogeny focuses on clade 7. We decided to identify our isolates as *P. europaea*-

like because *P. abietivora* is a very recently described species initially supported by a

single isolate (Li et al. 2019). Given their high sequence similarity (0.5% out of 3884 base

pairs sequenced), it is probable that isolates were identified as *P. europaea* in studies

preceding the formal description of *P. abietivora* and now would need to be reconsidered

as *P. abietivora*. Molnar et al. (2020) reported the identification of *P. abietivora* isolates

from the collection of the Pennsylvania department of agriculture. They could track the

first isolation of *P.abietivora* from *A. fraseri* and *Tsuga canadensis* as far as 1989. They

also discuss the high similarity in the sequences with *P. europaea*, especially for the ITS

(Molnar et al. 2020). More recently, Bily et al. (2022) isolated *P. abietivora* from multiple

deciduous trees, which would fit with the *P. europaea* hosts already described. They also

observed genetic variation between their strains and the *P. abietivora* type strain for the

loci they sequenced (Bily et al. 2022). In California, a survey reported the isolation of

Phytophthora sp. cadmea, baited from soil of restoration areas. This undescribed taxon

has high sequence similarity with *P. europaea* and *P. abietivora* (Bourret et al. 2020).

Based on those observations, we suggest caution be used when considering *P. europaea*

and *P. abietivora* as different species. While we agree that these species may represent

diverging lineages, more extensive population genomics and phenotypic studies may be

necessary to untangle the relationship between them.

It is interesting to note that to this date *P. europaea* is not reported as associated

with *Abies* species in Europe. Moreover, it is considered as a weak pathogen of deciduous

trees (Jung et al. 2017). In North America, Bily et al. (2022) reported that *P. abietivora*

strains recently collected on deciduous trees also seemed to be weak pathogens to the

deciduous trees species they challenged with their strains. Considering this information,

an interesting hypothesis could be that *P. europaea* was introduced to North America

from Europe through plant nursery trade like other species from this genus (Goss et al.

2009; Moralejo et al. 2009). After spreading throughout North America, it could have

invaded the Christmas tree plantation environment and started diverging, becoming the

P. abietivora that was recently described. Population genomic studies of *P. europaea /P.*

abietivora strains from Europe and North America could help us confirm this scenario.

The fulfillment of Koch's postulates with three *P. europaea*-like isolates obtained

from diseased trees confirmed that this species is able to infect both *A. balsamea* and *A.*

fraseri seedlings. However, disease progression and survival curves suggest that the

infection in *A. balsamea* progresses slower than in *A. fraseri*, suggesting that the *P.*

europaea-like isolates are more aggressive on *A. fraseri*. The sensitivity of *A. fraseri* to

PRR is well documented for other *Phytophthora* species (Benson and Grand 2000;

Hinesley et al. 2000; Quesada-Ocampo et al. 2009) and could explain this difference in

survival throughout of our greenhouse experiment. It is difficult to explain why our

reisolation attempts from root tissues were unsuccessful when the isolates' DNA could

be recovered in root tissues. One hypothesis could be that our surface sterilization was

too harsh, but our methodology did not differ much from what is usually used in such

experiments (Bily et al. 2022; Pettersson et al. 2017). Although we could not directly

satisfy the criteria of the fourth postulate, the fact that we only detected the inoculated

isolates' DNA in inoculated seedlings showing symptoms of PRR confirmed that the

inoculated isolates did indeed cause the disease observed in our seedlings. Our

experiment also showed that *P. europaea*-like isolates could infect seedlings naturally

without prior damage to the trees.

This study provides a sample of the Phytophthora diversity present in the

Christmas tree cultures of the province of Québec. For many of the species reported here,

the association with either *A. fraseri* or *A. balsamea* is novel information and many were

not reported yet in Canada. More investigation will be needed to evaluate the impact of

these potential pathogens as Koch's postulates were only tested on the *P. europaea*-like

isolates. We also present strong evidence that our *P. europaea*-like isolates are root rot

pathogens of balsam and Fraser firs. The information we provide will be helpful in the

development of targeted detection tools and pathogen management strategies that will

help producers lower the risks of PRR outbreaks in their plantations. Other

complementary work with metagenomics analysis and high throughput analysis is also in

process.

<u>Acknowledgments</u>

The authors would like to thank Jean-François Légaré, Éric Dussault, Christian Lacroix,

Dominique Choquette and Jacinthe Drouin for their contributions to fieldwork and

sampling; Marie-Krystel Gauthier, Marie-Josée Bergeron and Véronique Lévesque-

Tremblay for their precious technical advice and Michel Gravel, Marquis Goupil, Renald

Gilbert, Claire Gilbert, Larry Downey, Christian Vanasse, Serge Vaillancourt, Yohan

Blanchette, Junior Beloin, Gérald Couture, Jacques Fortin and David Gouin for giving us

access to their plantations.

-Conflict of interest

The authors declare no conflict of interest.

512   Litterature Cited

Aghighi, S., Burgess, T. I., Scott, J. K., Calver, M., and Hardy, G. E. S. J. 2016. Isolation

and pathogenicity of *Phytophthora* species from declining *Rubus anglocandicans*.

Plant Pathology 65:451-461.

Balci, Y., and Halmschlager, E. 2003. Incidence of *Phytophthora* species in oak forests

in Austria and their possible involvement in oak decline. Forest Pathology 33:157-

174.

Balci, Y., Balci, S., Eggers, J., MacDonald, W. L., Juzwik, J., Long, R., and Gottschalk, K.

W. 2006. First Report of *Phytophthora europaea* in Oak Forests in the Eastern and

North-Central United States. Plant Disease 90:827-827.

Balci, Y., Balci, S., Eggers, J., MacDonald, W. L., Juzwik, J., Long, R. P., and Gottschalk,

K. W. 2007. *Phytophthora* spp. Associated with Forest Soils in Eastern and North-

Central U.S. Oak Ecosystems. Plant Disease 91:705-710.

Bates, R. M., Sellmer, J. C., and Despot, D. A. 2004. Postharvest Characteristics of

Canaan Fir and Fraser Fir Christmas Trees. HortScience HortSci 39:1674-1676.

Benson, D. M., and Grand, L. F. 2000. Incidence of *Phytophthora Root* Rot of Fraser Fir

in North Carolina and Sensitivity of Isolates of *Phytophthora cinnamomi* to

Metalaxyl. Plant Disease 84:661-664.

Bienapfl, J. C., and Balci, Y. 2014. Movement of *Phytophthora* spp. in Maryland's Nursery

Trade. Plant Disease 98:134-144.

Bily, D., Nikolaeva, E., Olson, T., and Kang, S. 2022. *Phytophthora* spp. Associated with

Appalachian Oak Forests and Waterways in Pennsylvania, with *P. abietivora* as a

Pathogen of Five Native Woody Plant Species. Plant Disease 106:1143-1156.

Bourret, T. B., Aram, K., Mehl, H. K., Rizzo, D. M., Rooney-Latham, S., Swiecki, T. J.,

Bernhardt, E. A., and Frankel, S. J. 2020. Ten new provisional species of

Phytophthora and *Nothophytophthora* from California. Pages 46-47 in:

Proceedings of the seventh sudden oak death science and management

symposium: healthy plants in a world with Phytophthora. Gen. Tech. Rep. PSW-

GTR-268.

Brasier, C. 2013. Plant imports, Phytophthoras, and forest degradation. Pages 1-2 in: In:

Frankel, SJ; Kliejunas, JT; Palmieri, KM; Alexander, JM tech. coords. 2013.

Proceedings of the sudden oak death fifth science symposium. Gen. Tech. Rep.

PSW-GTR-243. Albany, CA: US Department of Agriculture, Forest Service, Pacific

Southwest Research Station: 1-2.

Brasier, C. M., Hamm, P. B., and Hansen, E. M. 1993. Cultural characters, protein

patterns and unusual mating behaviour of *Phytophthora gonapodyides* isolates

from Britain and North America. Mycological Research 97:1287-1298.

Charif, D., and Lobry, J. R. 2007. SeqinR 1.0-2: A Contributed Package to the R Project

for Statistical Computing Devoted to Biological Sequences Retrieval and Analysis.

Pages 207-232 in: Structural Approaches to Sequence Evolution: Molecules,

Networks, Populations. U. Bastolla, M. Porto, H. E. Roman and M. Vendruscolo,

eds. Springer Berlin Heidelberg, Berlin, Heidelberg.

Chastagner, G., Riley, K., and Hamm, P. 1990. Susceptibility of *Abies* spp. to seven

Phytophthora spp. Phytopathology 80:887.

Chastagner, G. A., and Benson, D. M. 2000. The Christmas Tree: Traditions, Production,

and Diseases. Plant Health Progress 1:15.

Chastagner, G. A., Hamm, P. B., and Riley, K. L. 1995. Symptoms and *Phytophthora* spp.

associated with root rot and stem canker of Noble fir Christmas trees in the Pacific

Northwest. Plant Disease 79:290-293.

Eklund, A., Trimble J. 2021. beeswarm: The Bee Swarm Plot, an Alternative to Stripchart.

Elmhirst, J. 2017. Canadian plant disease survey. Page 71. The Canadian

Phytopathological Society, Canada.

Farr, D. F., Rossman, A.Y. 2022. Fungal Databases, U.S. National Fungus Collections.

ARS USDA, ed.

Frampton, J., and Benson, D. M. 2012. Seedling resistance to *Phytophthora cinnamomi*

in the genus *Abies*. Annals of Forest Science 69:805-812.

Goss, E. M., Larsen, M., Chastagner, G. A., Givens, D. R., and Grünwald, N. J. 2009.

Population Genetic Analysis Infers Migration Pathways of *Phytophthora ramorum*

in US Nurseries. PLOS Pathogens 5:e1000583.

Grand, L., and Lapp, N. 1974. *Phytophthora cinnamomi* root rot of Fraser Fir in North

Carolina. Plant Disease Reporter 58:318-320.

Hamm, P. B., and Hansen, E. M. 1982. Pathogenicity of *Phytophthora* species to Pacific

Northwest Conifers. European Journal of Forest Pathology 12:167-174.

Hansen, E. M., Reeser, P., Sutton, W., and Brasier, C. M. 2015. Redesignation of

Phytophthora taxon Pgchlamydo as *Phytophthora chlamydospora* sp. nov. North

American Fungi 10:1-14.

Hardham, A. R. 2001. The cell biology behind *Phytophthora* pathogenicity. Australasian

Plant Pathology 30:91-98.

Hinesley, L. E., Parker, K. C., and Benson, D. M. 2000. Evaluation of Seedlings of Fraser,

Momi, and Siberian Fir for Resistance to *Phytophthora cinnamomi*. HortScience

HortSci 35:87-88.

Hoover, B. K., and Bates, R. M. 2013. Susceptibility of Fraser, Canaan, and Nordmann

Fir to Root Rot Incited by *Phytophthora cactorum* and *Phytophthora drechsleri*.

HortTechnology hortte 23:44-50.

International Collection Of Microorganisms (ICMP). 2022. Specimen data. Accessed

through Systematics Collection Data.

Judelson, H. S., and Blanco, F. A. 2005. The spores of *Phytophthora*: weapons of the

plant destroyer. Nature Reviews Microbiology 3:47-58.

Jung, T., Hansen, E. M., Winton, L., Osswald, W., and Delatour, C. 2002. Three new

species of *Phytophthora* from European oak forests. Mycological Research

106:397-411.

Jung, T., Orlikowski, L., Henricot, B., Abad-Campos, P., Aday, A., Aguín Casal, O.,

Bakonyi, J., Cacciola, S., Cech, T., and Chavarriaga, D. 2016. Widespread

Phytophthora infestations in European nurseries put forest, semi-natural and

horticultural ecosystems at high risk of *Phytophthora* diseases. Forest Pathology

46:134-163.

Jung, T., Chang, T. T., Bakonyi, J., Seress, D., Pérez-Sierra, A., Yang, X., Hong, C.,

Scanu, B., Fu, C. H., Hsueh, K. L., Maia, C., Abad-Campos, P., Léon, M., and

Horta Jung, M. 2017. Diversity of *Phytophthora* species in natural ecosystems of

Taiwan and association with disease symptoms. Plant Pathology 66:194-211.

Kampstra, P. 2008. Beanplot: A Boxplot Alternative for Visual Comparison of

Distributions. Journal of Statistical Software, Code Snippets 28:1 - 9.

Kassambra, A., Kosinski, M., Biecek P. 2021. survminer: Drawing Survival Curves using

'ggplot2'.

Kondo, T., Otsubo, K., and Kageyama, K. 2022. *Phytophthora* rot of alpine delphinium

caused by *Phytophthora* sp. kelmania. Journal of General Plant Pathology 88:145-

149.

Kuhlman, E. G., and Hendrix, F. 1963. *Phytophthora* root rot of Fraser Fir. Plant Disease

Reporter 47:552-553.

Kumar, S., Stecher, G., Li, M., Knyaz, C., and Tamura, K. 2018. MEGA X: Molecular

Evolutionary Genetics Analysis across Computing Platforms. Mol Biol Evol

35:1547-1549.

Li, D.-W., Schultes, N. P., LaMondia, J. A., and Cowles, R. S. 2019. *Phytophthora*

abietivora, A New Species Isolated from Diseased Christmas Trees in Connecticut,

U.S.A. Plant Disease 103:3057-3064.

Madden, L. V., Hughes, G., and Van Den Bosch, F. 2007. The study of plant disease

epidemics. Am Phytopath Society.

McKeever, K. M., and Chastagner, G. A. 2016. A Survey of *Phytophthora* spp. Associated

with *Abies* in U.S. Christmas Tree Farms. Plant Disease 100:1161-1169.

Mitcham-Butler, E. J., Eric Hinesley, L., and Pharr, D. M. 1988. Effects of Harvest Date,

Storage Temperature, and Moisture Status on Postharvest Needle Retention of

Fraser Fir. Journal of Environmental Horticulture 6:1-4.

Molnar, C., Nikolaeva, E., Kim, S., Olson, T., Bily, D., Kim, J.-E., and Kang, S. 2020.

Phytophthora Diversity in Pennsylvania Nurseries and Greenhouses Inferred from

Clinical Samples Collected over Four Decades. Microorganisms 8:1056.

Moralejo, E., Pérez-Sierra, A. M., Álvarez, L. A., Belbahri, L., Lefort, F., and Descals, E.

2009. Multiple alien *Phytophthora* taxa discovered on diseased ornamental plants

in Spain. Plant Pathology 58:100-110.

Ogle, D. 2017. Package 'FSA'. CRAN Repos:1-206.

Pettersson, M., Frampton, J., and Sidebottom, J. 2017. Influence of *Phytophthora* root rot

on planting trends of Fraser fir Christmas trees in the Southern Appalachian

Mountains. Tree Plant. Notes 60:4-11.

Pettersson, M., Frampton, J., Rönnberg, J., Brurberg, M. B., and Talgø, V. 2019.

Presence of *Phytophthora* species in Swedish Christmas tree plantations.

European Journal of Plant Pathology 153:1221-1236.

Quesada-Ocampo, L. M., Fulbright, D. W., and Hausbeck, M. K. 2009. Susceptibility of

Fraser Fir to *Phytophthora capsici*. Plant Disease 93:135-141.

R Core Team. 2013. R: A language and environment for statistical computing.

Reed, S. E., English, J. T., and Muzika, R.-M. 2019. *Phytophthora* Species Detected in

Two Ozark Forests with Unusual Patterns of White Oak Mortality. Plant Disease

103:102-109.

Reeser, P. W., Sutton, W., Hansen, E. M., Remigi, P., and Adams, G. C. 2011.

Phytophthora species in forest streams in Oregon and Alaska. Mycologia 103:22-

35.

Robideau, G. P., De Cock, A. W., Coffey, M. D., Voglmayr, H., Brouwer, H., Bala, K.,

Chitty, D. W., Désaulniers, N., Eggertson, Q. A., Gachon, C. M., Hu, C. H., Küpper,

F. C., Rintoul, T. L., Sarhan, E., Verstappen, E. C., Zhang, Y., Bonants, P. J.,

Ristaino, J. B., and Lévesque, C. A. 2011. DNA barcoding of oomycetes with

cytochrome c oxidase subunit I and internal transcribed spacer. Mol Ecol Resour

11:1002-1011.

Shafizadeh, S., and Kavanagh, J. A. 2005. Pathogenicity of *Phytophthora* species and

Pythium undulatum isolated from *Abies procera* Christmas trees in Ireland. Forest

Pathology 35:444-450.

Sims, L. L., Sutton, W., Reeser, P., and Hansen, E. M. 2015. The *Phytophthora* species

assemblage and diversity in riparian alder ecosystems of western Oregon, USA.

Mycologia 107:889-902.

Statistics Canada. 2016. Table 32-10-0421-01 Christmas tree production, Census of

Agriculture.

Statistics Canada. 2021. Table 32-10-0249-01 Land use, Census of Agriculture.

Talgø, V., Herrero, M., Toppe, B., Klemsdal, S., and Stensvand, A. 2006. First Report of

Root Rot and Stem Canker Caused by *Phytophthora cambivora* on Noble Fir

(Abies procera) for Bough Production in Norway. Plant Disease 90:682-682.

Talgø, V., Herrero, M. L., Toppe, B., Klemsdal, S. S., and Stensvand, A. 2007.

Phytophthora Root Rot and Stem Canker Found on Nordmann and Subalpine Fir

in Norwegian Christmas Tree Plantations. Plant Health Progress 8:29.

Talgø, V., Schmitz, S., Chandelier, A., Brurberg, M., and Thomsen, I. 2017. Baiting for

Phytophthora in waterways associated with Christmas tree production in Norway,

Belgium and Denmark. NIBIO BOOK 3:80-82.

Tamura, K., and Nei, M. 1993. Estimation of the number of nucleotide substitutions in the

control region of mitochondrial DNA in humans and chimpanzees. Molecular

Biology and Evolution 10:512-526.

Therneau, T. M., and Grambsch, P. M. 2000. Testing Proportional Hazards. Pages 127-

152 in: Modeling Survival Data: Extending the Cox Model. Springer New York,

New York, NY.

Tkaczyk, M., Milenković, I., Nowakowska, J. A., Borys, M., Kałuski, T., Gawlak, M., Czyż,

M., and Oszako, T. 2017. Morphological and molecular identification of

Phytophthora species isolated from the rhizosphere of declining oak trees in

Krotoszyn Plateau. Genetika 49:203-215.

USDA National Agricultural Statistics Service. 2021. Cut Christmas trees, acres in

production.

Wan, J. S. H., McDougall, K. L., and Liew, E. C. Y. 2020. The susceptibility of seven

threatened species to *Phytophthora gregata* and the aetiology of the disease

caused by it. Australian Journal of Botany 68:595-601.

Fig. 1. Multiple *Phytophthora* species were recovered in plantations and natural forests: Maximum likelihood analysis of the recovered isolates and reference taxa based on concatenated ITS, Cox1, β-Tubulin, Nad1, and HSP90 sequences. *Pythium undulatum* is used as an outgroup. The bootstrap test was conducted with 1,000 replicates bootstrap

693 values >80% are indicated on the nodes. The scale bar shows the number of expected

694 substitutions per site. References are represented in italic and isolates obtained in this

695 study are represented in bold.

Fig. 2. Rapid onset of PRR symptoms lead to seedlings death in inoculated groups: (**A,**

B) Disease progression curve in Fraser and balsam fir seedlings represented as the

percentage of the plant affected by symptoms of PRR for each seedling. The replicates'

data was averaged for each tested isolate and uninoculated controls (Ctrl); darkened

areas represent 95% confidence interval and letters indicate significant differences in

AUDPC (two-sided Dunn's test). (**C**) Survival probability varies between inoculated and

uninoculated (Ctrl) seedlings; grey horizontal lines represent the timing of each 48-hours

flooding period and letters indicate significant differences in survival curves (Log-rank

test).

Fig. 3. Inoculation with *P. europaea*-like isolates affects growth in both Fraser and balsam

fir seedlings: (**A, B**) Beanplots comparing height distributions between the initial time point

(inoculation week, t_0) and the final time point (12 weeks post-inoculation, t_{12}) for each

isolate inoculated and the uninoculated controls (Ctrl). Letters indicate significant

differences in height increments (Dunn's test). (C, D) Beanplots comparing collar diameter

distribution between t_0 and t_{12} for each isolate inoculated and the uninoculated controls

(Ctrl). Letters indicate significant differences in diameter increments (Dunn's test). (E, F)

Beanplots of dry root mass distribution at t_{12} for each isolate inoculated and the

uninoculated controls (Ctrl). Letters indicate significant differences in dry root weight

(Dunn's test). Black bars on the plots illustrate the mean of the data. Symbols represent

individual data points for each time point.