Tropomyosin 1-I/C co-ordinates kinesin-1 and dynein motors during *oskar* mRNA transport

Simone Heber[1†], Mark A. McClintock[2†], Bernd Simon[3], Janosch Hennig[3,4], Simon L. Bullock[2*], Anne Ephrussi[1*]

[1]   Developmental Biology Unit, European Molecular Biology Laboratory, 69117 Heidelberg, Germany

[2]   Division of Cell Biology, MRC Laboratory of Molecular Biology, Cambridge CB2 0QH, United Kingdom

[3]   Structural and Computational Biology Unit, European Molecular Biology Laboratory, 69117 Heidelberg, Germany

[4]   Biochemistry IV, Biophysical Chemistry, University of Bayreuth, 95447 Bayreuth, Germany

[†]These authors contributed equally to this work

[*]Correspondence: sbullock@mrc-lmb.cam.ac.uk, anne.ephrussi@embl.org

Abstract

Dynein and kinesin motors mediate long-range intracellular transport, translocating towards microtubule minus and plus ends, respectively. Cargoes often undergo bidirectional transport by binding to both motors simultaneously. However, it is not known how motor activities are coordinated in such circumstances. In *Drosophila*, sequential activities of the dynein-dynactin-BicD-Egalitarian (DDBE) complex and of kinesin-1 deliver *oskar* mRNA from nurse cells to the oocyte, and within the oocyte to the posterior pole. Here, through *in vitro* reconstitution, we show that Tm1-I/C, a Tropomyosin-1 isoform, links kinesin-1 in an inactive state to DDBE-associated *oskar* mRNA. NMR spectroscopy, small-angle X-ray scattering and structural modeling indicate that Tm1-I/C suppresses kinesin-1 activity by stabilizing its autoinhibited conformation, thus preventing a tug-of-war between the opposite polarity motors until kinesin-1 is activated in the oocyte. Our work reveals a novel strategy ensuring sequential activity of microtubule motors.

Introduction

Motor protein driven transport along cytoskeletal tracks is important for subcellular localization of many cargoes, including organelles, vesicles and RNAs. Whereas myosin motors transport cargoes on actin filaments[1], cytoplasmic dynein and kinesin motors are responsible for translocation along microtubules, driving minus-end-directed and plus-end-directed movements, respectively. The cargoes for these motors, however, often travel bidirectionally, requiring the coordinated activity of the two motor classes. For instance, in neurons, mRNAs[2,3] and mitochondria[4] are shuttled bidirectionally to ensure their constant supply to distal regions. Because dynein and kinesin motors are often simultaneously bound to the same cargo[5,6], their activities must be tightly coordinated to allow for efficient transport rather than a futile tug-of-war.

Individual motors appear to be in an autoinhibited state by default, with activation driven by binding of adaptor proteins and cargoes. For instance, dynein is activated by binding of dynactin and one of a number of coiled-coil containing activating adaptors[7,8,9,10,11]. In the case of mRNA transport in *Drosophila*, dynein uses Bicaudal-D (BicD) as the activating adaptor, which is linked to double-stranded mRNA localization signals by the RNA-binding protein Egalitarian (Egl)[12,13]. Interaction of Egl and BicD with each other, as well as with the motor are enhanced by binding the mRNA signal, revealing a role for the mRNA in controlling motor activity[12,13,14,15].

Kinesin-1 is a tetramer consisting of two Kinesin heavy chains (Khc) and two Kinesin light chains (Klc)[16]. Khc contains an N-terminal globular motor domain, a coiled-coil stalk domain and a C-terminal disordered tail domain (Figure 1a) and is autoinhibited by folding of the tail region of the heavy chain back on its motor domain. This interaction has been proposed to involve pivoting of the coiled-coil stalk at a hinge region, which allows docking of a conserved isoleucine-alanine-lysine (IAK) motif in the tail onto the motor domain, thereby locking the complex in a conformation that prevents processive movement[17,18,19,20]. The autoinhibited conformation is stabilized by Klc[21,22], which binds to the Khc stalk domain[23,24] and is important for recognition of a range of cargoes[25].

However, the mechanisms of regulation in dual motor transport remain elusive, with only a small number of adaptor proteins that coordinate both motors having been reported. For example, the mitochondrial transport adaptor TRAK2 forms a complex with both kinesin-1 and dynein-dynactin and links these two motors functionally[26,27]. UNC-16, the *C. elegans* homolog of the lysosome adaptor JIP3/Syd, was also shown to link kinesin-1 and dynein, allowing for

plus-end-directed transport of dynein by kinesin-1[28,29]. Similarly, the dynein adaptor Hook3 can

activate and link KIF1C (a kinesin-3 motor) and dynein[30,31].

oskar (*osk*) mRNA localization in the *Drosophila* egg chamber is a paradigm for the study of

RNA transport and motor regulation. Motor-based delivery of *osk* to the posterior pole of the

developing oocyte drives abdominal patterning and germline formation in the embryo[32].

Trafficking of *osk* mRNA to this site is mediated by the successive activities of the dynein[33]

and kinesin-1[34.35] motors, and requires motor switching. In the germline syncytium, the oocyte

is transcriptionally quiescent and relies on mRNAs provided by the 15 interconnected nurse

cells for its development[36]. At this stage, microtubule minus ends are nucleated in the oocyte

and extend into the nurse cells. Upon transcription in the nurse cells, *osk* mRNA is transported

along the polarized microtubule cytoskeleton into the oocyte by dynein in complex with

dynactin, BicD and Egl[12,33,37]. During mid-oogenesis, polarity of the microtubule cytoskeleton

changes dramatically, with microtubule plus ends becoming enriched at the oocyte posterior.

At this point, Khc translocates *osk* to the posterior pole[38]. Khc was previously shown to function

in *Drosophila* oocytes largely independently of Klc[39,40], raising the question of how Khc is linked

to *osk* and how motor activity is regulated.

Kinesin-mediated transport of *osk* RNA requires the unique I/C isoform of tropomyosin-1, Tm1-

I/C (hereafter referred to as Tm1)[39,40,41,42]. Tm1 binds to a non-canonical but conserved cargo

binding region in the Khc tail and stabilizes the interaction of the motor with RNA, suggesting

a function as an adaptor[43]. Both Khc and Tm1 are already loaded onto the *osk*

ribonucleoprotein particle (RNP) shortly after nuclear export in the nurse cells, although the

motor only appears to become active in the oocyte during mid-oogenesis[42]. Similarly, dynein

remains associated with *osk* RNPs during Khc-mediated transport within the oocyte, but is

inactivated by displacement of Egl by the Staufen (Stau) protein[44,45].

How the two motors are linked simultaneously to *osk* RNPs, and how cargo-bound Khc is

inhibited during dynein-mediated transport into the oocyte has been unclear. Here, we show

that Tm1 inhibits Khc activity by stabilizing its autoinhibited conformation through a novel

mechanism involving the regulatory tail domain. In addition, Tm1 links Khc to the dynein-

transported *osk* RNP, thereby allowing co-transport of inactive Khc on *osk* RNA by dynein. *In*

vivo, such a mechanism would avoid a tug-of-war between the two motors during the dynein-

dependent delivery of *osk* RNPs to the oocyte whilst ensuring that Khc is available on these

structures when it is time for this motor to move the RNPs to the posterior of the oocyte. With

its cargo-binding and motor regulatory functions, we propose that Tm1 functions as a non-

canonical light chain for kinesin-1 in the *Drosophila* oocyte.

Results

Tm1 can transport RNA with Khc *in vitro* but impairs Khc motility

In order to test the ability of Khc, Tm1 and the Khc-Tm1 complex to transport RNA, we performed total internal reflection fluorescence (TIRF) microscopy-based *in vitro* motility assays[15] using fluorescently labeled versions of recombinant, full-length Khc (Khc FL), full-length Tm1 (Tm1 FL) (Figure 1a, b) and *in vitro* transcribed *osk* 3'UTR RNA. At concentrations that allowed single-particle tracking (6.25 nM of each component), we observed transport of *osk* by Khc along microtubules in both the presence and absence of Tm1, albeit at low frequency (Figure 1c). In the absence of Tm1, 4.8 % (4/84) of Khc transport events co-localized with the *osk* 3'UTR signal. This value did not change significantly in the presence of Tm1 (2.8 %, 1/36; Figure 1d), despite 55 % (20/36) of Khc signals co-localizing with Tm1 signals. Correcting for the efficiency of labeling of Tm1 with the fluorophore (57 %) indicated that ~95 % of microtubule-bound Khc molecules were bound to Tm1. The high occupancy of Khc with Tm1 in these experiments is consistent with the strong binding between their interacting domains observed previously[43]. Mass photometry confirmed high occupancy of Khc with Tm1 at nanomolar concentrations (Figure S1), as well as the proposed 2:1 stoichiometry of the Khc-Tm1 complex[43]. We conclude from these experiments that Khc can associate with *osk* RNA but that the Khc-Tm1 complex does not have substantially higher affinity for the transcript in our assay conditions.

We next examined the effect of Tm1 on Khc motility. Without Tm1 present in the assay, microtubule-associated Khc complexes moved with a mean velocity of 0.221 μm s^{-1}. In the presence of Tm1, the mean velocity of Khc runs reduced to 0.068 μm s^{-1} (Figure 1e). These data reveal an inhibitory effect of Tm1 on Khc motility.

In addition to the 3'UTR, an RNA structure in the *osk* coding sequence, which is called the spliced *oskar* localization element (SOLE), is important for Khc-mediated transport of *osk in vivo*[46,47]. We tested if an RNA including the SOLE could stimulate motility of Khc in the presence of Tm1 by replacing the *osk* 3'UTR in our assay with a 135-nt RNA (*osk* min) comprising the SOLE and the oocyte entry signal of *osk* (Figure S2a), which resides in the 3'UTR and promotes dynein-mediated transport[48]. The inhibitory effect of Tm1 on Khc motility was also observed in the presence of *osk* min (Figure S2b-f). Thus, Tm1 inhibits motility of Khc in both the presence and absence of the SOLE element.

Tm1 inhibits Khc motility via the Khc regulatory tail

In order to elucidate the molecular mechanism behind Tm1-mediated inhibition of Khc, we set out to identify the region of Khc that is involved in this process. Regulatory functions have been attributed to an auxiliary, ATP-independent microtubule-binding domain (AMB) and the autoinhibitory IAK motif within the Khc tail (Figure 1a)[19,49,50]. Whereas the AMB allows for microtubule-sliding by Khc[51], the IAK motif inhibits Khc's ATPase activity[17,19], and thus processive motility, by binding to the Khc motor domain and locking the two Khc heads in a conformation that prevents neck-linker undocking and ADP-release[18]. We tested whether the regulatory tail is responsible for Tm1-mediated inhibition of Khc by assessing the effect of Tm1 on the motility of a tailless Khc truncation (Khc910; Figure 2a). As expected of a Khc without its regulatory tail, Khc910 was hyperactive in the *in vitro* motility assay, exhibiting more processive runs (Figure 2c and Table S1), an increased likelihood of processive movement upon binding microtubules (Figure 2d), and faster motility (Figure 2e and Table S1) than the full-length motor complex. However, the presence of Tm1 did not inhibit Khc910 motility (Figure 2c-e and Table S1). The same effects were observed when experiments were performed with *osk* min instead of *osk* 3'UTR (Figure S2c-f). These observations demonstrate that the regulatory tail of Khc is required for Tm1-mediated inhibition of motility.

To assess the relevance of the autoinhibitory IAK motif in the Khc tail for Tm1-mediated inhibition, we deleted this feature whilst leaving the AMB intact. To this end, we produced Khc940, which is truncated C-terminal to the AMB domain. Khc940 was hyperactive (in terms of the number of processive runs, fraction of Khc binding events that resulted in processive motility, and velocity) to a degree comparable to that of Khc910 (Figure 2b-e). However, despite the fact that a smaller fraction of Khc940 co-localized with Tm1 as compared with Khc FL and Khc910 (Figure S3), Khc940 was sensitive to inhibition by Tm1, as the number of runs, fraction of processive Khc, and Khc velocities were reduced in its presence (Figure 2b-e and Table S1). We conclude that Khc inhibition by Tm1 does not depend on the IAK motif and therefore involves another region of the Khc tail.

Khc inhibition by Tm1 is mediated by the Khc AMB domain

To test if the AMB domain in the Khc tail is required for inhibition by Tm1, we produced a Khc mutant in which only the AMB domain (aa 911-938) was deleted (KhcΔAMB) and tested its motility with and without Tm1 (Figure 3a). KhcΔAMB did not have an increased frequency or likelihood of processive movement or higher velocity compared to Khc FL (Figure 3b-d and Table S2), indicating that autoinhibition is not disrupted in this protein. Strikingly, the frequency, likelihood of motility, and velocity of processive KhcΔAMB movements were not decreased by Tm1 (Figure 3a-d and Table S2), despite the proportion of complexes associated with Tm1

171   not differing significantly from what was observed for Khc FL or Khc910 (Figure S4). Taken

172   together, these results reveal that the AMB domain of Khc mediates inhibition by Tm1 but is

173   not necessary for Tm1 binding.

The N-terminal domain of Tm1 is required for its inhibitory function

We next investigated which features of Tm1 mediate Khc inhibition. We previously showed

that a truncated Tm1 protein (1-335 of the 441-amino-acid full-length protein) comprising the

disordered N-terminal domain and the minimal Khc-binding region of the coiled-coil domain

can suppress *osk* RNA mislocalization in *Drosophila* oocytes that lack the endogenous Tm1

protein[43]. We therefore compared the effect of Tm1 1-335 to Tm1 FL in the motility assay

(Figure 4a) and found that both proteins inhibit Khc motility to a similar extent (Figure 4b-d

and Table S3). We also observed that the fraction of Tm1-bound Khc did not differ significantly

between Tm1 1-335 and Tm1 FL (Figure S5). These data show that the N-terminal region of

Tm1 is sufficient to inhibit Khc.

To address whether the Khc-binding domain of Tm1 (aa 252-334) is also sufficient to produce

an inhibitory effect we performed motility assays with this protein. We compared the effects of

untagged, unlabeled Tm1 FL and Tm1 252-334 (Figure 4e) on Khc motility. Whereas Tm1 FL

again decreased the number of processive runs of Khc, we observed no negative effect of

Tm1 252-334 on motility (Figure 4e, f, Figure S5d-f and Table S3). These results suggest that

inhibition of Khc requires the unstructured, N-terminal domain of Tm1.

Tm1 binding affects the disordered region of the regulatory Khc tail

Next, we set out to better understand the molecular mechanism of Khc inhibition by Tm1.

Because the inhibitory effect of Tm1 is mediated by the auxiliary microtubule-binding region

(AMB) within the Khc tail, we reasoned that Tm1 could alter microtubule binding by the tail

domain, perhaps by providing an additional interaction with microtubules. In order to test this

notion, we performed microtubule-binding assays[52] with recombinant Tm1 proteins. Tm1 was

incubated with polymerized microtubules before pelleting of microtubules, together with bound

protein, by ultracentrifugation. As a positive control, we confirmed that both Khc FL and Khc

tail partially partitioned into the microtubule-containing pellet fraction (Figure S6a). A small

fraction of Tm1 1-335 also partitioned into the pellet, indicating that the N-terminal or coiled-

coil domain of Tm1 can interact with microtubules. However, we did not observe increased

partitioning of the Khc tail (Khc 855-975) into the microtubule fraction in the presence of Tm1

1-335 (Figure S6b). This observation indicates that association with Tm1 does not alter the

degree of binding of the Khc tail to microtubules.

In the crystal structure of the Khc tail bound to Tm1, the Khc coiled coil extends into the AMB

domain (last visible aa: 916)[43]. Attempts to crystallize the same Khc construct alone were

unsuccessful, raising the possibility that the isolated motor protein is more flexible. It is
conceivable that the extended coiled coil of Khc is induced upon Tm1 binding, which could
reduce flexibility of the tail, potentially locking it in a conformation that impairs Khc activity. To
test this hypothesis, we probed the structure of the Khc alternative cargo-binding domain and
tail (Khc 855-975) in solution by NMR spectroscopy. The ^{1}H,^{15}N-HSQC NMR spectrum of the
Khc tail showed 34 resonances, each corresponding to a backbone N-H correlation (Figure
5a). Backbone assignment experiments and resulting secondary chemical shifts revealed that
the visible peaks are disordered and correspond to aa 937-975, the very C-terminal part of the
Khc tail that contains the auto-inhibitory IAK motif. For all tested coiled-coil constructs of Khc,
resonances for aa 855-936 were broadened beyond detection due to short transversal
relaxation times. The unfavorable relaxation behavior of H-N resonances in coiled-coil
structures is caused by slow rotational tumbling of the axis perpendicular to the helix
orientation which is, in the case of Khc-Tm1, potentially further slowed down by transient
intermolecular oligomerization. The absence of resonances for aa 855-936 is thus consistent
with this entire part of the protein forming an extended coiled-coil structure, even in the
absence of Tm1. Our data therefore suggest that the AMB domain is part of the structured
coiled coil even in absence of Tm1.
224
To further investigate the effect of Tm1-binding on the disordered tail, we performed NMR
titration experiments to detect any effects of Tm1 binding on the tail region. Upon titration of
increasing amounts of Tm1, there were intensity decreases for several resonances in ^{1}H,^{15}N-
HSQC spectra of Khc 855-975 (Figure 5b, c). The affected residues correspond to the IAK
motif and its flanking regions. This effect could be observed both in the absence and presence
of a U10 RNA (Figure 5c, d), and depended on the N-terminal domain of Tm1 because the
Tm1 coiled-coil (aa 252-334) construct did not affect the spectra (Figure 5e). These findings
are in agreement with our earlier findings that Tm1 inhibits Khc motility in the absence of RNA
and that the N-terminal region of Tm1 is required for its inhibitory effect. These intensity
decreases for the IAK motif and its proximal residues indicate an induced conformational
change or a second, transient interaction of the Tm1 N-terminal domain with this region.

Tm1 does not affect the interaction of the Khc tail with the isolated Khc motor domain

The above results led us to hypothesize that Tm1, upon binding to the coiled coil of Khc that
includes the AMB domain, influences the interaction of the Khc tail and motor domains. To test
this notion, we performed NMR titration experiments in which the Khc motor domain (aa Khc
1-365) was added to the tail domain prior to Tm1 1-335 titration. Titration of Khc 1-365 to
Khc 855-975 caused intensity losses for several resonances. As expected, the resonances
corresponding to the IAK motif (aa 941-948), as well as its flanking residues (aa 936-962),

were affected (Figure 5f, g). Some resonances in this region also showed chemical shift
perturbations (Figure 5f, h). Taken together, these data confirm an interaction between Khc
855-975 (tail) and Khc 1-365 (motor). Additional titration of Tm1 1-335 to this Khc tail-motor
complex caused further intensity decreases of the same resonances (Figure 5f, g), showing
that Tm1 1-335 can bind to the Khc tail when it is associated with the motor domains. We
conclude that binding of the Khc tail to Tm1 and the Khc motor are not mutually exclusive.
250
We therefore tested whether Tm1 can directly affect the interaction of the Khc motor and tail
domains in GST-pulldown assays (Figure S7). When used as bait, the GST-tagged Khc tail
domain (Khc 855-975) simultaneously captured the Khc motor domain (Khc 1-365) and
Tm1 1-335, consistent with our NMR-based evidence that formation of a trimeric complex is
possible. However, the fractions of Khc 1-365 and Tm1 1-335 pulled down by GST-Khc 855-
975 were low, indicating a rather weak or transient interaction. Upon titration of Tm1 1-335,
the amount of motor domain associating with the tail domain was not affected. Thus, we
conclude that Tm1 does not alter the ability of the isolated Khc motor domain to interact with
the isolated Khc tail domain.

Tm1 alters the auto-inhibited conformation of Khc

We next reasoned that Tm1 binding might stabilize the autoinhibited conformation of the full-
length Khc by favoring additional interactions within the coiled-coil stalk domain, likely including
the AMB domain. In order to obtain information about Khc's predominant conformation in its
free and Tm1-bound states, we measured small-angle X-ray scattering (SAXS) of Khc and the
Khc-Tm1 complex (Figure 6a). A dimensionless Kratky plot[53,54] of both scattering profiles
(Figure 6b) shows that Khc and Khc-Tm1 are extended molecules with flexible elements. To
identify structures that can explain the scattering curves of Khc and Khc-Tm1, we generated
a large pool of Khc and Khc-Tm1 models by connecting coordinates of the structured parts
that were generated by AlphaFold2[55,56]. Based on AlphaFold2 predictions, the Khc stalk
contains four coiled-coil regions (cc 1-4) connected by flexible hinge regions (hinges 2-4). The
stalk is connected to the motor domain's neck coiled coil by another flexible region (hinge 1)
(Figure S8a) We subsequently randomized the connecting linker and tail regions to create a
large pool of possible structures. In a second step, the C-terminal tail of one chain was pulled
into the binding pocket of the motor domain resulting in a second pool of structures with
random orientation of the stalk region and the IAK peptide bound to the motor domain dimer.
Both pools of structures were sorted by the quality of the fit to the SAXS data.
277
For both Khc and the Khc-Tm1 complex, the structures with the lowest χ^2 values (i.e. best
quality fits) all exhibited compact conformations with considerably lower than average values
for the radius of gyration (Rg). Since the autoinhibited conformations were more compact than

the initial pool with randomized linker conformations, there were on average more structures
with low χ^2 values in the pools of structures with the IAK motif bound to the motor domains
(Figure S8b). Inspection of the structural models that best fit the scattering curves indicated
that the arrangement of the coiled-coils is different for Khc and Khc-Tm1. Looking at the angles
(Figure 6c-d) of consecutive coiled-coil regions of Khc models, we observed that the
conformation of the hinge regions (hinge 2: aa 586-603, hinge 3: aa 707-710 and hinge 4:
aa 839-843, Figure S8a) leads to bent structures of the coiled-coil regions cc1-cc2 and cc3-
cc4 and elongated cc2-cc3 orientation, resulting in a triangular shape of the molecule (Figure
6c, Figure S8d). In this conformation, the N- and C-terminal parts of the stalk are separated,
and the AMB domain, which mediates inhibition by Tm1, is not in proximity to other stalk
regions (Figure 6c, Figure S8d-e). For Khc-Tm1, the stalk folds at hinge 3 (Figure 6c, Figure
S8d), bringing N- and C-terminal halves of the stalk in proximity, with the AMB domain oriented
parallel to the N-terminal stalk region (Figure S8f). Fits with single models are a crude
approximation because in solution, Khc likely is in a dynamic equilibrium of a multitude of
autoinhibited and more open conformations. Nevertheless, such single models reflect a
possible predominant or average conformation of the protein. These data therefore suggest
that binding of Tm1 to Khc results in a shift of the conformational equilibrium towards different,
more compact structures of the Khc stalk domain. The autoinhibited conformation of Khc is
likely stabilized by interactions within the Tm1-bound stalk in addition to the motor-tail
interaction. This is in agreement with our findings that the Khc AMB domain in complex with
Tm1 can inhibit Khc independently of the IAK motif.

Tm1 links inactive Khc to dynein RNPs for minus-end-directed *osk*
transport

Khc and Tm1 are loaded onto *osk* RNPs in nurse cells prior to dynein-driven transport into the
oocyte[42]. We reasoned that during this process, enhanced inhibition of Khc by Tm1 would
favor minus-end-directed movement by preventing a tug-of-war between dynein and Khc that
could result in counterproductive movements of RNPs away from the oocyte. To assess if this
is the case, we set out to reconstitute *osk* RNPs containing Khc, Tm1 and dynein *in vitro*.
309
Since dynein and kinesin require different conditions for optimal motility, we used a modified
assay buffer that allows movement of both motor classes[58]. To exclude the possibility that Tm1
promotes minus-end-directed movement of *osk* RNPs by stimulating dynein activity directly,
we first assessed its effects on the motile properties of the minimal machinery for dynein-
mediated RNA transport (dynein, dynactin, BicD, Egalitarian (DDBE)) assembled with an *osk*
3'UTR fragment that is sufficient to initiate dynein movement *in vivo* (*osk* 3'UTR 2+3)[48]. While
Tm1 could be observed translocating with dynein and the *osk* RNA (Figure 7a), its presence
did not alter dynein velocity or run length on microtubules (Figure 7b, c and Table S4).

Furthermore, Tm1 had no effect on the percentage of microtubule-bound dyneins that undergo
directed motility or the frequency of dynein binding to microtubules or initiating transport along
them (Figure S9a-c and Table S4). Thus, Tm1 does not directly affect dynein motility.
Treatment of these reconstituted RNPs with RNase A after assembly dramatically reduced the
incidence of Tm1 association with DDBE-*osk* 3'UTR 2+3 (Figure 7d, e), indicating that Tm1 is
linked to the dynein machinery via *osk* RNA rather than through protein-protein interactions.
This observation is consistent with the finding that Tm1 does not directly influence DDBE
activity.
326
We next assessed the behavior of Khc in the presence of DDBE-*osk* 3'UTR 2+3 RNPs but
without Tm1. In these conditions, dynein and associated *osk* RNA exhibited long movements
toward the minus ends of microtubules (Figure 8a), as shown by polarity marking of the tracks
(Figure S10). Of the relatively few Khc runs that were observed, only ~25 % were directed
towards the microtubule plus end and those were seldomly associated with *osk* RNA (Figure
8a-c and Table S5). The remainder moved towards the minus end in association with dynein
and the RNA (Figure 8a-c), indicating some capacity of Khc to associate with DDBE-*osk* RNA
complexes. Khc in these complexes appeared to retain some activity, demonstrated by
occasional plus-end-directed movements after dissociating from DDBE-*osk* RNPs
translocating in the minus end direction (9/127 complexes, 7.1 %; Figure S11). Inclusion of
Tm1 in the assay caused a 12-fold increase in the incidence of minus-end-directed Khc
transport events (Figure 8a, b and Table S5), which overlapped with dynein and *osk* RNA
signals (Figure 8c). These data reveal that Tm1 promotes formation of dual-motor *osk* RNPs
in which dynein activity is dominant over that of kinesin, resulting in an overwhelming minus-
end-directed bias in Khc movement. Treatment of these *osk* RNPs with RNase A strongly
suppressed the Tm1-induced association of Khc with minus-end-directed DDBE-
osk 3'UTR 2+3 RNA complexes (Figure S12), supporting the notion that Tm1 achieves this
effect through interactions with *osk* RNA.
345
To determine if Tm1 additionally suppresses the activity of Khc within minus-end-directed
DDBE-*osk* RNPs, we compared dynein velocities and the distances traveled by Khc
toward microtubule minus ends in the presence and absence of Tm1. Consistent with an
inhibitory effect on Khc activity, Khc-bound DDBE-*osk* RNA complexes moved ~50 %
more rapidly towards the minus end in the presence of Tm1 (Figure 8d and Table S5).
Moreover, Tm1 allowed Khc to associate with minus-end-directed DDBE-*osk* RNA
complexes for distances on microtubules that were ~65 % greater than in its absence
(Figure 8e and Table S5), and significantly reduced the incidence of Khc directional
switches (6/848 complexes, 0.7 %; $p < 0.0001$ compared to in the absence of Tm1 (Fisher's
exact test)). These observations suggest that Tm1 limits engagements of Khc with

microtubules that counter minus-end-directed motion. Collectively, these results support a

model in which Tm1 recruits Khc to the DDBE-*osk* complex in an inactive state thus avoiding

a tug-of-war between the opposite polarity motors.

Discussion

Tm1-I/C, an atypical RNA-binding isoform of tropomyosin 1, was recently implicated as an RNA adaptor for kinesin-1 transport[41,42]. By interacting directly with Khc's cargo-binding domain, Tm1 was shown to enhance RNA binding by Khc[43]. Our current study has revealed a previously unknown role of Tm1 in *osk* mRNP transport. We show that Tm1 negatively regulates Khc activity, which we propose occurs by inducing a conformational change in the Khc stalk domain that stabilizes the inhibitory Khc motor-tail interaction (Figure 9a). We demonstrate that Tm1 also links Khc in an inactive state to the dynein transport machinery. This work involves the first reconstitution of a native cargo-adaptor-motor complex that recapitulates regulation of both dynein and kinesin motors. *In vivo*, the mechanism we uncover would allow dynein-mediated transport of *osk* RNPs from the nurse cells to the oocyte to proceed without a tug-of-war with Khc whilst positioning the plus-end-directed motor on the RNPs for when it needs to drive their posteriorward movement within the mid-oogenesis oocyte (Figure 9b).

Kinesin-1 autoinhibition mechanisms

Despite decades of research, kinesin-1 autoinhibition is not fully understood. This is due in part to a paucity of structural information for full length kinesin-1. Whereas crystal structures of kinesin motor domains in different states are available, the conformations of the coiled-coil stalk in both the active and autoinhibited state of the motor remain elusive. While an extended conformation of the coiled-coil stalk was proposed to exist in active kinesin-1[59], this notion was challenged by subsequent studies that indicated an at least partially folded structure of the tail within the active complex[21,60]. A recent study involving structure prediction and single-particle negative stain electron microscopy questions if folding occurs at the flexible hinge region 2 and suggests a different conformation for the Khc stalk[61].

It has been proposed that the interaction of the IAK motif within the tail with the motor domain plays a key role in kinesin-1 autoinhibition. Consistent with this notion, we observed hyperactivity of *Drosophila* Khc motility when the IAK motif was deleted. However, it was recently shown that Klc inhibits Khc activity independently of the IAK-motor interaction[22]. We also find that the IAK motif is not needed for inhibition of Khc by Tm1. Instead, we find that the Khc AMB domain, a region outside the IAK motif previously shown to be essential for *osk* mRNA localization[62], contributes to inhibition in Khc-Tm1 complexes. Our work therefore identifies a novel role for the AMB domain.

Using NMR spectroscopy, we show that the IAK-motor interaction does not interfere with binding of regulating factors such as Tm1. Our SAXS data-driven structure modeling provides

insights into possible conformations of a full length kinesin-1. The Khc stalk adopts different
conformations when in complex with Tm1 (Figure 6c, d). This could explain why the IAK-motor
domain interaction, despite being necessary for autoinhibition, is insufficient to fully inhibit Khc.
It is conceivable that additional interactions in the rearranged, Tm1-bound stalk regions are
required to maintain the autoinhibited, folded conformation. This is in line with our own and
previous observations that the IAK-motor domain interaction is rather weak[17] and the report in
a recent preprint that interactions within the coiled-coil stalk and between stalk and motor
contribute to kinesin-1 autoinhibition[63]. Presumably, binding of Tm1 drives structural
rearrangements of the stalk that compact the molecule and stabilize the autoinhibited
conformation to robustly shut down Khc motility. We propose a model in which several weak
interactions – the motor-IAK interaction and interactions within the Tm1-bound coiled-coil stalk,
including the AMB domain – act synergistically to achieve this stable inhibited conformation of
Khc *in vivo* (Figure 9a).
409
With its functions in cargo-binding and motor regulation, we speculate that Tm1 acts as an
alternative Klc in the *Drosophila* germline to regulate processes which were previously thought
not to require a Klc[40]. Previously, PAT1 was shown to act as a positive regulator of Khc and
have redundant functions with Klc in the *Drosophila* germline [39]. As correct RNA localization
during oogenesis is critical for development, positive regulators such as PAT1 and negative
regulators such as Tm1 may have replaced Klc in order to provide nuanced regulation of Khc
activity. In the future, it will be interesting to determine whether Klc is also dispensable in other
tissues that use Tm1 for RNA localization, including the follicle cells of the egg chamber[64] and
the larval neuromuscular junction[65].
419

Co-ordination of kinesin-1 and dynein motors on *osk* RNPs

Although many cargo types have been shown to be transported by the action of dynein and
kinesin-1 motors, the mechanisms of motor regulation in the context of dual motor transport
have been elusive. We have identified one of very few known examples of a factor that links
dynein and kinesin-mediated transport. Furthermore, to our knowledge, Tm1 is the first
example of a protein that modulates dynein and kinesin behavior in a reconstituted dual-motor
assembly on a native cargo.
427

Several studies have reported that active dynein and kinesin motors engage in a tug-of-war
when artificially coupled[66,67,68,69,70,71]. However, the relevance of such mechanical antagonism
in localizing physiological cargoes for these motors is unresolved[6]. Although we also observe
motor opposition in reconstituted RNPs containing DDBE and Khc, our observation that Tm1
supports efficient dynein-mediated RNA localization through robust inhibition of Khc highlights

the importance of regulatory factors in addition to mechanical coupling in cargo-adaptor-motor
complexes.

Despite the potential for antagonistic engagement of dynein and kinesin on shared cargoes,
the motors can exhibit a puzzling co-dependence, in which strongly impairing the activity of
one motor type can inhibit the activity of the other[6]. Nonetheless, enhanced inhibition of Khc
by Tm1 in our reconstituted RNPs demonstrates the importance of negative kinesin regulation
in enabling robust dynein activity. Supporting the *in vivo* relevance of this regulatory logic,
kinesin-1 activating IAK mutations were recently shown to impair dynein-mediated transport
processes in *Aspergillus nidulans*[72]. Collectively, these observations point to complex interplay
between opposite polarity microtubule motors during bidirectional transport. Further
reconstitutions of dual-motor systems on native cargoes should reveal generalities of dynein-
kinesin crosstalk, as well as any cargo-specific mechanisms.

Potential kinesin-1 activation mechanisms in the female germline

Our study provides mechanistic insight into two critical aspects of *osk* mRNA transport – the
assembly of the dual motor complex and how Khc activity is suppressed during dynein-
mediated delivery of the transcript from the nurse cells to the oocyte. However, it is not known
how Tm1-mediated inhibition of Khc is alleviated at later stages of oogenesis in the oocyte.
One candidate to fulfill this role is Ensconsin. This protein is required for posterior *osk*
localization[73], as well as organelle transport in *Drosophila* neurons[74] and is homologous to
MAP7, which was recently shown to be required for full activity of the human kinesin-1 complex
in vitro[22,75]. Other factors that may relieve Khc inhibition in the oocyte include the exon junction
complex (EJC), which, together with the SOLE RNA structure, is essential for *osk* transport to
the posterior of the oocyte[46,76]. As Tm1 remains bound to the *osk* mRNP throughout transport
and is required for Khc-mediated translocation of the mRNA to the posterior of the oocyte[42,43],
it is likely that the activating factor(s) induces a conformational change in the Khc-Tm1 complex
rather than dissociation of Tm1. Investigating the molecular mechanisms that underpin these
dynamics is likely to inform our understanding of how kinesin-1 activity is orchestrated in other
transport systems.

Figure 1: A TIRF microscopy-based *in vitro* motility assay reveals an inhibitory effect of Tm1 on Khc motility. a Scheme of domain architecture of *Drosophila* Kinesin heavy chain (Khc) and atypical Tropomyosin 1 isoform I/C (Tm1). **b** Scheme of *in vitro* motility assay. Preincubated Khc, Tm1 and RNA mixes were diluted to 6.25 nM of each component, and injected into an imaging chamber with pre-immobilised microtubules on a passivated glass surface. **c** Example kymographs showing processive movements of AlexaFluor647-labeled SNAP-Khc FL (labeling efficiency: 79.75 %), Cy3-labeled *osk* 3'UTR and AlexaFluor488-labeled Tm1 FL (labeling efficiency: 57 %). Microtubule plus and minus ends are oriented toward the right and left of each kymograph, respectively. **d** Fraction of Khc transporting *osk* 3'UTR in absence and presence of AlexaFluor488-labeled Tm1 in the assembly mix. Data points represent individual microtubules (84 and 36 complexes analyzed per condition). Statistical significance was evaluated with an unpaired t-test. ns = not significant. **e** Histograms of velocity distributions of Khc

475 in absence or presence of Tm1. Statistical significance was evaluated with an unpaired t-test.
476 ****p<0.0001.

Figure 2: Tm1 inhibits Khc motility *in vitro* via Khc's regulatory tail. a, b Example kymographs showing processive movements of AlexaFluor647-labeled SNAP-Khc910 (a) and Khc940 (b) ± AlexaFluor488-labeled Tm1 FL. Microtubule plus and minus ends are oriented toward the right and left of each kymograph, respectively. **c, d** Number of processive events of Khc FL, Khc940 and Khc910 (c) and fraction of microtubule-bound Khc complexes that are processive ± Tm1 FL in the assembly mix

(d). Data points show values for individual microtubules (between 36 and 438 complexes analyzed per
condition). Error bars: SD. Statistical significance was evaluated with an unpaired t-test. ns = not
significant, **$p < 0.01$, *$p < 0.05$. Labeling efficiencies for Khc FL, Khc910 and Khc940 were 79.8 %, 54 %
and 78 %, respectively, so that total run numbers are underestimated. **e** Average velocities of Khc FL
and Khc910 ± Tm1 FL in the assembly mix. Data points represent individual runs. Statistical significance
was evaluated with an unpaired t-test. ****$p < 0.0001$.

Figure 3: Khc AMB domain mediates Tm1 inhibition of Khc. a Example kymographs showing processive movements of AlexaFluor647-labeled SNAP-KhcΔAMB ± AlexaFluor488-labeled Tm1 FL in the assembly mix. Microtubule plus and minus ends are oriented toward the right and left of each kymograph, respectively. **b, c** Processive events of Khc FL, Khc910 and KhcΔAMB ± Tm1 FL (b) and fraction of processive Khc events (c). Data points show values for individual microtubules (between 24 and 595 complexes analyzed per condition). Error bars: SD. Statistical significance was evaluated with an unpaired t-test. ****p<0.0001, ***p<0.001, **p<0.01. Labeling efficiencies for Khc FL, Khc910 and KhcΔAMB were 79.8 %, 54 % and 78.8 %, respectively, so that total run numbers are underestimated. **d** Average velocities of Khc FL, Khc910 and KhcΔAMB ± Tm1 FL. Data points represent individual runs. Statistical significance was evaluated with an unpaired t-test. ****p<0.0001.

Figure 4: The Tm1 N-terminal region is required for Khc inhibition. a Example kymographs showing processive movements of AlexaFluor647-labeled SNAP-Khc FL ± AlexaFluor488-labeled Tm1 FL or Tm1 1-335 in the assembly mix. **b, c** Processive events of Khc FL ± Tm1 FL or Tm1 1-335 in the assembly mix (b) and fraction of processive Khc events (c). Data points show values for individual microtubules (between 44 and 335 complexes analyzed per condition). Error bars: SD. Statistical significance was evaluated with an unpaired t-test. ****p<0.0001, ***p<0.001. **d** Average velocities of Khc FL ± Tm1 FL or Tm1 1-335 in the assembly mix. Data points represent individual runs. Statistical

510   significance was evaluated with an unpaired t-test. ****$p < 0.0001$. **e** Example kymographs showing

511   processive movements of AlexaFluor647-labeled SNAP-Khc FL ± Tm1 FL or Tm1 252-334. **f** Processive

512   events of Khc FL ± Tm1 FL or Tm1 252-334 in the assembly mix. Error bars: SD. Statistical significance

513   was evaluated with an unpaired t-test with Welch's correction. **$p < 0.01$. Microtubule plus and minus

514   ends are oriented toward the right and left of each kymograph, respectively (a and e).

Figure 5: Tm1 binding affects the free and motor-bound regulatory tail of Khc. a ^{1}H,^{15}N-HSQC spectrum of the Khc tail construct Khc 855-975 with backbone assignments indicated. **b, c** ^{1}H,^{15}N-HSQC titration experiment with U10 RNA and Tm1 1-335. Overlaid spectra (b) and plotted intensity ratios of individual residues. RNA was added before Tm1. **d** Intensity ratios of a titration experiment with Tm1 1-335 and U10 RNA in which Tm1 was added before RNA. **e** Intensity ratios of a titration experiment with

522 U10 RNA and Tm1 252-334 in which RNA was added before Tm1. **f - h** ^{1}H,^{15}N-HSQC titration
523 experiment of the Khc tail Khc 855-975 with the motor domain Khc 1-365 and Tm1 1-335. Overlaid
524 spectra (f), plotted intensity ratios (g) and chemical shift perturbation plot (h) of individual residues.
525 Khc 1-365 was added before Tm1 1-335.

Figure 6: Khc conformation is altered upon Tm1 binding. a Experimental SAXS curves overlaid with theoretical SAXS curves of best Crysol fits. **b** Dimensionless Kratky plot of scattering profiles of Khc and the Khc-Tm1 complex compared to a globular protein (lysozyme), a disordered protein with short secondary structure elements (SASDB[57] CY4) and a fully disordered extended protein (SASDB[57] A57) to illustrate the differences between Khc and Khc-Tm1 and the contributions of ordered and disordered regions. **c** Best fits for Khc and for Khc-Tm1. Khc models are shown in shades of blue with the AMB domains marked in red, Tm1 is shown in pink. cc 1-4 denote the four coiled-coil regions of the Khc stalk domain. **d** Schemes showing the orientations of Khc's stalk coiled coils in the structures with best Crysol fits. Angles between coiled coils are averages from the five best-fitting models.

Figure 7: Tm1 does not affect dynein activity in *osk* RNPs. a Kymographs showing translocation of dynein and *osk* 3'UTR 2+3 in the presence and absence of Tm1 as part of minimal minus-end-directed RNPs containing dynein, dynactin, BicD, and Egalitarian (DDBE). Co-translocation of Tm1 is observed when present in the assay. **b, c** Average velocities (b) and run lengths (c) of DDBE-*osk* 3'UTR 2+3 complexes assembled with and without Tm1. 1-cumulative frequency distributions of measured run lengths were fit as one-phase decays with the decay constant τ ± 95 % confidence interval shown. Hollow triangles represent empirical bin values used for fitting. **d** Kymographs showing reduced association of Tm1 with motile DDBE-*osk* 3'UTR 2+3 RNPs after treatment with RNase A. **e** Fraction of moving DDBE-*osk* 3'UTR 2+3 RNPs that colocalized with Tm1 signal with and without RNase A

547 treatment. Dots represent values for individual complexes (b) or average values of all motility on
548 individual microtubules (e). Microtubule plus and minus ends are oriented toward the right and left of
549 each kymograph, respectively (a and d). For all plots, mean ± SD is shown. Statistical significance was
550 determined by a Kruskal-Wallis ANOVA test (b), Mann-Whitney test (c), or Welch's t-test (e). ns: not
551 significant. ****: $p < 0.0001$.

Figure 8: Tm1 promotes association of inhibited Khc with minus-end-directed DDBE-*osk* RNPs.
a Kymographs showing motile behavior of DDBE (via TMR-dynein), AlexaFluor647-Khc, and AlexaFluor488 *osk* 3'UTR 2+3 in the presence and absence of unlabeled Tm1. Plus- and minus-end-directed Khc movements in the absence of Tm1 are indicated by white and yellow arrowheads, respectively. Microtubule plus and minus ends are oriented toward the right and left of each kymograph, respectively. **b** Frequency of processive Khc movements towards microtubule minus ends (towards left of x-axis, purple) and plus ends (towards right of x-axis, magenta) in the presence and absence of Tm1. **c** Cumulative fraction of plus- and minus-end-directed Khc signals that co-migrated with all combinations of other labeled assay components in the presence and absence of Tm1. **d** Average

velocities of minus-end-directed DDBE-*osk* 3'UTR 2+3 complexes with co-migrating Khc signal in the presence and absence of Tm1. **e** Distance traveled by Khc towards the minus end in the presence and absence of Tm1. 1-cumulative frequency distributions of measured run lengths were fit as one-phase decays with the decay constant $\tau \pm 95\,\%$ confidence interval shown. Hollow triangles represent empirical bin values used for fitting. Dots represent average values for all motility on individual microtubules (b) or values for individual complexes (d). For all plots mean ± SD is shown. Statistical significance was determined by Welch's t-test (b) or Mann-Whitney test (d, e). ****: p<0.0001.

Figure 9: a Model of Tm1-mediated stabilization of the auto-inhibited conformation of Khc. Whereas both the motor domain-IAK peptide interaction and interactions within the Tm1-bound, rearranged Khc stalk region, including the AMB domain, can inhibit Khc partially, synergy between these weak interactions stabilizes the inhibiting head-tail interaction. **b** Model of Tm1's role in dynein-mediated transport of *osk*. By linking Khc to the dynein-transported *osk* RNA and keeping Khc inhibited, Tm1 allows the first, dynein-mediated transport step from the nurse cells to the oocyte to proceed without a tug-of-war between the motors. These events also position Khc on the RNP for the second, Khc-mediated transport step.

Materials and Methods

Molecular Cloning

Khc FL for *in vitro* studies was cloned with a HisSUMO-SNAP-3C-tag into MCS1 between BamHI and HindIII sites of the pFastBacDual vector by InFusion cloning (Takara). The Khc mutant constructs used for *in vitro* studies were generated by site-directed mutagenesis of the Khc FL plasmid using primers that either inserted a stop codon after aa 910 (Khc910) or aa 940 (Khc940) or deleted aa 911-938 (KhcΔAMB) followed by DpnI digest of the template plasmid and transformation of the product in *E. coli* XL1-blue cells. For co-expression of Khc FL and Tm1, the Tm1 coding sequence was inserted into MCS2 of pFastBacDual-HisSUMO-SNAP-3C-Khc between SmaI and KpnI sites by InFusion cloning (Takara). Tm1 FL for motility assays was cloned with a HisSUMO-SNAP-tag between SacI and HindIII sites of the pET11 vector by InFusion cloning (Takara). The SNAP-Tm1 1-335 truncation was made by site-directed mutagenesis of the SNAP-Tm1 FL plasmid using primers that inserted a stop codon after aa 335 as described above.

Protein expression and purification

His6-SUMO-SNAP-3C-tagged Khc FL, Khc910 or ΔIAK and ΔAMB truncations and co-expressed Khc-Tm1 complexes were expressed using the Bac-to-Bac Baculovirus expression system (Gibco). 1 L of *Sf21* insect cells at 1×10^6 cells mL^{-1} shaking culture were infected with 10 mL of P1 virus stock. After expression for 3 d at 27.5 °C, cells were harvested by centrifugation at 1000 g for 20 min at 4°C. The pellets were lysed in a dounce tissue grinder in lysis buffer (20 mM Tris-HCl at pH 7.5, 500 mM NaCl, 1 mM MgCl$_2$, 0.1 mM ATP, 2 mM DTT, 5 % glycerol, 0.2 % Tween-20). The lysates were cleared by centrifugation and the soluble protein fraction was affinity-purified on a HisTrap Excel column (GE Healthcare). After elution in a 0–300 mM imidazole gradient, the HisSUMO-fusion tag was cleaved either by Senp2 (SNAP-tagged proteins for fluorescent labeling) or 3C protease (unlabeled proteins) digest during dialysis in GF150 buffer (25 mM HEPES/KOH pH 7.3, 150 mM KCl, 1 mM MgCl$_2$, 0.1 mM ATP, and 2 mM DTT) for 16 h before further purification by anion exchange chromatography on a HiTrap Q HP column (GE Healthcare), followed by size exclusion chromatography (SEC) on a Superdex 200 10/300 Increase column (GE Healthcare).

His6-SUMO-tagged and His6-SUMO-SNAP-tagged Tm1 protein constructs and GST-Khc 1-365 were expressed in BL21-CodonPlus(DE3)-RIL cells (Stratagene) by isopropyl β-D-1-thiogalactopyranoside (IPTG) induction for 16 h at 18 °C. Cells were grown in Luria-Bertani (LB) medium (0.5 % [w/v] yeast extract [MP], 1 % [w/v] tryptone [MP], 0.5 % [w/v] NaCl at pH 7.2) supplemented with antibiotics (100 μg mL^{-1} ampicillin or 10 μg mL^{-1} kanamycin and

34 µg mL^{-1} chloramphenicol). After harvesting, the pellets were lysed by a microfluidizer

processor (Microfluidics) in 20 mM Tris-HCl (pH 7.5), 500 mM NaCl, 5 % glycerol, 0.01 % NP-

40, and 40 mM imidazole buffer supplemented with protease inhibitor cocktail (Roche) and

2 mM DTT. The lysates were subsequently cleared by centrifugation. HisSUMO-tagged

proteins were affinity-purified by Ni-IMAC on a HisTrap column (Cytiva) and eluted with an

imidazole gradient (40–300 mM). After cleavage of the His6-SUMO-tag with Senp2 protease,

cleaved tags and protease were removed by a second, subtractive Ni-IMAC on a HisTrap

column (GE Healthcare). The proteins were further purified on either a 10/300 Superose 6

Increase GL or a HiLoad 16/600 Superdex 200 SEC column in either GF150 buffer for motility

assays or NMR buffer (25 mM NaPO$_4$ pH 6.5, 150 mM NaCl, and 0.5 mM TCEP) for NMR

experiments.

GST-tagged Khc 1-365 was affinity-purified on a GSTrap column (Cytiva) and glutathione

gradient (1-20 mM). The protein was further purified on a HiLoad 16/600 Superdex 200 SEC

column in GF150 buffer.

SNAP-tagged Khc and Tm1 proteins were fluorescently labeled with SNAP-Cell TMR Star,

SNAP-Surface Alexa Fluor 488 or SNAP-Surface Alexa Fluor 647 at 15 °C for 90 min in GF150

buffer with a 1.5x excess of dye. Free dye was removed by a subsequent desalting step on a

PD-10 desalting column (Cytiva). Labeling efficiencies were determined by measuring protein

concentration at 280 nm and dye concentration at the dye's excitation maximum. For

monomeric protein, labeling efficiency was calculated as $c_{dye}/c_{protein}$. For dimeric proteins,

labeling efficiency was calculated based on monomer labeling efficiencies as $(L_1 \times U_2) + (U_1 \times L_2) + (L_1 \times L_2)$

$L_2) + (L_1 \times L_2)$ with L1 = fraction of monomer 1 labeled, L2 = fraction of monomer 2 labeled, U1

= fraction of monomer 1 unlabeled, U2 = fraction of monomer 2 unlabeled.

Recombinant human dynein and *Drosophila* Egl-BicD complexes were expressed in *Sf9* insect

cells with the dynein complex fluorescently labeled via a SNAPf tag fused to the N-terminus of

Dynein heavy chain, as described previously[15]. Native porcine dynactin was purified from brain

tissue as described previously[15].

RNA *in vitro* transcription

The *osk* RNA constructs used for the motility assays were synthesized *in vitro* using the

MEGAscript T7 Transcription Kit (Ambion). The RNA was transcribed from a PCR product

amplified with a T7 promote-containing forward primer and containing an *osk* mRNA FL, *osk*

3'UTR or a ~529-nt region of the *osk* 3'UTR previously defined as region 2+3 (*osk* 3'UTR 2+3)

that is sufficient to promote the localization of reporter transcripts to the developing oocyte[48].

The template for *osk* min was generated from three overlapping primers by PCR and

subsequently amplified again with a T7 promoter containing forward primer. RNA was
transcribed from the final PCR product. For fluorescent labeling of the *osk* 3'UTR and *osk* min
transcripts, Cy3- or Cy5-UTP was included in the transcription reaction with a 4-fold excess of
unlabeled UTP, yielding transcripts with several labels per RNA molecule. For *osk* 3'UTR 2+3,
transcripts were fluorescently labeled with AlexaFluor488-UTP with a 5-fold excess of
unlabeled UTP. Transcripts were subsequently purified by phenol/chloroform extraction and
passed through a G50 mini Quick Spin RNA desalting column (Sigma-Aldrich) to remove
unincorporated nucleotides before sodium acetate/ethanol precipitation. The precipitated RNA
was resuspended in nuclease-free water and stored at -20 °C or -70 °C.

Polymerization and stabilization of microtubules

Microtubules were polymerized from porcine tubulin (Cytoskeleton Inc., Denver, CO) and
labeled with fluorophores and biotin by stochastic incorporation of labeled dimers into the
microtubule lattice. Mixes of 2.5 µM unlabelled tubulin, 0.17 µM Hilyte488- or AMCA-tubulin,
and 0.3 µM biotin-tubulin were incubated in BRB80 (80 mM PIPES pH 6.85, 2 mM $MgCl_2$,
0.5 mM EGTA, 1 mM DTT) with 0.67 mM GMPCPP (Jena Bioscience, Jena, Germany)
overnight at 37 °C. Polymerized microtubules were pelleted in a room temperature table top
centrifuge at 18,400 x g for 8.5 min, and washed once with pre-warmed (37 °C) BRB80. After
pelleting once more, the microtubules were gently resuspended in 300 µL pre-warmed (37 °C)
BRB80 containing 20-40 µM paclitaxel (taxol; Sigma-Aldrich).
672
Polarity marked microtubules with HiLyte 488 tubulin enriched at minus ends were made
according to the principle described previously[77]. Bright microtubule seeds were polymerized
at 37 ºC for 10 minutes in a mix containing 2.5 µM unlabelled porcine tubulin (5-fold excess
over fluorescent tubulin), 0.5 µM HiLyte 488 tubulin, and 0.3 µM biotin-tubulin in BRB80 with
0.67 mM GMPCPP. After 10 minutes, this mixture was diluted 1:1 with a tubulin polymerization
mix containing 12.5 µM unlabelled tubulin and 0.3 µM biotin-tubulin in BRB80 with 0.67 mM
GMPCPP for final tubulin concentrations of 7.5 µM unlabelled tubulin (30-fold excess over
fluorescent tubulin), 0.25 µM HiLyte 488 tubulin, and 0.3 µM biotin-tubulin. The final tubulin
mix was incubated at 37 ºC for an additional 2 h to allow growth of relatively dim microtubule
plus ends before harvesting and taxol stabilization as described above.

In vitro single-molecule reconstitution assays, TIRF microscopy and image analysis

TIRF-based single-molecule reconstitution assays were performed as described previously[15].
Briefly, taxol-stabilized porcine microtubules were immobilized in imaging flow chambers by
streptavidin-based linkages to biotin-PEG passivated cover slips. Khc RNP assay mixes were

assembled at 500 nM of each component. Assemblies of the dynein transport machinery with
Khc and Tm1 contained 100 nM dynein, 200 nM dynactin, 500 nM Egl-BicD, 1 µM RNA,
500 nM Khc (or Khc storage buffer as a control) and 2 µM Tm1 (or Tm1 storage buffer as a
control), and were incubated on ice for 1-2 h prior to imaging. The RNP assay mixes were then
diluted to concentrations suitable for imaging of single molecules (6.25 nM for Khc RNP
experiments, 5 nM dynein and 25 nM Khc for experiments containing both motors) in BRB12
(Khc experiments) or modified dynein motility buffer[58] (30 mM HEPES-KOH pH 7.0, 50 mM
KCl, 5 mM $MgSO_4$, 1 mM EGTA pH 7.5, 1 mM DTT, 0.5 mg mL^{-1} BSA - experiments including
DDBE) supplemented with 1 mg mL^{-1} α-casein, 20 µM taxol, 2.5 mM MgATP and an oxygen
scavenging system (1.25 µM glucose oxidase, 140 nM catalase, 71 mM 2-mercaptoethanol,
25 mM glucose final concentrations) to reduce photobleaching, and applied to the flow
chamber for imaging by TIRF microscopy.
700
For assays in which RNase A was added, assemblies were diluted into modified dynein motility
buffer as described above with either 1 µL of 0.2 µg/mL RNase A (final concentration of
9.5 ng/mL) diluted in motility buffer, or 1 µL of motility buffer alone added in addition. Such low
concentrations of RNase A were used to ensure mild digestion of the RNA (which was
monitored during acquisition via the fluorescent RNA signal) that would eliminate any RNA-
based connections between protein components whilst limiting digestion of the predominantly
double-stranded localization signal important for activating DDBE motility[48]. These mixes were
incubated at room temperature for 5 min before application to the imaging chambers.
709
710   For each chamber, a single multi-color acquisition of 500 frames was made at a frame rate of
711   ~2 frames s^{-1} and 150 ms exposure (1-2 colors) or 1.2-1.4 frames s^{-1} and 50 ms exposure (3
712   colors). Images of Khc-Tm1 were acquired using a Leica GSD TIRF microscope system. All
713   images involving the dynein machinery were acquired on a Nikon TIRF system using
714   Micromanager control software[78] as described previously[15].
715
Binding and motility of RNP components in the *in vitro* reconstitution assays were manually
analyzed by kymograph in FIJI[79] as described previously[15]. Statistical analysis and plotting
were performed using GraphPad Prism version 9.1.1 for MacOS, (GraphPad Software, San
Diego, USA).
As we noticed some variability of Khc activities between different assays, we repeated all
appropriate control conditions for each experiment on the same day and used these
controls as comparison for our analysis. Results were confirmed on multiple days in
duplicates or triplicates of the experiments.

Microtubule binding assays

For microtubule binding assays, microtubules were polymerized from 5 mg mL^{-1} porcine tubulin (Cytoskeleton Inc., Denver, CO) at 37 °C for 20 min in BRB80 supplemented with 1 mM GTP and then diluted 1:10 in BRB80 + 20 µM taxol. Proteins of interest were incubated at indicated concentrations with 20 µL of this microtubule solution for 30 min at RT in a final volume of 50 µL, before soluble and microtubule-bound fractions were segregated by ultracentrifugation at 80,000 g for 30 min over a 500 µL 30 % sucrose cushion. Soluble fractions were supplemented with 1x SDS-loading dye. Microtubule pellets were washed with BRB80 + taxol before redissolving in 50 µL 1x SDS-loading dye. Soluble and pellet fractions were analyzed by SDS-PAGE.

GST-pulldown competition assay

To test whether Tm1 competes with the Khc motor-tail interaction, 5 µM GST-Khc 855-975 as bait and an indicated excess of prey protein Khc 1-365 were incubated with increasing concentrations of Tm1 1-335 as competitor for Khc 855-975 binding in a total volume of 100 µL in GF150 buffer (25 mM HEPES/KOH pH 7.3, 150 mM KCl, 1 mM MgCl$_2$, 2 mM DTT). After incubation for 40 min on ice, the mixture was added to pre-equilibrated 50 µL Glutathione Sepharose 4B beads (GE Healthcare) and proteins were allowed to bind for 2 h at 4 °C on a nutating shaker. Beads were washed with 4 x 500 µL and 1 x 100 µL GF150 + 0.01 % NP-40. Bound proteins were eluted from the beads by boiling in 1x SDS-loading dye for 5 min at 95 °C. Input, wash and elution samples were analyzed by SDS-PAGE.

Mass photometry

Mass photometry data were acquired on a ReFeyn OneMP from co-expressed, co-purified Khc-Tm1 complexes at a sample concentration of 20 nM in PBS. Data acquisition was performed using AcquireMP, with data analysis performed in DiscoverMP (Refeyn Ltd, v1.2.3).

NMR spectroscopy

NMR measurements were performed at 298 K on a Bruker Avance III NMR spectrometer with magnetic field strengths corresponding to proton Larmor frequencies of 600 MHz equipped with a cryogenic triple resonance gradient probe head. NMR sample concentration was 50 µM for titration experiments and 177 µM for backbone assignment experiments. Experiments for backbone assignments were performed on ^{13}C,^{15}N-labeled samples using conventional triple-resonance experiments (HNCO, HNCA, CBCA(CO)NH, HN(CO)CA and HNCACB)[80]. All spectra were acquired using the apodization weighted sampling scheme[81] and processed using NMRPipe[82]. Resonance assignments were done with the program Cara.

758 For titrations, RNA and Tm1 were added to [15]N-labeled Khc 855-975 at indicated ratios and a
759 [1]H,[15]N-HSQC was recorded for each titration point. Peak intensity ratios were derived using
760 NMRView[83] and corrected for dilution. The extent of amide [1]H-[15]N chemical shift perturbations
761 (CSPs) in free versus bound Khc 855-975 were calculated according to Williamson[84] to
762 compensate for frequency range differences between [15]N and [1]H dimensions.

SAXS data acquisition and structure modeling

SAXS data on full length Khc in isolation and in complex with full length Tm1 in 25 mM
HEPES/KOH pH 7.3, 150 mM KCl, 1 mM MgCl$_2$ and 2 mM DTT at concentrations of 0.5, 1.0
and 2.0 mg mL^{-1} were recorded at 20 °C on BM29 beamline, ESRF Grenoble, France. Ten
frames with an exposure time of 1 s/frame were recorded in batch mode using an X-ray
wavelength of 0.992 Å and a sample distance of 2.81 m to the Pilatus2M detector. Data
reduction and buffer subtraction was performed by the automatic processing pipeline[85]. R_g and
$I(0)$ were determined using the Guinier approximation using Primus/autorg[86].
771
The scattering curve was subsequently compared to pools of structures generated by
connecting AlphaFold2[55,56] models of the Khc head domain (residues 1-378) homodimer
connected to four stretches of coiled-coil homodimers (residues 403-585, 604-706, 711-838
and 844-936 respectively). The linker regions between the structured parts were modified by
random phi/psi rotations in monomer A of the homodimer followed by a short four-step
simulated annealing energy minimization to restore proper bond geometries in monomer B,
for which all corresponding residues were rotated together with monomer A during the linker
randomization. The C-termini of monomer A and monomer B were subsequently randomized
independently. A total of 5000 structures were generated in this way. In a second step, the
structure of the motor homodimer was replaced by a homodimer (residues 1-344) bound to a
peptide from the C-terminus of molecule A (residues 934-953) containing the IAK motif, which
was followed by simulated annealing energy minimization to restore proper bond geometry.
During the simulated annealing energy minimization steps, the structures of the motor and
coiled-coil models or motor/peptide were unaltered by using a harmonic energy term to keep
the coordinates close to the starting coordinates in the template structure[87]. The calculations
were performed using CNS1.2[88] within the ARIA1.2 framework[89]. Structure pools for the Khc
homodimer in complex with Tm1 were generated in a similar way with an AlphaFold2-modeled
α-helix of Tm1 (residues 258-386) bound to the last coiled-coil stretch of Khc (residues 844-
936) in a position derived from the X-ray structure of the Khc-Tm1 complex[43] and additional
randomization of all residues outside the α-helix of Tm1. Each of the 2x5000 models were
fitted to the SAXS scattering intensities using CRYSOL[86].

Competing interest statement

The authors declare no competing interests.

Acknowledgements

We thank Lyudmila Dimitrova-Paternoga for reagents, the EMBL Protein Expression and Purification Core Facility, especially Karine Lapouge, the EMBL Advanced Light Microscopy Facility, especially Marko Lampe, and the EMBL Chemical Biology Core Facility, especially David Will, for their support. We thank the Bio-SAXS beamline at European Synchrotron Radiation Facility Grenoble, BM29. We thank Petra Pernot (ESRF), Jérôme Kieffer (ESRF) and Cy Jeffries (EMBL Hamburg) for discussions. S.H. was supported by the EMBL Interdisciplinary Postdoctoral fellowship (EIPOD) Programme under Marie Curie Cofund Actions MSCA-COFUND-FP (664726) and Deutsche Forschungsgemeinschaft (DFG)-Forschergruppe 2333 grant (EP37/4-1) to A.E.. Work in the laboratories of J.H. and A.E. was supported by funding from the DFG via the priority program SPP1935 to J.H. and A.E. (EP37/3-1 and EP37/3-2) and the EMBL. Work in S.L.B.'s group is supported by the Medical Research Council (MRC), as part of United Kingdom Research and Innovation (also known as UK Research and Innovation) [MRC file reference number MC_U105178790]. M.A.M. is supported by a project grant from the BBSRC (BB/T00696X/1). For the purpose of the MRC open access policy, the authors have applied a CC-BY public copyright license to any Author Accepted Manuscript version arising.

Author contributions

S.H., M.A.M., A.E., and S.L.B. conceived the study. S.H., M.A.M. and B.S. performed the experiments. S.H., M.A.M., B.S and J.H. analyzed the data. S.L.B. and A.E. supervised the study. S.H. drafted the manuscript. All authors contributed to the writing of the manuscript.

References

1. Titus, M. A. Myosin-Driven Intracellular Transport. *Cold Spring Harb. Perspect. Biol.* **10**, (2018).

2. Doyle, M. & Kiebler, M. A. Mechanisms of dendritic mRNA transport and its role in synaptic tagging. *EMBO J.* **30**, 3540–3552 (2011).

3. Rodrigues, E. C., Grawenhoff, J., Baumann, S. J., Lorenzon, N. & Maurer, S. P. Mammalian Neuronal mRNA Transport Complexes: The Few Knowns and the Many Unknowns. *Front. Integr. Neurosci.* **15**, 692948 (2021).

4. Misgeld, T. & Schwarz, T. L. Mitostasis in neurons: maintaining mitochondria in an extended cellular architecture. *Neuron* **96**, 651–666 (2017).

5. Welte, M. A. Bidirectional transport along microtubules. *Curr. Biol.* **14**, R525-37 (2004).

6. Hancock, W. O. Bidirectional cargo transport: moving beyond tug of war. *Nat. Rev. Mol. Cell Biol.* **15**, 615–628 (2014).

7. McKenney, R. J., Huynh, W., Tanenbaum, M. E., Bhabha, G. & Vale, R. D. Activation of cytoplasmic dynein motility by dynactin-cargo adapter complexes. *Science* **345**, 337–341 (2014).

8. Schlager, M. A., Hoang, H. T., Urnavicius, L., Bullock, S. L. & Carter, A. P. In vitro reconstitution of a highly processive recombinant human dynein complex. *EMBO J.* **33**, 1855–1868 (2014).

9. Zhang, K. *et al.* Cryo-EM Reveals How Human Cytoplasmic Dynein Is Auto-inhibited and Activated. *Cell* **169**, 1303-1314.e18 (2017).

10. Reck-Peterson, S. L., Redwine, W. B., Vale, R. D. & Carter, A. P. The cytoplasmic dynein transport machinery and its many cargoes. *Nat. Rev. Mol. Cell Biol.* **19**, 382–398 (2018).

11. Olenick, M. A. & Holzbaur, E. L. F. Dynein activators and adaptors at a glance. *J. Cell Sci.* **132**, (2019).

12. Dienstbier, M., Boehl, F., Li, X. & Bullock, S. L. Egalitarian is a selective RNA-binding protein linking mRNA localization signals to the dynein motor. *Genes Dev.* **23**, 1546–1558 (2009).

13. Bullock, S. L. & Ish-Horowicz, D. Conserved signals and machinery for RNA transport in Drosophila oogenesis and embryogenesis. *Nature* **414**, 611–616 (2001).

14. Sladewski, T. E. *et al.* Recruitment of two dyneins to an mRNA-dependent Bicaudal D transport complex. *eLife* **7**, (2018).

15. McClintock, M. A. *et al.* RNA-directed activation of cytoplasmic dynein-1 in reconstituted transport RNPs. *eLife* **7**, (2018).

16. Lawrence, C. J. *et al.* A standardized kinesin nomenclature. *J. Cell Biol.* **167**, 19–22 (2004).

17. Stock, M. F. *et al.* Formation of the compact conformer of kinesin requires a COOH-terminal heavy chain domain and inhibits microtubule-stimulated ATPase activity. *J. Biol. Chem.* **274**, 14617–14623 (1999).

18. Kaan, H. Y. K., Hackney, D. D. & Kozielski, F. The structure of the kinesin-1 motor-tail complex reveals the mechanism of autoinhibition. *Science* **333**, 883–885 (2011).

19. Hackney, D. D. & Stock, M. F. Kinesin's IAK tail domain inhibits initial microtubule-stimulated ADP release. *Nat. Cell Biol.* **2**, 257–260 (2000).

20. Verhey, K. J. & Hammond, J. W. Traffic control: regulation of kinesin motors. *Nat. Rev. Mol. Cell Biol.* **10**, 765–777 (2009).

21. Cai, D., Hoppe, A. D., Swanson, J. A. & Verhey, K. J. Kinesin-1 structural organization and conformational changes revealed by FRET stoichiometry in live cells. *J. Cell Biol.* **176**, 51–63 (2007).

22. Chiba, K., Ori-McKenney, K. M., Niwa, S. & McKenney, R. J. Synergistic autoinhibition and activation mechanisms control kinesin-1 motor activity. *Cell Rep.* **39**, 110900 (2022).

23. Diefenbach, R. J., Mackay, J. P., Armati, P. J. & Cunningham, A. L. The C-terminal region of the stalk domain of ubiquitous human kinesin heavy chain contains the binding site for kinesin light chain. *Biochemistry* **37**, 16663–16670 (1998).

24. Verhey, K. J. *et al.* Light chain-dependent regulation of Kinesin's interaction with microtubules. *J. Cell Biol.* **143**, 1053–1066 (1998).

25. Pernigo, S., Lamprecht, A., Steiner, R. A. & Dodding, M. P. Structural basis for kinesin-1:cargo

recognition. *Science* **340**, 356–359 (2013).

26.   Fenton, A. R., Jongens, T. A. & Holzbaur, E. L. F. Mitochondrial adaptor TRAK2 activates and
functionally links opposing kinesin and dynein motors. *Nat. Commun.* **12**, 4578 (2021).

27.   Canty, J. T., Hensley, A. & Yildiz, A. Mitochondrial TRAK adaptors coordinate dynein and
kinesin motility. *BioRxiv* (2021) doi:10.1101/2021.07.30.454553.

28.   Arimoto, M. *et al.* The Caenorhabditis elegans JIP3 protein UNC-16 functions as an adaptor to
link kinesin-1 with cytoplasmic dynein. *J. Neurosci.* **31**, 2216–2224 (2011).

29.   Celestino, R. *et al.* JIP3 interacts with dynein and kinesin-1 to regulate bidirectional organelle
transport. *J. Cell Biol.* **221**, (2022).

30.   Kendrick, A. A. *et al.* Hook3 is a scaffold for the opposite-polarity microtubule-based motors
cytoplasmic dynein-1 and KIF1C. *J. Cell Biol.* **218**, 2982–3001 (2019).

31.   Siddiqui, N. *et al.* PTPN21 and Hook3 relieve KIF1C autoinhibition and activate intracellular
transport. *Nat. Commun.* **10**, 2693 (2019).

32.   Ephrussi, A. & Lehmann, R. Induction of germ cell formation by oskar. *Nature* **358**, 387–392
(1992).

33.   Clark, A., Meignin, C. & Davis, I. A Dynein-dependent shortcut rapidly delivers axis
determination transcripts into the Drosophila oocyte. *Development* **134**, 1955–1965 (2007).

34.   Zimyanin, V. L. *et al.* In vivo imaging of oskar mRNA transport reveals the mechanism of
posterior localization. *Cell* **134**, 843–853 (2008).

35.   Brendza, R. P., Serbus, L. R., Duffy, J. B. & Saxton, W. M. A function for kinesin I in the
posterior transport of oskar mRNA and Staufen protein. *Science* **289**, 2120–2122 (2000).

36.   Roth, S. & Lynch, J. A. Symmetry breaking during Drosophila oogenesis. *Cold Spring Harb.*
Perspect. Biol. **1**, a001891 (2009).

37.   Navarro, C., Puthalakath, H., Adams, J. M., Strasser, A. & Lehmann, R. Egalitarian binds
dynein light chain to establish oocyte polarity and maintain oocyte fate. *Nat. Cell Biol.* **6**, 427–
435 (2004).

38.   Trcek, T. & Lehmann, R. Germ granules in Drosophila. *Traffic* **20**, 650–660 (2019).

39.   Loiseau, P., Davies, T., Williams, L. S., Mishima, M. & Palacios, I. M. Drosophila PAT1 is
required for Kinesin-1 to transport cargo and to maximize its motility. *Development* **137**, 2763–
2772 (2010).

40.   Palacios, I. M. & St Johnston, D. Kinesin light chain-independent function of the Kinesin heavy
chain in cytoplasmic streaming and posterior localisation in the Drosophila oocyte.
Development **129**, 5473–5485 (2002).

41.   Veeranan-Karmegam, R., Boggupalli, D. P., Liu, G. & Gonsalvez, G. B. A new isoform of
Drosophila non-muscle Tropomyosin 1 interacts with Kinesin-1 and functions in oskar mRNA
localization. *J. Cell Sci.* **129**, 4252–4264 (2016).

42.   Gáspár, I., Sysoev, V., Komissarov, A. & Ephrussi, A. An RNA-binding atypical tropomyosin
recruits kinesin-1 dynamically to oskar mRNPs. *EMBO J.* **36**, 319–333 (2017).

43.   Dimitrova-Paternoga, L. *et al.* Molecular basis of mRNA transport by a kinesin-1-atypical
tropomyosin complex. *Genes Dev.* **35**, 976–991 (2021).

44.   Gáspár, I. *et al.* An RNA-based feed-forward mechanism ensures motor switching in oskar
mRNA transport. *BioRxiv* (2021) doi:10.1101/2021.04.24.441269.

45.   Sanghavi, P., Laxani, S., Li, X., Bullock, S. L. & Gonsalvez, G. B. Dynein associates with oskar
mRNPs and is required for their efficient net plus-end localization in Drosophila oocytes. *PLoS*
ONE **8**, e80605 (2013).

46.   Ghosh, S., Marchand, V., Gáspár, I. & Ephrussi, A. Control of RNP motility and localization by
a splicing-dependent structure in oskar mRNA. *Nat. Struct. Mol. Biol.* **19**, 441–449 (2012).

47.   Simon, B., Masiewicz, P., Ephrussi, A. & Carlomagno, T. The structure of the SOLE element of
oskar mRNA. *RNA* **21**, 1444–1453 (2015).

48.   Jambor, H., Mueller, S., Bullock, S. L. & Ephrussi, A. A stem-loop structure directs oskar
mRNA to microtubule minus ends. *RNA* **20**, 429–439 (2014).

49.   Seeger, M. A. & Rice, S. E. Microtubule-associated protein-like binding of the kinesin-1 tail to
microtubules. *J. Biol. Chem.* **285**, 8155–8162 (2010).

50.   Navone, F. *et al.* Cloning and expression of a human kinesin heavy chain gene: interaction of
the COOH-terminal domain with cytoplasmic microtubules in transfected CV-1 cells. *J. Cell*

Biol. **117**, 1263–1275 (1992).

51.   Jolly, A. L. *et al.* Kinesin-1 heavy chain mediates microtubule sliding to drive changes in cell
shape. *Proc Natl Acad Sci USA* **107**, 12151–12156 (2010).

52.   Goode, B. L. & Feinstein, S. C. Identification of a novel microtubule binding and assembly
domain in the developmentally regulated inter-repeat region of tau. *J. Cell Biol.* **124**, 769–782
(1994).

53.   Kikhney, A. G. & Svergun, D. I. A practical guide to small angle X-ray scattering (SAXS) of
flexible and intrinsically disordered proteins. *FEBS Lett.* **589**, 2570–2577 (2015).

54.   Receveur-Brechot, V. & Durand, D. How random are intrinsically disordered proteins? A small
angle scattering perspective. *Curr. Protein Pept. Sci.* **13**, 55–75 (2012).

55.   Jumper, J. *et al.* Highly accurate protein structure prediction with AlphaFold. *Nature* **596**, 583–
589 (2021).

56.   Evans, R. *et al.* Protein complex prediction with AlphaFold-Multimer. *BioRxiv* (2021)
doi:10.1101/2021.10.04.463034.

57.   Kikhney, A. G., Borges, C. R., Molodenskiy, D. S., Jeffries, C. M. & Svergun, D. I. SASBDB:
Towards an automatically curated and validated repository for biological scattering data.
Protein Sci. **29**, 66–75 (2020).

58.   Fumagalli, L. *et al.* C9orf72-derived arginine-containing dipeptide repeats associate with
axonal transport machinery and impede microtubule-based motility. *Sci. Adv.* **7**, (2021).

59.   Hisanaga, S. *et al.* The molecular structure of adrenal medulla kinesin. *Cell Motil. Cytoskeleton*
12, 264–272 (1989).

60.   Friedman, D. S. & Vale, R. D. Single-molecule analysis of kinesin motility reveals regulation by
the cargo-binding tail domain. *Nat. Cell Biol.* **1**, 293–297 (1999).

61.   Weijman, J. F. *et al.* Molecular architecture of the autoinhibited kinesin-1 lambda particle. *Sci.*
Adv. **8**, (2022).

62.   Williams, L. S., Ganguly, S., Loiseau, P., Ng, B. F. & Palacios, I. M. The auto-inhibitory domain
and ATP-independent microtubule-binding region of Kinesin heavy chain are major functional
domains for transport in the Drosophila germline. *Development* **141**, 176–186 (2014).

63.   Tan, Z. *et al.* Autoinhibited kinesin-1 adopts a hierarchical folding pattern. *BioRxiv* (2023)
doi:10.1101/2023.01.26.525761.

64.   Cassella, L. & Ephrussi, A. Subcellular spatial transcriptomics identifies three mechanistically
different classes of localizing RNAs. *Nat. Commun.* **13**, 6355 (2022).

65.   Gardiol, A. & St Johnston, D. Staufen targets coracle mRNA to Drosophila neuromuscular
junctions and regulates GluRIIA synaptic accumulation and bouton number. *Dev. Biol.* **392**,
153–167 (2014).

66.   Derr, N. D. *et al.* Tug-of-war in motor protein ensembles revealed with a programmable DNA
origami scaffold. *Science* **338**, 662–665 (2012).

67.   Gicking, A. M. *et al.* Kinesin-1, -2, and -3 motors use family-specific mechanochemical
strategies to effectively compete with dynein during bidirectional transport. *eLife* **11**, (2022).

68.   D'Souza, A. I., Grover, R., Monzon, G. A., Santen, L. & Diez, S. Vesicles driven by dynein and
kinesin exhibit directional reversals without external regulators. *BioRxiv* (2022)
doi:10.1101/2022.09.27.509758.

69.   Belyy, V. *et al.* The mammalian dynein-dynactin complex is a strong opponent to kinesin in a
tug-of-war competition. *Nat. Cell Biol.* **18**, 1018–1024 (2016).

70.   Elshenawy, M. M. *et al.* Cargo adaptors regulate stepping and force generation of mammalian
dynein-dynactin. *Nat. Chem. Biol.* **15**, 1093–1101 (2019).

71.   Feng, Q., Gicking, A. M. & Hancock, W. O. Dynactin p150 promotes processive motility of DDB
complexes by minimizing diffusional behavior of dynein. *Mol. Biol. Cell* **31**, 782–792 (2020).

72.   Qiu, R., Zhang, J. & Xiang, X. Kinesin-1 autoinhibition facilitates the initiation of dynein cargo
transport. *J. Cell Biol.* **222**, (2023).

73.   Sung, H.-H. *et al.* Drosophila ensconsin promotes productive recruitment of Kinesin-1 to
microtubules. *Dev. Cell* **15**, 866–876 (2008).

74.   Barlan, K., Lu, W. & Gelfand, V. I. The microtubule-binding protein ensconsin is an essential
cofactor of kinesin-1. *Curr. Biol.* **23**, 317–322 (2013).

75.   Ferro, L. S. *et al.* Structural and functional insight into regulation of kinesin-1 by microtubule-

associated protein MAP7. *Science* **375**, 326–331 (2022).

76.   Hachet, O. & Ephrussi, A. Splicing of oskar RNA in the nucleus is coupled to its cytoplasmic
localization. *Nature* **428**, 959–963 (2004).

77.   Amrute-Nayak, M. & Bullock, S. L. Single-molecule assays reveal that RNA localization signals
regulate dynein-dynactin copy number on individual transcript cargoes. *Nat. Cell Biol.* **14**, 416–
423 (2012).

78.   Edelstein, A., Amodaj, N., Hoover, K., Vale, R. & Stuurman, N. Computer control of
microscopes using µManager. *Curr. Protoc. Mol. Biol.* **Chapter 14**, Unit14.20 (2010).

79.   Schindelin, J. *et al.* Fiji: an open-source platform for biological-image analysis. *Nat. Methods* **9**,
676–682 (2012).

80.   Sattler, M. Heteronuclear multidimensional NMR experiments for the structure determination of
proteins in solution employing pulsed field gradients. *Prog. nucl. magn. reson. spectrosc.* **34**,
93–158 (1999).

81.   Simon, B. & Köstler, H. Improving the sensitivity of FT-NMR spectroscopy by apodization
weighted sampling. *J. Biomol. NMR* **73**, 155–165 (2019).

82.   Delaglio, F. *et al.* NMRPipe: a multidimensional spectral processing system based on UNIX
pipes. *J. Biomol. NMR* **6**, 277–293 (1995).

83.   Johnson, B. A. & Blevins, R. A. NMR View: A computer program for the visualization and
analysis of NMR data. *J. Biomol. NMR* **4**, 603–614 (1994).

84.   Williamson, M. P. Using chemical shift perturbation to characterise ligand binding. *Prog. nucl.*
magn. reson. spectrosc. **73**, 1–16 (2013).

85.   Kieffer, J. *et al.* New data analysis for BioSAXS at the ESRF. *J. Synchrotron Radiat.* **29**, 1318–
1328 (2022).

86.   Franke, D. *et al.* ATSAS 2.8: a comprehensive data analysis suite for small-angle scattering
from macromolecular solutions. *J. Appl. Crystallogr.* **50**, 1212–1225 (2017).

87.   Simon, B., Madl, T., Mackereth, C. D., Nilges, M. & Sattler, M. An efficient protocol for NMR-
spectroscopy-based structure determination of protein complexes in solution. *Angew. Chem.*
Int. Ed **49**, 1967–1970 (2010).

88.   Brunger, A. T. Version 1.2 of the Crystallography and NMR system. *Nat. Protoc.* **2**, 2728–2733
(2007).

89.   Linge, J. P., Habeck, M., Rieping, W. & Nilges, M. ARIA: automated NOE assignment and
NMR structure calculation. *Bioinformatics* **19**, 315–316 (2003).

90.   Gruber, A. R., Lorenz, R., Bernhart, S. H., Neuböck, R. & Hofacker, I. L. The Vienna RNA
websuite. *Nucleic Acids Res.* **36**, W70-4 (2008).

Supplementary Material

Tropomyosin 1-I/C co-ordinates kinesin-1 and dynein motors during *oskar* mRNA transport

Simone Heber[1][†], Mark A. McClintock[2][†], Bernd Simon[3], Janosch Hennig[3,4], Simon L. Bullock[2]*,
Anne Ephrussi[1]*

1   Developmental Biology Unit, European Molecular Biology Laboratory, 69117 Heidelberg, Germany

2   Division of Cell Biology, MRC Laboratory of Molecular Biology, Cambridge CB2 0QH, United Kingdom

3   Structural and Computational Biology Unit, European Molecular Biology Laboratory, 69117 Heidelberg, Germany

4   Biochemistry IV, Biophysical Chemistry, University of Bayreuth, 95447 Bayreuth, Germany

[†]These authors contributed equally to this work

*Correspondence: sbullock@mrc-lmb.cam.ac.uk, anne.ephrussi@embl.org

1040

1041

Figure S1: Mass photometry measurements of the Khc-Tm1 complex at 20 nM concentration. The
sample comprises a mixture of 47 % particles of 224 kDa, corresponding to the Khc dimer (theoretical
MW = 221.2 kDa), 40 % particles of 266 kDa, corresponding to a Khc dimer bound to a Tm1 monomer
(theoretical MW = 269.2 kDa), and 13 % of the particles represent larger species. This indicates that in
solution, ~50 % of Khc is bound to Tm1 at steady-state at concentrations below the K_D, confirming the
2:1 stoichiometry of the Khc-Tm1 complex[43].

1048

Figure S2: a RNAfold [90] MFE prediction of *osk* min RNA. Both SOLE and OES are predicted as stem-loop structures, indicating that in the context of *osk* min, the LEs adopt their native folds. **b, c** Example kymographs showing processive movements of AlexaFluor647-labeled SNAP-Khc FL (b) or the tailless

Khc910 (c) in absence or presence of Tm1. Microtubule plus and minus ends are oriented toward the

right and left of each kymograph, respectively. **d** Processive events of Khc FL and Khc910 in absence

or presence of Tm1. In this and other figures, black or white circles indicate proteins that were present

or absent from the experiment, respectively. Data points show values for individual microtubules

(between 11 and 178 complexes analyzed per condition). Error bars: SD. Statistical significance was

evaluated with an unpaired t-test. ****$p<0.0001$. **e** Average velocities of Khc FL and Khc910 in absence

or presence of Tm1. Data points represent individual runs. Statistical significance was evaluated with

an unpaired t-test. ****$p<0.0001$ **f, g** Histograms of the velocity distributions of Khc FL (f) and Khc910

(g) in absence or presence of Tm1.

Figure S3: a All microtubule (MT)-binding events (processive, static and diffusive) observed for Khc FL,
Khc910 and Khc940 ± Tm1 FL. **b** Fraction of observed Khc events colocalizing with *osk* 3'UTR and **c**
fraction of observed Khc events colocalizing with Tm1, for Khc FL, Khc910 and Khc940. Data points
show values for individual microtubules. Statistical significance was evaluated with an unpaired t-test.
*p<0.1 **p<0.01.
1068

1070

Figure S4: a All microtubule (MT)-binding events (processive, static and diffusive) observed for Khc FL, Khc910 and KhcΔAMB ± Tm1 FL. **b** Fraction of observed Khc events colocalizing with *osk* 3'UTR and **c** fraction of observed Khc events colocalizing with Tm1, for Khc FL, Khc910 and KhcΔAMB. Data points show values for individual microtubules. Statistical significance was evaluated with an unpaired t-test. **p<0.01, ***p<0.001, ****p<0.0001.

1076
Figure S5: a All microtubule (MT)-binding events (processive, static and diffusive) observed for Khc FL
± Tm1 FL or Tm1 1-335. **b** Fraction of observed Khc events colocalizing with *osk* 3'UTR and **c** fraction
of observed Khc events colocalizing Tm1 FL or Tm1 1-335. Labeling efficiencies for Tm1 FL and Tm1
1-335 were 57 %, and 58 %, respectively, such that co-localization results are comparable. **d** All
microtubule (MT)-binding events (processive, static and diffusive) observed for Khc FL ± Tm1 FL or
Tm1 252-334. **e** Fraction of processive events of Khc FL ± Tm1 FL or Tm1 252-334. **f** Average velocities
of Khc FL ± Tm1 FL or Tm1 252-334 in the assembly mix. Data points show values for individual
microtubules. Statistical significance was evaluated with an unpaired t-test. *p<0.1 **p<0.01
****p<0.0001.

1086
Figure S6: a Microtubule binding assay with Tm1 FL and Tm1 1-335. Khc FL served as a positive
control. Both Tm1 FL and Tm1 1-335 partition partially into the microtubule pellet fraction. 10 µg of each
protein were incubated with polymerized microtubules prior to segregation of soluble and microtubule-
bound fractions by ultracentrifugation over a 30 % sucrose cushion. BSA was added to all samples as
a negative control. To control for microtubule-independent pelleting of the proteins, controls without
microtubules were performed. Soluble and microtubule pellet fractions were analyzed by SDS-PAGE.
b Microtubule binding assay with Khc 855-975 and Tm1 1-335. Khc 855-975 partially partitions into the
microtubule pellet both in absence and presence of increasing concentrations of Tm1 1-335. 25 µM Khc
855-975 were incubated with Tm1 1-335 at ratios 1:0.5, 1:1 and 1:2 and polymerized microtubules
before proceeding as described above.

Figure S7: GST-pulldown competition assays to assess the effect of Tm1 on Khc tail - motor interaction. **a** 5 µM or **b** 10 µM GST-Khc 855-975 as bait and 50 µM Khc 1-365 as prey. Tm1 1-335 was added at increasing concentrations (5 to 15 µM) as a competitor for binding to GST-Khc 855-975.

Figure S8: a Template model of Khc used for randomization. Coordinates of the structured parts (motor
domain, residues 1-378 and the four coiled-coil regions, residues 403-585, 604-706, 711-838 and 844-
936, respectively) were generated by AF2 [55,56] and connected with the coordinates of the missing linker
residues. **b** χ^2 distribution of all Crysol fits to the Khc (left) and Khc-Tm1 (right) scattering curves. **c** Best
(upper panel) and mediocre (lower panel) fits of Khc (left) and Khc-Tm1 (right) structural models to the
experimental scattering curves. **d** Five best fits for Khc (left) and for Khc-Tm1 (right) aligned. **e** Close-
ups of the AMB domains in the four best fitting models for Khc. **f** Close-ups of the AMB domains in the
four best fitting models for Khc-Tm1. Khc models are shown in shades of blue with the AMB domains
marked in red, Tm1 is shown in shades of pink.

1113

Figure S9: a-c Fraction of dynein binding events that underwent processive motility (a) and frequencies of processive dynein movement (b) and dynein binding to microtubules (c) in the presence and absence of Tm1 FL. Dots represent average values for individual microtubules. In each plot, the mean ± SD is shown. Statistical significance was determined by t-test. ns: not significant.

TMR-Dynein
A488-*osk* 3'UTR 2+3/
HiLyte 488 tubulin
A647-Khc FL
Merge

TMR-DDBE ●
A488-*osk* 3'UTR 2+3 ●
A647-Khc FL ●
Tm1 FL ●

30 s

5 µm

Microtubule minus ends enriched
in HiLyte 488 tubulin

Figure S10: Kymographs showing the Tm1-induced co-translocation of dynein, *osk* 3'UTR 2+3 RNA, and Khc toward microtubule minus ends, which are labeled by enrichment of fluorescent (HiLyte 488) tubulin dimers. Microtubule plus and minus ends are oriented toward the right and left of each kymograph, respectively.

TMR-DDBE ●
A488-*osk* 3'UTR 2+3 ●
A647-Khc FL ●
Tm1 FL ○

1125

Figure S11: Kymographs showing instances of bidirectional Khc movements in the absence of Tm1. Directional switches of Khc (white arrowheads) correlate with DDBE-*osk* 3'UTR 2+3 association and dissociation events. Microtubule plus and minus ends are oriented toward the right and left of each kymograph, respectively.

1130

1131

1132

Figure S12: a Kymographs showing reduced association of Khc with motile DDBE-*osk* 3'UTR 2+3
RNPs in the presence of Tm1 after treatment with RNase A. Microtubule plus and minus ends are
oriented toward the right and left of each kymograph, respectively. **b** Fraction of moving DDBE-*osk*
3'UTR 2+3 RNPs that colocalized with Khc signal with and without RNase A treatment in the presence
of Tm1. Dots represent average values of all motility on individual microtubules. Mean ± SD is shown.
Statistical significance was determined by Welch's t-test. ****: p<0.0001.

Supplementary Tables

1140

	MT binding events [min^{-1} µm^{-1}]	Processive events [min^{-1} µm^{-1}]	Fraction of processive events	Average velocity [µm s^{-1}]
Khc FL N=83	0.13 ± 0.03	0.11 ± 0.03	0.8415	0.183 ± 0.131
+ **Tm1 FL** N=36	0.11 ± 0.06	0.09 ± 0.07	0.8182	0.044 ± 0.422
Khc910 N=438	0.55 ± 0.26	0.55 ± 0.27	1	0.614 ± 0.180
+ **Tm1 FL** N=403	0.68 ± 0.31	0.68 ± 0.31	0.9952	0.560 ± 0.202
Khc940 N=382	0.61 ± 0.23	0.60 ± 0.22	0.9828	0.546 ± 0.260
+ **Tm1 FL** N=97	0.20 ± 0.06	0.15 ± 0.06	0.7327	0.073 ± 0.180

1141   **Table S1** Motility parameters of Khc FL, Khc910 and Khc940

	MT binding events [min⁻¹ μm⁻¹]	Processive events [min⁻¹ μm⁻¹]	Fraction of processive events	Average velocity [μm s⁻¹]
Khc FL N=142	0.07 ± 0.15	0.05 ± 0.02	0.7696	0.429 ± 0.300
+ Tm1 FL N=24	0.02 ± 0.01	0.01 ± 0.01	0.3114	0.159 ± 0.188
Khc910 N=384	0.47 ± 0.10	0.47 ± 0.10	0.8180	0.630 ± 0.169
+ Tm1 FL N=595	0.73 ± 0.11	0.72 ± 0.11	0.8955	0.649 ± 0.164
KhcΔAMB N=65	0.06 ± 0.02	0.05 ± 0.02	0.9929	0.507 ± 0.325
+ Tm1 FL N=293	0.12 ± 0.05	0.10 ± 0.04	0.9947	0.548 ± 0.266

1143   **Table S2** Motility parameters of KhcΔAMB compared to Khc FL and Khc910

	MT binding events [min^{-1} µm^{-1}]	Processive events [min^{-1} µm^{-1}]	Fraction of processive events	Average velocity [µm s^{-1}]
Khc FL N=335	0.23 ± 0.08	0.16 ± 0.06	0.6847	0.32 ± 0.245
+ **Tm1 FL** N=149	0.21± 0.07	0.09 ± 0.04	0.4449	0.229 ± 0.280
+ **Tm1 1-335** N=170	0.13 ± 0.03	0.06 ± 0.04	0.4376	0.125 ± 0.165
Khc FL N=84	0.07 ± 0.02	0.06 ±0.02	0.9524	0.171 ± 0.212
+ **Tm1 FL** N=44	0.04 ± 0.01	0.03 ± 0.01	0.7137	0.188 ± 0.243
+ **Tm1 252-334** N=171	0.08 ± 0.03	0.07 ± 0.03	0.8694	0.303 ± 0.188

1145   **Table S3** Motility parameters of Khc FL in presence of different Tm1 constructs.

	Total dynein binding frequency [µm⁻¹ min⁻¹]	Processive dynein frequency [µm⁻¹ min⁻¹]	Processive dynein fraction	Average dynein velocity [µm s⁻¹]	Average dynein run length [µm]
DDBE-*osk* 3'UTR 2+3 -Tm1 FL	1.16 ± 0.20 N=20 microtubules (2152 complexes)	0.67 ± 0.14 N=20 microtubules (1277 complexes)	0.58 ± 0.07 N=20 microtubules (2152 complexes)	0.292 ± 0.308 N=1277 complexes	4.26 ± 3.90 N=1277 complexes
DDBE-*osk* 3'UTR 2+3 +Tm1 FL	1.27 ± 0.24 N=20 microtubules (2207 complexes)	0.75 ± 0.19 N=20 microtubules (1295 complexes)	0.58 ± 0.06 N=20 microtubules (2207 complexes)	0.320 ± 0.334 N=1295 complexes	4.25 ± 3.66 N=1295 complexes
DDBE-*osk* 3'UTR 2+3 +Tm1 FL (Tm1-bound only)	N/A	N/A	N/A	0.304 ± 0.310 N=211 complexes	N/A

Table S4 Motility parameters of DDBE-*osk* 3'UTR 2+3 in presence and absence of Tm1 FL (Related to Figure 7 and S9). Mean ± SD are shown.

	Frequency of Khc transport events toward plus ends [μm^{-1} min^{-1}]	Frequency of Khc transport events toward minus ends [μm^{-1} min^{-1}]	Average Khc-bound dynein velocity [μm s^{-1}]	Average distance traveled by Khc toward minus ends [μm]
DDBE-*osk* 3'UTR 2+3 +Khc -Tm1 FL	0.005 ± 0.007 N=52 microtubules (29 complexes)	0.014 ± 0.013 N=52 microtubules (89 complexes)	0.236 ± 0.223 N=109 complexes	3.32 ± 3.82 N=109 complexes
DDBE-*osk* 3'UTR 2+3 +Khc +Tm1 FL	0.005 ± 0.008 N=36 microtubules (20 complexes)	0.179 ± 0.064 N=36 microtubules (822 complexes)	0.348 ± 0.281 N=913 complexes	4.92 ± 4.79 N=913 complexes

1150 **Table S5** Motility parameters of DDBE-*osk* 3'UTR 2+3 and Khc in presence and absence of Tm1 FL
1151 (Related to Figure 8). Mean ± SD are shown.

1152 Plasmids

CONSTRUCTS	RELEVANT INFORMATION	SOURCE
pFastBacDual-HisSUMO-SNAP-3C-Khc	Amp^r, p10/Polyhedrin promoter	This study
pFastBacDual-HisSUMO-SNAP-3C-Khc910	Amp^r, p10/Polyhedrin promoter	This study
pFastBacDual-HisSUMO-SNAP-3C-Khc940	Amp^r, p10/Polyhedrin promoter	This study
pFastBacDual-HisSUMO-SNAP-3C-Khc ΔAMB	Amp^r, p10/Polyhedrin promoter	This study
pFastBacDual-HisSUMO-SNAP-3C-Khc-Tm1	Amp^r, p10/Polyhedrin promoter	This study
pGEX6.1-Khc 855-975	Amp^r, tac promoter	This study
pETM11-His6-SUMO-SNAP-Tm1-FL	Kan^r, T7 promoter, lac operator	This study
pETM11-His6-SNAP-SUMO-Tm1 1-335	Kan^r, T7 promoter, lac operator	This study
pETM11-His6-SUMO-Tm1-FL	Kan^r, T7 promoter, lac operator	[43]
pETM11-His6-SUMO-Tm1 1-335	Kan^r, T7 promoter, lac operator	[43]

CONSTRUCTS	RELEVANT INFORMATION	SOURCE

pETM11-His6-SUMO-Tm1 252-334	Kan[r], T7 promoter, lac operator	[43]
pDyn1 (pAceBac1-ZZ-TEV-SNAPf-Dynein heavy chain - DYNC1H1)	Gen[r], Polyhedrin promoter	[8]
pDyn2 (pIDC-Dynein intermediate chain - DYNC1/2, Dynein light intermediate chain - DYNC1LI2, Tctex - DYNLT1, Robl - DYNLRB1, LC8 - DYNLL1)	Chlor[r], Polyhedrin promoter	[8]
pAceBac1-Egl-TEV-ZZ	Gen[r], Polyhedrin promoter	[15]
pIDC-BicD	Chlor[r], Polyhedrin promoter	[15]

Table S6 Plasmids

RNA sequences

osk **3'UTR**

GUUGGGUUCUUAAUCAAGAUACAUAUAUGCAAAUUUUGACUGGGCUGGCACC

GGAACCGACAAAAUAAGAACUUUUUGAUGAUUUUACGAUUUACGCUGAUGGA

UCGCUGCUUUUACGGAAUUCGCUUAGUUUUAAUAUGUUUUAUAUGUAGUAUG

UUCUCUGUCUUUGUUUAUUUAUAUGUUCGUGCACUUGUCCUAGUCCAUUAUU

GUAUAUUAUUGUGUGUUUUGUGUUCUAUGUUAGAUUUAAACUUCUCAAUUUU

UCGCUGUCUGUGAUUUGUUUUGCCAAUGCCAUUGAUUUUCUGCACACUUUUU

GCUGCUAUCCCAAAAGCUGUGUAAAAUAAUCAAAUGCAAAAUAAGCGCAAGCA

GCUGAAAACUUCUCUUCAAACUUUUCCGCUUUUCCCAAAACCAUUUUGCUUU

GAAAUCUGUUUUUACCAAAUCAGAUUAAACUGCAAAAUGGAACUUAAAUGCAA

AUCAUUGCAAUGCUUAUAAACUGUUUUUUGUUCUAUAUACUUUUGUGUGGGU

CAAAAUUCGGCAUGCUCCUGUAUCACACAACCUGCCACUUGCCCUUAAAAAG

AAGGGCGCAGUGGGCGUGGUACGUAUACAUAUGAGCCAUGCUGCAUUUUGG

CCGUAAUGAAAUGCACUGCUUUACUUGGAAAAUUCGCUUGCACAAAUCAAC

GCCGCGGCUGAUUUAUUAUUGAUGUGCUCAAGCAAAUUCAAGUGAAGCAUUU

GCGCGAUUUUCGUCUUUCUGUUUCCGUUUGCAAAAAGUUUAUAAAAUGCUU

ACACUCUGCUGCAGACACGCCAACCGGAAGUGCGCACUAAGCGCUUGUUUGU

AGCACAGUGUAGAAUUCUGGCGUAAUUUACAGCUCUACUUUAAAGUCUUCUA

GAUAGCUAUCUACUAUUUAUAAACUUAUUUAUUGUCUUGAAUGUAUGUUAAU

UGUAUGUAUUGAUGGUGAUCACGUUUUUUUUGUCCUAUAACAAGCUGCAAUG

UAAAAUCCAAAAAAAAUGAAAAAUAAAAUAAAAGGGAAAUCAAUGAUUUUG

UUUGCUCUAAGCUAUGUUCAAAGUAACUGG

Total length: 1074 nt

1178

osk min

CCACGCCCAGAAUGGAAUCA<mark style="background-color: yellow">CGAUAUCGAGCAUCAAGAGUGAAUAUCG</mark>AAAA

CUGACGGGCAAUCCAU<mark style="background-color: cyan">UGGAAAAUUCGCUUGCACAAAAUCAACGCCGCGGCU</mark>

<mark style="background-color: cyan">GAUUUAUUAUUGAUGUGCUCAAGCAAAUUCA</mark>

Total length: 135 nt

<mark style="background-color: yellow">SOLE</mark>

<mark style="background-color: cyan">OES</mark>

1186

1187

osk 3'UTR regions 2+3

UGUUCUAUAUACUUUUGUGUGGGUCAAAAUUCGGCAUGCUCCUGUAUCACAC

AACCUGCCACUUGCCCUUAAAAAGAAGGGCGCAGUGGGCGUGGUACGUAUAC

AUAUGAGCCAUGCUGCAUUUUGGCCGUAAUGAAAUGCACUGCUUUACUUGG

AAAAUUCGCUUGCACAAAAUCAACGCCGCGGCUGAUUUAUUAUUGAUGUGCU

CAAGCAAAUUCAAGUGAAGCAUUUGCGCGAUUUUCGUCUUUCUGUUUCCGUU

UGCAAAAAGUUUAUAAAAUGCUUACACUCUGCUGCAGACACGCCAACCGGAA

GUGCGCACUAAGCGCUUGUUUGUAGCACAGUGUAGAAUUCUGGCGUAAUUUA

CAGCUCUACUUUAAAGUCUUCUAGAUAGCUAUCUACUAUUUAUAAACUUAUUU

AUUGUCUUGAAUGUAUGUUAAUUGUAUGUAUUGAUGGUGAUCACGUUUUUUU

UGUCCUAUAACAAGCUGCAAUGUAAAAUCCAAAAAAAAAUGAAAAAUAAAAU

AAAAGG

Total length: 530 nt