Taxonomic identification accuracy from BOLD and GenBank databases using over a thousand insect barcodes from Colombia

Short title: Taxonomic identification accuracy from BOLD and GenBank databases using insect barcodes

Nathalie Baena-Bejarano[1]*, Catalina Reina[2], Diego Esteban Martínez-Revelo[3, 4], Claudia A. Medina[1], Eduardo Tovar[1], Sandra Uribe-Soto[4], Jhon Cesar Neita-Moreno[1], Mailyn A. Gonzalez[1]*

[1]Instituto de Investigación de Recursos Biológicos Alexander von Humboldt, Bogotá, Colombia.

[2]ICA - Instituto Colombiano Agropecuario, Soledad, Colombia.

[3]Asociación GAICA, Pasto, Nariño, Colombia.

[4]Grupo de Investigación en Sistemática Molecular, Universidad Nacional de Colombia, sede Medellín, Medellín, Colombia.

* Corresponding authors

E-mails: ntbaena@gmail.com (NBB), magonzalez@humboldt.org.co (MAG)

Abstract

Recent declines of insect populations at high rates have resulted in the need to develop a quick method to determine their diversity and to process massive data for the

identification of species of highly diverse groups. A short sequence of DNA from COI is

widely used for insect identification by comparing it against sequences of known

species. Repositories of sequences are available online with tools that facilitate

matching of the sequences of interest to a known individual. However, the performance

of these tools can differ. Here we aim to assess the accuracy in identification of insect

taxonomic categories from two repositories, BOLD Systems, and GenBank. This was

done by comparing the sequence matches between taxonomist identification and the

suggested identification from the platforms. We used 1160 sequences in eight orders of

insects from Colombia. After the comparison, we reanalyzed the results from a

representative subset of the data from the subfamily Scarabaeinae (Coleoptera).

Overall, BOLD systems outperformed GenBank, and the performance of both engines

differed by orders and other taxonomic categories (species, genus and family). Higher

rates of accurate identification were obtained at family and genus level. The accuracy

was higher in BOLD for the orders Coleoptera at family, for Coleoptera and Lepidoptera

at genus, and species level. The other orders performed similarly in both repositories.

Moreover, the Scarabaeinae subset showed that in this group species were correctly

identified when BOLD match percentage was above 93.4% and a total of 85% of the

samples were correctly assigned to a taxonomic category. These results accentuate the

great potential of the identification engines to place insects accurately into their

respective taxonomic categories based on DNA barcodes and highlight the reliable use

of BOLD Systems for insect identification in the absence of a large reference database

for a highly diverse country.

Introduction

More than a million species of plants and animals are in danger of extinction according

to the latest global assessment of the intergovernmental platform on biological diversity

and ecosystem services [1]. Declines in biodiversity and projections of a sixth mass

extinction [2] have placed the impact of such losses on the agendas of governments all

over the world [3]. Although insects constitute one of the most diverse groups on the

planet, many species remain undiscovered and recent evidence suggests the decline of

their populations at high rates [4,5]. To date, the risk of extinction of 12,161 species of

insects has been evaluated according to the criteria of the IUCN red list and 2,291

species are considered "threatened" around the world [6].

Colombia is known as a biodiversity hotspot in the world. Although insects are

the most studied class of animals in the country [7], research involving their genetic

diversity is still currently underrepresented. Countries around the world are

implementing massive sampling of insects with barcode generation under the premise

that barcodes will allow us to assess biodiversity at a higher speed [8]. Barcodes in

taxonomic works are showing an incredibly hidden diversity of insects with descriptions

of hundreds of new species facilitated by DNA, morphology, and natural history [9,10].

As a megadiverse country, Colombia has great challenges to complete the inventory of

its biodiversity and then the sustainable use of its resources. In particular, accelerating

species identification using an integrative taxonomy is a priority.

Assessing biodiversity via sampling and DNA barcoding has been successfully

applied in areas where coordinated sampling and barcoding facilitated the creation of

extensive libraries for a particular region. An example of such collaboration is described

in Morinière et al [8] where the Bavarian State Collection of Zoology (ZSM), the

Barcoding Fauna Bavarica (BFB), and the German Barcode of Life (GBOL) project

generated a library of 120,000 reliably identified species in Coleoptera, Diptera,

Hymenoptera, and Lepidoptera for Germany and Central Europe. A closer example in

the Neotropics is Costa Rica which has been leading the Bioalfa project and within the

last 15 years has generated 45,000 insect species barcodes recorded [11]. Added to

this initiative are other countries across the globe such as Canada, Ecuador, Sweden,

and Singapore [9,12–16]. Moreover, the development and implementation of effective

methods for species monitoring are important for assessing biodiversity, and overall

health of ecosystems [8]. The species richness of insects in terrestrial ecosystems

facilitates their use as bioindicators for measuring biodiversity and tracking the effects of

changes in environmental conditions. The challenges faced by biodiversity declines

have resulted in a need to generate new strategies for monitoring that are capable of

generating vast amounts of data in a rapid and cost-effective manner. One such

strategy that has been gaining traction in recent years has been the implementation of

DNA barcoding.

Barcode sequences are generated from a standardized region that consists of

short DNA sequences (generally between ~300 to ~700 bp). The mitochondrial

Cytochrome c oxidase subunit I (COI) is most often used for insect barcoding. A DNA

barcoding repository is built from sampling the DNA of individuals that are retained as

vouchers, from which a photograph is taken. In the case of insects, most often a leg is

removed from the specimen and used to obtain the DNA. In some cases, whole bodies

of the samples are used when the genetic material available is limited by the size of the

specimen. Although non-destructive DNA extraction methods are preferred to keep the

morphology for future reference, photographs are suggested for facilitating identification

before this step. This approach requires the creation of a large DNA reference database

where the identification of unknown specimens is achieved through the information

confirmed and curated in the database. The idea behind is that unknown samples could

be identified by matching their sequences to the species that are curated from the

database. The Barcode of Life Data Systems BOLD (http://www.boldsystems.org) [17]

is a platform specifically developed for this type of information. The BOLD platform

provides the identifications match based on BINs (Barcode Index Numbers) [18] that are

automatically generated through algorithms for the specimens. Although unidentified

species coded in BINs can help estimating diversity, the identification of species is also

possible using the Basic Local Alignment Tool (BLAST) from GenBank [19,20]. BLAST

allows matching sequences to a query sequence. The search consists of comparing

nucleotide sequences hosted in the database and to provide statistical significance of

these matches. The main goal from BLAST is not necessarily to provide species

identification because the same principle can be used for the search of gene families;

uses of identification databases are vast in biology. Scientists frequently use these two

platforms to identify species from barcodes. An assessment of these two main public

repositories from curated samples across taxa found a slightly better performance from

GenBank vs BOLD for insects (n=17) at species and genus level identification without

statistical significance, and more even results between repositories for plants (n=61)

and macro-fungi (n=16) [21]. Later, a revision of this insect data found the results to be

more similar than previously reported for this group [22]. Accurate identification of

species and overall composition in ecosystems based on sampling is a key factor for

conservation initiatives. Therefore, evaluating the performance of these platforms

assigning specimens to a specific taxonomic category level has a significant impact on

the advancement of other fields.

Here, we analyze a large inventory of over 1,000 insect barcodes for Colombia to

the database which includes the first strepsipteran barcode from Colombia. We aim to

compare the matches of the taxonomist identification to species, genus, and family level

with the search engines of BOLD and BLAST service from GenBank in order to provide

suggestions and next steps for the use of these tools. We assess the accuracy of the

species identification engine from BOLD by using a subset of insects corresponding to

Scarabaeinae subfamily (Coleoptera: Scarabaeidae).

Materials and Methods

Collection Locality/Geography

Insect samples were collected from ten departments of Colombia through different

projects (Colombia BIO, Santander BIO). A majority of the samples (81.0%) came from

three departments of Colombia (Antioquia, Vichada, and Santander), the spatial data of

the localities were processed in ArcGIS 10.2 (License E300 04/26/2013) (Fig 1). The

elevation of the samples covered a wide range with the lowest sampling occurring at 56

masl and the highest at 3563 masl. Although samples were obtained with seven

different entomological collecting methods (malaise, pitfall, light trap, hand-picked,

entomological net, Van Someren-Rydon trap and folding net), nine malaise traps

accounted for 62.6% of the total collected material, followed by pitfall 17.2%, and light

trap 8.1%.

Fig 1. Map of Colombia showing collecting localities in black dots. Ocean base

map from Esri, GEBCO, NOAA, National Geographic, DeLorme, HERE, Geonames.org,

and other contributors. Hillshade map based on SRTM30_PLUS v8 [23].

Taxa description

This dataset includes insects from the orders Coleoptera, Diptera, Hymenoptera,

Lepidoptera, Hemiptera, Odonata, Psocodea and Strepsiptera, from which Coleoptera

is the most represented order in the dataset (35%). The specimens were identified from

morphology by students working on their thesis and by specialists. As a result of varying

expertise, we provide an identification that goes from order to species level (S1 Table);

additionally in Scarabaeinae, identification was done to morphospecies level following

González-Alvarado et al [24]. Currently, few samples (7%) have been identified to lower

levels from molecular data by BOLD Data Manager. The specimens were deposited at

collections with 999 samples between the Entomological and Tissue collection of the

Institute Alexander von Humboldt (IAvH-E and IAvH-CT), Museo Javeriano de Historia

Natural (MPUJ) at Pontificia Universidad Javeriana with 63, Museo Francisco Luis

Gallego (MEFLG) at Universidad Nacional de Colombia, Sede Medellin with 55, and

Colección Entomológica Universidad de Antioquia (CEUA) with the remaining 43

samples. Records of some samples are also available for search in SIB Colombia.

DNA sampling and extraction

One leg was removed from each insect, when possible, but some samples required the

use of the full body to provide enough DNA. All samples were sent to the molecular

laboratory of Instituto Humboldt at CIAT, Palmira. Extraction protocols were conducted

following the high throughput DNA isolation from Ivanova et al [25]. The concentration of

selected DNA samples was estimated by quantification using the NanoDrop 1000

(Thermo Fisher Scientific). Amplification of cytochrome c oxidase I (COI) fragments

(658 bp) took place by polymerase chain reaction (PCR) with a laboratory prepared

master mixture which contained 2 μL of template DNA (~10-50 ng), 1X Taq buffer

((NH4)2SO4), 200 μM of each deoxynucleoside triphosphate, 2 mM MgCl2, 0.2 μM of

each primer, 0.4 μg/μL bovine serum albumin, and 1 U of Taq DNA polymerase. A

cocktail of four primers was used: LCO1490 5'-GGTCAACAAATCATAAAGATATTGG-

3', HCO2198 5'- TAAACTTCAGGGTGACCAAAAAATCA-3' [26], Lep-F1 5'-

ATTCAACCAATCATAAAGATATTGG-3' and Lep-R1 5'-

TAAACTTCTGGATGTCCAAAAAATCA-3' [27]. Primers were selected following

Canadian Centre for DNA Barcoding (CCDB) suggested primers for taxonomic groups.

The PCR cycling conditions consisted of a first cycle of denaturation at 94 °C for 3 min;

followed by 5 cycles of 94 °C for 30 s, 45 °C for 40 s, and 72 °C for 1 min; then 35

cycles of 94 °C for 30 s, 51 °C for 40 s, and 72 °C for 1 min; and a final extension cycle

at 72 °C for 5 min. PCR products were visualized by electrophoresis on 1.5% agarose

gels stained with SYBR Safe (Thermo Fisher Scientific) and using 1X TBE. The

ExoSAP-IT protocol was used to clean PCR products, after which Sanger sequencing of

amplicons were carried out at The Universidad Nacional de Colombia and the University

of the Andes. Sequences were assembled and edited in Geneious 10.0.9 2017

(Auckland, New Zealand) [28]. All files were uploaded in BOLD systems and are found

in the dataset "insects-all" (DS-CBIH2020; Dataset available at: DOI will be made

available after acceptance).

Data analysis

BOLD Batch ID engine

To identify samples to species level, we ran BOLD's Identification Engine in

datasets created for insect orders in July 2020. The identification engine allows the user

to select the search within the COI Species Database or COI Full Database. We

selected COI Species Database which will match the sequence against an identified

species as opposed to individuals without identification to species level. We kept filters

as default for a minimum overlap of 300 bp and sequence length ≥ 100 bp. We ran the

analysis with 97% similarity, a percentage that accounts for within species variation and

error of the sequence [8]. Under the criteria selected of 97% numerous species ID were

not obtained. We decided to lower the criteria and include matches using the 80%

similarity filter due to the lack of matches at 97%. The 80% similarity is the minimum

filter allowed in BOLD. A match was accepted as correct when the identification by the

taxonomist was in agreement by the suggested identification by BOLD. The ID search

engine provides a list of maximum 100 closest specimens, and we selected the first

suggestion from the list that has the highest percentage match and base pair overlap.

We included an alternative match when the suggestion was done towards a sequence

from our project and ran statistical tests separately to prevent circular analysis of

results. Matches were checked at species, genus, and family level.

BLAST

We used BLAST from GenBank in Geneious 10.0.9 2017 by default to look for

matches of the sequences. The parameters included were the database search from

Nucleotide collection, program set as the MegaBlast, results displayed as a Hit table,

and retrieve Matching region with Maximum hits of 100. The search was set to provide a

list of 100 closest nucleotide sequences, from which we selected the first suggestion

that obtained the highest Bit-score or Max score (name when the BLAST search is done

from the GenBank website). Most of the time the E value was 0 (or closest to zero) and

the Query cover 100%. Matches were checked at species, genus and family level.

The identification analyses in both approaches (BOLD and Blast) followed the

assumption that species from the databases are correctly identified; however, this

approach might not be accurate in cases where the most similar species were wrongly

identified from the databases.

Dung beetles (Scarabaeinae) case study

The Scarabaeinae subfamily was selected due to the high representation of this

group in the samples of Coleoptera (42%) and from the total insect samples (15%). In

order to evaluate the accuracy of species identification suggested by BOLD, we

consulted taxonomists with expertise in the identification of the Scarabaeidae family.

They were asked to reanalyzed the results and mention if they agree with the species

and genus identification suggested by BOLD and to explain why, as follows: Correctly

Identified Genus, Correctly Identified Species, Probable Correctly Identified Species,

Incorrect Identification (Table 1). When available, specimens deposited at the IAVH-E

collection and photographs uploaded to BOLD were checked for specimens to

corroborate the taxonomy, as field notes from the collectors.

Table 1. List of categories used by the taxonomist to define the identification

results suggested by BOLD with the Scarabaeinae subfamily.

Category	Definition
Correctly identified genus	Agreement between the taxonomist and BOLD at the genus level
Correctly identified species	Agreement between the taxonomist and BOLD at the species level
Probable correctly identified species	Agreement between the taxonomist morphospecies and the species suggested by BOLD (the genus matched and probable species)
Incorrect identification	Disagreement between the taxonomist and BOLD (the genus or species did not match)

Statistical analysis

Chi-square analyses were performed between match results for BOLD and

GenBank at the studied taxonomic levels (species, genera and family) to evaluate if

there was a significant difference between the platforms performance. An additional

series of Chi-square analyses were performed to identify differences in the platforms by

each order of insect at the species, genus, and family level. We adjusted our

significance threshold at each taxonomic level based on the number of orders

compared using the Bonferroni Correction. Flagged records were removed from the

analysis.

The non-parametric Kruskal-Wallis test in PAST, Version 4.05 [29] was used to

determine any significant differences of medians for BOLD % match of the correct

Scarabaeinae categories against incorrect identification. Assumptions for an ANOVA

were not fulfilled. Dunn's post hoc was carried out to identify differences between

categories.

Results

Description of the data

A total of 1160 sequences of insects from Colombia were generated since 2019. From

this set, 1088 sequences were barcode compliant and placed in 708 BINS (Table 2). Of

these 708 BINS, 500 were identified as unique (Fig 2). The most frequent BINS were all

in Coleoptera, BOLD:ADN3981 *Paulosawaya* sp. (11 specimens), BOLD:ADO6609

Paulosawaya ursina (Blanchard, 1850) (10 specimens), and BOLD:ADJ9394

Ptilodactylidae (9 specimens). From the specimens with barcode compliant 33 were

flagged as problematic records in most instances due to suspected contamination,

specimen mix-up or misidentification.

Fig 2. Summary of Barcode index numbers (BINS) obtained by orders of insects.

A total of 500 BINs were identified as unique and 208 BINs as non-unique in the

platform BOLD Systems.

Table 2. Summary of specimens and barcodes data by orders after removal of flagged records.

Order	Specimens #	Flagged record	Family count	Specimen count IDed Family	Unknown	Genera count	Specimen count IDed Genus	Species count	Specimen count IDed Species	Barcode compliant	Species count	BINS	Unique Bins	Non-Unique Bins
Coleoptera	411	14	25	397	0	57	271	63	117	369	57	218	184	34
Diptera	277	14	24	257	6	4	22	0	N/A	257	0	165	116	49
Hemiptera	82	4	11	78	0	0	N/A	0	N/A	68	0	58	56	2
Hymenoptera	201	2	10	199	0	16	47	2	3	176	2	142	112	30
Lepidoptera	166	0	8	161	5	57	154	79	128	163	73	104	18	86
Odonata	17	0	6	17	0	14	17	13	15	16	13	15	8	7
Psocodea	5	0	4	5	0	0	N/A	N/A	N/A	5	N/A	5	5	0
Strepsiptera	1	0	1	1	0	1	1	0	N/A	1	0	1	1	0
Total	**1160**	**34**	**89**	**1115**	**11**	**149**	**512**	**157**	**263**	**1055**	**145**	**708**	**500**	**208**

Coleoptera

S1 Table and S2 Table; Barcodes were obtained from 369 out of 411

coleopteran specimens and placed in 218 BINS, from which 184 (84.4%) were identified

as unique in BOLD. The samples were assigned to 25 families, 57 genera and 63

species.

BOLD search engine: The search at 80% provided identifications for a total of

395 specimens. A match in agreement with the taxonomist was found for 337 (85%)

specimens at the family level. For genus, the search resulted in 189 (70%) specimens

of 271 and for species in 57 (49%) specimens of 117 total.

BLAST: The search provided identifications for a total of 379 specimens. A match

in agreement with the taxonomist was found for 294 (78%) specimens at the family

level. For genus, the search resulted in 83 (33%) specimens of 255 and for species in

20 (20%) specimens of 101 total.

Diptera

S1 Table and S2 Table; Barcodes were obtained from 257 out of 277 dipteran

specimens and placed in 165 BINS, from which 116 (70.3%) were identified as unique

in BOLD. The samples were assigned to 24 families and 4 genera. There were 6

specimens not assigned to any category.

BOLD search engine: The search at 80% provided identifications for a total of

245 specimens. A match in agreement with the taxonomist was found for 202 (82%)

specimens at the family level, and for genus, 16 of 18 specimens (89%) matched the

identification of the taxonomist. No species were identified by the taxonomist.

BLAST: The search provided identifications for a total of 241 specimens. A match

in agreement with the taxonomist was found for 184 (76%) specimens at the family

level, and for genus 13 of 14 specimens (93%) matched the identification of the

taxonomist. No species were identified by the taxonomist.

Hemiptera

S1 Table and S2 Table; Barcodes were obtained from 68 out of 82 hemipteran

specimens and placed in 58 BINS, from which 56 (97%) were identified as unique in

BOLD. The samples were assigned to 11 families.

BOLD search engine: The search at 80% provided identifications for a total of 77

specimens. A match in agreement with the taxonomist was found for 66 (86%)

specimens at the family level. No genera or species were identified by the taxonomist.

BLAST: The search provided identifications for a total of 67 specimens. A match

in agreement with the taxonomist was found for 55 (82%) specimens at the family level.

No genera or species were identified by the taxonomist.

Hymenoptera

S1 Table and S2 Table; Barcodes were obtained from 176 out of 201

hymenopteran specimens and placed in 142 BINS, from which 112 (78.9%) were

identified as unique in BOLD. The samples were assigned to 10 families, 16 genera and

2 species.

BOLD search engine: The search at 80% provided identifications for a total of

199 specimens. A match in agreement with the taxonomist was found for 193 (97%)

specimens at the family level. At the genus level, the search resulted in 46 (98%) of 47

specimens and 3 of 3 (100%) specimens at the species level.

BLAST: The search provided identifications for a total of 182 specimens. A match

in agreement with the taxonomist was found for 176 (97%) specimens at the family

level. At the genus level, the search resulted in 38 (97%) of 39 specimens and 0 of 3

specimens at the species level.

Lepidoptera

S1 Table and S2 Table; Barcodes were obtained from 163 out of 166

lepidopteran specimens and placed in 104 BINS, from which 18 (17%) were identified

as unique in BOLD. The samples were assigned to 8 families, 57 genera and 79

species, 5 not assigned to any category.

BOLD search engine: The search at 80% provided identifications for a total of

161 specimens. A match in agreement with the taxonomist was found for 147 (91%)

specimens at the family level. For genus, the search resulted in 140 (91%) specimens

of 154 and for species in 86 (68%) specimens of 127 total.

BLAST: The search provided identifications for a total of 154 specimens. A match

in agreement with the taxonomist was found for 143 (93%) specimens at the family

level. For genus, the search resulted in 117 (79%) specimens of 148 and for species in

49 (49%) specimens of 99 total.

Odonata

S1 Table and S2 Table; Barcodes were obtained from 16 out of 17 Odonata specimens and placed in 15 BINS, from which 8 (53.3%) were identified as unique in BOLD. The samples were assigned to 6 families, 14 genera and 13 species.

BOLD search engine: The search at 80% provided identifications for a total of 17 specimens. A match in agreement with the taxonomist was found for 17 (100%) specimens at the family level. For genus, the search resulted in 13 (76%) specimens of 17 and for species in 6 (46%) specimens of 13 total.

BLAST: The search provided identifications for a total of 17 specimens. A match in agreement with the taxonomist was found for 17 (100%) specimens at the family level. For genus, the search resulted in 9 (56%) specimens of 16 and for species in 3 (20%) specimens of 15 total.

Psocodea

S1 Table and S2 Table; Barcodes were obtained from 5 out of 5 psocodean specimens and placed in 5 BINS, each identified as unique in BOLD (100%). The samples were assigned to 4 families.

BOLD search engine: The search at 80% provided identifications for a total of 5 specimens. A match in agreement with the taxonomist was found for 2 (40%) specimens at the family level. No genera or species were identified by the taxonomist.

BLAST: The search provided identifications for a total of 4 specimens. A match in agreement with the taxonomist was found for 1 (25%) specimen at the family level. No genera or species were identified by the taxonomist.

Strepsiptera

S1 Table and S2 Table; A barcode was obtained from one male specimen and placed in 1 new BIN. The sample was identified to the genus *Caenocholax*.

BOLD search engine: The search at 80% provided identification for the 1 specimen. A match in agreement with the taxonomist was found for it (100%) at the family and genus level. No species were identified by the taxonomist.

BLAST: The search provided identification for the 1 specimen. A match in agreement with the taxonomist was found for it (100%) at the family and genus level. No species were identified by the taxonomist.

Performance of BOLD and Blast (GenBank) identification based on taxonomist match

The performance of BOLD vs BLAST for identification of specimens was significantly greater with BOLD for the pooled comparison of all species ($x2 = 29.5$, df = 1, $P < 0.001$), genera ($x2 = 67.7$, df = 1, $P < 0.001$), and families ($x2 = 8.3$, df = 1, $P = 0.004$). Comparisons of the performance of BOLD and BLAST by Orders at the different taxonomic levels revealed significantly more matches with BOLD (Fig 3), at the family level (Fig 3A) in Coleoptera ($x2 = 7.7$, df = 1, $P = 0.006$), at genus level (Fig 3B) in Coleoptera ($x2 = 72.8$, df = 1, $P < 0.001$) and Lepidoptera ($x2 = 8.4$, df = 1, $P = 0.004$) and at the species level (Fig 3C) in Coleoptera ($x2 = 19.8$, df = 1, $P < 0.001$) and Lepidoptera ($x2 = 7.7$, df = 1, $P = 0.006$). No differences were found for the other orders tested at the different taxonomic categories. Same results were found when analyses were performed with matches to data from our project, the only comparison that

changed was at the genus level where no significant differences were found for

Lepidoptera (S3 Table).

Fig 3. Comparison by order of performance of BOLD Systems and BLAST for

identification of specimens in percentages. Correct matches for BOLD Batch ID

engine at 80% are shown in dark gray. Correct matches from MegaBlast search by

default (GenBank) are shown in light gray. *Is next to orders where Chi-square analyzed

with Bonferroni correction was significant. The sample size is displayed on top of each

column. A) by families. B) by genera C) by species.

Case of study Scarabaeinae subfamily (Coleoptera:

Scarabaeidae)

A total of 172 specimens were analyzed. The samples in the subfamily

Scarabaeinae were assigned to 16 genera and 27 species (S1 Table).

BOLD search engine: The search at 80% provided identifications for the 172

specimens (Fig 4A, S2 Table). A match in agreement with the taxonomist was found for

82 (80%) specimens of 103 at the genus level. For species, the search resulted in 55

(78%) specimens of 69 total. Based on this information, these records were classified

as 137 (80%) in agreement with the taxonomist and 35 (20%) in disagreement (Table 3,

Fig 4A).

Table 3. Summary of matches between the taxonomist and BOLD identifications for the subfamily of dung

beetles Scarabaeinae. The first section (I) corresponds to the initial matches performed in agreement with the taxonomist

after the search engine in BOLD at 80%. The second section (II) in grey includes the categories used by the taxonomist to

reevaluate the identifications provided by BOLD in (I) and the final matches in agreement.

	Categories	Count	%	Matches BOLD & Taxonomist	Count	Total	%	Difference % (disagreement with taxonomist)
(I) BOLD search engine identifications				Genus	82	103	80	
				Species	55	69	78	
				Sum agreement	**137**	172	**80**	20
(II) Taxonomist corroborated identifications – *de novo*	Correctly identified genus	52	30	Genus	52	73	71	
	Correctly identified species	77	45	Species	95	99	96	
	Probable correctly identified species	18	10					
	Incorrect identification	25	15	**Sum agreement**	**147**	172	**85**	15

Fig 4. Results of the reanalysis performed by the taxonomist in the subfamily of dung beetles Scarabaeinae (Coleoptera: Scarabaeidae). A) Comparison between the identification (ID) matches before (I, BOLD IDs) and after (II, taxonomist – *de novo* IDs) reanalyzing the data. B) Percentage of the categories assigned by the taxonomist to the identifications suggested with BOLD's search engine at 80%.

Then, we corroborated *de novo* the identification for these 172 records of dung beetle with a taxonomist (Fig 4B). The *de novo* specimens were assigned to the same 16 genera and 27 species (70 specimens), in addition to 31 morphospecies (genus and code "H", 97 specimens) (Table 4 and S4 Table). The new review resulted in 95 records (55%) correctly matched the taxonomist species identification or was deemed likely to be the species (Correctly identified species and Probable correctly identified species categories). At the genus level, 52 records (30%) correctly matched the taxonomist genus identification but either were not accurately matched with the species or the species identification was not provided (Correctly identified genus category). There were 25 records (15%) at the genus level which did not match the identification of the taxonomist and arguments against BOLD identification were provided by the taxonomist (Incorrect identification category) (S4 Table). Therefore, the records previously classified as 137 (80%) increased to 147 (85%) in agreement with the taxonomist and were reduced from 35 (20%) to 25 (15%) in disagreement based on the BOLD search engine at 80% (Table 3, Fig 4A).

Table 4. List of Scarabaeinae species identified by BOLD in agreement with the taxonomist's *de novo* identification. Morphospecies with genus and code "H" are included [24]. Only categories for correct species and probably correct species are presented here.

Process ID	BIN	Current ID	% Match	Overlap (bp)	Taxonomist's *de novo* ID	Match Species BOLD	Match BIN	Match notes
CBIHM686-18	BOLD:ACW4694	*Ateuchus*	93.4	568	*Ateuchus aeneomicans*	*Ateuchus aeneomicans*	BOLD:AAW8107	Correct species
CBIHI151-18	BOLD:ADL3091	*Canthon juvencus*	99.85	567	*Canthon juvencus*	*Canthon juvencus*	BOLD:ADL3091	Correct species
CBIHM706-18	BOLD:ADL3091	*Canthon juvencus*	99.85	567	*Canthon juvencus*	*Canthon juvencus*	BOLD:ADL3091	Correct species
CBIHM705-18	BOLD:ADL3091	*Canthon juvencus*	99.69	566	*Canthon juvencus*	*Canthon juvencus*	BOLD:ADL3091	Correct species
CBIHM633-18		*Canthon*	98.2	563	*Sylvicanthon aequinoctialis* [30]	*Canthon aequinoctialis*	BOLD:ABA2649	Correct species
CBIHM627-18	BOLD:ABA2649	*Canthon*	98.01	561	*Sylvicanthon aequinoctialis*	*Canthon aequinoctialis*	BOLD:ABA2649	Correct species
CBIHM635-18	BOLD:ABA2649	*Canthon*	97.55	570	*Sylvicanthon aequinoctialis*	*Canthon aequinoctialis*	BOLD:ABA2649	Correct species
CBIHM636-18	BOLD:ABA2649	*Canthon*	97.55	570	*Sylvicanthon aequinoctialis*	*Canthon aequinoctialis*	BOLD:ABA2649	Correct species
CBIHM632-18	BOLD:ABA2649	*Canthon*	97.25	570	*Sylvicanthon aequinoctialis*	*Canthon aequinoctialis*	BOLD:ABA2649	Correct species
CBIHM631-18	BOLD:ABA2647	*Canthon*	100	570	*Sylvicanthon aequinoctialis*	*Canthon aequinoctialisASolis1*	BOLD:ABA2647	Correct species
CBIHM637-18	BOLD:ABA2647	*Canthon*	100	570	*Sylvicanthon aequinoctialis*	*Canthon aequinoctialisASolis1*	BOLD:ABA2647	Correct species
CBIHM638-18	BOLD:ABA2647	*Canthon*	100	570	*Sylvicanthon aequinoctialis*	*Canthon aequinoctialisASolis1*	BOLD:ABA2647	Correct species
CBIHM634-18	BOLD:ABA2647	*Canthon*	99.85	570	*Sylvicanthon aequinoctialis*	*Canthon aequinoctialisASolis1*	BOLD:ABA2647	Correct species
CBIHM629-18	BOLD:ABA2649	*Canthon*	98.17	570	*Sylvicanthon aequinoctialis*	*Canthon aequinoctialisASolis3*	BOLD:ABA2649	Correct species
CBIHM628-18	BOLD:ABA2649	*Canthon*	98.08	570	*Sylvicanthon aequinoctialis*	*Canthon aequinoctialisASolis3*	BOLD:ABA2649	Correct species
CBIHM639-18	BOLD:ABA2649	*Canthon*	97.86	570	*Sylvicanthon aequinoctialis*	*Canthon aequinoctialisASolis3*	BOLD:ABA2649	Correct species
CBIHM747-18	BOLD:ACW3356	*Canthon subhyalinus*	100	570	*Canthon subhyalinus*	*Canthon subhyalinus*	BOLD:ACW3356	Correct species
CBIHM673-18	BOLD:ACW3356	*Canthon subhyalinus*	100	570	*Canthon subhyalinus*	*Canthon subhyalinus*	BOLD:ACW3356	Correct species
CBIHM674-18	BOLD:ACW3356	*Canthon subhyalinus*	100	570	*Canthon subhyalinus*	*Canthon subhyalinus*	BOLD:ACW3356	Correct species

CBIHM727-18	BOLD:ACW3356	*Canthon subhyalinus*	100	570	*Canthon subhyalinus*	*Canthon subhyalinus*	BOLD:ACW3356	Correct species
CBIHM744-18	BOLD:ACW3356	*Canthon subhyalinus*	100	567	*Canthon subhyalinus*	*Canthon subhyalinus*	BOLD:ACW3356	Correct species
CBIHM718-18	BOLD:ACW3356	*Canthon subhyalinus*	99.24	570	*Canthon subhyalinus*	*Canthon subhyalinus*	BOLD:ACW3356	Correct species
CBIHM716-18	BOLD:ACW3356	*Canthon subhyalinus*	98.93	570	*Canthon subhyalinus*	*Canthon subhyalinus*	BOLD:ACW3356	Correct species
CBIHM717-18	BOLD:ACW3356	*Canthon subhyalinus*	98.93	570	*Canthon subhyalinus*	*Canthon subhyalinus*	BOLD:ACW3356	Correct species
CBIHM741-18	BOLD:ACW3356	*Canthon*	100	570	*Canthon subhyalinus*	*Canthon subhyalinus*	BOLD:ACW3356	Correct species
CBIHI156-18	BOLD:ADO2303	*Canthon triangularis*	100	570	*Canthon triangularis*	*Canthon triangularis*	BOLD:ADO2303	Correct species
CBIHI157-18	BOLD:ADO2303	*Canthon triangularis*	100	570	*Canthon triangularis*	*Canthon triangularis*	BOLD:ADO2303	Correct species
CBIHM659-18	BOLD:ADL4806	*Coprophanaeus corythus*	97.24	540	*Coprophanaeus corythus*	*Coprophanaeus corythus*	BOLD:ACW3300	Correct species
CBIHM687-18	BOLD:ADL4607	*Coprophanaeus gamezi*	94.65	570	*Coprophanaeus gamezi*	*Coprophanaeus gamezi*	BOLD:ACW3623	Correct species
CBIHI019-17	BOLD:ADJ7028	*Deltochilum mexicanum*	95.26	570	*Deltochilum mexicanum*	*Deltochilum mexicanum*	BOLD:AAP9113	Correct species
CBIHI159-18	BOLD:ACW3574	*Dichotomius*	95.61	570	*Dichotomius agenor*	*Dichotomius agenor*	BOLD:ABA2814	Correct species
CBIHM648-18	BOLD:ACW3574	*Dichotomius*	95.61	570	*Dichotomius agenor*	*Dichotomius agenor*	BOLD:ABA2814	Correct species
CBIHM667-18	BOLD:ACW3574	*Dichotomius*	95.36	570	*Dichotomius agenor*	*Dichotomius agenor*	BOLD:ABA2814	Correct species
CBIHM666-18	BOLD:ACW3574	*Dichotomius*	95.28	570	*Dichotomius agenor*	*Dichotomius agenor*	BOLD:ABA2814	Correct species
CBIHM646-18	BOLD:ACW3574	*Dichotomius*	95.23	570	*Dichotomius agenor*	*Dichotomius agenor*	BOLD:ABA2814	Correct species
CBIHM647-18	BOLD:ACW3574	*Dichotomius*	95.21	567	*Dichotomius agenor*	*Dichotomius agenor*	BOLD:ABA2814	Correct species
CBIHM645-18	BOLD:ACW3574	*Dichotomius*	95.12	570	*Dichotomius agenor*	*Dichotomius agenor*	BOLD:ABA2814	Correct species
CBIHM665-18	BOLD:ACW3574	*Dichotomius*	95.12	570	*Dichotomius agenor*	*Dichotomius agenor*	BOLD:ABA2814	Correct species
CBIHI144-18	BOLD:AAC9021	*Digitonthophagus gazella*	100	570	*Digitonthophagus gazella*	*Digitonthophagus gazella*	BOLD:AAC9021	Correct species
CBIHM640-18	BOLD:ADL2649	*Eurysternus caribaeus*	96.48	570	*Eurysternus caribaeus*	*Eurysternus caribaeus*	BOLD:ABA7207	Correct species
CBIHI145-18	BOLD:ACW4232	*Eurysternus caribaeus*	100	570	*Eurysternus caribaeus*	*Eurysternus caribaeus*	BOLD:ACW4232	Correct species
CBIHI146-18	BOLD:ACW4232	*Eurysternus caribaeus*	100	570	*Eurysternus caribaeus*	*Eurysternus caribaeus*	BOLD:ACW4232	Correct species
CBIHI147-18	BOLD:ACW4232	*Eurysternus caribaeus*	100	570	*Eurysternus caribaeus*	*Eurysternus caribaeus*	BOLD:ACW4232	Correct species
CBIHI023-17	BOLD:ADJ7677	*Eurysternus foedus*	95.57	570	*Eurysternus foedus*	*Eurysternus foedus*	BOLD:ABU6799	Correct species
CBIHI030-17	BOLD:AAX0271	*Onthophagus acuminatus*	100	570	*Onthophagus acuminatus*	*Onthophagus acuminatusAS1*	BOLD:AAX0271	Correct species
CBIHM653-18	BOLD:ADL2546	*Eurysternus mexicanus*	100	570	*Eurysternus mexicanus*	*Eurysternus mexicanus*	BOLD:ADL2546	Correct species

CBIHM654-18	BOLD:ADL2546	*Eurysternus mexicanus*	100	570	*Eurysternus mexicanus*	*Eurysternus mexicanus*	BOLD:ADL2546	Correct species
CBIHM655-18	BOLD:ADL2546	*Eurysternus mexicanus*	100	570	*Eurysternus mexicanus*	*Eurysternus mexicanus*	BOLD:ADL2546	Correct species
CBIHM656-18	BOLD:ADL2546	*Eurysternus mexicanus*	100	570	*Eurysternus mexicanus*	*Eurysternus mexicanus*	BOLD:ADL2546	Correct species
CBIHM660-18	BOLD:ADL2546	*Eurysternus mexicanus*	100	570	*Eurysternus mexicanus*	*Eurysternus mexicanus*	BOLD:ADL2546	Correct species
CBIHM661-18	BOLD:ADL2546	*Eurysternus mexicanus*	100	570	*Eurysternus mexicanus*	*Eurysternus mexicanus*	BOLD:ADL2546	Correct species
CBIHM662-18	BOLD:ADL2546	*Eurysternus mexicanus*	100	570	*Eurysternus mexicanus*	*Eurysternus mexicanus*	BOLD:ADL2546	Correct species
CBIHM657-18	BOLD:ADL2546	*Eurysternus mexicanus*	99.83	570	*Eurysternus mexicanus*	*Eurysternus mexicanus*	BOLD:ADL2546	Correct species
CBIHM698-18	BOLD:ADL4924	*Malagoniella astyanax*	94.34	570	*Malagoniella astyanax*	*Malagoniella astyanax*	BOLD:ABA3615	Correct species
CBIHM701-18	BOLD:ACW3925	*Ontherus appendiculatus*	99	558	*Ontherus appendiculatus*	*Ontherus appendiculatus*	BOLD:ACW3925	Correct species
CBIHM703-18	BOLD:ACW3925	*Ontherus appendiculatus*	99	570	*Ontherus appendiculatus*	*Ontherus appendiculatus*	BOLD:ACW3925	Correct species
CBIHM699-18	BOLD:ACW3925	*Ontherus appendiculatus*	98.95	570	*Ontherus appendiculatus*	*Ontherus appendiculatus*	BOLD:ACW3925	Correct species
CBIHM702-18	BOLD:ACW3925	*Ontherus appendiculatus*	98.5	558	*Ontherus appendiculatus*	*Ontherus appendiculatus*	BOLD:ACW3925	Correct species
CBIHI160-18		*Onthophagus curvicornis*	99.82	552	*Onthophagus curvicornis*	*Onthophagus curvicornis*	BOLD:ADM9496	Correct species
CBIHI142-18	BOLD:ADM9497	*Onthophagus curvicornis*	98.17	570	*Onthophagus curvicornis*	*Onthophagus curvicornis*	BOLD:ADM9497	Correct species
CBIHI161-18	BOLD:ADM9497	*Onthophagus curvicornis*	98.17	570	*Onthophagus curvicornis*	*Onthophagus curvicornis*	BOLD:ADM9497	Correct species
CBIHI162-18	BOLD:ADM9496	*Onthophagus curvicornis*	99.82	567	*Onthophagus curvicornis*	*Onthophagus curvicornis*		Correct species
CBIHM719-18	BOLD:ADL5334	*Onthophagus lebasi*	99.84	569	*Onthophagus lebasi*	*Onthophagus lebasi*	BOLD:ADL5334	Correct species
CBIHM735-18	BOLD:ADL5334	*Onthophagus lebasi*	99.84	567	*Onthophagus lebasi*	*Onthophagus lebasi*	BOLD:ADL5334	Correct species
CBIHM726-18	BOLD:ADL5334	*Onthophagus lebasi*	99.35	567	*Onthophagus lebasi*	*Onthophagus lebasi*	BOLD:ADL5334	Correct species
CBIHM721-18	BOLD:ADL5334	*Onthophagus lebasi*	98.61	566	*Onthophagus lebasi*	*Onthophagus lebasi*	BOLD:ADL5334	Correct species
CBIHI152-18	BOLD:ADM9873	*Phanaeus bispinus*	96.94	570	*Phanaeus bispinus*	*Phanaeus bispinus*	BOLD:ACW4591	Correct species
CBIHI153-18	BOLD:ADN3451	*Phanaeus haroldi*	98.62	570	*Phanaeus haroldi*	*Phanaeus haroldi*	BOLD:ADN3451	Correct species
CBIHI154-18	BOLD:ADN3451	*Phanaeus haroldi*	98.62	570	*Phanaeus haroldi*	*Phanaeus haroldi*	BOLD:ADN3451	Correct species
CBIHM712-18	BOLD:ACW3786	*Sulcophanaeus leander*	99.83	570	*Sulcophanaeus leander*	*Sulcophanaeus leander*	BOLD:ACW3786	Correct species
CBIHM713-18	BOLD:ACW3786	*Sulcophanaeus leander*	99.68	570	*Sulcophanaeus leander*	*Sulcophanaeus leander*	BOLD:ACW3786	Correct species
CBIHM711-18	BOLD:ACW3786	*Sulcophanaeus leander*	99.66	570	*Sulcophanaeus leander*	*Sulcophanaeus leander*	BOLD:ACW3786	Correct species

CBIHM729-18	BOLD:ADL4592	*Uroxys micros*	100	504	*Uroxys micros*	*Uroxys micros*	BOLD:ADL4592	Correct species
CBIHM731-18	BOLD:ADL4592	*Uroxys micros*	100	501	*Uroxys micros*	*Uroxys micros*	BOLD:ADL4592	Correct species
CBIHM740-18	BOLD:ADL4592	*Uroxys micros*	100	504	*Uroxys micros*	*Uroxys micros*	BOLD:ADL4592	Correct species
CBIHM746-18	BOLD:ADL4592	*Uroxys micros*	99.83	513	*Uroxys micros*	*Uroxys micros*	BOLD:ADL4592	Correct species
CBIHM739-18	BOLD:ADL4592	*Uroxys micros*	99.16	494	*Uroxys micros*	*Uroxys micros*	BOLD:ADL4592	Correct species
CBIHI025-17	BOLD:ADK7115	*Canthidium*	95.39	567	*Canthidium* sp. 21H	*Canthidium centrale*	BOLD:ABU9359	Probably correct species
CBIHM650-18	BOLD:ACW2842	*Canthon*	94.97	567	*Canthon* sp. 06H	*Canthon cyanellus*	BOLD:ADN2169	Probably correct species
CBIHM651-18	BOLD:ACW2842	*Canthon*	94.97	567	*Canthon* sp. 06H	*Canthon cyanellus*	BOLD:ADN2169	Probably correct species
CBIHM663-18	BOLD:ACW2842	*Canthon*	94.9	570	*Canthon* sp. 06H	*Canthon cyanellus*	BOLD:ADN2169	Probably correct species
CBIHM664-18	BOLD:ACW2842	*Canthon*	94.9	570	*Canthon* sp. 06H	*Canthon cyanellus*	BOLD:ADN2169	Probably correct species
CBIHM649-18	BOLD:ACW2842	*Canthon*	94.85	567	*Canthon* sp. 06H	*Canthon cyanellus*	BOLD:ADN2169	Probably correct species
CBIHM742-18	BOLD:ACW4541	*Canthon*	96.83	570	*Canthon* sp. 09H	*Canthon caelius*	BOLD:ABU7368	Probably correct species
CBIHM668-18	BOLD:ACW4541	*Canthon*	96.62	567	*Canthon* sp. 09H	*Canthon caelius*	BOLD:ABU7368	Probably correct species
CBIHM669-18	BOLD:ACW4541	*Canthon*	96.62	567	*Canthon* sp. 09H	*Canthon caelius*	BOLD:ABU7368	Probably correct species
CBIHM670-18	BOLD:ACW4541	*Canthon*	96.62	567	*Canthon* sp. 09H	*Canthon caelius*	BOLD:ABU7368	Probably correct species
CBIHM730-18	BOLD:ACW4541	*Canthon*	96.62	567	*Canthon* sp. 09H	*Canthon caelius*	BOLD:ABU7368	Probably correct species
CBIHM732-18	BOLD:ACW4541	*Canthon*	96.6	570	*Canthon* sp. 09H	*Canthon caelius*	BOLD:ABU7368	Probably correct species
CBIHM820-20		*Canthon*	95.11	408	*Canthon* sp. 09H	*Canthon caelius*	BOLD:ABU7368	Probably correct species
CBIHM683-18	BOLD:ADL3950	*Canthon*	95.01	570	*Canthon* sp. 23H	*Canthon bispinus*	BOLD:ADR8659	Probably correct species
CBIHM684-18	BOLD:ADL3950	*Canthon*	95.01	570	*Canthon* sp. 23H	*Canthon bispinus*	BOLD:ADR8659	Probably correct species
CBIHI148-18	BOLD:AAX0252	*Onthophagus*	98.04	570	*Onthophagus* sp. 07H	*Onthophagus haematopus*	BOLD:AAX0252	Probably correct species
CBIHI150-18	BOLD:AAX0252	*Onthophagus*	98.04	570	*Onthophagus* sp. 07H	*Onthophagus haematopus*	BOLD:AAX0252	Probably correct species
CBIHI149-18	BOLD:AAX0252	*Onthophagus*	97.88	570	*Onthophagus* sp. 07H	*Onthophagus haematopus*	BOLD:AAX0252	Probably correct species

419 Correctly identified species had a percentage match range in BOLD from 93.4 to

420 100 (Fig 5). Probable correctly identified species ranged from 94.85 to 98.04. Correctly

421 identified genus range was between 87.7 and 100%. Incorrect identification

422 percentages range from 83.59 to 99.83. The categories were significantly different

423 (Kruskal-Wallis H = 73.4, P < 0.001). The correctly identified species (P < 0.001),

424 probable correctly identified species (P < 0.05) and correctly identified genus (P < 0.05)

425 categories were different from the incorrect identifications.

427 **Fig 5. Boxplot of Scarabaeinae case of study showing the range of BOLD**

428 **percentage matches for the categories: Correctly Identified Species, Probable**

429 **Correctly Identified Species, Correctly Identified Genus, and Incorrect**

430 **Identification.** The median is represented by a line, the mean is represented by an X

431 markers and outliers are the white dots.

432 # Discussion

433 BOLD systems outperformed BLAST providing more accurate insect identifications.

434 This overall pattern was found for all taxonomic levels evaluated (species, genus and

435 family). Although differences were found between taxa, BOLD consistently provided

436 higher identifications for the order Coleoptera at all taxonomic levels and for

437 Lepidoptera at genus and species level. For instance, Hymenoptera and Odonata

438 performed similarly for both repositories at all taxonomic levels, alike most of the other

439 orders at the family level when analyzed separately. These results are partially in

440 disagreement with Meiklejohn et al [21] which reported no significant differences

between BOLD and GenBank for insect samples identification at genus and species

level. A further review of their data showed that correct identifications from both

platforms were closer than estimated previously [22]. Although we report different

results in the performance of BOLD compared to GenBank in this study, a key factor to

consider is the magnitude of insect data involved. Meiklejohn et al [21], who studied 17

specimens in 12 orders at genus and species level using COI whereas we included

1160 specimens in eight orders at family, genus and species level. BOLD Systems is a

platform developed with the intention of providing species identification and in that

sense does not perform equally to GenBank, but both are the main public repositories

often used to provide identifications. Often BOLD is considered to contain data more

curated, but GenBank data quality is good overall [22,31]. Differences in identifications

of taxa can be considered due to differential taxonomic curation of the databases and

the lack of barcodes of reference from Colombia. However, we could not identify a clear

pattern with the lack of references. The most represented groups were Coleoptera and

Lepidoptera. The group that performed the best with identifications was Coleoptera and

it was also the order that provided the most unique BINS, in contrast with Lepidoptera

that provided the least unique BINS also outperformed GenBank at the species and

genus level, but no differences were found at family.

A great potential of Colombian insect biodiversity was found from BINs counting

with 70% of the sequences assigned to a unique BIN. Each BIN represents a cluster of

species or an operational taxonomic unit (OTU) that is compared against the database

assigning the barcodes to known clusters (BIN) or creating a new cluster (unique BIN)

[18]. Unique BINs were found across all the orders of insects and Coleoptera provided

the highest count of new BINs followed by Diptera and Hymenoptera, some of the

hyperdiverse orders of insects. However, Lepidoptera was the order with highest non-

unique BINs. This finding is likely attributable to a great effort of DNA sampling directed

to Lepidoptera that displays 4848 barcodes from the country with 1474 BINs [32]; in

comparison to Coleoptera with 318 sequences in 82 BINs, Diptera 2646 with 339 BINs

and Hymenoptera 2179 with 240 BINs [32].

Furthermore, our discovery of 500 unique BINs from our dataset of 1088

sequences showcases the need for further sampling and identification in order to add

more species into these databases and increase the accuracy of matches. However,

this would require the implementation of a systematic method of collecting samples

across Colombia, a process that could take a considerable length of time and amount of

labor. Of our seven collecting methods used in this study, the malaise traps accounted

for 62.6%, and we would suggest a targeted approach by placing these traps in

strategic areas across Colombia. We expect that a larger sample will continue providing

a higher number of unique BINs, but also, we see a necessary taxonomic effort to

match all of this genetic diversity with the insect species already described from

Colombia. Nevertheless, the implementation of DNA barcodes in the country will bring

an opportunity to recognize any hidden diversity. As it has been suggested that BINs

underestimate by a 10% the species due to lack of power within close species

[11,33,34]. Although Meier et al [35] suggested awareness on the use of BINS as

identification tool specially when use for species descriptions. Therefore, an integrative

approach with morphology and genetic diversity can boost the recognition of species.

A closer look to the subfamily Scarabaeinae displayed a great performance of

identification at different levels. The percentage of similarity for a match (BOLD match

percentage) to identify species correctly in Scarabaeinae was in a wide range between

93 to a 100%. Although correct matches were statistically significant the range was not

perfect, as shown at least three specimens were incorrectly identified with values higher

than 94%. These outliers do not reduce the potential of genetic identification tools. The

high percentages could be due to specimen mix-up. Overall, the identification success

rate of Scarabaeinae was 85% (30% towards genera and 55% towards species) when

combining the barcode results with a taxonomist effort (Table 3, Fig 4A). The species

identification improved from 78% to 96% (or in the overall from 30% to 55%) with this

taxonomist *de novo* identifications (Table 3, Fig 4A). Other studies of identification of

beetles using barcodes found species matches as high as 92.1% of their samples when

considered traditionally identified species and BINS and as high as 98.3% of the

samples when haplotypes were considered [36]. In comparison, our identification

percentage was lower with previous studies. These can be due to the lack of sequence

records of the species in the database. Pentisaari et al [36] suggested identification

failures due to difficulty with morphological characters and controversial taxonomic

status of species. Taxonomic input is crucial for the function of this tool, also if the

species is not represented in the database there is no possible match to occur. In fact, a

few suggestions provided by BOLD for incorrect species identifications were for close

related species or species within their group of species. For example, *Dichotomius*

andresi and *Dichotomius protectus* were matched with *Dichotomius satanas*, a

morphologically related species in the "*Dichotomius satanas* species group". Similarly,

Eurysternus hypocrita was identified as *Eurysternus olivaceus*, a species in the

"*Eurysternus velutinus* species group". The specimen identified as *Phanaeus pyrois*

was matched as *Phanaeus malyi*, a species that was long considered synonymous with

P. pyrois. Currently, *P. malyi* has been revalidated and registered in Colombia without a

specific location [37].

The distribution range of species could be an additional factor as observed with

the dung beetles data where BOLD identify all species correctly in six genera

(*Coprophanaeus, Digitonthophagus, Malagoniella, Onthophagus, Phanaeus* and

Sulcophanaeus), eight genera partially correct with some of the species identified

correctly but not all (*Ateuchus, Canthidium, Canthon, Deltochilum, Dichotomius,*

Eurysternus, Ontherus and *Uroxys*) and failed completely for the genera

Pseudocanthon (1 specimen) and *Scybalocanthon* (6 specimens). The correctly

identified species were obtained for species with a wide distribution and shared

between South America and Central America, such as *Canthon juvencus*, *Canthon*

subhyalinus, *Eurysternus caribaeus*, *Eurysternus foedus*, *Eurysternus mexicanus*

Onthophagus acuminatus, Onthophagus lebasi, and *Uroxys micros*, most likely due to

representativeness of COI sequences of these species from Costa Rica [38]. These

species are found across countries or ecosystems (e. g. dry and humid forests, natural

and disturbed ecosystems). In contrast, the identification was incorrect for some species

with restricted distributions in Colombia such as *Canthon arcabuquensis, Dichotomius*

andresi, *Uroxys cuprescens*, and *Deltochilum susanae,* the latest was first identified as

the morphospecies *Deltochilum* sp. 12H [39] and later described as a new species [40]

or species with Andean distributions as in *Dichotomius protectus*, *Eurysternus*

marmoreus and *Ontherus brevicollis* [41–46]. In the case of Scarabaeinae dung beetles,

the barcode results helped to confirm at least five species that were assigned as

morphospecies in the dung beetles of Colombia "Reference Collection" hosted at of

IAVH Entomological Collection. The diversity of dung beetle's subfamily Scarabaeinae

in Colombia exceeds by a large number of species that of neighboring countries such

as Panama, Ecuador or Brazil. In general, a defined and well-identified species in

Colombia has other species, morphologically similar and taxonomically closer, that have

been assigned as morphospecies. This is why the Reference Collection was created,

which so far houses about 85% percent of species of Colombia as morphospecies. We

see the barcode as a fundamental integrative taxonomy tool, which will help to outline

these morphospecies and define which of those are new species for science, and to

improve the final taxonomic list of this important group of beetles. Overall, there are still

challenges to overcome and more research is recommended in this topic. The reference

databases keep improving and updating their resources and tools for users [20,31].

BOLD provided reliable identifications for the tested groups of insects overcoming

challenges such as a small genetic reference of insects for a highly diverse country.

548 Acknowledgements

We are grateful to our funding sources: A Minciencias postdoctoral fellowship Fondo

Nacional de Financiamiento para la Ciencia, la Tecnología y la Innovación "Francisco

José de Caldas" call No. 848-2019 and Humboldt Institute (awarded to NBB), Colombia

BIO Agreement # FP 44842-109-2016 (IAvH No. 16-062) between Colciencias (now

Minciencias) and Humboldt Institute, and Santander BIO inter-agency Agreement #2243

(IAvH No. 17-199) with the General System of Royalties of the Santander government,

Colombia's National Planning Department (BPIN 2017000100046), Humboldt Institute

and Industrial University of Santander (BPIN 2017000100046). We would like to thank

the following researchers for their fieldwork efforts collecting the samples and/or for

identifications of samples: H Arenas, C Beltran, C Bota, J Cardenas, C Castro, A

Clavijo, Y Correa, A Diaz, D Espitia, G Fagua, A González, D Hincapie, A Lopera, J

Mendivil, R Morales, EE Palacio, A Parrales, P Pulido, JD Suaza, EA Tenorio, E Torres,

X. Villalba. We want to thank to the valuable Entomological collections where these

samples were deposited: the Entomological and Tissue collection of the Institute

Alexander von Humboldt (IAvH-E and IAvH-CT), Museo Francisco Luis Gallego

(MEFLG) at Universidad Nacional de Colombia, sede Medellín, Museo Javeriano de

Historia Natural (MPUJ) at Pontificia Universidad Javeriana and Colección

Entomológica Universidad de Antioquia (CEUA). Thank you to the IAvH Conservation

Genetic lab members: P Montoya, P Pulido, N Franco, V Sandoval, C Quiroga, and Y

Castañeda for their valuable comments and improvements to the manuscript.

569 Author Contributions

Conceptualization: Nathalie Baena-Bejarano, Mailyn Gonzalez.

Data curation: Nathalie Baena-Bejarano, Catalina Reina, Diego E. Martínez-Revelo.

Formal analysis: Nathalie Baena-Bejarano.

Investigation: Nathalie Baena-Bejarano, Diego E. Martínez-Revelo, Claudia A. Medina,

Jhon Cesar Neita, Catalina Reina, Eduardo Tovar, Sandra Uribe, Mailyn Gonzalez.

Methodology: Nathalie Baena-Bejarano, Catalina Reina, Diego E. Martínez-Revelo,

Claudia A. Medina, Eduardo Tovar, Mailyn Gonzalez.

Project administration: Nathalie Baena-Bejarano, Mailyn Gonzalez, Jhon Cesar Neita.

Supervision: Mailyn Gonzalez, Jhon Cesar Neita, Nathalie Baena-Bejarano.

Resources: Mailyn Gonzalez, Claudia A. Medina, Jhon Cesar Neita, Eduardo Tovar,

Diego E. Martínez-Revelo.

Writing – original draft Preparation: Nathalie Baena-Bejarano.

Writing – review & editing: Nathalie Baena-Bejarano, Diego E. Martínez-Revelo,

Claudia A. Medina, Catalina Reina, Eduardo Tovar, Jhon Cesar Neita, Mailyn Gonzalez,

Sandra Uribe.

References

1. Díaz S, Settele J, Brondízio E, Ngo H, Guèze M, Agard J, et al. IPBES (2019):

Summary for policymakers of the global assessment report on biodiversity and

ecosystem services of the Intergovernmental Science-Policy Platform on

Biodiversity and Ecosystem Services. Bonn, Germany: IPBES secretariat; 2019 p.

56. Available: https://www.ipbes.net/

2. Barnosky AD, Matzke N, Tomiya S, Wogan GOU, Swartz B, Quental TB, et al. Has

the Earth's sixth mass extinction already arrived? Nature. 2011;471: 51–57.

doi:10.1038/nature09678

3. CBD. Convention on Biological Diversity. 2019 [cited 2 Sep 2019]. Available:

https://www.cbd.int/

4. Janzen DH, Hallwachs W. Perspective: Where might be many tropical insects? Biol

Conserv. 2019;233: 102–108. doi:10.1016/j.biocon.2019.02.030

5. Sánchez-Bayo F, Wyckhuys KAG. Worldwide decline of the entomofauna: A review

of its drivers. Biol Conserv. 2019;232: 8–27. doi:10.1016/j.biocon.2019.01.020

6. IUCN. The IUCN Red List of Threatened Species. 2022 [cited 4 Sep 2022].

Available: https://www.iucnredlist.org

7. Arbeláez-Cortés E. Knowledge of Colombian biodiversity: published and indexed.

Biodivers Conserv. 2013;22: 2875–2906. doi:10.1007/s10531-013-0560-y

8. Morinière J, Cancian de Araujo B, Lam AW, Hausmann A, Balke M, Schmidt S, et

al. Species Identification in Malaise Trap Samples by DNA Barcoding Based on

NGS Technologies and a Scoring Matrix. Fontaneto D, editor. PLOS ONE.

2016;11: e0155497. doi:10.1371/journal.pone.0155497

9. Arias-Penna DC, Whitfield JB, Janzen DH, Hallwachs W, Dyer LA, Smith MA, et al.

A species-level taxonomic review and host associations of Glyptapanteles

(Hymenoptera, Braconidae, Microgastrinae) with an emphasis on 136 new reared

species from Costa Rica and Ecuador. ZooKeys. 2019;890: 1–685.

doi:10.3897/zookeys.890.35786

10. Sharkey MJ, Janzen DH, Hallwachs W, Chapman EG, Smith MA, Dapkey T, et al.

Minimalist revision and description of 403 new species in 11 subfamilies of Costa

Rican braconid parasitoid wasps, including host records for 219 species. ZooKeys.

2021;1013: 1–665. doi:10.3897/zookeys.1013.55600

11.   Janzen DH, Hallwachs W, Pereira G, Blanco R, Masis A, Chavarria MM, et al.

Using DNA-barcoded Malaise trap samples to measure impact of a geothermal

energy project on the biodiversity of a Costa Rican old-growth rain forest. Genome.

2020;63: 407–436. doi:10.1139/gen-2020-0002

12.   Geiger M, Moriniere J, Hausmann A, Haszprunar G, Wägele W, Hebert P, et al.

Testing the Global Malaise Trap Program – How well does the current barcode

reference library identify flying insects in Germany? Biodivers Data J. 2016;4:

e10671. doi:10.3897/BDJ.4.e10671

13.   Steinke D, Breton V, Berzitis E, Hebert PDN. The School Malaise Trap Program:

Coupling educational outreach with scientific discovery. PLOS Biol. 2017;15:

e2001829. doi:10.1371/journal.pbio.2001829

14.   Kutty SN, Wang W, Ang Y, Tay YC, Ho JKI, Meier R. Next-Generation identification

tools for Nee Soon freshwater swamp forest, Singapore. Gard Bull Singap.

2018;70: 155–173. doi:10.26492/gbs70(suppl.1).2018-08

15.   Karlsson D, Hartop E, Forshage M, Jaschhof M, Ronquist F. The Swedish Malaise

Trap Project: A 15 Year Retrospective on a Countrywide Insect Inventory.

Biodivers Data J. 2020;8: e47255. doi:10.3897/BDJ.8.e47255

16.   Srivathsan A, Lee L, Katoh K, Hartop E, Kutty SN, Wong J, et al. ONTbarcoder and

MinION barcodes aid biodiversity discovery and identification by everyone, for

everyone. BMC Biol. 2021;19: 217. doi:10.1186/s12915-021-01141-x

17. Ratnasingham S, Hebert PDN. bold: The Barcode of Life Data System

(http://www.barcodinglife.org). Mol Ecol Notes. 2007;7: 355–364.

doi:10.1111/j.1471-8286.2007.01678.x

18. Ratnasingham S, Hebert PDN. A DNA-Based Registry for All Animal Species: The

Barcode Index Number (BIN) System. Fontaneto D, editor. PLoS ONE. 2013;8:

e66213. doi:10.1371/journal.pone.0066213

19. Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ. Basic local alignment search

tool. J Mol Biol. 1990;215: 403–410. doi:10.1016/S0022-2836(05)80360-2

20. Schoch CL, Ciufo S, Domrachev M, Hotton CL, Kannan S, Khovanskaya R, et al.

NCBI Taxonomy: a comprehensive update on curation, resources and tools.

Database. 2020;2020: baaa062. doi:10.1093/database/baaa062

21. Meiklejohn KA, Damaso N, Robertson JM. Assessment of BOLD and GenBank –

Their accuracy and reliability for the identification of biological materials. Fugmann

SD, editor. PLOS ONE. 2019;14: e0217084. doi:10.1371/journal.pone.0217084

22. Pentinsaari M, Ratnasingham S, Miller SE, Hebert PDN. BOLD and GenBank

revisited – Do identification errors arise in the lab or in the sequence libraries?

Kuntner M, editor. PLOS ONE. 2020;15: e0231814.

doi:10.1371/journal.pone.0231814

23. Becker JJ, Sandwell DT, Smith WHF, Braud J, Binder B, Depner J, et al. Global

Bathymetry and Elevation Data at 30 Arc Seconds Resolution: SRTM30_PLUS.

Mar Geod. 2009;32: 355–371. doi:10.1080/01490410903297766

24. González-Alvarado AF, Torres E, Medina CA. Escarabajos coprófagos

(Coleoptera: Scarabaeidae: Scarabaeinae) de bosques secos colombianos en la

Colección Entomológica del Instituto Alexander von Humboldt. Biota Colomb.

2015;16: 88–95.

25. Ivanova NV, Dewaard JR, Hebert PDN. An inexpensive, automation-friendly

protocol for recovering high-quality DNA: Technical Note. Mol Ecol Notes. 2006;6:

998–1002. doi:10.1111/j.1471-8286.2006.01428.x

26. Folmer O, Black M, Hoeh W, Lutz R, Vrijenhoek R. DNA primers for amplification of

mitochondrial cytochrome c oxidase subunit I from diverse metazoan invertebrates.

Mol Mar Biol Biotechnol. 1994;3: 294–299.

27. Hebert PDN, Penton EH, Burns JM, Janzen DH, Hallwachs W. Ten species in one:

DNA barcoding reveals cryptic species in the neotropical skipper butterfly *Astraptes*

fulgerator. Proc Natl Acad Sci. 2004;101: 14812–14817.

doi:10.1073/pnas.0406166101

28. Kearse M, Moir R, Wilson A, Stones-Havas S, Cheung M, Sturrock S, et al.

Geneious Basic: An integrated and extendable desktop software platform for the

organization and analysis of sequence data. Bioinformatics. 2012;28: 1647–1649.

doi:10.1093/bioinformatics/bts199

29. Hammer O, Harper DAT, Ryan PD. PAST: Paleontological Statistics Software

Package for Education and Data Analysis. Palaeontol Electron. 4: 9.

30. Cupello M, Vaz-de-Mello FZ. A monographic revision of the Neotropical dung

beetle genus Sylvicanthon Halffter & Martínez, 1977 (Coleoptera: Scarabaeidae:

Scarabaeinae: Deltochilini), including a reappraisal of the taxonomic history of

'Canthon sensu lato.' Eur J Taxon. 2018. doi:10.5852/ejt.2018.467

31. Madden MJL, Young RG, Brown JW, Miller SE, Frewin AJ, Hanner RH. Using DNA

barcoding to improve invasive pest identification at U.S. ports-of-entry. Labra M,

editor. PLOS ONE. 2019;14: e0222291. doi:10.1371/journal.pone.0222291

32. BOLD systems public data portal - record list Database. 2022 [cited 21 Jul 2022].

Available: https://www.boldsystems.org/

33. Janzen DH, Hallwachs W. DNA barcoding the Lepidoptera inventory of a large

complex tropical conserved wildland, Area de Conservacion Guanacaste,

northwestern Costa Rica. Genome. 2016;59: 641–660. doi:10.1139/gen-2016-0005

34. Janzen DH, Hallwachs W, Blandin P, Burns JM, Cadiou J-M, Chacon I, et al.

Integration of DNA barcoding into an ongoing inventory of complex tropical

biodiversity. Mol Ecol Resour. 2009;9: 1–26. doi:10.1111/j.1755-

0998.2009.02628.x

35. Meier R, Blaimer BB, Buenaventura E, Hartop E, Rintelen T, Srivathsan A, et al. A

re-analysis of the data in Sharkey et al.'s (2021) minimalist revision reveals that

BINs do not deserve names, but BOLD Systems needs a stronger commitment to

open science. Cladistics. 2022;38: 264–275. doi:10.1111/cla.12489

36. Pentinsaari M, Hebert PDN, Mutanen M. Barcoding Beetles: A Regional Survey of

1872 Species Reveals High Identification Success and Unusually Deep

Interspecific Divergences. Fontaneto D, editor. PLoS ONE. 2014;9: e108651.

doi:10.1371/journal.pone.0108651

37. Moctezuma V, Halffter G. Taxonomic revision of the Phanaeus endymion species

group (Coleoptera: Scarabaeidae), with the descriptions of five new species. Eur J

Taxon. 2021;747: 1–71. doi:10.5852/ejt.2021.747.1333

38. Solís Á, Kohlmann B. Checklist and distribution atlas of the Scarabaeinae

(Coleoptera: Scarabaeidae) of Costa Rica. Zootaxa. 3482: 1–32.

doi:10.11646/zootaxa.3482.1.1

39. Martínez-Revelo DE, Castro-Moreno C, Medina CA. Escarabajos coprófagos de la

cuenca alta y media del río Bita, Vichada, Colombia. Biota Colomb. 2018;19: 226–

235. doi:10.21068/c2018.v19n01a15

40. González-Alvarado A, Vaz-de-Mello FZ. Towards a comprehensive taxonomic

revision of the Neotropical dung beetle subgenus Deltochilum (Deltohyboma) Lane,

1946 (Coleoptera: Scarabaeidae: Scarabaeinae): Division into species-groups.

Zhang F, editor. PLOS ONE. 2021;16: e0244657.

doi:10.1371/journal.pone.0244657

41. Génier F. A revision of the Neotropical genus Ontherus Erichson (Coleoptera:

Scarabaeidae, Scarabaeinae). Mem Entomol Soc Can. 1996;128: 3–170.

doi:10.4039/entm128170fv

42. Camero Rubio E, Lobo JM. The Distribution of the Species of *Eurysternus* Dalman,

1824 (Coleoptera: Scarabaeidae) in America: Potential Distributions and the

Locations of Areas to be Surveyed. Trop Conserv Sci. 2012;5: 225–244.

doi:10.1177/194008291200500210

43. Camero Rubio E, Lobo JM. Distribución conocida y potencial de las especies del

género Eurysternus Dalman, 1824 (Coleoptera: Scarabaeidae) de Colombia. Bol

SEA. 2010; 257–264.

44. Sarmiento-Garcés R, Amat-García G. Escarabajos del género Dichotomius Hope

1838 (Scarabaeidae: Scarabaeinae) en Colombia. Fauna de Colombia. Monografía

4. Bogotá, Colombia: Universidad Nacional de Colombia; 2014.

doi:10.13140/2.1.4692.9926

45. Medina C, González A. Escarabajos coprófagos de la subfamilia Scarabaeinae. In:

Pizano C, García H, editors. El Bosque Seco Tropical en Colombia. Instituto de

Investigación de Recursos Biológicos Alexander von Humboldt (IAvH); pp. 195–

213.

46. Chamorro W, Marin-Armijos D, Asenjo A, Vaz-De-Mello FZ. Scarabaeinae dung

beetles from Ecuador: a catalog, nomenclatural acts, and distribution records.

ZooKeys. 2019;826: 1–343. doi:10.3897/zookeys.826.26488

Supporting information captions

S1 Table. Project dataset with the insect records.

S2 Table. Records of insects by orders with matches from BOLD Batch ID engine

and BLAST (GenBank).

S3 Table. Chi-squared tests. A) Dataset analyses including BOLD's first suggestion

with the highest percentage match and base pair overlap. This dataset includes

matches to sequences on our project. B) Alternative dataset analyses including next

higher percentage match and base pair overlap. This excludes matches to sequences

from our project. *Is next to orders where Chi-square was significant. **Is next to orders

where Chi-square analyzed with Bonferroni correction was significant.

S4 Table. List of Scarabaeinae in other categories as determined by the

taxonomist.

Figure

Figure

Figure

A

B

Figure

Figure