Changes in protein dynamical parameters derived from quasi-elastic neutron scattering spectra

induced by coherent scattering: A molecular dynamics simulation study

Tatsuhito Matsuo[†]

Institute for Quantum Life Science, National Institutes for Quantum Science and Technology,

4-9-1 Anagawa, Chiba, 263-8555, Japan

[†] Corresponding author: Tatsuhito Matsuo, Ph.D.

Tel: +81 (0)70 3943 3439

Email: matsuo.tatsuhito@qst.go.jp

Abstract

Protein dynamics at the sub-nanosecond timescale and the Å length-scale has widely been studied

using quasi-elastic neutron scattering (QENS). In almost all QENS studies, where the measured

QENS spectra contain both incoherent and coherent scattering contributions, analysis of the QENS

spectra is carried out under the assumption that the contribution of coherent scattering is negligible

whereas a study using polarization analysis has shown that the coherent scattering accounts for

more than 10% of the total scattering intensity. The effects of coherent scattering on the values of

dynamical parameters of proteins obtained by analysis of QENS spectra remain unclear. Here,

molecular dynamics (MD) simulation on hen egg white lysozyme was used to investigate this issue.

QENS spectra containing only incoherent scattering and those containing both incoherent and

coherent scattering were calculated from the MD trajectory. Dynamical parameters were then

extracted from the two simulated QENS spectra. Comparison of the resultant dynamical parameters

has shown that the error in the values of the dynamical parameters induced by coherent scattering

is at most 6%. This error is unlikely to significantly affect the results of QENS studies that investigate

the relative changes in protein dynamics caused by different physicochemical conditions such as

temperature unless dynamical parameters need to be determined with high precision at the

absolute scale.

Introduction

Quasi-elastic neutron scattering (QENS) is an experimental technique that has extensively been used to study dynamical behavior of proteins at the sub-nanosecond timescale and the Å length-scale [1–6]. The incoherent neutron scattering cross-section of a hydrogen (H) atom is more than 40 times larger than its isotope, deuterium, and any other atoms found in proteins [7], and about half of the total number of atoms constituting protein molecules are H atoms. As a result, experimentally measured QENS spectra are dominated by scattering of H atoms of the protein molecules. Because H atoms are distributed almost uniformly in space, dynamical information studied by QENS reflects the motions of H atoms averaged over all the H atoms in protein molecules.

Dynamical information is extracted from the experimentally measured QENS spectra $S_{exp}(\vec{Q}, \omega)$, where $\vec{Q}$ and ω are the momentum transfer and the energy transfer of neutrons upon scattering, respectively, which is called dynamic structure factor. In typical QENS measurements without polarization analysis, which is a technique that can separate coherent and incoherent scattering [8], it is assumed that the scattering contribution of the coherent dynamic structure factor $S_{coh}(\vec{Q}, \omega)$ is negligible and thus $S_{exp}(\vec{Q}, \omega)$ equals to the incoherent dynamic structure factor $S_{inc}(\vec{Q}, \omega)$, for which various theoretical models and phenomenological fitting equations are formulated [1]. On the other hand, Gaspar et al. have experimentally shown by polarization analysis that there exists more than 10% of Q-dependent coherent contribution over a wide Q range in $S_{exp}(\vec{Q})$, where $S_{exp}(\vec{Q}, \omega)$ was

integrated with regards to ω [9]. Although incoherent scattering is the major component of $S_{exp}(\vec{Q})$,

this seminal work has raised a concern about the validity of the dynamical information derived from

the QENS study without polarization analysis. Therefore, it is of critical importance to investigate to

what extent coherent scattering might affect dynamical parameters extracted from $S_{exp}(\vec{Q}, \omega)$

compared with those extracted from $S_{inc}(\vec{Q}, \omega)$.

In this study, this issue was investigated by simulating $S_{inc}(\vec{Q}, \omega)$ and $S_{coh}(\vec{Q}, \omega)$ based on a MD

trajectory of hen egg white lysozyme (HEWL). Dynamical parameters that are typically obtained and

discussed in QENS studies were extracted from the simulated $S_{inc}(\vec{Q}, \omega)$ and $S_{exp}(\vec{Q}, \omega)$ (= $S_{inc}(\vec{Q}, \omega)$

+ $S_{coh}(\vec{Q}, \omega)$). The resultant dynamical parameters were compared to reveal the possible changes in

these values induced by coherent scattering.

Materials and methods

Molecular dynamics simulation

In this study, one of the trajectories of a hen egg white lysozyme (HEWL) molecule (PDB ID: 1AKI)

generated previously was used [10]: The software GROMACS (version 2018.3) [11] was used to

perform molecular dynamics (MD) simulation with the AMBER03 force field [12]. The heavy water

model TIP3P-HW [13] was employed as a solvent molecule. In a cubic box, a HEWL molecule was

placed and solvated such that all edges of the simulation box were $\geq$ 10 Å away from the protein

surface. Counter ions were added to neutralize the charges of the entire system. van der Waals force

and electrostatic interactions were cut off at 8.0 Å [12].

All the exchangeable (labile) hydrogen atoms of HEWL were replaced with deuterium by doubling

the mass of the corresponding hydrogen atoms as in previous studies [14–16]. Whereas this simple

procedure cannot explain an experimentally observed molecular behavior occurring at a long time

scale (400 ns) [17], significant errors were not seen up to 50 ns from the start of the simulation. As

a simulation time, 6 ns was chosen such that the simulation time was short enough not to generate

large errors in protein dynamics and at the same time long enough to extract dynamical parameters

obtained by QENS.

The system was first energy-minimized by the steepest descent method, followed by NVT and NPT

equilibrations, which were carried out for 100 ps each with a time step of 2 fs. The resultant structure

was subjected to a production run for 6 ns at 300 K and at 1 atm, and the first 5 ns of the trajectory

was discarded to extract part of the trajectory which have reached a stable mean square

displacement (MSD). Translational and rotational motions of the whole molecule were removed

from the MD trajectory so that only intramolecular motions are observed. The trajectory was saved

every 2 ps.

Calculation of incoherent and coherent dynamic structure factors from the MD trajectory

The incoherent dynamic structure factor $S_{inc}(\vec{Q}, \omega)$ and the coherent dynamic structure factor $S_{coh}(\vec{Q},$

$\omega)$ were calculated from the MD trajectory through the incoherent intermediate scattering function

$I_{inc}(\vec{Q}, t)$ and the coherent intermediate scattering function $I_{coh}(\vec{Q}, t)$, respectively, as follows:

$I_{inc}(\vec{Q}, t) = \sum_{i=1}^{N} b_{i,inc}^{2} \langle \exp\{i\vec{Q} \cdot (\vec{r_i}(t) - \vec{r_i}(0))\} \rangle,$ (1)

$I_{coh}(\vec{Q}, t) = \sum_{i=1}^{N} \sum_{j=1}^{N} b_{i,coh} b_{j,coh} \langle \frac{\sin(\vec{Q} \cdot (\vec{r_i}(t) - \vec{r_j}(0)))}{\vec{Q} \cdot (\vec{r_i}(t) - \vec{r_j}(0))} \rangle = \sum_{i=j}^{N} b_{i,coh}^{2} \langle \frac{\sin(\vec{Q} \cdot (\vec{r_i}(t) - \vec{r_j}(0)))}{\vec{Q} \cdot (\vec{r_i}(t) - \vec{r_j}(0))} \rangle +$

$2 \sum_{i>j}^{N} b_{i,coh} b_{j,coh} \langle \frac{\sin(\vec{Q} \cdot (\vec{r_i}(t) - \vec{r_j}(0)))}{\vec{Q} \cdot (\vec{r_i}(t) - \vec{r_j}(0))} \rangle,$ (2)

where N is the total number of atoms constituting a protein molecule, and $b_{i,inc}$ and $b_{i,coh}$ are the

incoherent and coherent scattering lengths of the i-th atom, respectively, which are tabulated in

Table 1. $\vec{r_i}(t)$ is the position vector of the i-th atom at time t, which is given by a MD trajectory,

and $\langle \ \rangle$ denotes the thermal average. In the case of solution samples, scattering intensity is

orientationally averaged since protein molecules are randomly oriented, and the orientational

average and the thermal average are independent. The vectors $\vec{Q}$ and $\vec{r_i}(t) - \vec{r_j}(0)$ are thus

treated as their modulus $Q = |\vec{Q}|$ and $|\vec{r_i}(t) - \vec{r_j}(0)|$, respectively.

Table. 1 Incoherent and coherent scattering lengths of atoms used in this study. The values are taken

from the literature [1].

Atom type	b_{coh} [fm]	b_{inc} [fm]
H	-3.742	25.217
D	6.674	4.033
C	6.6484	0
N	9.36	1.98
O	5.805	0
S	2.847	0

$S_{inc}(Q, \omega)$ and $S_{coh}(Q, \omega)$ are the time Fourier transforms of $I_{inc}(Q, t)$ and $I_{coh}(Q, t)$, respectively,

$S_{inc}(Q, \omega) = \int_{0}^{\infty} I_{inc}(Q, t) I_{res}(Q, t) \cos \omega t \, dt,$ (3)

$S_{coh}(Q, \omega) = \int_0^\infty I_{coh}(Q, t) I_{res}(Q, t) \cos \omega t \, dt,$ (4)

where $I_{res}(Q, t)$ is an instrumental resolution function, which defines the time-window of

observation. In this study, this function was represented by a Gaussian function as,

$I_{res}(Q, t) = \exp\left\{-\frac{1}{2}\left(\frac{t}{\tau_{res}}\right)^2\right\},$ (5)

where τ_{res} is the resolution time, which was set to 10 ps in this study. This corresponds to the

energy resolution (FWHM) of 77 µeV. Since $I_{res}(Q, t)$ drops to almost zero at around t = 50 [ps],

$I_{inc}(Q, t)$ and $I_{coh}(Q, t)$ were calculated up to t = 50 [ps] using the MD trajectory with the time

length of 600 ps.

As stated above, the experimentally-observed dynamic structure factor $S_{exp}(Q, \omega)$ is the sum of $S_{inc}(Q$

$\omega)$ and $S_{coh}(Q, \omega)$ of protein molecules, i.e., $S_{exp}(Q, \omega) = S_{inc}(Q, \omega) + S_{coh}(Q, \omega)$. Relative contribution

of incoherent and coherent scattering to $S_{exp}(Q, \omega)$ was investigated through the scattering intensity

$S_{inc}(Q)$ and $S_{coh}(Q)$, where $S_{inc}(Q\ \omega)$ and $S_{coh}(Q, \omega)$ are integrated in the energy direction:

$S_{inc}(Q) = \int_{-\infty}^\infty S_{inc}(Q, \omega) d\omega$ (6)

$S_{coh}(Q) = \int_{-\infty}^\infty S_{coh}(Q, \omega) d\omega$ (7)

$S_{coh}(Q)$ corresponds to a scattering curve obtained by small-angle neutron scattering. Because

$S_{inc}(Q\ \omega)$ and $S_{coh}(Q, \omega)$ were simulated in the energy range between -0.5 meV and 0.5 meV, the

integrations in Eqs. 6 and 7 were conducted in this energy range.

Analysis of the dynamic structure factors

Dynamical parameters that are typically obtained by QENS measurements were extracted by fitting

the simulated $S_{exp}(Q, \omega)$ and $S_{inc}(Q, \omega)$ with the following phenomenological equation:

$S_{exp\,or\,inc}(Q, \omega) = C(Q)S_{loc}(Q, \omega) \otimes R(Q, \omega) + B(Q)$

$= C(Q)[A(Q)\delta(\omega) + (1 - A(Q))L_{loc}(Q, \omega)] \otimes R(Q, \omega) + B(Q),$ $\quad\quad\quad$ (8)

where $C(Q)$ is a scaling factor between the fit and the simulated spectra, and $S_{loc}(Q, \omega)$ represents

the contribution from local atomic motions in a protein molecule. $R(Q, \omega)$ and $B(Q)$ are a

resolution function and a background, respectively. $L_{loc}(Q, \omega)$ is a Lorentzian function describing

local atomic motions, which is described as

$L_{loc}(Q, \omega) = \frac{1}{\pi}\frac{\Gamma_{loc}(Q)}{\omega^2 + \Gamma_{loc}(Q)^2},$ $\quad\quad\quad$ (9)

where $\Gamma_{loc}(Q)$ is the half width at the half maximum (HWHM) of the Lorentzian. The term

including $\delta(\omega)$ is the elastic component and that including $L_{loc}(Q, \omega)$ is the quasi-elastic

component.

Analysis of Q^2-dependence of $\Gamma_{loc}(Q)$ provides the detailed features of the corresponding diffusive

motions. $\Gamma_{loc}(Q)$, which was obtained by the fitting of the simulated QENS spectra using Eq. 8, was

found to be approximated by the jump-diffusion model [18]:

$\Gamma_{loc}(Q) = \frac{D_{loc}Q^2}{1 + D_{loc}Q^2\tau_{loc}},$ $\quad\quad\quad$ (10)

where τ_{loc} and D_{loc} are the residence time and the jump-diffusion coefficient of local atomic

motions, respectively.

Q-dependence of $A(Q)$, the elastic incoherent structure factor (EISF), provides information on the

geometry of local atomic motions. The EISF was fitted by the following equation that describes a

diffusion-inside-a-sphere model [19]:

$A(Q) = p_0 + (1 - p_0)\left(\frac{3j_1(Qa)}{Qa}\right)^2,$ (11)

where p_0 is the immobile fraction of atoms, motions of which are too slow to be resolved at the

energy resolution employed while $(1 - p_0)$ denotes a fraction of atoms resolved at the energy

resolution employed. The latter atoms undergo diffusive motions within a sphere of a radius of a.

$j_1(x)$ is the first-order spherical Bessel function of the first kind.

Results and Discussion

First, $S_{inc}(Q)$ and $S_{coh}(Q)$ of HEWL calculated from the MD trajectory were compared to investigate

the relative fraction of incoherent and coherent scattering in $S_{exp}(Q)$, which corresponds to the

experimentally obtained S(Q) of proteins ($S_{exp}(Q) = S_{inc}(Q) + S_{coh}(Q)$). The results are shown in Fig.

1 (a). It is seen that $S_{inc}(Q)$ is independent of Q values, which is consistent with experimental studies

[9,20,21] while $S_{coh}(Q)$ shows a strong Q-dependence with small peaks at around Q = 0.6 and 1.4

[$Å^{-1}$]. $S_{exp}(Q)$ thus shows slightly larger intensity below Q = 1.0 [$Å^{-1}$] than above Q = 1.0 [$Å^{-1}$], which

is a tendency also observed in $S_{exp}(Q)$ obtained from the experimental QENS spectra on another

protein troponin [21]. $S_{coh}(Q)$ corresponds to the so-called small-angle neutron scattering (SANS)

curve, which is used for structural analysis of proteins, and hence it was compared with a SANS

curve of a crystal structure of HEWL calculated using the program CRYSON [22] (Fig. 1 (b)). The

resultant SANS curve agrees quite well with the simulated $S_{coh}(Q)$, suggesting the validity of the

calculation procedure used in this study. Fig. 1 (c) shows the relative fraction of incoherent and

coherent scattering in the simulated $S_{exp}(Q)$. it is found that the fraction of incoherent scattering

depends on Q values, as reported previously using polarization analysis [9], and it reaches 80–90%

as Q increases. Accordingly, the fraction of coherent scattering is 40% at $Q = 0.4$ [Å^{-1}], above which

analysis of QENS spectra is generally carried out, and then decreases to 10–20% at higher Q values.

Next, the shapes of the simulated $S_{inc}(Q, \omega)$ and $S_{coh}(Q, \omega)$ were directly compared, which has not

previously been reported neither experimentally nor theoretically. For this purpose, $S_{coh}(Q, \omega)$ was

normalized such that its peak intensity at $\omega = 0$ [meV] is equal to that of $S_{inc}(Q, \omega)$. The results are

shown in Fig. 2 for $Q = 0.5, 1.0$, and 1.5 [Å^{-1}]. It is seen that both spectra have essentially almost the

same shape. On the other hand, when one takes a closer look at these spectra as shown in the

insets of Fig. 2, the degree of broadening of the spectra is slightly different between $S_{inc}(Q, \omega)$ and

$S_{coh}(Q, \omega)$ and this difference shows Q-dependence. To investigate the effects of these coherent

scattering contributions on dynamical parameters derived from QENS measurements, analyses of

the simulated $S_{exp}(Q, \omega)$ and $S_{inc}(Q, \omega)$ were carried out as a next step.

The simulated $S_{exp}(Q, \omega)$ and $S_{inc}(Q, \omega)$ were fitted using the phenomenological equation of Eq. 8. A

fitting example is shown in Fig. 3, which shows that both spectra are reproduced in good quality by

this equation. The resultant Q^2-dependences of the widths of the Lorentzian ($\Gamma_{loc}(Q)$) and Q-

dependences of the EISFs are summarized in Fig. 4. The $\Gamma_{loc}(Q)$ was found to be fitted by the jump-

diffusion model (Eq. 10) in both cases as shown in Fig. 4 (a) and (c). Moreover, both EISFs were also

reproduced by the diffusion-inside-a-sphere model (Eq. 11) as shown in Fig. 4 (b) and (d). Slight

differences in the broadening of the QENS spectra between $S_{inc}(Q, \omega)$ and $S_{coh}(Q, \omega)$, which was

shown in Fig. 2, resulted in almost identical values of the $\Gamma_{loc}(Q)$ and the EISF as shown in Fig. 4 (e)

and (f), respectively. Although the relative fraction of coherent scattering is 40% at $Q = 0.4$ [$\mathring{A}^{-1}$] (Fig.

1 (c)), this does not affect the $\Gamma_{loc}(Q)$ and the EISF values (Fig. 4 (e) and (f)). This is because the shape

of the simulated QENS spectra is very similar between $S_{inc}(Q, \omega)$ and $S_{coh}(Q, \omega)$.

Dynamical parameters obtained by the fitting of the simulated QENS spectra, i.e., the jump diffusion

coefficient (D_{loc}), the residence time (τ_{loc}), the radius of a sphere (a), and the immobile fraction of

atoms (p_0), are compared in Fig. 5. It is found that the D_{loc} value increases from $1.54 \pm 0.03 \times 10^{-5}$

cm^2/s to $1.63 \pm 0.03 \times 10^{-5}$ cm^2/s by the contribution of coherent scattering, and τ_{loc} decreases from

11.67 ± 0.07 ps to 11.42 ± 0.07 ps. The a value increases from 1.389 ± 0.004 Å to 1.438 ± 0.003 Å

while the p_0 value increases from 0.722 ± 0.001 to 0.737 ± 0.001 by introducing coherent scattering.

These results indicate that while the immobile fraction of atoms slightly increases, the contribution

of coherent scattering to QENS spectra apparently leads to slightly enhanced protein dynamics.

Using modern neutron spectrometers such as BL02 DNA at J-PARC [23], dynamical parameters can

now be estimated with a precision of 1–2% [21]. On the other hand, the differences in dynamical

parameters caused by coherent scattering discussed in the previous paragraph are 5.8%, 2.2%, 3.5%,

and 2.1% for D_{loc}, τ_{loc}, a, and p_0, respectively. Thus, differences in some dynamical parameters

induced by coherent scattering are beyond the precision of experiments and hence it is essential

to bear in mind that dynamical parameters extracted from QENS spectra measured without

polarization analysis might be varied by several percent compared with those that would be

obtained from purely incoherent QENS spectra, which do not contain the contribution of coherent

scattering. The present study thus suggests that in the case where dynamical parameters need to

be determined at the absolute scale with high precision, QENS measurements with polarization

analysis will be required as already proposed before [9]. On the other hand, in the case where

changes in dynamical behavior of proteins induced by temperature [24,25], pressure [26,27], and

other biological conditions [21,28], are investigated, the widths of the Lorentzian function and the

EISF values generally show much larger changes than those shown in Fig. 4 (e) and (f) as the sample

state is varied. In this case, a several percent of differences in dynamical parameters caused by

coherent scattering would have negligible effects on the conclusion of QENS studies on proteins.

236 To the author's knowledge, this is the first study to investigate the possible effects of coherent

237 scattering on the dynamical parameters obtained by QENS spectra on proteins. In summary,

238 analysis of the simulated QENS spectra, $S_{exp}(Q, \omega)$ and $S_{inc}(Q, \omega)$, has shown that despite the non-

239 negligible contribution of coherent scattering, which accounts for 10–40% depending on the Q-

240 range, the dynamical parameters extracted from the simulated $S_{inc}(Q, \omega)$ are modulated by at most

241 6% by coherent scattering. However, these differences in dynamical parameter values are unlikely

242 to affect the results of QENS studies to investigate the relative changes in protein dynamics caused

243 by different physicochemical conditions of the sample unless dynamical parameters need to be

244 determined with high precision at the absolute scale.

Figure 1. Q-dependencies of the S(Q) profiles of incoherent ($S_{inc}(Q)$) and coherent ($S_{coh}(Q)$)

scattering calculated from the MD trajectory of hen egg white lysozyme (HEWL) in D_2O. The profiles

shown are from only HEWL molecules and do not contain contribution from hydration water and

bulk water. (a): $S_{exp}(Q)$ denotes the sum of $S_{inc}(Q)$ and $S_{coh}(Q)$, which are shown in cyan and magenta,

respectively. This is the S(Q) profile that is measured in typical QENS measurements without nuclear

polarization. (b): Comparison of $S_{coh}(Q)$ calculated from the MD trajectory and that estimated from

the software CRYSON, which calculates a small-angle neutron scattering curve, using a HEWL

structure (PDB ID: 1AKI). (c) The relative fraction of coherent and incoherent scattering in $S_{exp}(Q)$

shown in (a) at each Q value.

Figure 2. Comparisons of incoherent dynamic structure factor $S_{inc}(Q, \omega)$ and coherent dynamic

structure factor $S_{coh}(Q, \omega)$ at Q = 0.5 [Å^{-1}] (a), 1.0 [Å^{-1}] (b), and 1.5 [Å^{-1}] (c). Note that $S_{coh}(Q, \omega)$ is

normalized such that its peak is equal to that of $S_{inc}(Q, \omega)$. The real fraction of $S_{coh}(Q, \omega)$ is as shown

in Fig. 1 (a) and (b). The insets show the magnified views of the corresponding spectra.

273　Figure 3. Examples of the fitting of simulated QENS spectra at Q = 1.0 [Å⁻¹]. (a): QENS spectra

274　containing both incoherent and coherent scattering, which is equal to those obtained from real

275　measurements without nuclear polarization. (b): QENS spectra arising from only incoherent

276　scattering. Grey filled squares represent the simulated QENS spectra, which are denoted as "Target".

277　The upper panels show the residuals between the "Target" and the fitted values.

Figure 4. Q- or Q^2-dependences of the widths ($\Gamma_{loc}(Q)$) of the Lorentzian function the EISF. Open

squares denote the Γ_{loc} values, which were estimated by the fitting of the simulated QENS spectra

containing both incoherent and coherent scattering labeled as "Inc + Coh" (a, b) or those containing

only incoherent scattering labeled as "Inc" (c, d). Solid lines denote the fits using Eq. 11 for the EISF

and Eq. 10 for the widths. Comparisons of Γ_{loc} and the EISF between "Inc + Coh" and "Inc" are shown

in (e) and (f), respectively.

Figure 5. Summary of dynamical parameters extracted from the simulated QENS spectra. Comparisons of the jump diffusion coefficient (a), the residence time (b), the radius of a sphere (c), and the immobile fraction of atoms (d) are shown. Parameter values obtained from the simulated QENS spectra containing both incoherent and coherent scattering are shown in cyan while those obtained from the QENS spectra containing only incoherent scattering are shown in magenta.

References

1. Bée, M. *Quasielastic Neutron Scattering*; Adam Hilger, Bristol and Philadelphia, 1988; ISBN 9780852743713.

2. Gabel, F.; Bicout, D.; Lehnert, U.; Tehei, M.; Weik, M.; Zaccai, G. Protein dynamics studied by neutron scattering. *Q. Rev. Biophys.* **2002**, *35*, 327–367, doi:10.1017/s0033583502003840.

3. Sakai, V.G.; Arbe, A. Quasielastic neutron scattering in soft matter. *Curr. Opin. Colloid Interface Sci.* **2009**, *14*, 381–390, doi:10.1016/j.cocis.2009.04.002.

4. Matsuo, T.; Peters, J. Sub-Nanosecond Dynamics of Pathologically Relevant Bio-Macromolecules Observed by Incoherent Neutron Scattering. *Life* **2022**, *12*, 1259, doi:https://doi.org/10.3390/life12081259.

5. Smith, J.C. Protein dynamics: Comparison of simulations with inelastic neutron scattering experiments. *Q. Rev. Biophys.* **1991**, *24*, 227–291, doi:10.1017/S0033583500003723.

6. Matsuo, T.; Peters, J. Chapter 2: Introduction to incoherent neutron scattering: a powerful technique to investigate the dynamics of bio-macromolecules. In *Analytical Techniques for the Elucidation of Protein Function*; Hojo, H., Sharma, R., Suetake, I., Eds.; John Wiley & Sons, Incorporated, 2022 ISBN 9781119886327.

7. Sears, V.F. Neutron scattering lengths and cross sections. *Neutron News* **1992**, *3*, 26–37, doi:10.1080/10448639208218770.

8. Schärpf, O. Polarization analysis techniques for quasielastic neutron scattering. *Phys. B Condens. Matter* **1992**, *182*, 376–388, doi:https://doi.org/10.1016/0921-4526(92)90041-P.

317   9.   Gaspar, A.M.; Busch, S.; Appavou, M.S.; Haeussler, W.; Georgii, R.; Su, Y.; Doster, W. Using
318   polarization analysis to separate the coherent and incoherent scattering from protein samples.
319   *Biochim Biophys Acta* **2010**, *1804*, 76–82, doi:10.1016/j.bbapap.2009.06.024.

320   10.   Matsuo, T. Effects of heavy water on protein dynamics studied by molecular dynamics
321   simulation: Focusing on dynamical parameters obtained by quasi-elastic neutron scattering.
322   *bioRxiv* **2022**, 2022.09.08.507213, doi:10.1101/2022.09.08.507213.

323   11.   Abraham, M.J.; Murtola, T.; Schulz, R.; Páll, S.; Smith, J.C.; Hess, B.; Lindahl, E. GROMACS: High
324   performance molecular simulations through multi-level parallelism from laptops to
325   supercomputers. *SoftwareX* **2015**, *1–2*, 19–25, doi:https://doi.org/10.1016/j.softx.2015.06.001.

326   12.   Duan, Y.; Wu, C.; Chowdhury, S.; Lee, M.C.; Xiong, G.; Zhang, W.; Yang, R.; Cieplak, P.; Luo, R.;
327   Lee, T.; et al. A point-charge force field for molecular mechanics simulations of proteins based
328   on condensed-phase quantum mechanical calculations. *J Comput Chem* **2003**, *24*, 1999–2012,
329   doi:10.1002/jcc.10349.

330   13.   Linse, J.B.; Hub, J.S. Three- And four-site models for heavy water: SPC/E-HW, TIP3P-HW, and
331   TIP4P/2005-HW. *J. Chem. Phys.* **2021**, *154*, doi:10.1063/5.0050841.

332   14.   Steinbach, P.J.; Loncharich, R.J.; Brooks, B.R. The effects of environment and hydration on
333   protein dynamics: A simulation study of myoglobin. *Chem. Phys.* **1991**, *158*, 383–394,
334   doi:https://doi.org/10.1016/0301-0104(91)87078-A.

335   15.   Guzzi, R.; Arcangeli, C.; Bizzarri, A.R. A molecular dynamics simulation study of the solvent
336   isotope effect on copper plastocyanin. *Biophys. Chem.* **1999**, *82*, 9–22, doi:10.1016/S0301-
337   4622(99)00097-6.

338   16.   Lawrence, C.P.; Skinner, J.L. Flexible TIP4P model for molecular dynamics simulation of liquid
339   water. *Chem Phys Lett* **2003**, *372*, 842–847, doi:10.1016/S0009-2614(03)00526-8.

340   17.   Agarwal, R.; Smith, M.D.; Smith, J.C. Capturing Deuteration Effects in a Molecular Mechanics
341   Force Field: Deuterated THF and the THF-Water Miscibility Gap. *J. Chem. Theory Comput.* **2020**,
342   *16*, 2529–2540, doi:10.1021/acs.jctc.9b01138.

343   18.   Singwi, K.; Sjölander, A. Diffusive Motions in Water and Cold Neutron Scattering. *Phys. Rev.*
344   **1960**, *119*, 863–871, doi:10.1103/PhysRev.119.863.

345   19.   Volino, F.; Dianoux, A.J. Neutron incoherent scattering law for diffusion in a potential of
346   spherical symmetry: general formalism and application to diffusion inside a sphere. *Mol. Phys.*
347   **1980**, *41*, 271–279, doi:10.1080/00268978000102761.

348   20.   Rusevich, L.; Embs, J.; Bektas, I.; Paulsen, H.; Renger, G.; Pieper, J. Protein and solvent dynamics
349   of the water-soluble chlorophyll-binding protein (WSCP). *EPJ Web Conf.* **2015**, *83*, 2016.

350   21.   Matsuo, T.; Tominaga, T.; Kono, F.; Shibata, K.; Fujiwara, S. Modulation of the picosecond
351   dynamics of troponin by the cardiomyopathy-causing mutation K247R of troponin T observed
352   by quasielastic neutron scattering. *Biochim Biophys Acta Proteins Proteom* **2017**, *1865*, 1781–
353   1789, doi:10.1016/j.bbapap.2017.09.007.

354   22.   Svergun, D.I.; Richard, S.; Koch, M.H.; Sayers, Z.; Kuprin, S.; Zaccai, G. Protein hydration in
355   solution: experimental observation by x-ray and neutron scattering. *Proc Natl Acad Sci U S A*

1998, *95*, 2267–2272.

23.   Shibata, K.; Takahashi, N.; Kawakita, Y.; Matsuura, M.; Yamada, T.; Tominaga, T.; Kambara, W.;

Kobayashi, M.; Inamura, Y.; Nakatani, T.; et al. The Performance of TOF near Backscattering

Spectrometer DNA in MLF, J-PARC. *JPS Conf. Proc.* **2015**, *8*, 36022, doi:10.7566/jpscp.8.036022.

24.   Stadler, A.M.; Digel, I.; Embs, J.P.; Unruh, T.; Tehei, M.; Zaccai, G.; Buldt, G.; Artmann, G.M. From

powder to solution: hydration dependence of human hemoglobin dynamics correlated to body

temperature. *Biophys. J.* **2009**, *96*, 5073–5081, doi:10.1016/j.bpj.2009.03.043.

25.   Stadler, A.M.; Demmel, F.; Ollivier, J.; Seydel, T. Picosecond to nanosecond dynamics provide a

source of conformational entropy for protein folding. *Phys Chem Chem Phys* **2016**, *18*, 21527–

21538, doi:10.1039/c6cp04146a.

26.   Peters, J.; Martinez, N.; Lehofer, B.; Prassl, R. Low-density lipoproteins investigated under high

hydrostatic pressure by elastic incoherent neutron scattering. *Eur Phys J E Soft Matter* **2017**, *40*,

68, doi:10.1140/epje/i2017-11558-8.

27.   Peters, J. High Hydrostatic Pressure–A Key Element to Investigate Molecular Dynamics in

Biosystems. *Front. Phys.* **2022**, *9*, 1–7, doi:10.3389/fphy.2021.801539.

28.   Matsuo, T.; Francesco, A. De; Peters, J. Molecular Dynamics of Lysozyme Amyloid Polymorphs

Studied by Incoherent Neutron Scattering. *Front. Mol. Biosci.* **2022**, *8*, 812096,

doi:10.3389/fmolb.2021.812096.

374