Experimental Diets Dictate the Metabolic Benefits of *Lactobacillus* Strains in Obesity

Authors

Ida Søgaard Larsen[1,9] , Béatrice S.-Y. Choi[1], Bandik Föh[2,3], Nanna Ny Kristensen[4], Adia Ouellette[1], Rune Falkenberg Haller[4], Peter Bjarke Olsen[4], Delphine Saulnier[5], Christian Sina[2], Benjamin A. H. Jensen[1,6,7,8], André Marette[1,7]

Affiliations

[1]Quebec Heart and Lung Institute, Faculty of Medicine, and Institute of Nutrition and Functional Foods (INAF), Laval University, Québec, Canada

[2]Institute of Nutritional Medicine, University of Lübeck, Lübeck, Germany

[3]Department of Medicine I, University Hospital Schleswig-Holstein, Lübeck, Germany

[4]Novozymes A/S, Denmark, Krogshoejvej 36, DK-2880 Bagsvaerd, Denmark

[5]Novozymes Berlin GmbH, Berlin, Gustav-Meyer Allee 25, 13355 Berlin, Germany

[6]Department of Biomedical Sciences, Faculty of Health and Medical Sciences, University of Copenhagen, Denmark

[7]Co-senior authors

[8] Lead corresponding author: Benjamin.jensen@sund.ku.dk

[9]Current address: Novozymes A/S, Denmark, Krogshoejvej 36, DK-2880 Bagsvaerd, Denmark

Author Contributions

BAHJ, ISL, NNK, AM and BSYC conceived and designed the studies. ISL, BAHJ, and BSYC carried out the *in vivo* studies. ISL, BSYC, BF, AO, RFH, PBO, and DS generated data. BAHJ designed the diets and supervised all parts of the study; AM and CS supervised parts of the study. ISL integrated the data and wrote the first drafts of the manuscript, which was revised by all the authors, who agreed upon the submitted manuscript.

Conflicts of interest

This study was partly funded by Novozymes A/S (NZ) which provided the bacterial strains and was granted a patent related to the findings (WO20084051). NZ contributed to study design and parts of the analyses but had no role in data integration, interpretation and conclusion.

Funding

BAHJ was supported by Lundbeck Foundation (grant number: R232-2016-2425) and Novo Nordisk Foundation (grant numbers: NNF17OC0026698; NNF21OC0066931). AM was supported by the Canadian Institutes of Health Research (grant numbers: FDN-143247 and MRT-168045) and was the holder of a Pfizer/CIHR partner Chair in the pathogenesis of insulin resistance and cardiovascular diseases. BSYC was funded by doctoral scholarships from Natural Sciences and Engineering Research Council (NSERC) and from the Sentinel North program. BF and CS were supported by the German research society (DFG) IRTG 1911 Project. CS was endowed Chair of Nutritional Medicine supported by Fresenius Kabi. DS was partly supported by a grant from the German Federal Ministry of Education and Research (BMBF) from the Good Bacteria and Bioactive in Industry (GOBI) Consortium (Grant number: BMBF 031B0074A).

Acknowledgment

We thank Joanie Dupont-Morissette, Christine Dallaire, Jacinthe Julien, Laurence Morin, and Jenny Rancourt-Mercier from IUCPQ, Laval University, for excellent technical assistance for *in vivo* protocols. Sincere gratitude to Lea B. S. Hansen for essential bioinformatic input. Additionally, we thank Torsten Schröder from the Institute of Nutritional Medicine for data generation as well as Jeffrey Schultchen from NZ for providing the probiotic strains.

Keywords

The manuscript includes

4 figures

4 supplementary figures

1 table

3 supplementary tables

Abstract

Background: Growing evidence supports the use of probiotics to prevent or mitigate obesity-related dysmetabolism and non-alcoholic fatty liver disease (NAFLD). However, frequent reports of responders versus non-responders to probiotic treatment warrant a better understanding of key modifiers of host-microbe interactions. The influence of host diet on probiotic efficacy, in particular against metabolic diseases, remains elusive.

Method: We fed C57BL6/J mice a low fat reference diet or one of two energy-matched high fat and high sucrose diets for 12 weeks; a classical high fat diet (HFD) and a customized fast food-mimicking diet (FFMD). During the studies, mice fed either obesogenic diet were gavaged daily with one of two probiotic *Lactobacillus* strains, namely *Limosilactobacillus reuteri* (*L. reuteri*) or *Lacticaseibacillus paracasei* subsp. *paracasei* (*L. paracasei*)*,* or vehicle.

Results: The tested probiotics exhibited a reproducible efficacy but dichotomous response according to the obesogenic diets used. Indeed, *L. paracasei* prevented weight gain, improved insulin sensitivity, and protected against NAFLD development in mice fed HFD, but not FFMD. Conversely, *L. reuteri* improved glucoregulatory capacity, reduced NAFLD development, and increased distal gut bile acid levels associated with changes in predicted functions of the gut microbiota exclusively in the context of FFMD-feeding.

Conclusion: We found that the probiotic efficacy of two *Lactobacillus* strains is highly dependent on experimental obesogenic diets. These findings highlight the need to carefully consider the confounding impact of diet in order to improve both the reproducibility of preclinical studies and their clinical research translatability.

Introduction

Since 1975 the global prevalence of overweight and obesity has nearly tripled to ~40% of the adult population (World Health Organization (WHO)., 2016). Obesity is a leading risk factor of cardiometabolic multimorbidity (Kivimäki et al., 2017) including type 2 diabetes (Chatterjee et al., 2017; Fonseca, 2009) and non-alcoholic fatty liver disease (NAFLD) (Fabbrini et al., 2010), also known as metabolic associated fatty liver disease (MAFLD) (Eslam et al., 2020), which affects ~25% of the population worldwide (Younossi et al., 2016). The etiology of obesity is highly complex, involving positive energy balance resulting from poor dietary habits and sedentarity (Hall and Guo, 2017) but also features hereditary and socioeconomic traits (Stunkard et al., 1990; Tyrrell et al., 2016). Although lifestyle interventions such as dietary restrictions may facilitate acute weight loss, this tends to decrease over time resulting in recurrent weight gain (Dulloo and Montani, 2015; Thonusin et al., 2020; Varkevisser et al., 2019; Venditti et al., 2008). Pharmacological therapies against obesity with proven clinical efficacy are emerging, however adverse effects such as nausea are frequently reported among users (Khalil et al., 2020). Surgical strategies, most notably gastric bypass, are more successful yet highly invasive and not spared from potential perioperative complications (Arterburn et al., 2020).

Probiotics - defined as "live microorganisms which when administered in adequate amounts confer a health benefit on the host" (FAO/WHO, 2001) - exhibit promising preclinical potential to reduce obesity, NAFLD, and insulin resistance. Probiotics are therefore considered a non-invasive yet potentially highly effective therapeutic approach against obesity and associated metabolic diseases (Delzenne et al., 2011; Wiciński et al., 2020). Still, evidence from clinical trials is inconsistent and often of low quality, as the probiotic field is loosely regulated (The Lancet Gastroenterology & Hepatology, 2019). This emphasizes a critical need for an enhanced molecular understanding of strain-specific capabilities (Krautkramer et al., 2020) including potential adverse effects, as reported recently for probiotic use in cancer immunotherapy (Spencer et al., 2021). Human studies with probiotic supplementation further reveal responders and non-responders as a common trait, both in terms of colonization and general host effects (Reid et al., 2010; Zmora et al., 2018). Whether the lack of consistent probiotic efficacy in previous studies relates to methodological, genetic and/or environmental factors, such as dietary patterns, remains largely unknown (Reid et al., 2010; Zmora et al., 2018).

A limited number of studies have reported efficacy discrepancies of *Lactobacilli* and other bacterial taxa in mice depending on the diet they consumed. Indeed, *L. helveticus* R0052 induced different effects on anxiety-like behavior depending on whether the mice consumed a fiber-rich chow or high-fat Western diet (Ohland et al., 2013), which was also the case for *L. plantarum* WCFS1's effects on bacterial gene

expression and host intestinal inflammation (Tachon et al., 2014). On a similar note, *Prevotella copri* appears to exhibit opposing effects on glucose metabolism in both humans and mice depending on dietary challenge, being either a fiber-rich low fat diet (Kovatcheva-Datchary et al., 2015) or a low fiber, high fat diet (Pedersen et al., 2016). Importantly, these studies were confounded by dietary differences in energy and fiber content. To our knowledge, no studies have yet examined probiotic efficacy in different but energy-matched obesogenic diets resulting in distinct metabolic phenotypes and disease severity in preclinical murine models. This is critically important to optimize the clinical translatability of preclinical research in the probiotic field.

In this study, we therefore evaluated the efficacy of two *Lactobacillus* strains in mice fed either a standard HFD commonly used in the field or a customized diet designed to exacerbate NAFLD and nonalcoholic steatohepatitis (NASH). We demonstrate that the efficacy of the tested *Lactobacillus* strains on preventing obesity, insulin resistance, and NAFLD development is entirely dependent on the obesogenic diet used, underlining the critical importance of considering the composition of experimental diets in preclinical studies for evaluating the effects of probiotics. Our work further highlights the need to carefully assess dietary patterns and nutritional factors as potential confounders of probiotic efficacy in clinical studies.

Materials & methods:

Animals study design and diets

Three independent studies with identical experimental outlines were carried out using male C57BL6/J mice (Jackson Lab) co-housed three mice per cage. Twelve days prior to the start of the study period the mice were changed from a chow diet to a compositionally defined low fat diet (LFD) during acclimatization. Experimental groups were stratified based on similar average and variance of baseline body weight, fat mass, and additionally for Study 1, 5h fasting blood was sampled for determinations of plasma glucose and insulin levels before starting the experimental diets (described in a later section) without disturbing the social hierarchy within cages (**Figure S1A-E**). During the 12-week study period mice were fed experimental diets from eight weeks of age receiving either a commonly used high fat diet (HFD, D12451, Ssniff Spezialdiäten (Study 1) or Research diets (Study 2), a customized fast food-mimicking diet (FFMD, D12079 mod. Ssniff Spezialdiäten (Study 1 and 3) or continued on the LFD consisting of dietary sources matched to the FFMD (**Fig 1A, Table 1**). Feed intake was monitored and exchanged three times a week and body weight was measured weekly. Body composition was assessed by magnetic resonance (MR) scan (Bruker) prior to, and then every four weeks during the study. Energy uptake was measured at week 11 of the study by monitoring the feed intake and fecal excretion during 24h in clean cages with bedding followed by energy excretion using a calorimeter. From the start of feeding experimental diets, mice were orally gavaged daily 4h into the light cycle with 100 µL of either vehicle (PBS, Gibco) or 10^8 CFU *Limosilactobacillus reuteri,* formerly known as *Lactobacillus reuteri,* DSM 32910 (*L. reuteri*) or *Lacticaseibacillus paracasei* subsp. *paracasei,* formerly classified as *Lactobacillus paracasei* subsp. *tolerans,* DSM 32851 (*L. paracasei*) freshly thawed from stocks stored at -80°C and diluted in PBS (Gibco). After twelve weeks, the mice were anaesthetized with isoflourane (Fresenius Kabi) and euthanized by cervical dislocation in alternating order after 5h of fasting from 2h into the light cycle. Cardiac puncture was performed using a 25G needle and 1 mL syringe coated with EDTA (Sigma-Aldrich). Blood was transferred to Eppendorf tubes containing 1 µL of DPP IV inhibitor (Millipore) and 1 µL of a protease inhibitor cocktail (Sigma) and kept on ice. The samples were centrifuged at 1000 rcf at 4 °C for 10 min and the plasma was placed on dry ice and transferred to -80 °C storage until further processing.

The weight of the liver, white adipose tissues (WATs), and cecum were measured, and the tissues immediately frozen in liquid nitrogen and stored at -80 °C until further processing. Tissues were dissected by the same operator and taken in the same order for all mice in each study. The small intestine (from stomach to cecum) and colon (from cecum to rectum) were kept on a Plexiglas plate cooled by underlying

wet ice throughout the handling time. Duodenum was considered the first 5 cm of the small intestine and the remaining small intestine tissue was divided into 3 parts of equal length. The proximal 3 cm of the distal 1/3 of the small intestine was discarded and the remaining tissue categorized as ileum. Content of the small intestine, cecum, and colon were isolated by mechanical pressure, frozen on dry ice, and subsequently stored at -80 °C. Tissues from the ileum and colon were snap-frozen in liquid nitrogen and stored at -80 °C. The animal studies were conducted in accordance with Canadian Council on Animal Care under the Laval University license 2017-086.

Assessment of glucose homeostasis, oral glucose tolerance, glucose-stimulated insulin, and C-peptide levels

Study 1 baseline glucose homeostasis was assessed in 5h fasted mice the day of study initiation. Blood glucose was measured by sampling blood from the tail vein using a glucometer (OneTouch Vario Flex, LifeScan) and plasma sampled using EDTA prepared capillary tubes (Sarstedt) kept on ice until blood samples were centrifuged for 10 min at 1000 rcf at 4 °C. Plasma insulin was quantified by Mouse Ultrasensitive Insulin ELISA (Alpco) using the manufacturer's protocol.

An oral glucose tolerance test was carried out in week 10 of the experimental protocol. Mice were fasted from 2h into the light cycle for 5h still receiving their usual oral daily gavage. Fasting blood glucose and plasma insulin were measured as described for baseline glucose homeostasis prior to oral gavage with 1.5 µg/g (Study 1 and 3) or 2.0 µg/g (Study 2) lean mass of dextrose. Blood glucose was measured from tail vein puncture at time points 15, 30, 60, 90, and 120 min after dextrose challenge. Blood samples for quantification of plasma insulin levels were sampled in EDTA prepared capillary tubes (Sarstedt) at time points 0, 15, 30, 60, and 120 min and for C-peptide at time points 0, 15, and 30 min post oral glucose challenge. Mice were subcutaneously injected with 0.5 mL saline (Hospira) after the procedure for rehydration. Blood samples were centrifuged for 10 min at 1000 rcf at 4 °C. Plasma insulin was quantified by Mouse ultrasensitive Insulin ELISA (Alpco) and C-peptide by Mouse C-peptide ELISA kit (Crystal Chem) following the manufacturers' protocols.

Liver triacylglycerol (TG) quantification

TG was quantified in snap-frozen liver tissue stored at -80 □C until cryo-grinding in liquid nitrogen and 50 ±5 mg tissue added 0.9 mL of a 2:1 chloroform:methanol solution and homogenized 1 min at 50 os/sec using a TissueLyser LT (Qiagen) with beads. The content was transferred to new tubes and added 0.3 mL methanol and incubated for 2h with rotation. The tubes were centrifuged for 15 min at 4000 rcf at 10°C followed by transfer of 0.825 mL supernatant to new tubes and 0.4 mL chloroform was added. The

samples were vortexed, 0.275 mL of 0.73% NaCl was added and samples were vortexed again for 30 sec followed by centrifugation for 3 min at 4000 rcf at 10°C. The upper phase was aspirated and the tubes rinsed with 0.8 mL Folch solution (Chloroform:Methanol:NaCl 0.58% (3:48:47)) repeatedly for three washes followed by addition of 2-5 drops of methanol until the solution appeared clear after vortexing. The liquid was evaporated using liquid nitrogen and the extracted lipids were resuspended in a solution of isopropanol with 10% TritonX-100 and TG levels quantified by Infinity Triglycerides Reagent (ThermoFisher) and calculated from serial diluted standard curves.

Lipidomics of liver tissue and plasma

Liver and plasma lipids were extracted in a solvent consisting 89.9:10:0.1 of Heptan/IPA/HCOOH and analyzed by liquid chromatography-mass spectrometry (LC-MS) (Thermo Scientific). The MS was operated in negative and positive ion mode electrospray ionization mode. Data was collected from 80-1500 m/z. Lipids were normalized to sample median by log transformation and Pareto scaling using MetaboAnalyst (Chong et al., 2019), which was additionally used to generate heatmaps of the top 40 or 100 features of each measure as indicated in figure legends.

Histopathological scoring of liver tissue

Liver tissue (lobus dexter medialis hepatis) was collected and fixed in 4% paraformaldehyde for three days. After paraffin embedding 3 µm sections were prepared and stained with hematoxylin and eosin (H&E) or Sirius Red. NAFLD activity was assessed blindly in H&E-stained cross-sections using an established scoring system for murine NAFLD (Liang et al., 2014). In short, the levels of hepatocellular hypertrophy, macrovesicular and microvesicular steatosis were determined at 40× to 100× magnification relative to the total liver area and scored as described in **Supplementary Table 1**. Hepatic inflammation was scored by counting the number of inflammatory foci per 1 mm² using a 100× magnification. Inflammatory foci were defined as clusters (not rows) of at least 5 inflammatory cells and the mean of five random fields was used for scoring, as previously described (Liang et al., 2014). Histopathological categories were based on previously described methodology (Liang et al., 2014) where liver slides are categorized with NAFLD if the steatosis score (based on the sum of micro-, macrovesicular steatosis, and hepatocellular hypertrophy scores) ≥ 1 with an inflammation score of 0. Mice are categorized with NASH if steatosis score ≥ 1 and an inflammation score ≥ 1. Livers are considered healthy if the steatosis score = 0 regardless of the inflammation score. Fibrosis was blindly assessed by the Ishak fibrosis stage using Sirius Red-stained cross-sections following previously published methods (Ishak et al., 1995). The scoring system introduced by Ishak *et al*. allows for a sensitive scoring of fibrosis that takes quantity and location of fibrosis into account (Goodman, 2007) and is described further in **Supplementary Table 2**.

Cytokine quantification in liver tissue

Cryo-grinded liver tissues were washed by adding 200 µL ice-cold PBS and protease inhibitors (Sigma-Aldrich), centrifugation at 16,000 x g for 20 sec at 4 °C and aspiration of the liquidphase. This wash step was repeated a total of 10 times. Afterwards 400 µL T-PER Tissue Protein Extraction Buffer (ThermoFisher) and beads were added and vortexed for 2 x 1 min at 50 os/sec. Samples were centrifuged at 4,000 rcf for 1 min at 4 °C and the supernatants transferred to new tubes that were centrifuged again at 13,000 rcf for 10 min at 4 °C. Protein concentration was measured in triplicates using Pierce BCA Protein Assay kit (ThermoFisher) following the manufacturer's instructions. Cytokine levels in samples normalized to the protein concentration were measured using ProcartaPlex Immunoassay (ThermoFisher) following the manufacturer's instructions on the Bio-Plex 200 system (Bio-Rad).

Bile acid (BA) characterization in liver tissue and cecum content

BAs in liver tissue and cecum content were extracted with methanol and 5 µL was injected onto a reverse-phased chromatography (1290 Infinity II, Agilent) and mass spectrometer (6546 Q-TOF/MS, Agilent). The MS was operated in negative ion mode electrospray ionization. Data were collected from 80-1600 m/z. The chromatographic part of the LC-MS-system was set up with a CSH-peptide C18 column (Waters) using a gradient system with Eluent A: water with 0.1% formic acid and Eluent B: Acetonitrile with 0.1% formic acid. A flow of 0.250 mL/min was used, starting at 99.5% Eluent A lowering to 1% within 19 min followed by a new gradient endpoint 99.5% Eluent A at 25.5 min. The gradient was kept isocratic until 30 min and the eluent returned to initial conditions at 26 min. All gradients were linear. Results were related to serial diluted standard curves.

Tissue gene expression by quantitative reverse transcription PCR (RT-qPCR)

RNA from snap-frozen ileum and liver tissues were extracted using the Quick-RNA Miniprep Plus kit (Zymo Research). cDNA was synthesized from 1.5 µg ileum or 2 µg liver RNA using High-Capacity cDNA Reverse Transcriptase kit (Applied Bioscience) following the manufacturer's protocol. qPCR was carried out using 4µl of cDNA, 5µl of Advanced qPCR MasterMix (Wisent Bioproducts) and 0.5µl of each primer (diluted at a concentration of 10 µM) in a total reaction volume of 10 µL with the following cycle setting: 95 °C for 2 minutes, (95 °C for 20 sec, 61.5 °C for 20 sec, 72 °C for 20 sec) x 40 ending with a melting curve: 65 °C to 95 °C. The targets were evaluated in each tissue and accepted in case of a unified peak from the melting curve and amplification efficiency of 100%±11 and $R^2 > 0.99$ from a standard curve. Relative expression was calculated by $2^{\Delta Cq}$ of target Cq to 18S Cq of the sample

accepting replicates with coefficient of variation < 0.02. Target primer sequences are reported in **Supplementary Table 3.**

Microbiota profiling by 16S rRNA gene amplicon sequencing

Fecal samples were collected prior to the start of the studies and every 4 weeks throughout the study period at the same time as body composition assessment 22-24h after the latest oral gavage with *Lactobacillus* or vehicle. During necropsy 2-5h after the latest oral gavage contents of small intestine and colon were collected. All samples were immediately placed on dry ice and stored at -80 °C until DNA extraction using the NucleoSpin 96 soil kit (Macherey-Nagel) following the manufacturer's instructions. Stocks of the administered *Lactobacilli* were processed similarly to the remaining samples. The 16S rRNA gene was amplified over 25 cycles using primers for the V3-V4 region with Illumina adaptors (S-D-Bact-0341-b-S-17: *5'-TCGTCGGCAGCGTCAGATGTGTATAAGAGACAGCCTACGGGNGGCWGCAG-3'and* S-D-Bact-0785-a-A-21: *5'-GTCTCGTGGGCTCGGAGATGTGTATAAGAGACAGGACTACHVGGGTATCTAATCC-3')*(Klindworth et al., 2013). Indices were added in a second PCR over 8 cycles with unique primer combinations using the Nextera XT Index Kit V2 (Illumina). The samples were pooled and cleaned using AMPure XP beads (Beckman Coulter) and the library was sequenced on an Illumina MiSeq desktop sequencer using the MiSeq Reagent Kit V3 (Illumina) for 2x 300 bp paired-end sequencing. The generation of an amplicon sequence variant (ASV) table was carried out as previously described (Larsen et al., 2021). The dataset had a median of 45,468 and a mean of 46,400 reads per sample with a standard deviation of 19,177. Alpha diversity measures and analysis of differential abundances were calculated based on rarefied data. Principal coordinate analysis (PCoA) of microbiota composition was carried out using weighted UniFrac distances. Functions of the microbiota were predicted using the Kyoto Encyclopedia of Genes and Genomes (KEGG) orthology generated by Picrust2 and illustrated with PCoAs using Bray-Curtis distances.

Software and Statistical analysis

The statistical analysis is specified in the figure legend of each figure. Statistical analysis by linear mixed-effects models was carried out in R using the lmerTest package (Kuznetsova et al., 2017) including study, batch, and cage as nested random effects. Statistical significance was illustrated by asterisks based on false discovery rate (fdr) corrected q-values. Two-way analysis of variance (ANOVA) were conducted to analyze the effects of treatments and diet in addition to Bonferroni multiple comparison test to respective vehicle-treated control on the same diet. Categorical data was tested for statistical differences by Kruskal-Wallis and Mann-Whitney tests with Dunn's post hoc for multiple comparisons testing. Statistical

analysis of lipidomics data was carried out by one-way ANOVA with Tukey's post-hoc analysis in the MetaboAnalyst software. The graphical abstract was created with BioRender.com. Bar charts were made using GraphPad Prism 9. Analysis of community-based statistical differences on gut microbiota composition and its predicted functions were carried out by permutational multivariate analysis of variance (PERMANOVA) by the R package vegan (Oksanen et al., 2017) using weighted UniFrac or Bray-Curtis distances, respectively, reporting an average p-value of 100 permutations. The glmer.nb function from the lme4 package (Bates et al., 2014) fitting a generalized linear mixed-effects model (GLMM) for the negative binomial family building on glmer was used for the analysis of differential abundance of bacterial phylae on rarefied data. The top 100 most abundant taxa on each taxonomy rank were used in the analysis. Cage was used as random effect in the model. Multiple testing correction was done with the fdr method. Parameters assessed in independent animal studies were shown in the same graph with experiment-dependent point shapes, indicated in figure legends when appropriate. Study 1 sample size (n) = 9 mice per group except for FFMD + *L. paracasei* group with n = 8. Study 2 n = 11 mice per group. Study 3 n = 12 mice per group except for FFMD + Vehicle group with n = 9. Asterisks identify statistically significant p-values where * = < 0.05, ** = < 0.01 and *** = < 0.001.

Results

Experimental diet determines *Lactobacillus* efficacy on body composition and insulin sensitivity

We ensured equal starting points between the experimental groups and fed the mice one of two energy-matched high fat, high sucrose diets or a low fat reference diet (LFD) for 12 weeks with daily oral gavage of either *L. reuteri, L. paracasei,* or vehicle (**Fig 1A**, **S1A-E, Table 1**). Despite similar feed energy content, vehicle-treated FFMD-fed mice exhibited enhanced body weight-, fat- and lean mass gain compared to their classical HFD-fed counterparts (**Fig 1B-C, Fig S1F**). This trait was closely associated with an increase in energy uptake (**Fig 1D**) and a similar trend for energy intake (p = 0.02, q = 0.11, **Fig 1E**). Of note, fat percentage, subcutaneous, epididymal and retroperitoneal white adipose tissues (WATs) were equally increased by FFMD and HFD, whereas FFMD augmented visceral mesenteric WAT deposition to a higher degree than HFD (**Fig S1G-K**). *L. paracasei* reduced body weight and fat mass gain in HFD-, but not in FFMD-fed mice (**Fig 1B-C, S1G**). This effect of *L. paracasei* on body composition was not explained by a reduction of energy intake or uptake (**Fig 1D-E**).

We next investigated the effect of the *Lactobacilli* on glucose homeostasis after 10 weeks of feeding the obesogenic diets. Importantly, while the classical HFD impaired glucose homeostasis and increased both blood glucose and glucose-stimulated insulin secretion (GSIS) during glucose challenge compared to LFD-fed mice, FFMD-feeding significantly aggravated all of the above (**Fig 1F-J**). Strikingly, FFMD-fed mice more than doubled their fasting insulin levels compared to HFD-fed counterparts (**Fig 1F**). Moreover, although fasting blood glucose was unaffected by the type of obesogenic diet, FFMD-fed mice failed to return to baseline blood glucose levels two hours post challenge accompanied by a 2-3 fold increase in GSIS, suggesting an early onset of frank diabetes in the latter dietary model (**Fig 1G-J**). Notably, *L. reuteri* lowered fasting insulin levels and improved glucose tolerance, with a concomitant trend towards diminished GSIS based on both insulin and c-peptide excursions exclusively in FFMD-fed mice (**Fig 1F-J**, **S1L-M**). In contrast, *L. paracasei* improved glucose tolerance in HFD-fed mice after high dose (**Fig S1N-O**) but not low dose (**Fig 1H-J**) glucose challenge. Collectively, these data demonstrates that the metabolic benefits of two *Lactobacilli* strains are highly dependent on the energy-dense obesogenic diets used.

NAFLD development was diet-dependently mitigated by *Lactobacilli*

As a major glucoregulatory organ, we next assessed hepatic lipid accumulation. Both liver weight *per se* and TG concentration were >3-fold higher in FFMD-fed mice compared to their HFD-fed counterparts (**Fig 2A-C**). In agreement with enhanced liver TGs, FFMD feeding altered the hepatic lipid profile (**S2A**),

and increased both histological prevalence and severity of diet-induced NAFLD, with ~90 % of the animals in the FFMD + Vehicle group having histopathologically-validated NASH, contrasting with only ~45 % of NASH observed in HFD + Vehicle mice (**Fig 2D-E**). Accordingly, FFMD-fed mice developed significantly increased hepatocellular hypertrophy and microvesicular steatosis, as well as excessive hepatic inflammation and moderate fibrosis (**Fig 2F-K, S2C-D**) all pointing towards aggravated liver damage (Tandra et al., 2011) as compared to their HFD-fed counterparts.

L. paracasei prevented diet-induced hepatic lipid accumulation in 17 out of 21 HFD-fed mice, presenting TG levels similar to those of LFD-fed reference mice (**Fig 2C**). A similar trajectory was noted for the hepatic lipid profile, where a subset of the HFD + *L. paracasei* group clustered together with their LFD-fed counterparts (**S2B**). The clustering of the HFD + *L. paracasei* group fully recapitulated the histological steatosis score and NAFLD distribution (**Fig 2D-E, S2B**). Thus, HFD + *L. paracasei* mice clustering together with LFD-fed reference mice had a blinded steatosis score of 0-1, whereas those clustering together with HFD + vehicle control mice obtained a steatosis score ≥4, hence corroborating that NAFLD severity and not dietary feeding dictated hepatic lipid profile (**Fig 2D-E, S2B**). Both *L. reuteri* and *L. paracasei*, on the other hand, partly prevented the massively increased liver tissue mass and TG concentration in FFMD-fed mice (**Fig 2B-C**). When assessing the histological steatosis score, we observed that *L. paracasei* reduced steatosis development exclusively in HFD-fed mice, while only *L. reuteri* was able to significantly reduce this phenotype in FFMD-fed mice (**Fig 2D**). The selective NAFLD-reducing effect of *L. paracasei* in HFD-fed animals was reflected by a strong trend of diminished hepatocellular hypertrophy (p = 0.06) and macrovesicular steatosis (**Fig 2F-G, K**) hence mirroring the observed hepatic TG concentrations (**Fig 2C**). The selective amelioration of NAFLD by *L. reuteri* in FFMD-fed mice was mainly associated with reduced hepatocellular hypertrophy and a partial improvement of microvesicular steatosis (**Fig 2F, H**). *L. paracasei* reduced inflammatory foci counts in HFD-fed mice supported by a similar tendency in hepatic cytokine levels (**Fig 1I, S2C-D**, PERMANOVA p = 0.089), pinpointing the beneficial NAFLD ameliorating capabilities of *L. paracasei* in HFD-, but not FFMD-fed mice.

L. reuteri diet-dependently altered intestinal bile acid levels

HFD and FFMD feeding distinctively impacted the plasma lipid profile (**Fig 3A**), where a majority of the altered lipids were regulated similarly to that observed in the liver (**Fig S2A**). Circulating levels of total cholesterol (**Fig 3B**) and TG (**Fig S3A**, p = 0.01, q = 0.06) were increased by FFMD compared to HFD while total free fatty acids (FA) were not significantly altered by the background diet (**Fig S3B**). A subset of the HFD-fed mice receiving *L. paracasei* clustered with the LFD reference group (**Fig 3A**), as seen

with hepatic lipids (**Fig S2B**) and numerically tended to reduce HFD-induced total cholesterol levels (**Fig 3B**, p = 0.19, q = 0.32). *L. reuteri* significantly increased the relative abundance of FA 18:2 exclusively in FFMD-fed mice (**Fig 3C**).

Due to the known influence of bile acids (BAs) on the gut-liver axis by the enterohepatic pathway we quantified specific BAs in liver tissue. *L. paracasei* tended to increase total liver BA levels in HFD-fed mice (p = 0.02, q = 0.11) and the BA precursor THCA (**Fig S3C-D**) associated with borderline higher levels of an unconjugated and and taurine-conjugated BAs (**Fig S3E-F**, p < 0.05, q < 0.1). Over 90% of BAs excreted into the intestine are recirculated in the small intestine (Inagaki et al., 2005) prompting us to quantify the remaining BAs in the cecum, which did not differ in weight between the groups (**Fig S3G**). FFMD increased the levels of unconjugated BAs in this intestinal segment compared to HFD as well as secondary BAs without affecting the total cecal BA concentration (**Fig 3D-E, S3H**). These increases were associated with higher expression of fibroblast growth factor (Fgf)15 in ileum tissue (**Fig 3G**), which is upregulated by BAs through intestinal Fxr (Inagaki et al., 2005). The indication of enhanced activation of the Fxr-Fgf15 enterohepatic pathway did, however, not result in repression of Cyp7a1 expression (**Fig S3I**), the rate-limiting enzyme of BA synthesis in the liver (Inagaki et al., 2005) thus suggesting a dysregulation of the pathway by FFMD feeding. Intestinal Fxr-activation by BAs has additionally been found to mediate Paneth cell defects (Liu et al., 2021), supported by numerically reduced expression of the host-defense protein RegIIIγ in FFMD-fed mice in our study (**Fig 3H**, p = 0.16, q = 0.82). The *Lactobacilli* did not significantly alter *Fgf15* or *RegIIIγ* expression on the respective diets (**Fig 3G-H**).

Despite the tendency of augmented hepatic BA levels in *L. paracasei* treated HFD-fed mice (**Fig S3C**), we did not observe changes in cecal BA levels in this group (**Fig 3E-G**). Conversely, *L. reuteri* increased cecal (**Fig 3D**) but not hepatic (**Fig S3C**) BA levels exclusively in FFMD-fed mice. Notably, only conjugated BAs were affected by *L. reuteri* (**Fig 3E-F**), suggesting a reduced re-uptake of the conjugated BA as proposed for other *L. reuteri* strains (Lu et al., 2022; Zhang et al., 2022).

Predicted functions of the gut microbiota correlated with disease severity, not macronutrient composition, and were modulated by probiotics

We collected fecal samples at weeks 0, 4, 8, and 12 to assess longitudinal changes in gut microbiota composition as a function of *Lactobacilli* and diet. At termination, we further collected the content of small intestine and colon. Microbiota composition was markedly affected by dietary macronutrient composition (FFMD and LFD clustering separately from HFD), in colonic and fecal samples**,** suggesting that dietary nutrients, such as proteins, carbohydrate or lipid sources, rather than lipid proportion or the

occurrence of obesity determined gut microbiota composition (**Table 1, Fig 4A-C, S4A-B**). This was despite equal baseline microbiota composition and with a transient increase of fecal alpha diversity measures from FFMD and LFD (**Fig S4C-E**).

Lactobacillus administration did not significantly alter the microbiota composition in small intestine or colon content sampled 2-5h after the latest oral gavage (**Fig 4A-B, D**). The fecal microbiota, sampled 22-24h after the latest gavage, was however transiently modified by *L. paracasei* in HFD-fed mice (**Fig S4A, D**). *L. reuteri* temporarily changed FFMD-induced fecal microbiota composition (**Fig S4B, D**), which tended to be modified by the end of the study (**Fig 4A-D)** and was significantly modified in a replicated study (**Fig S4F-H**). Investigation of differential abundances at all phylogenetic levels robustly modified by diet in the different studies revealed an increase in abundance of the phylum *Proteobacteria* by FFMD in colonic and fecal samples compared to LFD (p < 0.001, **Fig 4E, S4I-J**). The FFMD-induced *Proteobacteria* abundance in the distal intestinal segments was numerically reduced by *L. reuteri* and to a lesser degree *L. paracasei* in reproduced studies (**Fig 4E, S4I-J**).

To investigate whether colonization of the administered *Lactobacilli* in the intestinal tract was affected by diet we assessed the ASVs originating from the administered *L. reuteri* (ASV15) and *L. paracasei* (ASV5 and ASV31, **Fig 4F**). As expected, these ASVs were found in the small intestine of mice 2-5 hours after receiving the corresponding *Lactobacillus* strain (**Fig 4G**). No change from background signal was detected in colon content 2-5 hours, or feces 22-24 hours after oral gavage (**Fig 4G**), indicating negligible colonic colonization by the applied probiotics.

We next assessed the predicted functions of the gut microbiota by assessing KEGG orthology. Interestingly, while colon and fecal microbiota composition reflected ingested nutrients (**Fig 4A-C**), the predicted microbiota functionality more closely resembled the pathology of experimental mice (**Fig 4H-J**). As such, LFD-fed mice clustered in close vicinity to HFD-fed mice, despite being fed a diet matched to the FFMD (**Table 1, Fig 4H-J**). This observation was further supported by *L. paracasei* treated HFD-fed mice who exhibited an intermediary disease phenotype and likewise clustered between the LFD-fed mice and their vehicle-treated HFD-fed counterparts (**Fig 4H-J**). Despite the lack of clear and reproducible effects on gut microbiota composition from the gavage by the *Lactobacilli* strains there was a significant change in FFMD-induced predicted functions of the microbiota by *L. reuteri* in all assessed intestinal segments at termination (**Fig 4H-K**), again mirroring their partial protection against FFMD-induced liver pathology.

Discussion

We used mice fed two distinct but energy-matched high fat and high sucrose diets to test the preventive effects of two *Lactobacillus* strains on obesity and related metabolic disturbances. While both HFD and FFMD induced obesity and compromised glucoregulatory capacity, FFMD led to enhanced ectopic fat distribution to visceral fat and liver. This induced major phenotypic differences between the HFD and FFMD, notably on body weight gain, energy harvesting, glucose homeostasis, enterohepatic circulation of BAs as well as gut microbiota composition and predicted functionality. Accordingly, the greatest diet-specific impact of the customized FFMD was observed on NAFLD features including hepatocellular morphology, liver lipid and inflammatory profiles. The FFMD-induced liver pathology culminated in significant fibrotic remodeling in >75% of the mice, hence contrasting the results in HFD-fed mice where no fibrotic bridging was observed and only 1 of 9 mice exhibited pronounced portal fibrosis. These diet-specific effects can all potentially affect the ability of probiotics to alleviate obesity-related dysmetabolism and related diseases such as NAFLD. This has major physiological significance and clinical relevance given that the choice of experimental diets in preclinical investigation, and the dietary habits of subjects enrolled in clinical trials, are thus likely to influence the efficacy of probiotics. This could lead to the dismissal of promising probiotic candidates for mitigating obesity and related metabolic disorders due to false negative results in preclinical trials, a lack of clinical translatability when advancing to clinical phases where dietary macronutrient composition is not specifically controlled for, or - worst case - lead to the endorsement of probiotics with limited or potentially even unfavorable effects in combination with certain diets not adequately recapitulated in preceeding studies (Spencer et al., 2021; The Lancet Gastroenterology & Hepatology, 2019).

In our study, *L. paracasei* reduced obesity development associated with diminished NAFLD exclusively in the context of HFD-feeding without changing energy balance. As a result, 45% of the HFD-mice receiving *L. paracasei* had no histopathological signs of NAFLD, which was not prevented in any other HFD-fed mouse in the study. *L. reuteri* on the other hand improved glucose homeostasis and the more severe NAFLD phenotype only in FFMD-fed mice. This was associated with increased plasma FA 18:2 levels, reported to inversely correlate with human hepatocellular carcinoma (HCC)(Qiu et al., 2015) and a murine NASH-HCC induction (Vlock et al., 2020). *L. reuteri* additionally modified the FFMD-induced BA profile by augmenting conjugated BAs resulting in increased total BA levels in cecum. This may suggest increased energy excretion although this remained under the detection range of the methods in the current studies. Alternatively, the *L. reuteri* strain may dampen a FFMD-induced immune response by modifications of secondary BAs, as recently demonstrated by others (Campbell et al., 2020; Paik et al., 2022; Zhang et al., 2022), although this was not possible to assess further in the current study setup. It

should be noted that the metabolic benefits of *L. paracasei* were associated with reduced weight gain and fat percentage in the HFD model. However, the improved insulin sensitivity in *L. reuteri*-treated FFMD animals could not be linked to diminished weight gain suggesting different modes of action. Importantly, these diet-dependent effects of the probiotic candidates were highly reproducible between repetitive studies.

We also observed that the diet composition, rather than the amount of dietary fat and high glycemic carbohydrates, was the main driver of gut microbial community structures. This was examplified by a reproducible FFMD-induced augmentation of *Proteobacteria* phylum abundances, which has previously been linked to epithelial damage via disruption in anaerobiosis (Litvak et al., 2017) and NASH (Zhu et al., 2013). Interestingly, we and others previously observed *Proteobacteria* as the dominant bacterial phylum in extra-intestinal tissues of morbidly obese individuals with T2D compared to normoglycemic weight-matched counterparts (Anhê et al., 2020; Massier et al., 2020). Predicted functions of the microbiota, however, followed disease phenotype rather than diet composition. As such, the tested probiotic strains had negligible impact on microbiota composition, but not its predicted functions, in the sampled intestinal segments and showed minimal signs of colonization of the gastro-intestinal tract, supporting previous reports (Kristensen et al., 2016). In the study from Zmora *et al*, individual responses to probiotics were suggested to originate from genetic or baseline microbiota differences naturally found in humans (Zmora et al., 2018). Both factors were accounted for in our studies, where we show dietary differences alone were sufficient in determining probiotic efficacy.

It remains uncertain what caused the diet-dependent effects on probiotic efficacy observed in the present study. Possible explanations could relate to direct dietary influence where the probiotics are dependent on the utilization of certain dietary sources. Alternatively, diet-induced differences in gastrointestinal ecology, such as bile acid concentrations, could be influencing the probiotic effects; a trait that is warranted to investigate in future studies, just as a combination therapy may alleviate diets dependency. To this end, a combination of the *L. reuteri* and *L. paracasei* strains is currently investigated in an ongoing randomized, double-blinded, placebo-controlled clinical trial in prediabetic subjects (ClinicalTrials.gov identifier NCT04767789).

Differences in probiotic efficacy have been reported for *L. helveticus* R0052, *L. plantarum* WCFS1, and *P. copri* (Kovatcheva-Datchary et al., 2015; Ohland et al., 2013; Pedersen et al., 2016; Tachon et al., 2014) when comparing lean, chow-fed mice to obese, Western diet-fed mice. Our study demonstrates that strain-specific efficacies of *Lactobacilli* on body composition, insulin resistance, and NAFLD were

reproducible, yet highly influenced by the type of obesogenic diet, and thus substantially expanding the understanding of dietary influence on probiotic efficacy. While future studies are warranted to elucidate the generic implication of our findings, the dietary interactions described here may likely extend beyond the tested *Lactobacillus* strains. Potentially, this could aid in explaining empiric observations of responders versus non-responders to probiotics in general and might indicate a need for pre-treatment stratification of patients.

References

Anhê, F.F., Jensen, B.A.H., Varin, T. V, Servant, F., Van Blerk, S., Richard, D., Marceau, S., Surette, M., Biertho, L., Lelouvier, B., et al. (2020). Type 2 diabetes influences bacterial tissue compartmentalisation in human obesity. Nat. Metab. https://doi.org/10.1038/s42255-020-0178-9.

Arterburn, D.E., Telem, D.A., Kushner, R.F., and Courcoulas, A.P. (2020). Benefits and Risks of Bariatric Surgery in Adults. JAMA *324*, 879. https://doi.org/10.1001/jama.2020.12567.

Bates, D., Mächler, M., Bolker, B., and Walker, S. (2014). Fitting Linear Mixed-Effects Models using lme4. *67*. https://doi.org/10.18637/jss.v067.i01.

Campbell, C., McKenney, P.T., Konstantinovsky, D., Isaeva, O.I., Schizas, M., Verter, J., Mai, C., Jin, W., Guo, C., Violante, S., et al. (2020). Bacterial metabolism of bile acids promotes generation of peripheral regulatory T cells. Nature https://doi.org/10.1038/s41586-020-2193-0.

Chatterjee, S., Khunti, K., and Davies, M.J. (2017). Type 2 diabetes. Lancet *389*, 2239–2251. https://doi.org/10.1016/S0140-6736(17)30058-2.

Chong, J., Wishart, D.S., and Xia, J. (2019). Using MetaboAnalyst 4.0 for Comprehensive and Integrative Metabolomics Data Analysis. Curr. Protoc. Bioinforma. *68*, 1–128. https://doi.org/10.1002/cpbi.86.

Delzenne, N.M., Neyrinck, A.M., Bäckhed, F., and Cani, P.D. (2011). Targeting gut microbiota in obesity: effects of prebiotics and probiotics. Nat. Rev. Endocrinol. *7*, 639–646. https://doi.org/10.1038/nrendo.2011.126.

Dulloo, A.G., and Montani, J.-P. (2015). Pathways from dieting to weight regain, to obesity and to the metabolic syndrome: an overview. Obes. Rev. *16*, 1–6. https://doi.org/10.1111/obr.12250.

Eslam, M., Sanyal, A.J., George, J., Sanyal, A., Neuschwander-Tetri, B., Tiribelli, C., Kleiner, D.E., Brunt, E., Bugianesi, E., Yki-Järvinen, H., et al. (2020). MAFLD: A Consensus-Driven Proposed Nomenclature for Metabolic Associated Fatty Liver Disease. Gastroenterology *158*, 1999-2014.e1. https://doi.org/10.1053/j.gastro.2019.11.312.

Fabbrini, E., Sullivan, S., and Klein, S. (2010). Obesity and nonalcoholic fatty liver disease: Biochemical, metabolic, and clinical implications. Hepatology *51*, 679–689. https://doi.org/10.1002/hep.23280.

FAO/WHO (2001). Probiotics in food - Health and nutritional properties and guidelines for evaluation. Report of a Joint FAO/WHO expert consultation on evaluation of health and nutritional properties of probiotics in food including powder milk with live lactic acid bacter.

Fonseca, V.A. (2009). Defining and Characterizing the Progression of Type 2 Diabetes. Diabetes Care *32*, S151–S156. https://doi.org/10.2337/dc09-S301.

Goodman, Z.D. (2007). Grading and staging systems for inflammation and fibrosis in chronic liver diseases. J. Hepatol. *47*, 598–607. https://doi.org/10.1016/j.jhep.2007.07.006.

Hall, K.D., and Guo, J. (2017). Obesity Energetics: Body Weight Regulation and the Effects of Diet

Composition. Gastroenterology *152*, 1718-1727.e3. https://doi.org/10.1053/j.gastro.2017.01.052.

Inagaki, T., Choi, M., Moschetta, A., Peng, L., Cummins, C.L., McDonald, J.G., Luo, G., Jones, S.A., Goodwin, B., Richardson, J.A., et al. (2005). Fibroblast growth factor 15 functions as an enterohepatic signal to regulate bile acid homeostasis. Cell Metab. *2*, 217–225. https://doi.org/10.1016/j.cmet.2005.09.001.

Ishak, K., Baptista, A., Bianchi, L., Callea, F., De Groote, J., Gudat, F., Denk, H., Desmet, V., Korb, G., MacSween, R.N.M., et al. (1995). Histological grading and staging of chronic hepatitis. J. Hepatol. *22*, 696–699. https://doi.org/10.1016/0168-8278(95)80226-6.

Khalil, H., Ellwood, L., Lord, H., and Fernandez, R. (2020). Pharmacological Treatment for Obesity in Adults: An Umbrella Review. Ann. Pharmacother. *54*, 691–705. https://doi.org/10.1177/1060028019898912.

Kivimäki, M., Kuosma, E., Ferrie, J.E., Luukkonen, R., Nyberg, S.T., Alfredsson, L., Batty, G.D., Brunner, E.J., Fransson, E., Goldberg, M., et al. (2017). Overweight, obesity, and risk of cardiometabolic multimorbidity: pooled analysis of individual-level data for 120 813 adults from 16 cohort studies from the USA and Europe. Lancet Public Heal. *2*, e277–e285. https://doi.org/10.1016/S2468-2667(17)30074-9.

Klindworth, A., Pruesse, E., Schweer, T., Peplies, J., Quast, C., Horn, M., and Glöckner, F.O. (2013). Evaluation of general 16S ribosomal RNA gene PCR primers for classical and next-generation sequencing-based diversity studies. Nucleic Acids Res. *41*, 1–11. https://doi.org/10.1093/nar/gks808.

Kovatcheva-Datchary, P., Nilsson, A., Akrami, R., Lee, Y.S., De Vadder, F., Arora, T., Hallen, A., Martens, E., Björck, I., and Bäckhed, F. (2015). Dietary Fiber-Induced Improvement in Glucose Metabolism Is Associated with Increased Abundance of Prevotella. Cell Metab. *22*, 971–982. https://doi.org/10.1016/j.cmet.2015.10.001.

Krautkramer, K.A., Fan, J., and Bäckhed, F. (2020). Gut microbial metabolites as multi-kingdom intermediates. Nat. Rev. Microbiol. https://doi.org/10.1038/s41579-020-0438-4.

Kristensen, N.B., Bryrup, T., Allin, K.H., Nielsen, T., Hansen, T.H., and Pedersen, O. (2016). Alterations in fecal microbiota composition by probiotic supplementation in healthy adults: a systematic review of randomized controlled trials. Genome Med. *8*, 52. https://doi.org/10.1186/s13073-016-0300-5.

Kuznetsova, A., Brockhoff, P.B., and Christensen, R.H.B. (2017). lmerTest Package: Tests in Linear Mixed Effects Models. J. Stat. Softw. *82*, 1–26. https://doi.org/10.18637/jss.v082.i13.

Larsen, I.S., Jensen, B.A.H., Bonazzi, E., Choi, B.S.Y., Kristensen, N.N., Schmidt, E.G.W., Süenderhauf, A., Morin, L., Olsen, P.B., Hansen, L.B.S., et al. (2021). Fungal lysozyme leverages the gut microbiota to curb DSS-induced colitis. Gut Microbes *13*. https://doi.org/10.1080/19490976.2021.1988836.

Liang, W., Menke, A.L., Driessen, A., Koek, G.H., Lindeman, J.H., Stoop, R., Havekes, L.M., Kleemann., R., and Van Den Hoek, A.M. (2014). Establishment of a general NAFLD scoring system for rodent

models and comparison to human liver pathology. PLoS One *9*, 1–17. https://doi.org/10.1371/journal.pone.0115922.

Litvak, Y., Byndloss, M.X., Tsolis, R.M., and Bäumler, A.J. (2017). Dysbiotic Proteobacteria expansion: a microbial signature of epithelial dysfunction. Curr. Opin. Microbiol. *39*, 1–6. https://doi.org/10.1016/j.mib.2017.07.003.

Liu, T.C., Kern, J.T., Jain, U., Sonnek, N.M., Xiong, S., Simpson, K.F., VanDussen, K.L., Winkler, E.S., Haritunians, T., Malique, A., et al. (2021). Western diet induces Paneth cell defects through microbiome alterations and farnesoid X receptor and type I interferon activation. Cell Host Microbe *29*, 988-1001.e6. https://doi.org/10.1016/j.chom.2021.04.004.

Lu, M., Sun, J., Zhao, Y., Zhang, H., Li, X., Zhou, J., Dang, H., Zhang, J., Huang, W., Qi, C., et al. (2022). Prevention of High-Fat Diet-Induced Hypercholesterolemia by Lactobacillus reuteri Fn041 Through Promoting Cholesterol and Bile Salt Excretion and Intestinal Mucosal Barrier Functions. Front. Nutr. *9*, 1–14. https://doi.org/10.3389/fnut.2022.851541.

Massier, L., Chakaroun, R., Tabei, S., Crane, A., Didt, K.D., Fallmann, J., Von Bergen, M., Haange, S.B., Heyne, H., Stumvoll, M., et al. (2020). Adipose tissue derived bacteria are associated with inflammation in obesity and type 2 diabetes. Gut 1–11. https://doi.org/10.1136/gutjnl-2019-320118.

Ohland, C.L., Kish, L., Bell, H., Thiesen, A., Hotte, N., Pankiv, E., and Madsen, K.L. (2013). Effects of Lactobacillus helveticus on murine behavior are dependent on diet and genotype and correlate with alterations in the gut microbiome. Psychoneuroendocrinology *38*, 1738–1747. https://doi.org/10.1016/j.psyneuen.2013.02.008.

Oksanen, J., Blanchet, F.G., Kindt, R., Legendre, P., Minchin, P.R., O'Hara, R.B., Simpson, G.L., Solymos, P., Stevens, M.H.H., and Wagner, H. (2017). vegan: Community Ecology Package. R Packag. Ver. 2.4–3 https://doi.org/10.4135/9781412971874.n145.

Paik, D., Yao, L., Zhang, Y., Bae, S., D'Agostino, G.D., Zhang, M., Kim, E., Franzosa, E.A., Avila-Pacheco, J., Bisanz, J.E., et al. (2022). Human gut bacteria produce TH17-modulating bile acid metabolites.

Pedersen, H.K., Gudmundsdottir, V., Nielsen, H.B., Hyotylainen, T., Nielsen, T., Jensen, B.A.H., Forslund, K., Hildebrand, F., Prifti, E., Falony, G., et al. (2016). Human gut microbes impact host serum metabolome and insulin sensitivity. Nature *535*, 376–381. https://doi.org/10.1038/nature18646.

Qiu, J.-F., Zhang, K.-L., Zhang, X.-J., Hu, Y.-J., Li, P., Shang, C.-Z., and Wan, J.-B. (2015). Abnormalities in Plasma Phospholipid Fatty Acid Profiles of Patients with Hepatocellular Carcinoma. Lipids *50*, 977–985. https://doi.org/10.1007/s11745-015-4060-6.

Reid, G., Gaudier, E., Guarner, F., Huffnagle, G.B., Macklaim, J.M., Munoz, A.M., Martini, M., Ringel-Kulka, T., Sartor, B.R., Unal, R.R., et al. (2010). Responders and non-responders to probiotic

interventions. Gut Microbes *1*, 200–204. https://doi.org/10.4161/gmic.1.3.12013.

Spencer, C.N., McQuade, J.L., Gopalakrishnan, V., McCulloch, J.A., Vetizou, M., Cogdill, A.P., Khan, M.A.W., Zhang, X., White, M.G., Peterson, C.B., et al. (2021). Dietary fiber and probiotics influence the gut microbiome and melanoma immunotherapy response. Science *374*, 1632–1640. https://doi.org/10.1126/science.aaz7015.

Stunkard, A.J., Harris, J.R., Pedersen, N.L., and McClearn, G.E. (1990). The Body-Mass Index of Twins Who Have Been Reared Apart. N. Engl. J. Med. *322*, 1483–1487. https://doi.org/10.1056/NEJM199005243222102.

Tachon, S., Lee, B., and Marco, M.L. (2014). Diet alters probiotic Lactobacillus persistence and function in the intestine. Environ. Microbiol. *16*, 2915–2926. https://doi.org/10.1111/1462-2920.12297.

Tandra, S., Yeh, M.M., Brunt, E.M., Vuppalanchi, R., Cummings, O.W., Ünalp-Arida, A., Wilson, L.A., and Chalasani, N. (2011). Presence and significance of microvesicular steatosis in nonalcoholic fatty liver disease. J. Hepatol. *55*, 654–659. https://doi.org/10.1016/j.jhep.2010.11.021.

The Lancet Gastroenterology & Hepatology (2019). Probiotics: elixir or empty promise? Lancet Gastroenterol. Hepatol. *4*, 81. https://doi.org/10.1016/S2468-1253(18)30415-1.

Thonusin, C., Shinlapawittayatorn, K., Chattipakorn, S.C., and Chattipakorn, N. (2020). The impact of genetic polymorphisms on weight regain after successful weight loss. Br. J. Nutr. *124*, 809–823. https://doi.org/10.1017/S0007114520001968.

Tyrrell, J., Jones, S.E., Beaumont, R., Astley, C.M., Lovell, R., Yaghootkar, H., Tuke, M., Ruth, K.S., Freathy, R.M., Hirschhorn, J.N., et al. (2016). Height, body mass index, and socioeconomic status: mendelian randomisation study in UK Biobank. BMJ *352*, i582. https://doi.org/10.1136/bmj.i582.

Varkevisser, R.D.M., van Stralen, M.M., Kroeze, W., Ket, J.C.F., and Steenhuis, I.H.M. (2019). Determinants of weight loss maintenance: a systematic review. Obes. Rev. *20*, 171–211. https://doi.org/10.1111/obr.12772.

Venditti, E.M., Bray, G.A., Carrion-Petersen, M.L., Delahanty, L.M., Edelstein, S.L., Hamman, R.F., Hoskin, M.A., Knowler, W.C., and Ma, Y. (2008). First versus repeat treatment with a lifestyle intervention program: Attendance and weight loss outcomes. Int. J. Obes. *32*, 1537–1544. https://doi.org/10.1038/ijo.2008.134.

Vlock, E.M., Karanjit, S., Talmon, G., and Farazi, P.A. (2020). Reduction of polyunsaturated fatty acids with tumor progression in a lean non-alcoholic steatohepatitis- associated hepatocellular carcinoma mouse model. J. Cancer *11*, 5536–5546. https://doi.org/10.7150/jca.48495.

Wiciński, M., Gębalski, J., Gołębiewski, J., and Malinowski, B. (2020). Probiotics for the Treatment of Overweight and Obesity in Humans—A Review of Clinical Trials. Microorganisms *8*, 1148. https://doi.org/10.3390/microorganisms8081148.

World Health Organization (WHO). (2016). Obesity and overweight fact sheet.

Younossi, Z.M., Koenig, A.B., Abdelatif, D., Fazel, Y., Henry, L., and Wymer, M. (2016). Global epidemiology of nonalcoholic fatty liver disease—Meta-analytic assessment of prevalence, incidence, and outcomes. Hepatology *64*, 73–84. https://doi.org/10.1002/hep.28431.

Zhang, C., Fang, R., Lu, X., Zhang, Y., Yang, M., Su, Y., Jiang, Y., and Man, C. (2022). Lactobacillus reuteri J1 prevents obesity by altering the gut microbiota and regulating bile acid metabolism in obese mice. Food Funct. *13*, 6688–6701. https://doi.org/10.1039/D1FO04387K.

Zhu, L., Baker, S.S., Gill, C., Liu, W., Alkhouri, R., Baker, R.D., and Gill, S.R. (2013). Characterization of gut microbiomes in nonalcoholic steatohepatitis (NASH) patients: A connection between endogenous alcohol and NASH. Hepatology *57*, 601–609. https://doi.org/10.1002/hep.26093.

Zmora, N., Zilberman-Schapira, G., Suez, J., Mor, U., Dori-Bachash, M., Bashiardes, S., Kotler, E., Zur, M., Regev-Lehavi, D., Brik, R.B.-Z., et al. (2018). Personalized Gut Mucosal Colonization Resistance to Empiric Probiotics Is Associated with Unique Host and Microbiome Features. Cell *174*, 1388-1405.e21. https://doi.org/10.1016/j.cell.2018.08.041.

Table 1: Composition of diets

Ingredient	Protein content	Fat content	U	Low Fat Diet Control mixed Protein S9552-E026	Fast Food Mimicking Diet mixed Protein S9552-E024	High fat diet 45 kcal% Fat S9552-E060
Casein	87	<0.6	%	3,000	3,000	25,000
Poultry protein	71	10	%	8,500	8,500	——
Soyprotein isolate	88	0,4	%	4,900	4,900	——
Wheat gluten, Vital	76,5	1,7	%	2,300	2,300	——
Egg yolk	38	56	%	1,600	1,600	——
Egg white	79	0,1	%	1,600	1,600	——
Whey powder, 71 % Lactose	11,9	1	%	2,900	2,900	——
L-Cystine	72,2	——	%	0,300	0,300	0,300
Corn starch	——	——	%	45,260	5,200	7,000
Maltodextrin	——	——	%	8,000	8,000	11,000
Sucrose 1)	——	——	%	7,000	24,400	20,050
Fructose 1)	——	——	%	——	6,500	——
Cellulose powder	——	——	%	5,000	5,000	5,700
Vitamin premixture 2)	——	——	%	1,000	1,000	1,000
Mineral premixture + additions 2)	——	——	%	4,300	4,300	6,000
Choline Cl (50 %)	——	——	%	0,290	0,290	0,200
L-Carnitine	——	——	%	0,050	0,050	——
Cholesterol	——	——	%	——	0,150	0,150
Butylated hydroxytoluene	——	——	%	0,010	0,010	——
Butter fat, dehydrated	——	——	%	2,100	18,800	——
Corn oil	——	——	%	1,620	1,100	——
Linseed oil	——	——	%	0,270	0,100	——
Pork lard	——	——	%	——	——	20,800
Soybean oil	——	——	%	——	——	2,800
Proximate contents						
Crude protein			%	17,2	17,2	22,100
Crude fat			%	5,8	21,8	23,600
Crude fibre			%	5,0	5,1	5,700
Crude ash			%	4,8	4,8	5,400
Starch			%	43,5	5,0	6,700
Dextrin			%	7,9	7,9	10,900
Sugar (total)			%	10,2	34,0	21,100
Energy (Atwater)			MJ/kg	15,7	19,2	19,300
kcal% Protein				18	15	20,000

kcal% Fat				14	43	45,000
kcal% Carbohydrates				68	42	35,000
Lysine			%	0,95	0,95	1,83
Methionine			%	0,43	0,43	0,78
Methionine & Cystine			%	0,94	0,94	1,18
Threonine			%	0,62	0,62	0,97
Tryptophan			%	0,19	0,19	0,29
Arginine			%	0,91	0,91	0,86
Histidine			%	0,37	0,37	0,67
Valine			%	0,86	0,86	1,54
Isoleucine			%	0,71	0,71	1,25
Leucine			%	1,23	1,23	2,19
Phenylalanine			%	0,74	0,74	1,14
Phe + Tyr			%	1,29	1,29	2,31
Glycine			%	0,96	0,96	0,47
Glutamic acid			%	3,00	3,00	4,97
Aspartic acid			%	1,36	1,36	1,64
Proline			%	1,20	1,20	2,53
Serine			%	0,82	0,82	1,32
Alanine			%	0,84	0,84	0,66
Fatty acids, % in the diet						
C 4:0				0,08	0,72	——
C 6:0				0,05	0,47	——
C 8:0				0,03	0,26	——
C 10:0				0,06	0,56	0,03
C 12:0				0,07	0,64	0,05
C 14:0				0,24	2,00	0,29
C 16:0				1,15	5,50	5,34
C 18:0				0,36	1,97	2,92
C 20:0				0,02	0,04	0,07
C 16:1				0,13	0,43	0,62
C 18:1				1,68	5,16	9,42
C 18:2				1,21	1,21	3,45
C 18:3				0,17	0,17	0,37
Cholesterol			mg/kg	~440	~2340	~1,580
Calcium			%	0,82	0,82	0,92
Phosphorus			%	0,58	0,57	0,64
Sodium			%	0,39	0,38	0,20
Magnesium			%	0,1	0,1	0,23
Potassium			%	0,8	0,8	0,97

Iron			mg/kg	72	72	168
Manganese			mg/kg	22	22	95
Zinc			mg/kg	42	42	65
Copper			mg/kg	13	13	13
Iodine			mg/kg	0,31	0,31	1,16
Selenium			mg/kg	0,20	0,20	0,18
Vitamin A			IU/kg	15.900	20.500	15.000
Vitamin D3			IU/kg	1.550	1.550	1.500
Vitamin E			mg/kg	156	159	150
Vitamin C			mg/kg	30	30	30
Vitamin K			mg/kg	20	20	20
Vitamin B1			mg/kg	26	26	25
Vitamin B2			mg/kg	18	18	16
Vitamin B6			mg/kg	17	17	16
Vitamin B12			µg/kg	32	32	30
Niacin			mg/kg	53	53	47
Pantothenic acid			mg/kg	59	59	55
Folic acid			mg/kg	16	16	16
Biotin			µg/kg	355	355	300
Choline			mg/kg	1.380	1.380	920

Supplementary Table 1: Grading system for murine NAFLD activity score, adapted from Liang *et al* (Liang et al., 2014).

Histological feature	Score			
	0	1	2	3
Steatosis score				
Macrovesicular steatosis	< 5%	5-33%	33-66%	>66%
Microvesicular steatosis	< 5%	5-33%	33-66%	>66%
Hypertrophy	< 5%	5-33%	33-66%	>66%
Inflammation Score				
Number of inflammatory foci per field	<0.5	0.5-1.0	1.0-2.0	>2.0

Supplementary Table 2: Extended staging system for hepatic fibrosis as introduced by Ishak et al. (Ishak et al., 1995)

Histological changes	Score
No fibrosis	0
Fibrous expansion of some portal areas with or without short fibrous septa	1
Fibrous expansion of most portal areas with or without short fibrous septa	2
Fibrous expansion of most portal areas with occasional portal to portal (P-P) bridging	3
Fibrous expansion of portal areas with marked bridging (P-P as well as portal to central (P-C))	4
Marked bridging (P-P and/or P-C) with occasional nodules (incomplete cirrhosis)	5
Cirrhosis, probable or definite	6

Supplementary Table 3: RT-qPCR primer sequences

Target gene	Forward primer sequence	Reverse primer sequence
18S	GTAACCCGTTGAACCCCATT	CCATCCAATCGGTAGTAGCG
Fgf15	GAGGACCAAAACGAACGAAATT	ACGTCCTTGATGGCAATCG
Cyp7a1	AGCAACTAAACAACCTGCCAGTACTA	GTCCGGATATTCAAGGATGCA
RegIIIγ	CCGTGCCTATGGCTCCTATTG	GCACAGACACAAGATGTCCTG

Figure 1: Body composition and insulin sensitivity are diet-dependently affected by *Lactobacilli*

A) Experimental outline for all three studies with 12 weeks of feeding either High Fat Diet (HFD), Fast Food-mimicking Diet (FFMD) or Low Fat reference Diet (LFD) (matched to the FFMD in nutritional sources) with indicated total energy content in MJ/kg and energy distribution of macronutrients in percentage. Shades indicate mixed nutritional sources of the macronutrients. During the study, the mice were orally gavaged daily with either 10^8 CFU of either *Limosilactobacillus reuteri (L. reuteri)*, *Lacticaseibacillus paracasei* subsp. *paracasei* (*L. paracasei)*, or an equal volume of vehicle (PBS). **B)** Body weight in grams at the end of the study period. **C)** Fat mass in grams assessed by MR scan at the end of the study period. **D)** Energy uptake per cage monitored over 24h as the inverse percentage of energy excretion compared to energy intake assessed in week 11 of the study. **E)** Accumulated energy intake per cage during the study period as average intake per mouse. **F)** Plasma 5h fasting insulin assessed in week 10 of the study. **G)** Blood glucose levels of 5h fasted mice in week 10 of the study. **H)** Blood glucose values from Study 1 during oral glucose tolerance test (oGTT) with 1.5 µg/g lean mass dextrose in week 10. **I)** Corresponding plasma insulin levels during oGTT. **J)** Corresponding plasma C-peptide levels during oGTT. **B-G)** Bars indicate group mean ± SEM. Points represent individual mice; individual studies are identified by point shapes where circles indicate Study 1, triangles Study 2, and squares Study 3. Dashed line indicates LFD group mean with interquartile range shown in grey. Asterisks indicate fdr-corrected q-values <0.05 using linear mixed effects models comparing indicated groups. **H-J)** Lines indicate group mean ± SEM. Asterisks indicate p-values <0.05 using two-way ANOVA with multiple comparisons between vehicle-treated groups or *Lactobacillus* treated group to vehicle-treated group fed the same diet with Bonferroni post-hoc test.

Figure 2: Non-alcoholic fatty liver disease (NAFLD) mitigation by *Lactobacilli* depends on diet

A) Images of total livers from necropsy from representative mice in vehicle-gavaged dietary groups of Study 1 weighting respectively 1.01 g (LFD), 1.58 g (HFD) and 3.37 g (FFMD). **B)** Liver weight in grams at the end of the study. **C)** Liver triacylglyceride concentration as µg/mg tissue. **D)** Histological steatosis score from 0-9 blindly assessed in H&E stained liver slides. **E)** Histopathological NAFLD categories as relative distribution per group of: No NAFLD, NAFLD, and NASH based on steatosis and inflammation scores of H&E stained liver slides. **F)** Hepatocellular hypertrophy score from 0-3 blindly assessed in H&E-stained liver slides. **G)** Macrovesicular steatosis score from 0-3 blindly assessed in H&E-stained liver slides. **H)** Microvesicular steatosis score from 0-3 blindly assessed in H&E-stained liver slides. **I)** Average number of inflammatory foci in 5 high power fields (HPF). **J)** Histological fibrosis assessed by Ishak scoring of Sirius Red-stained liver slides from Study 1. **K)** H&E and Sirius Red

stained liver slides from Study 1 selected as representative by having the median value of the group in the histological steatosis score. **B-D, F-I)** Point shape represent individual mice in three individual studies where circles indicate Study 1, triangles Study 2, squares Study 3. **B-D,F-J)** Dashed line indicates mean of LFD group with interquartile range shown in grey. **B-C)** Bars indicate group mean ± SEM. Asterisks indicate fdr-corrected q-values <0.05 using linear mixed effects models comparing indicated groups. **D, F-J)** Bars indicate group median with interquartile range. Asterisks indicate p-values <0.05 by Kruskal-Wallis test with multiple comparisons between vehicle-treated groups or *Lactobacillus* group to vehicle-treated group fed the same diet with Dunn's post-hoc test.

Figure 3: *L. paracasei* and *L.reuteri* diet-dependently affect liver lipids and cecum bile acids (BAs)
A) Heatmap of the 40 most discriminating plasma lipids from end of study 1 assesed by lipidomics. Experimental groups are indicated by color and samples are distributed by hierical clustering. **B)** Total cholesterol levels in plasma. Point shape represent individual mice in three individual studies where circles indicate Study 1, triangles Study 2, squares Study 3. **C)** Relative plasma levels of free fatty acid (FA) 18:2 measured by lipidomics. **D)** Total measured BA concentration in cecum content. **E)** Concentration of unconjugated BA in cecum content as fold change to the LFD group. **F)** Concentration of conjugated BA in cecum content as fold change to the LFD group. **G)** Relative Fibroblast growth factor 15 (Fgf15) gene expression in ileum tissue by RT-qPCR. **H)** Relative Regenerating islet-derived protein 3-gamma (RegIIIγ) gene expression in ileum tissue by RT-qPCR. **B-H)** Bars indicate group mean ± SEM. Points represent individual data points. Asterisks indicate fdr-corrected q-values <0.05 using linear mixed effects models comparing indicated groups. **B-D, G-H)** Dashed line indicate mean of LFD group with interquartile range shown in grey.

Figure 4: Gut microbiota composition and predicted function are affected by *L. reuteri* in combination with FFMD.
A) Principal coordinate analysis (PCoA) of microbiota composition using weighted UniFrac distances of small intestine content sampled at the end of Study 1. Centroids indicate group average illustrated with 50% CI and smaller points indicate individual data points. **B)** As a for colon content. **C)** As a for fecal samples collected at week 12 of the study. **D)** PERMANOVA p-values comparing the microbiota composition between indicated groups on weighted UniFrac distances. Asterisks indicate p-values < 0.05. **E)** Relative abundance in % of *Proteobacteria* phylum in small intestine-, colon content, and fecal samples at the end of Study 1. Points indicate individual data points. Asterisks indicate p-values < 0.05 comparing the indicated groups. **F)** Distribution of amplicon sequence variants (ASVs) as relative abundances identified in *Limosilactobacillus reuteri* DSM 32910 (*L. reuteri*) or *Lacticaseibacillus paracasei* subsp. *paracasei*

DSM 32851 (*L. paracasei*) stocks. **G)** Relative abundance in % of ASV15, ASV5 and ASV31, respectively, from the administered *Lactobacillus* strains in small intestine-, colon content, and fecal samples collected 2-5h (small intestine and colon) or 22-24h after latest oral gavage at the end of the study. Points indicate individual data points and bars group mean with SEM. **H)** PCoA of predicted functions of the microbiota by KEGG orthology using Bray-Curtis distances of small intestine content sampled at the end of Study 1. Centroids indicate group average illustrated with 50% CI and smaller points indicate individual data points. **I)** As h for colon content. **J)** As h for fecal samples collected at week 12 of the study. **K)** PERMANOVA p-values comparing the predicted functions of the microbiota between indicated groups using Bray-Curtis distances. Asterisks indicate p-values < 0.05.

Supplementary figure S1: Baseline measures, final adipose tissue weights and replicated glucose tolerance tests

A) Baseline body weight in grams. **B)** Baseline fat mass in grams assessed by magnetic resonance (MR) scans. **C)** Baseline lean mass in grams assessed by MR scans. **D)** 5h fasting plasma insulin levels at baseline from Study 1. **E)** 5h fasting blood glucose (BG) levels at baseline from Study 1. **F)** Lean mass in grams assessed by MR scans at week 12 of the study. **G)** Fat percentage assessed by MR scans to total body weight at week 12 of the study. **H)** Weight of epididymal white adipose tissue (eWAT) at the end of the study. **I)** Weight of subcutaneous inguinal WAT (iWAT) at the end of the study. **J)** Weight of visceral retroperitoneal WAT (rpWAT) at the end of the study. **K)** Weight of visceral mesenteric WAT (mWAT) at the end of the study. **L)** Blood glucose values from Study 3 during oral glucose tolerance test (oGTT) with 2 µg/g lean mass glucose in week 10. **M)** Corresponding plasma insulin levels during oGTT shown in l. **N)** Blood glucose values from Study 2 during oral glucose tolerance test (oGTT) with 1.5 µg/g lean mass glucose in week 10. **O)** Corresponding plasma insulin levels during oGTT shown in n. **A-E)** Baseline measures prior study start while all groups were fed LFD and later divided into the indicated groups. **A-O)** Bars or lines indicate group mean ± SEM. Dashed line indicate mean of LFD group with interquartile range shown in grey. **A-K)** Points represent individual data points. Asterisks indicate fdr-corrected q-values <0.05 by linear mixed effects models comparing indicated groups. **A-C, F-K)** Point shape indicates three individual studies where rounds indicate Study 1, triangles Study 2, squares Study 3. **L-O)** Asterisks indicate p-values <0.05 using two-way ANOVA with multiple comparisons between vehicle-treated groups or *Lactobacillus* treated group to vehicle-treated group fed the same diet with Bonferroni post-hoc test.

Supplementary figure S2: Liver lipids and cytokine levels

A) Heatmap of 40 most discriminating lipids in liver tissue from end of study 1 assesed by lipidomics. Experimental group of each sample are indicated by color and distributed by hierarchical clustering. **B)** Heatmap of 100 most discriminating lipids of LFD and HFD fed groups in liver tissue from end of study 1 assesed by lipidomics. Experimental groups are indicated by color distributed by hierarchical clustering. **C)** Histological inflammation score from 0-3 blindly assessed from H&E-stained liver tissue. Asterisk indicates fdr-corrected q-value <0.05 using linear mixed effects models comparing indicated groups. **D)** Cytokine levels as fold change to LFD reference measured in liver tissue in Study 1. **C-D)** Bars indicate group mean ± SEM with points illustrating individual data points.

Supplementary figure S3: Plasma lipids , liver and cecum bile acids (BA)

A) Plasma triacylglycerol levels from the end of Study 1. **B)** Plasma free fatty acids (FFA) levels from the end of Study 1. **C)** Total measured BA concentration in liver tissue. **D)** Concentration of 3α,7α,12α-trihydroxycholestanoic acid (THCA) in liver tissue. **E)** Concentration of unconjugated BA, cholic acid (CA) -derivatives and chenodeoxycholic acid (CDCA) in liver tissue. **F)** Concentration of conjugated BA in liver tissue. **G)** Weight of full cecum in grams. Points represent individual mice in three individual studies indicated by point shape where rounds indicate Study 1, triangles Study 2, squares Study 3**. H)** Concentration of secondary bile acid lithocholic acid (LCA) in cecum content. **I)** Relative cholesterol 7 alpha-hydroxylase (Cyp7a1) gene expression in liver tissue by RT-qPCR. **A-I)** Bars indicate group mean ± SEM. Asterisks indicate fdr-corrected q-values <0.05 using linear mixed effects models comparing indicated groups. **A-D, G-I)** Dashed line indicates mean of LFD group with interquartile range shown in grey.

Supplementary figure S4: Baseline microbiota composition and alpha diversity assessments

A) Principal coordinate analysis (PCoA) of microbiota composition using weighted UniFrac distances of fecal samples collected in week 4 of Study 1. Centroids indicate group average illustrated with 50% CI and smaller points indicating individual data points. **B)** As a for fecal samples collected week 8. **C)** PCoA of baseline fecal samples collected prior to the start of Study 1 when all mice were fed LFD. Centroids indicate the later designated experimental groups. **D)** PERMANOVA p-values comparing the microbiota composition between indicated groups of Study 1 on weighted UniFrac distances. Asterisks indicate p-values < 0.05. **E)** Alpha diversity measures of observed species richness (upper panels) and Shannon's diversity index (lower panels) in small intestine and colon content from the end of the study as well as in fecal samples over time. **F)** PCoA of small intestine samples in all three studies indicated by number in the centroid. Centroid indicates group average per study projecting to individual data points. **G)** As f for

fecal samples collected at the end of the three studies. **H)** PERMANOVA p-values comparing the microbiota composition between indicated groups in individual studies on weighted UniFrac distances. Asterisk indicate p-values < 0.05. **I)** Relative abundance in % of phylum *Proteobacteria* in fecal samples and small intestine content at the end of Study 2. Boxplot indicates distribution within the groups and points indicate individual data points. Asterisk indicate p-value < 0.05 comparing the indicated groups. **J)** As i for Study 3.

Graphical abstract

Figure 1

Figure 2

Figure 3

A — Plasma lipids

Legend (heatmap):
- LFD + Vehicle
- HFD + Vehicle
- HFD + L. reuteri
- HFD + L. paracasei
- FFMD + Vehicle
- FFMD + L. reuteri
- FFMD + L. paracasei

Legend (bar graphs):
- HFD + Vehicle
- HFD + *L.reuteri*
- HFD + *L.paracasei*
- FFMD + Vehicle
- FFMD + *L.reuteri*
- FFMD + *L.paracasei*

B — Total cholesterol (mmol/L plasma)
C — FA 18:2 (Relative plasma conc.)
D — Cecal BA (mg/kg cecum content)
E — Unconjugated BA (Fold change to LFD): CA, CA1, CA2, CA3, CDCA
F — Conjugated BA (Fold change to LFD): TCA, TCDCA, TCDCA1, TCDCA2, TCDCA3, TαMCA, TβMCA, TωMCA
G — *Fgf15* (Relative expression in ileum)
H — *RegIIIγ* (Relative expression in ileum)

Figure 4

A Small intestine - Microbiota composition

B Colon - Microbiota composition

C Feces - Microbiota composition

D Microbiota composition

PERMANOVA p-value	Small intestine	Colon	Feces
HFD+Vehicle vs FFMD+Vehicle	**0.008 **	**0.001 ***	**0.004 **
HFD+Vehicle vs HFD+*L.reuteri*	0.20	0.82	0.30
HFD+Vehicle vs HFD+*L.paracasei*	0.42	0.62	0.69
FFMD+Vehicle vs FFMD+*L.reuteri*	0.12	0.08	0.10
FFMD+Vehicle vs FFMD+*L.paracasei*	0.64	0.82	0.88

E Small intestine / Colon / Feces

F Taxa summary

G ASV15 / ASV5 / ASV31

Legend:
- HFD + Vehicle
- HFD + *L.reuteri*
- HFD + *L.paracasei*
- FFMD + Vehicle
- FFMD + *L.reuteri*
- FFMD + *L.paracasei*

H Small intestine - Predicted functions

I Colon - Predicted functions

J Feces - Predicted functions

K Predicted functions

PERMANOVA p-value	Small intestine	Colon	Feces
HFD+Vehicle vs FFMD+Vehicle	**0.006 **	**0.017 *	**0.009 **
HFD+Vehicle vs HFD+*L.reuteri*	0.47	0.84	0.26
HFD+Vehicle vs HFD+*L.paracasei*	0.36	0.57	0.17
FFMD+Vehicle vs FFMD+*L.reuteri*	**0.04 *	**0.01 *	**0.04 *
FFMD+Vehicle vs FFMD+*L.paracasei*	0.55	0.44	0.99

Figure S1

Figure S2

A Liver lipids (All groups)

B Liver lipids (LFD and HFD groups)

C Inflammation score

D Hepatic cytokine levels

Figure S3

Figure S4