1 **Overruled by nature: A plastic response to an ecological regime**

2 **shift disconnects a gene and its trait**

4 F. Besnier[1], Ø. Skaala[1], V. Wennevik[1], F. Ayllon[1], K.R. Utne[1], P.T. Fjeldheim[1], K.

5 Andersen-Fjeldheim[1], S. Knutar[1], K.A. Glover[1,2]

7
8
9 [1] Institute of Marine Research, Bergen, Norway
10 [2] Institute of Biology, University of Bergen, Norway
11
12
13
14

Abstract

In Atlantic salmon, age at maturation represents a life history trait with a sex-specific trade-off between reproductive success and survival, mediated by the *vgll3* gene through sex-specific dominance. By genotyping ~1500 historical and contemporary samples of salmon from the river Etne with a 50k SNP-array, we identified three loci significantly associated with sea-age. These included *vgll3* and *six6*, which collectively explained 50% and 36% of the age of maturation for males and females in the 1983-1984 period respectively. In the same period, we also observed sex-specific epistasis, where the *six6* genotype only influenced age at maturation in males carrying the late-maturation *vgll3* allele. Strikingly, the combined influence of these genes was nearly absent in all samples from 2013-2016, while allelic frequencies at *vgll3* remained unchanged. Many salmon populations in the northeast Atlantic, including that from Etne, have been exposed to a major ecological regime shift in the marine environment in 2005. Because this is associated with consistently slower growth and higher age at maturation, we conclude that the regime shift has led to the sudden bypassing of the influence of *vgll3* and *six6* on maturation through growth-driven plasticity.

Introduction

Understanding the mechanisms by which organisms adapt to their environments is a central question in biology [1,2]. However, beyond the academic curiosity that motivates biologists to investigate how species evolved and adapted until now, accelerating climate change and ongoing habitat destruction catalyses a sense of urgency when considering the fate of many species in the future. As unprecedentedly fast environmental changes have occurred for many taxa during the last half century [3-5], the current mechanisms of adaptation of many species may not couple with the pace at which environmental change is now occurring, leading thus to population declines and extinctions. Consequently, the mechanisms of adaptation and their potential to remain effective in a context of rapidly changing environment are becoming increasingly important to understand [6-9].

In the context of environmental change, genes associated with sexual conflicts [10] are of particular interest. Sexual conflicts are associated with sex-specific selection and therefore expected to promote genetic diversity in the underlying genes [11], thus potentially helping maintain standing genetic variation on important fitness traits within the population. An illustration of this occurs in anadromous Atlantic salmon (*Salmo salar* L.) where a single gene (*vgll3*) is strongly associated with the age at maturation, a highly adaptative trait subject to sexual conflict [12,13]. In many species, age at maturation represents a trade-off, as late-maturation gives larger body sizes and thus higher reproductive success, but at the increased risk of dying before reproduction [14]. In Atlantic salmon, the optimum value for this trait differs between sexes, as males mature earlier and smaller whereas females benefit from later maturation and a larger body size resulting in more and larger eggs [14].

This sexual conflict is in part resolved by the mediation of *vgll3* [12] through sex-

specific dominance.

Studies of *vgll3* and age of maturation in Atlantic salmon have resulted in

inconstant estimations regarding the degree to which this gene influences the trait.

Several studies have demonstrated a very strong association in wild North European

populations [12,13,15,16], but in stark contrast, little or no association has been observed

in wild populations in North America [17] [18]. Furthermore, conflicting observations have

also been reported in in domesticated Norwegian farmed strains reared under

aquaculture conditions [19] [20]. This begs the question, why does the influence of *vgll3*

on age at maturation vary so greatly?

Salmonids have been exposed to a wide range of anthropological challenges

including habitat modifications since the Industrial Revolution [21]. Furthermore, during

the past two decades, most salmon populations in the North Atlantic have shifted

towards later maturation [22,23] and smaller size at age [23-26]. Although these changes

are thought to be caused by a regime-shift in oceanic conditions in 2005 [23], whether

they reflect rapid evolution or plastic responses is unknown. The same trends

towards later maturation have also been documented in the salmon population

inhabiting the river Etne on the west coast of Norway, where the proportion of fish

maturing after one year at sea dropped from 63% in the period 1983-84 to 34% in

2018-19 [27]. The main objective of the present study was to investigate whether the

observed changes in age at maturation were the result of phenotypic plasticity, or

alternatively, evolution in the gene(s) influencing this trait. To address this, we

genotyped historical (early 1980´s – pre regime shift) and contemporary (mid 2010´s

post regime shift) samples with a 50k panel of genome-wide SNPs.

Results

Temporal changes in age and size at maturation

Marine growth was compared between the historical (1983-84) and contemporary

(2013-16) samples (Fig. 1). During this period, the length of the fish during the first

winter at sea decreased significantly, as well the length of the two sea-winter (2SW)

adults, while the length of the one sea-winter (1SW) adults remained stable (Fig. 1).

During the same period, we also observed a trend towards later maturation

with a change from 46% to 8% 1SW in the female group, and from 75% to 56% in

the male group (Table 1).

Genome scans

The genome scan for age at maturation (Fig. 2) identified three loci displaying a

significant association in the historical samples, one on each of chromosomes

SSA09, SSA24 and SSA25. The genomic regions identified in SSA25 and SSA09

overlapped with loci previously described as major contributors to the sea age

variability in salmon: *vgll3* on SSA25 and *six6* located on SSA09. The degree of

association between age at maturation and these genomic regions was investigated

further in the historical and contemporary dataset.

***vgll3* and *six6* haplotypes**

Haplotypes were reconstructed in the historical and contemporary samples

separately, across the genomic regions in SSA25 and SSA9 that contained the *vgll3*

and s*ix6* genes respectively. On SSA25, two haplotypes were predominant in the

historical samples with frequencies of 56% and 39% (Table 2) and respectively

associated with early and late sea age. The mean sea age was 1.18 for the

homozygous haplotype "1221", and 2.46 for the homozygous haplotype "2112",

which will thus be referred to as *vgll3*-E and *vgll3*-L alleles from hereon. In the

contemporary samples, the same two haplotypes were found in almost identical

frequencies (df=1, χ^2=0.85, p=0.35) to the historical samples (59% and 39% for

vgll3-E and *vgll3*-L respectively). Indicating that no temporal change in haplotype

frequencies occurred at this locus during the three-decade period.

The mean age at maturation in each genotype class (Fig. 3) displayed a

strong association with an additive effect of *vgll3* on female sea age in the historical

data (Fig. 3.a), whereas association of *vgll3* almost disappeared in the contemporary

female data (Fig. 3.c). For the historical male data (Fig. 3b), a strong *vgll3*

association with a dominant *vgll3*-E allele was also detected, whereas the genetic

association was strongly reduced in the contemporary data (Fig. 3.d).

On SSA09, four main haplotype sequences linked to the *six6* gene were

predominant in both historical and contemporary samples (Table 2). With a

respective sea age of 1.3 and 1.4 years for the homozygous fish in the historical

samples, both "111" and "211" haplotypes were assigned to the early variant of *six6,*

whereas haplotypes "121" and "122", with a respective mean sea age of 1.9 and 2.1

years, were assigned to the late variant. Following this, we observed an increase in

frequency of the *six6* late variant, from 44% in the historical samples to 60% in the

contemporary samples (df=1 χ^2=111, p<$2.2.10^{-16}$). The mean age at maturation in

each genotype class (Fig. 3) showed an additive effect of *six6* in the historical female

data (Fig. 3.a), and weak *six6* association in the historical male samples (Fig. 3.b).

Cumulatively, *vgll3* and *six6* accounted for 50% of the model deviance in the

historical male dataset and 36% in the historical female dataset. In stark contrast, the

same loci only accounted for 7% and 3% of the model deviance in the contemporary

male and female datasets respectively.

Testing the interaction between these loci revealed significant departure from

additivity between *vgll3* and *six6* in the males of the historical samples. This

interaction is illustrated in Fig. 3.b where the *six6* genotype is only associated with

age at maturation for the samples that are homozygous for the *vgll3-L* allele. No

significant departure from additivity was observed in the historical female samples

(df=1 χ^2=0.3 p=0.18), nor in the contemporary male (df=1 χ^2=0.28 p=0.26) and

female (df=1 χ^2=0.07 p=0.32) samples.

Genetic admixture

The salmon population inhabiting the river Etne has been subject to 24%

introgression from domesticated farmed escapees [28-30]. We therefore tested whether

the observed temporal change in genetic architecture of age at maturation could be

linked to admixture with farmed fish. However, no interaction was detected between

individual admixture and *vgll3* genotype (χ^2=3.6, df=2, p=0.161), nor between

individual admixture and *six6* genotype (χ^2=2.7, df=2, p=0.24). We also checked the

allelic frequencies at the *six6* locus in the farmed samples to assess if the increase in

six6-L allele observed in the contemporary sample could be explained by admixture.

With 17% *six6-E* and 68% *six6-L* allele, haplotype frequencies in the farmed samples

were more like the contemporary samples than the historical. However, the observed

genetic admixture in this population, which is estimated to circa 20-25%, could only

explain a 6% increment in frequency of *six6*-L, which is less than half of the increase

observed between historical and contemporary samples.

Discussion

We document the sudden dissociation between age at maturation in a population of

Atlantic salmon and the genotypes of three loci, including the previously identified

candidate genes *vgll3* and *six6* [12,13,15]. This dissociation was observed in samples

separated by a 30-year time interval and was accompanied by an increase in the

frequency of late maturing allele at the *six6* locus, while no change was observed at

the *vgll3* locus. The salmon population in the river Etne has been subject to

introgression from domesticated salmon escaping from commercial fish farms, with

an average 24% of genetic admixture in the contemporary population [28-30]. However,

from the observed allelic frequencies at the *six6* locus, genetic admixture could not

explain alone the change in allelic frequencies between historical and contemporary

samples. In addition, the possibility that the dissociation could have been caused by

genetic introgression from domesticated salmon escapees was discarded by the

absence of interaction between genetic effects and admixture.

The combined effect of *vgll3* and *six6* was not fully additive as we detected a

significant epistatic interaction in the historical samples of males. *Six6* was

previously identified as a candidate locus associated with age at maturation [12], but

little is known about the genetic effects of this locus. The work presented here is the

first to document a sex-specific interaction between *vgll3* and *six6.*

In the river Etne, the frequency of late maturing fish was higher among

females than among males. This difference is assumed to be the result of adaptation

to sexual conflict where the advantage of maturing late is greater for females than for

males [14]. With the *vgll3-E* allele being dominant in males only, *vgll3* is believed to

play a key function in the resolution of sexual conflict in the optimal age of maturation

in salmon [12,13]. However, with the present results documenting the almost complete

dissociation between age at maturation and *vgll3* genotypes in parallel with a trend

towards later maturation strategy in both sexes simultaneously, the perennity of local

adaptation in the population inhabiting the river Etne can be questioned. Noteworthy,

sex-specific strategies of maturation seem to perdure in the population despite

vgll3's contribution being almost completely eradicated in the contemporary samples.

In fact, the difference in early-maturation frequencies between males and females

increases in the contemporary dataset, strongly suggesting that other mechanisms

are acting instead of, or in parallel with *vgll3*, to maintain sex specific age at

maturation. This hypothesis seems consistent with the description of multiple loci

associated with age at maturation [20].

The absence of association between *vgll3* and age at maturation in multiple

North American populations [17,18] already suggested that *vgll3* is not the only

regulator of sexual conflict for age of maturation in Atlantic salmon. The present

study confirms this observation and further shows that the relative influence of *vgll3*

is not stable in time. In the case of the population in the river Etne, our data reveals

that other unidentified factors also play a major role in maintaining the sex-specific

differences in maturation strategies when the link between age at maturation and

vgll3 is disconnected.

The observed trend towards slower marine growth and later maturation in the

population inhabiting the river Etne has also been reported in many other salmon

populations in the North Atlantic [22-26]. While the precise triggers and mechanisms

underpinning to the development of sexual maturation are not yet fully understood in

Atlantic salmon, slow growth and late maturation is consistent with the hypothesis of

size or perhaps growth-rate threshold as a determinant for salmon maturation [31-33].

Such an energy-budget threshold might also be genetically regulated through the

mediation of *vgll3*, which has been shown to be linked with cell fat regulation [34].

Following this hypothesis, slow growth due to environmental conditions, such as lack

of prey would lead to later maturation as a higher proportion of fish would fail to

reach the weight (or growth-rate) threshold for maturation after one single year at

sea.

Observations suggest important changes in the environmental conditions met

by salmon migrating from the river Etne in the period 1980-2010 [23]. The marine

migration pattern for salmon originating from the river Etne is not fully known,

however, the first phase is probably a northward migration through the southern

Norwegian Sea [35] where changes in the oceanographic conditions have been

reported in the last years. For example, water temperature increased by nearly 1˚C

at 50-200 m depth from early 1980's until 2021, mainly due to warmer water masses

flowing into the southern Norwegian Sea [36,37]. The early 1980's are also considered

as the end of "The great salinity anomaly", which was a period with large inflow of

cold and fresh Arctic water into the Norwegian Sea. The large proportion of Arctic

water entering the Norwegian Sea is correlated to increased productivity [38], and to

improved feeding conditions for post-smolts in the region [39]. Therefore, salmon

returning in 1983 and 1984 had probably been feeding in a very productive sea

during the initial post-smolt phase, whereas salmon returning to rivers in 2013-2016

had been feeding in a warm and saline Norwegian Sea. During this later period,

observed stomach fullness and condition factor for post-smolt sampled in the

Norwegian Sea were low [40]. In addition, the potential interspecific competition with

other pelagic fish for prey [41] was low in the early 1980's as the total biomass of

pelagic fish feeding in the Norwegian Sea in 1982-1983 was around 1/3 of the total

biomass in the period 2013-2016 [42,43] when Norwegian Spring-spawning herring had

not yet recovered from the collapse in the late 1960ies, and NEA-mackerel and blue

whiting stock biomasses were at low levels [43]. Many observations seem to confirm

that environmental conditions were less favorable for salmon growth in the last

decade than in the early 1980's. It is thus conceivable that the changes observed in

the river Etne are caused by environmental perturbations occurring over a large

region and affecting salmon populations as well as other organisms.

Under commercial aquaculture conditions, farmed Atlantic salmon typically

display early sexual maturation [33], and importantly, the relative influence of *vgll3* on

age at maturation appears to be largely bypassed [19]. Yet again, this response is sex-

specific as a study conducted under aquaculture conditions [19] reported that *vgll3* did

not show any correlation with age at maturation in females while displaying only

weak association in males [19]. The commercial farm strain used in the

aforementioned study, known as Mowi, stemmed from wild Norwegian salmon

populations in which *vgll3* was identified as a candidate gene for influencing age at

maturation [12,13]. The authors concluded that high calorie feed intake combined with

artificial light and temperature regimes as well as potential genetic or epigenetic

components, may alter the impact of *vgll3* on age at maturation [19]. Anecdotally, it is

also worth noting that despite multiple generations of directional selection against

early maturing fish in the domesticated farmed salmon, the genetic variability of *vgll3*

remains high in the Mowi strain [19]. It is thus possible that a rapid change in

environmental conditions, such as feed intake and therefore growth-rate, may

bypass the effect of *vgll3* without letting selection (artificial selection in this case)

alter the allelic frequencies of the gene.

We thus can hypothesize a two-threshold model where the effect of *vgll3* on

age at maturation is bypassed when abundance of feed resources is very high, as in

farming conditions, or when feed resources are very low and fish need to spend

more time at sea to acquire the necessary energy-reserves for maturation and

reproduction. However, when resource availability falls between both thresholds,

vgll3 may play a more important role in determining age at maturation. Inversely if

feed resources exceed the high threshold, or do not reach the low one, *vgll3* is

bypassed by environmental conditions, and therefore age at maturation is

determined by a combination of other genetic and environmental factors.

In the face of changing environmental conditions, one may expect the allelic

frequencies at the associated loci to change in response to the newly induced

selection pressure [15,16]. This is the case for *six6* where the temporal trend towards

later maturation is accompanied by an increase of the *six6*-L allele frequency. This

shift in allelic frequencies is likely due to positive selection on the *six6-L* allele, alone,

or in conjunction with admixture from domesticated salmon where *six6*-L was found

in higher frequency. In contrast, the *vgll3* allelic frequencies remained stable despite

major changes in age at maturation. This result seems to indicate a change in

environmental conditions, creating a situation where the influence of *vgll3* was

effectively bypassed before natural selection had time to operate, whereas selection

had the time to modify the allelic frequencies at the *six6* locus before the influence of

this gene was also bypassed. This hypothesis is strongly supported by the

observations of sudden change in age at maturation in many Atlantic salmon

populations inhabiting Northeast Atlantic rivers [23], including the population in the

river Etne [27], where a highly distinct and sudden drop in early marine growth was

reported in 2005 and subsequently, referred to as an ecological regime shift in the

northeast Atlantic ocean [23].

As this, and other salmon population in the Northeast Atlantic were subjected

to an ecological regime shift, leading to consistently reduced marine growth rates

and increased age at maturation, we conclude that growth-driven plasticity has

almost completely bypassed the combined influence of *vgll3* and *six6* within on age

at maturation, and furthermore, on resolving the sexual conflict for this trait in Atlantic

salmon. Despite the genetic dissociation between genes and phenotype, sex-specific

strategies of maturation age remained in the contemporary population, suggesting

that other genetic or environmental factors are acting instead of or in addition to *vgll3*

to maintain such sex-specific strategy.

Methods

Samples

This study is based on samples obtained from adult salmon captured in the river

Etne, in western Norway, 6o° N. This river is home to a salmon population of

typically 1000-2500 adults returning from the sea annually [44]. A permanent trapping

facility installed in the river has permitted sampling almost the entire adult spawning

population since 2014, which facilitated access to both an extensive set of DNA

samples, as well as to phenotypic and phenological data. For the present study, 797

wild adult salmon captured in the 1983-84 angling season were compared to 751

wild adult salmon captured in the upstream fish trap in the period 2013 to 2016 [28].

For the contemporary samples (2013-2016), an estimation of individual genetic

admixture was computed as the proportion of domestic ancestry in each individuals

genome (see Besnier, et al. [28] for details). In addition, a sample of 350 domesticated

farmed salmon escapees that were removed from the river in the period 1989-2012

(15 per year), were genotyped with the same set of markers.

309 Genotyping and sex determination

All samples were genotyped on a ThermoFisher Axiom 57K single nucleotide

polymorphism (SNP) array (NOFSAL03, 55735 markers) developed by Nofima

(Norwegian institute for applied research in food aquaculture and fisheries) in

collaboration with private aquaculture companies Mowi and SalmoBreed [28]. SNPs

were checked following the "Best Practice Workflow" on the Affymetrix axiom

analysis software (available at: https://www.thermofisher.com/no/en/home/technical-

resources/software-downloads.html). SNPs with call rates lower than 0.97 and

samples with call rates lower than 0.85 were discarded, whereas markers classified

as "PolyHighResolution" (High resolution in both homozygous and heterozygous

clusters) were conserved for further data analysis.

Fish were sexed by examining variants of the sdY gene [45,46]; *i.e.,* males were

identified based on the presence of exons 2 and 4. Samples were genotyped on an

Applied Biosystems ABI 3730 Genetic Analyser, and genotypes were called using

GeneMapper (Applied Biosystems, v. 4.0). The analysis of the sdY gene provides an

accurate identification of sex, although, a very low percentage of fish identified as

genetic males are phenotypic females due to carrying an inactive pseudo-copy of the

sdY gene with both exon 2 and 4 [47].

328 Genome scan for loci associated with age at maturation

Age at maturation was modeled as a binary trait consisting of early maturing fish,

i.e., fish returning after one sea-winter (1SW), also known as grisling, *vs.* late

maturing fish, *i.e.*, fish returning after two or more winters at sea (2^+SW). The

probability of maturing early was then modeled in a generalized linear model with

logit link function:

$$\log\left(\frac{P_{(e)}}{1-P_{(e)}}\right) = Gi + sex + year + e \qquad\qquad \text{(Model 1)}$$

Where $P_{(e)}$ is the probability of early maturing, G_i is the SNP genotype at locus *i, sex*

is a binary factor accounting for genetically determined sex, *year* is a factor

accounting for the sampling year, and *e* a normally distributed vector of residuals.

Significance of association between genotype and sea age was estimated by

comparing the deviance of Model 1 with the deviance of the model without genetic

effect. The difference was compared to a Chi-squared distribution with one degree of

freedom. Model 1 was fitted at each SNP available on the dataset, and the obtained

p-values were adjusted for multiple testing by following Bonferroni correction. All p-

values given in the genome scan result section are corrected for multiple testing.

Haplotyping

Haplotypes were reconstructed using the Phase 2.1 software [48] in the historical and

contemporary data separately. Two loci on SSA9 and SSA25 were more specifically

considered for haplotype reconstruction as they displayed high association with age

at maturation, and associated genes *vgll3* and *six6* were previously described in the

same two genomic regions [12,13,15]. On SSA25, a haplotype window consisting of four

polymorphic SNPs (AX-87309414, AX-172546510, AX-87309615, AX-87420691)

was reconstructed in the region spanning from 28.65 to 28.66 Mb containing *vgll3*

(https://www.ncbi.nlm.nih.gov/gene/106586514). On SSA09, a haplotype window

consisting of three SNPs (AX-172546568, AX-88029383, AX-87668000), spanning

from   24.86-24.95Mb   was   reconstructed   around   the   position   of   *six6.*

(https://www.ncbi.nlm.nih.gov/gene/106610974)

Statistical analyses

Genetic structure to age at maturation variation

Due to the sex specific dominance observed in *vgll3*, the genetic structure was

estimated for males and females separately. The probability of early maturation was

modeled as a response to additive and dominance genetic effects as follow:

$log \frac{P_{(e)}}{1-P_{(e)}} = a_i + d_i + year + e$ (Model 2)

Where $P_{(e)}$ is the probability of early maturing, a_i and d_i are respectively the additive

and dominance effects at locus *i*, *year* is a factor accounting for the sampling year,

and *e* a normally distributed vector of residuals. The model was fitted in a GLM with

logit link function in R [49], and the variance contribution was deduced from the

difference in model deviance between Model 2 and a model than only accounted for

capture year. Two loci interaction models were tested similarly by accounting for the

additional interaction parameters (a_i*a_j; a_i*d_j, a_j*d_i; d_i*d_j) between loci $_i$ and $_j$. In

contrast with the genome scan, the test for significance if the different genetic

parameters were not corrected for multiple testing. All p-values given with the

estimation of genetic effect are nominal values.

Comparison in age at maturation

Potential differences in age at maturation between historical and contemporary

samples was tested by chi-squared test on a contingency table reporting the number

of observed 1, 2 and 3+ sea winter adults within historical and contemporary

samples.

Size at age

Individual body length was recorded for every fish passing the trap in the

contemporary sample, whereas the adult length of the 1983-1984 fish were

calculated from reading scales samples. In addition, the growth at first sea winter

was estimated for both historical and contemporary samples by calculating length at

first sea winter from reading scales. The difference in length between samples from

1983-1984 and from 2013-2016 was tested with a two-sided t-test, separately for

each sea-age and sex categories.

Potential role of admixture

The salmon population in the river Etne has been subject to introgression from

domesticated salmon escaping from commercial fish farms, with an average 24% of

genetic admixture in the contemporary population [28-30]. Individual admixture with

domesticated salmon has already been correlated with earlier adult maturation in this

population [28]. Therefore, in order to account for the potential influence of admixture

on the temporal influence of loci on age at maturation in this population, we

investigated whether the change in genetic architecture of age at maturation could

be linked to genetic admixture.

A third model was fitted with the aim to evaluate a potential interaction

between admixture and *vgll3* or *six6* genotypes.

$log \frac{P_{(e)}}{1-P_{(e)}} = a_i * admix + d_i * admix + year + e$ (Model 3)

Where *admix* is the individual genetic admixture as computed in Besnier et al [28].

Data availability

405   Data for this study are available from the IMR.brage.unit.no public repository.

406   Link to data: https://imr.brage.unit.no/

References

1. Losos, J.B. Ecological character displacement and the study of adaptation. *PNAS* **97**, 5693-5695 (2000).
2. Andrew, R.L. *et al.* A road map for molecular ecology. *Mol. Ecol.* **22**, 2605-26 (2013).
3. Chevin, L.M., Lande, R. & Mace, G.M. Adaptation, plasticity, and extinction in a changing environment: towards a predictive theory. *PLoS Biol.* **8**, e1000357 (2010).
4. Pörtner, H.O. *et al.* IPCC, 2022: Climate Change 2022: Impacts, Adaptation, and Vulnerability. Contribution of Working Group II to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change. (2022).
5. Dullinger, S. *et al.* Extinction debt of high-mountain plants under twenty-first-century climate change. *Nat. Clim. Change* **2**, 619-622 (2012).
6. Candolin, U. & Heuschele, J. Is sexual selection beneficial during adaptation to environmental change? *Trends Ecol. Evol.* **23**, 446-52 (2008).
7. Kwan, L., Bedhomme, S., Prasad, N.G. & Chippindale, A.K. Sexual conflict and environmental change: trade-offs within and between the sexes during the evolution of desiccation resistance. *J. Genet.* **87**, 383–394 (2008).
8. Bandillo, N.B. *et al.* Dissecting the genetic basis of local adaptation in soybean. *Sci. Rep.* **7**, 17195 (2017).
9. García de Leániz, C. *et al.* A critical review of adaptive genetic variation in Atlantic salmon: implications for conservation. *Biol. Rev.* **82**, 173-211 (2007).
10. Parker, G.A. Sexual selection and sexual conflict. in *Sexual Selection and Reproductive Competition in Insects* (eds. BLUM, N.A. & BLUM, M.S.) 123-166 (Academic Press, Cambridge, 1974).
11. Rowe, L., Chenoweth, S.F. & Agrawal, A.F. The genomics of sexual conflict. *Am. Nat.* **192**, 274-286 (2018).
12. Barson, N.J. *et al.* Sex-dependent dominance at a single locus maintains variation in age at maturity in salmon. *Nature* **528**, 405-408 (2015).
13. Ayllon, F. *et al.* The *vgll3* locus controls age at maturity in wild and domesticated Atlantic salmon (*Salmo salar* L.) males. *PLoS Genet.* **11**, e1005628 (2015).
14. Fleming, I.A. & Einum, S. Reproductive Ecology: A Tale of Two Sexes in *Atlantic Salmon Ecology* (eds. Aas, Ø., Einum, S., Klemetsen, A. & Skurdal, J.) 33-65 (Blackwell Publishing, 2011).
15. Czorlich, Y., Aykanat, T., Erkinaro, J., Orell, P. & Primmer, C.R. Rapid sex-specific evolution of age at maturity is shaped by genetic architecture in Atlantic salmon. *Nat. Ecol. Evol.* **2**, 1800-1807 (2018).
16. Jensen, A.J. *et al.* Large-effect loci mediate rapid adaptation of salmon body size after river regulation. *Proc Natl Acad Sci U S A* **119**, e2207634119 (2022).
17. Boulding, E.G., Ang, K.P., Elliott, J.A.K., Powell, F. & Schaeffer, L.R. Differences in genetic architecture between continents at a major locus previously associated with sea age at sexual maturity in European Atlantic salmon. *Aquac.* **500**, 670-678 (2019).
18. Mohamed, A.R. *et al.* Polygenic and sex specific architecture for two maturation traits in farmed Atlantic salmon. *BMC Genom.* **20**, 139 (2019).
19. Ayllon, F. *et al.* The influence of *vgll3* genotypes on sea age at maturity is altered in farmed mowi strain Atlantic salmon. *BMC Genet.* **20**, 44 (2019).

20.   Sinclair-Waters, M. *et al.* Beyond large-effect loci: large-scale GWAS reveals a mixed
large-effect and polygenic architecture for age at maturity of Atlantic salmon. *Genet*
Sel Evol **52**, 9 (2020).

21.   Forseth, T. *et al.* The major threats to Atlantic salmon in Norway. *ICES J. Mar. Sci.* **74**,
1496-1513 (2017).

22.   Otero, J. *et al.* Contemporary ocean warming and freshwater conditions are related
to later sea age at maturity in Atlantic salmon spawning in Norwegian rivers. *Ecol.*
Evol. **2**, 2192-203 (2012).

23.   Vollset, K.W. *et al.* Ecological regime shift in the Northeast Atlantic Ocean revealed
from the unprecedented reduction in marine growth of Atlantic salmon. *Sci. Adv.*
8(2022).

24.   Quinn, T.P., McGinnity, P. & Cross, T.F. Long-term declines in body size and shifts in
run timing of Atlantic salmon in Ireland. *J. Fish Biol.* **68**, 1713-1730 (2006).

25.   Bal, G. *et al.* Evidence for long-term change in length, mass and migration phenology
of anadromous spawners in French Atlantic salmon *Salmo salar*. *J. Fish Biol.* **90**,
2375-2393 (2017).

26.   Todd, C.D., Hanson, N.N., Boehme, L., Revie, C.W. & Marques, A.R. Variation in the
post-smolt growth pattern of wild one sea-winter salmon (*Salmo salar* L.), and its
linkage to surface warming in the eastern North Atlantic Ocean. *J. Fish Biol.* **98**, 6-16
(2021).

27.   Harvey, A.C. *et al.* Time series covering up to four decades reveals major changes
and drivers of marine growth and proportion of repeat spawners in an Atlantic
salmon population. *Ecol. Evol.* **12**, e8780 (2022).

28.   Besnier, F. *et al.* Introgression of domesticated salmon changes life history and
phenology of a wild salmon population. *Evol. Appl.* **15**, 853-864 (2022).

29.   Glover, K.A. *et al.* Atlantic salmon populations invaded by farmed escapees:
quantifying genetic introgression with a Bayesian approach and SNPs. *BMC Genet.*
14, 74 (2013).

30.   Karlsson, S., Diserud, O.H., Fiske, P. & Hindar, K. Widespread genetic introgression of
escaped farmed Atlantic salmon in wild salmon populations. *ICES J. Mar. Sci.* **73**,
2488-2498 (2016).

31.   Rowe, D.K., Thorpe, J.E. & Shanks, A.M. Role of fat stores in the maturation of male
Atlantic salmon (*Salmo salar*) parr. *Can. J. Fish. Aquat. Sci.* **48**, 405-413 (1991).

32.   Simpson, A.L. Differences in body size and lipid reserves between maturing and
nonmaturing Atlantic salmon parr, *Salmo salar* L. *Can. J. Zool.* **70**(1992).

33.   Taranger, G.L. *et al.* Control of puberty in farmed fish. *Gen. Comp. Endocrinol.* **165**,
483-515 (2010).

34.   Halperin, D.S., Pan, C., Lusis, A.J. & Tontonoz, P. Vestigial-like 3 is an inhibitor of
adipocyte differentiation. *J. Lipid Res.* **54**, 473-81 (2013).

35.   Gilbey, J. *et al.* The early marine distribution of Atlantic salmon in the North-east
Atlantic: A genetically informed stock-specific synthesis. *Fish Fish.* **22**, 1274-1306
(2021).

36.   ICES. ICES Working Group on the Integrated Assessments of the Norwegian Sea
(WGINOR; outputs from 2020 meeting). 3. 114 pp. (2021).

37.   Skagseth, Ø. & Mork, K.A. Heat content in the Norwegian Sea, 1995–2010. *ICES J.*
Mar. Sci. **69**, 826-832 (2012).

38.   Skagseth, Ø. *et al.* Arctic and Atlantic waters in the Norwegian Basin, between year
500   variability and potential ecosystem implications. *Front. Mar. Sci.* **9**(2022).
501   39.   Utne, K.R. *et al.* Impacts of a changing ecosystem on the feeding and feeding
502   conditions for Atlantic salmon during the first months at sea. *Front. Mar. Sci.*
503   **9**(2022).
504   40.   Utne, K.R. *et al.* Poor feeding opportunities and reduced condition factor for salmon
505   post-smolts in the Northeast Atlantic Ocean. *ICES J. Mar. Sci.* **78**, 2844-2857 (2021).
506   41.   Utne, K.R. *et al.* Feeding interactions between Atlantic salmon (*Salmo salar*)
507   postsmolts and other planktivorous fish in the Northeast Atlantic. *Can. J. Fish. Aquat.*
508   *Sci.* **78**, 255-268 (2021).
509   42.   ICES. Working Group on Widely Distributed Stocks (WGWIDE). 3:95. 874 pp. (2021).
510   43.   ICES. Report of the Working Group on Widely Distributed Stocks (WGWIDE). (2008).
511   44.   Harvey, A.C., Tang, Y., Wennevik, V., Skaala, O. & Glover, K.A. Timing is everything:
512   Fishing-season placement may represent the most important angling-induced
513   evolutionary pressure on Atlantic salmon populations. *Ecol. Evol.* **7**, 7490-7502
514   (2017).
515   45.   Eisbrenner, W.D. *et al.* Evidence for multiple sex-determining loci in Tasmanian
516   Atlantic salmon (*Salmo salar*). *Heredity* **113**, 86-92 (2014).
517   46.   Yano, A. *et al.* An immune-related gene evolved into the master sex-determining
518   gene in rainbow trout, *Oncorhynchus mykiss*. *Curr. Biol.* **22**, 1423-8 (2012).
519   47.   Ayllon, F. *et al.* Autosomal sdY pseudogenes explain discordances between
520   phenotypic sex and DNA marker for sex identification in Atlantic salmon. *Front.*
521   *Genet.* **11**, 544207 (2020).
522   48.   Stephens, M., Smith, E.P. & Donnelly, P. A new statistical method for haplotype
523   reconstruction from population data. *Am. J. Hum. Genet.* **68**, 978–989 (2001).
524   49.   Team, R.C. R: A language and environment for statistical computing. in *R Foundation*
525   *for Statistical Computing* (Vienna, Austria, 2022).
526
527
528

Acknowledgements

This work was financed by the Norwegian Ministry of Trade, Industry and Fisheries. This authority paid no part in the study design nor interpretation of results. We would like to acknowledge the research institute Nina, and anglers, for donating some of the farmed salmon samples used in this study. We would like to acknowledge the river Etne owners association for continued collaboration and allowing capture and handling of fish and IMR staff involved in monitoring the population in the trap. We would like to thank Maria Quintela Sanchez for insightful comments on the earlier drafts of the manuscript, and Emily K. Glover for drawing the salmons used in Figure 1.

Authors' contributions

Ø.S and K.A.G conceived and designed the study.

P.T.F, K.A.-F and Ø.S conducted the fieldwork and sample collection.

S.K conducted the laboratory work.

F.B analysed and interpreted the data with critical input from V.W, K.R.U, K.A.G, F.A

F.B and K.A.G led the writing with contributions from all co-authors.

Competing interests

The authors declare no competing interests.

549 **TABLES**

551 **Table 1.** Percentage and numbers (in brackets) of fish displaying 1-2-3$^+$ age at

552 maturation, by sex, in the historical (H – 1983/84, N = 797) and contemporary (C –

553 2013-2016, N = 751) samples.

	1SW % (N)	2SW % (N)	3$^+$SW % (N)
Females_H	46 (141)	37 (114)	17 (51)
Females_C	8 (30)	63 (227)	28 (102)
Males_H	75 (311)	13 (56)	12 (48)
Males_C	58 (227)	33 (127)	9 (36)

554
555

	Haplotypes	Historical				Contemporary				Farmed	
		F		M		F		M		F	M
		%(N)	M.S.A	%(N)	M.S.A	%(N)	M.S.A	%(N)	M.S.A	%(N)	%(N)
vgll3	1221	$59_{(364)}$	1.2	$54_{(455)}$	1.1	$61_{(438)}$	2.2	$58_{(454)}$	1.4	$53_{(93)}$	$63_{(264)}$
	1222	$0_{(0)}$	NA	$0_{(1)}$	NA	$0_{(0)}$	NA	$0_{(1)}$	NA	$0_{(0)}$	$0_{(0)}$
	1111	$1.5_{(9)}$	NA	$2_{(17)}$	NA	$0_{(0)}$	NA	$0_{(0)}$	NA	$0_{(0)}$	$0_{(1)}$
	1112	$0_{(1)}$	NA	$0_{(0)}$	NA	$0_{(0)}$	NA	$0_{(1)}$	NA	$0_{(0)}$	$0_{(1)}$
	2221	$1_{(9)}$	NA	$2_{(18)}$	NA	$1_{(9)}$	NA	$1(13)$	NA	$3_{(5)}$	$5_{(24)}$
	2222	$0_{(0)}$	NA	$0.5_{(3)}$	NA	$0_{(3)}$	NA	$0_{(0)}$	NA	$0_{(0)}$	$0_{(0)}$
	2111	$1.5_{(5)}$	NA	$0.5_{(4)}$	NA	$0_{(0)}$	NA	$1_{(0)}$	NA	$0_{(0)}$	$0_{(1)}$
	2112	$37_{(230)}$	2.5	$41_{(344)}$	2.5	$38_{(272)}$	2.5	$40_{(309)}$	2.2	$44_{(76)}$	$31_{(128)}$
six6	111	$41_{(256)}$	1.4	$40_{(337)}$	1.2	$23_{(166)}$	2.3	$25_{(195)}$	1.6	$16_{(29)}$	$14_{(65)}$
	112	$0_{(3)}$	NA	$1_{(8)}$	NA	$0_{(0)}$	NA	$0_{(0)}$	NA	$0_{(0)}$	$0_{(0)}$
	121	$24_{(146)}$	2.5	$27_{(225)}$	1.5	$31_{(227)}$	2.4	$35_{(275)}$	1.5	$42_{(76)}$	$44_{(194)}$
	122	$21_{(127)}$	2.3	$19_{(160)}$	2.0	$29_{(209)}$	2.6	$26_{(203)}$	1.4	$26_{(48)}$	$26_{(115)}$
	211	$12_{(73)}$	1.6	$12_{(100)}$	1.2	$11_{(80)}$	2.3	$10_{(80)}$	2.0	$2_{(3)}$	$2_{(8)}$
	221	$1_{(8)}$	NA	$1_{(11)}$	NA	$4_{(28)}$	NA	$3_{(25)}$	NA	$14_{(26)}$	$14_{(62)}$
	222	$1_{(5)}$	NA	$0_{(1)}$	NA	$2_{(12)}$	NA	$1_{(2)}$	NA	$0_{(0)}$	$0_{(0)}$

FIGURE LEGENDS

Figure 1: Fish length at different life stages, back calculated from scale reading. The size of the drawings of fish on the figure are proportional to the measured length. Fish length and sea age distribution are compared between historical and contemporary samples for females and males separately. The length at first sea winter (immature fish) decreased from 47.2 to 41.7 cm (t=14.1, df=500, p<2.10^{-16}) for the females and from 45.4 to 41.0 cm (t=13.2, df=711, p<2.10^{-16}) for the males. During the same period, the length of the 1SW adult females remained stable, averaging 55 cm (t=0.89, df=37, p=0.37), whereas length increased marginally from 55.6 to 56.8 cm for the 1SW males (t=2.7, df=441, p=7.10^{-2}). For the 2SW adult fish however, average length declined from 79.4 to 75.3 cm for females (t=5.5, df=202, p=9.10^{-8}) and from 80.6 to 72.6 cm for males (t=5.4, df=85, p=4.10^{-7}). This trend for smaller size at age was also accompanied by a trend for delayed maturation with a strong decline of the frequency of the fish maturing after one winter at sea from 46% to 8% for the females (χ^2=120, df=1, p<2.10^{-16}), and from 75% to 44% for the females.

Figure 2: scan for SNP association with age at maturation. Females from the historical samples (**a**) display a particularly high association between age at maturation and chromosome 25 (p<1.10^{-25}) and chromosome 9 (p<1.10^{-10}). Males from the historical samples (**b**) also display a high association on chromosome 25 (p<1.10^{-30}). In the females of the contemporary samples (**c**), no locus was associated with the trait, whereas in the males from the same period (**d**) chromosome 25 was associated with age at maturation (p<1.10^{-7}) however not as strongly as in the historical samples.

589
590

Figure 3: mean sea age of the samples for each class of genotypes at *vgll3* and *six6*. In the females from the historical samples (**a**) *vgll3 and six6* had an additive effect on sea age (df=1 χ^2=24 p<2.10^{-16}) and (df=1 χ^2=7.4 p=8.10^{-9}). The males from the historical samples (**b**) displayed a significant dominance of the *vgll3-E allele* (df=1χ^2=12.8 p<2.10^{-16}), as well as an additive effect of six6 (df=1 χ^2=2.2, p=$6.3.10^{-4}$) which is dependent on the genetic background at *vgll3* as a result of epistasis (df=1 χ^2=10.3 p=0.001). The contemporary female samples (**c**) displayed a weak additive effect of *vgll3* (df=1 χ^2=1.0 p=$2.7.10^{-4}$), whereas contemporary males (**d**) displayed dominance of *vgll3*-E (df=1 χ^2=2.6 p=$6.9.10^{-4}$

1983–84
Females

47 cm

55 cm

79 cm

46
54

2013–16
Females

41 cm

55 cm

75 cm

92
8

Immature fish: first
winter at sea

Adult 1SW

Adult 2SW

Sea age distribution %

1983–84
Males

45 cm

55 cm

80 cm

75
25

2013–16
Males

41 cm

56 cm

72 cm

56
44

Immature fish: first
winter at sea

Adult 1SW

Adult 2SW

Sea age distribution %