1 **Overruled by nature: A plastic response to an ecological regime**

2 **shift disconnects a gene and its trait**

3 F. Besnier[1*], Ø. Skaala[1], V. Wennevik[1], F. Ayllon[1], K.R. Utne[1], P.T. Fjeldheim[1], K.

4 Andersen-Fjeldheim[1], S. Knutar[1], K.A. Glover[1,2]

5
6 [1] Institute of Marine Research, Bergen, Norway
7 [2] Institute of Biology, University of Bergen, Norway
8 *Corresponding author
9

Abstract

In Atlantic salmon, age at maturation is a life history trait ruled by a sex-specific trade-off between reproductive success and survival. Following an ecological regime shift in 2005, many North Atlantic salmon populations currently display smaller size at age and delayed age at maturation. However, whether this change reflects rapid evolution or plastic response is unknown. Some 1500 historical and contemporary salmon from river Etne (Western Norway) genotyped at 50k SNPs revealed three loci significantly associated with age at maturation. These included *vgll3* and *six6*, which collectively explained 36 to 50% of the age at maturation variation in the 1983-1984 period. Strikingly, the combined influence of these genes was nearly absent in all samples from 2013-2016, despite allelic frequencies at *vgll3* remaining unchanged. We conclude that the regime shift has led to the sudden bypassing of the influence of *vgll3* and *six6* on maturation through growth-driven plasticity.

Introduction

Understanding the mechanisms by which organisms adapt to their environments is a central question in biology (Losos 2000; Andrew *et al.* 2013). However, beyond the academic curiosity that motivates biologists to investigate how species evolved and adapted until now, accelerating climate change and ongoing habitat destruction catalyses a sense of urgency when considering the fate of many species in the future. As unprecedentedly fast environmental changes have occurred for many taxa during the last half century (Chevin *et al.* 2010; Dullinger *et al.* 2012; Pörtner *et al.* 2022), the current mechanisms of adaptation of many species may not couple with the pace at which environmental change is now occurring, leading thus to population declines and extinctions. Consequently, the mechanisms of adaptation and their potential to remain effective in a context of rapidly changing environment are becoming increasingly important to understand (García de Leániz *et al.* 2007; Candolin & Heuschele 2008; Kwan *et al.* 2008; Bandillo *et al.* 2017).

In the context of environmental change, genes associated with sexual conflicts (Parker 1974) are of particular interest. Sexual conflicts are associated with sex-specific selection and therefore expected to promote genetic diversity in the underlying genes (Rowe *et al.* 2018), thus potentially helping maintain standing genetic variation on important fitness traits within the population. An illustration of this occurs in anadromous Atlantic salmon (*Salmo salar* L.) where a single gene (*vgll3*) is strongly associated with the age at maturation, a highly adaptative trait subject to sexual conflict (Ayllon *et al.* 2015; Barson *et al.* 2015). In many species, age at maturation represents a trade-off, as late-maturation gives larger body sizes and thus higher reproductive success, but at the increased risk of dying before reproduction (Fleming & Einum 2011). In Atlantic salmon, the optimum value for this

trait differs between sexes, as males mature earlier and smaller whereas females

benefit from later maturation and a larger body size resulting in more and larger eggs

(Fleming & Einum 2011). This sexual conflict is in part resolved by the mediation of

vgll3 (Barson *et al.* 2015) through sex-specific dominance.

Studies of *vgll3* and age of maturation in Atlantic salmon have resulted in

inconstant estimations regarding the degree to which this gene influences the trait.

Several studies have demonstrated a very strong association in wild North European

populations (Ayllon *et al.* 2015; Barson *et al.* 2015; Czorlich *et al.* 2018; Jensen *et al.*

2022), but in stark contrast, little or no association has been observed in wild

populations in North America (Boulding *et al.* 2019; Mohamed *et al.* 2019).

Furthermore, conflicting observations have also been reported in domesticated

Norwegian farmed strains reared under aquaculture conditions (Ayllon *et al.* 2019;

Sinclair-Waters *et al.* 2020). This begs the question, why does the influence of *vgll3*

on age at maturation vary so greatly?

Salmonids have been exposed to a wide range of anthropological challenges

including habitat modifications since the Industrial Revolution (Forseth *et al.* 2017).

Furthermore, during the past two decades, most salmon populations in the North

Atlantic have shifted towards later maturation (Otero *et al.* 2012; Vollset *et al.* 2022)

and smaller size at age (Quinn *et al.* 2006; Bal *et al.* 2017; Todd *et al.* 2021; Vollset

et al. 2022). Although these changes are thought to be caused by a regime-shift in

oceanic conditions in 2005 (Vollset *et al.* 2022), whether they reflect rapid evolution

or plastic responses is unknown. The same trends towards later maturation have

also been documented in the salmon population inhabiting the river Etne on the west

coast of Norway, where the proportion of fish maturing after one year at sea dropped

from 63% in the period 1983-84 to 34% in 2018-19 (Harvey *et al.* 2022). The main

objective of the present study was to investigate whether the observed changes in

age at maturation were the result of phenotypic plasticity, or alternatively, evolution

in the gene(s) influencing this trait. To address this, we genotyped historical (early

1980´s – pre regime shift) and contemporary (mid 2010´s post regime shift) samples

with a 50k panel of genome-wide SNPs. This approach revealed the dissociation

between age at maturation and two loci, *vgll3* and *six6,* that explained 36 to 50% of

the variation in the 1983-1984 period, but only 7% in the most recent samples.

Material and Methods

Samples

This study is based on samples obtained from adult salmon captured in the river

Etne, in western Norway, 6o° N. This river is home to a salmon population of

typically 1000-2500 adults returning from the sea annually (Harvey *et al.* 2017). A

permanent trapping facility installed in the river has permitted sampling almost the

entire adult spawning population since 2014, which facilitated access to both an

extensive set of DNA samples, as well as to phenotypic and phenological data. For

the present study, 797 wild adult salmon captured in the 1983-84 angling season

were compared to 751 wild adult salmon captured in the upstream fish trap in the

period 2013 to 2016 (Besnier *et al.* 2022). For the contemporary samples (2013-

2016), an estimation of individual genetic admixture was computed as the proportion

of domestic ancestry in each individuals genome (see Besnier *et al.* (2022) for

details). In addition, a sample of 350 domesticated farmed salmon escapees that

were removed from the river in the period 1989-2012 (15 per year), were genotyped

with the same set of markers.

Genotyping and sex determination

All samples were genotyped on a ThermoFisher Axiom 57K single nucleotide polymorphism (SNP) array (NOFSAL03, 55735 markers) developed by Nofima (Norwegian institute for applied research in food aquaculture and fisheries) in collaboration with private aquaculture companies Mowi and SalmoBreed (Besnier *et al.* 2022). SNPs were checked following the "Best Practice Workflow" on the Affymetrix axiom analysis software (available at: https://www.thermofisher.com/no/en/home/technical-resources/software-downloads.html). SNPs with call rates lower than 0.97 and samples with call rates lower than 0.85 were discarded, whereas markers classified as "PolyHighResolution" (High resolution in both homozygous and heterozygous clusters) were conserved for further data analysis.

Fish were sexed by examining variants of the sdY gene (Yano *et al.* 2012; Eisbrenner *et al.* 2014); *i.e.,* males were identified based on the presence of exons 2 and 4. Samples were genotyped on an Applied Biosystems ABI 3730 Genetic Analyser, and genotypes were called using GeneMapper (Applied Biosystems, v. 4.0). The analysis of the sdY gene provides an accurate identification of sex, although, a very low percentage of fish identified as genetic males are phenotypic females due to carrying an inactive pseudo-copy of the sdY gene with both exon 2 and 4 (Ayllon *et al.* 2020).

Genome scan for loci associated with age at maturation

Age at maturation was modeled as a binary trait consisting of early maturing fish, *i.e.*, fish returning after one sea-winter (1SW), also known as grisling, *vs.* late maturing fish, *i.e.*, fish returning after two or more winters at sea (2$^+$SW). The

probability of maturing early was then modeled in a generalized linear model with logit link function:

$$\log\left(\frac{P_{(e)}}{1-P_{(e)}}\right) = Gi + sex + year + e \qquad \text{(Model 1)}$$

Where $P_{(e)}$ is the probability of early maturing, G_i is the SNP genotype at locus i, *sex* is a binary factor accounting for genetically determined sex, *year* is a factor accounting for the sampling year, and *e* a normally distributed vector of residuals. Significance of association between genotype and sea age was estimated by comparing the deviance of Model 1 with the deviance of the model without genetic effect. The difference was compared to a Chi-squared distribution with one degree of freedom. Model 1 was fitted at each SNP available on the dataset, and the obtained p-values were adjusted for multiple testing by following Bonferroni correction. All p-values given in the genome scan result section are corrected for multiple testing.

Haplotyping

Haplotypes were reconstructed using the Phase 2.1 software (Stephens *et al.* 2001) in the historical and contemporary data separately. Two loci on SSA9 and SSA25 were more specifically considered for haplotype reconstruction as they displayed high association with age at maturation, and associated genes *vgll3* and *six6* were previously described in the same two genomic regions (Ayllon *et al.* 2015; Barson *et al.* 2015; Czorlich *et al.* 2018). On SSA25, a haplotype window consisting of four polymorphic SNPs (AX-87309414, AX-172546510, AX-87309615, AX-87420691) was reconstructed in the region spanning from 28.65 to 28.66 Mb containing *vgll3* (https://www.ncbi.nlm.nih.gov/gene/106586514). On SSA09, a haplotype window consisting of three SNPs (AX-172546568, AX-88029383, AX-87668000), spanning

from 24.86-24.95Mb was reconstructed around the position of *six6*.

(https://www.ncbi.nlm.nih.gov/gene/106610974)

Statistical analyses

Genetic structure to age at maturation variation

Due to the sex specific dominance observed in *vgll3*, the genetic structure was

estimated for males and females separately. The probability of early maturation was

modeled as a response to additive and dominance genetic effects as follow:

$log \frac{P_{(e)}}{1-P_{(e)}} = a_i + d_i + year + e$ (Model 2)

Where $P_{(e)}$ is the probability of early maturing, a_i and d_i are respectively the additive

and dominance effects at locus *i*, *year* is a factor accounting for the sampling year,

and *e* a normally distributed vector of residuals. The model was fitted in a GLM with

logit link function in R (Team 2022), and the variance contribution was deduced from

the difference in model deviance between Model 2 and a model than only accounted

for capture year. Two loci interaction models were tested similarly by accounting for

the additional interaction parameters between loci $_i$ and $_j$. In contrast with the

genome scan, the test for significance if the different genetic parameters were not

corrected for multiple testing. All p-values given with the estimation of genetic effect

are nominal values.

Comparison in age at maturation

Potential differences in age at maturation between historical and contemporary

samples was tested by chi-squared test on a contingency table reporting the number

of observed 1, 2 and 3+ sea winter adults within historical and contemporary

samples.

Size at age

Individual body length was recorded for every fish passing the trap in the

contemporary sample, whereas the adult length of the 1983-1984 fish were

calculated from reading scales samples. In addition, the growth at first sea winter

was estimated for both historical and contemporary samples by calculating length at

first sea winter from reading scales. The difference in length between samples from

1983-1984 and from 2013-2016 was tested with a two-sided t-test, separately for

each sea-age and sex categories.

Potential role of admixture

The salmon population in the river Etne has been subject to introgression from

domesticated salmon escaping from commercial fish farms, with an average 24% of

genetic admixture in the contemporary population (Glover *et al.* 2013; Karlsson *et al.*

2016; Besnier *et al.* 2022). Individual admixture with domesticated salmon has

already been correlated with earlier adult maturation in this population (Besnier *et al.*

2022). Therefore, in order to account for the potential influence of admixture on the

temporal influence of loci on age at maturation in this population, we investigated

whether the change in genetic architecture of age at maturation could be linked to

genetic admixture.

A third model was fitted with the aim to evaluate a potential interaction

between admixture and *vgll3* or *six6* genotypes.

$$log \frac{P_{(e)}}{1-P_{(e)}} = a_i * admix + d_i * admix + year + e \qquad \text{(Model 3)}$$

Where *admix* is the individual genetic admixture as computed in Besnier et al

(Besnier *et al.* 2022) .

Results

Temporal changes in age and size at maturation

Marine growth was compared between the historical (1983-84) and contemporary (2013-16) samples. During this period, the length of the fish during the first winter at sea decreased significantly, as well the length of the two sea-winter (2SW) adults, while the length of the one sea-winter (1SW) adults remained stable (Table 1).

During the same period, we also observed a trend towards later maturation with a strong decline of the frequency of the fish maturing after one winter at sea from 46% to 8% for the females (χ^2=120, df=1, p<2.10^{-16}), and from 75% to 58% for the males (χ^2=25, df=1, p=4.10^{-7}) (Table 2).

Genome scans

The genome scan for age at maturation (Fig.1) identified three loci displaying a significant association in the historical samples, one on each of chromosomes SSA09, SSA24 and SSA25. The genomic regions identified in SSA25 and SSA09 overlapped with loci previously described as major contributors to the sea age variability in salmon: *vgll3* on SSA25 and *six6* located on SSA09. The degree of association between age at maturation and these genomic regions was investigated further in the historical and contemporary dataset.

vgll3 and _six6_ haplotypes

Haplotypes were reconstructed in the historical and contemporary samples separately, across the genomic regions in SSA25 and SSA9 that contained the *vgll3* and s*ix6* genes respectively. On SSA25, two haplotypes were predominant in the

historical samples with frequencies of 56% and 39% (Table 3) and respectively

associated with early and late sea age. The mean sea age was 1.18 for the

homozygous haplotype "1221", and 2.46 for the homozygous haplotype "2112",

which will thus be referred to as *vgll3*-E and *vgll3*-L alleles from hereon. In the

contemporary samples, the same two haplotypes were found in almost identical

frequencies (df=1, χ^2=0.85, p=0.35) to the historical samples (59% and 39% for

vgll3-E and *vgll3*-L respectively), indicating that no temporal change in haplotype

frequencies occurred at this locus during the three-decade period.

The mean age at maturation in each genotype class (Fig. 2) displayed a

strong association with an additive effect of *vgll3* on female sea age in the historical

data (Fig. 2.A, Table.S1), whereas association of *vgll3* almost disappeared in the

contemporary female data (Fig. 2.C, Table.S1). For the historical male data (Fig. 2B,

Table.S1), a strong *vgll3* association with a dominant *vgll3*-E allele was also

detected, whereas the genetic association was strongly reduced in the contemporary

data (Fig. 2.D, Table.S1).

On SSA09, four main haplotype sequences linked to the *six6* gene were

predominant in both historical and contemporary samples (Table 3). With a

respective sea age of 1.3 and 1.4 years for the homozygous fish in the historical

samples, both "111" and "211" haplotypes were assigned to the early variant of *six6,*

whereas haplotypes "121" and "122", with a respective mean sea age of 1.9 and 2.1

years, were assigned to the late variant. Following this, we observed an increase in

frequency of the *six6* late variant, from 44% in the historical samples to 60% in the

contemporary samples (df=1 χ^2=111, p<2.2.10^{-16}). The mean age at maturation in

each genotype class (Fig. 2) showed an additive effect of *six6* in the historical female

data (Fig. 2.A, Table.S2), and historical male data (Fig. 2.B Table.S2). In contrast,

we didn't observe any association between *six6* and sea age in the contemporary

data (Fig. 2.C,D Table.S2).

Cumulatively, *vgll3* and *six6* accounted for 50% of the model deviance in the

historical male dataset and 36% in the historical female dataset. In stark contrast, the

same loci only accounted for 7% and 3% of the model deviance in the contemporary

male and female datasets respectively.

Testing the interaction between these loci revealed significant departure from

additivity between *vgll3* and *six6* in the males of the historical samples (Table.S3).

This interaction is illustrated in Fig. 2.b where the *six6* genotype is only associated

with age at maturation for the samples that are homozygous for the *vgll3-L* allele. No

significant departure from additivity was observed in the historical female (Table.S3).

Genetic admixture

The salmon population inhabiting the river Etne has been subject to introgression

from domesticated farmed escapees. We therefore tested whether the observed

temporal change in genetic architecture of age at maturation could be linked to

admixture with farmed fish. When predicting age at maturation from the joined effect

of admixture and genotype, no interaction was detected between individual

admixture and *vgll3* genotype (χ^2=3.6, df=2, p=0.161), nor between individual

admixture and *six6* genotype (χ^2=2.7, df=2, p=0.24). We also checked the allelic

frequencies at the *six6* locus in the farmed samples to assess if the increase in *six6-*

L allele observed in the contemporary sample could be explained by admixture. With

17% *six6-E* and 68% *six6-L* allele, haplotype frequencies in the farmed samples

were more like the contemporary samples than the historical. However, the observed

genetic admixture in this population, which is estimated to circa 24%, could only

explain a 6% increment in frequency of *six6*-L, which is less than half of the increase

observed between historical and contemporary samples.

Discussion

We document the sudden dissociation between age at maturation in a population of

Atlantic salmon and the genotypes of three loci, including the previously identified

candidate genes *vgll3* and *six6* (Ayllon *et al.* 2015; Barson *et al.* 2015; Czorlich *et al.*

2018). This dissociation was observed in samples separated by a 30-year time

interval and was accompanied by an increase in the frequency of late maturing allele

at the *six6* locus, while no change was observed at the *vgll3* locus. The salmon

population in the river Etne has been subject to introgression from domesticated

salmon escaping from commercial fish farms, with an average 24% of genetic

admixture in the contemporary population (Glover *et al.* 2013; Karlsson *et al.* 2016;

Besnier *et al.* 2022). However, genetic admixture could not explain alone the change

in allelic frequencies observed at *six6,* nor the dissociation between the genes and

age at maturation.

The combined effect of *vgll3* and *six6* was not fully additive as we detected a

significant epistatic interaction in the historical samples of males. *Six6* was

previously identified as a candidate locus associated with age at maturation (Barson

et al. 2015), but little is known about the genetic effects of this locus. The work

presented here is the first to document a sex-specific interaction between *vgll3* and

six6.

In the river Etne, the frequency of late maturing fish was higher among

females than among males. This difference is assumed to be the result of adaptation

to sexual conflict where the advantage of maturing late is greater for females than for

males (Fleming & Einum 2011). With the *vgll3-E* allele being dominant in males only,

vgll3 is believed to play a key function in the resolution of sexual conflict in the

optimal age of maturation in salmon (Ayllon *et al.* 2015; Barson *et al.* 2015).

However, with the present results documenting the almost complete dissociation

between age at maturation and *vgll3* genotypes in parallel with a trend towards later

maturation strategy in both sexes simultaneously, the perennity of local adaptation in

the population inhabiting the river Etne can be questioned. Noteworthy, sex-specific

strategies of maturation seem to perdure in the population despite *vgll3*'s

contribution being almost completely eradicated in the contemporary samples. In

fact, the difference in early-maturation frequencies between males and females

increases in the contemporary dataset, strongly suggesting that other mechanisms

are acting instead of, or in parallel with *vgll3*, to maintain sex specific age at

maturation. This hypothesis seems consistent with the description of multiple loci

associated with age at maturation (Sinclair-Waters *et al.* 2020).

The absence of association between *vgll3* and age at maturation in multiple

North American populations (Boulding *et al.* 2019; Mohamed *et al.* 2019) already

suggested that *vgll3* is not the only regulator of sexual conflict for age of maturation

in Atlantic salmon. The present study confirms this observation and further shows

that the relative influence of *vgll3* is not stable in time. In the case of the population in

the river Etne, our data reveals that other unidentified factors also play a major role

in maintaining the sex-specific differences in maturation strategies when the link

between age at maturation and *vgll3* is disconnected.

The observed trend towards slower marine growth and later maturation in the

population inhabiting the river Etne has also been reported in many other salmon

populations in the North Atlantic (Quinn *et al.* 2006; Otero *et al.* 2012; Bal *et al.* 2017;

Todd *et al.* 2021; Vollset *et al.* 2022). While the precise triggers and mechanisms

underpinning to the development of sexual maturation are not yet fully understood in

Atlantic salmon (Mobley *et al.* 2021), slow growth and late maturation is consistent

with the hypothesis of size or perhaps growth-rate threshold as a determinant for

salmon maturation (Rowe *et al.* 1991; Simpson 1992; Taranger *et al.* 2010). Such an

energy-budget threshold might also be genetically regulated through the mediation of

vgll3, which has been shown to be linked with cell fat regulation (Halperin *et al.*

2013) or six6 which has been linked to stomach fullness and prey composition

(Aykanat *et al.* 2020). Following this hypothesis, slow growth due to environmental

conditions, such as lack of prey would lead to later maturation as a higher proportion

of fish would fail to reach the weight (or growth-rate) threshold for maturation after

one single year at sea.

Observations suggest important changes in the environmental conditions met

by salmon migrating from the river Etne in the period 1980-2010 (Vollset *et al.* 2022).

The marine migration pattern for salmon originating from the river Etne is not fully

known, however, the first phase is probably a northward migration through the

southern Norwegian Sea (Gilbey *et al.* 2021) where changes in the oceanographic

conditions have been reported in the last years. For example, water temperature

increased by nearly 1˚C at 50-200 m depth from early 1980's until 2021, mainly due

to warmer water masses flowing into the southern Norwegian Sea (Skagseth & Mork

2012; ICES 2021a). The early 1980's are also considered as the end of "The great

salinity anomaly", which was a period with large inflow of cold and fresh Arctic water

into the Norwegian Sea. The large proportion of Arctic water entering the Norwegian

Sea is correlated to increased productivity (Skagseth *et al.* 2022), and to improved

feeding conditions for post-smolts in the region (Utne *et al.* 2022). Therefore, salmon

returning in 1983 and 1984 had probably been feeding in a very productive sea

during the initial post-smolt phase, whereas salmon returning to rivers in 2013-2016

had been feeding in a warm and saline Norwegian Sea. During this later period,

observed stomach fullness and condition factor for post-smolt sampled in the

Norwegian Sea were low (Utne *et al.* 2021a). In addition, the potential interspecific

competition with other pelagic fish for prey (Utne *et al.* 2021b) was low in the early

1980's as the total biomass of pelagic fish feeding in the Norwegian Sea in 1982-

1983 was around 1/3 of the total biomass in the period 2013-2016 (ICES 2008,

2021b) when Norwegian Spring-spawning herring had not yet recovered from the

collapse in the late 1960ies, and NEA-mackerel and blue whiting stock biomasses

were at low levels (ICES 2008). Many observations seem to confirm that

environmental conditions were less favorable for salmon growth in the last decade

than in the early 1980's. It is thus conceivable that the changes observed in the river

Etne are caused by environmental perturbations occurring over a large region and

affecting salmon populations as well as other organisms.

Under commercial aquaculture conditions, farmed Atlantic salmon typically

display early sexual maturation (Taranger *et al.* 2010), and importantly, the relative

influence of *vgll3* on age at maturation appears to be largely bypassed (Ayllon *et al.*

2019). Yet again, this response is sex-specific as a study conducted under

aquaculture conditions (Ayllon *et al.* 2019) reported that *vgll3* did not show any

correlation with age at maturation in females while displaying only weak association

in males (Ayllon *et al.* 2019). The commercial farm strain used in the aforementioned

study, known as Mowi, stemmed from wild Norwegian salmon populations in which

vgll3 was identified as a candidate gene for influencing age at maturation (Ayllon *et*

al. 2015; Barson *et al.* 2015). The authors concluded that high calorie feed intake

combined with artificial light and temperature regimes as well as potential genetic or

epigenetic components, may alter the impact of *vgll3* on age at maturation (Ayllon *et*

al. 2019). Anecdotally, it is also worth noting that despite multiple generations of

directional selection against early maturing fish in the domesticated farmed salmon,

the genetic variability of *vgll3* remains high in the Mowi strain (Ayllon *et al.* 2019). It is

thus possible that a rapid change in environmental conditions, such as feed intake

and therefore growth-rate, may bypass the effect of *vgll3* without letting selection

(artificial selection in this case) alter the allelic frequencies of the gene.

We thus can hypothesize a two-threshold model where the effect of *vgll3* on

age at maturation is bypassed when abundance of feed resources is very high, as in

farming conditions, or when feed resources are very low and fish need to spend

more time at sea to acquire the necessary energy-reserves for maturation and

reproduction. When resource availability falls between both thresholds, *vgll3* may

play a more important role in determining age at maturation. Inversely if feed

resources exceed the high threshold, or do not reach the low one, *vgll3* is bypassed

by environmental conditions, and therefore age at maturation is determined by a

combination of other genetic and environmental factors.

In the face of changing environmental conditions, one may expect the allelic

frequencies at the associated loci to change in response to the newly induced

selection pressure (Czorlich *et al.* 2018; Jensen *et al.* 2022). This is the case for *six6*

where the temporal trend towards later maturation is accompanied by an increase of

the *six6*-L allele frequency. This shift in allelic frequencies is likely due to positive

selection on the *six6-L* allele, alone, or in conjunction with admixture from

domesticated salmon where *six6*-L was found in higher frequency. In contrast, the

vgll3 allelic frequencies remained stable despite major changes in age at maturation.

This result seems to indicate a change in environmental conditions, creating a

situation where the influence of *vgll3* was effectively bypassed before natural

selection had time to operate, whereas selection had the time to modify the allelic

frequencies at the *six6* locus before the influence of this gene was also bypassed.

This hypothesis is strongly supported by the observations of sudden change in age

at maturation in many Atlantic salmon populations inhabiting Northeast Atlantic rivers

(Vollset *et al.* 2022), including the population in the river Etne (Harvey *et al.* 2022),

where a highly distinct and sudden drop in early marine growth was reported in 2005

and subsequently, referred to as an ecological regime shift in the northeast Atlantic

ocean (Vollset *et al.* 2022).

As this, and other salmon population in the Northeast Atlantic were subjected

to the same ecological regime shift, leading to consistently reduced marine growth

rates and increased age at maturation, we conclude that growth-driven plasticity has

almost completely bypassed the combined influence of *vgll3* and *six6* within on age

at maturation, and furthermore, on resolving the sexual conflict for this trait in Atlantic

salmon. Together with the interaction between *vgll3* and *six6* described in the

historical data, the dissociation between genes and age at maturation represents an

original finding that changes our understanding of the genetic architecture of age at

maturation in Atlantic salmon.

Acknowledgments

This work was financed by the Norwegian Ministry of Trade, Industry and Fisheries.

This authority paid no part in the study design nor interpretation of results. We would

like to acknowledge the research institute Nina, and anglers, for donating some of

the farmed salmon samples used in this study. We would like to acknowledge the

river Etne owners association for continued collaboration and allowing capture and

handling of fish and IMR staff involved in monitoring the population in the trap. We

would like to thank María Quintela for insightful comments on the earlier drafts of the

manuscript, and Emily K. Glover for drawing the salmons used in Figure 1.

429 **Data availability**

430   Data for this study are available from the IMR.brage.unit.no public repository.

431   Link to data: https://imr.brage.unit.no/

References

1.
Andrew, R.L., Bernatchez, L., Bonin, A., Buerkle, C.A., Carstens, B.C., Emerson, B.C. *et al.* (2013). A road map for molecular ecology. *Mol. Ecol.*, 22, 2605-2626.

2.
Aykanat, T., Rasmussen, M., Ozerov, M., Niemela, E., Paulin, L., Vaha, J.P. *et al.* (2020). Life-history genomic regions explain differences in Atlantic salmon marine diet specialization. *J Anim Ecol*, 89, 2677-2691.

3.
Ayllon, F., Kjaerner-Semb, E., Furmanek, T., Wennevik, V., Solberg, M.F., Dahle, G. *et al.* (2015). The *vgll3* locus controls age at maturity in wild and domesticated Atlantic salmon (*Salmo salar* L.) males. *PLoS Genet.*, 11, e1005628.

4.
Ayllon, F., Solberg, M.F., Besnier, F., Fjelldal, P.G., Hansen, T.J., Wargelius, A. *et al.* (2020). Autosomal sdY pseudogenes explain discordances between phenotypic sex and DNA marker for sex identification in Atlantic salmon. *Front. Genet.*, 11, 544207.

5.
Ayllon, F., Solberg, M.F., Glover, K.A., Mohammadi, F., Kjaerner-Semb, E., Fjelldal, P.G. *et al.* (2019). The influence of *vgll3* genotypes on sea age at maturity is altered in farmed mowi strain Atlantic salmon. *BMC Genet.*, 20, 44.

6.
Bal, G., Montorio, L., Rivot, E., Prevost, E., Bagliniere, J.L. & Nevoux, M. (2017). Evidence for long-term change in length, mass and migration phenology of anadromous spawners in French Atlantic salmon *Salmo salar*. *J. Fish Biol.*, 90, 2375-2393.

7.
Bandillo, N.B., Anderson, J.E., Kantar, M.B., Stupar, R.M., Specht, J.E., Graef, G.L. *et al.* (2017). Dissecting the genetic basis of local adaptation in soybean. *Sci. Rep.*, 7, 17195.

8.
Barson, N.J., Aykanat, T., Hindar, K., Baranski, M., Bolstad, G.H., Fiske, P. *et al.* (2015). Sex-dependent dominance at a single locus maintains variation in age at maturity in salmon. *Nature*, 528, 405-408.

9.
Besnier, F., Ayllon, F., Skaala, Ø., Solberg, M.F., Fjeldheim, P.T., Anderson, K.C. *et al.* (2022). Introgression of domesticated salmon changes life history and phenology of a wild salmon population. *Evol. Appl.*, 15, 853-864.

10.
Boulding, E.G., Ang, K.P., Elliott, J.A.K., Powell, F. & Schaeffer, L.R. (2019). Differences in genetic architecture between continents at a major locus previously associated with sea age at sexual maturity in European Atlantic salmon. *Aquac.*, 500, 670-678.

11.
Candolin, U. & Heuschele, J. (2008). Is sexual selection beneficial during adaptation to environmental change? *Trends Ecol. Evol.*, 23, 446-452.

12.
Chevin, L.M., Lande, R. & Mace, G.M. (2010). Adaptation, plasticity, and extinction in a changing environment: towards a predictive theory. *PLoS Biol.*, 8, e1000357.

13.

Czorlich, Y., Aykanat, T., Erkinaro, J., Orell, P. & Primmer, C.R. (2018). Rapid sex-specific

evolution of age at maturity is shaped by genetic architecture in Atlantic salmon.

Nat. Ecol. Evol., 2, 1800-1807.

14.

Dullinger, S., Gattringer, A., Thuiller, W., Moser, D., Zimmermann, N.E., Guisan, A. *et al.*

(2012). Extinction debt of high-mountain plants under twenty-first-century climate

change. *Nat. Clim. Change*, 2, 619-622.

15.

Eisbrenner, W.D., Botwright, N., Cook, M., Davidson, E.A., Dominik, S., Elliott, N.G. *et al.*

(2014). Evidence for multiple sex-determining loci in Tasmanian Atlantic salmon

(*Salmo salar*). *Heredity*, 113, 86-92.

16.

Fleming, I.A. & Einum, S. (2011). Reproductive Ecology: A Tale of Two Sexes In: *Atlantic*

Salmon Ecology (eds. Aas, Ø, Einum, S, Klemetsen, A & Skurdal, J). Blackwell

Publishing, pp. 33-65.

17.

Forseth, T., Barlaup, B.T., Finstad, B., Fiske, P., Gjøsæter, H., Falkegård, M. *et al.* (2017). The

major threats to Atlantic salmon in Norway. *ICES J. Mar. Sci.*, 74, 1496-1513.

18.

García de Leániz, C., Fleming, I.A., Einum, S., Verspoor, E., Jordan, W.C., Consuegra, S. *et al.*

(2007). A critical review of adaptive genetic variation in Atlantic salmon: implications

for conservation. *Biol. Rev.*, 82, 173-211.

19.

Gilbey, J., Utne, K.R., Wennevik, V., Beck, A.C., Kausrud, K., Hindar, K. *et al.* (2021). The early

marine distribution of Atlantic salmon in the North-east Atlantic: A genetically

informed stock-specific synthesis. *Fish Fish.*, 22, 1274-1306.

20.

Glover, K.A., Pertoldi, C., Besnier, F., Wennevik, V., Kent, M. & Skaala, Ø. (2013). Atlantic

salmon populations invaded by farmed escapees: quantifying genetic introgression

with a Bayesian approach and SNPs. *BMC Genet.*, 14, 74.

21.

Halperin, D.S., Pan, C., Lusis, A.J. & Tontonoz, P. (2013). Vestigial-like 3 is an inhibitor of

adipocyte differentiation. *J. Lipid Res.*, 54, 473-481.

22.

Harvey, A.C., Skaala, O., Borgstrom, R., Fjeldheim, P.T., Andersen, K.C., Rong Utne, K. *et al.*

(2022). Time series covering up to four decades reveals major changes and drivers of

marine growth and proportion of repeat spawners in an Atlantic salmon population.

Ecol. Evol., 12, e8780.

23.

Harvey, A.C., Tang, Y., Wennevik, V., Skaala, O. & Glover, K.A. (2017). Timing is everything:

Fishing-season placement may represent the most important angling-induced

evolutionary pressure on Atlantic salmon populations. *Ecol. Evol.*, 7, 7490-7502.

24.

ICES (2008). Report of the Working Group on Widely Distributed Stocks (WGWIDE).

25.

ICES (2021a). ICES Working Group on the Integrated Assessments of the Norwegian Sea
(WGINOR; outputs from 2020 meeting). 3. 114 pp.
26.
ICES (2021b). Working Group on Widely Distributed Stocks (WGWIDE). 3:95. 874 pp.
27.
Jensen, A.J., Hagen, I.J., Czorlich, Y., Bolstad, G.H., Bremset, G., Finstad, B. *et al.* (2022).
Large-effect loci mediate rapid adaptation of salmon body size after river regulation.
Proc. Natl. Acad. Sci. U. S. A., 119, e2207634119.
28.
Karlsson, S., Diserud, O.H., Fiske, P. & Hindar, K. (2016). Widespread genetic introgression of
escaped farmed Atlantic salmon in wild salmon populations. *ICES J. Mar. Sci.*, 73,
2488-2498.
29.
Kwan, L., Bedhomme, S., Prasad, N.G. & Chippindale, A.K. (2008). Sexual conflict and
environmental change: trade-offs within and between the sexes during the evolution
of desiccation resistance. *J. Genet.*, 87, 383–394.
30.
Losos, J.B. (2000). Ecological character displacement and the study of adaptation. *PNAS*, 97,
5693-5695.
31.
Mobley, K.B., Aykanat, T., Czorlich, Y., House, A., Kurko, J., Miettinen, A. *et al.* (2021).
Maturation in Atlantic salmon (Salmo salar, Salmonidae): a synthesis of ecological,
genetic, and molecular processes. *Reviews in Fish Biology and Fisheries*, 31, 523-571.
32.
Mohamed, A.R., Verbyla, K.L., Al-Mamun, H.A., McWilliam, S., Evans, B., King, H. *et al.*
(2019). Polygenic and sex specific architecture for two maturation traits in farmed
Atlantic salmon. *BMC Genom.*, 20, 139.
33.
Otero, J., Jensen, A.J., L'Abee-Lund, J.H., Stenseth, N.C., Storvik, G.O. & Vollestad, L.A.
(2012). Contemporary ocean warming and freshwater conditions are related to later
sea age at maturity in Atlantic salmon spawning in Norwegian rivers. *Ecol. Evol.*, 2,
2192-2203.
34.
Parker, G.A. (1974). Sexual selection and sexual conflict. In: *Sexual Selection and*
Reproductive Competition in Insects (eds. BLUM, NA & BLUM, MS). Academic Press,
Cambridge, pp. 123-166.
35.
Pörtner, H.O., Roberts, D.C., Tignor, M., Poloczanska, E.S., Mintenbeck, K., Alegría, A. *et al.*
(2022). IPCC, 2022: Climate Change 2022: Impacts, Adaptation, and Vulnerability.
Contribution of Working Group II to the Sixth Assessment Report of the
Intergovernmental Panel on Climate Change.
36.
Quinn, T.P., McGinnity, P. & Cross, T.F. (2006). Long-term declines in body size and shifts in
run timing of Atlantic salmon in Ireland. *J. Fish Biol.*, 68, 1713-1730.
37.
Rowe, D.K., Thorpe, J.E. & Shanks, A.M. (1991). Role of fat stores in the maturation of male
Atlantic salmon (*Salmo salar*) parr. *Can. J. Fish. Aquat. Sci.*, 48, 405-413.

38.
Rowe, L., Chenoweth, S.F. & Agrawal, A.F. (2018). The genomics of sexual conflict. *Am. Nat.*,
192, 274-286.
39.
Simpson, A.L. (1992). Differences in body size and lipid reserves between maturing and
nonmaturing Atlantic salmon parr, Salmo salar L. *Can. J. Zool.*, 70.
40.
Sinclair-Waters, M., Odegard, J., Korsvoll, S.A., Moen, T., Lien, S., Primmer, C.R. *et al.* (2020).
Beyond large-effect loci: large-scale GWAS reveals a mixed large-effect and polygenic
architecture for age at maturity of Atlantic salmon. *Genet Sel Evol*, 52, 9.
41.
Skagseth, Ø., Broms, C., Gundersen, K., Hátún, H., Kristiansen, I., Larsen, K.M.H. *et al.* (2022).
Arctic and Atlantic waters in the Norwegian Basin, between year variability and
potential ecosystem implications. *Front. Mar. Sci.*, 9.
42.
Skagseth, Ø. & Mork, K.A. (2012). Heat content in the Norwegian Sea, 1995–2010. *ICES J.
Mar. Sci.*, 69, 826-832.
43.
Stephens, M., Smith, E.P. & Donnelly, P. (2001). A new statistical method for haplotype
reconstruction from population data. *Am. J. Hum. Genet.*, 68, 978–989.
44.
Taranger, G.L., Carrillo, M., Schulz, R.W., Fontaine, P., Zanuy, S., Felip, A. *et al.* (2010).
Control of puberty in farmed fish. *Gen. Comp. Endocrinol.*, 165, 483-515.
45.
Team, R.C. (2022). R: A language and environment for statistical computing. In: *R
Foundation for Statistical Computing* Vienna, Austria.
46.
Todd, C.D., Hanson, N.N., Boehme, L., Revie, C.W. & Marques, A.R. (2021). Variation in the
post-smolt growth pattern of wild one sea-winter salmon (*Salmo salar* L.), and its
linkage to surface warming in the eastern North Atlantic Ocean. *J. Fish Biol.*, 98, 6-16.
47.
Utne, K.R., Díaz Pauli, B., Haugland, M., Jacobsen, J.A., Maoileidigh, N., Melle, W. *et al.*
(2021a). Poor feeding opportunities and reduced condition factor for salmon post-
smolts in the Northeast Atlantic Ocean. *ICES J. Mar. Sci.*, 78, 2844-2857.
48.
Utne, K.R., Skagseth, Ø., Wennevik, V., Broms, C.T., Melle, W. & Thorstad, E.B. (2022).
Impacts of a changing ecosystem on the feeding and feeding conditions for Atlantic
salmon during the first months at sea. *Front. Mar. Sci.*, 9.
49.
Utne, K.R., Thomas, K., Jacobsen, J.A., Fall, J., Maoiléidigh, N.Ó., Broms, C.T. *et al.* (2021b).
Feeding interactions between Atlantic salmon (*Salmo salar*) postsmolts and other
planktivorous fish in the Northeast Atlantic. *Can. J. Fish. Aquat. Sci.*, 78, 255-268.
50.
Vollset, K.W., Urdal, K., Utne, K.R., Thorstad, E.B., Sægrov, H., Raunsgard, A. *et al.* (2022).
Ecological regime shift in the Northeast Atlantic Ocean revealed from the
unprecedented reduction in marine growth of Atlantic salmon. *Sci. Adv.*, 8.
51.

Yano, A., Guyomard, R., Nicol, B., Jouanno, E., Quillet, E., Klopp, C. *et al.* (2012). An immune-
related gene evolved into the master sex-determining gene in rainbow trout,
Oncorhynchus mykiss. *Curr. Biol.*, 22, 1423-1428.
623
624
625

Competing interests

629   The authors declare no competing interests.

630 **TABLES**

632 **Table 1.** Comparison of fish length between historical and contemporary samples

633 during first winter at sea (immature fish), adult maturing after one sea winter (1SW),

634 and adult maturing after 2 winters at sea (2SW).

	phenotype	Mean value (cm) 1983-1984	Mean value (cm) 2013-2016	df	t	p
Females	First sea winter length	47.2	41.7	500	14	<0.001
	1SW adult length	55.1	55.0	37	0.89	0.37
	2SW adult length	79.4	75.3	202	5.50	<0.001
Males	First sea winter length	45.4	41.0	711	13.2	<0.001
	1SW adult length	55.6	56.8	441	2.7	0.007
	2SW adult length	80.6	72.6	85	5.4	<0.001

639 **Table 2.** Percentage and numbers (in brackets) of fish displaying 1-2-3$^+$ sea winter

640 (SW) of age at maturation, by sex, in the historical (H – 1983/84, N = 797) and

641 contemporary (C – 2013-2016, N = 751) samples.

	1SW % (N)	2SW % (N)	3$^+$SW % (N)
Females_H	46 (141)	37 (114)	17 (51)
Females_C	8 (30)	63 (227)	28 (102)
Males_H	75 (311)	13 (56)	12 (48)
Males_C	58 (227)	33 (127)	9 (36)

642
643

Table 3. Observed occurrences of each haplotype per sex and associated Mean Sea Age (M.S.A) in the historical, contemporary

and farmed samples. (M.S.A is calculated as mean sea age of the samples that are homozygous for the haplotype.)

		Historical				Contemporary				Farmed	
	Haplotypes	**F**		**M**		**F**		**M**		**F**	**M**
		%(N)	**M.S.A**	**%(N)**	**M.S.A**	**%(N)**	**M.S.A**	**%(N)**	**M.S.A**	**%(N)**	**%(N)**
vgll3	1221	$59_{(364)}$	1.2	$54_{(455)}$	1.1	$61_{(438)}$	2.2	$58_{(454)}$	1.4	$53_{(93)}$	$63_{(264)}$
	1222	$0_{(0)}$	NA	$0_{(1)}$	NA	$0_{(0)}$	NA	$0_{(1)}$	NA	$0_{(0)}$	$0_{(0)}$
	1111	$1.5_{(9)}$	NA	$2_{(17)}$	NA	$0_{(0)}$	NA	$0_{(0)}$	NA	$0_{(0)}$	$0_{(1)}$
	1112	$0_{(1)}$	NA	$0_{(0)}$	NA	$0_{(0)}$	NA	$0_{(1)}$	NA	$0_{(0)}$	$0_{(1)}$
	2221	$1_{(9)}$	NA	$2_{(18)}$	NA	$1_{(9)}$	NA	$1(13)$	NA	$3_{(5)}$	$5_{(24)}$
	2222	$0_{(0)}$	NA	$0.5_{(3)}$	NA	$0_{(3)}$	NA	$0_{(0)}$	NA	$0_{(0)}$	$0_{(0)}$
	2111	$1.5_{(5)}$	NA	$0.5_{(4)}$	NA	$0_{(0)}$	NA	$1_{(0)}$	NA	$0_{(0)}$	$0_{(1)}$
	2112	$37_{(230)}$	2.5	$41_{(344)}$	2.5	$38_{(272)}$	2.5	$40_{(309)}$	2.2	$44_{(76)}$	$31_{(128)}$
six6	111	$41_{(256)}$	1.4	$40_{(337)}$	1.2	$23_{(166)}$	2.3	$25_{(195)}$	1.6	$16_{(29)}$	$14_{(65)}$
	112	$0_{(3)}$	NA	$1_{(8)}$	NA	$0_{(0)}$	NA	$0_{(0)}$	NA	$0_{(0)}$	$0_{(0)}$
	121	$24_{(146)}$	2.5	$27_{(225)}$	1.5	$31_{(227)}$	2.4	$35_{(275)}$	1.5	$42_{(76)}$	$44_{(194)}$
	122	$21_{(127)}$	2.3	$19_{(160)}$	2.0	$29_{(209)}$	2.6	$26_{(203)}$	1.4	$26_{(48)}$	$26_{(115)}$
	211	$12_{(73)}$	1.6	$12_{(100)}$	1.2	$11_{(80)}$	2.3	$10_{(80)}$	2.0	$2_{(3)}$	$2_{(8)}$
	221	$1_{(8)}$	NA	$1_{(11)}$	NA	$4_{(28)}$	NA	$3_{(25)}$	NA	$14_{(26)}$	$14_{(62)}$
	222	$1_{(5)}$	NA	$0_{(1)}$	NA	$2_{(12)}$	NA	$1_{(2)}$	NA	$0_{(0)}$	$0_{(0)}$

648
649
FIGURE LEGENDS

Figure 1: **Scan for SNP association with age at maturation.** Red dots

represent SNPs with significant association with sea age (p<0.01 after

correction for multiple tests). Females from 1983-84 (**A**), males from

1983-84 (**B**), females from 2013-16 (**C**) and males from 2013-16 (**D**).

Figure 2: **Mean sea age of the samples for each class of genotypes at**

***vgll3* and *six6*.** Females from 1983-84 (**A**), males from 1983-84 (**B**),

females from 2013-16 (**C**) and males from 2013-16 (**D**).

661
662
663