Title: A new gene encoding a cytosolic glutamine synthetase in pine is linked to developing tissues

Authors: José Miguel Valderrama-Martín[1,2], Francisco Ortigosa[1], Juan Carlos Aledo[2], Concepción Ávila[1], Francisco M. Cánovas[1], Rafael A. Cañas[2]*.

Adresses: [1]Grupo de Biología Molecular y Biotecnología de Plantas, Departamento de Biología Molecular y Bioquímica, Universidad de Málaga, Campus Universitario de Teatinos, 29071-Málaga, Spain
[2]Integrative Molecular Biology Lab, Universidad de Málaga, Campus Universitario de Teatinos, 29071-Málaga, Spain

Correspondence: *Rafael A. Cañas, Departamento de Biología Molecular y Bioquímica, Facultad de Ciencias, Universidad de Málaga, Campus Universitario de Teatinos s/n, E-29071 Málaga, Spain. e-mail: rcanas@uma.es Phone numer: +34-925-131673 Fax number: +34-952-132041.

Authors e-mails: José Miguel Valderrama-Martín (jmvalderrama@uma.es); Francisco Ortigosa (fortigosa@uma.es); Juan Carlos Aledo (caledo@uma.es); Concepción Ávila (cavila@uma.es); Francisco M. Cánovas (canovas@uma.es); Rafael A. Cañas (rcanas@uma.es).

ORCID IDs: JMVM (0000-0003-1350-899X); FO (0000-0002-7517-7594); JCA (0000-0002-3497-9945); CA (0000-0001-8817-7529); FMC (0000-0002-4914-2558); and RAC (0000-0001-9727-5585)

SUMMARY

The enzyme glutamine synthetase (EC 6.3.1.2) is mainly responsible for the incorporation

of inorganic nitrogen into organic molecules in plants. In the present work, a new pine *GS1*

(*PpGS1b.2*) gene was identified, showing a high sequence identity with the *GS1b.1* gene

previously characterized in conifers. Phylogenetic analysis revealed that the presence of

PpGS1b.2 is restricted to the genera *Pinus* and *Picea* and is not found in other conifers.

Gene expression data suggest a putative role of *PpGS1b.2* in plant development, similar to

other *GS1b* genes from angiosperms, suggesting evolutionary convergence. The

characterization of GS1b.1 and GS1b.2 at the structural, physicochemical, and kinetic

levels has shown differences even though they have high sequence homology. Alterations

in the kinetic characteristics produced by the site-directed mutagenesis approach carried out

in this work strongly suggest an implication of amino acids at positions 264 and 267 in the

active center of pine GS1b.1 and GS1b.2. Therefore, the amino acid differences between

GS1b.1 and GS1b.2 would support the functioning of both enzymes to meet distinct plant

needs.

KEYWORDS: Biochemistry, development, glutamine synthetase, kinetic parameters,

nitrogen metabolism, physicochemical properties, conifer, *Pinus pinaster*

INTRODUCTION

Nitrogen (N) is an essential element, a constituent of the main biomolecules and a limiting factor for plant growth (Hirel and Krapp, 2021). N is assimilated from ammonium into organic molecules by the glutamine synthetase (GS, EC 6.3.1.2)/glutamate synthase (GOGAT, EC 1.4.7.1) cycle. Ammonium is first incorporated into glutamate to form glutamine in an ATP-dependent reaction catalyzed by the GS enzyme (Heldt and Piechulla, 2011), and then this glutamine together with 2-oxoglutarate is used to produce two glutamate molecules by the GOGAT enzyme (Bernard and Habash, 2009). Studies have shown that up to 95% of ammonium is assimilated via the GS/GOGAT cycle (Lea *et al*., 1999) for the formation of glutamine and glutamate, which, in turn, will be used to produce all N-containing biomolecules in the plant (Forde and Lea, 2007; Bernard and Habash, 2009).

The GS enzyme has been widely studied in plants since it is directly responsible for the incorporation of inorganic N into organic molecules. Recently, three different lineages of *GS* genes have been identified in seed plants: *GS1a* and *GS1b* encode cytosolic enzymes, and *GS2* encodes a plastid-located enzyme (Valderrama-Martín *et al*., 2022). The three *GS* gene lineages are present in cycads and *Ginkgo biloba*, as well as basal angiosperms. Nevertheless, no GS2 genes have been found in other gymnosperms, such as conifers and gnetales, and no *GS1a* genes have been found in modern angiosperms, including monocot and eudicotyledon species (Valderrama-Martín *et al*., 2022). In general, GS1b is encoded by a small multigene family, while GS1a and GS2 are usually encoded by a single nuclear gene (James *et al*., 2018; Valderrama-Martín *et al*., 2022).

GS2 and GS1a are associated with photosynthetic organs (Blackwell *et al.*, 1987; Ávila *et al.*, 2001), and their expression is regulated by light conditions (Cantón *et al.*, 1999; Gómez-Maldonado *et al.*, 2004a; Valderrama-Martín *et al.*, 2022). Indeed, GS2 and GS1a are considered to play a fundamental role in the assimilation of the ammonium released during photorespiration and nitrate photoassimilation processes (Wallsgrove *et al.*, 1987; Blackwell *et al.*, 1987; Cantón *et al.*, 1999; Tegeder and Masclaux-Daubresse, 2017). In this sense, new evidence suggests that the *GS2* gene may arose through a gene duplication

from a *GS1a* gene in a common ancestor of cycads, ginkgo, and angiosperms (Valderrama-Martín *et al.*, 2022).

GS1b corresponds to the GS1 isoenzyme traditionally studied in model angiosperms. Although this lineage is represented by a unique gene in most of the gymnosperms, in ginkgo and angiosperms, GS1b is represented by a small multigenic family. These genes have different expression patterns depending on the organ and physiological conditions accounting for their different functions (Hirel and Krapp, 2021). These enzymes have been described as a key components of plant nitrogen use efficiency, with essential roles in processes such as senescence (Thomsen *et al.*, 2014), amino acid catabolism, primary assimilation, and different stress responses (Bernard and Habash, 2009). The different genes of this lineage are differentially regulated by developmental state, tissue, nutritional status, and external stimuli (Thomsen et al. 2014; Hirel and Krapp, 2021). Finally, several studies have focused on the enzymatic characterization of GS from angiosperms and gymnosperms (Sakakibara *et al.*, 1996; de la Torre *et al.*, 2002; Ishiyama *et al.*, 2004a; Ishiyama *et al.*, 2004b; Ishiyama *et al.*, 2006; Yadav, 2009; Zhao *et al.*, 2014; Castro-Rodríguez *et al.*, 2015) to define a more accurate role landscape for the different GS isoforms.

Some GS1b isoforms are directly related to developmental processes and have been associated with grain yield in crops. *AtGS1.1* and *AtGS1.2* from *Arabidopsis thaliana* are involved in seed production and germination (Guan *et al.*, 2015). *AtGS1.1* has also been described to be involved in root development during seed germination and *AtGS1.2* plays a role in rosette development (Lothier *et al.*, 2011; Guan *et al.*, 2015). Indeed, a recent study over of *AtGS1.1*, *AtGS1.2* and *AtGS1.3 Arabidopsis* mutants suggested synergistic roles for these genes in plant growth and development (Ji *et al.*, 2019). In cereals, enzymes of this GS lineage are involved in seed yield and plant development, such as *GS1;3* from *Oryza sativa* and *Hordeum vulgare*, which play roles in seed maturation and germination (Goodall *et al.*, 2013; Fujita *et al.*, 2022). Thus, overexpressing lines of *HvGS1.1* showed an improvement in grain yield (Gao *et al.*, 2019). Rice mutants lacking the *OsGS1;1* gene presented reduced grain filling and growth (Tabuchi *et al.*, 2005), although the same phenotype was present in rice lines overexpressing *OsGS1;1* (Bao *et al.*, 2014). In addition, rice lines grown in culture chambers and overexpressing *OsGS1;1* presented an increase in

spikelet yield. Rice mutants for *OsGS1b;2* also presented a depletion in the number of tillers (Funayama *et al*., 2013), and *Sorghum bicolor* lines overexpressing *GS1* genes exhibited the opposite phenotype (Urriola and Rathore, 2015). Studies in *Zea mays* using mutant lines for *ZmGS1b.3* and *ZmGS1b*.4 have shown the roles of these genes in kernel number and size, respectively (Martin *et al*., 2006). Transgenic lines of *Phaseolus vulgaris* overexpressing GS1 also showed earlier flower and seed development, while overexpressing GS1 lines of wheat showed an increase in grain weight (Habash *et al*., 2001). Moreover, a recent study on wheat indicated that *TaGS1.1* and *TaGS1.3* are mainly expressed in embryos and grain transport tissues, where these isoforms synergistically carry out ammonium assimilation (Wei *et al*., 2021).

In conifers, only one isoform of the GS1b family has been identified to date. The unique GS1b identified in conifers has been suggested to play an essential role in N remobilization to developing organs (Suárez *et al*., 2002). Previous works in pine have shown that GS1b is involved in the canalization of ammonium into glutamine during seed germination and the early developmental stages of seedlings (Ávila *et al*., 2001), which could be important for the loss of seed dormancy (Schneider and Gifford, 1994). Indeed, the roles of GS1b in seed development and germination are also supported by its expression patterns associated with the vascular system of zygotic and somatic pine embryos at different developmental stages and by its expression in procambium cells of pine zygotic embryos (Pérez-Rodríguez *et al*., 2005). Moreover, the expression of this isoenzyme has been suggested to be controlled by gibberellic acid, a phytohormone involved in many aspects of plant growth and development (Gómez-Maldonado *et al*., 2004b).

In this work, a new gene encoding a cytosolic GS (*PpGS1b.2*) was identified in maritime pine (*Pinus pinaster*). This gene was discovered through sequence searches in transcriptomic data from isolated tissues through laser capture microdissection (Cañas *et al*., 2017). Orthologs of this gene have also been identified in the genomes of other conifers, and phylogenetic analysis has revealed that *PpGS1b.2* belongs to the *GS1b* lineage. Although this new *GS1* gene presents a high sequence homology to the already known *PpGS1b*, hereafter *PpGS1b.1*, *PpGS1b.2* showed low expression levels with characteristic and localized tissue expression. The expression patterns suggest that this new gene could play a specific role during plant development, mainly during embryo

development, as has been shown for other *GS1b* genes in angiosperms. Furthermore, a

detailed comparative analysis of the kinetic properties of the isoenzymes GS1b.1 and

GS1b.2 and single/double-point mutants of both isoforms support distinct functions for

these enzymes in pine.

RESULTS

Sequence and phylogenetic analyses

A new cytosolic *GS* gene was identified in a transcriptomic analysis of tissues isolated
using laser capture microdissection (Cañas *et al.*, 2017). At the amino acid sequence level,
the new GS presents 80.85% and 92.68% identity with PpGS1a and PpGS1b, respectively
(Figure 1A). Despite the high identity between the coding sequences of the new gene and
PpGS1b, the promoter regions of both genes are very distinct (Figure S1A). The lengths of
the three pine GS proteins are very similar, with 357 residues for PpGS1a, 355 for PpGS1b
and 357 for the new protein (Figure 1A). However, the calculated isoelectric points were
more different between the pine GS proteins, being 6.21 in the case of PpGS1a, 5.73 for
PpGS1b and 5.36 for the new protein.

A phylogenetic analysis confirmed the classification of the GS from seed plants into three
main groups, GS2, GS1a and GS1b, in line with previously reported results (Valderrama-
Martín *et al.*, 2022) (Figure 1B). As expected, no GS2 sequence was detected in conifers,
but only those of GS1a and GS1b (Figure 1B). The new GS isoform was grouped within
the conifer GS1b sequences; thus, the gene coding this new GS1b isoenzyme has been
named *PpGS1b.2*. Orthologs of *PpGS1b.2* have also been detected in other members of the
Pinaceae family of the genera *Pinus* and *Picea* but not in the rest of the conifers included in
this analysis (Figure 1B).

Gene expression analyses

The expression of *GS* genes in *P. pinaster* has been analyzed in different tissues and
conditions to establish a framework that allows us to unravel the potential role of *PpGS1b.2*
by comparing its expression pattern to other *GS* genes in maritime pine.

The expression profiles were analyzed in embryos and seedlings during the initial
developmental stages (Figure 2A). *PpGS1a* expression was high in cotyledons and needles,
lower in hypocotyls and nearly undetectable in roots and embryos except for germinated
embryos. *PpGS1b.1* and *PpGS1b.2* expression patterns in embryos were very similar, with
a peak of expression in germinated embryos. In seedlings, the expression was ubiquitous in

all organs for both genes, although *PpGS1b.2* expression levels were lower than those of

PpGS1b.1, between 5- and 10-fold. This expression pattern was different when isolated

tissues were considered (Figure 2B). *PpGS1b.1* was expressed at high levels throughout the

plant, especially in the root cortex, where the expression was 40 times that shown by this

gene in the other samples. However, *PpGS1b.2* expression was very localized, mainly in

the shoot apical meristem, emerging needles, developing root vascularization and root

meristem. Expression was almost undetectable in the rest of the tissues analyzed. Finally,

the expression of *PpGS1a* was detected only in the three photosynthetic tissues: the

mesophyll of young needles, the mesophyll of cotyledons and the hypocotyl cortex.

The seasonal expression of the three *GS* genes has also been quantified in needles from

adult trees (Figure 3A). *PpGS1a* showed the highest expression, followed by *PpGS1b.1*,

which was expressed between 10 and 30 times less than *PpGS1a*. The expression levels of

PpGS1b.2 were very low compared to those of other *GSs*. The expression patterns of the

three genes in different whorls were as before, with higher levels in the first months of the

year and lower levels at the end of the year. There was a remarkable exception for whorl 0

in May, the first harvesting month for the needles that emerged during the sampling year.

PpGS1b.2 exhibited an expression peak in whorl 0 in May. In contrast, *PpGS1a* had its

lowest expression, and *PpGS1b.1* was expressed at similar levels to the other whorls. The

relative abundance of *PpGS1b.2* transcripts was still one and two orders of magnitude

lower than those of *PpGS1b.1* and *PpGS1a*, respectively. According to these results, the

expression levels of the three genes were also analyzed in buds and emerging needles

(Figure 3B-D). The expression of *PpGS1a* was almost undetectable in buds, but its

expression rapidly increased in nascent needles by the end of the month. *PpGS1b.1*

expression remained almost invariable in both organs with a similar expression pattern. The

levels of *PpGS1b.2* were higher in the buds and decreased from Day 14 to 28 when the

expression was similar in buds and emerging needles. The relative abundance of *PpGS1b.1*

transcripts was still higher than that of *PpGS1b.2*.

GS gene expression has also been analyzed at different developmental stages, including

juvenile and mature xylem and phloem, as well as the male and female reproductive

structures, different root zones and different stages of zygotic embryo development (Figure

4). In all those samples, *PpGS1a* expression was barely detectable. An example of *PpGS1a*

expression is shown for phloem, xylem, and male and female strobili, with very low levels

(< 0.04), even in female strobilus with an expression peak (< 0.08) (Figure 4A). *PpGS1b.1*

expression was the highest observed thus far among the *GS* genes analyzed in vascular

tissues and strobili (Figure 4A). Interestingly, *PpGS1b.2* expression was almost

undetectable in vascular tissues, but its levels peaked in the male strobilus (approximately

0.28), opposite to what occurred with *PpGS1b.1* in that organ. In root samples, *PpGS1b.1*

and *PpGS1b.2* presented a similar expression pattern, with increased expression in lateral

roots and root tips, although the expression levels for *PpGS1b.1* were approximately 80-

fold higher than that shown by *PpGS1b.2* (Fig, 4B). Finally, in zygotic embryos, the

expression levels of both genes were significantly higher in the precotyledonary and early

cotyledonary stages, where *PpGS1b.2 levels* were higher than those shown by *PpGS1b.1*

(Figure 4C). However, this ratio of the expression of both genes was reversed in the later

stages of development in cotyledonary and mature embryos. Nevertheless, the differences

in expression between the two genes were not statistically significant in either case.

Protein structure prediction and physicochemical and kinetic properties

Very few differences were observed between the GS1b.1 and GS1b.2 subunit structures due

to the similarity of their amino acid sequences (Figure5A,B). Both proteins presented a

predicted decameric structure formed by two pentameric rings with small differences in

structure and the disposition of the subunits in the quaternary structure (Figure S2A,B).

However, the thermodynamic stability of GS1b.1 monomers was three times higher than

that of GS1b.2 monomers (Table 1). The *in silico* replacement of residues of the GS1b.1

and GS1b.2 amino acid sequences displayed some differences in the structural stability of

both enzymes (Figure S3). Some of the amino acids used for this analysis did not cause any

notable effects on the structure or destabilized both proteins equally. However, several

amino acids gave rise to large differences in the free energy of folding. Specifically, the

inclusion of arginine or glutamate around position 280 produced a great destabilization of

the structure of GS1b.2 but not of GS1b.1. Some of these amino acids also caused great

destabilization of GS1b.2 when substituted at position 148 but did not have the same effect

in GS1b.1. In fact, only isoleucine and arginine produced marked effects on the structural

stability of GS1b.1. As small differences in the structure suggested that there might be changes in the physicochemical and kinetic properties of both enzymes, a functional comparison of the recombinant isoforms of GS1b.1 and GS1b.2 was performed (Figure 5, S4; Tables 1, 2).

Both isoforms were tested over a wide pH range; GS1b.1 maximum activity was reached at pH 6.5 while that of GS1b.2 maximum activity was reached at pH 6 (Figure 5C). The activity of both enzymes increased with the reaction temperature, reaching the maximum activity at 42 ℃ (Figure 5D). These data have allowed the calculation of the activation energy (Ea) for each enzyme (Table 1). The Ea was different for both enzymes: the Ea of GS1b.1 was 39.9 kJ/mol, and the Ea values of GS1b.2 for its elemental reaction steps were 46.1 kJ/mol and 18.7 kJ/mol, with a break point at 24 ℃. Regarding the thermal stability, GS1b.1 was very stable, only decreasing its activity at 60 ℃ after 5 minutes of preincubation, although it never completely lost its activity, even after 20 min at 60 ℃ (Figure 5E). However, GS1b.2 showed a decreased activity even after 5 minutes of preincubation at 45 ℃ with almost a total loss of activity after 5 minutes at 60 ℃ (Figure 5E).

GS1b.1 and GS1b.2 showed distinctive behaviors for ammonium and glutamate (Figure 5F,G). GS1b.2 exhibited substrate inhibition for ammonium (K_i 22.57 mM). The affinities of both enzymes for ammonium were high (GS1b.1 K_m 0.12 mM and GS1b.2 K_m 0.21 mM). However, the V_{max} was 5.88 times higher for GS1b.1 (Table 2). Regarding to glutamate, GS1b.1 showed substrate inhibition at high concentrations (K_i 84.51 mM), while GS1b.2 presented positive cooperativity. In both cases, the affinity was very low (GS1b.1 K_m 64.15 mM and GS1b.2 EC50 48.63 mM), with large differences in the V_{max} values of both enzymes (GS1b.1 101.6 nkat/mg protein and GS1b.2 7.66 nkat/mg protein) (Table 2). GS1b.1 and GS1b.2 showed equal behavior for Mg_2^+, with positive cooperativity and similar affinity (EC50 values of 14.49 and 10.87 mM, respectively) but different V_{max} values (71.32 and 5.64 nkat/mg protein, respectively) (Figure S4, Table 2). Finally, the affinities for ATP were high and similar for both enzymes (K_m of 0.18 and 0.29 mM for GS1b.1 and GS1b.2, respectively), with a higher V_{max} for GS1b.1 (24.96 nkat/mg protein) than for GS1b.2 (7.39 nkat/mg protein). However, there was substrate inhibition for GS1b.1 at moderate levels of ATP (K_i 5.88 mM).

Analysis of mutant proteins

To determine the roles that certain residues could play in GS activity, mutants of GS1b.1

and GS1b.2 were obtained by exchanging amino acids at positions 264 and 267. These

residues belong to a region that accumulates a significant number of differences between

the two isoforms and is important for stability, as shown by the *in silico* substitution

analysis (Figure S3). Additionally, these residues have been selected based on their charge

and structural differences between both GSs. The amino acid swapping at positions 264 and

267 seemed to produce only slight changes in the subunit arrangement, even in the double

mutant. Calculation of hydrogen bonds revealed interactions between residues 264 and 267

with those present at positions 261, 263, 265 and 268. These residues were analyzed in

detail, and only small differences in their arrangements could be observed (Figure 6A-G,

S5). The quaternary structures of the mutants also showed no significant differences when

compared (Figure S6) and the thermodynamic stability of the monomers was similar to that

of the WT (Table 1).

Compared to WT, none of the optimal pH values were affected in any of the mutants tested,

except for GS1b.2E264K, where the optimum was reached at pH 7 (Figure S7A), and the

double mutants, where the optimum pH was 6 for both enzymes (Figure 6H, S7).

A slight increase in the optimal temperature (45 °C) was detected in all mutants except for

GS1b.2E264K, which experienced a large change in its optimal temperature (30 °C) (Figure

6I, S7B). Although the activity patterns in response to reaction temperature were similar in

the mutants with respect to the WT enzymes, the activity was slightly higher at all

temperatures in the GS1b.1K267H single and GS1b.2 double mutants. In the case of

GS1b.1 K264E and GS1b.2 H267K, the activity was higher at temperatures above the

optimum (45 °C). Finally, the GS1b.1 double mutant retains considerable activity levels

(>40%) even at very low reaction temperatures, such as 4 °C (Figure 6I, S7B). Ea was

barely affected (Table 1) in GS1b.1K264E (34.8 kJ/mol). In contrast, the GS1b.1 double

mutant Ea was strongly affected (15.2 kJ/mol), and GS1b.1 K267H showed different Ea

values for its elemental reaction steps (35.2 kJ/mol and 6.7 kJ/mol), similar to GS1b.2 WT.

However, GS1b.2 E264K presented a unique Ea for its reaction (39.9 kJ/mol), and different

Ea values were detected for the elemental reaction steps of GS1b.2 H267K (33.7 kJ/mol

and 10.3 kJ/mol) and the GS1b.2 double mutant (28.4 kJ/mol and 6.7 kJ/mol). Interestingly,

all GS1b.1 mutants experienced decreases in their thermostability compared to that of the

WT, and only GS1b.2H267K showed an increased thermostability compared to GS1b.2

WT (Figure 6J, S7C).

GS1b.1 behavior regarding ammonium was only modified in the GS1b.1K267H mutant,

which showed substrate inhibition for ammonium (K_i 13.14 mM). Furthermore, the affinity

was increased in this mutant, GS1b.2H267K, and both double mutants (K_m between 0.02

and 0.09 mM). Meanwhile, all the GS1b.2 mutants lost substrate inhibition by ammonium,

and all exhibited normal hyperbolic saturation (Figure 6K, S8, Table 2). Regarding

glutamate, GS1b.1K264E lost substrate inhibition, now presenting normal hyperbolic

saturation with an increase in its affinity (K_m 2.2 mM) accompanied by a reduction in V_{max}

(16.82 nkat/mg protein). Additionally, none of the mutants in 267 and double mutants

reached saturation and seemed to have lost affinity for this substrate, as occurred with Mg_2^+

in all the mutants except for GS1b.2E264K (Figure S9, S10, Table 2). GS1b.1 mutants

exhibited substrate inhibition by ATP, but only the double mutants of GS1b.1 lost substrate

inhibition by ATP and presented a normal hyperbolic saturation for this substrate (Figure

S11, Table 2). Interestingly, all GS1b.2 mutants presented inhibition by ATP (K_i ranging

from 5.06 to 8.76 mM), in contrast to the hyperbolic Michaelis-Menten saturation exhibited

by the WT (Figure S11, Table 2).

DISCUSSION

The phylogenetic analysis carried out in this work (Figure 1) grouped the new GS isoform (GS1b.2) within the conifer GS1b.1 group. Furthermore, the identification of *GS1b.2* in the genome, its different promoter sequences, including different TF binding sites (Figure S1B), and its different gene expression patterns rule out the possibility that it is an allelic variant of *PpGS1b.1* (HF548531.1), suggesting that *PpGS1b.2* (KU641799.1; KU641800.1) is likely the result of a gene duplication. The presence of *GS1b.2* in members of the genera *Pinus* and *Picea* indicates (Figure 1) that this gene duplication should have taken place in a common ancestor of these two groups but not of the entire *Pinaceae* family since orthologs of *GS1b.2* have not been identified in other conifers. Gene duplication is very common in plants (De Smet and Van de Peer, 2012), and it could lead to the acquisition of new functions (neofunctionalization) or simply to redundant activity to maintain the correct metabolic flux, as occurs with GS in *Populus* and rice (Yamaya and Kusano, 2014; Castro-Rodríguez *et al.*, 2015), contributing to metabolic homeostasis (Moreira *et al.*, 2022). In fact, the GS1b family in angiosperms has been extended by gene duplication so that different isoenzymes can play nonredundant or synergistic roles within the plant, as proposed for *Arabidopsis GS1* genes (Ji *et al.*, 2019).

To explore the possible neofunctionalization of this new *GS* after gene duplication, the expression patterns of the maritime pine *GS* genes were analyzed in different organs and tissues (Figure 2-4). *PpGS1b.2* appears to be expressed primarily in developing organs and tissues and is tightly regulated throughout embryonic development. This contrasts with *PpGS1b.1* expression, which was high in all analyzed samples. This could indicate a strong regulation of *PpGS1b.2* at both the localization and expression levels, suggesting a specialized function. The expression of *PpGS1b.2* is consistent with the association of some *GS1b* isogenes with plant developmental processes in angiosperms (Habash *et al.*, 2001; Tabuchi *et al.*, 2005; Martin *et al.*, 2006; Lothier *et al.*, 2011; Funayama *et al.*, 2013; Goodall *et al.*, 2013; Bao *et al.*, 2014; Guan *et al.*, 2015; Urriola and Rathore, 2015; Gao *et al.*, 2019; Ji *et al.*, 2019; Wei *et al.*, 2021; Fujita *et al.*, 2022). These data suggest an evolutionary convergence that has led to the emergence of GS1b isoforms with similar roles in different plant species. The expansion of the GS1b family in certain conifers

supports that GS1b diversification in angiosperms responds to different plant needs

associated with N assimilation (Hirel and Krapp, 2021). In pine, *GS1b.1* has also been

associated with this function due to its expression during zygotic and somatic embryo

development (Pérez-Rodríguez *et al*., 2005). All these expression data pose different

hypotheses about the role of this new isoenzyme: a) GS1b.2 could support GS1b.1 activity

in developing tissues with a high demand for glutamine or assimilated N; and b) GS1b.2

could play a specific role in certain developing tissues.

To explore the differential roles of GS1b.1 and GS1b.2 in maritime pine, the structure, as

well as the physicochemical and kinetic properties of both enzymes, were analyzed.

Modeling of both maritime pine GS1b isoforms reports small differences between GS1b.1

and GS1b.2 when their tertiary and quaternary structures were compared (Figure 5A,B;

S2). However, any minor difference in subunit arrangements could be of great importance

since the GS active site is formed by the N- and C-terminal domains of adjacent subunits

(Llorca *et al*., 2006).

Although GS1b.1 and GS1b.2 are very similar in their primary sequences and structures,

quite a few differences have been found in their properties. The thermodynamic stability of

GS1b.1 was three times higher than that shown by GS1b.2 (Table 1). Both isoenzymes

present similar values (approximately 63 kJ/mol) for the change in Gibbs free activation

energy ($\Delta G^{\ddagger}$), but their kinetic response to temperature changes below and above 24^{o}C may

be very different (Table 1). For the new isoform, $\Delta G^{\ddagger}$ and the rate-limiting step are

dominated by different activation parameters at different operating temperatures: $\Delta H^{\ddagger}$ for

temperatures below 24^{o}C and T$\Delta S^{\ddagger}$ for temperatures above 24^{o}C. In contrast, GS1b.1

showed a nonvariable activation energy throughout the whole range of temperatures

assayed (Table 1). These differences in dominant activation parameters could reflect

functional differences between the two active sites, as has been previously suggested for

glutamine synthetase isoforms from other sources (Wedler and Horn, 1976).

The optimum pH levels for GS1b.1 and GS1.b2 are 6.5 and 6, respectively (Table 1; Figure

5C), similar to those of GS1b.2 and GS1b.3 from poplar (Castro-Rodríguez *et al*., 2015).

Interestingly, these optimal pH values are lower than the cytosolic pH (7.1-7.5) (Zhou *et*

al., 2021), which could be a mechanism to avoid enzyme inhibition by the acidification

process associated with GS activity and ammonium (Hachiya *et al.*, 2021). The optimum temperature for both enzymes (42 ºC) (Table 1; Figure 5D) is very similar to that shown by GS1b isoenzymes in other plants (Zhao *et al.*, 2014, Castro-Rodríguez *et al.*, 2015). However, both GS1b enzymes had exceptional thermostability compared to other GS1b enzymes of plants (Figure 5E) (Sakakibara *et al.*, 1996; Zhao *et al.*, 2014; Castro-Rodríguez *et al.*, 2015). Concerning glutamate and ATP, GS1b.1 exhibited substrate inhibition behavior, as previously observed for *Arabidopsis* GLN1;3 (Table 2; Figure 5G) (Ishiyama *et al.*, 2004b). These inhibitions are consistent with the role of GS1b.1 in primary nitrogen assimilation in pine and its high expression since high levels of glutamate and ATP, outside of their homeostatic ranges, could indicate metabolic and energetic problems in the cell that may result in unnecessary or detrimental large-scale nitrogen assimilation. Interestingly, GS1b.2 exhibited positive cooperativity for glutamate (Table 2; Figure 5G) and showed substrate inhibition for ammonium (Table 2; Figure 5F). The positive cooperativity mechanism provides high sensitivity to fluctuating substrate concentrations (Levitzki and Koshland, 1976), enabling GS1b.2 to respond rapidly to changes in glutamate availability. In this case, the inhibition of GS1b.2 by ammonium could lead to control of the levels of the final product or to a specific function on the signaling pathway of one of its substrates. This is because both the end product and the substrate of the GS/GOGAT cycle, glutamate and ammonium, have been reported to play roles in plant growth and development (Qiu *et al.*, 2020; Ortigosa *et al.*, 2021), where GS could act as an integrating link for both signaling pathways. Interestingly, glutamate has been described to play important roles in seed germination (Kong *et al.*, 2015), root architecture (Forde, 2014; López-Bucio *et al.*, 2019) and pollen germination and pollen tube growth (Michard *et al.*, 2011; Wudick *et al.*, 2018), among other functions (Qiu *et al.*, 2020). Ammonium has been shown recently to modulate plant root architecture in pine seedlings (Ortigosa *et al.*, 2022). Therefore, based on the *PpGS1b.2* expression patterns, the kinetic characteristics toward glutamate, and previous works, this enzyme could be involved in developmental processes. Furthermore, this could also be a mechanism to avoid high GS activity levels when ammonium is in excess, which could lead to excessive cytosol acidification (Hachiya *et al.*, 2021) of sensitive cells in developing tissues.

The structural, physicochemical, and kinetic analysis carried out in this work on the mutant

enzymes showed some differences from the WT isoforms, but almost none of them

achieved a complete exchange of the properties between GS1b.1 and GS1b.2. The

mutations tested in this work did not greatly affect the protein structure, either in the

surroundings of the exchanged amino acids and the subunit structure (Figure 6A-G) or in

the quaternary structure (Figure S6), which could explain why the thermodynamic stability

of the mutants was not compromised in any case (Table 1). Although all the mutants

presented alterations in the activity levels at the different pH values and temperatures

analyzed in comparison with the WT, only GS1b.1K264E,K267H and GS1b.2E264K

produced variations in the optimal pH, and only GS1b.2E264K presented a considerable

variation in its optimum temperature (Figure S7B) and Ea (Table 1). In fact, among all the

mutants, GS1b.2E264K presented the greatest number of changes in physicochemical

properties. In fact, this could indicate that none of these amino acids have strong

involvement in these enzyme properties or, perhaps, that the changes that can produce these

mutations are being buffered by other residues.

Interestingly, these mutations had large effects on the kinetic properties (Table 2; Figure

6K; S8-S11). The results suggest that these residues are involved in ammonium affinity.

Although it has been described that the presence of glutamine and serine at positions 49 and

174, respectively, is essential for the high affinity for ammonium in *Arabidopsis* GS

(Ishiyama *et al*., 2006), these residues are not present in either GS1b.1 or GSb1.2 of *P*.

pinaster. Previous kinetic studies have shown the presence of high-affinity GS isoforms

that either do not have this combination of amino acids or have none of them (Sakakibara *et*

al., 1996; de la Torre *et al*., 2002; Yadav, 2009; Zhao *et al*., 2014; Castro-Rodríguez *et al*.,

2015). These previous works and the current results support the hypothesis proposed by

Castro-Rodríguez *et al*. (2015), indicating that key residues determining GS behavior for

ammonium may vary between plant species.

Mutations have produced a great number of changes in the behavior of these enzymes

against their substrates and in their kinetic parameters. However, a reversal has only been

achieved for ATP in double mutants, suggesting that the differences in these properties are

due to the collaborative efforts of several residues, probably those that differ between the

two enzymes. This may indicate that GS1b.1 and GS1b.2 have undergone evolutionary

selection so that the two enzymes satisfy different plant needs, with only minor changes in

their amino acid sequences. This hypothesis is also supported by the differences between

the two enzymes at the structural stability level (Figure S3). When introduced at certain

positions, some amino acids had a large effect on the protein stability of one isoform but

not the other. The region between amino acids 260-300 of GS1b.2 was particularly affected

by the introduction of some amino acids, but none of these substitutions appear to produce

similar effects on GS1b.1. In fact, these data suggest that the two enzymes are probably

undergoing different evolutionary paths.

EXPERIMENTAL PROCEDURES

Sequence identification and phylogenetic analyses

416
417

The phylogenetic analysis was made using protein sequences of plant GS that were
obtained from online public databases or assembled from transcriptomic data contained in
the SRA database at the NCBI except for *Pinus pinaster* sequences that were cloned and
sequenced in the present work (Table S1). For the sequence obtaining, the procedure
presented in Valderrama-Martin *et al*. (2022) was followed. Briefly, *tblastn* was used in
BLAST searches (Altschul *et al.*, 1990) using GS1b.1 from *Pinus taeda* as the query.
Transcriptomic assemblies were made in the web platform Galaxy (Afgan *et al.*, 2018).
Raw reads were trimmed using *trimmomatic* (Bolger *et al.*, 2014) and assembled with
Trinity (Grabherr *et al.*, 2011). Database identifiers, names and species for the different
GS sequences are presented in Table S1. All protein sequences used in the present work
are available in Dataset S1.

The sequence data set was composed of 96 GS proteins. The phylogenetic analysis was
mainly focused on conifer GS sequences. The alignment and phylogenetic analysis were
conducted as described in Valderrama-Martin *et al*. (2022) using MEGA version 11
(Tamura *et al.*, 2021). The alignment was conducted with *muscle* (Edgar, 2004). The
phylogenetic analysis was carried out through a maximum-likelihood estimation with
complete deletion of gaps, the missing data, and the Jones–Taylor–Thornton amino acid
substitution model (Jones *et al.*, 1992). Nearest-neighbor interchange was used for tree
inference. The initial tree was constructed using the NJ/BioNJ method. The phylogeny test
was performed using the bootstrap method with 1000 replications. The GS sequences of
Chlamydomonas reinhardtii were used as outer group. The distance matrix and the original
tree in Newick format are available in Datasets S2 and S3. The original tree was visualized
with the Interactive Tree of Life web tool (Letunic and Bork, 2019).

Protein structure prediction and modeling

For the 3D modeling and structure predictions of *P. pinaster* GS1b.1 and GS1b.2 individual
subunits, Alphafold (Jumper *et al*., 2021; Varadi *et al*., 2022) through ColabFold (Mirdita

et al., 2022) has been used. ColabFold allows faster protein structure prediction by
integrating MMseqs2 for multiple sequence alignments and AlphaFold2, but it does not
allow the structure prediction of large protein subunits or complexes. The quaternary
structure prediction has been achieved using Alphafold's models as input for the Galaxy
Package, a combination of several programs that have been designed based on sequence
and structure information together with physical chemistry principles (Shin *et al*., 2014).
The models obtained from ColabFold were employed for the comparison and graphic
representation of the protein structure in PyMOL (https://pymol.org/2/) and in Jmol
(http://www.jmol.org/). Jmol was also used for the calculation of the hydrogen bonds.
Quaternary structure models obtained with Alphafold and the Galaxy Package has been
used in PyMol for the structure analysis and comparison of the models. The
thermodynamic stability of the monomers has been determined using models obtained in
AlphaFold together with the "foldx.mut()" function of the "ptm" R package (Aledo, 2021).

Plant material

Maritime pine seeds (*P. pinaster* Aiton) from *Sierra Segura y Alcaraz* (Albacete, Spain)
(ES17, Ident. 09/10) were provided by the *Red de Centros Nacionales de Recursos*
Genéticos Forestales of the Spanish *Ministerio para la Transición Ecológica y el Reto*
Demográfico with the authorization number ESNC103. Pine seeds were imbibed for 48 h in
water with aeration to induce germination. Seeds were germinated in vermiculite. Seedlings
were grown in plant growth chambers (Aralab Fitoclima 1200, Rio de Mouro, Portugal)
under 16 h light photoperiod, a light intensity of 125 µmol m^{-2} s^{-1}, a constant
temperature of 23 °C, 50% relative humidity and watered twice a week with distilled
water. Embryo and seedling samples were harvested at different stages: dry, post-
imbibition and germinated (0.5 cm of emerged radicle) embryos; and one-week-old from
emergence (Stage 1) and one-month-old from emergence seedlings (Stage 2). At the
harvest, seedlings were divided into their different organs. For the measure of *GS* gene
expression in different sections of roots, 2 months-old seedlings were used. The samples
were immediately frozen in liquid N and stored at -80 ℃ until powdering with a mixer mill
MM400 (Retsh, Haan, Germany) and further analyses were conducted.

Plant material and cDNA to analyze *GS* gene expression levels in maritime pine tissues

from one-month-old seedlings were previously obtained by Cañas *et al*. (2017). RNA

samples from 14 tissues isolated through laser capture microdissection were employed. The

cDNA was synthesized and amplified as described by Cañas *et al*. (2014).

Samples from Cañas *et al*. (2015) were used to analyze *GS* gene expression in needles of

adult trees. Briefly, needle whorls corresponding to the annual growth of a single year were

harvested from different 25 years old *P. pinaster* specimens at *Los Reales de Sierra*

Bermeja (Estepona, Spain). Whorls were named from 0 to 3 referring to the year of

appearance of that whorl. Whorl 0 was first collected in May when it was completely

formed. Samples were collected each month throughout 2012, were immediately frozen in

liquid nitrogen and stored at –80°C until their utilization for RNA extraction. Buds and

nascent needles were collected from the same adult specimens once a week during April of

2013. For gene expression analyses three different trees were employed.

Juvenile and mature phloem, together with male and female strobili were harvested from 25

to 35-year-old maritime pines located *at Los Reales de Sierra Bermeja* (Estepona, Spain)

Juvenile xylems were collected from the last 5 internodes in the crown and mature xylem

from the base of the trunk of 28 to 31-year-old maritime pines from *Los Reales de Sierra*

Bermeja by removing bark and phloem and scraping with a sterile blade. (Villalobos,

2008). All the tissues were frozen immediately using liquid nitrogen and storage at -80 ºC

until use

Zygotic embryos from *P. pinaster* were obtained from a single maritime pine seed orchard

(PP-VG-014, Picard, Saint-Laurent-Médoc, France) and collected at different

developmental stages (Avila et al., 2022). All samples were frozen in liquid nitrogen and

stored at −80°C until use.

RNA extraction and RT-qPCR

Total RNA from maritime pine samples was extracted following Canales et al. (2012).

RNA concentration and purity (A260/A280) was then quantified using a NanoDrop© ND-

1000 spectrophotometer (ThermoFisher Scientific, Walthman, MA, USA). The integrity of

the RNA was checked by electrophoresis. iScript Reverse Transcription Supermix (Bio-

Rad, Hercules, CA, USA) was used for the reverse transcription of 500 ng of total RNA of

each sample in a final volume reaction of 10µL including 2µL of reaction buffer and 0,5µL

of reverse transcriptase enzyme with the following conditions in a thermal cycler with the

following conditions: 30 min at 42°C; 10 min at 65°C; hold at 4°C.

For the RT-qPCR analysis, three biological samples were used with three technical

replicates each. The qPCR was carried out using 5 μL SsoFastTM EvaGreen® Supermix

(Bio-Rad, Hercules, CA, USA), 10 ng of cDNA, 20 pmol of each primer in a total reaction

volume of 10 μL on a C1000[TM] Thermal Cycler with a CFX384[TM] Touch Realm-Time PCR

Detection System (Bio-Rad, Hercules, CA, USA) with the following conditions: initial

denaturation step at 95°C 2 min; 40 cycles of denaturation at 95 °C 5 s and elongation at 60

ºC 20 s. Finally, a melt curve was developed from 65 to 95 ºC with increments of 0.5 ºC

each 5 s. Two maritime pine *saposin-like aspartyl protease* and *RNA binding protein* genes

were used as reference for results normalization (Granados et al., 2016). Expression data

have been analyzed using the *qpcR* R library and the MAK3 model (Ritz and

Spiess,☐2008). The primers used for RT-qPCR assays are presented in Table S2.

Cloning, mutagenesis, recombinant expression, and purification of GS1b.1 and

GS1b.2

In the search for new *GS* genes in conifers, 3 genes in *P. pinaster* have been identified in

transcriptome databases that were named as *PpGS1a*, *PpGS1b.1* and *PpGS1b.2*. The cDNA

of the three genes were amplified by PCR using iProof HF Master Mix (Bio-Rad, Hercules,

CA, USA) and cloned into the pJET1.2 vector (ThermoFisher Scientific, Walthman, MA,

USA) following the manufacturers' instructions. The used primers were designed from

sequences obtained from the maritime pine transcriptome assembled in Cañas *et al*. (2017).

Primers are shown in Table S2. *PpGS1a* was obtained from amplified cDNA of emerging

needles (EN) isolated in Cañas *et al*. (2017). *PpGS1b.1* and *PpGS1b.2* were obtained from

amplified cDNA of developing root cortex (DRC) isolated in Cañas *et al*. (2017).

For protein recombinant expression, the CDS of wild type (WT) *PpGS1b.1* and *PpGS1b.2*

were subcloned into pET30a vector (Merck, Darmstadt, Germany) including a N-terminal

6xHis-tag by PCR. For this task, AseI and XhoI sites were added to *PpGS1b.1* 5´and

3´ends respectively while NdeI and XhoI sites were added to *PpGS1b.2* 5´and 3´endings

respectively. These restriction sites along with the 6xHis-tag were introduced by PCR.

Used primers are listed in Table S2. The plasmid and PCR product were then cut using the

appropriate restriction enzymes and the PCR product was inserted into the plasmid using

T4 DNA ligase.

Plasmids were transformed and expressed in the *Escherichia coli* strain BL-21 (DE3) RIL

cells (Agilent, Santa Clara, CA, USA). For protein expressions, the bacterial clones were

grown at 37 ºC and 180 rpm in an orbital shaker with 500 mL of Luria-Bertani medium

supplemented with kanamycin (0.05 mg/mL) and chloramphenicol (0.034 mg/mL). When

the optical density (OD) reached a 0.5-0.6 value at 600 nm, cultures were tempered and

isopropyl-β-D-thiogalactoside was added to a final concentration of 1 mM to induce protein

expression. Once the isopropyl-β-D-thiogalactoside was supplied the cultures were

incubated at 25 ºC and 120 rpm for 5 hours, the cells were collected by centrifugation. The

bacterial pellet was resuspended in 5 mL of buffer A (Tris 50 mM pH 8; NaCl 300 mM;

imidazole 250 mM) with 4 mg of lysozyme and incubated for 30 min in ice, bacteria were

then lysed by ultrasonication with 20 pulses of 5 seconds at 20% amplitude with 5 seconds

rest between pulses in a Branson Sonifier® Digital SFX 550 (Branson Ultrasonics, CT,

USA). The soluble fraction was clarified by centrifugation (1620 x g at 4 ºC for 30 min).

Proteins from the soluble fraction were purified by affinity chromatography with Protino

Ni-TED PackedColumns2000 (Macherey-Nagel, Düren, Germany) based on the His-tag

tail. The soluble fraction from bacterial lysate was loaded in a column previously

equilibrated with buffer A. Protein elution was performed by adding buffer B (Tris 50 mM

pH8; NaCl 300 mM; imidazole 250 mM) and a total of 9 mL of eluate was recovered in 1

mL fractions. Collected fractions were quantified by Bradford (Bradford, 1976) and

analyzed on SDS-page and western-blot using GS-specific antibodies obtained from rabbit

(Figure S12) (Cantón *et al.*, 1996). Fractions containing the proteins were concentrated

with Amicon® Ultra-15 Centrifugal Filters Ultracel®-100K (Merck-Millipore, Burlington,

Massachusetts, State of Virginia) with 100 kDa pores and the resulting concentrate was

stored in 50% (v/v) glycerol at -20 ºC for later kinetic measurements and physicochemical

analyses.

Site-directed mutagenesis

Considering characteristics and properties of differing amino acids between GS1b.1 and

GS1b.2, residues at position 264 and 267 were selected to be shifted between both

isoenzymes. Site-directed mutagenesis was carried out following Edelheit *et al*. (2009). The

wild type CDS from those sequences included on the pET30a vector were amplified by

PCR using two reverse-complementary primers (Table S2) that already included the

mutation to be introduced. The primers were used separately in a PCR reaction using 50 or

500ng of plasmid and 10 pmol of each primer. The final products of both reactions were

then mixed and hybridized. The PCR products were checked out in an agarose gel and

purified using NucleoSpin® Gel and PCR Clean-up (Macherey-Nagel, Düren, Germany).

Finally, the PCR product was digested with FastDigest® DpnI (ThermoFisher Scientific,

Walthman, MA, USA) to degrade the vector used as template for the amplification.

Physicochemical assays

Physicochemical properties were determined by conducting the transferase assay as

described in Cánovas *et al*. (1991). Reactions were carried out in 96 well microtiter plates

with a final reaction volume of 150 µL. The reaction mix contained 90.6 mM MOPS pH 7,

20 mM arsenate, 2.93 mM $MnCl_2$, 60 mM NH_2OH and 0.4 mM ADP. When determining

the optimal pH level for the activity of the different isoforms, different buffers were used

instead when determining the optimal pH level for the activity of the different isoforms:

acetate (4.5-5); MES (6-6.5); HEPES (7-7.5); Tris (8-8.5); and sodium carbonate (9-10).

The reaction was initiated by adding glutamine in a final concentration of 120 mM and,

after 15 minutes of incubation at 37 ºC, 150 µL of STOP solution (10% $FeCl_3$ • 6 H_2O in

HCl 0.2 N; 24% trichloroacetic acid and 5% HCl) was added to stop the reaction. Finally,

the plate was centrifugated for 3 minutes at 3220 x g and 100 µL of the reaction volume

were withdrawn for its absorbance measurement at 540 nm in a PowerWave HY (BioTek,

Winooski, VT; USA) plate lector. For thermostability characterization, proteins were
preincubated at different times and temperatures before adding the reaction mix.

Kinetic assays

For the quantification of the kinetic properties, biosynthetic assays were carried out as
described by Gawronski and Benson (2004) with some modifications. Reactions were
conducted in 96 wells microtiter plates in a final volume of 100 µL. GS activity was
determined as a function of NADH absorbance depletion at 340 nm in a coupled reaction
using lactate-dehydrogenase (LDH, EC 1.1.1.27) and pyruvate kinase (PyrK, EC 2.7.1.40).
The following reaction mix was used: 50 mM Hepes pH 7, 10 mM $MgCl_2$, 60 mM NH_4Cl,
250 mM glutamate, 6.25 mM ATP, 1 mM phosphoenolpyruvate, 0.6 mM NADH, 1U
PyRK and 1U LDH. Reactions were pre-incubated for 5 minutes at 37 ºC and the GS
activity was initiated by adding different concentrations of the substrate that was being
analyzed. Reactions were developed for 40 min at 37 ºC with shaking and absorbance
measurement at 340 nm each minute. Analysis of the kinetic characteristics of GS1b.1 WT,
GS1b.2 WT and their mutants were performed with GraphPad Prism 8.0.0 (GraphpPad,
San Diego, CA, USA).

ACCESSION NUMBERS

The cDNA sequence data have been submitted to the GenBank database under accession
numbers KU641797 (PpGS1a); KU641798, HF548531.1 (PpGS1b.1); and KU641796,
KU641799.1, KU641800.1 (PpGS1b.2).

ACKNOWLEDGEMENTS

The authors are grateful to Professor Francisco Ruiz Cantón for the kind supply of adult
tree samples, to Jean François Trontin for the kind supply of embryo samples and to José
Miguel Granados for his help during needle harvesting. This work was funded by *Spanish*
Ministerio de Economía y Competitividad, grant number BIO2015-73512-JIN
MINECO/AEI/FEDER, UE, and *Ministerio de Ciencia e Innovación*, grant number
PID2021-125040OB-I00, MICINN, FEDER, UE. This work was also supported by *Junta*
de Andalucía, grant number P20_00036 PAIDI 2020/FEDER, UE; and the *Universidad de*

Málaga grant B4-2021-01 (Ayudas Plan Propio). JMVM was supported by a scholarship

from the Spanish *Ministerio de Educación y Formación Profesional* (FPU17/03517). FO

was supported by a grant from the Universidad de Málaga (*Programa Operativo de Empleo*

Juvenil vía SNJG, UMAJI11, FEDER, FSE, *Junta de Andalucía*) and funds of the research

group BIO-114 (Biología Molecular y Biotecnología) from PAIDI, Junta de Andalucía.

AUTHOR CONTRIBUTIONS

JMVM, FO and RAC have performed the experiments. RAC performed the phylogenetic

analysis. JMVM and JCA have made the *in silico* structural protein analyses. JMVM and

RAC have written the manuscript. FO, JCA, CA and FMC made additional contributions

and edited the manuscript. JMVM and RAC have planned and designed the research. RAC,

CA and FMC were responsible of the funding acquisition.

CONFLICT OF INTEREST

The authors declare that they have no conflicts of interest in relation to the content of this

manuscript.

DATA AVAILABILITY

Constructions are available from the corresponding authors upon request. The rest of data

supporting the findings of this study are available within the paper and within its

supplementary materials published online.

SHORT LEGENDS FOR SUPPORTING INFORMATION

Dataset S1. Protein sequences used for phylogenetic analysis.

Dataset S2. Phylogenetic distance matrix.

Dataset S3. Original tree resulted from phylogenetic analysis in Newick format.

Table S1. GSs used for phylogenetic analysis.

Table S2. List of primers.

648 **Figure S1. Promoter region analysis of *PpGS1b.1* and *PpGS1b.2*.** For promoter

649 comparison, 1916 and 2408 nucleotides upstream of start codon of *PpGS1b.1* and

650 *PpGS1b.2* were recovered from genomic data, respectively (Sterck et al., 2022). **A.**

651 Sequences alignment of the promoter region. Position 1995 corresponds to -1 nt before start

652 codon. **B.** Putative transcription factor binding sites identified using the PlantRegMap

653 prediction tool (Tian et al., 2020).

654 **Figure S2. GS1b.1 WT and GS1b.2 WT quaternary structure. A.** GS1b1.1 WT

655 quaternary structure. **B.** GS1b.2 WT quaternary structure.

656 **Figure S3. Structural stability of GS1b.1 and GS1b.2 against amino acid substitution.**

657 The presented amino acids have been substituted in each position of GS1b.1 and GS1b.2

658 amino acid sequences. The differences in the folding free energy between WT and mutant

659 ($\Delta\Delta G$) for GS1b.1 and GS1b.2 are compared (square plots). The rectangular plots represent

660 the difference between GS1b.1 $\Delta\Delta G$ and GS1b.2 $\Delta\Delta G$ (Y axis) for each position (X axis).

661 **Figure S4. Representation of kinetic characteristics of GS1b.1 WT and GS1b.2 WT**

662 **for magnesium and ATP. A.** Magnesium. **B.** ATP.

663 **Figure S5. Subunit structure of the GS1b.1 and GS1b.2 mutants. A.** GS1b.1K264E. **B.**

664 GS1b.1K267H. **C.** GS1b.2E264K. **D.** GS1b.2H267K. Amino acids exchanged, and amino

665 acids associated with them by hydrogen bonds are represented in dark magenta. Amino

666 acids from 330 to the end of the protein are represented in green.

667 **Figure S6. Quaternary structure of the GS1b.1 and GS1b.2 mutants. A.** GS1b.1K264E.

668 **B.** GS1b.1K267H. **C.** GS1b.1K264E,K267H. **D.** GS1b.2E264K. **E.** GS1b.2H267K. **F.**

669 GS1b.2E264K,H267K.

670 **Figure S7. Physicochemical properties of the GS1b.1 and GS1b.2 mutants.** Activity of

671 the GS1b.1K264E, GS1b.1K267H, GS1b.2E264K, GS1b.2H267K have been tested at

672 different pH levels (**A**) and temperatures (**B**). Their thermal stability has been also

673 characterized (**C**).

674 **Figure S8. Representation of kinetic characteristics of GS1b.1 and GS1b.2 single**

675 **mutants for ammonium.**

REFERENCES

Afgan, E., Baker, D., Batut, B., van den Beek, M., Bouvier, D., Cech, M. *et al.* (2018) The Galaxy platform for accessible, reproducible and collaborative biomedical analyses: 2018 update. *Nucleic Acids Research*, **46**, W537-W544. https://doi.org/10.1093/nar/gky379

Aledo, J.C. (2021) ptm: an R package for the study of methionine sulfoxidation and other post-translational modifications. *Bioinformatics*, **37**, 3979–3980. https://doi.org/10.1093/bioinformatics/btab348

Altschul, S.F., Gish, W., Miller, W., Myers, E.W. & Lipman D.J. (1990) Basic local alignment search tool. *Journal of Molecular Biology*, **215**, 403-410. https://doi.org/10.1016/S0022-2836(05)80360-2

Ávila, C., Suárez, M.F., Gómez-Maldonado, J. & Cánovas, F.M. (2001) Spatial and temporal expression of two cytosolic glutamine synthetase genes in Scots pine: functional implications on nitrogen metabolism during early stages of conifer development. *The Plant Journal*, **25**, 93-102. https://doi.org/10.1046/j.1365-313x.2001.00938.x

Ávila, C., Llebrés, M.T., Castro-Rodríguez, V., Lobato-Fernández, C., Reymond, I., Harvengt, L. et al. (2022) Identification of metabolic pathways differentially regulated in somatic and zygotic embryos of maritime pine. *Frontiers in Plant Science*, **13**, 877960. https://doi.org/10.3389/fpls.2022.877960

Bao, A., Zhao, Z., Ding, G., Shi, L., Xu, F. & Cai, H. (2014) Accumulated expression level of cytosolic *glutamine synthetase 1* gene (*OsGs1;1* or *OsGS1;2*) alter plant development and the carbon-nitrogen metabolic status in rice. *PLoS ONE*, **9**, e95581. https://doi.org/10.1371/journal.pone.0095581

Bernard, S.M. & Habash, D.Z. (2009) The importance of cytosolic glutamine synthetase in nitrogen assimilation and recycling. *New Phytologist*, **182**, 608–620. https://doi.org/10.1111/j.1469-8137.2009.02823.x

Blackwell, R.D., Murray, A.J.S. & Lea, P.J. (1987) Inhibition of photosynthesis in barley with decreased levels of chloroplastic glutamine synthetase activity. *Journal of Experimental Botany*, **196**, 1799–1809. https://doi.org/10.1093/jxb/38.11.1799

Bolger, A.M., Lohse, M. & Usadel, B. (2014) Trimmomatic: a flexible trimmer for Illumina Sequence Data. *Bioinformatics*, **30**, 2114-2120. https://doi.org/10.1093/bioinformatics/btu170

Bradford, M.M. (1976) A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. *Analytical Biochemistry*, **72**, 248-254. https://doi.org/10.1006/abio.1976.9999

Canales, J., Rueda-López, M., Craven-Bartle, B., Ávila, C. & Cánovas, F.M. (2012) Novel insights into regulation of asparagine synthetase in conifers. *Frontiers in Plant Science*, **3**, 100. https://doi.org/10.3389/fpls.2012.00100

Cánovas, F.M., Cantón, F.R., Gallardo, F., García-Gutiérrez, A. & de Vicente, A. (1991) Accumulation of glutamine synthetase during early development of maritime pine seedlings (*Pinus pinaster*). *Planta*, **185**, 372-378. https://doi.org/10.1007/BF00201059

Cantón, F.R., García-Gutiérrez, A., Crespillo, R. & Cánovas, F.M. (1996) High-level expression of *Pinus sylvestris* glutamine synthetase in *Escherichia coli*. Production of polyclonal antibodies against the recombinant protein and expression studies in pine seedlings. *FEBS Letters*, **393**, 205-210. https://doi.org/10.1016/0014-5793(96)00886-1

Cantón, F.R., Suárez, M.F., Josè-Estanyol, M. & Cánovas, F.M. (1999) Expression analysis of a cytosolic glutamine synthetase gene in cotyledons of Scots pine seedlings: developmental, light regulation and spatial distribution of specific transcripts. *Plant Molecular Biology*, **404**, 623-34. https://doi.org/10.1023/a:1006219205062

Cañas, R.A., Canales, J., Gómez-Maldonado, J., Ávila, C. & Cánovas, F.M. (2014) Transcriptome analysis in maritime pine using laser capture microdissection and 454 pyrosequencing. *Tree physiology*, **34**, 1278-1288. https://doi.org/10.1093/treephys/tpt113

Cañas, R.A., Canales, J., Muñoz-Hernández, C., Granados, J.M., Ávila, C., García-Martín, M.L. et al. (2015) Understanding developmental and adaptive cues in pine through metabolite profiling and co-expression network analysis. *Journal of Experimental Botany*, **66**, 3113-3127. https://doi.org/10.1093/jxb/erv118

Cañas, R.A., Li, Z., Pascual, M.B., Castro-Rodríguez, V., Ávila, C., Sterck, L. et al. (2017) The gene expression landscape of pine seedling tissues. *The Plant Journal*, **91**, 1064-1087. https://doi.org/10.1111/tpj.13617

Castro-Rodríguez, V., García-Gutiérrez, A., Cañas, R.A., Pascual, M.B., Ávila, C. & Cánovas, F.M. (2015) Redundancy and metabolic function of the glutamine synthetase gene family in poplar. *BMC Plant Biology*, **15**, 20. https://doi.org/10.1186/s12870-014-0365-5

de la Torre, F., García-Gutiérrez, A., Crespillo, R., Cantón, F.R., Ávila, C. & Cánovas, F.M. (2002) Functional expression of two pine glutamine synthetase genes in bacteria reveals that they encode cytosolic holoenzymes with different molecular and catalytic properties. *Plant and Cell Physiology*, **43**, 802-809. https://doi.org/10.1093/pcp/pcf094

De Smet, R. & Van de Peer, Y. (2012) Redundancy and rewiring of genetic networks following genome-wide duplication events. *Current Opinion in Plant Biology*, **15**, 168-176. https://doi.org/10.1016/j.pbi.2012.01.003

Edelheit, O., Hanukoglu, A. & Hanukoglu, I. (2009) Simple and efficient site-directed mutagenesis using two single-primer reactions in parallel to generate mutants for protein structure-function studies. *BMC Biotechnology*, **9**, 61. https://doi.org/10.1186/1472-6750-9-61

Edgar, R.C. (2004) MUSCLE: multiple sequence alignment with high accuracy and high throughput. *Nucleic Acids Research*, **32**, 1792-1797. https://doi.org/10.1093/nar/gkh340

Forde, B.G. & Lea, P.J. (2007) Glutamate in plants: metabolism, regulation, and signaling. *Journal of Experimental Botany*, **58**, 2339-2358. https://doi.org/10.1093/jxb/erm121

Forde, B.G. (2014) Glutamate signalling in roots. *Journal of Experimental Botany*, **65**, 779-787. https://doi.org/10.1093/jxb/ert335

Fujita, T., Beier, M.P., Tabuchi-Kobayashi, M., Hayatsu, Y., Nakamura, H., Umetsu-Ohashi, T. et al. (2022) Cytosolic glutamine synthetase GS1;3 is involved in rice grain

ripening and germination. *Frontiers in Plant Science*, **13**, 835835. https://doi.org/10.3389/fpls.2022.835835

Funayama, K., Kojima, S., Tabuchi-Kobayashi, M., Sawa, Y., Nakayama, Y., Hayakawa, T. et al. (2013) Cytosolic glutamine synthetase 1;2 is responsible for the primary assimilation of ammonium in rice roots. *Plant and Cell Physiology*, **54**, 934-943. https://doi.org/10.1093/pcp/pct046

Gao, Y., de Bang, T.C. & Schjoerring, J.K. (2019) Cisgenic overexpression of cytosolic glutamine synthetase improves nitrogen utilization efficiency in barley and prevents grain protein decline under elevated CO_2. *Plant Biotechnology Journal*, **17**, 1209-1221. https://doi.org/10.1111/pbi.13046

Gawronski, J.D. & Benson, D.R. (2004) Microtiter assay for glutamine synthetase biosynthetic activity using inorganic phosphate detection. *Analytical Biochemistry*, **327**, 114-118. https://doi.org/10.1016/j.ab.2003.12.024

Gómez-Maldonado, J., Ávila, C., Barnestein, P., Crespillo, R. & Cánovas, F.M. (2004a) Interaction of cist-acting elements in the expression of a gene encoding cytosolic glutamine synthetase in pine seedlings. *Physiologia Plantarum*, **121**, 537-545. https://doi.org/10.1111/j.1399-3054.2004.00353.x

Gómez-Maldonado, J., Cánovas, F.M. & Ávila, C. (2004b) Molecular analysis of the 5'-upstream of a gibberellin-inducible cytosolic glutamine synthetase gene (*GS1b*) expressed in pine vascular tissue. *Planta*, **218**, 1036-1045. https://doi.org/10.1007/s00425-003-1185-2

Goodall, A.J., Kumar, P. & Tobin, A.K. (2013) Identification and expression analyses of cytosolic glutamine synthetase genes in barley (*Hordeum vulagre* L.). *Plant and Cell Physiology*, **54**, 492-505. https://doi.org/10.1093/pcp/pct006

Grabherr, M.G., Haas, B.J., Yassour, M., Levin, J.Z., Thompson, D.A., Amit, I. et al. (2011) Full-length transcriptome assembly from RNA-seq data without a reference genome. *Nature Biotechnology*, **29**, 644-652. https://doi.org/10.1038/nbt.1883

Guan, M., Moller, I.S. & Schjoerring, J.K. (2015) Two cytosolic glutamine synthetase isoforms play specific roles for seed germination and seed yield structure in *Arabidopsis*. *Journal of Experimental Botany*, **66**, 203-212. https://doi.org/10.1093/jxb/eru411

Granados, J.M., Ávila, C., Cánovas, F.M. & Cañas, R.A. (2016) Selection and testing of reference genes for accurate RT-qPCR in adult needles and seedlings of maritime pine. *Tree Genetic & Genomes*, **12**, 60-75. https://doi.org/10.1007/s11295-016-1018-7

Habash, D.Z., Massiah, A.J., Rong, H.L., Wallsgrove, R.M. & Leigh, R.A. (2001) The role of cytosolic glutamine synthetase in wheat. *Annals of Applied Biology*, **138**, 83-89. https://doi.org/10.1111/j.1744-7348.2001.tb00087.x

Hachiya, T., Inaba, J., Wakazaki, M., Sato, M., Toyooka, K., Miyagi, A., et□al. (2021) Excessive ammonium assimilation by plastidic glutamine synthetase causes ammonium toxicity in *Arabidopsis thaliana*. *Nature Communications*, **12**, 4944. https://doi.org/10.1038/s41467-021-25238-7

Heldt, H. & Piechulla, B. (2011) *Plant Biochemestry*: fourth edition. San Diego: Academic Press, Elsevier.

Hirel, B. & Krapp, A. (2021) Nitrogen utilization in plants I biological and agronomic importance. In: Jez, J. (Ed.) *Encyclopedia of biological chemistry III*, Third edition. Amsterdam: Elsevier, pp. 127-140.

Ishiyama, K., Inoue, E., Tabuchi, M., Yamaya, T. & Takahashi, H. (2004a) Biochemical background and compartmentalized functions of cytosolic glutamine synthetase for active ammonium assimilation in rice roots. Plant and Cell Physiology, 45, 1640-1647. https://doi.org/10.1093/pcp/pch190

Ishiyama, K., Inoue, E., Watanabe-Takahashi, A., Obara, M., Yamaya, T. & Takahashi, H. (2004b) Kinetic properties and ammonium-dependent regulation of cytosolic isoenzymes of glutamine synthetase in *Arabidopsis*. *The Journal of Biological Chemistry*, **279**, 16598-16605. https://doi.org/10.1074/jbc.M313710200

Ishiyama, K., Inoue, E., Yamaya, T. & Takahashi, H. (2006) Gln49 and Ser174 residues play critical roles in determining the catalytic efficiencies of plant glutamine synthetase. *Plant and Cell Physiology*, **47**, 299-303. https://doi.org/10.1093/pcp/pci238

James, D., Borphukan, B., Fartyal, D., Achary, V.M.M. & Reddy, M.K. (2018) Transgenic manipulation of glutamine synthetase: A target with untapped potential in

various aspects of crop improvement. In: Gosal SS, Wani SH, editors. *Biotechnology of Crop Improvement.* Cham: Springer International Publishing AG, pp 367–416.

Ji, Y., Li, Q., Liu, G., Selvaraj, G., Zheng, Z. & Wei, Y. (2019) Roles of cytosolic glutamine synthetase in *Arabidopsis* development and stress responses. *Plant and Cell Physiology*, **60**, 657-671. https://doi.org/10.1093/pcp/pcy235

Jmol. An open-source Java viewer for chemical structures in 3D. http://www.jmol.org/. Accessed August 2022.

Jones, D.T., Taylor, W.R. & Thornton, J.M. (1992) The rapid generation of mutation data matrices from protein sequences. *Computer Applications in the Biosciences*, **8**, 275-282. https://doi.org/10.1093/bioinformatics/8.3.275

Jumper, J., Evans, R., Pritzel, A., Green, T., Figurnov, M., Ronneberger, O. et al. (2021) Highly accurate protein structure prediction with AlphaFold. *Nature*, **596**, 583-589. https://doi.org/10.1038/s41586-021-03819-2

.**Kong, D., Ju, C., Parihar, A., Kim, S., Cho, D. & Kwak, J.M.** (2015) *Arabidopsis* glutamate receptor homolog3.5 modulates cytosolic Ca^{2+} level to continue effect of abscisic acid in seed germination. *Plant Physiology*, **167**, 1630-1642. https://doi.org/10.1104/pp.114.251298

Lea, P.J. & Ireland, R.J. (1999) Nitrogen metabolism in higher Plants. In: Singh BK, editor. *Plant Amino Acids*. New York: Marcel Dekker Inc, pp. 1-47.

Letunic, I. & Bork, P. (2019) Interactive Tree Of Life (iTOL) v4: recent updates and new developments. *Nucleic Acids Research,* **47**, W256-W259. https://doi.org/10.1093/nar/gkz239

Levitzki, A. & Koshland, D.E. Jr (1976) The role of negative cooperativity and half-of-the-sites reactivity in enzyme regulation. *Current Topics in Cellular Regulation*, **10**, 1-40. https://doi.org/10.1016/b978-0-12-152810-2.50008-5

Llorca, O., Betti, M., González, J.M., Valencia, A., Márquez, A.J. & Valpuesta, J.M. (2006) The three-dimensional structure of an eukaryotic glutamine synthetase: Functional implications of its oligomeric structure. *Journal of Structural Biology*, **156**, 469-479. https://doi.org/10.1016/j.jsb.2006.06.003

López-Bucio, J.S., de la Cruz, H.R. & Guevara-García, A.A. (2019) Glutamate sensing in plants. In: Ramakrishna A, Roshchina VV, editors. *Neurotransmitters in plants: perspectives and applications*. Raton: CRC Press, pp. 231–140.

Lothier, J., Gaufichon, L., Sormani, R., Lemaître, T., Azzopardi, M., Morin, H. et al. (2011) The cytosolic glutamine synthetase GLN1;2 plays a role in the control of plant growth and ammonium homeostasis in *Arabidopsis* rosettes when nitrate supply is not limiting. *Journal of Experimental Botany*, **62**, 1375-1390. https://doi.org/10.1093/jxb/erq299

Michard, E., Lima, P.T., Borges, F., Silva, A.C., Portes, M.T., Carvalho, J.E. et al. (2011) Glutamate receptor-like genes form Ca^{2+} channels in pollen tubes and are regulated by pistil D-serine. *Science*, **332**, 434-437. https://doi.org/10.1126/science.1201101

Martin, A., Lee, J., Kichey, T., Gerentes, D., Zivy, M., Tatout, C. et al. (2006) Two cytosolic glutamine synthetase isoforms of maize are specifically involved in the control of grain production. *The Plant Cell*, **18**, 3252-3274. https://doi.org/10.1105/tpc.106.042689

Mirdita, M., Schütze, K., Moriwaki, Y., Heo, L., Ovchinnikov, S. & Steinegger, M. (2022) ColabFold: making protein folding accessible to all. *Nature Methods*, **19**, 679-682. https://doi.org/10.1038/s41592-022-01488-1

Moreira, E., Coimbra, S. & Melo, P. 2022. Glutamine synthetase: an unlikely case of functional redundancy in *Arabidopsis thaliana*. *Plant Biology*, **24**, 713-720. https://doi.org/10.1111/plb.13408

Pérez-Rodríguez, M.J., Suárez, M.F., Heredia, R., Avila, C., Breton, D., Trontin, J.F. et al. (2005) Expression patterns of two glutamine synthetase genes in zygotic and somatic pine embryos support specific roles in nitrogen metabolism during embryogenesis. *New Phytologist*, **169**, 35-44. https://doi.org/10.1111/j.1469-8137.2005.01551.x

Qiu, X.M., Sun, Y.Y., Ye, X.Y. & Li, Z.G. (2020) Signaling role of glutamate in plants. *Frontiers in Plant Science*, **13**, 1743. https://doi.org/10.3389/fpls.2019.01743

Ortigosa, F., Lobato-Fernández, C., Shikano, H., Ávila, C., Taira, S., Cánovas, F.M. et al. (2022) Ammonium regulates the development of pine roots through hormonal crosstalk

and differential expression of transcription factors in the apex. *Plant, Cell and Environment*, **45**, 915-935. https://doi.org/10.1111/pce.14214

Ritz, C. & Spiess, A.N. (2008) qpcR: an R package for sigmoidal model selection in quantitative real-time polymerase chain reaction analysis. *Bioinformatics*, **24**, 1549-1551. https://doi.org/10.1093/bioinformatics/btn227

Sakakibara, H., Shimizu, H., Hase, T., Yamazaki, Y., Takao, T., Shimonishi, Y. et al. (1996) Molecular identification and characterization of cytosolic isoforms of glutamine synthetase in maize roots. *The Journal of Biological Chemistry*, **271**, 29561-29568. https://doi.org/10.1074/jbc.271.47.29561

Schneider, W.L. & Gifford, D.J. (1994) Loblolly pine seed dormancy. I. The relationship between protein synthesis and the loss of dormancy. *Physiologia Plantarum*, **90**, 246-252. https://doi.org/10.1111/j.1399-3054.1994.tb00384.x

Shin, W.H., Lee, G.R., Heo, L., Lee, H. & Seok, C. (2014) Prediction of protein structure and interaction by GALAXY protein modeling programs. *BioDesign*, **2**, 1-11.

Sterck, L., de María, N., Cañas, R.A., de Miguel, M., Perdiguero, P., Raffin, A. et al. (2022) Maritime Pine Genomics in Focus. In: De La Torre, A.R. (eds) *The Pine Genomes. Compendium of Plant Genomes*. Springer, Cham, 67-123.

Suárez, M.F., Ávila, C., Gallardo, F., Cantón, F.R., García-Gutiérrez, A.G., Claros, M.G. et al. (2002) Molecular and enzymatic analysis of ammonium assimilation in woody plants. *Journal of Experimental Botany*, **53**, 891-904. https://doi.org/10.1093/jexbot/53.370.891

Tabuchi, M., Sugiyama, K., Ishiyama, K., Inoue, E., Sato, T., Takahashi, H. et al. (2005) Severe reduction in growth rate and grain filling of rice mutants lacking *OsGS1;1* a cytosolic glutamine synthetase1;1. *The Plant Journal*, **42**, 641-651. https://doi.org/10.1111/j.1365-313X.2005.02406.x

Tamura, K., Stecher, G. & Kumar, S. (2021) MEGA11: molecular evolutionary genetics analysis version 11. *Molecular Biology and Evolution*, **38**, 3022-3027. https://doi.org/10.1093/molbev/msab120

Tegeder, M. & Masclaux-Daubresse, C. (2017) Source and sink mechanisms of nitrogen transport and use. *New Phytologist*, **217**, 35-53. https://doi.org/10.1111/nph.14876

Thomsen, H.C., Eriksson, D., Møller, I.S. & Schjoerring, J.K. (2014) Cytosolic glutamine synthetase: a target for improvement of crop nitrogen use efficiency? *Trends in Plant Science*, **19**, 656–663. https://doi.org/10.1016/j.tplants.2014.06.002

Tian, F., Yang, D.C., Meng, Y.Q., Jin, J. & Gao, G. (2020) PlantRegMap: charting functional regulatory maps in plants. *Nucleic Acids Research*, **48**, D1104-D1113. https://doi.org/10.1093/nar/gkz1020

Urriola, J. & Rathore, K.S. (2015) Overexpression of a glutamine synthetase gene affects growth and development in sorghum. *Transgenic Research*, **24**, 397-407. https://doi.org/10.1007/s11248-014-9852-6

Valderrama-Martín, J.M., Ortigosa, F., Ávila, C., Cánovas, F.M., Hirel, B., Cantón, F.R. et al. (2022) A revised view on the evolution of glutamine synthetase isoenzymes in plants. *The Plant Journal*, **110**, 946-960. https://doi.org/10.1111/tpj.15712

Varadi, M., Anyango, S., Deshpande, M., Nair, S., Natassia, C., Yordanova, G. et al. (2022) AlphaFold protein structure database: massively expanding the structure coverage of protein-sequence space with high-accuracy models. *Nucleic Acids Research*, **50**, D439-D444. https://doi.org/10.1093/nar/gkab1061

Villalobos, D.P. (2008) Aproximación genómica al estudio de la formación de la madera en los pinos. Universidad de Málaga, Málaga (Spain). Doctoral dissertation.

Wallsgrove, R.M., Turner, J.C., Hall, N.P., Kendall, A.C. & Bright, S.W.J. (1987) Barley mutants lacking chloroplast glutamine synthetase-Biochemical and genetic analysis. *Plant Physiology*, **83**, 155-158. https://doi.org/10.1104/pp.83.1.155

Wedler, F.C. & Horn, B.R. (1976) Catalytic mechanisms of glutamine synthetase enzymes. Studies with analogs of possible intermediates and transition states. *Journal of Biological Chemistry*, **251**, 7530-7538. https://doi.org/10.1016/S0021-9258(17)32883-1

Wei, Y., Xiong, S., Zhang, Z., Meng, X., Wang, L., Zhang, X. et al. (2021) Localization, gene expression, and functions of glutamine synthetase isoenzymes in wheat grain (*Trticum*

aestivum L.). *Frontiers in Plant Science*, **12**, 580405. https://doi.org/10.3389/fpls.2021.580405

Wudick, M.M., Portes, M.T., Michard, E., Rosas-Santiago, P., Lizzio, M.A., Nunes, C. O. et al. (2018) CORNICHON sorting and regulation of GLR channels underlie pollen tube Ca^{2+} homeostasis. *Plant Science*, **360**, 533-536. https://doi.org/10.1126/science.aar6464

Yadav, S.K. (2009) Computational structural analysis and kinetic studies of a cytosolic glutamine synthetase from *Camellia sinensis* (L.) O. Kuntze. *Protein Journal*, **28**, 428-434. https://doi.org/10.1007/s10930-009-9210-3

Yamaya, T. & Kusano, M. (2014) Evidence supporting distinct functions of three cytosolic glutamine synthetases and two NADH-glutamate synthases in rice. *Journal of Experimental Botany*, **65**, 5519-5525. https://doi.org/10.1093/jxb/eru103

Zhao, W., Yang, J., Tian, Y., Fu, X., Zhu, B., Xue, Y. et al. (2014) Expression, purification, and characterization of recombinant mangrove glutamine synthetase. *Molecular Biology Reports*, **41**, 7575-7583. https://doi.org/10.1007/s11033-014-3649-9

Zhou, J.Y., Hao, D.L. & Yang, G.Z. (2021) Regulation of cytosolic pH: the contributions of plant plasma membrane H^+-ATPases and multiple transporters. *International Journal of Molecular Sciences*, 22, 12998. https://doi.org/10.3390/ijms222312998

TABLES

Table 1. Physicochemical properties of the wild type PpGS1b.1 and PpGS1b.2 and their mutated versions.

Property	GS1b.1WT	GS1b.2WT	GS1b.1K264E	GS1b.2E264K	GS1b.1K267H	GS1b.2H267K	GS1b.1 K264E,K267H	GS1b.2 E264K,H267K
Optimum temperature	42	42	45	30	45	45	45	45
Optimum pH	6.5	6	6.5	7	6.5	6	6	6
Monomer ΔG folding (kcal/mol)	-9.7	-2.7	-10.1	-2.3	-9.2	-2.7	-10.0	-2.3
Ea (kJ/mol)	39.8	46.1 / 18.7	34.8	39.9	35.2 / 6.7	33.7 / 10.3	15.2	28.4 / 6.7
$\Delta G^{\ddagger}$ (kJ/mol)	63.6	62.6 / 62.8	63.5	62.3	62.9 / 62.2	63.1 / 62.4	62.7	62.1 / 62.0
$\Delta H^{\ddagger}$ (kJ/mol)	36.9	43.6 / 16.1	32.3	37.4	32.7 / 4.3	31.2 / 7.9	12.7	25.9 / 4.2
$T\Delta S^{\ddagger}$ (kJ/mol)	-26.7	-19.0 / -46.7	-31.2	-24.9	-30.2 / -57.9	-31.9 / -54.5	-50.0	-36.2 / -57.8
Break (ºC)		24			34	35		27.5

Table 2. Kinetic properties of the wild type PpGS1b.1 and PpGS1b.2 and their mutated versions.

Substrate	GS1b.1WT	GS1b.2WT	GS1b.1K264E	GS1b.2E264K	GS1b.1K267H	GS1b.2H267K	GS1b.1 K264E,K267H	GS1b.2 E264K,H267K
NH_4^+	Vmax 38.15	Vmax 6.48	Vmax 21.01	Vmax 16.36	Vmax 79.3	Vmax 58.26	Vmax 36.05	Vmax 56.88
	Km 0.12	Km 0.21	Km 0.16	Km 0.16	Km 0.06	Km 0.05	Km 0.02	Km 0.09
		Ki 22.57			Ki 13.14			
Glu	Vmax 101.6	Vmax 7.66	Vmax 16.82	Vmax 15.69	Non saturated	Non saturated	Non saturated	Non saturated
	Km 64.18	nH 1.311	Km 2.20	Km 18.41				
	Ki 84.51	EC50 48.63						
Mg^{2+}	Vmax 71.32	Vmax 5.64	Non saturated	Vmax 25.43	Non saturated	Non saturated	Non saturated	Non saturated
	nH 1.66	nH 2.33		nH 1.61				
	EC50 14.49	EC50 10.87		EC50 26.78				
ATP	Vmax 24.96	Vmax 7.39	Vmax 29.73	Vmax 27.40	Vmax 59.82	Vmax 28.57	Vmax 38.69	Vmax 100.6
	Km 0.18	Km 0.29	Km 0.11	Km 0.21	Km 0.06	Km 0.39	Km 0.09	Km 0.42
	Ki 5.88		Ki 3.19	Ki 8.23	Ki 5.85	Ki 8.76		Ki 5.06

EC50 in mM; Ki in mM; Km in mM; nH is dimensionless; Vmax in nkat/mg protein. EC50, the concentration of substrate that produces a half-maximal reaction rate; Ki, dissociation constant for substrate binding; Km, Michaelis-Menten constant; nH, Hill slope; Vmax, maximal reaction rate.

FIGURE LEGENDS

Figure 1. Protein alignment and evolutionary analysis by Maximum Likelihood method. A. Protein alignment of maritime pine GSs. PpGS1a sequence is showed as reference, dots highlight conserved residues in the three sequences. **B.** The evolutionary history was inferred by using the Maximum Likelihood method and JTT matrix-based model (Jones et al., 1992). The tree with the highest log likelihood (-12198.89) is shown. The percentage of trees in which the associated taxa clustered together is shown next to the branches. Initial tree(s) for the heuristic search were obtained automatically by applying Neighbor-Join and BioNJ algorithms to a matrix of pairwise distances estimated using the JTT model, and then selecting the topology with superior log likelihood value. The tree is drawn to scale, with branch lengths measured in the number of substitutions per site. This analysis involved 96 amino acid sequences. All positions containing gaps and missing data were eliminated (complete deletion option). There was a total of 348 positions in the final dataset. Evolutionary analyses were conducted in MEGA11 (Tamura et al., 2021). Numbers close to the branches shown bootstrap values. The first two letters of the sequence names correspond to the genera and species listed in Table S1. Golden tree branches correspond to GS2 sequences; blue branches to GS1a sequences; and red branches to GS1b sequences. Discontinuous lines in GS1b branches highlight the new sequences found in *Pinus* and *Picea* genera. Red dots shown the sequences from *Pinus* and *Picea* genera.

Figure 2. *GS* gene expression in maritime pine seedlings. A. Expression levels of *GS* genes of maritime pine during germination and initial seedling development. Stage 1 (S1) corresponds to seedlings with active mobilization of reserves from megagametophyte to the seedling (one-week-old from emergence). Stage 2 (S2) corresponds to seedlings without megagametophyte and developing the first new needles (one-month-old from emergence). **B.** Gene expression levels of GS in tissues from one-month-old seedlings (Cañas et al., 2017). AM, shoot Apical Meristem; EN, Emerging Needles; YNM, Young Needles Mesophyll tissue; YNV, Young Needles Vascular tissue; CM, Cotyledon Mesophyll tissue; CV, Cotyledon Vascular tissue; HC, Hypocotyl Cortex; HV, Hypocotyl Vascular tissue; HP, Hypocotyl Pith; RC, Root Cortex; RV, Root Vascular tissue; DRC, Root Developing Cortex; DRV, Root Developing Vascular tissue; RM, Root apical Meristem. Letters above

the columns highlight the statistical significance ($P<0.05$) in a Tukey post-hoc test after an ANOVA analysis. Error bars show SE with n=3.

Figure 3. Seasonal *GS* expression profiles in pine needles from adult trees. A. Expression levels of *GS* genes were determined in needles from maritime pine along a year. Each needle whorl corresponds to the annual growth of a single year, the whorls were named by numbers, from 0 to 3 being this the oldest whorl. The whorl 0 corresponds to needles emerged in the same year of harvesting. For supplementary information see Cañas et al. (2015). Asterisks above the data points highlight the statistical significance ($P<0.05$) between needle whorls in a specific month in a Tukey post-hoc test after an ANOVA analysis. Error bars show SE with n=3. **B.** Expression levels of *GS* genes in buds and developing needles during the first 21 days of emergence. Letters above the data points highlight the statistical significance ($P<0.05$) in a Tukey post-hoc test after an ANOVA analysis. Error bars show SE with n=3. **C.** Picture of buds and male strobilus in April during the first harvesting time. **D.** Picture of buds and emerging needles in May at four harvesting point (21 days).

Figure 4. *GS* expression levels in different developing tissues. A. Gene expression levels of *PpGS1a*, *PpGS1b.1* and *PpGS1b.2* in different tissues of adult trees: juvenile and mature phloem; juvenile and mature xylem; and male and female strobili. **B.** Gene expression of *PpGS1b.1* and *PpGS1b.2* in different parts of the root from one-month-old seedlings: primary root, lateral roots, and root tip. **C.** Genes expression of *PpGS1b.1* and *PpGS1b.2* in different developmental stages of zygotic embryos: PC (pre-cotyledonary stage); EC (early cotyledonary stage), C (cotyledonary stage) and M (mature embryo). Letters above the columns highlight the statistical significance ($P<0.05$) in a Tukey post-hoc test after an ANOVA analysis. Error bars show SE with n=3.

Figure 5. Enzymatic characterization of recombinant GS1b.1 and GS1b.2 isoforms. A. Comparison of GS1b.1 and GS1b.2 subunit structure GSb1.1 is represented in green and GS1b.2 in cyan. The region that presented most differences between GS1b.1 and GS1b.2 (amino acids from 125 to 150) are represented in red and pink respectively. **B.** Root mean square deviation (RMSD) values between GS1b.1 and GS1b.2 monomers structure. **C.** Enzyme activity at different assay pH (from 4.5 to 10) for GS1b.1 (red line) and GS1b.2

(blue line). **D.** Enzyme activity at different assay temperature (from 5 to 70°C) for GS1b.1 (red line) and GS1b.2 (blue line). **E.** Thermal stability of GS1b.1 and GSb1.2 at different temperatures (37, 42, 45, 53 and 60°C) after different preincubation times (from 0 to 20 min). **F.** Kinetics of GS1b.1 and GS1b.2 for ammonium. **G.** Kinetics of GS1b.1 and GS1b.2 for glutamate. Error bars show the SD. Mean values are composed with at least three independent determinations.

Figure 6. Characterization of mutated GS1b.1 and GS1b.2 proteins. Disposition of the amino acids, either those that have been exchanged and those associated with them by hydrogen bonds in the GS1b.1 K264E (**A**), GS1b.2 E264K (**B**), GS1b.1 K267H (**C**) and GS1b.2 H267K (**D**) mutants. Alpha carbons of the amino acids are represented in pink. **E.** Amino acid region affected by mutations. Subunit structure of the GS1b.1 (**F**) and GS1b.2 (**G**) double mutant. Amino acids exchanged and amino acids associated with them by hydrogen bonds are represented in dark magenta. Amino acids from 330 to the end of the protein are represented in green. **H.** Comparison of the physicochemical properties of the GS1b.1 WT and its double mutant. **I.** Comparison of the physicochemical properties of the GS1b.2 WT and its double mutant. **J.** Thermal stability of the double mutants at different temperatures (37, 42, 45, 53 and 60°C) after different preincubation times (from 0 to 20 min). **K.** Kinetics of GS1b.1 and GS1b.2 double mutants for ammonium. Error bars show the SD. Mean values are composed with at least three independent determinations.

A

PpGS1a

PpGS1b.1

PpGS1b.2

Whorl 0
Whorl 1
Whorl 2
Whorl 3

B

PpGS1a

PpGS1b.1

PpGS1b.2

Bud
Needles

C

Bud

Male strobilus

D

Bud

Emerging needle

A

B

C

D

E

PpGS1b.1 GG IKVIK KAIEK
PpGS1b.2 GG IEVIH KAIEK

264   267

F

G

H

I

J GS1b.1 K264E,K267H

GS1b.2 E264K,H267K

K GS1b.1 K264E,K267H   GS1b.2 E264K,H267K