Contrasted reaction norms of wheat yield in pure vs mixed stands explained by tillering plasticities and shade avoidance

Authors: Meije Gawinowski[1,2,3], Jérôme Enjalbert[1], Paul-Henry Cournède[2], Timothée Flutre[1]

[1] Université Paris-Saclay, INRAE, CNRS, AgroParisTech, GQE - Le Moulon, 91190 Gif-sur-Yvette, France
[2] Université Paris-Saclay, CentraleSupélec, MICS, 91190 Gif-sur-Yvette, France
[3] Université Paris Cité, 75006 Paris, France

Corresponding authors:
Meije Gawinowski, meije.gawinowski@inrae.fr
Timothée Flutre, timothee.flutre@inrae.fr

Abstract

- **Context**

Mixing cultivars is an agroecological practice of crop diversification, increasingly used for cereals. The yield of such cereal mixtures is higher on average than the mean yield of their components in pure stands, but with a large variance. The drivers of this variance are plant-plant interactions leading to different plant phenotypes in pure and mixed stands, i.e phenotypic plasticity.

- **Objectives**

The objectives were (i) to quantify the magnitude of phenotypic plasticity for yield in pure versus mixed stands, (ii) to identify the yield components that contribute most to yield plasticity, and (iii) to link such plasticities to differences in functional traits, height and earliness.

- **Methods**

A new experimental design based on a precision sowing allowed phenotyping each cultivar in mixture, at the level of individual plants, for above-ground traits throughout growth. Eight commercial cultivars of *Triticum aestivum* L. were grown in pure and mixed stands in field plots repeated for two years (2019-2020, 2020-2021) with contrasted climatic conditions and with nitrogen fertilization, fungicide and weed removal management strategies. Two quaternary mixtures were assembled with cultivars contrasted either for height or earliness.

- **Results**

Compared to the average of cultivars in pure stands, the height mixture strongly underyielded over both years (-29%) while the earliness mixture overyielded the second year (+11%) and slightly underyielded the first year (-8%). The second year, the magnitude of plasticity in grain weight was significantly and positively associated with their relative yield differences in pure stands (R^2=0.51). The plasticity in spike number was the strongest contributor to yield plasticity (~56%), driven by even stronger but opposed plasticities in both tiller emission and regression. For both years, the plasticity in tiller emission was significantly, positively associated with the height differentials between cultivars in mixture (R^2=0.43 in 2019-2020 and 0.17 in 2020-2021).

- **Conclusions**

Plasticity in the early recognition of potential competitors for resources is a major component of genotype strategies in mixtures, as shown here for tillering dynamics. Our results also highlighted a link between plasticity in tiller emission and height differential in mixture. Both height and tillering dynamics displayed plasticities typical of the shade avoidance syndrome.

● **Implications**

Both the new experimental design and decomposition of plasticities developed in this study open avenues to better study plant-plant interactions in agronomically-realistic conditions. This study also contributed a unique, plant-level data set allowing the calibration of process-based plant models to explore the space of all possible mixtures.

Keywords: plant-plant interactions, phenotypic plasticity, tillering, cultivar mixtures, bread wheat, yield components, shade avoidance syndrome

1. Introduction

1.1. Crop diversification with cultivar mixtures and overyielding

Besides the large increase in crop production due to the massive use of pesticides and fertilizers (Evenson and Gollin, 2003), the high-input farming system of industrialized countries comes at a cost. Indeed, pesticides were found to be harmful for the environment (IPBES, 2019), human health (Expertise collective, 2022), soil (Lal, 2007) and water (Moss, 2008), endangering food security in the long term. In contrast, crop diversification is promoted in the framework of the agroecological transition (Altieri, 1989; Gliessman, 2014) by means of different practices, such as cultivar mixtures for annual crops (Barot *et al.*, 2017), with the goal of taking advantage of niche complementarity and facilitation (Brooker *et al.*, 2015).

Cultivar mixtures, a first step towards crop diversification, are known to provide, on average, advantages in terms of disease reduction (Wolfe, 1985; Mundt, 2002). The benefits on yield are usually quantified with the comparison index named "relative yield total" (RYT) which equals the total yield of the mixture divided by the yield of the cultivars in pure stands weighted by their sowing proportions in the mixture (Weigelt and Jolliffe, 2003; Reiss and Drinkwater, 2018). We focused on wheat because as the most important crop in the world, representing 15% of cultivated areas, any diversification effort will have a major impact. For the case study of winter bread wheat, Borg *et al.* (2018) showed that cultivar mixtures have a positive RYT on average, of +2% (i.e overyielding). However, they also showed that RYT displays a large variability, of ± 40%. If the mechanisms driving disease control in mixtures are well studied (Borg *et al.*, 2018; Pélissier *et al.*, 2023), how cultivars interact in mixtures under non-limiting conditions for water, nitrogen and diseases is still understudied. Indeed, there still is a large variability in RYT between mixtures, of ± 20% in the same environment (data from Forst *et*

al., 2019). Since even in non-limiting conditions for water, nitrogen and diseases, the variability of RYT is still misunderstood, we chose to focus on these conventional conditions, where light can still be a major limiting factor.

1.2. Differences in relative yields between mixtures and importance of plasticity

Deciphering the causes of RYT variability is therefore critical. First, wheat cultivars are known to display a large variability for yield (grain weight per unit area), some outperforming the others in a given environment (Slafer, Savin and Sadras, 2014). Moreover, a cultivar performance can change when observed *between* stands, in pure vs. mixed stand, whether positively or negatively. This change in performance and related traits is referred to as phenotypic plasticity, understood here as the tendency for phenotypic variation to be influenced by the environment (Hallgrimsson *et al.*, 2019). Here the change in neighboring plants between pure and mixed stands can be considered as the main environmental variable. Moreover, our reaction norm of interest here is the difference in mean plant behavior between environments (pure versus mixed stands) for a given trait and genotype (Guntrip and Sibly, 1998). Regarding the large variability in RYT, such gaps between observed and expected mixture yields correspond to phenotypic plasticity between pure and mixed stands. However, the extent of phenotypic plasticity in yield and its components, and the magnitude of plasticity differences between cultivars, remain largely unexplored, as well as the underlying traits and processes. It is also imperative to go beyond the analysis of reaction norms for single traits separately given the importance of phenotypic integration that can lead to compensation between traits (Pigliucci and Preston, 2003). Beyond plant density, grain yield is decomposed into several components, spike density, grain number per spike and thousand kernel weight, that display

contrasted phenotypic plasticities (Sadras and Slafer, 2012). Among these components, a specific importance must be given to tillering as it is the most plastic trait in response to competition for light in winter bread wheat (Lecarpentier, 2017).

1.3. Hypotheses, rationale and design

In this paper we present an original field study of bread winter wheat cultivar mixtures at plant scale, enabling us to assess the plasticity of different life-history traits, whether as the difference or the ratio of a given trait in pure vs mixed stands. The objectives were to assess the magnitude of phenotypic plasticity between stand for yield and identify the yield components contributing the most to yield plasticity, the final aim being to link such plasticities to differences between cultivars in terms of functional traits, here height and earliness.

First, we hypothesized (H1) that a cultivar's plasticity in pure vs mixed stands was associated with the difference between pure-stand yield and the mean of the pure-stand yields of the other cultivars with which it was mixed. Indeed, in field trial conditions where light is the main limiting factor, if a local cultivar had a very different pure-stand yield than the mean of the other cultivars with which it was mixed, we would expect higher yield plasticity in mixture (whether positive or negative). Second, as the presence of neighbors is perceived early by plants, notably through light-quality signals (Smith, 2000; Demotes-Mainard *et al.*, 2016), we hypothesized (H2) that differences in neighbor plant-plant interactions between pure and mixed stands were more likely to disproportionately affect the first yield component to be established, namely spike number, during the tillering dynamics. Third, we hypothesized (H3) that height differences in mixtures during the crop cycle (rather than at maturity) were positively associated with the perception of light availability, and hence correlated with plasticity in tillering dynamics. A prerequisite to quantify such plasticities in yield and its components is the identification of each cultivar in the mixture, which is challenging and rarely done. We

used genotype-centered precision sowing to address this issue. In mixed stands, each plant of

a given genotype was sowed to be surrounded by plants of other genotypes, unlike mixtures in

alternate ranks. To integrate a sufficient genetic diversity and make the continuous monitoring

of individual plants feasible, notably their tillering dynamics in both pure and mixed stands,

we designed an experiment with eight winter bread wheat genotypes. Moreover, we wanted

our experiment to be representative of French farmers cultivating mixtures. The experiments

were thus made under field conditions rather than in outdoor trays or inside a greenhouse. We

also chose to study two quaternary mixtures as it is a typical mixture size used by French

farmers (Mille *et al.*, 2006). To contribute to the questions raised by Borg *et al.* (2018), two

mixtures were assembled, one contrasted for earliness and the other for height.

2. Materials and Methods

2.1. Plant material

Among a panel of 210 European cultivars of winter bread wheat (Touzy *et al.*, 2019), eight

were chosen with contrasted heading dates and heights at maturity (all French cultivars except

one German, Table S1): Accroc, Aubusson, Bagou, Belepi, Bergamo, Boregar, Expert and

Kalahari. Two four-way mixtures were assembled, knowing that French farmers mix on

average four to five cultivars (Mille *et al.*, 2006). The first one, the earliness mixture, was

composed of Accroc, Aubusson, Bergamo and Expert, the first two having earlier heading dates

than the two others. The second one, the height mixture, was composed of Bagou, Belepi,

Boregar and Kalahari, with Belepi and Kalahari known to be taller at maturity. Both mixtures

were assembled in equi-proportions, i.e the proportion of each cultivar at sowing was 25%.

2.2. Field trial

A field trial was conducted at Le Moulon, Gif-sur-Yvette, France (latitude 48.710192, longitude 2.160042, altitude 161m), from sowing on October 30, 2019 to harvest on July 20, 2020. It was fully replicated once, with sowing on November 5, 2020, and harvest on August 2, 2021. Each year, the whole trial was a rectangle of 8m wide and 12m long, and comprised 12 "nano-plots" ($0.88m \times 2.32m = 2m^2$ for pure stands and $0.88m \times 1.68m = 1.5m^2$ for mixed stands): 8 pure stands and 4 mixed stands (two replicates per mixture). All nano-plots were sown as a regular lattice made of 12 ranks, with 30 rows for pure stands and 22 rows for mixed stands. The pure stands had more rows to allow for destructive sampling all along the crop cycle. Both row and rank spacings measured 8 cm corresponding to a sowing density of 160 seeds.m^{-2}, allowing both a strong interaction between plants, and sufficient tillering plasticity (multiple tillers per plant). The areas of the nano-plots, similarly to other studies (Chen *et al.*, 2021), were chosen to allow for the non-destructive phenotyping of individual plants in the center of the plots without damaging other plants. The spatial distribution of each mixture (Figure S1) was obtained by randomizing each genotype (Lieng, Richardt and Dodgson, 2012), and seeds were sown at the center of PVC rings of 0.2cm thickness, 5.5cm radius and 5cm depth (Figure S2) to facilitate the phenotyping and harvest of individual plants without disturbing their growth. In terms of agronomic management, seeds were treated prior to sowing with Celest Extra, and fungicides were applied during the crop cycle: the first year 250 L/ha of Sportak + 1L/ha of Tebucur on 2020/04/03, 2L/ha of Rivexo on 2020/05/18 and 1L/ha of Caramba Star on 2020/05/29, and the second year 2L/ha of Rivexo on 2021/05/12 and 1L/ha of Caramba Star on 2021/06/10. Nitrogen fertilization was applied twice, at the BBCH30 and BBCH55 stages, with respectively 70 and 40 kg/ha. The preceding crop for the 2019-2020 trial was a fallow, and it was alfalfa for the 2020-2021 trial. Corridors between nano-plots were regularly weeded by hand. Average daily temperature, rainfall and global radiation were

measured by a local weather station and monitoring data was retrieved from the INRAE

CLIMATIK platform (https://agroclim.inrae.fr/climatik/, in French) managed by the AgroClim

laboratory of Avignon, France. The average daily temperature between October and August

was 12.1°C in 2019-2020 and 10.8°C in 2020-2021, cumulative rainfall was 617 mm in 2019-

2020 and 665 mm in 2020-2021, and cumulative global radiation was 4261.22 $MJ.m^{-2}$ in 2019-

2020 and 3447.19 $MJ.m^{-2}$ in 2020-2021.

2.3. Phenotyping

For each year, from January until June, multiple traits were phenotyped once a month for a

total of six time points. In pure stands, 36 plants were destructively sampled at each time point

to record height (PH, in cm), tiller number (TN) and above-ground dry biomass (PW, in g)

after a 48h-drying at 60°C. In mixed stands, height and tiller number were recorded non-

destructively at each time point for each individual plant. Phenological stages BBCH30 (ear at

1cm) and BBCH55 (half of ear emerged above the flag leaf ligule; heading) were determined

based on plants in pure stands (Table S2). At harvest all remaining plants were sampled to

record the following variables at the individual-plant scale: for plant i, shoot dry weight (SW_i,

in g), spike dry weight (SPW_i, in g), spike number (SN_i), grain number (GN_i) and grain weight

(GW_i). From these were then computed the grain number per spike ($GNpS_i = GN_i / SN_i$) and

the thousand kernel weight ($TKW_i = GW_i / GN_i * 1000$, in g). To account for border effects,

the data from all plants sown along a border of a nano-plot, whether in pure or mixed stand,

were discarded, i.e only the data from inner plants were analyzed. As a result, at a given time

point, the data from 20 (respectively 59 to 90) plants were analyzed for each cultivar in pure

(resp. mixed) stand.

2.4. Statistical analysis

2.4.1. Replicates of mixtures

Each year, each mixture was cultivated in two replicate nano-plots. For each year separately, the effect of the replicate was assessed per genotype in each mixture with a linear regression for each of the phenotyped variables (PH and TN per time points; SW, SPW, SN, GN, GW, PW, HI, GNpS and TKW at harvest). As such replicate effects were not significant ($\alpha=5\%$) in the vast majority of the cases (data not shown), the following analyses were done using the data from both replicates without a replicate effect.

2.4.2. Yield

The yield of a given cultivar, whether in pure or mixed stand, was computed by dividing the mean grain weight per plant of this cultivar by the area of one plant in the stand computed as the inverse of the sowing density ($1/160=0.00625$ m^2). For French official variety trials, called Value for Cultivation, Use and Sustainability (VCUS) testing, the GEVES accounts for border effects using inter-row and inter-plot distances which typically increases the microplot area used for yield calculation by 20%. Hence in our case, in addition to discarding the data from border plants, the area of one (inner) plant corrected for borders equaled 0.00750 m^2.

2.4.3. Relative yield totals

The relative yield total of a given mixed stand (Reiss and Drinkwater, 2018) was computed as $RYT = Y_{mix} / (p_1 \cdot Y_{pure,1} + p_2 \cdot Y_{pure,2} + p_3 \cdot Y_{pure,3} + p_4 \cdot Y_{pure,4})$ where Y_{mix} was the grain weight per plant averaged over all plants in the mixed stand, p_g was the proportion of plants in the mixed stand being of genotype g, and $Y_{pure,g}$ was the grain weight per plant averaged over all plants in the pure stand of the given genotype. A value above 1 (respectively below 1)

indicates overyielding (respectively underyielding). The RYT was also expressed in percentage: 100 x (RYT-1), so that a positive value (respectively negative) indicates overyielding (resp. underyielding).

2.4.4. Dominance rankings between genotypes

A genotype will be said to *dominate* another within a given stand when its plant grain weight is higher, either if observed in parallel pure plots, or in the same mixture plot. Dominance rankings between genotypes of a given mixture were defined based on Tukey's range test on average plant grain weight using the R package multcomp (Hothorn, Bretz and Westfall, 2008). This test was applied per mixture and year. Pairwise differences between genotypes were declared significant if their p-values were below 0.05, followed by a letter-based representation (Piepho, 2004). Dominance rankings were established in ascending order based on this letter representation and decimal ranks were given to cultivars belonging to two groups. The same procedure was also applied on the pure-stand data for all genotypes of a given mixture.

2.4.5. Tests of phenotypic plasticity (reaction norms)

For each genotype per year, the mean equality of different traits between pure and mixed stands was tested. Our first index of phenotypic plasticity, $PPI_1(X)$, hence was the difference between the mean value of a trait X in mixed (μ_{mixed}) vs pure (μ_{pure}) stands: $PPI_1(X) = \mu_{mixed} - \mu_{pure}$. The null hypothesis was that the mean of the trait in the pure stand was equal to its mean in the mixed stand, meaning no plasticity in mixtures. Graphically, a line can be drawn from μ_{pure} to μ_{mixed} (Figure 1) and corresponds to what is called a "reaction norm". Moreover, the sign and significance of the slope of a reaction norm are the same as the sign and significance of PPI_1. The hypothesis was tested separately for different traits. A bootstrap algorithm was applied, assuming neither equal variances between pure and mixed stands nor the normality of trait

values per stand. One thousand bootstrap samples were generated to approximate the distribution of the test statistic under the null hypothesis, and a p-value was computed based on this distribution. No correction was applied to correct for multiple testing over all traits and years as justified in an exploratory context (Rothman, 1990; Heller, 2011).

2.4.6. Test of the relation between plant grain weight plasticity and plant grain weight difference between neighbors

As a second index of phenotypic plasticity PPI_2, the plant grain weight plasticity of a given cultivar f in a given year was quantified using the opposite of the relative competition intensity (fifth index indicated as nr. 3 in table 1 of Weigelt and Jolliffe, 2003), i.e., $PPI_2(GW_f) = 100 \times (GW_{mix,f} - GW_{pure,f}) / GW_{pure,f}$, where $GW_{s,f}$ is the mean grain weight of the focal cultivar f in stand s (pure or mixed) in the given year. The same year, the relative plant grain weight difference between the focal genotype and its neighbor cultivars, noted $\Delta GW_{pure,fy}$, was quantified as the pure-stand plant grain weight of the focal cultivar f in comparison with the mean of the pure-stand plant grain weights of its three neighbor cultivars in the mixture, noted $mGW_{pure,f}$, with $\{n \in G \setminus f\}$ being the set of neighbors cultivars and G the set of all cultivars in the mixture:

$$mGW_{pure,f} = 1/3 \cdot \Sigma_{n \in G \setminus f} \; GW_{pure,n}$$

$$\Delta GW_{pure,f} = 100 \times (\, | \, GW_{pure,f} - \, mGW_{pure,f} \, | \,) / mGW_{pure,f}$$

The absolute value of the second plasticity index $PPI_2(GW)$ was then regressed on the absolute value of $\Delta GW_{pure,f}$, per year, using a linear model:

$$\left|PPI_2(GW_f)\right| = \beta_0 + \beta_1 \cdot \left|\Delta GW_{pure,f}\right| + \epsilon_f \text{ with } \epsilon_f \sim N(0, \sigma^2)$$

or a quadratic model:

$$\left|PPI_2(GW_f)\right| = \beta_0 + \beta_1 \cdot \left|\Delta GW_{pure,f}\right| + \beta_2 \cdot \left|\Delta GW_{pure,f}^2\right| + \epsilon_f \text{ with } \epsilon_f \sim N(0, \sigma^2).$$

Both models were compared per year using the likelihood ratio test as computed by the anova function in R.

2.4.7. Decomposition of plant grain weight plasticity into the plasticity of its components

The grain weight of a plant i, GW_i, can be decomposed into the product SN_i x $GNpS_i$ x SKW_i using the single kernel weight $SKW_i=TKW_i/1000$. For a given genotype in a given year, we started by summing each variable over all n_s harvested plants (s indicating the stand, pure or mixed) to get $X_{s,tot}$ where X is one of GW, SN and GN, and we then computed $GNpS_{s,tot} = GN_{s,tot} / SN_{s,tot}$ and $SKW_{s,tot} = GW_{s,tot} / GN_{s,tot}$, where $SN_{s,tot}$ is the average of spike number over al n_s harvested plants. As a third index of phenotypic plasticity (PPI_3) between pure and mixed stands for a given trait of a given genotype in a given year, we used the natural log of the ratio of the trait value in mixture $X_{m,tot}$ over the trait value in pure stand $X_{p,tot}$ such as $PPI_3(X)=log(X_{m,tot}/X_{p,tot})$. This index was computed for GW, SN, GNpS and SKW. Thanks to the property of the log turning multiplication into addition, the index for plant grain weight $PPI_3(GW)$ can be decomposed as the sum $PPI_3(SN) + PPI_3(GNpS) + PPI_3(SKW)$. The contribution of each component plasticity to plant grain weight plasticity were compared based on their absolute value. As an example, $|PPI_3(X_1)| > |PPI_3(X_2)|$ was interpreted as trait X_1 plasticity contributing more than trait X_2 plasticity to grain weight plasticity. Moreover, note the non-linearity of the log, so that an absence of plant grain weight plasticity, $PPI_3(GW)=0$ may be due to $PPI_3(SN)=-1$, $PPI_3(GNpS)=+1$ and $PPI_3(SKW)=0$, in which the reduction of 63% of SN in mixture compared to pure stand is compensated by an increase of 172% of GNpS.

2.4.8. Plasticity in tiller emission and regression

Spike number (SN) was decomposed into a sum of two metrics, $SN = MTN + (-NRT)$ where MTN was the maximal number of tillers and NRT the number of regressed tillers. Mean equality of each metric between pure and mixed stands was tested for each genotype and year separately with the same bootstrap approach as for testing phenotypic plasticity with PPI_1 above (see 2.4.5.).

2.4.9. Correlation between tiller plasticity and height differential in mixture

For each year y, the height differential in mixture between genotype g and the other genotypes was computed at each date d (in days after sowing) between sowing and tiller cessation: $\Delta PH_{yg,mix}(d) = PH_{yg,mix}(d) - (\sum_{j \neq g} PH_{yj,mix}(d)) / 3$. This height differential during tiller emission, when the maximal number of tillers (MTN) is reached, was compared with the second plasticity index in pure versus mixed stands for the maximal number of tillers (MTN) computed as $PPI_2(MTN)$. Similarly, the height differential was also computed at each date d between tiller cessation and maturity. This index of height differential during tiller regression was compared with the second plasticity index in pure versus mixed stands for the number of regressed tillers (NRT) computed as $PPI_2(NRT)$. For the analysis of the tiller-regression phase, we fitted three linear models, all with the same response variable $PPI_2(NRT)$, but differing in their explanatory variables: the first model with only the height differential, the second model adding the plasticity index for tiller emission $PPI_2(MTN)$, and the third model with also the interaction between both explanatory factors. The best model was selected according to a likelihood ratio test.

Analyses were conducted in R (R Core Team, 2020) and figures were produced with the ggplot2 package (Wickham, 2009). Supplementary information, data and code that support the findings of this study are openly available at the INRAE space of the *Recherche Data Gouv* repository https://doi.org/10.57745/LZS8SU.

3. Results

3.1. Contrasted yields between the two years

The average yield in our field trials of winter bread wheat (see subsection 2.4.2) equaled 9.3 t.ha^{-1} (11.15 t.ha^{-1} before correction for border effects), in the range of yields in farmers' fields observed in Northern France (Brisson *et al.*, 2010), and it was associated with an average ear density of 593 spikes.m^{-2}. The first year (10.31 t.ha^{-1}; 630 spikes.m^{-2}) was more productive than the second one (8.32 t.ha^{-1}; 555 spikes.m^{-2}). Looking at pure stands, this yield decrease was especially strong for all four cultivars of the earliness mixture (-17%) and two cultivars (Bagou, Boregar) of the height mixture (-32%). Regarding mixed stands, the total yield of the height mixture sharply decreased the second year compared to the first (-41%) whereas the yield of earliness mixture remained stable (-1%). The lower yields in the second year were associated with lower global radiations starting after BBCH30 as well as lower rainfall between BBCH30 and BBCH77 and higher rainfall after BBCH77 (Figure S3).

3.2. Significant plant grain weight plasticity in pure versus mixed stands led to dominance relationships in mixtures

In the four mixture-year combinations, the plant grain weight observed in the mixture deviated from the expectation based on the plant grain weights in pure stands and the sowing proportions, as shown by the reaction norms between pure and mixed stands (Figure 1). Three RYTs (see subsection 2.4.3) were negative, ranging from -40% to -8%, and one was positive at +11%. The earliness mixture performed globally well, with a positive RYT the second year (ear density of 577 spikes.m^{-2} both years), whereas the height mixture was underyielding both years (with 687 spikes.m^{-2} in 2019-2020 and 529 spikes.m^{-2} in 2020-2021).

Figure 1: Reaction norms of mean grain weight per plant between pure and mixed stands per cultivar for both mixtures in both years. Vertical segments represent confidence intervals at 95%. Letters indicate pairwise differences between cultivars in pure or mixed stands. Line types indicate if the plasticity index PPI$_1$ is significantly different from zero or not.

Dominance relationships among cultivars were defined by testing pairwise plant grain weight differences and turning them into rankings (see subsection 2.4.4 and details in Table S3).

Although trivially done for the pure stands, this was made possible in the mixed stands thanks

to our experimental design. For three mixture-year combinations, all cultivars yielded similarly

in pure stands but differently in the mixture (Figure 1). In the fourth case (earliness mixture in

2020-2021), the ranking of one cultivar among the four changed between pure and mixed

stands. Nevertheless, overall, neither cultivar rankings nor mean plant grain weights were

significantly correlated between pure and mixed stands (Figure S4).

Measuring plant grain weight in mixtures per cultivar also allowed us to compute a first index

of phenotypic plasticity (PPI_1) between pure and mixed stands for each cultivar (see subsection

2.4.5). Out of the sixteen cultivar-year combinations, 7 (44%) displayed a significant plasticity

(Figure 1). Among these, only two had positive reaction norms, Aubusson and Expert in the

earliness mixture in 2020-2021, hence explaining the overyielding of this mixture.

The magnitude of plant grain weight plasticity in pure versus mixed stand for a given cultivar,

using a second phenotypic plasticity index $|PPI_2|$ was regressed on the absolute difference

between its grain weight in pure stand and the average grain weight of the other cultivars in

pure stand with which it was mixed (see subsection 2.4.6 and Figure 2). The association in the

first year was not significant at $\alpha=5\%$ (slope p-value = 0.35 and $R^2 = 0.14$). The second year

however, the association was significant at $\alpha=5\%$ (slope p-value = 0.046 and $R^2 = 0.51$) with a

quantile-quantile plot of residuals being reasonable given the small sample size. The second

year, even though a quadratic model could provide a fit visually more appealing, the linear

model was retained based on the likelihood ratio test (p-value = 0.21).

Figure 2: Plasticity index |PPI$_2$| between pure and mixed stands for plant grain weight (GW)
per cultivar and year versus the absolute difference between the pure-stand plant grain weight
of the focal cultivar and the mean of the pure-stand grain weights of the other cultivars in the
mixture. Line types indicate if the slope of the regression lines is significantly different than
zero or not.

3.3. Plasticity in both tillering emission and regression drives plasticity in spike number which explained most of grain weight plasticity

Given the cases of significant phenotypic plasticity in grain weight displayed by some cultivars,
we proposed a new way to decompose this plasticity into the phenotypic plasticity of its
components using a third index PPI$_3$ (see subsection 2.4.7, Figure 3 for 2019-2020, Figure S5

for 2020-2021). Out of the sixteen cultivar-year combinations, the seven with significant grain

weight plasticity always had a significant plasticity for at least one of its components.

Moreover, among the nine other combinations with no grain weight plasticity, five had at least

one plastic component. Overall, the plasticity in spike numbers was the main contributor, in

five out of seven cases in 2019-2020 (notably for three cultivars in the height mixture), and

also six out of eight cases in 2020-2021. Moreover, the plasticities in grain weight components

were strongly correlated, although with a change of sign between years (Figure S6).

Figure 3: Values of phenotypic plasticity (as the ratio mixed/pure) in log scale of plant grain
weight (GW) and its components, spike number (SN), grain number per spike (GNpS) and
single kernel weight (SKW) in 2019-2020. Note that the y-axis is in log scale, allowing for the
additivity of the components. Stars indicate significant differences in pure versus mixed stands
for a given variable, genotype and year.

1 Spike number is the result of tillering, a dynamical process whose particularity is to be non-

2 monotonic. Thanks to our experimental design, we were able to phenotype this process for the

3 first time at the level of individual plants in field conditions for both pure and mixed stands, as

4 exemplified for the height mixture in 2019-2020 (Figure 4; see Figure S7 for the other cases).

5
6

7
8 Figure 4: Average tillering dynamics from individual plant data all along the crop cycle for
9 each cultivar of the height mixture in 2019-2020 for both pure and mixed stands. Error bars
10 correspond to 95% confidence intervals.

11
12

In qualitative terms, the two phases of tillering dynamics, emission and regression separated

by a plateau, were clearly visible for both pure and mixed stands. However, these dynamics

looked quantitatively different between both stands. Beyond their outcome, spike number, the

tillering dynamics themselves were also plastic. Indeed, our two metrics, the maximum number

of tillers (MTN) for the emission phase and the number of regressed tillers (NRT) for the

regression phase, showed several significant cases of plasticity between stands, as exemplified

for the earliness mixture in 2019-2020 (see subsection 2.4.8, Figure 5 for the height mixture in

2019-2020 and Figure S8 for the other cases). Furthermore, the plasticities of these metrics

could compensate for each other (Figure 5), sometimes completely, hence leading to no

plasticity in the spike number, or only partially hence leading to spike number plasticity.

Reaction norms for the proportion of regressed tillers (NRT/MTN) is available in

supplementary material (Figure S9).

Figure 5: Reaction norms of number of axes per plant during tiller emission (MTN) and
regression (-NRT) between pure and mixed stands per cultivar for the earliness mixture in
2019-2020. Vertical segments represent confidence intervals at 95%. Letters indicate pairwise
differences between cultivars in pure or mixed stands. Line types indicate the statistical
significance of the reaction norms.

15 3.4. Dynamics of height differentials explain tillering plasticity

16 Knowing that tillering dynamics are influenced by the perception of a light-quality signal, we

17 monitored the height dynamics of individual plants (Figure 6 and Figure S10), testing for

18 phenotypic plasticity between pure and mixed stands at each time point (Figure S11).

Figure 6: Mean plant height (in cm) throughout growth for the different cultivars grown in
mixed stands for both mixtures in 2019-2020. Vertical segments represent confidence intervals
at 95%.

In the earliness mixture in 2019-2020, the two cultivars chosen for their earliness, Accroc and

Aubusson, were indeed taller than their neighbors, Bergamo and Expert, early in the crop cycle

because the elongation of their main stem started earlier. In the height mixture in 2019-2020

the two cultivars chosen for their higher height at maturity, Belepi and Kalahari, were indeed

taller at the end of the growth cycle.

Moreover, during the phase of tiller emission, the height differentials per cultivar (in

comparison with the others) in the mixtures were significantly associated with their plasticity

in maximum tiller number (see subsection 2.4.9 and Figure 7). The slope of the linear

regression was positive for both years: 6.742 ± 1.581 in 2019-2020 (p-value 2.69×10^{-4},

$R^2 = 0.43$) and 4.870 ± 1.957 in 2020-2021 (p-value 0.0186, $R^2 = 0.17$). We also tested this

relation during the phase of tiller regression (Figure S12) while also controlling for the

plasticity during the phase of tiller emission. For both years, the plasticity in tiller regression

was significantly associated with both height differential and plasticity on tiller emission.

2 Figure 7: Plasticity of the maximum tiller number (MTN) versus height differentials in mixture
3 computed during the tiller emission phase. Line types indicate if the slope of the regression
4 lines is significantly different than zero or not.
5

4. Discussion

We proposed a new experimental design and analytical framework to study, in field trials, the

reaction norms of cereals in pure versus mixed stands. We applied them to scrutinize the

development and productivity of eight bread wheat genotypes, in pure stand and into two

quaternary mixtures, over two years. The main advantage of the new design consisted in giving

access to the behavior of each genotype in a mixture. As a result, it allowed the study of yield

phenotypic plasticity per genotype between pure and mixed stands, as exemplified by our

results using three indices of phenotypic plasticity. We notably proposed a new index to study

such a plasticity by decomposing it into the plasticities of its components. Although our design

and framework came with limits discussed below, they delivered new results summarized right after.

4.1. Limits of the experimental design

Due to the precision of our experiment and the constraint to do non-destructive measurements in mixtures, we used PVC rings in mixtures to better separate the plants and facilitate measurements, and we still performed destructive measurements in pure stands. This could have introduced a bias in mixture measurements that we considered negligible although we did not assess it.

As motivated in the introduction, by applying nitrogen fertilization as well as fungicides, and by removing weeds manually, we focused our experiments on conditions under which light availability was the main limiting resource underlying competition between plants. Water stress was also not deemed impactful enough to require irrigation. Yet, abiotic and biotic factors have an impact on the outcome of plant-plant interactions. Resources such as nutrients and water are known to trigger competition mechanisms other than those for light (Craine and Dybzinski, 2013). As a consequence, the RYTs of specific cultivar mixtures may change from over to underyielding, and inversely, depending on the timing and intensity of such stress factors. Moreover some crop traits may be advantageous to limit weed growth (Wicks *et al.*, 2004; Lazzaro *et al.*, 2019) but, as a corollary, they may also limit the growth of other cultivars in a mixture. Overall, our results may change under strong abiotic or biotic stress, but the proposed framework could nevertheless quantify plasticity in yield components under any stress scenario.

Among the studied yield components, spike number results from tillering dynamics, known to be strongly dependent on plant density (Darwinkel, 1978). We hence chose for our field trials a density of 160 plants.m^{-2} allowing strong interactions between plants during the tillering

process while maintaining a sufficient number of spikes per plant at harvest (3.57 on average over the two years). This density is slightly lower than the range advised to French farmers (200-400 plants.m^{-2}) but it is not abnormal: Abichou, de Solan and Andrieu, (2019) qualified as "usual" densities between 170 and 250 plants.m^{-2}. Nevertheless, other plant densities are likely to have an impact on the magnitude and sign of phenotypic plasticities and this impact is worth investigating further, as well as the effect of inter-row distance (Abichou, de Solan and Andrieu, 2019). Finally, we chose eight elite cultivars to compose two quaternary mixtures but the use of more diverse genotypes (eg. landraces) and other mixtures of other orders (eg. binary) may produce different results.

4.2. Significant reaction norms of yield components between pure and mixed stands

Plant grain weight displayed a large variation between pure and mixed stands. Over the four mixture-year combinations, only one favorable case (RYT>1) was observed, contradicting the usual overyielding of mixtures on average (Borg *et al.*, 2018). The observed variation in RYT was due to the phenotypic plasticity of cultivars, with both positive and negative reaction norms from pure to mixed stand that affected the dominance relationships between cultivars. The results also confirmed our first hypothesis: the second year, the magnitude of plant grain weight plasticity for a given cultivar was significantly positively associated with the difference of its plant grain weight with the three other cultivars, observed in pure stand. The fact that this association was not significant the first year could be explained by the better yield potential of the first year meteorological conditions, leading to much smaller yield differences between cultivars in pure stands, in link with the stress gradient hypothesis (Malkinson and Tielbörger, 2010). Nevertheless, the yield difference among cultivars in pure stands can be regarded as an

indicator of asynchronous production trends, also called asynchrony, a notion with increasing interest for the assembly of cultivars in mixtures (Valencia *et al.*, 2020; Weih *et al.*, 2021). Thanks to a new decomposition, we then showed that plant grain weight plasticity resulted from the plasticities of its component that can cancel or amplify each other. We confirmed our second hypothesis of a main contribution of spike number plasticity, similarly to what was observed in rice (Jennings and Aquino, 1968). Plasticity in spike number was itself driven by plasticity in both the tiller emission and regression phases that, importantly, could compensate for each other. In agreement with our third hypothesis, the height differentials of cultivars in mixture were significantly associated with the tillering plasticities, notably for the maximal number of tillers. Finally, our results highlighted the importance of the environmental conditions, with several relationships changing sign from one year to the next.

Overall our results showed that the most integrative trait of interest, plant grain weight, can be stabilized in mixture over contrasted years, as well as its main component, spike number, through the net result of cancelation-amplification of the underlying determinants.

4.3. Plasticity in tillering

In general, phenotypic plasticity of a given trait can be classified as advantageous, neutral or disadvantageous depending on the relationships with fitness (Alpert and Simms, 2002). Moreover, the plasticity in a given trait can be qualified as *active* in the sense of "anticipatory, and often highly integrated, phenotypic changes in response to some environmental cue or signal", whereas the plasticity in another trait would be best viewed as *passive*, i.e., as a "consequence of the environment, such as stunted growth owing to low resource levels" (Forsman, 2015). In the case of small-grain cereals in general, and wheat in particular, the phenotypic plasticities of yield and its components have been widely studied, but mostly in pure stands (Sadras and Slafer, 2012; Sadras and Rebetzke, 2013; Slafer, Savin and Sadras,

2014). As a consequence, little is known in mixed stands about either the impact of competition between cultivars on these plasticities, or the other way around, i.e the impact of these plasticities on the outcomes of the competition between cultivars. Moreover, a discussion of active versus passive plasticity of each yield component is lacking, especially in heterogeneous stands such as cultivar mixtures for which our study specifically contributes.

Given the early detection of light signals (R:FR), their integration as a bet on future competition (Schmitt and Wulff, 1993) and the consequences in terms of architectural changes (Casal, 1988; Evers *et al.*, 2006; Xie, Mayes and Sparkes, 2016), plasticity in tillering can be characterized as active. As the outcome for a plant of an early bet, here on tillering, was more consequential on plant grain weight than would be a late bet, such as on thousand kernel weight, improving the odds of the bet on tillering would justify maintaining the capacity to detect variation in light cues and respond appropriately to them. Moreover, we observed a hierarchy of plant grain weight yield-component plasticities in pure versus mixed stands in agreement with known results in the literature obtained when comparing stress versus no-stress conditions (Sadras and Slafer, 2012). In the majority of cases, the contribution of the plasticity in spike number to plant grain weight plasticity was higher than the ones from the plasticities of grains per spike and thousand kernel weight. The plasticity hierarchy in plant grain weight components can hence be interpreted as reflecting differences between active and passive plasticity. Moreover, because of the processes involved in responding to specific signals, active plasticity can come at a cost (DeWitt, Sih and Wilson, 1998). Due to the sequential building of plant grain weight components in annual plants, it can further be hypothesized that the tradeoff between the cost of active plasticity and its advantages will be more beneficial for early processes than late ones, as illustrated by the tillering dynamics and the R:FR signaling.

We also observed phenotypic plasticity in height during plant growth and it displayed a convergence pattern consistent with the shade avoidance syndrome, as already documented in

barley varietal mixtures (Dahlin *et al.*, 2020). Because this syndrome is associated with more biomass being allocated to vegetative structure to the detriment of reproductive ones, such a plasticity in height is considered harmful in crops, with ongoing efforts to reduce it in wheat (Wille *et al.*, 2017). A reduced plasticity in stem elongation was for instance observed in modern, smaller cultivars in comparison to older, taller ones for oat (Semchenko and Zobel, 2005) and wheat (Weiss *et al.*, 2009). Interestingly, we observed a similar convergence pattern in terms of tillering dynamics (Figure 4) as well as a significant relationship between the plasticity in tiller emission and height differential in mixture (Figure 7). This possibly reflects a causal interplay between height and tillering due to a shared mechanism, as exemplified by the pleiotropic effect of Rht genes (Xu *et al.*, 2023).

4.4. Significance for assembly rules and breeding strategies

Regarding assembly rules, farmers often avoid mixing cultivars of different height or earliness (Borg *et al.*, 2018). However, our results show that the tallest cultivars at maturity or the earliest ones are not always dominating in the mixture. For instance, Kalahari had the highest final height in the height mixture but it did not dominate whatever the year. Similarly, Accroc and Aubusson were the earliest cultivars in the earliness mixture, especially the first year, but their dominance ranks this year were respectively second and third. Phenotypic data such as final height and heading date, easily recorded in pure stand trials and largely made available to farmers, may hence not be that informative to predict the outcome of competition in varietal mixtures, but this remains to be tested on a larger data set in terms of genotypes, mixtures and years.

Moreover, farmers in France base their choices of wheat cultivars on the results of field trials, notably with different management intensities (Loyce *et al.*, 2012), yield being a major

determinant and currently all trials only involve pure stands. As shown in Figure 2, yield differences between cultivars in pure stands could be informative of the magnitude of yield plasticity in mixtures, although the strength of the association depends on the year. The wealth of pure-stand yield data could hence be of interest to guide farmers in their assembly for mixtures. Given the potential benefits, this will have to be further tested on a larger data set.

Furthermore, our study highlights the importance of monitoring the plasticity of key traits such as tillering to decipher the behavior and performance of cultivars in mixtures. However, the observed patterns of plasticity in both tillering phases notably depended on the year (genotype-by-genotype-by-year interactions, GxGxE), which complicates the definition of mixture assembly rules. Hence, knowing the yield variability of cultivars in pure stands over various environments, although useful (Wuest, Peter and Niklaus, 2021), may not be sufficient. For instance, the plant grain weight in pure stands can be similar over years, while differing significantly in mixtures (eg.cv. Belepi and Kalahari). Complex interactions between plants in mixture and climate make mixture performances hard to predict. The plant-plant interactions in mixtures (GxG) may strongly depend on abiotic conditions (GxGxE), which leads to complex interactions hard to disentangle.

Although it may be relevant to select against plasticity in height due to the shade avoidance syndrome (Wille *et al.*, 2017), such a strategy may ultimately be counterproductive if plasticities in height and tillering have partially shared genetic architecture. Barot *et al.* (2017) suggested that plasticity in some traits may be worth keeping: tillering may be one such example.

To breed for mixture yield, Litrico and Violle (2015) proposed to maximize the phenotypic variance in agronomic traits between mixture components, while minimizing it for interaction traits. They defined agronomic traits as the ones at least partially determining agronomic value (e.g., grain yield), and interaction traits as those involved in interactions with neighbors. As

1 our study demonstrates, tillering fits both these definitions, hence questioning the applicability

2 of this framework.

4.5.   Perspectives and open questions

Because of the detailed monitoring of individual plants in mixed stands, it was only feasible to

experiment with eight cultivars and two quaternary mixtures. Despite the small number of

mixtures, the observed RYTs were representative of the variability in the literature (Borg *et al.*,

2018). Binary mixtures are expected to ease the study of interactions, at the cost of restricting

the number of observed cultivars. We made the choice to experiment with mixtures more

complex than binary ones, to stick to the reality of farmers' practices, but with more complex

interactions to disentangle. Furthermore, similarly to other studies (Dahlin *et al.*, 2020), we

ignored the below-ground organs, even though root traits can contribute to a functional

understanding of mixing effects (Montazeaud *et al.*, 2018). However, adding this layer of

phenotyping to our experimental design would have required the usage of a field-based, non-

destructive   root   phenotyping   protocol   which   remains   to   be   developed.

Our analysis of reaction norms for plant grain weight and how they are explained by phenotypic

traits such as height and earliness is strongly based on ecophysiological knowledge. Beyond

the active plasticity for tillering by the R:FR sensing, more and more other signals involved in

plant-plant communication are described (Subrahmaniam *et al.*, 2018; Ninkovic *et al.*, 2019),

and they certainly deserve better attention to understand the outcomes of plant interactions in

mixtures.

Predicting mixture productivities will require a better knowledge of plant development in both

pure and mixed stands, notably regarding the tillering process. Due to the phenotypic plasticity

between   stands,   the   genotype-by-genotype   interactions,   and   their   dependency   on

environmental conditions, observing mixtures will not be dispensable. From our results, a

challenge would consist in phenotyping at high throughput the maximum number of tillers at stage BBCH30, possibly at several densities. In pure stands a UAV-based strategy seems feasible (Wu *et al.*, 2022), but transposing it in mixture at large scale will require the development of a precision sower. Another strategy is to develop functional-structural plant models with an explicit tillering process (Blanc *et al.*, 2021), allowing an *in-silico* exploration of mixture behavior after calibration on data as the ones we produced here. Combining both strategies, i.e., UAV-based phenotyping and process-based modeling, is already feasible for wheat in pure stands (Jégo, Pattey and Liu, 2012; Yang *et al.*, 2021), but remains to be explored in mixed stands.

5. Conclusion

Crop diversification is a critical leverage for the agroecological transition, and intra-specific diversification at the field scale, using cultivar mixtures, is increasingly used by farmers to stabilize yields and control diseases. However, there still is a knowledge gap on how to design efficient crop mixtures and further investigations are needed. We therefore developed an experimental and analytical framework to better study the plant-plant interactions underlying mixture performance. Based on eight wheat cultivars, we assembled two cultivar mixtures contrasted for height and earliness. This allowed us to decipher the importance of trait plasticity, and more specifically the early and active plasticity of tillering driving spike number which is major yield component. Extending such an experimental study and coupling it to ecophysiological modeling and on-farm experiments will allow a better understanding of the complex genetic-by-environment interactions, and design resilient mixtures in a less predictable agro-climatic context.

Acknowledgements

The authors thank D. Tropée and G. Vieceli for their valuable help and support with the field experiments, as well as T. Randrianarisoa, N. Vazeux-Blumental, H. Belcram, A. Postec, L. Malicet-Chebbah, B. Rouger, E. Forst, G. Van Frank, E. Blanc, A. Merbhène, M. Turbet-Delof, I. Goldringer, A. Hospital, M. Colas, F. Ammar, A. Sidik Meite, K. Ménard and N. Tinomme. The authors thank T. Moittie (Asur Plant Breeding), M. Balduz (Lemaire Deffontaines), C. Duquet and M. Prevost (Limagrain Europe), C. Michelet (RAGT2n) and S. Caiveau (Syngenta France SAS) for providing the seeds. The authors thank R. Rincent for providing the adjusted means of heading date and final height from the BreedWheat panel. The authors thank V. Allard and P. Roumet for fruitful discussions on topics related to this work, as well as the three anonymous reviewers for their particularly constructive feedback. This work was supported by the doctoral school FIRE of the Learning Planet Institute and the BAP department from INRAE (PerfoMix project).

Literature Cited

Abichou, M., de Solan, B. and Andrieu, B. (2019) 'Architectural Response of Wheat Cultivars to Row Spacing Reveals Altered Perception of Plant Density', *Frontiers in Plant Science*, 10. Available at: https://doi.org/10.3389/fpls.2019.00999.

Alpert, P. and Simms, E.L. (2002) 'The relative advantages of plasticity and fixity in different environments: when is it good for a plant to adjust?', *Evolutionary Ecology*, 16(3), pp. 285–297. Available at: https://doi.org/10.1023/A:1019684612767.

Altieri, M.A. (1989) 'Agroecology: a new research and development paradigm for world agriculture', *Agriculture, Ecosystems & Environment*, 27(1/4), pp. 37–46.

Barot, S. *et al.* (2017) 'Designing mixtures of varieties for multifunctional agriculture with the help of ecology. A review', *Agronomy for Sustainable Development*, 37(2), p. 13. Available at: https://doi.org/10.1007/s13593-017-0418-x.

Blanc, E. *et al.* (2021) 'Functional–Structural Plant Modeling Highlights How Diversity in Leaf Dimensions and Tillering Capability Could Promote the Efficiency of Wheat Cultivar Mixtures', *Frontiers in Plant Science*, 12, p. 734056. Available at: https://doi.org/10.3389/fpls.2021.734056.

Borg, J. *et al.* (2018) 'Unfolding the potential of wheat cultivar mixtures: A meta-analysis perspective and identification of knowledge gaps', *Field Crops Research*, 221, pp. 298–313. Available at: https://doi.org/10.1016/j.fcr.2017.09.006.

Brisson, N. *et al.* (2010) 'Why are wheat yields stagnating in Europe? A comprehensive data analysis for France', *Field Crops Research*, 119(1), pp. 201–212. Available at: https://doi.org/10.1016/j.fcr.2010.07.012.

Brooker, R.W. *et al.* (2015) 'Improving intercropping: a synthesis of research in agronomy, plant physiology and ecology', *New Phytologist*, 206(1), pp. 107–117. Available at: https://doi.org/10.1111/nph.13132.

Casal, J.J. (1988) 'Light quality effects on the appearance of tillers of different order in wheat (Triticum aestivum)', *Annals of Applied Biology*, 112(1), pp. 167–173. Available at: https://doi.org/10.1111/j.1744-7348.1988.tb02052.x.

Chen, J. *et al.* (2021) 'Diversity increases yield but reduces harvest index in crop mixtures', *Nature Plants*, 7(7), pp. 893–898. Available at: https://doi.org/10.1038/s41477-021-00948-4.

Craine, J.M. and Dybzinski, R. (2013) 'Mechanisms of plant competition for nutrients, water and light', *Functional Ecology*. Edited by D. Robinson, 27(4), pp. 833–840. Available at: https://doi.org/10.1111/1365-2435.12081.

Dahlin, I. *et al.* (2020) 'Plasticity of barley in response to plant neighbors in cultivar mixtures', *Plant and Soil*, 447(1–2), pp. 537–551. Available at: https://doi.org/10.1007/s11104-019-04406-1.

Darwinkel, A. (1978) 'Patterns of tillering and grain production of winter wheat at a wide range of plant densities.', *Netherlands Journal of Agricultural Science*, 26(4), pp. 383–398. Available at: https://doi.org/10.18174/njas.v26i4.17081.

Demotes-Mainard, S. *et al.* (2016) 'Plant responses to red and far-red lights, applications in horticulture', *Environmental and Experimental Botany*, 121, pp. 4–21. Available at: https://doi.org/10.1016/j.envexpbot.2015.05.010.

DeWitt, T.J., Sih, A. and Wilson, D.S. (1998) 'Costs and limits of phenotypic plasticity', *Trends in Ecology & Evolution*, 13(2), pp. 77–81. Available at: https://doi.org/10.1016/S0169-5347(97)01274-3.

Evenson, R.E. and Gollin, D. (2003) 'Assessing the Impact of the Green Revolution, 1960 to 2000', *Science*, 300(5620), pp. 758–762. Available at: https://doi.org/10.1126/science.1078710.

Evers, J.B. *et al.* (2006) 'Cessation of Tillering in Spring Wheat in Relation to Light Interception and Red : Far-red Ratio', *Annals of Botany*, 97(4), pp. 649–658. Available at: https://doi.org/10.1093/aob/mcl020.

Expertise collective (ed.) (2022) *Effects of pesticides on health – New data*. INSERM, EDP Sciences. Available at: https://doi.org/10.1051/978-2-7598-2721-3.

Forsman, A. (2015) 'Rethinking phenotypic plasticity and its consequences for individuals, populations and species', *Heredity*, 115(4), pp. 276–284. Available at: https://doi.org/10.1038/hdy.2014.92.

Forst, E. *et al.* (2019) 'A generalized statistical framework to assess mixing ability from incomplete mixing designs using binary or higher order variety mixtures and application to wheat', *Field Crops Research*, 242, p. 107571. Available at: https://doi.org/10.1016/j.fcr.2019.107571.

Gliessman, S. (2014) *Agroecology: The Ecology of Sustainable Food Systems*. CRC Press.

Guntrip, J. and Sibly, R.M. (1998) 'Phenotypic plasticity, genotype-by-environment interaction and the analysis of generalism and specialization in Callosobruchus maculatus', *Heredity*, 81(2), pp. 198–204. Available at: https://doi.org/10.1046/j.1365-2540.1998.00354.x.

Hallgrimsson, B. *et al.* (2019) 'The developmental-genetics of canalization', *Seminars in Cell & Developmental Biology*, 88, pp. 67–79. Available at: https://doi.org/10.1016/j.semcdb.2018.05.019.

Heller, R. (2011) 'Discussion of "Multiple Testing for Exploratory Research" by J. J. Goeman and A. Solari', *Statistical Science*, 26(4). Available at: https://doi.org/10.1214/11-STS356A.

Hothorn, T., Bretz, F. and Westfall, P. (2008) 'Simultaneous Inference in General Parametric Models', *Biometrical Journal*, 50(3), pp. 346–363.

IPBES (2019) *Global assessment report on biodiversity and ecosystem services of the Intergovernmental Science-Policy Platform on Biodiversity and Ecosystem Services.* Zenodo. Available at: https://doi.org/10.5281/ZENODO.3831673.

Jégo, G., Pattey, E. and Liu, J. (2012) 'Using Leaf Area Index, retrieved from optical imagery, in the STICS crop model for predicting yield and biomass of field crops', *Field Crops Research*, 131, pp. 63–74. Available at: https://doi.org/10.1016/j.fcr.2012.02.012.

Jennings, P.R. and Aquino, R.C. (1968) 'STUDIES ON COMPETITION IN RICE. III. THE MECHANISM OF COMPETITION AMONG PHENOTYPES', *Evolution*, 22(3), pp. 529–542. Available at: https://doi.org/10.1111/j.1558-5646.1968.tb03990.x.

Lal, R. (2007) 'Anthropogenic Influences on World Soils and Implications to Global Food Security', in *Advances in Agronomy*. Elsevier, pp. 69–93. Available at: https://doi.org/10.1016/S0065-2113(06)93002-8.

Lazzaro, M. *et al.* (2019) 'Unraveling diversity in wheat competitive ability traits can improve integrated weed management', *Agronomy for Sustainable Development*, 39(1), p. 6. Available at: https://doi.org/10.1007/s13593-018-0551-1.

Lecarpentier, C. (2017) *Plasticité de l'architecture aérienne du blé en réponse à la compétition pour la lumière au sein de cultures pures ou d'associations variétales : caractérisation expérimentale et développement d'un modèle 3D*. Paris-Saclay.

Lieng, H., Richardt, C. and Dodgson, N.A. (2012) 'Random Discrete Colour Sampling'. The Eurographics Association. Available at: https://doi.org/10.2312/COMPAESTH/COMPAESTH12/081-087.

Litrico, I. and Violle, C. (2015) 'Diversity in Plant Breeding: A New Conceptual Framework', *Trends in Plant Science*, 20(10), pp. 604–613. Available at: https://doi.org/10.1016/j.tplants.2015.07.007.

Loyce, C. *et al.* (2012) 'Growing winter wheat cultivars under different management intensities in France: A multicriteria assessment based on economic, energetic and environmental indicators', *Field Crops Research*, 125, pp. 167–178. Available at: https://doi.org/10.1016/j.fcr.2011.08.007.

Malkinson, D. and Tielbörger, K. (2010) 'What does the stress-gradient hypothesis predict? Resolving the discrepancies', *Oikos*, 119(10), pp. 1546–1552. Available at: https://doi.org/10.1111/j.1600-0706.2010.18375.x.

Mille, B. *et al.* (2006) 'Assessing four-way mixtures of winter wheat cultivars from the performances of their two-way and individual components', *European Journal of Plant Pathology*, 114(2), pp. 163–173. Available at: https://doi.org/10.1007/s10658-005-4036-0.

Montazeaud, G. *et al.* (2018) 'Crop mixtures: does niche complementarity hold for belowground resources? An experimental test using rice genotypic pairs', *Plant and Soil*, 424(1–2), pp. 187–202. Available at: https://doi.org/10.1007/s11104-017-3496-2.

Moss, B. (2008) 'Water pollution by agriculture', *Philosophical Transactions of the Royal Society B: Biological Sciences*, 363(1491), pp. 659–666. Available at: https://doi.org/10.1098/rstb.2007.2176.

Mundt, C.C. (2002) 'Performance of wheat cultivars and cultivar mixtures in the presence of Cephalosporium stripe', *Crop Protection*, 21(2), pp. 93–99. Available at: https://doi.org/10.1016/S0261-2194(01)00067-9.

Ninkovic, V. *et al.* (2019) 'Who is my neighbor? Volatile cues in plant interactions', *Plant Signaling & Behavior*, 14(9), p. 1634993. Available at: https://doi.org/10.1080/15592324.2019.1634993.

Pélissier, R. *et al.* (2023) 'A major genetic locus in neighbours controls changes of gene expression and susceptibility in intraspecific rice mixtures', *New Phytologist*, 238(2), pp. 835–844. Available at: https://doi.org/10.1111/nph.18778.

Piepho, H.-P. (2004) 'An Algorithm for a Letter-Based Representation of All-Pairwise Comparisons', *Journal of Computational and Graphical Statistics*, 13(2), pp. 456–466. Available at: https://doi.org/10.1198/1061860043515.

Pigliucci, M. and Preston, K. (2003) 'Phenotypic integration: studying the ecology and evolution of complex phenotypes', *Ecology Letters*, 6(3), pp. 265–272. Available at: https://doi.org/10.1046/j.1461-0248.2003.00428.x.

R Core Team (2020) *R: A Language and Environment for Statistical Computing*. Vienna, Austria: R Foundation for Statistical Computing. Available at: https://www.R-project.org/.

Reiss, E.R. and Drinkwater, L.E. (2018) 'Cultivar mixtures: a meta-analysis of the effect of intraspecific diversity on crop yield', *Ecological Applications*, 28(1), pp. 62–77. Available at: https://doi.org/10.1002/eap.1629.

Rothman, K.J. (1990) 'No adjustments are needed for multiple comparisons', *Epidemiology (Cambridge, Mass.)*, 1(1), pp. 43–46.

Sadras, V.O. and Rebetzke, G.J. (2013) 'Plasticity of wheat grain yield is associated with plasticity of ear number', *Crop and Pasture Science*, 64(3), p. 234. Available at: https://doi.org/10.1071/CP13117.

Sadras, V.O. and Slafer, G.A. (2012) 'Environmental modulation of yield components in cereals: Heritabilities reveal a hierarchy of phenotypic plasticities', *Field Crops Research*, 127, pp. 215–224. Available at: https://doi.org/10.1016/j.fcr.2011.11.014.

Schmitt, J. and Wulff, R.D. (1993) 'Phytochrome and Plant Competition', *Trends in Ecology and Evolution*, 8(2), pp. 47–51.

Semchenko, M. and Zobel, K. (2005) 'The effect of breeding on allometry and phenotypic plasticity in four varieties of oat (Avena sativa L.)', *Field Crops Research*, 93(2–3), pp. 151–168. Available at: https://doi.org/10.1016/j.fcr.2004.09.019.

Slafer, G.A., Savin, R. and Sadras, V.O. (2014) 'Coarse and fine regulation of wheat yield components in response to genotype and environment', *Field Crops Research*, 157, pp. 71–83. Available at: https://doi.org/10.1016/j.fcr.2013.12.004.

Smith, H. (2000) 'Phytochromes and light signal perception by plants—an emerging synthesis', *Nature*, 407(6804), pp. 585–591. Available at: https://doi.org/10.1038/35036500.

Subrahmaniam, H.J. *et al.* (2018) 'The genetics underlying natural variation of plant-plant interactions, a beloved but forgotten member of the family of biotic interactions', *The Plant Journal*, 93(4), pp. 747–770. Available at: https://doi.org/10.1111/tpj.13799.

Touzy, G. *et al.* (2019) 'Using environmental clustering to identify specific drought tolerance QTLs in bread wheat (T. aestivum L.)', *Theoretical and Applied Genetics*, 132(10), pp. 2859–2880. Available at: https://doi.org/10.1007/s00122-019-03393-2.

Valencia, E. *et al.* (2020) 'Synchrony matters more than species richness in plant community stability at a global scale', *Proceedings of the National Academy of Sciences*, 117(39), pp. 24345–24351. Available at: https://doi.org/10.1073/pnas.1920405117.

Weigelt, A. and Jolliffe, P. (2003) 'Indices of plant competition', *Journal of Ecology*, 91(5), pp. 707–720. Available at: https://doi.org/10.1046/j.1365-2745.2003.00805.x.

Weih, M. *et al.* (2021) 'Grain Yield Stability of Cereal-Legume Intercrops Is Greater Than Sole Crops in More Productive Conditions', *Agriculture*, 11(3), p. 255. Available at: https://doi.org/10.3390/agriculture11030255.

Weiss, A. *et al.* (2009) 'Quantifying phenotypic plasticity using genetic information for simulating plant height in winter wheat', *NJAS: Wageningen Journal of Life Sciences*, 57(1), pp. 59–64. Available at: https://doi.org/10.1016/j.njas.2009.10.001.

Wickham, H. (2009) *ggplot2: Elegant Graphics for Data Analysis*. Springer-Verlag New York. Available at: http://ggplot2.org.

Wicks, G.A. *et al.* (2004) 'Winter Wheat Cultivar Characteristics Affect Annual Weed Suppression', *Weed Technology*, 18(4), pp. 988–998. Available at: https://doi.org/10.1614/WT-03-158R1.

Wille, W. *et al.* (2017) 'Reducing shade avoidance responses in a cereal crop', *AoB PLANTS*. Edited by M.B. Jackson, 9(5). Available at: https://doi.org/10.1093/aobpla/plx039.

Wolfe, M.S. (1985) 'The Current Status and Prospects of Multiline Cultivars and Variety Mixtures for Disease Resistance', *Annual Review of Phytopathology*, 23(1), pp. 251–273. Available at: https://doi.org/10.1146/annurev.py.23.090185.001343.

Wu, F. *et al.* (2022) 'Estimation of Winter Wheat Tiller Number Based on Optimization of Gradient Vegetation Characteristics', *Remote Sensing*, 14(6), p. 1338. Available at: https://doi.org/10.3390/rs14061338.

Wuest, S.E., Peter, R. and Niklaus, P.A. (2021) 'Ecological and evolutionary approaches to improving crop variety mixtures', *Nature Ecology & Evolution* [Preprint]. Available at: https://doi.org/10.1038/s41559-021-01497-x.

Xie, Q., Mayes, S. and Sparkes, D.L. (2016) 'Optimizing tiller production and survival for grain yield improvement in a bread wheat × spelt mapping population', *Annals of Botany*, 117(1), pp. 51–66. Available at: https://doi.org/10.1093/aob/mcv147.

Xu, D. *et al.* (2023) 'Dissecting pleiotropic functions of the wheat Green Revolution gene *Rht-B1b* in plant morphogenesis and yield formation', *Development*, 150(20), p. dev201601. Available at: https://doi.org/10.1242/dev.201601.

Yang, T. *et al.* (2021) 'Plant phenomics & precision agriculture simulation of winter wheat growth by the assimilation of unmanned aerial vehicle imagery into the WOFOST model', *PLOS ONE*. Edited by A. Chemura, 16(10), p. e0246874. Available at: https://doi.org/10.1371/journal.pone.0246874.

Supplementary Material

Table S1

Mixture	Cultivar	Maintainer (registration year, country)	Heading (DOY)	Final height (cm)	Awns
Earliness mixture	Accroc	RAGT (2010, FR)	128.4	87.02	Yes
	Aubusson	Limagrain (2001, FR)	129.9	87.03	No
	Bergamo	RAGT (2012, FR)	145.3	92.17	No
	Expert	Syngenta (2007, FR)	144.1	90.93	No
Height mixture	Bagou	Asur Plant Breeding (2007, FR)	136.7	82.97	No
	Belepi	Lemaire Deffontaines (2013, FR)	138.1	100.93	No
	Boregar	RAGT (2008, FR)	140.7	84.08	Yes
	Kalahari	Limagrain (2010, DE)	141.9	115.19	No

Table S1 - Composition of both mixtures, as well as heading date (since January 1, DOY=day of the year) and height at maturity of the cultivars as obtained from a field trial from the BreedWheat project.

Table S2

Mixture	Cultivar	Sowing date		Ear 1cm date		Heading date		Harvest date	
		2019	2020	2020	2021	2020	2021	2020	2021
Earliness mixture	Accroc	10/31	11/05	03/10	04/13	04/28	05/25	07/20	08/02
	Aubusson			03/14	04/09	05/02	05/31		
	Bergamo			03/27	04/13	05/08	05/31		
	Expert			04/03	04/17	05/19	05/31		
Height mixture	Bagou			03/17	04/13	05/19	06/02		
	Belepi			04/08	04/20	05/17	06/02		
	Boregar			04/11	04/21	05/18	06/02		
	Kalahari			03/28	04/19	05/17	06/02		

4
5
6 Table S2 – Dates of sowing and harvest for plants grown in pure and mixed stands, as well as
7 the dates of the stages ear 1cm (BBCH30) and heading (BBCH55) for plants grown in pure
8 stands (format Month/Day).
9
10

Table S3

Mixture	Year	Cultivar	Letters from the Tukey tests		Dominance rank in mixed stands	Dominance status in mixed stands
			pure	mixed		
Earliness mixture	2019-2020	Accroc	a	b	2	dominated
		Aubusson	a	ab	1.5	dominating
		Bergamo	a	b	2	dominated
		Expert	a	a	1	dominating
	2020-2021	Accroc	a	c	3	dominated
		Aubusson	a	a	1	dominating
		Bergamo	a	bc	2.5	dominated
		Expert	a	ab	1.5	dominating
Height mixture	2019-2020	Bagou	a	b	2	dominating
		Belepi	a	a	1	dominating
		Boregar	a	c	3	dominated
		Kalahari	a	bc	2.5	dominated
	2020-2021	Bagou	b	a	1	dominating
		Belepi	a	a	1	dominating
		Boregar	b	b	2	dominated
		Kalahari	ab	ab	1.5	~dominated

Table S3 - Table with Tukey classes and matching dominance ranks for each genotype over the two growing years.

Figure S1

	1	2	3	4	5	6	7	8	9	10	11	12	
C	Ber	Aub	Acc	Ber	Exp	Acc	Aub	Exp	Aub	Ber	Ber	Exp	22
D	Exp	Exp	Acc	Aub	Aub	Exp	Ber	Acc	Exp	Aub	Acc	Exp	21
E	Acc	Aub	Exp	Exp	Ber	Acc	Aub	Aub	Ber	Aub	Exp	Ber	20
F	Exp	Acc	Ber	Aub	Aub	Exp	Acc	Ber	Exp	Acc	Aub	Acc	19
G	Acc	Acc	Aub	Exp	Acc	Ber	Exp	Aub	Acc	Exp	Ber	Exp	18
H	Exp	Ber	Ber	Aub	Aub	Acc	Exp	Ber	Acc	Exp	Acc	Aub	17
A	Aub	Acc	Exp	Acc	Ber	Exp	Aub	Acc	Aub	Acc	Ber	Exp	16
B	Ber	Acc	Acc	Exp	Ber	Acc	Aub	Exp	Ber	Acc	Aub	Ber	15
C	Aub	Exp	Ber	Aub	Aub	Ber	Exp	Aub	Aub	Acc	Aub	Exp	14
D	Ber	Aub	Exp	Acc	Exp	Aub	Ber	Exp	Exp	Acc	Ber	13	
E	Acc	Acc	Aub	Ber	Acc	Acc	Aub	Aub	Acc	Aub	Exp	12	
F	Exp	Aub	Ber	Aub	Aub	Exp	Ber	Acc	Exp	Acc	Ber	11	
G	Ber	Acc	Acc	Acc	Exp	Aub	Acc	Ber	Aub	Ber	Exp	10	
H	Exp	Acc	Ber	Aub	Ber	Exp	Ber	Aub	Aub	Acc	Ber	9	
A	Aub	Exp	Exp	Acc	Acc	Exp	Aub	Exp	Acc	Ber	Aub	Exp	8
B	Ber	Ber	Aub	Ber	Aub	Ber	Acc	Acc	Exp	Aub	Acc	Ber	7
C	Acc	Acc	Exp	Acc	Acc	Exp	Ber	Exp	Ber	Ber	Acc	Aub	6
D	Ber	Exp	Ber	Aub	Exp	Aub	Aub	Acc	Aub	Ber	Ber	Exp	5
E	Aub	Acc	Aub	Aub	Ber	Acc	Ber	Acc	Exp	Acc	Aub	Exp	4
F	Acc	Ber	Exp	Acc	Exp	Aub	Acc	Ber	Exp	Aub	Ber	Exp	3
G	Exp	Aub	Aub	Acc	Aub	Ber	Exp	Aub	Acc	Exp	Acc	Ber	2
H	Acc	Exp	Ber	Aub	Exp	Aub	Acc	Ber	Aub	Ber	Ber	Exp	1

A) Earliness mixture   B) Height mixture

Figure S1 – Unscaled schematic spatial distribution of cultivars in "nano-plots" for A) the earliness mixture and B) the height mixture for both years. Here, ranks are from 1 to 12 (x-axis) and rows are from 1 to 22 (y-axis). In each mixture there is one color per cultivar.

Figure S2

A) At sowing   B) After emergence

Figure S2 - Images of "nano-plots" of mixtures with PVC rings in 2019-2020.

Figure S3

Figure S3 - Cumulative daily global radiation, mean temperature and rainfall for both years, in comparison with their respective average from 2000 to 2019. Ribbons correspond to approximate ranges of the stages "ear 1cm", "heading" and "maturity" that correspond respectively to BBCH30, BBCH55 and BBCH92.

Figure S4

11   A) Colour by genotype

B) Colour by plasticity significance

C) Using mean grain weight, Pearson correlations of 0.51 in 2019-2020 (p-value = 0.19)
and -0.26 in 2020-2021 (p-value = 0.54)

Figure S4 - Correlation between dominance ranks (A, B; overall Pearson correlation coefficient
at -0.19, p-value = 0.48) or mean grain weights (C; overall Pearson correlation coefficient at
0.26, p-value = 0.33) in pure versus mixed stands.

Figure S5

Figure S5: Log values of phenotypic plasticity of grain weight (GW) and its components, spike number (SN), grain number per spike (GNpS) and thousand kernel weight (TKW) in 2020-2021. Stars indicate significant differences in pure versus mixed stands for a given variable, genotype and year.

1 Figure S6

4 A) 2019-2020

B) 2020-2021

Figure S6: Distributions along the diagonal of the phenotypic plasticity in pure versus mixed
stands for grain weight (GW) and its components, spike number (SN), grain number per spike
(GNpS) and single kernel weight (SKW) in A) 2019-2020 and B) 2020-2021. Pairwise
scatterplots are below the diagonal, in green for cultivars of the earliness mixture and purple
for cultivars of the height mixture with corresponding linear regressions for each mixture and
both (in red). Pearson correlations are above the diagonal, with stars to indicate significance at
the 5% level.

1 Figure S7

2
3
4   A) Mixture 1 in 2019-2020
5

6
7   B) Mixture 1 in 2020-2021

2
3
4   C) Mixture 2 in 2020-2021
5

6   Figure S7 - Average tillering dynamics from individual plant data all along the crop cycle for
7   each cultivar of A) Earliness mixture in 2019-2020, B) Earliness mixture in 2020-2021, C)
8   Height mixture in 2020-2021 for both pure and mixed stands. Error bars correspond to 95%
9   confidence intervals.

1 # Figure S8

4   A) Mixture 1 in 2020-2021

6   B) Mixture 2 in 2019-2020

2   C) Mixture 2 in 2020-2021

4 Figure S8: Reaction norms of the maximum tiller number (MTN) and the number of regressed
5 tillers (NRT) during the emission and regression phases of axes per plant between pure and
6 mixed stands per cultivar of the earliness mixture in 2019-2020 (A), as well as the height
7 mixture in 2019-2020 (B) and 2020-2021 (C). Vertical segments represent confidence intervals
8 at 95%. Letters indicate pairwise differences between cultivars in pure or mixed stands. Line
9 types indicate the statistical significance of the reaction norms.

Figure S9

A) 2019-2020

B) 2020-2021

Figure S9: Reaction norms of the proportion of regressed tillers (pNRT) between pure and mixed stands per cultivar of the earliness and height mixtures in 2019-2020 (A) and 2020-2021 (B). Vertical segments represent confidence intervals at 95%. Letters indicate pairwise differences between cultivars in pure or mixed stands. Line types indicate the statistical significance of the reaction norms.

Figure S10

Figure S10: Plant height (in cm) throughout growth for the different cultivars grown in mixed stands for both mixtures in 2020-2021. Vertical segments represent confidence intervals at 95%.

Figure S11

A) Earliness mixture 2019-2020

B) Earliness mixture 2020-2021

C) Height mixture 2019-2020

D) Height mixture 2020-2021

Figure S11: Reaction norms of plant height all along the crop cycle between pure and mixed

stands per cultivar for both mixtures and years. Line types indicate the statistical significance

of the phenotypic plasticities. "DAS" stands for "days after sowing".

1 Figure S12

3 Figure S12: Plasticity of the number of regressed tillers (NRT) versus height differentials in
4 mixture computed during the regression phase. Line types indicate if the slopes of the
5 regression lines are significantly different than zero or not.