1 Differential retention of Pfam domains creates long-term evolutionary trends
2

3 Jenny James[1,2], Paul Nelson[1,3], Joanna Masel[1,*]
4
5 [1]Dpt. Ecology & Evolutionary Biology, University of Arizona, Tucson AZ, USA
6 [2]Department of Ecology and Genetics, Uppsala University, Sweden
7 [3]Present address: Empress Therapeutics, Cambridge MA, USA
8 [*] corresponding author

9
Abstract

Protein domains that emerged more recently in evolution have higher structural disorder and greater
clustering of hydrophobic residues along the primary sequence. It is hard to explain how selection
acting via descent with modification could act so slowly as not to saturate over the extraordinarily
long timescales over which these trends persist. Here we hypothesize that the trends were created by a
higher level of selection which differentially affects the retention probabilities of protein domains
with different properties. This hypothesis predicts that loss rates should depend on disorder and
clustering trait values. To test this, we inferred loss rates via maximum likelihood for animal Pfam
domains, after first performing a set of stringent quality control methods to reduce annotation errors.
Intermediate trait values, matching those of ancient domains, are associated with the lowest loss rates,
making our results difficult to explain with reference to previously described homology detection
biases. Our results support the hypothesis that differential domain loss slowly weeds out those protein
domains that have non-optimal levels of disorder and clustering. The same preferences also shape
differential diversification of Pfam domains, further impacting proteome composition.

24
25
Introduction

27
The possibility that evolution could result in long-term directional change over time has enduring
appeal, but there are few well-documented examples. It has proven difficult to rigorously assess
purported universal tendencies toward increases complexity and size over time (Gregory 2008;
McShea 1991). Perhaps the best documented example of a long-term trend is that some taxonomic
groups show a tendency toward increasing body size over time, aka 'Cope's rule' (Cope 1885; Heim
et al. 2015; Payne et al. 2009). Interestingly, this trend does not seem to be caused by directional
selection slowly exerting a preference for larger individuals, but rather by the differential
diversification of larger body size clades (Heim et al. 2015). This is unsurprising, because directional
selection among individuals of the same species works quickly, making it an unlikely explanation for
such a slow directional trend.
38
New examples of long-term trends were recently observed in the properties of protein domains (e.g.
intrinsic structural disorder) as a function of how much time has elapsed since their birth, with
considerable work performed to exclude artifactual explanations (Foy et al. 2019; James et al. 2021).
We find it more plausible that some bias in the evolutionary process is responsible for these trends,
rather than that the external environments shaping birth have moved in such a consistent direction
over such long timescales. One hypothesis is that regardless of time of birth, all domains were born
with properties biased toward promoting *de novo* gene birth (Wilson et al. 2017), and since then have
had different amounts of time to evolve away from that starting point, toward properties that are more
optimal for correct folding (Bucciantini et al. 2002; Chiti and Dobson 2017). This hypothesis is
supported by the fact that the amino acid frequencies characteristic of newborn animal domains also
make the expression of a random peptide more benign in *Escherichia coli* (Kosinski et al. 2022). But
it is very hard to explain why directional selection on amino acid frequencies has been so slow to take
full effect. We know that the evolution of amino acid frequencies can be rapid, as it is able to keep
pace with the rapid evolution of nucleotide composition (Brbic et al., 2015).
53
54 Here we consider the possibility that instead of being caused by bias during descent with
55 modification, long-term trends as a function of age are due to the differential retention vs. loss of

protein domains with different properties. We hypothesize that domains are born with a wide range of biochemical properties. From this initial diversity, "fitter" domains, i.e. those that duplicate more often and/or are lost less often, are more likely to have descendants in extant organisms, and thus to appear in the older domain age classes.

Under this hypothesis, we expect a protein property that shows a long-term trend with age to be predictive of loss rate. One observed trend, extending all the way back to the last universal common ancestor, is that younger domains tend to have 'clustered' rather than interspersed hydrophobic acids along their primary sequence (Foy et al. 2019; James et al. 2021). A second trend, restricted to the evolutionary history of animals, is that younger domains tend to have a higher level of intrinsic structural disorder (ISD), primarily due to trends in the frequencies of the 20 amino acids (James et al. 2021). Here we test whether ISD and hydrophobic clustering predict the loss rate of a domain.

What we are proposing is that the relevant level of selection may not be the one that selects among alternative alleles of the same gene (Fig. 1, top), but instead the level that selects among alternative domains capable of fulfilling the same cellular functions as genes duplicate, differentiate, and are lost (Fig.1, bottom). Rapid loss of recently born protein-coding genes has previously been postulated (Palmieri, Kosiol, and Schlötterer 2014; Schlötterer 2015; Tautz and Domazet-Lošo 2011). The number of domain copies appears to be surprisingly changeable, with fairly extreme expansions and contractions observed even over short evolutionary timescales (Hahn, Han, and Han 2007; Moore and Bornberg-Bauer 2012), leaving plenty of scope for differential loss to be a biologically significant influence.

Figure 1. Alternative mechanisms/levels of directional selection over evolutionary time. Directional evolution in a trait with cohort age, represented by a change from dark to pale blue, can occur either via (top) descent with modification during which selected (pale blue) alleles replace other alleles, or (bottom) differential retention in which an initial diversity of Pfams is differentially retained. Differential diversification (middle) combines tendencies for duplication and tendencies for loss. Note that this is a levels of selection situation; a Pfam is a clade of protein-coding sequences and their component alleles, and the same figure could be interpreted with individual interests in lieu of allele interests, and clade interests in lieu of Pfam interests.

We focus on protein domains rather than on complete genes. We take domain annotations from the
Pfam database (El-gebali et al. 2019), and refer to domains simply as Pfams. Protein-coding domains
are sometimes thought of as functional units of protein sequence that are able to fold independently
(Ponting and Russell 2002). However, Pfams are annotated by HMMER3 on the basis of evolutionary
rather than functional relatedness (Finn, Clements, and Eddy 2011), and as such are fundamental units
of protein sequence homology. This makes them easier to work with than genes, which commonly
contain multiple different Pfams of a variety of ages (Bagowski, Bruins, and Velthuis 2010;
Bornberg-Bauer and Albà 2013).
97
We infer loss rates by maximizing the likelihood of observed presence / absence of at least one Pfam
instance across all species in a phylogeny. As well as this central focus on total loss rates (Fig. 1,
bottom), we perform a complementary analysis of differential diversification (Fig. 1, middle) by
taking the mean number of distinct instances of Pfams within a species. Both presence/absence data
and number-of-instances data are vulnerable to both false positives and false negatives during
homology detection. Our previous age assignment procedure (James et al. 2021) used a keyword
based approach to remove Pfams that were likely incorrectly annotated in the focal genome due to
contamination (Breitwieser et al. 2019; Lu and Salzberg 2018; Salzberg 2017), a problem that if
uncorrected will result in nonsensical loss rates for affected Pfams. James et al. (2021) also removed
Pfams that exhibited an implausible phylogenetic distribution, which could have resulted from either
contamination or horizontal gene transfer. Here we go further, reducing the rate of false negatives by
checking all six reading frames of contigs for unannotated Pfam instances, and then reconsidering
whether the phylogenetic distribution of the Pfam is implausible. Second, we infer total loss rates
jointly with a rate of false positive hits. Finally, we consider the hypothesis of homology detection
bias while interpreting results.
113
114   We ask whether ISD and clustering scores predict the rate of total loss of a Pfam in a lineage, and also
115   the number of Pfam instances per species. We analyse animal lineages only, because the ISD trends as
116   a function of Pfam age found by (James et al. 2021) were specific to animals.
117

118   **Results**
119
We used the phylostratigraphy dataset of James et al., restricted to the 6841 Pfams annotated in at
least two out of our of 343 animal species, and already subject to a number of filters used by James et
al. We scanned all six reading frames of intergenic regions to find species that contained unannotated
instances of these Pfams. We also performed additional quality controls to remove likely viral or other
contaminants on the basis of incoherent distribution across the tree (see Methods), resulting in a
dataset of 6700 Pfams. We then estimated the rate of total loss of each Pfam over the animal species
in our dataset (see Methods). During this maximum likelihood procedure, we iteratively removed
Pfam presence data for clades (mostly single species) where they are more likely to be false positive
hits or horizontal gene transfer events.
129
130   The median inferred loss rate of a Pfam in the animal phylogeny was 0.0009/MY (1st and 3rd quartiles
131   of 0.00021 and 0.0035), i.e. 0.9 losses per 100,000 years. Note that this is not the rate at which a
132   single instance of a Pfam is lost, but rather the rate at which all instances of a Pfam are lost from a
133   lineage. Unsurprisingly, Pfams with a higher mean number of instances per genome tend to have a
134   lower rate of total loss (Supplementary fig. 1, adjusted R^2 = 0.30, p < 1e-16).
135

136
Figure 2: Pfam loss rates and copy number depend, often non-monotonically, on ISD and clustering
values. The lowest loss rates are seen for Pfams with a mean ISD of 0.18 (A) and mean clustering
value of 0.85 (B). Pfams with a low ISD have a higher number of instances per genome, in a more
monotonic relationship (C). Pfams with a mean clustering value of 0.85 have the highest number of
instances per genome. Each point represents a Pfam. Violin plots are a visual guide only, based on
dividing the data into groups of equal range of ISD (A,C) or clustering (B, D). Loess regression
curves are shown in dark blue, with 95% confidence intervals shown in grey. In A) and B), horizontal
bands of points near the bottom represent 0, 1, 2, etc. observed loss events, while in C) and D)
horizontal bands represent Pfams present in only a single copy per genome. In B) and D), for better
visualization, the x-axis has been truncated at 2.5.

147
<u>Pfams with a mean ISD of 0.18 are least often lost</u>
Pfams with high ISD have high loss rates (adjusted R^2 = 0.023, slope = 1.35, p = 4e-36), which is the
predicted direction in order for their relationship to explain the animal phylostratigraphy trend (Fig.
3A). Note that the Box-Cox transformation of loss rates makes the units of the slope in these linear
models hard to interpret. Interestingly, this relationship is not just non-linear, but also non-monotonic
(loess regression in Fig. 2A). A regression model that includes a breakpoint at ISD = 0.18 (95% CI
0.15-0.21), representing the minimum loss rate, explains substantially more of the variance in loss rate
among Pfams (adjusted R^2 = 0.030, slopes -2.00 and 3.86), and fits the data significantly better than a
model without a breakpoint (p = 6e-12), better than a model with zero slope for ISD values higher
than the breakpoint (p = 1e-45), and better than a model with zero slope for ISD values lower than the
breakpoint (p = 0.0002).

159
160 The existence of a breakpoint rather than a monotonic relationship contradicts predictions from the
161 homology detection bias hypothesis. That hypothesis expects overestimation of loss rates for high ISD
162 Pfams whose homologs are more difficult to detect. While this could drive the overall relationship we
163 found in which Pfams with high ISD have high loss rates, it cannot explain why loss rates also rise
164 with low ISD <0.18.

165
Our results are instead consistent with selection acting to preferentially remove Pfams at either end of
the spectrum of ISD. In further support of this, there is less variability in ISD within older Pfam

cohorts (Fig. 4A, slope = -4.1-5 ISD/MY, adjusted R^2 = 0.88, p = 3e-22, using linear regression
weighted by number of Pfams per age cohort) than within younger Pfam cohorts. Ancient Pfams
(those born over 2000MYA) that are still around today have a mean ISD of 0.21 (median 0.20), at the
upper edge of the 95% confidence interval for the breakpoint representing minimum loss rate. These
two observations further support the hypothesis that differential loss drives the long term trend in ISD
as a function of evolutionary age observed in James et al (2021) and Foy et al. (2019). The fact that
ancient Pfams may have ISD a little above the breakpoint might be noise in the data, or it might
indicate that the trend of falling ISD with age (shown in Fig. 4A) has not yet fully saturated.
176
<u>Pfams with moderately interspersed hydrophobic residues are least often lost</u>
The hydrophobic residues of younger domains are more clustered (i.e. have higher clustering) than
those of old domains (James et al. 2021). This directional trend could be explained mostly simply if
Pfams with higher clustering were differentially lost. However, as with ISD, our data strongly support
a non-monotonic relationship between clustering and loss rate (Figure 2B, losses/MY is two-
parameter Box-Cox transformed prior to analyses). The best fitting linear model for clustering
incorporated a breakpoint with a clustering value of 0.81 (95% confidence interval: 0.77 to 0.85)
corresponding to the lowest loss rate. This model has an adjusted R^2 of 0.020, and fits the data
significantly better than a model without a breakpoint (p = 1e-25), than a model with zero slope for
the lower clustering values (p = 8e-28), and than a model with zero slope for the higher clustering
values (p = 2e-29). Clustering is not known to affect homology detection, but even if it did, it is again
not clear how homology detection bias could produce a non-monotonic relationship.
189
A clustering value of 1 corresponds to a random distribution of hydrophobic residues, while the
optimal clustering value of 0.81 represents interspersion of hydrophobic residues. Similar to the ISD
results, this breakpoint value corresponds closely to the average clustering score of the most ancient
Pfams (over 2000 MY old), which have mean and median clustering scores of 0.89 and 0.88,
respectively. The fact that this is higher than the upper end of our 95% confidence interval for the
breakpoint suggests that the trend in clustering values with age has not yet saturated. The y-intercept
of Fig. 3B shows that Pfams are born with clustering scores a little above 1. Because Pfams tend to be
born with relatively random amino acid placement corresponding to a clustering score of ~1 that is so
much higher than the optimum value of 0.81, differential Pfam retention can explain the progressive
drop in clustering scores with Pfam age. In further support of the hypothesis that differential loss is
responsible for the trend in clustering with age, there is more variability in clustering within younger
Pfam cohorts (Fig. 4B, slope of standard deviation in clustering with age = -5.5 × 10^{-5}, adjusted R^2 =
0.60, p = 2e-10, using linear regression weighted by number of Pfams per age cohort).
203
<u>Results are robust to Pfam age as a confounding factor</u>
When ascertaining the causal effects of ISD and clustering on loss rates, Pfam age has the potential to
be a confounding factor in our analyses, for two reasons. Firstly, if older Pfams also have lower ISD
and less clustered hydrophobic amino acids (James et al. 2021) for reasons other than differential loss
due to these properties, this could potentially drive a spurious relationship between Pfam properties
and loss rates. As expected, we observe lower loss rates for older Pfams in our data (Adjusted R^2 =
0.050, p = 2e-76, supplementary figure 2). Secondly and more subtly, Pfams of identical age share the
same speciation events, which enable Pfam losses to be observed. This results in phylogenetic
structure in the data. A further complication is that the fact that older Pfams have survived to be
observed might create ascertainment bias toward estimated loss rates that are lower than the true loss
propensity.
215
To control for both indirect correlations and phylogenetic structure, we used mixed-effect linear
regression models, including age as a random rather than a quantitative effect. Note that each branch
of the tree corresponds to a possible Pfam age, making age a discrete quantitative variable.
Empirically, we found that random effects models are better supported model choices: for example,
for models of loss rate and ISD, R^2 = 0.33 if we include age as a random effect, and R^2 = 0.058 if we
include age as a linear predictor, where R^2 here is the conditional R^2 describing the variance explained
by the entire model. AIC scores for the respective models are 23580 and 24432. Similarly, for models

of loss rate and clustering, a model that includes age as a random effect model has $R^2 = 0.34$, while a
fixed effect model has $R^2 = 0.05$. AIC values are 23603 and 24476, respectively.
225
The non-monotonic relationship between loss rate and ISD remains statistically supported in the
random effect model ($p = 0.0003$), with the model having a marginal R^2, which represents the
variance explained in loss rate just by the fixed effect of ISD, of 0.0030, and a slope of -1.17 for
values of ISD below 0.18, and 0.64 for values above. The non-monotonic relationship with clustering
also remained supported in the random effect model ($p = 0.02$), with a marginal R^2 of 0.00054, with a
slope of -0.24 for values of clustering below 0.81, and 0.097 for values of clustering above 0.81.
While controlling for Pfam age causes marginal R^2 values to drop dramatically, the causal direction is
unclear. I.e., differences in ISD and clustering might drive changes in loss rates as a function of Pfam
age, in which case controlling for age would remove a good portion of the genuine biological effect.
However, the statistical significance we find indicates that Pfam age alone is insufficient to be the sole
driver of the dependence of loss rates on ISD and clustering.
237
<u>Differences in loss rates are small enough to produce slow change</u>
Even without controlling for phylogenetic structure/age, the effect sizes for loss and clustering on ISD
are clearly small. In most contexts, this would cast doubt on their biological relevance. However, our
motivating question is to discover what mechanism might have a small enough effect size such that its
steady action could continue for billions of years before saturating. We therefore next use simulations
to ask whether the effects of ISD and clustering on loss are of approximately the right magnitude to
explain the relationship between these variables and Pfam age, as observed in James et al. (2021).
245
We simulate a set of Pfams initialized with the distribution of ISD or clustering values observed in
young animal Pfams, and subject them to differential loss at rates given by our fitted regression
models with breakpoints. This produces directional trends in ISD and clustering on the pertinent
timescale (Fig. 3C, 3D). The relationship between ISD (Fig. 3A) and clustering (Fig. 3B) is visually
extremely similar to that generated in our simulations (Figs. 3C and 3D). Our simple single-branch
single loss event simulation ignores many complexities of the actual phylogenetic tree, but is
sufficient to show that the effect sizes of ISD and clustering on loss rates are of approximately the
right magnitude to explain phylostratigraphy trends.
254

255
Figure 3. Observed phylostratigraphy trends in ISD (A) and Clustering (B) values occur on similar
timescales as in our simulations of differential loss rates (C and D). In A) and B) each box represents
all Pfams of a particular age category, with the widths of the boxes proportional to the square root of
the number of Pfams in the group. Boxplots show the median, upper, and lower quartiles of the data,
while the whiskers represent the largest and smallest data value above and below the interquartile
range multiplied by 1.5. Grey points show the values for individual Pfams. The dark blue line is the
loess regression slope of the effect of age on ISD (A) and clustering (B), with the 95% confidence
interval shown in grey. In C and D, we plot the mean of ISD (C) and clustering (D), with error bars
showing the standard deviation. Timepoints all use the same simulations and so are not independent,
I.e. the number of domains in our simulations decreases with time. Dashed lines indicate the values of
ISD and clustering corresponding to the lowest loss rates.
267
268

269
Figure 4. Younger Pfam cohorts have higher variance within their ISD (A) and clustering (B)
distributions, a trend whose timescale is captured in our simulations (C, D). A) and B) Point size is
scaled by n = number of Pfams in that category. Plotted in dark blue is the weighted linear regression
slope (weighted by number of observations per category). Grey lines indicate the 95% confidence
intervals on the regression. In A), for ISD, weighted R^2 = 0.60, p = 2e-10. In B), for clustering,
weighted R^2 = 0.60, p = 2e-10. In C and D) the standard deviation of ISD (C) and clustering (D)
values are plotted for cohorts of simulated Pfams.
277
<u>ISD and clustering also predict mean number of Pfam instances</u>
The mean number of instances of the Pfam per genome is an alternative metric of evolutionary
success. Not only is it likely an indirect proxy of total loss rate (supplementary fig. 1), but it also
shapes proteome composition in its own right, independently of its effect on total loss rate (Figure 1,
middle vs. bottom). Unfortunately, mean number of instances per genome is expected to depend on
genome annotation quality, which is likely to vary across the animal species in our dataset. We have
at least partly addressed this problem by restricting analysis to genomes deemed "Complete". In
agreement with our loss rate results, Pfams with a higher mean number of instances per genome have
lower levels of ISD (Figure 2C, R^2 = 0.0069, p = 7e-11). Pfams with a higher mean number of
instances per genome also have low levels of hydrophobic clustering (adjusted R^2 = 0.0013, p =
0.0017). A non-linear model for the effect of ISD on mean number instances is only marginally
supported over a linear model (p = 0.049 in model comparison with linear relationship, breakpoint =
0.57, 95% confidence intervals: 0.42, -.73), unlike the clear case for a non-linear relationship with loss
rates. However, for hydrophobic clustering, we again find support for a non-linear relationship
(Figure 2D, adjusted R^2 = 0.0099), with a highly supported breakpoint at the same value of clustering
(0.85, 95% confidence intervals: 0.79 - 0.90). A model with a clustering breakpoint fits the data
significantly better than one without (p = 1e-13).
295
296 **Discussion**
297
The persistence of trends in biochemical properties of protein domains (ISD and hydrophobic
clustering) over such strikingly long timespans has been an unresolved puzzle. Our results suggest
that at birth, Pfam domains have a wide range of properties, and are then differentially retained vs.

lost as a function of these properties. Specifically, optimal retention is achieved with somewhat low
ISD values close to 0.18, and with interspersion of hydrophobic amino acids corresponding to a
clustering value of 0.8. The oldest cohort of Pfams has mean ISD and clustering close to these values,
and there is less variance within old Pfam cohorts than within young Pfam cohorts. Simulations show
that the effect sizes of differential retention as a function of ISD and clustering are of the right order
of magnitude (i.e. small enough) to explain how phylostratigraphy trends can persist for such long
evolutionary times without saturating. Conditional on retention, similar values of ISD and clustering
also predict the mean number of domain instances per genome, magnifying the impact of these biases
on proteome-wide composition.
310
This study implements high-throughput phylostratigraphy; while we implemented a number of novel
measures to reduce the frequency of false positive and false negative homologs, our dataset will not
be free of error. We focus on Pfam domains, for which homology detection is performed using the
more sensitive program HMMER rather than BLASTP (El-gebali et al. 2019; Finn, Clements, and
Eddy 2011). Concerns about homology detection bias usually focus on its potential for causing some
rapidly evolving Pfams to be annotated as too young (Moyers and Zhang 2017; Weisman, Murray,
and Eddy 2020). However, Pfam age is not a central focus of our analysis. The potential issue here is
rather that homology detection bias might explain why Pfams with high ISD appear to be lost more
often and have fewer instances per genome (a similar bias for clustering has not been documented).
However, the known difficulty in detecting homology given higher ISD cannot explain our observed
non-monotonic dependence of loss rates and instance number on ISD.
322
To detect as many true homologs as possible, instead of relying on existing genome annotations, we
scan all six reading frames for hits. Each Pfam was as a result included in a mean of 7.35 and a
median of 4 new animal species, out of 343. However, HMMER can produce false positives (Pearson,
Li, and Lopez 2017); while use of curated Pfam seeds might reduce this problem, it will not
completely eliminate it. We minimized impact by co-inferring false positive (or horizontally
transferred) hits, with a higher rate for those found in our 6-frame scan, at the time of our inference of
loss rate. Overall, these data quality measures resulted in a substantial decrease in the mean loss rate
of approximately 14%. Another key quality control was to remove some Pfams altogether as potential
contaminants not from the genomes under study. Our dataset and methodology represent a new
standard for work on protein domain evolution, which, while imperfect, has resulted in a higher
quality dataset for further analysis. Nevertheless, given the residual likely presence of false positive
and false negative Pfam hits, we expect our results on total loss of a Pfam to be more robust to
genome annotation issues than the number of Pfam instances per genome. However, we note that our
instance number and loss rate results qualitatively agree.
337
While protein domains no doubt experience idiosyncratic selection pressures associated with their
specific structures and functions within organisms, our findings suggest a more universal tendency in
which Pfams with ISD and clustering values close to the 'optimum' are differentially retained in the
long term. A candidate for a universal cause is that the same biophysical properties that promote
correct folding also promote toxic aggregation, posing a universal dilemma for proteins (Bucciantini
et al. 2002; Chiti and Dobson 2017). Foy et al. (2019) proposed that young genes tend to address this
dilemma via a more "primitive" strategy, with high ISD representing a sacrifice of folding but safety
from misfolding. The slow evolution of hydrophobic interspersion was proposed to eventually
provide an alternative method for avoiding toxic aggregation that allows for tighter protein folds (Foy
et al. 2019). Our findings suggest that the emergence of a tight protein fold might instead depend on
good luck in the properties of the newborn protein sequence from which it derives. This idea is
supported by lattice protein simulations designed to capture frustration between folding and
aggregation, in which there is an abundance of lower fitness peaks, and where sequences born with
high hydrophobicity (the lattice protein equivalent of low ISD) are more likely to find a high fitness
peak (Bertram and Masel 2020).
353
354 While selection among alternative alleles of the same domain (i.e. descent with modification) is often
355 the default explanation for molecular evolution trends (fig. 1 top), we provide evidence for selection

acting on a higher level, among protein domains, highlighting the importance of differential retention and diversification of domains as an evolutionary force in protein sequence evolution (fig. 1 middle and bottom). Selection acting on multiple different levels to affect a trait does not necessarily result in conflict between the levels (Lewontin 1970). This higher level is analogous to the dependence of speciation and extinction rates on the traits of species (see, for example (Aguilée et al. 2018)). An interesting parallel to our findings is the work of Heim et al. (2015), who found that long-term trends do not seem to be caused by directional selection, but rather by the differential diversification of certain clades. Here we find that the same broad category of explanation applies to long-term trends in protein properties as a function of age since they originated.

The study of long-term directional change in evolution, and the level of selection at which it occurs, has historically focused on species-level traits such as morphology. But there are several advantages of conducting such studies on proteins. Firstly, because Pfam domains are easily identified using HMMER, they are highly tractable for use in the study of higher levels of selection. In particular, it is easier to count how many instances there are of a Pfam in a genome than it is to identify all species in a clade.

Secondly, it is impossible to distinguish between e.g. high extinction rates vs. low speciation rates from observations of extant species distributions alone (Louca and Pennell 2020). An inherent difficulty is that we cannot observe extinct species once they are gone; fossil data is required to provide additional information (Mongiardino Koch, Garwood, and Parry 2021). However, when a domain undergoes complete loss in one lineage, we can directly infer that loss through comparison to other lineages in which it was not lost. Sophisticated methods developed to study speciation and extinction rates (Fitzjohn 2012) could be adapted to take advantage of this during the study of domain duplication and loss rates. Future studies could examine not just how loss rates depend on trait values, but also account for systematic variation in duplication and loss rates among clades, or over time. The study of protein domain diversification and loss is interesting not only to advance our understanding of the evolution of the proteome, but also as a powerful new study system for those interested more broadly in the causes of long-term directional trends in evolution, and in levels of selection.

Methods

Pfam selection

We begin with the Pfam phylostratigraphy dataset of James et al. (2021). Briefly, for a species to be included in this dataset, a "Complete" annotated genome had to be available from NCBI, and the species had to be present in Timetree (Hedges, Dudley, and Kumar 2006). We further restricted this dataset to those Pfams that were present in a minimum of 2 animal species, and for which both ISD and clustering scores could be calculated, resulting in a dataset of 6841 Pfams, and their distributions over a phylogenetic tree of 343 animal species. These Pfams had previously been subject to two quality control filtering steps: a keyword-based screening designed to exclude possible contaminants, and an additional z-score screen to exclude Pfams with species distributions indistinguishable from chance. Genes and Pfams annotated as mitochondrial were also already excluded from our starting database.

Pfam ages were initially taken from James et al. (2021), based on a species tree of 435 eukaryote species, with additional resolution among the two most ancient Pfam cohorts taken from (Weiss et al. 2016). While we calculate loss rates only within the animal species tree, our analysis includes Pfams that originated prior to animals and that are therefore also found in other taxa.

False negatives

To detect potentially missing Pfam instances in unannotated genes, including those missed due to frameshift sequencing errors, we translated all regions not annotated as coding sequences (CDS) in all

six frames. For this purpose, we downloaded annotated genomes of the 343 animal species in our
dataset from NCBI, with access dates by species varying between May and July 2019. We removed
sequences annotated as "CDS" and any sequences of "N" more than 3 nucleotides long. We then
translated the resulting nucleic acid sequences in all six reading frames, not requiring a start codon for
initiation but ending each polypeptide at stop codons, similar to the procedure implemented by
Deutekom et al. (2019), and ran amino acid sequences through InterProScan version 5.33-72.0. We
record all novel Pfam hits for each species. This resulted in each Pfam being included in a mean of
7.35 and a median of 4 new animal species. Although this six-frame search procedure generates false
positives at an elevated rate relative to annotated Pfams, these false positive rates remain modest (see
below). To avoid inflation of mean instances / genome by pseudogenes, we only correct putative false
negatives in cases where the species in question would otherwise have zero instances of the Pfam.
421
422   <u>False positives</u>
423
Excluding contaminant Pfams
As in James et al. (2021), we exclude any Pfam that was both present in fewer than half the species in
the original dataset (i.e. not restricted to animals), and had a species distribution that was
indistinguishable from chance. Such Pfams are likely to be contaminants. Briefly, for each number of
species n, we performed 20,000 simulations in which we placed Pfams in n randomly selected
species. We then inferred the number of total losses by parsimony, and used the mean and variance
for the resulting distribution of simulated numbers of total losses to calculate z scores for each
empirically observed distribution of n Pfam instances. To remove Pfams with species distributions
indistinguishable from chance, James et al. excluded Pfams with z scores < -2. Here, we used an
additional screen where we excluded any Pfams that had z scores < -2 either with or without the
intergenic hits flagged as possible false negatives as described above. This resulted in the further
removal of 141 Pfams, leaving a dataset of 6700 Pfams.
436
Excluding false positive Pfam presence while estimating loss rates by maximum likelihood
We jointly estimate the total loss rate of each Pfam by maximum likelihood, together with the identity
of false positive or horizontally transferred hits/instances. First consider the case of no false positive
or horizontally transferred hits. Using branch lengths obtained from Timetree, we assume that loss
events are independent, making the probability of loss of Pfam i with loss rate λ_i per million years on
branch j of length t_j equal to $1 - \exp\left(-t_i\lambda_j\right)$. The likelihood that Pfam i is completely lost on branches
j and retained in at least one copy on branches k, given total loss rate λ_i, is:

$$L_\lambda(\lambda_i) = \prod_k \exp\left(-t_k\lambda_i\right) \prod_j \left(1 - \exp(-t_j\lambda_i)\right)$$

444
We first provisionally assign losses by assuming a Pfam arose on the branch basal to the
monophyletic clade that includes all species with the Pfam, and is lost on the branches that explain the
current distribution with fewest number of losses, i.e. we use Dollo parsimony (see figure 5). We then
use Newton's method to find the loss rate that maximizes the log likelihood, initialized at the loss rate
calculated as the number of losses/total branch length of the Pfam tree. Because multiple loss events
can occur on the same branch, the estimated ML loss rate estimate is generally very close but not
identical to the initialization value (parsimony mean and median: 0.0058, 0.00088, ML mean and
median: 0.0067, 0.00094). Occasionally, false positives / horizontal transfer events that are not yet
accounted for trigger runaway inference of absurdly high ML loss rates at this stage. When the ML
loss rate > 1 per million years), we therefore reset the loss rate to the initialization value originally
estimated under parsimony, with associated likelihood, prior to proceeding.
456

457
Figure 5) Inferring loss events for a Pfam by Dollo parsimony. Dollo parsimony assumes a Pfam
arose exactly once, on the branch basal to the monophyletic clade that includes all species with the
Pfam, and is lost on the branches that explain the current distribution with the fewest number of
losses. Species with at least one instance of the Pfam are indicated with checks, and those without by
crosses. We initialize the Pfam loss rate as 1/sum of unshaded branch lengths, and then refine this
coarse Dollo-parsimony based estimate through maximum likelihood techniques that allow for false
positive/horizontally transferred instances.
465
466 Next, to remove false positive / horizontally transferred Pfam presence data, we use a recursive
467 empirical Bayesian procedure. We calculated two prior probabilities: f_c, that the annotated presence
468 of a Pfam in a species or clade is a false positive, and f_i, that a Pfam presence discovered only by our
469 six reading frame scan of intergenic sequences is a false positive. These were initialized at 10^{-6} and
470 0.2, respectively.
471
472 If P_C is the set of species annotated as having the Pfam from CDS observations, and P_i is the set of
473 species annotated as having the Pfam only by us from the InterProScan intergenic search, the
474 likelihood of observing the data if the species sets F_C and F_i are false positives is:

$$L = L_\lambda(\lambda_i)f_c^{|F_c|}(1 - f_c)^{|P_c-F_c|}f_i^{|F_i|}(1 - f_i)^{|P_i-F_i|}$$

(1)

475
476 where λ_i is re-estimated by maximum likelihood for each species set F_C and F_i, and absolute
477 magnitude notation indicates the number of species in the set.
478
To find the sets F_C and F_i that maximize likelihood, we first group Pfam presence observations into
the largest possible monophyletic clades where all species putatively possess at least one copy of the
Pfam of interest. We then begin by "flipping" each such monophyletic clade, i.e. designating all
species in a clade as being false positive hits or horizontal transfer events, and determining whether
the likelihood improves. The false positive designation that improves the likelihood the most is kept,
and then the process is repeated. In these subsequent iterations, we also test whether flipping a false
positive back to a true positive improves likelihood. The process ends when no single flip improves
the likelihood. After applying this process to all Pfams, we use the resulting set of false positive
exclusion to estimate the two false positive probabilities. Using this as a revised empirical prior, we
then repeat the procedure until no further change in the false positive rates is observed. Our method
rapidly converges within 4 iterations, to a false positive rate of 0.00021 for Pfams previously
annotated as coding, and 0.031 for Pfams annotated here by us based on presence in putatively
intergenic regions. We calculate the mean number of Pfam instances per animal species within the set
of species annotated as true positives at the end of the procedure described above.
493
494 Our procedures to include presence data when Pfams were previously unannotated, and to exclude
495 presence data as described above, had a noticeable quantitative impact on our data. The combined
496 effect of these quality control measures caused the mean and median loss rate to fall from 0.0075 and

497 0.00091 to 0.0065 and 0.00090. The mean and median number of Pfam instances per species rose
498 from 5.9 and 2.2 to 6.2 and 2.4.
499
500 For some young Pfams (appearing after divergence with plants), our procedures to reduce false
501 negative and false positive errors in presence annotation within the animal species tree lead us to re-
502 estimate Pfam age. All of the Pfam loss rate and age data is available online at: https://github.com/j-e-
503 james/DomainLossRate.
504
ISD
We calculated predicted ISD using IUPred 2 (Dosztányi et al. 2005; Mészáros, Erdös, and Dosztányi
2018), which scores each amino acid between 0 and 1, with a low score indicating low disorder. In
order to obtain a single estimate for each specific Pfam instance, we averaged over all amino acids.
Per Pfam scores were then calculating by finding the mean over all copies of the Pfam in our dataset
(i.e. animal species with more Pfam instances had higher weight).
511
Clustering
Briefly, we calculated clustering as a kind of normalized index of dispersion: the variance in the
number of the most hydrophobic amino acids (Leu, Ile, Val, Phe, Met and Trp) among blocks of 6
consecutive amino acids compared to the mean number, normalized so as to be comparable across
Pfams of different lengths and hydrophobicities. If the length of a Pfam was not a multiple of 6, we
took the average of all possible amino acid frames of 6, truncating the ends appropriately. For more
detail, see Foy et al (2019) and James et al (2021). A clustering value of 1 indicates random
distribution of hydrophobic residues, with higher values indicating that hydrophobic residues tend to
be arranged in clusters along the sequence, and lower values indicating that hydrophobic residues are
more evenly distributed along the sequence than would be expected by chance. As for ISD, per Pfam
scores were then calculated as the mean over all instances of each Pfam in our animal dataset.
523
Statistics
All plotting and statistical analysis was performed in R. Linear regression models that incorporate
breakpoints were implemented using the segmented package. We used the packages lme4, MuMIn,
and segmented to conduct models, and ggplot2 to create plots. Prior to linear regression, we
transformed loss rate using an optimized two parameter Box-Cox transform ($\lambda = 0.0054$, $\lambda_2 = 6.1\text{e-}06$)
calculated using the R package geoR. We transformed mean instance number using a Box-Cox
transform ($\lambda = -0.61$). We transform after rather than before taking the mean because the natural units
for scoring impact on the proteome (Fig. 1 middle) are linear. Histograms comparing the distribution
of untransformed versus transformed loss rates, in addition to Q-Q plots illustrating the requirement
for transformation in our linear regression models with ISD, are shown in supplementary figure 3.
534
Throughout this work, we report the adjusted R^2 values for our linear regression model results, as
implemented in the programming language R, which uses the Wherry formula (Wherry 1931). This R^2
accounts for the number of explanatory variables included in the model. Nested models were
compared using the anova function in base R. We conducted mixed effects models in R using the lmer
package, and calculated corresponding pseudo R^2 values using the R package MumIn. This package
returns marginal R^2 values, which can be interpreted as the variance explained by fixed effects in the
model, and conditional R^2 values, which can be interpreted as the variance explained by both the fixed
and random effects in the model (Nakagawa and Schielzeth 2013). R scripts and data required to
replicate our analysis are available at: https://github.com/j-e-james/DomainLossRate.
544
Simulations
We performed simulations to assess the biological relevance of our inferred loss rates, using custom
python code available at https://github.com/j-e-james/DomainLossRate. We initialize these
simulations by sampling, with replacement, the Pfams that we estimate to be younger than 100 MY
old, to produce an initial set of 5000 Pfams, each represented by its ISD and clustering values. We
then assign each Pfam the loss rate predicted from either its ISD value, or its clustering value, on the
basis of our corresponding regression model. For each Pfam, we then sample the time to a single loss

from an exponential distribution, with scale parameter = 1/loss rate. We then track the mean and standard deviation of the population of survivors over time.

Acknowledgements

We thank Cecile Ané for helpful discussions that prompted a more complex project that we abandoned in favor of this simpler approach. We thank Nathan Aviles for helpful discussions about fancier statistical approaches that we didn't use. We thank all of the participants at the 2019 "Workshop on Long-Term Trends in Evolution" for stimulating discussions. Both this work and the workshop were primarily supported by the John Templeton Foundation (60814). This work was also supported by the National Institutes of Health (GM-104040), and in its final stages, by the John Templeton Foundation (62220) and a Wenner-Gren postdoctoral stipendium.

References

Aguilée, Robin, Fanny Gascuel, Amaury Lambert, and Regis Ferriere. 2018. "Clade Diversification Dynamics and the Biotic and Abiotic Controls of Speciation and Extinction Rates." *Nature Communications* 9 (3013). https://doi.org/10.1038/s41467-018-05419-7.

Bagowski, Christoph P, Wouter Bruins, and Aartjan J W Velthuis. 2010. "The Nature of Protein Domain Evolution : Shaping the Interaction Network." *Current Genomics* 11: 368–76.

Basu, Malay Kumar, Liran Carmel, Igor B. Rogozin, and Eugene V. Koonin. 2008. "Evolution of Protein Domain Promiscuity in Eukaryotes." *Genome Research* 18 (3): 449–61. https://doi.org/10.1101/gr.6943508.

Bertram, Jason, and Joanna Masel. 2020. "Evolution Rapidly Optimizes Stability and Aggregation in Lattice Proteins despite Pervasive Landscape Valleys and Mazes." *Genetics* 214 (4): 1047-1057. https://doi.org/10.1101/776450.

Bornberg-Bauer, Erich, and M. Mar Albà. 2013. "Dynamics and Adaptive Benefits of Modular Protein Evolution." *Current Opinion in Structural Biology* 23 (3): 459–66. https://doi.org/10.1016/j.sbi.2013.02.012.

Breitwieser, Florian P., Mihaela Pertea, Aleksey V. Zimin, and Steven L. Salzberg. 2019. "Human Contamination in Bacterial Genomes Has Created Thousands of Spurious Proteins." *Genome Research* 29 (6): 954–60. https://doi.org/10.1101/gr.245373.118.

Bucciantini, Monica, Elisa Giannoni, Fabrizio Chiti, Fabiana Baroni, Niccolò Taddei, Giampietro Ramponi, Christopher M. Dobson, and Massimo Stefani. 2002. "Inherent Toxicity of Aggregates Implies a Common Mechanism for Protein Misfolding Diseases." *Nature* 416 (6880): 507–11. https://doi.org/10.1038/416507a.

Chiti, Fabrizio, and Christopher M. Dobson. 2017. "Protein Misfolding, Amyloid Formation, and Human Disease: A Summary of Progress Over the Last Decade." *Annual Review of Biochemistry* 86 (1): 27–68. https://doi.org/10.1146/annurev-biochem-061516-045115.

Cope, E. D. 1885. "On the Evolution of the Vertebrata, Progressive and Retrogressive." *The American Naturalist* 19 (2): 140–48.

Deutekom, Eva S., Julian Vosseberg, Teunis J.P. Van Dam, and Berend Snel. 2019. "Measuring the Impact of Gene Prediction on Gene Loss Estimates in Eukaryotes by Quantifying Falsely Inferred Absences." *PLoS Computational Biology* 15 (8): e1007301. https://doi.org/10.1371/journal.pcbi.1007301.

Dosztányi, Zsuzsanna, Veronika Csizmók, Péter Tompa, and István Simon. 2005. "The Pairwise Energy Content Estimated from Amino Acid Composition Discriminates between Folded and Intrinsically Unstructured Proteins." *Journal of Molecular Biology* 347 (4): 827–39. https://doi.org/10.1016/j.jmb.2005.01.071.

El-gebali, Sara, Jaina Mistry, Alex Bateman, Sean R Eddy, Simon C Potter, Matloob Qureshi, Lorna J Richardson, et al. 2019. "The Pfam Protein Families Database in 2019" *Nucleic Acids Research* 47 (D1): D427–D432. https://doi.org/10.1093/nar/gky995.

Finn, Robert D., Jody Clements, and Sean R. Eddy. 2011. "HMMER Web Server: Interactive Sequence Similarity Searching." *Nucleic Acids Research* 39 (Web Server issue): W29–W37. https://doi.org/10.1093/nar/gkr367.

Fitzjohn, Richard G. 2012. "Diversitree: Comparative Phylogenetic Analyses of Diversification in R."
Methods in Ecology and Evolution 3 (6): 1084–92. https://doi.org/10.1111/j.2041-
210X.2012.00234.x.
Foy, Scott G, Benjamin A Wilson, Jason Bertram, Matthew H J Cordes, and Joanna Masel. 2019. "A
Shift in Aggregation Avoidance Strategy Marks a Long-Term Direction to Protein Evolution."
Genetics 211 (4): 1345–55. https://doi.org/10.1534/genetics.118.301719.
Gregory, T. Ryan. 2008. "Evolutionary Trends." *Evolution: Education and Outreach* 1 (3): 259–73.
https://doi.org/10.1007/s12052-008-0055-6.
Hahn, Matthew W., Mira V. Han, and Sang Gook Han. 2007. "Gene Family Evolution across 12
Drosophila Genomes." *PLoS Genetics* 3 (11): 2135–46.
https://doi.org/10.1371/journal.pgen.0030197.
Hedges, S. Blair, Joel Dudley, and Sudhir Kumar. 2006. "TimeTree: A Public Knowledge-Base of
Divergence Times among Organisms." *Bioinformatics* 22 (23): 2971–72.
https://doi.org/10.1093/bioinformatics/btl505.
Heim, Noel A., Matthew L. Knope, Ellen K. Schaal, Steve C. Wang, and Jonathan L. Payne. 2015.
"Cope's Rule in the Evolution of Marine Animals." *Science* 347 (6224): 867–70.
https://doi.org/10.1126/science.1260065.
James, Jennifer E., Sara M. Willis, Paul G. Nelson, Catherine Weibel, Luke J. Kosinski, and Joanna
Masel. 2021. "Universal and Taxon-Specific Trends in Protein Sequences as a Function of Age."
ELife 10: e57347. https://doi.org/10.7554/eLife.57347.
Kosinski, Luke J, Nathan R Aviles, Kevin Gomez, and Joanna Masel. 2022. "Random Peptides Rich
in Small and Disorder-Promoting Amino Acids Are Less Likely to Be Harmful." *Genome
Biology and Evolution* 14 (6): 1–14. https://doi.org/10.1093/gbe/evac085.
Lewontin, R. C. 1970. "Units of Selection." *Annual Review of Ecology and Systematics* 1: 1–18.
https://doi.org/10.1016/B978-008045405-4.00810-7.
Louca, Stilianos, and Matthew W. Pennell. 2020. "Extant Timetrees Are Consistent with a Myriad of
Diversification Histories." *Nature* 580 (7804): 502–5. https://doi.org/10.1038/s41586-020-2176-
1.
Lu, Jennifer, and Steven L. Salzberg. 2018. "Removing Contaminants from Databases of Draft
Genomes." *PLoS Computational Biology* 14 (6): e1006277.
https://doi.org/10.1371/journal.pcbi.1006277.
McShea, Daniel W. 1991. "Complexity and Evolution: What Everybody Knows." *Biology and
Philosophy* 6 (3): 303–24. https://doi.org/10.1007/BF00132234.
Mészáros, Bálint, Gábor Erdös, and Zsuzsanna Dosztányi. 2018. "IUPred2A: Context-Dependent
Prediction of Protein Disorder as a Function of Redox State and Protein Binding." *Nucleic Acids
Research* 46 (Web Server issue): W329–37. https://doi.org/10.1093/nar/gky384.
Mongiardino Koch, Nicolás, Russell J. Garwood, and Luke A. Parry. 2021. "Fossils Improve
Phylogenetic Analyses of Morphological Characters." *Proceedings of the Royal Society B* 288:
20210044. https://doi.org/10.1098/rspb.2021.0044.
Moore, Andrew D., and Erich Bornberg-Bauer. 2012. "The Dynamics and Evolutionary Potential of
Domain Loss and Emergence." *Molecular Biology and Evolution* 29 (2): 787–96.
https://doi.org/10.1093/molbev/msr250.
Moyers, Bryan A., and Jianzhi Zhang. 2017. "Further Simulations and Analyses Demonstrate Open
Problems of Phylostratigraphy." *Genome Biology and Evolution* 9 (6): 1519–27.
https://doi.org/10.1093/gbe/evx109.
Nakagawa, Shinichi, and Holger Schielzeth. 2013. "A General and Simple Method for Obtaining R2
from Generalized Linear Mixed-Effects Models." *Methods in Ecology and Evolution* 4 (2): 133–
42. https://doi.org/10.1111/j.2041-210x.2012.00261.x.
Nasir, Arshan, Kyung Mo Kim, and Gustavo Caetano-Anollés. 2014. "Global Patterns of Protein
Domain Gain and Loss in Superkingdoms." *PLoS Computational Biology* 10 (1): e1003452.
https://doi.org/10.1371/journal.pcbi.1003452.
Palmieri, Nicola, Carolin Kosiol, and Christian Schlötterer. 2014. "The Life Cycle of Drosophila
Orphan Genes." *ELife* 3: e01311. https://doi.org/10.7554/elife.01311.
Payne, Jonathan L., Alison G. Boyer, James H. Brown, Seth Finnegan, Micha Kowalewski, Richard
A. Krause, S. Kathleen Lyons, et al. 2009. "Two-Phase Increase in the Maximum Size of Life

over 3.5 Billion Years Reflects Biological Innovation and Environmental Opportunity." *Proceedings of the National Academy of Sciences of the United States of America* 106 (1): 24–27. https://doi.org/10.1073/pnas.0806314106.

Pearson, William R., Weizhong Li, and Rodrigo Lopez. 2017. "Query-Seeded Iterative Sequence Similarity Searching Improves Selectivity 5 – 20-Fold." *Nucleic Acids Research* 45 (7): e46. https://doi.org/10.1093/nar/gkw1207.

Ponting, Chris P., and Robert R. Russell. 2002. "The Natural History of Protein Domains." *Annual Review of Biophysics and Biomolecular Structure* 31 (1): 45–71. https://doi.org/10.1146/annurev.biophys.31.082901.134314.

Salzberg, Steven L. 2017. "Horizontal Gene Transfer Is Not a Hallmark of the Human Genome." *Genome Biology* 18 (85). https://doi.org/10.1186/s13059-017-1214-2.

Schlötterer, Christian. 2015. "Genes from Scratch--the Evolutionary Fate of de Novo Genes." *Trends in Genetics* 31 (4): 215–19. https://doi.org/10.1016/j.tig.2015.02.007.

Tautz, Diethard, and Tomislav Domazet-Lošo. 2011. "The Evolutionary Origin of Orphan Genes." *Nature Reviews Genetics* 12 (10): 692–702. https://doi.org/10.1038/nrg3053.

Vieira-Silva, Sara, and Eduardo P.C. Rocha. 2008. "An Assessment of the Impacts of Molecular Oxygen on the Evolution of Proteomes." *Molecular Biology and Evolution* 25 (9): 1931–42. https://doi.org/10.1093/molbev/msn142.

Weisman, Caroline M., Andrew W. Murray, and Sean R. Eddy. 2020. "Many, but Not All, Lineage-Specific Genes Can Be Explained by Homology Detection Failure." *PLoS Biology* 18 (11): e3000862. https://doi.org/10.1371/journal.pbio.3000862.

Weiss, Madeline C., Filipa L. Sousa, Natalia Mrnjavac, Sinje Neukirchen, Mayo Roettger, Shijulal Nelson-Sathi, and William F. Martin. 2016. "The Physiology and Habitat of the Last Universal Common Ancestor." *Nature Microbiology* 1: 16116. https://doi.org/10.1038/nmicrobiol.2016.116.

Wherry, R. J. 1931. "A New Formula for Predicting the Shrinkage of the Coefficient of Multiple Correlation." *The Annals of Mathematical Statistics* 2 (4): 440–57.

Wilson, Benjamin A., Scott G. Foy, Rafik Neme, and Joanna Masel. 2017. "Young Genes Are Highly Disordered as Predicted by the Preadaptation Hypothesis of de Novo Gene Birth." *Nature Ecology and Evolution* 1 (6): 0146. https://doi.org/10.1038/s41559-017-0146.