Extensive Dysregulation of SLK Splicing in Cancers Impacts

Metastasis

Ying-Qun Yang [1,2, #], Yue Hu [1,2, #], Si-Rui Zhang [1,2, #], Jie-Fu Li [1], Jia-Wen

Guan[1], Wen-Jing Zhang [3], Yu Sun [3], Xiao-Yan Feng [1], Jing Sun[1], Yun Yang [1],

Zefeng Wang [1, *], Huan-Huan Wei [1, *]

[1] CAS Key Laboratory of Computational Biology, Shanghai Institute of Nutrition

and Health, Chinese Academy of Sciences, Shanghai 200031, China

[2] University of Chinese Academy of Sciences, Beijing 100049, China

[3] Institute of Cancer Stem Cell, Dalian Medical University, Dalian 116044,

China

[#] These authors contributed equally to this work

*Corresponding to: weihuanhuan@picb.ac.cn (H-H. Wei),

wangzefeng@picb.ac.cn (Z. Wang).

Abstract

RNA splicing control is a pivotal aspect of gene regulation and is closely associated with cancer development. From a pan-cancer transcriptome investigation in the splicing layer, we discovered a critical cancer-associated alternative splicing (AS) event at exon 13 of SLK which produces two isoforms, SLK-L and SLK-S. The splicing is dramatically shifted towards SLK-L across multiple prevalent cancer types. We demonstrated that SLK-L plays an essential role in cancer development, especially in metastasis both in cells and in mice, whereas splicing toward SLK-S inhibits cancer development. RNA-seq revealed the two SLK isoforms play different roles in pathways related to cell migration. Furthermore, different SLK isoforms demonstrate varying binding affinities to certain cell junction markers, in part indicating the AS of SLK contributes to cancer cell migration. In addition, the splicing factor Rbfox2 was identified to specifically inhibit the inclusion of exon 13 by binding intron 12 of SLK. Collectively, our study innovatively uncovers the biological consequences and underlying mechanisms for one of the most mis-spliced genes in cancer, highlighting its potential significance in cancer diagnosis and treatment.

Keywords: SLK, Alternative splicing, Cancer, Metastasis, Cell junction, Migration.

Introduction

RNA splicing control is a critical layer of gene regulation in addition to transcriptional and epigenetic regulation. Extensive alteration of splicing is considered an established hallmark of cancers [1-8]. Studying splicing dysregulation in cancers is a challenge due to the large noise, as well as the diverse and complicated mechanisms. We previously established a transcriptome-wide identification of cancer-specific splicing events across several tumors through analysis of transcriptome sequencing data from The Cancer Genome Atlas (TCGA) [9]. This method was utilized to analyze multiple prevalent forms of cancer that have an adequate number of paired normal samples in this study. As a result, we identified a critical cancer-associated splicing event at the exon 13 of STE20-like protein kinase (SLK), which produces two isoforms, SLK-L and SLK-S. Impressively, the splicing is dramatically shifted towards SLK-L among thousands of cancer patients across multiple cancer types.

SLK is a serine/threonine kinase and belongs to the five germinal center kinase families. It contains several conserved domains, including a kinase domain, a predicted central unstructured region, and a C-terminal coiled-coil domain. Previous studies discovered that SLK was involved in several cellular processes such as cell cycle progression by phosphorylating PLK1 and promoting G2/M phase transformation [10, 11], or activating the phosphorylation of p38 mitogen-activated protein kinase (MAPK) through ASK1 [12, 13]. Over the twenty years, SLK has emerged as a regulator of the cytoskeleton [14]. It has been shown to activate the Rho family of GTPases, which in turn regulate

cytoskeletal dynamics and cell migration [15]. It impacts the migration of mouse

fibroblasts by directly interacting with LIM domain-binding protein 1 (Ldb1) [16],

and is involved in the epithelial to mesenchymal transition (EMT) process [17-19]

and the regulation of focal adhesion turnover [20] through its kinase activity [15].

Interestingly, several recent studies revealed that the ability of SLK for

regulating cellular functions appears to be kinase activity independent,

suggesting that it may be an important scaffold for signal transduction

pathways[14]. For instance, in fibroblasts, either the expression of an SLK

variant lacking kinase activity or depletion of SLK can result in growth arrest,

indicating that both kinase-dependent and -independent functions of SLK are

necessary [17]. Functions of Slik, the homolog of SLK in *Drosophila*, are

dependent on its proper subcellular localization, which is mediated through its

C-terminal ATH domain independent of its activation status [21]. These results

suggested that the alternative 13[th] exon in the ATH domain of SLK may have

important functions. However, the underlying mechanisms remain unknown.

Accumulating evidence has shown that *SLK* plays important role in

cellular functions associated with cancer development [14, 22, 23]. However, few

studies report and investigate the splicing switch of *SLK* in cancers. The

prevalent AS alteration of *SLK* indicates that the two isoforms may play

different roles in cancers. Furthermore, the extensively dysregulated splicing

events across cancer types suggest the possibility of universal functional and

mechanistic implications. Therefore, uncovering the functions and

mechanisms of abnormal splicing of *SLK* in cancers will provide insights into

the cancer-associated key AS events with therapeutic potential and clinical

values.

Results

88

89

90 ***SLK* splicing is significantly and extensively altered across multiple**
91 common types of cancer

92   Using the previously established method [9], we performed a pan-cancer

93 investigation of global AS dysregulation by comparing the Percent Spliced In

94 (PSI) value between tumor samples and their adjacent normal controls (Fig.

95 1a). For a reliable comparison, we selected the most prevalent forms of cancer

96 that have a sufficient number of paired normal samples. As a result, we

97 identified a substantial number of splicing events that were differentially

98 expressed in cancer. Our findings indicate that approximately 70% of splicing

99 events are altered across different cancer types (Fig. S1a). This observation

100 further suggests that the shared spliced genes might potentially serve as an

101 underlying biological mechanism that promotes cancer development and

102 progression. Additionally, we observed that such shared genes altering in

103 more than 10 cancer types are enriched in cancer-related migration processes

104 such as actin filament organization, cell junction assembly, and cell-substrate

105 adhesion (Fig. 1b). Of particular interest, we have discovered notable and

106 comprehensive changes in the splicing of SLK gene in 12 different types of

107 cancer for the first time (Fig. 1b and S1a). Besides, in nine cancer types, SLK

108 showed a significant shift toward the long isoform with PSI significantly

109 increased compared to that in normal samples, which indicates AS of SLK play

110 key roles during cancer development (Fig. 1c).

111   The human *SLK* gene is located on chromosome 10 and consists of 19

112 exons. The SLK protein contains the N-terminal kinase catalytic domain

113 (encoded by exons 1 to 8), an interdomain linker, and the C-terminal coiled-coil

domain responsible for protein-protein interaction (encoded by exons 11 to 19).

The alternative inclusion of exon 13 in *SLK* pre-mRNA produces two isoforms,

the long or short isoform (SLK-L and SLK-S), which differs at a fragment of 31

amino acids in the C-terminal coiled-coil domain (Fig. 1d). The comparison of

the protein structures predicted by Alphafold (https://alphafold.ebi.ac.uk/, [24])

demonstrates that the exon 13 encoded sequence locates in the terminus of

the long α-helix inside the coiled-coil domain (Fig. 1e), suggesting a

kinase-independent activity between SLK-L and SLK-S. The structure of SLK

also indicates its potential role as scaffolding protein.

SLK AS event is conserved in mammals (Fig. S1b), indicating functional

importance. We further investigated the expression of SLK isoforms in various

tissues using Genotype-Tissue Expression (GTEx) data [25]. The exon count

analysis showed a preference for SLK-S expression in tissues with slow

self-renewal (e.g., brain), whereas the SLK-L is preferably expressed in

tissues with rapid proliferation and self-renewal (e.g., skin). In particular, the

non-dividing muscular tissue of the esophagus mainly contains the SLK-S

isoform, whereas the fast-renewal mucosa membrane of the esophagus

contains the predominant SLK-L isoform (Fig. S1c). Such pattern is largely

consistent with the *SLK* splicing changes in tumors that contain many actively

dividing cells (Fig. 1c), implying that SLK-L may play a role in promoting tumor

development. Moreover, the mechanism of this dysregulation may play

general roles across cancer types.

138
139
140
Figure 1 Pan-cancer transcriptome analysis in the splicing layer revealed SLK
AS alters dramatically in multiple types of human cancer. **a,** Analysis pipeline
to identify AS events differentially spliced in cancers versus normal tissues. **b,**
GO enriched network of overlapped AS genes in more than 10 cancer types. **c,**
The SLK splicing ratios across various cancer types in the TCGA database.
PSI: percent spliced in. **d,** Schematics of human SLK pre-mRNA and protein
isoforms produced from SLK AS. the 13[th] exon is included in SLK-L while
skipped in SLK-S. **e,** The comparison of the predicted structure of SLK-L
(yellow) and SLK-S (blue) isoform by Alphafold. The alternative spliced exon
13[th] is highlighted in red.
151

Alternative Splicing of SLK affects cancer cell proliferation and migration

To further study the function of AS in SLK, we used siRNAs that were designed to specifically target SLK-L (binding to the 13th exon of SLK) or SLK-S (binding to the junction of the 12th and 14th exons) in colon cancer HCT116 cells (Fig. S2a-S2b). The highly invasive HCT116 cells have a normal diploid karyotype with the PSI of the SLK between 0.4-0.7, which is ideal to examine the change of AS ratio in both directions. We found that the cells with SLK-S knockdown showed similar cellular phenotypes compared to the control cells, whereas cells with SLK-L knockdown showed a significant reduction in cell proliferation, anchorage-dependent colony formation, migration, and invasion (Fig. S2c-S2f), suggesting that the two isoforms have distinct cellular functions.

Since siRNA-mediated gene knockdown is transient and often leads to off-target effects, resulting in residual expression of the targeted gene (Fig. s2b). we used CRISPR technology to establish HCT116 stable cell lines that specifically express SLK-L or SLK-S to ensure the reliability and stability of the functional experiments (Fig. 2a-2b). Consistent with the siRNA knockdown results, HCT116 cells with only SLK-S showed a significant reduction in cell proliferation (Fig. 2c), colony formation (Fig. 2d), cell migration (Fig. 2e), and cell invasion (Fig. 2f), again suggesting that SLK-L deficiency may suppress cancer progression. As a comparison, the cells with SLK-S depletion exhibited normal or slightly increased cell growth and cell migration (Fig. 2e-2f, Fig. S2e-S2f), suggesting that SLK-L isoform is required for the robust proliferation and migration of cancer cells.

For the purpose of determine whether the splicing switch of SLK were

authentically responsible for the observed cellular phenotypes, we

re-expressed the isoforms separately or in combination in the SLK knockout

(SLK-KO) HCT116 cells generated with CRISPR-Cas technology (Fig. 3a-3b).

Our results showed that cells with re-expressed SLK-L isoform exhibited

restored cell proliferation and migration/invasion, while cells with re-expressed

SLK-S isoform were similar to SLK-KO cells (see Fig. 3c-3f). These findings

provide evidence that the SLK-L isoform promotes the proliferation and

migration of cancer cells like an oncogene.

Figure 2 The cancer cells stably express SLK isoforms were generated by
CRISPR-cas and exhibited different proliferation and migration abilities. **a,**

HCT116 cells expressing different SLK isoforms were constructed by CRISPR. **b,** RNA and protein levels of SLK under CRISPR genome editing. **c-f,** Cancer cell progression was measured from triplicate experiments, with p values calculated by one-way ANOVA and Tukey's test. **c,** various SLK cells were grown for 6 days, with cell numbers counted every day (n=3) to measure cell growth **d,** anchorage-dependent colony formation of different SLK cell lines (n=3; scale bar:1 cm). The planting efficiency was quantified with ImageJ. **e,** *In vitro* migration of different SLK cell lines was analyzed by wound healing assay (n=3). **f,** cell invasion was measured by transwell assay (n=3; scale bar: 200 µm). Cell number was quantified with ImageJ. *p<0.05, **p<0.01, ***p<0.001, ****p<0.0001, ns, not significant. The chart represents mean±SD.

Figure 3 Cancer progression was restored by recovering the expression of SLK isoforms. **a,** Construction of SLK knockout cell line by CRISPR. **b,** SLK

isoforms with various tags were re-expressed in SLK knockout cells, with the
RNA and protein levels of SLK detected accordingly. **c-g,** Cancer cell
development was measured as described in Fig. 2.

SLK splicing switch affects cancer growth and metastasis in mice

Based on the cell function experiment, we investigated the effect of dysregulated SLK AS on cancer progress *in vivo*. We generated xenograft mice using HCT116 cells expressing SLK-WT, SLK-L, SLK-S, or SLK-KO. Nude mice were inoculated with a low (1×10^6 cells, Fig. 4a) or high (5×10^6 cells, Fig. S3A) number of cells in two separate experiments. As expected, the size and weight of the subcutaneous tumors increased rapidly in the SLK-WT and SLK-L groups while they were in smaller size and weight in the SLK-S group (Fig. 4a-4b and Fig. S3a-3c). Moreover, in the high-dosage experiment, the mice inoculated with SLK-WT and SLK-L exhibited lower survival in comparison to the SLK-S group (Fig. S3a). These results indicated that SLK-L contributes to cancer development, and suggested the splicing switch to SLK-S had the potential for cancer suppression. Surprisingly, the SLK knockout cells grew faster than SLK-S cells (Fig. S3a), indicating an undiscovered complementary mechanism upon SLK knockout.

Interestingly, in situ tumorigenicity assay exhibited that SLK-L cells demonstrated a similar phenotype compared with SLK-WT cells, which is consistent with *in vitro* experiments. However, we noticed that in the *in vitro* experiments, SLK-L cells demonstrated higher metastasis capabilities than SLK-WT cells (Fig. 2 and 3). Considering metastasis is the main cause of cancer-related deaths [26], and the emerged functions of SLK as a central regulator of cytoskeletal dynamics [14, 27] and cell migration [28, 29], we hypothesized that SLK splicing switch has critical effects on cancer metastasis. To testify our hypothesis, we developed a tail vein injection mouse model to examine metastatic colonization and subsequent growth of different SLK cells. Different groups of SLK cells were stably transduced with viral vectors

encoding firefly luciferase and injected into mice via the lateral tail vein. The

bioluminescence of the tumor cells was monitored and measured to quantify

changes in metastatic burden over time. For reliability, the experiment was

performed in two immune deficiency mouse strains, the BALB/c nude mice

(Fig. S3d) and C-NKG mice (Fig. 4d and S3e). In BALB/c nude mice (Fig. S3d),

four in five mice injected with the SLK-L cells demonstrated lung and other

organ metastasis before the endpoint (7 weeks post-injection) while only one

and zero mice exhibited metastasis in SLK-WT and SLK-S group, respectively

(Fig. S3d). In C-NKG mice, lung and other organ metastasis was observed in

all SLK groups, with SLK-L cells demonstrating significant higher metastatic

burden and fatality than SLK-WT and SLK-S cells (Fig. 4d and Fig. S3e). The

faster metastasis formation and growth in C-NKG mice than in BALB/c mice

mainly contributed to the higher level of immune deficiency of C-NKG mice,

considering that depletion of NK cells in mice increases metastasis [30].

Figure 4 Different groups of SLK cells exhibited distinct impacts on cancer
development. **a-c,** Xenograft tumors were generated in BALB/c nude mice by
subcutaneous injection with SLK-WT, SLK-L, or SLK-S in **b** and **c**, respectively,
with *p* values calculated by Dunnett's test. **d-e,** Metastasis of SLK cell groups
was investigated by the tail vein metastasis model. Different SLK cell groups
stably expressing firefly luciferase were injected into the lateral tail vein of
immune deficiency mice and the metastasis was monitored by
bioluminescence imaging every week post injection **(see also Fig. S3)**. **d,** The
bioluminescence images were demonstrated in the two replicates of C-NKG
mice on day 14. **e,** Box plot of the mean $\log_{10}$ transformed luciferase signal
measured for each mouse in d. Statistical significance was calculated using a
Student's *t* test; *p<0.05, **p<0.01, ***p<0.001, ****p<0.0001, ns, not
significant.

RNA-seq revealed SLK splicing switch influence key components in cell junction and cell migration

Cell and mouse experiments indicated that the different isoforms of SLK had significant impact on cell migration and metastasis. We next sought to determine how splicing switch of SLK isoforms decide cancer progression. Though accumulating studies unveiled that SLK plays functions in EMT and cytoskeleton thereby influences cell migration, there is still a lack of systematic investigation on the functions of SLK and its isoforms. To understand the underlying mechanism of SLK splicing switch in a systematic manner, we further examine the biological functions of SLK isoforms by analyzing RNA-seq data of different SLK cell lines (Fig. 6a). We discovered many differentially expressed genes are enriched in cancer-related biological process, such as the Wnt signaling pathway and tissue migration (Fig. 6b). There are significant overlaps between these differentially expressed genes across different SLK cell types (Fig. 6c), indicating SLK isoforms are involved in similar network yet play distinct roles. Among them, L vs S and KO vs WT shared higher overlap than other comparisons (Fig. 6c). However, the genes differently expressed in L vs S is significantly enriched in pathways related to cancer development while the genes in KO vs WT may involved in other cellular functions (Fig. 6b). These findings further indicate the cancer-associate function of SLK attributes to the regulation in the splicing level other than the expression level. In addition, by comparing the SLK-L and SLK-S isoforms, we were able to account for the changes primarily caused by the splicing switch. Within these differential genes from SLK-L and SLK-S cell lines, we identified two EMT-related gene sets using GSEA. The GSEA also reveals slight enrichment of EMT gene sets in all comparisons (Fig. S5). Interestingly, SLK-S and SLK-KO exhibited a

different trend of enrichment compared to the SLK-L. This observation

suggests the involvement of different SLK isoforms in the process of cell

migration.

Meanwhile, the Wnt signaling pathway can induce the expression of the

key transcription factor Twist, which can repress the expression of epithelial

markers such as E-cadherin, occludins, and claudins. On the contrary, it will

induce expression of mesenchymal markers such as N-cadherin, fibronectin,

and vimentin [31]. Consistently, the expression of Twist cannot be detected in

SLK-S cells but is moderately expressed in SLK-WT and SLK-L cells (Fig 6f),

again demonstrated that the two isoforms play distinct roles in EMT and

metastasis. To further identify potential gene targets of SLK in cancer

metastasis, we retrieved all differentially expressed genes involved in cell

junction assembly, Wnt signaling pathway, and regulation of epithelial cell

proliferation, and found that many cell junction markers have been influenced.

Collectively, RNA-seq of different SLK isoforms revealed the isoforms execute

distinct functions in cellular pathways related to cancer in a systematic

manner.

Figure 6 Switch of SLK isoforms change EMT-related genes **a,** RNA-seq of indicated SLK cell lines. **b,** Gene ontology enrichment for differentially expressed genes from SLK-KO vs SLK-WT, SLK-L vs SLK-S, SLK-L vs SLK-WT and SLK-S vs SLK-WT. **c**, Overlap of differential genes from different comparisons. **d-e,** GSEA was performed to rank differentially expressed genes for SLK-L and SLK-S. **f,** Heatmap of overlapped differential genes in GO:0016055 "Wnt signaling pathway", GO:0034329 "cell junction assembly", GO:0002064 "epithelial cell development", and GO:0060485 "mesenchyme development".

SLK splicing switch influences cell migration in part through their different binding abilities to EMT and cell junction markers

Based on the experimental results and RNA-seq data, we consider that AS of SLK is closely intertwined with EMT and cell junction regulation, thus influencing cell migration. Considering that SLK-L has extra 31 amino acids which form a longer coiled-coil domain by the structural prediction and comparison by Alphafold (Fig. 1e), it indicates that the functional difference between the two isoforms attributes to the binding activity rather than the kinase activity. Therefore, there are two plausible explanations of how SLK isoforms can affect functions. First, the distinct isoforms may exhibit divergent cellular localization, enabling discrepant biological functions. Alternatively, the two SLK isoforms may have similar cellular localization, but act as antagonists and have differential interactions with corresponding targets, ultimately influencing distinct cascades. To investigate the mechanism, we first sought to determine whether the extra amino acids can affect the subcellular localization of SLK isoforms. By delivering different tagged SLK isoforms into the cells and detecting the immunofluorescence localization, we found both SLK isoforms locate near the cell membrane (Fig. 7a), suggesting their function difference is irrelevant to cellular localization.

As SLK isoform switch is closely linked to cell migration, we next tested several biomarkers of EMT such as E-cadherin, occludin, N-cadherin, and vimentin that were differentially expressed based on RNA-seq data to assess their expression in different SLK cell lines (Fig. 7b). As a result, occludin and vimentin were detected to be differentially expressed in SLK-L versus SLK-S stable cell lines. SLK-S cells have a higher expression level of occludin than

SLK-L cells (Fig. 7c) whereas vimentin has a lower expression level in SLK-S

cells than in SLK-L cells.

Occludin is considered a staple of tight junctions and performs vital

functions in EMT, which contributes to cancer metastasis. To address whether

SLK isoforms interact differentially with occludin, we test their interaction both

through immunoprecipitation and localization. Immunoprecipitation showed

that SLK isoforms had different binding capabilities with occludin. The SLK-S

had a stronger binding ability to occludin, while the SLK-L was barely bound to

occludin (Fig. 7c). Furthermore, the immune colocalization demonstrated that

both SLK isoforms and occludin localized around the cell membrane (Fig. 7d),

indicating the possible interaction between SLK and occludin.

Given cancer cells with low occludin expression are more likely to migrate

and invade [32], we assess the relationship between occludin expression level

and cell development in this study by overexpressing occludin in SLK-L cells

(Fig. 7e) or knockdown occludin in SLK-S cells (Fig. 7f). Consistently, the

overexpression of occludin in SLK-L cells could inhibit cell migration and

invasion (Fig. 7g), whereas knockdown occludin in SLK-S cells promoted cell

migration and invasion (Fig. 7j). The findings suggest that SLK-L could function

as an oncogenic protein by disrupting the interaction between SLK-S and

occludin. The overexpression of the SLK long isoform may destabilize tight

junctions (TJs), resulting in irregular cell fate. These results support the notion

that high levels of SLK-L expression are associated with carcinogenicity and

the promotion of cancer metastasis, providing valuable insight into this

phenomenon.

Figure 7 SLK isoforms can influence cell migration partially through their different binding abilities to cell junction markers. **a,** SLK isoforms demonstrate same cellular localization. **b,** The expression level of EMT markers in different SLK cell lines. **c,** SLK isoforms showed varied binding abilities with occludin. **d,** The subcellular localization of SLK and Occludin. Scale bar:20um. **e-j,** Regulation occludin affect cell migration and invasion in indicated SLK cell lines. **e,** Overexpression occludin in SLK-L cells. **f,** Knockdown the expression of occlidin in SLK-S cells. Cell migration (**g,i;** n=3), and invasion (**h,j;** n=3; scale bar:200μm).*P<0.05, **P<0.01, ***P<0.001, ****P<0.0001, ns:not significant. One-way ANOVA followed by Tukey's test was used in **g-j**. The chart represents mean±SD.

The RNA binding protein Rbfox2 regulates SLK alternative splicing

We next proceeded to investigate the regulatory mechanism responsible for the AS of SLK. In addition to the regulation by the core spliceosome [33], AS is also controlled by the cooperation of multiple regulatory *cis*-elements that recruit *trans*-acting splicing factors to affect proximal splice sites (SS) through multifarious mechanisms [33-36]. Since the AS of SLK exhibited significant effects on cancer development, we aim to investigate how SLK splicing is regulated. We downloaded RBP binding peaks from the ENCODE eCLIP-seq dataset [37] and identified the potential splicing factors that could regulate the splicing of SLK 13[th] exon. As a result, three splicing factors were discovered to have binding peaks near the 12[th] intron and the 13[th] exon of SLK, including Rbfox2, QKI, and HNRNPA1 (Fig. 5a and S4a). We overexpressed these splicing factors to test their function. Rbfox2 overexpression is found to change the AS of SLK while QKI and HNRNPA1 exhibited less influence (Fig. S4b). In addition, the RNA immunoprecipitation (RIP) assay confirmed the binding of Rbfox2 to SLK RNA (Fig. 5b). Rbfox2 is a member of the RNA-binding Fox proteins (Rbfox) family which regulates splicing by binding to the motif (5'-UGCAUGU-3') on the pre-RNAs [38-41]. The expression of Rbfox2 is found to decrease in the cancer types in which the PSI of SLK increased (Fig. 5c).

To test whether Rbfox2 inhibits SLK splicing by binding to its pre-RNA, we checked the sequence of SLK RNA and found two Rbfox2 binding motifs in the 12[th] intron. Then we constructed SLK splicing reporters with the wildtype or mutated Rbfox2 binding motifs to assess the regulatory mechanism of Rbfox2 (Fig. 5d). Subsequently, we tested the splicing of SLK reporters in cells undergoing Rbfox2 overexpression or knockdown. As expected, in Rbfox2-overexpressed cells, AS of the wild-type SLK reporter tended to skip

the 13th exon, producing more SLK-S. However, the skipping ability of the

mutated reporter was attenuated, leaving the splicing ratio similar to the mock

control (Fig. 5e). On the contrary, the SLK-WT reporter tended to retain the

13th exon in Rbfox2 knockdown cells, resulting in the AS shift towards SLK-L

whereas the splicing of mutated reporters remained unchanged with Rbfox2

deficiency (Fig. 5f). The reporter gene assay confirmed that Rbfox2 regulates

SLK alternative splicing by binding to the motif sequence 5'-UGCAUG-3' in the

12th intron to promote the skipping of SLK 13th exon, resulting in lower PSI and

decreased SLK long isoform.

In addition, we tested the regulation of Rbfox2 to endogenous AS of SLK.

Rbfox2 was either overexpressed or knocked down in cancer cells to regulate

the AS. The results were consistent with that of the SLK reporter. Endogenous

13th exon of SLK skipped after overexpression of Rbfox2 as compared to

nonspecific and negative controls (Fig. 5g, left), whereas it was included after

knockdown of Rbfox2 by siRNA (Fig. 5g, right). Collectively, the results

demonstrated that Rbfox2 can regulate SLK AS by binding to its recognition

motif on the SLK 12th intron.

Figure 5 Rbfox2 regulates the splicing of SLK. **a,** Rbfox2 binding peaks was
detected in front of the SLK exon 13[th]. **b,** The RIP assay showed the binding of
Rbfox2 to SLK RNA. **c,** Rbfox2 expression level in tumors versus normal
tissues across various cancer types from TCGA data. **d,** The minigene
reporter of SLK wildtype and mutants were constructed. **e-f.** Rbfox2 regulates
SLK splicing through its binding motif. **e,** Rbfox2 overexpression inhibited SLK
minigene splicing. **f,** Rbfox2 knockdown promoted SLK minigene splicing. **g,**
Rbfox2 regulates splicing of the endogenous *SLK* gene. RNA loading control:
GAPDH; protein loading control: GAPDH or Tubulin. The chart represents
mean±SD. *P<0.05, **P<0.01, ***P<0.001, ****P<0.0001. Two-way ANOVA
was applied to analyze the difference among means in E and F while one-way
ANOVA was used in G.

Discussion

438
439
More than 90% of human genes undergo alternative splicing [42]. After

eliminating noises caused by tissue-specific splicing, numerous gene-splicing

alterations have been detected in a large amount of cancer patients [9].

Abnormal splicing is considered a molecular hallmark of cancer and thus worth

efforts in investigating the underlying mechanisms, the cellular consequences,

and the possibility of manipulation, for cancer treatment. In this study, we have

demonstrated that the significant and extensive alterations across multiple

cancer types are involved in actin filament organization, cell junction assembly

and cell-substrate adhesion, which suggests a universal functional and

mechanistic implications. We demonstrated SLK AS is one of the crucial

splicing events that are dysregulated in cancer and splicing switch to SLK-S

results in cancer inhibition.

As a protein kinase, previous studies of SLK focused on its catalytic

activities. Recently, a few studies have demonstrated that some cellular

functions of SLK appears to be kinase activity independent, suggesting that it

may be an important scaffold for signal transduction pathways [14]. In some

cases, overexpression of scaffolding proteins often leads to the titration of

signaling components, while their knockdown interferes with the formation of

signaling complexes, resulting in similar phenotypes [43]. Our study of the AS of

SLK by coincidence shed light on these discoveries, by implying the possibility

that this may not be a regulation of the splicing level rather than the expression

level.

Interestingly, in the animal experiments of testing AS of SLK, our study

found that SLK knockout cell lines showed similar proliferation rate in the nude

mice compared to SLK-WT though knocking out SLK is embryonic lethal. On

one hand, it indicates an undiscovered complementary mechanism upon SLK

knockout. On the other hand, it implies that targeting the expression of SLK

may be therapeutically ineffective. In contrast, we discovered the SLK-S cells

produced smaller subcutaneous tumors in tumorigenesis experiments and

significant less metastasis burden than other SLK groups. Collectively, these

findings indicate that AS of the *SLK* gene has the potential to be a target for

tumor diagnosis and therapy, which need further validation.

We have discovered that occludin may have different binding abilities to

different SLK isoforms, However, the precise mechanism of how they interact

with each other remained for further exploration. Most occludin functions were

initially   attributed   to   conformational   changes   following   selective

phosphorylation [44], hence more solid validations between SLK AS and

occludin are required.

The mechanisms of the SLK splicing were found to be comprehensive.

The AS of SLK was proved to be regulated by Rbfox2 through the binding

motif 5'-UGCAUG-3' on SLK pre-RNA. However, the expression of Rbfox2

was not simultaneously altered in some cancer types compared with SLK

splicing (Fig. 5c), implying the AS of SLK is regulated in a complicated manner.

Moreover, though overexpression of Rbfox2 can suppress SLK exon 13[th]

splicing, its regulation did not exhibit significant tumor inhibition in vitro (data

not shown). In addition, Rbfox2 is a ubiquitous RNA binding protein that have

various RNA targets other than SLK, thus therapeutics strategies targeting

Rbfox2 to regulate SLK alternative does not have promising outlook.

Nevertheless, Rbfox2 form a large assembly of splicing regulators (LASR) [45, 46],

thus the components of LASR also have the potential in affecting SLK splicing.

In summary, the expression levels of SLK isoforms showed drastic

variations in tumor cells and normal cells. We found that different isoforms of

the *SLK* gene had different biological functions. SLK long isoform is crucial for

cancer maintenance and development. SLK isoforms might affect cell

functions by the different binding abilities of occludin. Rbfox2 inhibited SLK

exon 13 splicing through its binding motif. Collectively, our study uncovers the

biological consequences and underlying mechanisms for one of the most

mis-spliced genes in cancer, highlighting its potential significance in cancer

diagnosis and treatment. By taking *SLK* as an example, we hope to draw

attention to the importance of splicing regulation in cancer.

Materials and Methods

Plasmid construction

The CDS sequences of SLK-L and SLK-S were amplified from the cDNA of HCT116 cells. The two SLK isoforms were cloned into the pcDNA3.1 overexpression vector with the HA or V5 tag sequence. The CDS regions of SLK isoforms with the HA or Flag tag or the firefly luciferase CDS were inserted after the CMV promoter of the pCDH backbone plasmid to create lentivirus vectors, respectively. The Rbfox2 CDS region with the HA tag was cloned into the pcDNA3.1 backbone to produce Rbfox2 overexpression vectors.

For the SLK minigene reporter, we cloned SLK exon 12, 13, 14, a 337-nt sequence with 5'-SS, a 290-nt sequence containing branch point and 3'-SS of intron 12, and a fragment of intron 13, into the pcDNA3.1 plasmid backbone. The SLK mutation vectors were constructed by replacing the Rbfox2 binding motif 5'-TGCATGT-3' with 5'-ACGTACA-3'.

To generate stable cell lines with SLK-S or SLK knockout, we designed guide RNAs (gRNAs) targeting specific sites of SLK genome DNA and cloned them into the CRISPR-Cas9 plasmid. For cancer cell lines selectively expressed SLK-L, an additional template of exon13-14 or 12,14 were supplied as the donor DNA at the target site to allow the cells to repair the double-strand break through homology-directed repair (HDR) pathways (Fig. 2a, Fig. 3a and Table S1).

Co-immunoprecipitation

293T or HCT116 cells were seeded in a 15-cm dish and transfected with HA-tagged SLK isoforms and V5-tagged candidate proteins. The PBS-washed

cells were used after transfection for 48 h. The lysis buffer (P0013, Beyotime)

was then added into the cells and the supernatants were collected after

centrifuging at 15000 rpm at 4 °C for 30 min. Next, 10% supernatant was

retained as input and the pallet was resuspended. Primary antibodies (HA-Tag

Mouse mAb #2367 and V5-Tag Rabbit mAb #13202, CST) were added and

incubated overnight. The antibodies were diluted at the ratio of 1:100.

Subsequently, 50 μl protein A/G agarose beads were added to each tube and

incubated at room temperature for 1 h. Following centrifugation at 2500 rpm for

5 min, the supernatants were removed, and the agarose beads were

resuspended in 500 μl IP buffer and washed at room temperature 3 times.

After the washing step, 50 μl of 2×SDS protein loading buffer was added, and

the denatured proteins were obtained by incubation at 98 °C for 10 min.

RNA Immunoprecipitation

Cells grown in a 15-cm dish (90% confluence) were digested with 3 mL of

0.25% trypsin at 37 °C for 2 min. Next, 5 mL of complete medium was added to

the dish to terminate digestion. The collected cells were subjected to RNA

immunoprecipitation   following   the   Abcam   protocol

(https://www.abcam.com/epigenetics/rna-immunoprecipitation-rip-protocol).

Cancer cell development and function

Cell proliferations were analyzed by cell counting or CCK-8 assay. For cell

counting, 5000 cells were seeded per well on 24-well plates, and the cell

number was counted at 24, 48, 72, and 96 h after incubation. For the CCK-8

assay, 1000 cells were seeded per well on 96-well plates, and the Cell

Counting Kit-8 (CCK-8, 40203ES60, Yeason) was used to determine the cell

proliferation.

A colony formation assay was carried out to test the tumor formation

ability of cancer cells. In brief, 2000 cells were seeded onto a 6-well plate in a

complete medium with 10% FBS for one week. The cells were then dyed with

crystal violet for 30 min before washing with PBS. The plating efficiency was

calculated using colony numbers divided by seeded cell numbers.

Cell migration ability was measured through wound healing experiments.

Cells were seeded onto a 6-well plate. After reaching 90% confluence, the

cells were scratched with a 200-μl pipette tip. The images were captured at 0,

12, 24, and 48hr after scratching. The average value of wound closure was

quantified by Image J software. Experiments were performed in triplicates.

Transwell invasion assay was used to determine the invasion ability of

cells. Matrigel Matrix (356234, Corning) was thawed at 4 °C and mixed with

medium without FBS at the ratio of 1:8. Next, 50 μl of the mixture was added to

each Transwell and incubated at 37 °C for 1 h. Subsequently, 1×10^5 cells

(2×10^5 cells in Fig 3F) were seeded in each Transwell in 200 μl medium

without FBS, and 500 μl medium with 10% FBS was added as an inducer in

the 24-well-plate. The cells were cultured at 37 °C for 24 h. The matrix and

media were removed, and the cells were incubated with 4% paraformaldehyde

(PFA) for 30 min after washing with PBS. After removing PFA, the cells were

washed with PBS twice. The cells were then stained with crystal violet for 30

min and de-stained with PBS twice before imaging. Cells were counted from

the captured images by using Image J software. Each experiment was

repeated three times.

Lentivirus packaging and infection

HEK 293T cells were plated at 50% confluence in a 10-cm dish before

transfection with plasmids. The envelope plasmid pMD2.G, the packaging

plasmid psPAX2, and the transfer plasmids were co-transfected into HEK

293T cells by Lipofectamine 3000. The mediums with lentivirus from

transfected cells were collected after 48 h of transfection and filtered through a

0.22-μm membrane.

Immunofluorescence assay

293T cells were seeded in a Nunc^{TM} dish (150680, Thermo) and

transfected with HA-tagged SLK isoforms and V5-tagged occludin (for Fig 6C)

or SLK different isoforms with HA or V5 tag (for Fig S5). Cells were incubated

with 4% PFA for 1h. After being washed with PBS, cells were incubated with

1mL 0.25% Triton^{TM} X-100 (9036-19-5, Meck) per dish for 10 min. Then cells

were washed with PBS twice and incubated with HA and V5 antibodies

(dilution ratio: 1:1000) at 4°C overnight. Alexa Fluor® 594 (ab 150080, Abcam)

and Alexa Fluor® 488 (ab 150077, Abcam) were Incubated with cells at room

temperature for 1h. The nuclei were stained with DIPA (P36985, Invitrogen™)

and observed and photographed under Zeiss LSM880.

Xenograft tumor formation

We conducted two biological replicants of tumorigenesis experiments in

nude mice. In the first replicant, 5×10^6 cells were resuspended in PBS and

injected subcutaneously into a BALB/c-nude mouse (10 mouse/group). The

body weight and tumor volume were measured twice a week after injection. In

SLK-WT and SLK-L groups, several mice were sacrificed before the endpoint

due to the fast-growing tumors and the following tissue rupture. In the second

biological replicant, 1×10^6 cells were resuspended in PBS and injected

subcutaneously into a BALB/c nude mouse. The tumor volume was measured

every three days. The experiment was terminated 33 days after injection. The

tumor volume was calculated as volume $(mm^3) = L \times W^2/2$, where L is the

largest diameter and W is the smallest diameter of the tumor. After the

experiment, the mice were euthanized, and the tumor was dissected.

Tail vein metastasis experiment

Tail vein metastasis experiments were carried out in both BALB/c nude

mice and C-NKG mice at Cyagen Co., Ltd (SuZhou, China). Briefly,

luciferase-labeled HCT116 cells (SLK-WT, SLK-L, or SLK-S) were

resuspended in 100 μL PBS at concentrations of 1.0×10^6 or 2.0×10^6 and

injected into the lateral tail veins of seven-week-old C-NKG or BALB/c nude

female mice between 18-22 g, respectively. The mice were anesthetized and

injected intraperitoneally with D-luciferin (Yeasen, China) for a dose of 150

mg/kg on the indicated day post-injection. Metastasis was quantified by total

flux (photons/second) of bioluminescence in the lungs on a PerkinElmer IVIS

Lumina III *in vivo* imaging system.

Bioinformatic analyses

To identify the cancer-related alternative splicing from TCGA, we

downloaded the analyzed data from the published database (IDEAS), which

calculate the PSI value with rMATs. We filtered out cancer types with fewer

than 20 normal tissue counterparts to ensure robust and reliable estimation of

PSI changes. The Wilcox test was applied to determine the significance

between cancer and normal groups. For alternative splicing events shown in

Figure S1a, we only included the significantly changed splicing events with ≥10

read counts, and ≥5 observations in normal samples. We downloaded the level

3 RNA-seq data from TCGA (http://gdac.broadinstitute.org/), and the splicing

events shared by more than 10 cancer types were analyzed in the GO

enrichment network.

CLIP-seq analysis

The processed signal files for eCLIP-seq of all RBPs were downloaded

from the ENCODE consortium. To visualize binding peaks of the indicated

RBPs, we utilized the R packages 'Gviz' and 'rtracklayer'. For Figure 5a, the

HepG2 and K562 data were downloaded from the ENCODE consortium, and

the HEK293 data were obtained from the GEO database (GSE77629). All

eCLIP-seq data were aligned to GENCODE hg38 v39 using STAR and then

visualized using karyoploteR.

RNA-seq analysis of SLK cell types

For the RNA-seq analysis, we used trim galore to remove the adapter

from sequencing data, followed by gene expression calculation using RSEM

based on GENCODE hg38 v39. Gene count data were imported using the

tximport package and normalized by the TMM method from NOISeq. NOISeq

was used to obtain differentially expressed genes from different cell lines, with

q=0.8 and |M| > 0.5 using in NOISeq analysis. ClusterProfiler package was

used to perform GO enrichment analysis, and ggplot2 was applied to visualize

the results. The Venn Diagram was constructed using the 'VennDiagram'

package. Gene sets that were differentially expressed in both L vs S and L vs

WT, L vs S and S vs WT, and L vs S and KO vs WT were further investigated

as potential targets. The gene expression profiles of these genes were plotted

using 'heatmap'.

Structure prediction of SLK-L and SLK-S

Parafold [47], a paralyzed accelerated version of alphafold2 [24] was applied to

predict the structure of SLK-L and SLK-S. The input is the RNA sequence of

the open reading frame of each isoform. Deepmind monomer model sets were

employed for generating the prediction. The top 5 ranked results were

investigated according to Alphafold default evaluation.

Data Availability

eCLIP data of RBPs and the processed signal files (in bigwig format) were

downloaded from ENCODE consortium (https://www.encodeproject.org).

RNA- seq data during the current study are available in the NODE data

depository. Custom computer code and original data are available upon

request.

Acknowledgments

We thank Prof. Yong-Bo Wang of Fudan University and Prof. Yang Wang of Dalian Medical University for the manuscript revision. We thank Ms. Yun Jiang, Dr. Yujie Chen, Ms. Nianhao Cheng, and Ms. Shuang Han for their assistance with the experiments. We are grateful for the experimental support of the UliSchwarz Quantitative Biology Core Facility in Bio-Med Big Data Center, SINH, CAS. This work is supported by the National Natural Science Foundation of China (32171294) and Shanghai Municipal Natural Science Foundation (20ZR1467300) to H-H. Wei. It is also supported by the National Natural Science Foundation of China (31730110, 91940303, and 32030064), the National Key Research and Development Program of China (2018YFA0107602), and the Strategic Priority Research Program of the Chinese Academy of Sciences (XDB38040100) to Z. Wang.

Author contributions

H-H. Wei and Z. Wang conceived the project. H-H Wei designed the experiments and interpreted the results. Y-Q. Yang, H-H. Wei and J. Sun carried out molecular, biochemical, and cell function experiments. Y. Hu and S-R. Zhang analyzed the eCLIP-seq, TCGA data, and RNA-seq data and performed other bioinformatic analysis. H-H. Wei was responsible for the tail vein injection experiment. W-J. Zhang and Y. Sun performed the mice xenograft experiment. J-F. Li and J-W. Guan performed protein structure prediction. X-Y. Feng generated the CRISPR HCT116 cell lines. H-H. Wei, Y-Q. yang and Y. Hu wrote the manuscript. Y. Yang helped to interpret the data.

Conflict of interest

686   The authors declare that they have no conflict of interest.

687   **References**

688   1.   Coltri, P.P., dos Santos, M.G.P. & da Silva, G.H.G. Splicing and cancer:
689   Challenges and opportunities. *Wires Rna* **10** (2019).
690   2.   Dong, X. & Chen, R. Understanding aberrant RNA splicing to facilitate
691   cancer diagnosis and therapy. *Oncogene* **39**, 2231-2242 (2020).
692   3.   Wang, B.D. & Lee, N.H. Aberrant RNA Splicing in Cancer and Drug
693   Resistance. *Cancers (Basel)* **10** (2018).
694   4.   Oltean, S. & Bates, D.O. Hallmarks of alternative splicing in cancer.
695   *Oncogene* **33**, 5311-5318 (2014).
696   5.   Wang, Y. *et al.* The Splicing Factor RBM4 Controls Apoptosis,
697   Proliferation, and Migration to Suppress Tumor Progression. *Cancer Cell*
698   **26**, 374-389 (2014).
699   6.   Qi, Y. *et al.* A splicing isoform of TEAD4 attenuates the Hippo-YAP
700   signalling to inhibit tumour proliferation. *Nat Commun* **7**, ncomms11840
701   (2016).
702   7.   Martinez-Montiel, N., Rosas-Murrieta, N.H., Anaya Ruiz, M.,
703   Monjaraz-Guzman, E. & Martinez-Contreras, R. Alternative Splicing as a
704   Target for Cancer Treatment. *Int J Mol Sci* **19** (2018).
705   8.   Cherry, S. & Lynch, K.W. Alternative splicing and cancer: insights,
706   opportunities, and challenges from an expanding view of the
707   transcriptome. *Genes Dev* **34**, 1005-1016 (2020).
708   9.   Tsai, Y.S., Dominguez, D., Gomez, S.M. & Wang, Z. Transcriptome-wide
709   identification and study of cancer-specific splicing events across multiple
710   tumors. *Oncotarget* **6**, 6825-6839 (2015).
711   10.   O'Reilly, P.G. *et al.* The Ste20-like kinase SLK is required for cell cycle
712   progression through G2. *J Biol Chem* **280**, 42383-42390 (2005).
713   11.   Ellinger-Ziegelbauer, H. *et al.* Ste20-like kinase (SLK), a regulatory kinase
714   for polo-like kinase (Plk) during the G2/M transition in somatic cells.
715   *Genes Cells* **5**, 491-498 (2000).

12. Hao, W. *et al.* Induction of apoptosis by the Ste20-like kinase SLK, a
germinal center kinase that activates apoptosis signal-regulating kinase
and p38. *J Biol Chem* 281, 3075-3084 (2006).
13. Pryce, B.R. *et al.* Muscle-specific deletion of SLK/Stk2 enhances p38
activity and myogenesis in mdx mice. *Biochim Biophys Acta Mol Cell Res*
1868, 118917 (2021).
14. Garland, B., Delisle, S., Al-Zahrani, K.N., Pryce, B.R. & Sabourin, L.A.
The Ste20-like kinase - a Jack of all trades? *J Cell Sci* 134 (2021).
15. Guilluy, C. *et al.* Ste20-related kinase SLK phosphorylates Ser188 of
RhoA to induce vasodilation in response to angiotensin II Type 2 receptor
activation. *Circ Res* 102, 1265-1274 (2008).
16. Storbeck, C.J. *et al.* The Ldb1 and Ldb2 transcriptional cofactors interact
with the Ste20-like kinase SLK and regulate cell migration. *Mol Biol Cell*
20, 4174-4182 (2009).
17. Conway, J., Al-Zahrani, K.N., Pryce, B.R., Abou-Hamad, J. & Sabourin,
L.A. Transforming growth factor beta-induced epithelial to mesenchymal
transition requires the Ste20-like kinase SLK independently of its catalytic
activity. *Oncotarget* 8, 98745-98756 (2017).
18. Baron, K.D. *et al.* Recruitment and activation of SLK at the leading edge of
migrating cells requires Src family kinase activity and the LIM-only protein
4. *Biochim Biophys Acta* 1853, 1683-1692 (2015).
19. Machicoane, M. *et al.* SLK-dependent activation of ERMs controls
LGN-NuMA localization and spindle orientation. *J Cell Biol* 205, 791-799
(2014).
20. Quizi, J.L. *et al.* SLK-mediated phosphorylation of paxillin is required for
focal adhesion turnover and cell migration. *Oncogene* 32, 4656-4663
(2013).
21. Panneton, V. *et al.* Regulation of Catalytic and Non-catalytic Functions of
the Drosophila Ste20 Kinase Slik by Activation Segment Phosphorylation.
J Biol Chem 290, 20960-20971 (2015).
22. Roovers, K. *et al.* The Ste20-like kinase SLK is required for ErbB2-driven
breast cancer cell motility. *Oncogene* 28, 2839-2848 (2009).

23. Johnson, T.M., Antrobus, R. & Johnson, L.N. Plk1 activation by Ste20-like
kinase (Slk) phosphorylation and polo-box phosphopeptide binding
assayed with the substrate translationally controlled tumor protein (TCTP).
Biochemistry **47**, 3688-3696 (2008).

24. Tunyasuvunakool, K. *et al.* Highly accurate protein structure prediction for
the human proteome. *Nature* **596**, 590-596 (2021).

25. Consortium, G.T. The GTEx Consortium atlas of genetic regulatory effects
across human tissues. *Science* **369**, 1318-1330 (2020).

26. Chaffer, C.L. & Weinberg, R.A. A perspective on cancer cell metastasis.
Science **331**, 1559-1564 (2011).

27. Wagner, S., Flood, T.A., O'Reilly, P., Hume, K. & Sabourin, L.A.
Association of the Ste20-like kinase (SLK) with the microtubule. Role in
Rac1-mediated regulation of actin dynamics during cell adhesion and
spreading. *J Biol Chem* **277**, 37685-37692 (2002).

28. Al-Zahrani, K.N., Baron, K.D. & Sabourin, L.A. Ste20-like kinase SLK, at
the crossroads. *Cell Adhesion & Migration* **7**, 1-10 (2013).

29. Wagner, S. *et al.* FAK/src-family dependent activation of the Ste20-like
kinase SLK is required for microtubule-dependent focal adhesion turnover
and cell migration. *PLoS One* **3**, e1868 (2008).

30. Reticker-Flynn, N.E. *et al.* Lymph node colonization induces
tumor-immune tolerance to promote distant metastasis. *Cell* **185**,
1924-1942 e1923 (2022).

31. Shapiro, I.M. *et al.* An EMT-driven alternative splicing program occurs in
human breast cancer and modulates cellular phenotype. *PLoS Genet* **7**,
e1002218 (2011).

32. Arnouk, H., Yum, G. & Shah, D. Cripto-1 as a Key Factor in Tumor
Progression, Epithelial to Mesenchymal Transition and Cancer Stem Cells.
Int J Mol Sci **22** (2021).

33. Matera, A.G. & Wang, Z. A day in the life of the spliceosome. *Nat Rev Mol*
Cell Biol **15**, 108-121 (2014).

34. Sveen, A., Kilpinen, S., Ruusulehto, A., Lothe, R.A. & Skotheim, R.I.
Aberrant RNA splicing in cancer; expression changes and driver
mutations of splicing factor genes. *Oncogene* **35**, 2413-2427 (2016).

35. Biamonti, G., Catillo, M., Pignataro, D., Montecucco, A. & Ghigna, C. The
alternative splicing side of cancer. *Semin Cell Dev Biol* 32, 30-36 (2014).

36. Wang, Y., Bao, Y., Zhang, S. & Wang, Z. Splicing dysregulation in cancer:
from mechanistic understanding to a new class of therapeutic targets. *Sci*
China Life Sci 63, 469-484 (2020).

37. Consortium, E.P. An integrated encyclopedia of DNA elements in the
human genome. *Nature* 489, 57-74 (2012).

38. Choi, S., Sa, M., Cho, N., Kim, K.K. & Park, S.H. Rbfox2 dissociation from
stress granules suppresses cancer progression. *Exp Mol Med* 51, 1-12
(2019).

39. Braeutigam, C. *et al.* The RNA-binding protein Rbfox2: an essential
regulator of EMT-driven alternative splicing and a mediator of cellular
invasion. *Oncogene* 33, 1082-1092 (2014).

40. Venables, J.P. *et al.* RBFOX2 Is an Important Regulator of Mesenchymal
Tissue-Specific Splicing in both Normal and Cancer Tissues. *Molecular*
and Cellular Biology 33, 396-405 (2013).

41. Huh, G.S. & Hynes, R.O. Regulation of alternative pre-mRNA splicing by a
novel repeated hexanucleotide element. *Genes Dev* 8, 1561-1574 (1994).

42. Pan, Q., Shai, O., Lee, L.J., Frey, B.J. & Blencowe, B.J. Deep surveying of
alternative splicing complexity in the human transcriptome by
high-throughput sequencing. *Nat Genet* 40, 1413-1415 (2008).

43. Duff, D. & Long, A. Roles for RACK1 in cancer cell migration and invasion.
Cell Signal 35, 250-255 (2017).

44. Blasig, I.E. *et al.* Occludin protein family: oxidative stress and reducing
conditions. *Antioxid Redox Signal* 15, 1195-1219 (2011).

45. Ying, Y. *et al.* Splicing Activation by Rbfox Requires Self-Aggregation
through Its Tyrosine-Rich Domain. *Cell* 170, 312-323 e310 (2017).

46. Damianov, A. *et al.* Rbfox Proteins Regulate Splicing as Part of a Large
Multiprotein Complex LASR. *Cell* 165, 606-619 (2016).

47. Zhong, B. *et al.* in International Conference on High Performance
Computing in Asia-Pacific Region Workshops 1-9 (2022).

812